Identification of bis-benzylisoquinoline alkaloids as SARS-CoV-2 entry

inhibitors from a library of natural products in vitro

Running title: SARS-CoV-2 entry inhibitors

Chang-Long He[1, #], Lu-Yi Huang[1, #], Kai Wang[1, #], Chen-Jian Gu[2], Jie Hu[1], Gui-Ji

Zhang[1], Wei Xu[2], You-Hua Xie[2,*], Ni Tang[1,*], Ai-Long Huang[1,*]

[1]Key Laboratory of Molecular Biology for Infectious Diseases (Ministry of

Education), Institute for Viral Hepatitis, Department of Infectious Diseases, The

Second Affiliated Hospital, Chongqing Medical University, Chongqing, 400010,

China

[2]Key Laboratory of Medical Molecular Virology (MOE/NHC/CAMS), Department of

Medical Microbiology and Parasitology, School of Basic Medical Sciences, Shanghai

Medical College, Fudan University

[#]These authors contributed equally to this work.

***Corresponding authors:**

Ai-Long Huang, **Ni Tang**, Key Laboratory of Molecular Biology for Infectious

Diseases (Ministry of Education), Institute for Viral Hepatitis, Department of

Infectious Diseases, The Second Affiliated Hospital, Chongqing Medical University,

Chongqing, China. Phone: 86-23-68486780, Fax: 86-23-68486780, E-mail:

ahuang@cqmu.edu.cn (A.L.H.), nitang@cqmu.edu.cn (N.T.); **You-Hua Xie**, Key

Laboratory of Medical Molecular Virology (MOE/NHC/CAMS), School of Basic

Medical Sciences, Shanghai Medical College, Fudan University, e-mail,

yhxie@fudan.edu.cn

Abstract

Coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) is a major public health issue. To screen for antiviral

drugs for COVID-19 treatment, we constructed a SARS-CoV-2 spike (S) pseudovirus

system using an HIV-1-based lentiviral vector with a luciferase reporter gene to

screen 188 small potential antiviral compounds. Using this system, we identified nine

compounds, specifically, bis-benzylisoquinoline alkaloids, that potently inhibited

SARS-CoV-2 pseudovirus entry, with EC_{50} values of 0.1–10 μM. Mechanistic studies

showed that these compounds, reported as calcium channel blockers (CCBs),

inhibited Ca^{2+}-mediated membrane fusion and consequently suppressed coronavirus

entry. These candidate drugs showed broad-spectrum efficacy against the entry of

several coronavirus pseudotypes (SARS-CoV, MERS-CoV, SARS-CoV-2 [S-D614

and S-G614]) in different cell lines (293T, Calu-3, and A549). Antiviral tests using

native SARS-CoV-2 in Vero E6 cells confirmed that four of the drugs

(SC9/cepharanthine, SC161/hernandezine, SC171, and SC185/neferine) reduced

cytopathic effect and supernatant viral RNA load. Among them, cepharanthine

showed the strongest anti-SARS-CoV-2 activity. Collectively, this study offers new

lead compounds for coronavirus antiviral drug discovery.

Introduction

The novel coronavirus reported in 2019 (2019-nCoV), officially named severe acute

respiratory syndrome coronavirus 2 (SARS-CoV-2), is a new type of coronavirus

belonging to the genus *Betacoronavirus*. It is a single-stranded RNA virus with a

genome of approximately 29 Kb and with a high pathogenicity and high infectivity.[1-3]

As of 8 December 2020, there were nearly 68 million confirmed cases of coronavirus

disease 2019 (COVID-19) globally, including more than 1.5 million confirmed deaths,

and the number of countries, areas, or territories with cases had reached 220.[4]

Although some countries have contained the pandemic, the number of confirmed

cases and deaths is expected to rise globally. Currently, no licensed specific antiviral

drugs are available to combat the infection. The development of effective viral

inhibitors and antibody-based therapeutics for COVID-19 is a high global priority.

The SARS-CoV-2 RNA genome encodes 29 structural and non-structural

proteins, including spike (S), envelope (E), membrane (M), nucleocapsid (N) proteins,

and the ORF1a/b polyprotein.[5] The S glycoprotein embedded as a trimer in the virus

envelope is a key structural protein that mediates SARS-CoV-2 entry into host cells.[6]

SARS-CoV-2 S protein comprises two functional subunits, of which S1 is responsible

for binding to host-cell receptors and S2 mediates the fusion of viral and cell

membranes. Binding of the receptor-binding domain (RBD) located in the S1 subunit

of SARS-CoV-2 S protein to the receptor angiotensin converting enzyme 2 (ACE2) is

crucial for viral entry into cells.[7-9] Blockade of SARS-CoV-2 entry has become an

attractive therapeutic strategy as it prevents the initiation of any steps towards viral

replication. To date, antibodies blocking S–ACE2 interaction have proven to be the

most effective for COVID-19 treatment.[10-12] However, their high cost limits their

broad application. Thus, small molecule-based drugs targeting virus entry steps are

highly anticipated.

Efforts to find SARS-CoV-2 entry inhibitors have led to the identification of

several drug candidates. Lipopeptides targeting the HR1 domain in S2 significantly

inhibited SARS-CoV-2 entry.[13-15] Host proteases furin, TMPRSS2, and cathepsin B/L

help cleave S, thus facilitating the entry process. Inhibition of their activities by

small-molecule inhibitors such as camostat or E-64d blocks SARS-CoV-2

infection.[16-18] Apilimod, an inhibitor of the lipid kinase PIKfyve, inhibits viral

infection by preventing the release of viral contents from endosomes.[19] Although

some progress has been made toward SARS-CoV-2 entry inhibitors, there is still more

work to be done.

In this study, we successfully constructed a high-titer SARS-CoV-2 S

pseudovirus system using an HIV-1-based lentiviral vector with a luciferase reporter

gene for the screening of specific viral entry inhibitors. Based on this classic

pseudovirus model, a small potential antiviral compound library was screened. We

identified several compounds, especially, bis-benzylisoquinoline alkaloids, that

potently inhibited SARS-CoV-2 pseudovirus entry, with EC_{50} values of 0.1–10 μM.

Subsequently, we preliminarily explored the underlying mechanisms of the hit

compounds, and their antiviral activity against SARS-CoV-2 was confirmed in Vero

E6 cells with authentic virus. We identified compounds with effective antiviral

activity, thereby providing new lead compounds for coronavirus antiviral drug

discovery.

Results

Screening of SARS-CoV-2 S-G614 entry inhibitors in 293T-ACE2 cells

The S protein variant D614G (a single change in the genetic code; D = aspartic acid, G = glycine) carrying the amino acid mutation D614G has become the most prevalent form in the global pandemic and has been associated with increased infectivity of SARS-CoV-2.[20-22] Using a luciferase-expressing pseudovirus encoding SARS-CoV-2 S (G614) protein, a library of 188 chemical compounds (Supplementary Table S1, most were natural compounds) was screened in 293T-ACE2 cells to find novel anti-SARS-CoV-2 entry inhibitors. A workflow chart of screening is shown in Fig. 1a. 293T-ACE2 cells were pretreated with 20 μM of each compound for 1 h and then, a SARS-CoV-2 S-G614 pseudovirus (3.8×10^4 copies) mixture containing 20 μM of each compound was added to the cells. A known entry inhibitor, E-64d (targeting host cathepsin B/L), was used as a positive control and DMSO as a negative control. Luciferase activity was measured to assess the inhibitory effect of the 188 compounds on S-G614 pseudovirus entry 72 h post-infection. After a preliminary screening, 41 compounds associated with a relative infection rate <30% (Fig. 1b) were identified. Half of these displayed cytotoxicity at 20 μM. We selected 19 compounds with low cytotoxicity for further testing. The chemical structures of the 19 compounds are shown in Supplementary Fig. S1. The selectivity index values in terms of cytotoxicity were greater than 2, suggesting that they may be safe *in vivo*. As the S-G614 pseudovirus was generated in the envelope-defective HIV-1 backbone, vesicular stomatitis virus (VSV) pseudovirus was used as a control to exclude compounds that have anti-HIV activities. Through a filtering strategy, we identified nine compounds

(SC9, SC161, SC171, SC182, SC183, SC184, SC185, SC186, and SC187) that

specifically inhibited S-G614 pseudovirus entry. Except SC171, all these compounds

were bis-benzylisoquinoline alkaloids. The EC_{50} values of the nine compounds

against S-G614 pseudovirus were below 10 μM, but above 20 μM in VSV-G

pseudovirus (Table 1, Supplementary Fig. S2).

Investigation of the treatment timing and antiviral efficacy in different cell lines

Next, we analyzed the relationship between the antiviral efficacy of the nine selected

compounds against S-G614 pseudovirus and the timing of treatment. We divided the

S-G614 pseudovirus entry into three stages: pretreatment, pseudovirus entry, and

pseudovirus post-entry. In total, nine experimental groups were set up for each

compound, including eight treatment groups (A–G) and a control group (Fig. 2a).

Importantly, pretreatment with each compound (group B in Fig. 2) significantly

inhibited S-G614 pseudovirus infection. In the pseudovirus entry stage (group C in

Fig. 2), the compounds exerted similar suppressive effects. However, in the

pseudovirus post-entry stage (group D in Fig. 2), none of the compounds showed any

inhibitory effect. These results indicated that all nine candidates blocked pseudovirus

infection by targeting host factors at the entry stage.

Hoffmann *et al*. highlight that cell lines mimicking important aspects of

respiratory epithelial cells should be used when analyzing the anti-SARS-CoV-2

activity.[23] Hence, we determined their EC_{50} values against S-G614 pseudovirus in

Calu3, A549, and 293T-ACE2 cells (Fig. 3a–i). As shown in Fig. 3, the EC_{50} values of

SC9 (Fig. 3a), SC161 (Fig. 3b), SC171 (Fig. 3c), SC182 (Fig. 3d), and SC185 (Fig.

3g) were below 5 μM in all three cell lines. Unfortunately, the EC_{50} values of SC183

(Fig. 3e), SC184 (Fig. 3f), SC186 (Fig. 3h), and SC187 (Fig. 3i) did not reach the

threshold we had set and thus were excluded from subsequent experiments.

Inhibition of entry of pseudovirus bearing coronavirus S protein by the

candidate compounds

Considering that SARS-CoV-2, SARS-CoV, and MERS-CoV all belong to the genus

Betacoronavirus, we next determined whether the remaining compounds have

broad-spectrum antiviral effects against these coronaviruses, using pseudovirus-based

inhibition assays. We constructed S-D614, S-SARS, and S-MERS pseudoviruses

using the same lentiviral system as S-G614, and then determined the EC_{50} values of

SC9 (cepharanthine, Fig. 4a), SC161 (hernandezine, Fig. 4b), SC171 (Fig. 4c), SC182

(tetrandrine, Fig. 4d), and SC185 (neferine, Fig. 4e) against these pseudoviruses in

293T cells expressing ACE2 or DPP4. Interestingly, SC9, SC161, SC171, and SC185

exhibited highly potent pan-inhibitory activity against coronavirus S-pseudotyped

viruses. The EC_{50} values of the five compounds are shown in Fig. 4f. As SARS-CoV

and SARS-CoV-2 have been reported to enter host cells via binding to ACE2, and

while DPP4 is critical for MERS-CoV entry, it could be ruled out that the five

compounds interfere with ACE2 to block pseudovirus entry.

Interaction between the target compounds and the SARS-CoV-2 RBD domain

Next, we used competitive ELISAs and thermal shift assays to determine whether

these five compounds interact with the SARS-CoV-2 RBD domain. SBP1, one of the

recombinant ACE2 polypeptides, exhibited a weak ability to inhibit the entry of

S-G614 pseudovirus and a weak RBD-binding ability (Fig. 5a, 5b), whereas the

interaction between SC9, SC161, SC171, or SC185 and RBD was negligible (Fig. 5b).

Thermal shift assay results showed that the ΔT_M values of SC161, SC185, and SBP1

were 0 °C, 0.6 °C, and 1.3 °C, respectively (Fig. 5c–d), indicating that SC161 and

SC185 could not stabilize the RBD domain. Thus, the blockade of virus entry by these

candidate compounds is not related to interaction with the SARS-CoV-2 RBD

domain.

Depletion of Ca^{2+} blocks the entry of SARS-CoV-2 S-G614 pseudovirus

Our data indicated that the above five compounds may target host cells to inhibit

SARS-CoV-2 entry. Therefore, we explored host factors via which these five

compounds inhibit virus entry in 293T-ACE2 cells. We examined whether these

compounds perturb cell fusion mediated by SARS-CoV-2 S protein, the initiation

process for virus entry. Compared to DMSO, SC9, SC161, SC182, and SC185 at 5

μM potently inhibited SARS-CoV-2 S-mediated membrane fusion of 293T cells, and

the cell-cell fusion rate in the SC171 treatment group was 40% (Fig. 6a, 6b). Previous

studies have shown that the calcium ion (Ca^{2+}) plays a critical role in SARS-CoV or

MERS-CoV S-mediated fusion to host cell entry.[24,25] Supplementation of the culture

medium with 2 mM $CaCl_2$ enhanced S-G614 pseudovirus entry ability by 2-fold

compared with calcium-free medium (Fig. 6c). In contrast, the entry-blocking effect

of the compounds sharply decreased in calcium-depleted medium (Fig. 6d–e).

Moreover, treatment with 20 μM of BAPTA-AM, a calcium-chelating agent used to

deplete intracellular Ca^{2+}, significantly decreased the entry capacity of S-G614

pseudovirus by 70% (Fig. 6f and 6g). More importantly, when cells were pretreated

with 20 μM BAPTA-AM in combination with one of the four compounds, among

which three bi-benzylisoquinoline alkaloids have been reported as calcium channel

blockers (CCBs),[26-30] the EC_{50} values of these compounds against S-G614

pseudovirus decreased by more than 10-fold (Fig. 6h–i). Intracellular cholesterol

levels increase upon CCB treatment,[31,32] and upregulation of cholesterol biosynthesis

results in blockade of SARS-CoV-2 infection.[33] Consistent herewith, intracellular

cholesterol levels increased in 293T-ACE2 cells pretreated with 5 μM SC9, SC161,

SC171, and SC185 (Fig. 6j). These data indicated that blockade of S-G614

pseudovirus entry by SC9, SC161, SC171, and SC185 mainly depends on calcium

homeostasis.

Cell entry inhibition by the compounds in authentic SARS-CoV-2 infected Vero

E6 cells

The antiviral activities of SC9, SC161, SC171, and SC185 were confirmed in Vero E6

cells infected with native SARS-CoV-2. Changes in cell structure and morphology

caused by SARS-CoV-2 were observed 48 h post-infection. As shown in Fig. 7a,

remdesivir (5 μM), used as a positive control, completely inhibited

SARS-CoV-2-induced cytopathogenic effect (CPE) on Vero E6 cells. SC161

(hernandezine), SC171, and SC185 (neferine) partly inhibited CPE at 10 μM, whereas

SC9 (cepharanthine) completely inhibited CPE at 10 μM. Cell supernatants were

collected at 48 h post-infection for RT-qPCR quantification of SARS-CoV-2 genomic

RNA. The RNA level in the remdesivir (5 μM) treatment group was 0.01% of that in

the DMSO control group (Fig. 7b). The RNA levels in the 10 μM SC9, SC161,

SC171, and SC185 treatment groups were 0.08%, 70.27%, 43.55%, and 76.98%,

respectively, of that in the DMSO control group (Fig. 7c). The results showed that

these compounds inhibited SARS-CoV-2 to varying degrees and may be useful as

leads for SARS-CoV-2 therapeutic drug development.

Discussion

Viral entry into host cells is one of the most critical steps in the life cycle of a virus.

Since blockade of the entry process is a promising therapeutic option for COVID-19,

research attention has been focused on the discovery of viral entry inhibitors. To block

the entry of SARS-CoV-2, the RBD–ACE2 interaction or viral fusion must be

abrogated. To date, a few entry inhibitors against SARS-CoV-2 have been

reported.[19,34-37] ACE2 is the key host receptor mediating SARS-CoV-2 entry. Host

factors including HMGB1 (an ACE2 expression regulator), calcium channel, and

vacuolar ATPases have also been reported to contribute to the infectivity of

SARS-CoV-2.[33,38,39] Pharmacological inhibition of these crucial host factors provides

another potential therapeutic approach. Although entry SARS-CoV-2 inhibitor

development is very attractive, no candidates have progressed into clinical trials.

To address this need, we screened 188 compounds to identify SARS-CoV-2

(S-G614) entry inhibitors in 293T-ACE2 cells. Nineteen compounds, including 11

bis-benzylisoquinoline alkaloids (SC9, SC161, SC180-188) and an analog of VE607

(SC171) were identified. VE607, previously reported as a SARS entry inhibitor with

an IC_{50} of 3 μM,[40] did not block SARS-CoV-2 entry at 20 μM in our assay. Among the

19 hits, nine compounds (SC9, SC161, SC171, SC182–187) with relatively high

activity, low cytotoxicity, and good specificity were selected for subsequent analyses.

The efficacy of the nine compounds in different SARS-CoV-2 entry stages was

examined to clarify whether they interfere with S-ACE2 interaction by targeting S or

host factors. The compounds showed effective entry blockade only in the pretreatment

stage, and did not bind to the S RBD, indicating that they target host factors.

SARS-CoV-2 enters TMPRSS2-negative 293T-ACE2 cells via endocytosis, and

TMPRSS2 inhibitors do not block SARS-CoV-2 infection in this cell line.[17] Besides

in 293T cells, our hits also inhibited SARS-CoV-2 infection in the TMPRSS2-positive

lung epithelial cell lines Calu-3 and A549, indicating that they do not target

TMPRSS2.

ACE2 is the common host receptor of SARS-CoV-2 and SARS-CoV, whereas

MERS-CoV infection relies on DPP4.[41,42] Interestingly, our findings demonstrated

that bis-benzylisoquinoline alkaloids (SC9/cepharanthine, SC161/hernandezine,

SC182/tetrandrine, and SC185/neferine) and the VE607 analog SC171 could potently

block infection by SARS-CoV-2, SARS-CoV, and MERS-CoV pseudoviruses.

Therefore, our compounds do not target ACE2 or DPP4 and thus may target some

other host factors. Recent studies have demonstrated a crucial role of Ca^{2+} in viral

fusion of Ebola, SARS-CoV, and MERS-CoV. Ca^{2+} coordinates with negatively

charged residues, such as aspartic acid and glutamic acid, within the viral fusion

peptide. Depletion of extracellular or especially, intracellular Ca^{2+} significantly

reduced infection by these viruses.[24,25,43-45] We noted that the identified

bis-benzylisoquinoline alkaloids had been reported as CCBs[26-29]. CCBs, originally

used to treat cardiovascular diseases, are supposed to have a high potential to treat

SARS-CoV-2 infections[46]. In this study, the bis-benzylisoquinoline CCBs abolished

S–ACE2-mediated membrane fusion in 293T-ACE2 cells. Calcium-free medium or

intracellular Ca^{2+} chelation significantly diminished SARS-CoV-2 pseudovirus

infection, suggesting that Ca^{2+} is required for SARS-CoV-2 entry. Upon pretreatment

with BAPTA-AM, the bis-benzylisoquinoline CCBs had >10-fold higher EC_{50} values

than those without BAPTA-AM pretreatment. These results corroborated that the

efficacy of these inhibitors depends on cellular calcium homeostasis. Additionally,

upregulation of cholesterol biosynthesis has been revealed as a common mechanism

underlying viral resistance. Perturbation of the cholesterol biosynthesis pathway with

the CCB amlodipine reduced viral infection[33]. Consistent herewith, the

bis-benzylisoquinoline CCBs upregulated intracellular cholesterol, which likely

contributes to infection inhibition.

In conclusion, we reported a set of bis-benzylisoquinoline alkaloids as

coronavirus entry inhibitors. These inhibitors effectively protected different cell lines

(293T, Calu-3, and A549) from infection by different coronaviruses (SARS-CoV,

MERS-CoV, SARS-CoV-2 [S-D614 and S-G614]) *in vitro*. The compounds block

host calcium channels, thus inhibiting Ca^{2+}-mediated fusion and suppressing virus

entry. Considering the effectiveness of CCBs in the control of hypertension, our study

provided clues to support that CCBs may be useful in treating coronavirus infection in

patients with hypertension.

Acknowledgments

We would like to thank Prof. Cheguo Cai (Wuhan University, Wuhan, China) for providing the pNL4-3.Luc.R-E- plasmid. This work was supported by the Key Laboratory of Infectious Diseases, CQMU, 202001, the Science and Technology Research Program of Chongqing Municipal Education Commission (KJQN202000418 to L-Y.H.), Emergency Project from the Science & Technology Commission of Chongqing (cstc2020jscx-fyzx0053 to A-L.H.), the Emergency Project for Novel Coronavirus Pneumonia from the Chongqing Medical University (CQMUNCP0302 to K.W.), the Leading Talent Program of CQ CSTC (CSTCCXLJRC201719 to N.T.), and a Major National Science & Technology Program grant (2017ZX10202203 to A-L.H.) from the Science & Technology Commission of China.

Author contributions

A.L.H., N.T., and Y.H.X. and designed and directed the study. C.L.H., L.Y.H., K.W., J.H., and G.J.Z. constructed the pesudoviruses and screened the compounds. C.J.G. and W.X. performed authentic SARS-CoV-2 assays. All authors reviewed the manuscript and consented to the description of author contribution.

Competing interests: The authors declare no competing interests.

Materials and Methods

Plasmids

The codon-optimized gene encoding SARS-CoV-2 spike (S) protein (GenBank:

QHD43416) with 19 amino acids deletion at the C-terminal was synthesized by Sino

Biological Inc (Beijing, China), and cloned it into the pCMV3 vector between the

restriction enzyme *Kpn*I and *Xba*I sites (denoted as pS-D614). The recombinant

plasmid pS-D614 was used as template, and the D614G mutant S-expressing plasmid

(denoted as pS-G614) was constructed by site-directed mutagenesis. SARS-CoV

S-expressing plasmid (Cat: VG40150-ACGLN, named as pS-SARS) and MERS-CoV

S-expressing plasmid (Cat: VG40069-CF, named as pS-MERS) were obtained from

Sino Biological Inc (Beijing, China). The VSV-G-expressing plasmid pMD2.G was

donated by Prof. Ding Xue from Tsinghua University (Beijing, China). The HIV-1

NL4-3 ΔEnv Vpr luciferase reporter vector (pNL4-3.Luc.R-E-) constructed by N.

Landau was donated by Prof. Cheguo Cai from Wuhan University (Wuhan, China).

Human ACE2 and DPP4 expression plasmids were derived from GeneCopoeia

(Guangzhou, China).

Cell lines and cell culture

HEK 293T, A549, and Calu3 cells were purchased from the American Type Culture

Collection (ATCC, Manassas, VA, USA). Cells were cultured at 37 ºC and 5% CO_2

atmosphere in Dulbecco's modified Eagle medium (DMEM; Hyclone, Waltham, MA,

USA) containing 10% fetal bovine serum (FBS; Gibco, Rockville, MD, USA), 100

mg/mL of streptomycin, and 100 units/mL of penicillin. HEK 293T cells transfected

with human ACE2 and DPP4 (293T-ACE2, 293T-DPP4) were cultured under the

same conditions with the addition of G418 (0.5 mg/mL) to the medium.

Antigens and antibodies

The RBD domain of SARS-CoV-2 S protein (His-tag) were synthesized by Prof.

Xuefei Cai at Key Laboratory of Molecular Biology for Infectious Diseases (Ministry

of Education), Chongqing Medical University. The anti-RBD monoclonal antibody

was kindly provided by Prof. Aishun Jin from Chongqing Medical University.

RBD-binding   peptide   SBP1   derived   from   human   ACE2   α   helix   1

(IEEQAKTFLDKFNHEAEDLFYQS) was synthesized by GenScript (Nanjing,

China).

Compounds and reagents

Custom compound library containing 188 small molecules, remdesivir, and aloxistatin

(E-64d) were purchased from MedChemExpress (HY-L027) and Chemdiv.

BAPTA-AM (T6245) was purchased from TargetMol. All the compounds were

dissolved in dimethyl sulfoxide (DMSO) at a stock concentration of 20 mM.

Cell cytotoxicity assay

The CellTiter 96® AQ$_{ueous}$ One Solution Cell Proliferation Assay (G3582, Promega,

USA) was used to assess cell viability according to the product's description. Briefly,

HEK 293T cells were dispensed into 96-well plate (2×10^4 cells/well), cultured in

medium containing gradient concentrations of the compound for 72 hours at 37 ℃ in

a humidified 5% CO_2 incubator. The cells were incubated with 100 µl fresh medium

after removal of the medium. Then 20 µl of CellTiter 96® AQueous One Solution

Reagent was added into each sample well and incubating the plate was incubated at

37°C for 1-4 hours in a humidified 5% CO_2 atmosphere. The absorbance at 490 nm

was measured using a microplate reader (Synergy H1, BioTek, USA).

Pseudoviruse production and quantification

Pseudotyped viruses were produced as previously described[47]. Briefly, 5×10^6 HEK

293T cells were co-transfected with 6 µg each of pNL4-3.Luc.R-E- and recombinant

plasmid (pS-SARS, pS-MERS, pS-D614, or pS-G614) using Lipofectamine 3000

Transfection Reagent (Invitrogen, Rockville, MD) according to the manufacturer's

instructions. After 48 h transfection, pseudotyped viruses expressing S-SARS,

S-MERS, S-D614, and S-G614 spike protein were harvested, centrifuged and filtered

through 0.45 µm filters, and subsequently stored at -80°C. 293T cells were

co-transfected with pNL4-3.Luc.R-E- and pMD2.G plasmid to collect the VSV-G

pseudovirus.

The copies of the pseudovirus were expressed as numbers of viral RNA genomes

per mL of viral stock solution and determined using RT-qPCR with primers and a

probe that targeting LTR. Sense primer: 5′-TGTGTGCCCGTCTGTTGTGT-3′,

anti-sense   primer: 5′-GAGTCCTGCGTCGAGAGAGC-3′,   probe:

5′-FAM-CAGTGGCGCCCGAACAGGGA-BHQ1-3′. Briefly, viral RNAs were

extracted according to the manufacturer's instructions with TRIzol reagent

(Invitrogen). Then, the TaqMan One-Step RT-PCR Master Mix Reagents (Applied

Biosystems, Thermo Fisher) was used to amplify total RNAs. pNL4-3.Luc.R-E-

vector with certain copies was used to generate standard curves. All the pseudotyped

viruses were titrated to the uniform titer (copies/mL) for the following research.

Compound screening

For pseudovirus-based inhibition assay, 188 compounds were screened via luciferase

activity. HEK 293T cells cultured in 96-well plate (2×10^4 cells/well) were incubated

with each compound (20 μM) for 1 hour and were infected with the same amount of

pseudovirus (3.8×10^4 copies in 50 µL). The cells were replaced to fresh DMEM

medium 8 h post-infection. Cells were lysed by 30 µl lysis buffer (Promega, Madison,

WI, USA) at 72 h post-infection to measure RLU with luciferase assay reagent

(Promega, Madison, WI, USA) according to the product description. All data were

performed at least three times and expressed as means ± standard deviations (SDs).

Cell-cell fusion assay

HEK 293T effector cells were transfected with plasmid pAdTrack-TO4-GFP encoding

green fluorescent protein (GFP) or pS-G614 encoding the corresponding

SARS-CoV-2 S protein. 293T-ACE2 cells were used as target cells. At 8 h

post-transfection, the effector cells were washed twice with PBS and were pretreated

with compounds or DMSO as control for another 16h. Subsequently, the effector cells

were overlaid on target cells at a ratio of approximately one S-expressing cell to two

receptor-expressing cells with about 90% confluent. After a 4-hour coculture, images

of syncytia were captured with an inverted fluorescence microscope (Nikon eclipse Ti,

Melville, NY).

For quantification of cell-cell fusion, three fields were randomly selected in each

well to count the fused and unfused cells. The fused cells were at least twice as large

as the unfused cells, and the fluorescence intensity was weaker in fused cells since

GFP diffusion from one effector cell to target cells. The percentage of cell-cell fusion

was calculated as: [(number of the fused cells/number of the total cells) $\times$ 100%].

Competitive ELISA

The recombinant RBD proteins derived from SARS-CoV-2 were coated on 96-well

microtiter plate (50ng/well) at 4°C overnight. After blocked with blocking buffer (5%

FBS and 2% BSA in PBS) for 1 hour at 37°C, serial dilution solutions of compounds,

ACE2 peptide SBP1 or DMSO were added into the plates and incubated at 37°C for 1

hour. Plates were washed five times with phosphate-buffered saline, 0.05% Tween-20

(PBST) to remove the free drug or DMSO. The wells were incubated with mouse

anti-RBD monoclonal antibody (1:1000 dilution) for 1 hour at 37°C, and then washed

with PBST five times and incubated with Horseradish peroxidase (HRP)-conjugated

goat anti-mouse antibody (Abmart, Shanghai, China) for 1 hour at 37°C. TMB

substrate was added and incubated for 15 minutes at 37°C for color development,

finally the absorbance at 450 nm was measured by a microplate reader.

Differential scanning fluorimetry

Briefly, purified His-RBD was diluted to 200 μg/mL in PBS buffer containing Sypro

Orange 5× (ThermoFisher) in a 96-well white PCR plate. Compounds and ACE2

derived peptide SBP1 were added at a final concentration of 20 μM. All samples were

tested in triplicate with a Bio-Rad RT-PCR system. The samples were first

equilibrated at 25 °C for 3 min, then heated from 25 °C to 85 °C with a step of 1 °C

per 1 min, and the fluorescence signals were continuously collected using CFX

Maestro. T_m values were calculated as previously described[48].

Functional analysis of Ca^{2+} in pseudovirus infectivity

293T-ACE2 cells were seeded in 96-well plate ($2x10^4$ cells/well) and incubated at

37 °C for 8h. For extracellular calcium depletion assays, cells were washed three

times using PBS with or without Ca^{2+}. Calcium-free medium with or without 2 mM

calcium chloride and 5 μM compounds were added and incubated for 1h at 37°C.

Next, S-G614 pseudovirus was added to the cells for 8 h at 37°C. For intracellular

calcium depletion assays, cells were washed three times using PBS with or without

Ca^{2+}, DMEM with BAPTA-AM (20 μM) or DMSO and gradient concentrations of

compounds were added and incubated for 2 h at 37°C. In the following, S-G614

pseudovirus was added to infect the cells for 8 h at 37°C. For both types of assays,

complete medium was then added after removed the medium. The cells were

measured by luciferase activity.

Cellular cholesterol measurement

Cellular cholesterol was measured using the Cholesterol/Cholesterol Ester-Glo™

Assay (J3190, Promega, USA) according to the product's description. In short, HEK

293T-ACE2 cells were assigned to 96-well plates (4×10^4 cells/well) and cultured for

24 hours at 37 °C with 5 μM compounds. DMSO was used as negative control. The

medium in the 96-well plate was removed and cells was washed twice with 200 μl

PBS. Then, 50 μl of cholesterol lysis solution was added, shaking the plate carefully

and incubating for 30 minutes at 37°C. Following, 50 μl of cholesterol detection

reagent was added with esterase or without esterase to all wells and the plate was

incubated at room temperature for 1 hour. Finally, recording luminescence with a

plate-reading luminometer. Total cholesterol concentration was calculated by

comparing the luminescence of samples and controls under the same condition.

Cytopathic effect (CPE) assay and quantification of SARS-CoV-2 infection

Vero E6 cells were dispensed into 96-well plate (4.0×10^4 cells/well), pre-treated with

medium containing compounds or DMSO for 1 hour at 37 ℃. Then 60 μl DMEM

medium which supplemented with compounds or DMSO was replaced immediately,

the same amount of SARS-CoV-2 (100 $TCID_{50}$/well) was added and incubated for 1

hour at 37 ℃. The mixture was removed and the cells were washed twice with PBS,

and cultured with 100 μl fresh medium for 48 hours at 37 ℃ with 5% CO_2

atmosphere. Cytopathic effects (CPE) induced by the virus was observed using

microscope at 48 h post-inoculation. Cell culture supernatant was collected at 48 h

post-infection for viral RNA quantification using the novel coronavirus real-time

RT-PCR Kit (Shanghai ZJ Bio-Tech Co, Ltd, Shanghai, China). Remdesivir (5 μM)

was used as positive control in the experiment.

Statistical analyses

Data were analyzed using GraphPad Prism version 6.0 software and were presented as

means ± SD. Statistical significance was determined using ANOVA for multiple

comparisons. Student's t test was applied to compare the two groups. Differences with

P values < 0.05 were deemed statistically significant.

Data availability

Data supporting the results of this study can be obtained from the author upon request

References

1   Coronaviridae Study Group of the International Committee on Taxonomy of,

V. The species Severe acute respiratory syndrome-related coronavirus:

classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* **5**, 536-544

(2020).

2   Liu, Y. *et al.* Aerodynamic analysis of SARS-CoV-2 in two Wuhan hospitals.

Nature **582**, 557-560 (2020).

3   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of

probable bat origin. *Nature* **579**, 270-273 (2020).

4   Organization, W. H. *Coronavirus disease (COVID-19) situation reports*,

<https://www.who.int/emergencies/diseases/novel-coronavirus-2019> (2020).

5   Wang, K. *et al.* Longitudinal dynamics of the neutralizing antibody response

to SARS-CoV-2 infection. *Clin Infect Dis* (2020).

6   Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion

conformation. *Science* **367**, 1260-1263 (2020).

7   Yan, R. *et al.* Structural basis for the recognition of SARS-CoV-2 by

full-length human ACE2. *Science* **367**, 1444-1448 (2020).

8   Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain

bound to the ACE2 receptor. *Nature* **581**, 215-220 (2020).

9   Shang, J. *et al.* Structural basis of receptor recognition by SARS-CoV-2.

Nature **581**, 221-224 (2020).

10   Cao, Y. *et al.* Potent neutralizing antibodies against SARS-CoV-2 identified by

high-throughput single-cell sequencing of convalescent patients' B cells. *Cell*

(2020).

11 Du, S. *et al.* Structurally Resolved SARS-CoV-2 Antibody Shows High

Efficacy in Severely Infected Hamsters and Provides a Potent Cocktail Pairing

Strategy. *Cell* (2020).

12 Wan, J. *et al.* Human-IgG-Neutralizing Monoclonal Antibodies Block the

SARS-CoV-2 Infection. *Cell Rep* **32**, 107918 (2020).

13 Zhu, Y., Yu, D., Yan, H., Chong, H. & He, Y. Design of potent membrane

fusion inhibitors against SARS-CoV-2, an emerging coronavirus with high

fusogenic activity. *J Virol* (2020).

14 Xia, S. *et al.* Fusion mechanism of 2019-nCoV and fusion inhibitors targeting

HR1 domain in spike protein. *Cell Mol Immunol* (2020).

15 Xia, S. *et al.* Inhibition of SARS-CoV-2 (previously 2019-nCoV) infection by

a highly potent pan-coronavirus fusion inhibitor targeting its spike protein that

harbors a high capacity to mediate membrane fusion. *Cell Res* **30**, 343-355

(2020).

16 Hoffmann, M. *et al.* Nafamostat Mesylate Blocks Activation of SARS-CoV-2:

New Treatment Option for COVID-19. *Antimicrob Agents Chemother* **64**

(2020).

17 Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and

TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**,

271-280 e278 (2020).

18   Riva, L. *et al.* Discovery of SARS-CoV-2 antiviral drugs through large-scale

compound repurposing. *Nature* (2020).

19   Kang, Y. L. *et al.* Inhibition of PIKfyve kinase prevents infection by Zaire

ebolavirus and SARS-CoV-2. *Proc Natl Acad Sci U S A* (2020).

20   Korber, B. *et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that

D614G Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827 e819

(2020).

21   Yurkovetskiy, L. *et al.* Structural and Functional Analysis of the D614G

SARS-CoV-2 Spike Protein Variant. *Cell* (2020).

22   Jiang, X. *et al.* Bimodular effects of D614G mutation on the spike

glycoprotein of SARS-CoV-2 enhance protein processing, membrane fusion,

and viral infectivity. *Signal Transduction and Targeted Therapy* **5** (2020).

23   Maisonnasse, P. *et al.* Hydroxychloroquine use against SARS-CoV-2 infection

in non-human primates. *Nature* **585**, 584-587 (2020).

24   Lai, A. L., Millet, J. K., Daniel, S., Freed, J. H. & Whittaker, G. R. The

SARS-CoV Fusion Peptide Forms an Extended Bipartite Fusion Platform that

Perturbs Membrane Order in a Calcium-Dependent Manner. *J Mol Biol* **429**,

3875-3892 (2017).

25   Straus, M. R. *et al.* Ca(2+) Ions Promote Fusion of Middle East Respiratory

Syndrome Coronavirus with Host Cells and Increase Infectivity. *J Virol* **94**

(2020).

26   Low, A. M. *et al.* Plant alkaloids, tetrandrine and hernandezine, inhibit

calcium-depletion stimulated calcium entry in human and bovine endothelial

cells. *Life Sci* **58**, 2327-2335 (1996).

27   Leung, Y. M., Berdik, M., Kwan, C. Y. & Loh, T. T. Effects of tetrandrine and

closely related bis-benzylisoquinoline derivatives on cytosolic Ca2+ in human

leukaemic HL-60 cells: a structure-activity relationship study. *Clin Exp*

Pharmacol Physiol **23**, 653-659 (1996).

28   John P. Felix *et al.* Bis(benzylisoquinoline) Analogs of Tetrandrine Block

L-Type Calcium Channels: Evidence for Interaction at the Diltiazem-Binding

Site. *Biochemistry* (1992).

29   Anselmi, E. *et al.* Selective inhibition of calcium entry induced by

benzylisoquinolines in rat smooth muscle. *J Pharm Pharmacol* **44**, 337-343

(1992).

30   Liu, Q. Y., Karpinski, E., Rao, M. R. & Pang, P. K. Tetrandrine: a novel

calcium channel antagonist inhibits type I calcium channels in neuroblastoma

cells. *Neuropharmacology* **30**, 1325-1331 (1991).

31   Ranganathan, S., Harmony, J. A. & Jackson, R. J. Effect of Ca2+ blocking

agents on the metabolism of low density lipoproteins in human skin fibroblasts.

Biochem Biophys Res Commun **107**, 217-224 (1982).

32   Mori, S., Ito, H. & Yamamoto, K. Effects of calcium antagonists on low

density lipoprotein metabolism in human arterial smooth muscle cells. *Tohoku*

J Exp Med **154**, 329-333 (1988).

33   Daniloski, Z. *et al.* Identification of required host factors for SARS-CoV-2

infection in human cells. *Cell* (2020).

34   Tharappel, A. M., Samrat, S. K., Li, Z. & Li, H. Targeting Crucial Host

Factors of SARS-CoV-2. *ACS Infect Dis* **6**, 2844-2865 (2020).

35   Xiu, S. *et al.* Inhibitors of SARS-CoV-2 Entry: Current and Future

Opportunities. *J Med Chem* (2020).

36   Yang, L. *et al.* Identification of SARS-CoV-2 entry inhibitors among already

approved drugs. *Acta Pharmacol Sin* (2020).

37   Yang, C. *et al.* Salvianolic acid C potently inhibits SARS-CoV-2 infection by

blocking the formation of six-helix bundle core of spike protein. *Signal*

Transduct Target Ther **5**, 220 (2020).

38   Wei, J. *et al.* Genome-wide CRISPR screens reveal host factors critical for

SARS-CoV-2 infection. *Cell* (2020).

39   Danta, C. C. Calcium Channel Blockers: A Possible Potential Therapeutic

Strategy for the Treatment of Alzheimer's Dementia Patients with

SARS-CoV-2 Infection. *ACS Chem Neurosci* **11**, 2145-2148 (2020).

40   Kao, R. Y. *et al.* Identification of novel small-molecule inhibitors of severe

acute respiratory syndrome-associated coronavirus by chemical genetics.

Chem Biol **11**, 1293-1299 (2004).

41   Wang, N. *et al.* Structure of MERS-CoV spike receptor-binding domain

complexed with human receptor DPP4. *Cell Res* **23**, 986-993 (2013).

42   Lu, G. *et al.* Molecular basis of binding between novel human coronavirus

MERS-CoV and its receptor CD26. *Nature* **500**, 227-231 (2013).

43 Nathan, L. *et al.* Calcium Ions Directly Interact with the Ebola Virus Fusion

Peptide To Promote Structure-Function Changes That Enhance Infection. *ACS*

Infect Dis **6**, 250-260 (2020).

44 Das, D. K. *et al.* Conformational changes in the Ebola virus membrane fusion

machine induced by pH, Ca2+, and receptor binding. *PLoS Biol* **18**, e3000626

(2020).

45 Gunaratne, G. S., Yang, Y., Li, F., Walseth, T. F. & Marchant, J. S.

NAADP-dependent Ca2+ signaling regulates Middle East respiratory

syndrome-coronavirus pseudovirus translocation through the endolysosomal

system. *Cell Calcium* **75**, 30-41 (2018).

46 Straus, M. R., Bidon, M., Tang, T., Whittaker, G. R. & Daniel, S. FDA

approved calcium   channel blockers   inhibit SARS-CoV-2 infectivity in

epithelial lung cells. *Biorxiv* (2020).

47 Hu, J. *et al.* Development of cell-based pseudovirus entry assay to identify

potential viral entry inhibitors and neutralizing antibodies against

SARS-CoV-2. *Genes Dis* (2020).

48 Niesen, F. H., Berglund, H. & Vedadi, M. The use of differential scanning

fluorimetry to detect ligand interactions that promote protein stability. *Nat*

Protoc **2**, 2212-2221 (2007).

Fig. 1 Screening of compounds active against SARS-CoV-2 S-G614 pseudovirus

in 293T-ACE2 cells. a Schematic diagram of the screening method and workflow.

The screening method is shown on the left, and the selection criteria for hits are

outlined on the right side. **b** Scatter plot of primary screening of 188 compounds

against S-G614 infection. Inhibition ratios for all drugs obtained in a preliminary

screening are represented by scattered points. Red dots indicate the 41 compounds

with an inhibition rate ≥70%. DMSO and E-64d were used as a negative and positive

control, respectively.

Fig. 2 Effect of treatment timing on the efficacy of nine selected compounds on

SARS-CoV-2 S-G614 pseudovirus entry. a Treatment timing diagrams for the nine

selected compounds. HEK 293T cells were treated with the nine compounds or

DMSO before, during, or after S-G614 pseudovirus entry. Eight treatment conditions

(A–G) were tested for each compound. **b–j** Inhibitory effects of **(b)** SC9, **(c)** SC161,

(d) SC171, **(e)** SC182, **(f)** SC183, **(g)** SC184, **(h)** SC185, **(i)** SC186, and **(j)** SC187 on

the entry of S-G614 pseudovirus at different treatment time points. All experiments

were repeated at least three times.

Fig. 3 Efficacy of nine compounds in Calu3, 293T-ACE2, and A549 cell lines.

Inhibitory effects of **(a)** SC9, **(b)** SC161, **(c)** SC171, **(d)** SC182, **(e)** SC183, **(f)** SC184,

(g) SC185, **(h)** SC186, and **(i)** SC187 against S-G614 infection in the three cell lines.

All experiments were repeated at least three times.

Fig. 4 Dose-response curves for five broad-spectrum inhibitors of five types of coronaviruses. a–e Inhibition curves of five selected compounds **(a)** SC9, **(b)** SC161, **(c)** SC171, **(d)** SC182, **(e)** SC185 on S-D614, S-G614, S-SARS, S-MERS, and VSV-G pseudoviruses. **f** EC_{50} values of five selected compounds against entry of different pseudoviruses. All experiments were repeated at least three times.

Fig. 5 Interactions between four selected compounds and the SARS-CoV-2 RBD

domain. a Inhibitory effect of the ACE2 peptide SBP1 at 20 μM on the invasion of

S-G614 pseudovirus. **b** Binding of pre-coated His-RBD with compounds at 20 μM as

determined by competitive ELISA. SBP1 and DMSO were used as a positive and

negative control, respectively. **c** Melting curves for His-RBD (10 μM) incubated with

20 μM of each compound tested by differential scanning fluorimetry. **d** T_M values of

His-RBD incubated with SC161, SC185, or SBP1. All experiments were repeated at

least three times.

Fig. 6 Investigation of the mechanism of the selected compounds. a Inhibitory

effect of SC9, SC161, SC171, SC182, and SC185 at 5 μM on SARS-CoV-2 S

mediated cell-cell fusion. **b** Cell–cell fusion rate in the presence of 5 μM SC9, SC161,

SC171, SC182, or SC185. The fusion rate in a DMSO-treated group was set as 100%.

c Effect of extracellular Ca^{2+} depletion on S-G614 pseudovirus entry. **d** Effect of 5

μM of each compound on S-G614 pseudovirus entry into 293T-ACE2 cells in medium

supplemented with 2 mM $CaCl_2$. **e** Effect of 5 μM of each compound on S-G614

pseudovirus entry in calcium-depleted culture medium. **f** Effect of intracellular Ca^{2+}

depletion using 20 μM BAPTA-AM on S-G614 pseudovirus entry. DMSO was used

as a control. **g** Cell viability assay of 293T-ACE2 cells treated with increasing

concentrations of BAPTA-AM (0–50 μM). **h** Inhibition curves of the compounds

against S-G614 pseudovirus entry in the presence of 20 μM BAPTA-AM. **i** EC_{50}

values of the compounds for S-G614 pseudovirus entry in the presence or absence of

20 μM BAPTA-AM. **j** Cholesterol (normalized to total protein) in 293T-ACE2 cells

treated with the compounds or DMSO for 24 h. All experiments were repeated at least

three times.

Fig. 7 Antiviral activities of four selected compounds against authentic

SARS-CoV-2 in Vero E6 cells. a We assessed the inhibitory effect of the compounds

on native SARS-CoV-2 infection by observing their cytopathogenic effects. SC9,

SC161, SC171, and SC185 were tested at 0.1 μM, 1 μM, 10 μM, and DMSO and

remdesivir (5 μM) were used as a negative and positive control, respectively. **b** The

relative RNA level in the DMSO treatment group was set to 100%, and that in the

positive control remdesivir (5 μM) treatment group was 0.01%. **c** The relative RNA

levels in the SC9, SC161, SC171, and SC185 (10 μM) treatment groups were 70.27%,

43.55%, 76.98% and 0.08%, respectively. At 1 and 0.1 μM, relative RNA levels did

not significantly decrease. All experiments were repeated at least three times.