Title

The dorsal hippocampus is required for the formation of long-term duration memories

in rats

Running head

Dorsal hippocampus and duration memory

Authors

Toshimichi Hata[1], Tatsuya Yamashita[1], and Taisuke Kamada[2, 3]

Affiliations

[1]Faculty of Psychology, Doshisha University, 1-3 Tatara Miyakodani, Kyotanabe,

Japan 610-0394

[2]Graduate School of Psychology, Doshisha University, 1-3 Tatara Miyakodani,

Kyotanabe, Japan 610-0394

[3]Present affiliation: *Medical Innovation Center, Graduate School of Medicine, Kyoto*

University, Sakyo-ku, Kyoto, Japan 606-8507

Corresponding author

Toshimichi Hata E-mail: thata@mail.doshisha.ac.jp

Acknowledgments

We would like to thank Editage (www.editage.com) for English language editing.

Manuscript information

Pages: 36

Tables: 1

Figures: 4

Abstract: 244 words

ABSTRACT

Interval timing—the perception of durations mainly in seconds or minutes—is a

ubiquitous behavior in organisms. Animal studies have suggested that the hippocampus

plays an essential role in duration memory; however, the memory processes involved are

unclear. To clarify the role of the dorsal hippocampus in the acquisition of long-term

duration memories, we adapted the "time-shift paradigm" to a peak-interval procedure.

After a sufficient number of training sessions with an initial target duration (20 s), the rats

underwent "shift sessions" with a new target duration (40 s) under a muscimol (0.5 μg

per side) infusion into the bilateral dorsal hippocampus. The memory of the new target

duration was then tested in drug-free "probe sessions," including the empty trials in which

any lever presses were not reinforced. In the probe sessions, the mean peak function of

the muscimol group was located leftward to the control group, but these two functions

were superimposed on the standardized time axis, suggesting a scalar property. In the

session-by-session analysis, the mean peak time (an index of timing accuracy) of the

muscimol group was lower than that of the control group in the probe sessions, but not in

the shift sessions. In the trial-by-trial analysis, the mean start time and stop times (indices

of timing accuracy) were lower than in the control group in the probe sessions. These

findings suggest that the dorsal hippocampus is required for the formation of long-term

duration memories within the range of interval timing.

Keywords: *Interval timing, Time perception, Peak interval, Dorsal hippocampus,*

Long-term memory, Short-term memory, Time-shift paradigm

1. Introduction

Interval timing—the perception, estimation, and discrimination of durations in seconds, minutes, or hours—is a fundamental ability in many species to perform functions such as foraging and decision making (Buhusi & Meck, 2005). For instance, the nectar-feeding birds Amakihi (*Loxops virens*) periodically revisit the flowers from which they had previously collected nectar for efficient foraging (Kamil, 1978). Pigeons and rats underestimate the value of delayed reinforcement as a function of the duration of the delay (Mazur, 1987; Richards, Mitchell, & de Wit, 1997; Green, Myerson, Holt, Slevin, & Estle, 2004). For humans, instant noodles would be soggy if we could not estimate the interval it takes for them to be ready. Thus, interval timing is a ubiquitous behavior for organisms.

Animal studies have shown that hippocampal dysfunction alters interval timing behavior (for review, Lee, Thavabalasingam, Alushaj, Cavdaroglu, & Ito, 2019). The "classic effects" (Yin & Troger, 2011) of hippocampal or hippocampus-related regions dysfunctions on interval timing are to alter the accuracy of the estimated durations—a leftward shift of the response rate function during the peak-interval (PI) procedure (Hata & Okaichi, 1998; Meck, 1988; Meck, Church, & Olton, 1984; Olton Meck, & Church, 1987; Tam, Jennings, & Bonardi, 2015; Yin & Meck, 2014) and an interresponse time (IRT) distribution in the differential reinforcement of low rate (DRL) schedules (Jaldow & Oakley, 1990). These findings suggest that hippocampal lesions do not impair the interval timing itself (i.e., the regulation of behavior depending on the passage of time), while lesions alter the length of the estimated durations.

Other lines of evidence from trace conditioning, electrophysiology, and computational models suggest that the hippocampus and hippocampal neurons are involved in the

memory or coding of durations (see Table 1 in Lee et al., 2019). In trace conditioning, an

unconditioned stimulus (US) is delivered after the termination of a conditioned stimulus

(CS). The CS offset – US onset interval (CS–US interval) is usually set in the millisecond-

to-second range. Many studies using a second CS–US interval range (i.e., range of the

interval timing) have reported that hippocampal lesions, especially on the dorsal parts,

impair the acquisition and retention of trace fear conditioning (McEchron, Bouwmeester,

Tseng, Weiss, & Disterhoft, 1998; McEchron, Tseng, & Disterhoft, 2000; Quinn,

Oommen, Morrison, & Fanselow, 2002; Quinn, Loya, Ma, & Fanselow, 2005; Lin &

Honey, 2011; Raybuck & Lattal, 2011; Sellami, Abed, Brayda-Bruno, Etchamendy,

Valério, Oulé, … Marighetto, 2017). Lee et al. (2019) stated that trace conditioning

studies largely substantiate the hippocampus in representations of temporal duration

memory. The hippocampal "time cell," which fires at a specific moment in a temporally

structured experience, has received much attention in the last decade (Pastalkova, Itskov,

Amarasingham, & Buzsaki, 2008; MacDonald, Carrow, Place, & Eichenbaum, 2013;

Saltz, Tiganj, Khasnabish, Kohley, Sheehan, Howard, & Eichenbaum, 2016; for review,

Eichenbaum, 2014). Moreover, recent computational studies suggest that the temporal

distance between the to-be-learned time criterion and the time of the peak activity of each

time cell provides an error signal to correct the synaptic weight (Oprisan, Aft, Buhusi, &

Buhusi, 2018), and the memory of the duration is stored in the hippocampus (Oprisan,

Buhusi, & Buhusi, 2018). Taken together, these recent findings suggest that the

hippocampus plays an important role in duration memory.

However, these studies are insufficient to determine whether the hippocampus plays a

vital role in the acquisition of duration memories. First, studies on trace conditioning did

not selectively focus on duration memory. Thus, the impairment in the acquisition of trace

conditioning following hippocampal dysfunction can be explained not only by the

inability to acquire the memory of the duration but also by other impairments, such as the

disappearance of the memory trace of the CS presentation, or impairment of the

continuation of the timing proposed in the ICAT model of temporal processing (Petter,

Lusk, Hesslow, & Meck, 2016). Second, due to the nature of electrophysiological

correlational studies, it cannot be concluded in principle that the activity of the time cell

is the source of the duration memory. Lesion or inactivation studies should be performed

to answer this question. Third, models, including computational models, must be

supported by experimental observations. Taken together, an experimental design that

improves these deficiencies should be designed to determine whether the hippocampus

plays an important role in the acquisition of duration memory.

To meet these requirements, we applied the "time-shift paradigm" (Hata, 2011) to the

Peak-interval (PI) procedure. In the time-shift paradigm, after sufficient training with the

initial target duration (e.g., 20 s), the target duration was changed to 40 s in a "shift

session." In previous studies, a drug was injected into the striatum (MacDonald, Cheng,

& Meck, 2012) or plasticity-related gene expression levels in the striatum were evaluated

(Dallérac, Graupner, Knippenberg, Martinez, Tavares, Tallot, … Doyère, 2017) in the

shift session to clarify the relationship between the acquisition of duration memory and

neuronal processes. We believe that this paradigm can dissociate the effects of a drug

with regard to the acquisition of the duration memory from its impact on other processes

(e.g., task rules, which may be included in the early stage of learning during the first target

duration).

An important modification of the time-shift paradigm in our current experiment is to

adopt the probe sessions provided from the day after the shift sessions to evaluate the

effects of the drug infused during the shift sessions. As mentioned above, some indirect

evidence suggests the involvement of the dorsal hippocampus in duration memory.

However, the processes or stages of temporal memory in which the hippocampus is

involved remains unclear. An early study that applied the time-shift paradigm to the PI

procedure did not report impairments in the acquisition of the second duration memory

in hippocampal lesioned rats (Meck, 1988). However, this finding does not necessarily

mean that the hippocampus does not play a crucial role in acquiring duration memory.

This result may be explained by the notion that the hippocampus is important for the

acquisition of long-term, but not short-term, duration memories. In general, the PI

procedure is comprised of two types of trials— the "food" and "empty" trials. The food

trials have a discrete fixed interval (FI) schedule to allow learning of the target durations.

Empty trials are used to test the extent of duration memory. These trials are presented in

random order. Only data from the empty trials are usually analyzed. In empty trials, the

subjects can access not only the long-term duration memories that are acquired through

the previous sessions, but also the short-term duration memories that are retrieved from

the recent food trials within the session. In such situations, even if the hippocampal lesion

impairs the formation of long-term duration memory, the animals can behave adaptively

in the shift sessions, depending on the intact short-term duration memory. In fact, in a

previous study, an intra-dorsal hippocampal infusion of muscimol, an agonist of GABA

A receptors, impaired the intermediate-term (5 min) memory, but not the short-term (10

s) memory in a delayed non-matching to sample (DNMTS) task in the eight-arm radial

maze (Lee & Kesner, 2003). If probe sessions that are comprised of only empty trials are

prepared after the shift sessions, then the within-session short-term memory is no longer

unavailable in the sessions. All the available memories are limited to long-term memories

acquired through the previous sessions. In such situations, the impairment of the

acquisition of the long-term duration memory assumed by the hippocampal dysfunction

in the shift sessions should come to the surface in the probe sessions. To detect this

possibility, probe sessions provided from the day following the shift sessions were used

to evaluate the effect of the drug.

Therefore, the purpose of the present study was to clarify the role of the dorsal

hippocampus in the acquisition of long-term duration memory by applying the time-shift

paradigm to the PI procedure with probe sessions. After learning the first target duration

(20 s), the rats underwent PI training for a new target duration (40 s) during shift sessions

under the pharmacological inhibition of the bilateral dorsal hippocampus. The memory

of the new target duration was then tested using drug-free probe sessions, including empty

trials only. We hypothesized that an intra-hippocampal infusion of muscimol during the

shift sessions would impair the acquisition of long-memory of the second target duration.

We expected that the effects of the drug would be detected in the probe sessions, but not

in the shift sessions.

2. Methods

2.1 Subjects

Sixteen experimentally naïve, 10-week-old male Wistar albino rats (Shimizu

Laboratory Supplies, Kyoto, Japan) were used, but one animal died during surgery. Each

rat was individually housed in a stainless steel cage. The rats were depriving food for the

first day of shaping; hence, their body weight decreased to 85% of their free-feeding

weight at the start of the behavioral procedure. To compensate for the natural growth, the

body weight of the rats was increased at a rate of 5 g per week. Water was available *ad*

libitum. The temperature of the breeding room was maintained at 22 °C ± 2 °C. The light-

dark cycle was 12:12 h, and the light phase started at 8:00 A.M. All experimental sessions

were executed during the light phase. All experimental procedures were approved by the

Doshisha Committee of Animal Experiments (A17075).

2.2 Apparatus

Eight identical operant chambers were individually installed in a sound-attenuating box

with a 25 W LED light on the ceiling. Each chamber ($220 \times 165 \times 200$ mm in DWH) was

equipped with a non-retractable lever on the left side (40 mm from the floor) and a food

cup (15 mm from the floor) in the center of the front wall. The floor was made with

stainless steel bars with a 3 mm diameter, each bar 10 mm apart. A self-made application

developed with XOJO® (XOJO Inc. Austin, TX, USA) was run on two Macintosh

PowerBook Air computers (Apple, Cupertino, CA, USA) to control the experiment and

collect data. This was interfaced with two programmable controllers (SYSMAC CPM1A-

40CDR-A-V1, OMRON, Kyoto, Japan) and two USB I/O controllers to control the eight

chambers (RBIO-2U, Kyoritsu Electronic Industry, Osaka, Japan).

2.3 Procedure

Habituation. After handling for 5 min per day for 5 days, each animal was trained on

a 20 s fixed time (FT) schedule for five sessions, one session per day. In each session, 30

food pellets (F0021-J, Bio-serv, Flemington, NJ, USA) were delivered separately every

20 s, irrespective of animal behavior. The house light was continuously turned on during

the session.

Shaping. Rats were trained to press the lever under a continuous reinforcement

schedule until they earned 60 pellets from their lever presses or after 60 min had passed

in each session, whichever came first. The house light was turned on during the session.

The acquisition criteria were that they earned 60 pellets within 60 min in three successive

days.

PI 20 s training (sessions 1-30). The rats were then trained on the PI procedure with

one session per day. A session consisted of two types of trials—food and empty trials—

in a random order separated by intertrial intervals (ITIs) of 40 ± 10 s. The ratio of

food:empty trials was 42:18 in sessions 1-21 and 15:15 in sessions 22-30. Both types of

trials started with the onset of the house light and tone (signal stimuli) and finished with

their offsets. In a food trial, the first lever press after 20 s from the start of the trial was

reinforced with a food pellet, and then the trial finished (discrete trial FI 20 s). An empty

trial lasted for 60 s without any reinforcements for any lever presses. If a lever press was

emitted during the last five seconds of an ITI, the ITI was extended for 10 s from the

press.

Surgery. Rats were anesthetized with 2-3% isoflurane (2-3 L/min flow rate, MK-A110,

Muromachi, Tokyo, Japan) and fixed on a stereotaxic frame (Kopf Instruments, Tujunga,

CA, USA). A guide cannula (C232G-3.8/SPC, Plastics One, Roanoke, VA, USA)

attached to a dummy cannula (C232DC-3.8/SPC, Plastics One) was lowered into each

hemisphere of the dorsal hippocampus. The target coordinates of the tip of the guide

cannula based on a brain map (Paxinos & Watson, 1998) were as follows: -3.6 mm

anteroposterior to the bregma, ± 1.9 mm mediolateral, and -2.2 mm dorsoventral from the

skull surface. The cannula assembly was fixed to the skull surface using dental cement

and two small screws. The dummy cannula was inserted into the guide cannula and fixed

using a dust cap (303DC, Plastics One). Rats were allowed to recover for ≥5 d after the

surgery. For a few days during the early recovery period, an antibiotic agent (Mycillinsol,

Meiji, Tokyo, Japan) was applied to the wound once per day to avoid infections.

PI 20 s retraining (sessions 31-37). After the recovery period, rats were retrained using

the PI procedure (seven sessions) similar to the training phase. The ratio of the food:empty

trials was 15:15. The last two retraining sessions were defined as baseline sessions (Figure

2A).

Shift session (sessions 38-39). Rats were assigned to one of two groups—the artificial

cerebrospinal fluid (aCSF) or muscimol (Mus) group—to ensure that the performances

during the last three sessions of the retraining phase were as even as possible. Immediately

before starting each session, the dummy cannula was replaced with an internal cannula

(C232I-3.8/SPC, Plastics One), extending 1 mm from the tip of the guide cannula for

infusion. The muscimol solution for Mus rats (0.5 µg/µL, 1 µL per side) or aCSF for

aCSF rats (1 µL) was infused into the bilateral dorsal hippocampus at a flow rate of 0.5

µL/min for 2 min using a gastight syringe (100 µL, Hamilton, Reno, NV, USA) and

microsyringe pump (ESP-32, Eicom, Kyoto, Japan). The cannula was left in place for 1

min to allow dispersion of the solution, which was then replaced with a dummy cannula.

The dummy cannula was again inserted into the guide cannula and fixed with a dust cap.

The significant changes in the procedure from the previous sessions were that lever press

behavior was reinforced with a 40 s FI during food trials, and that the duration of the

empty trials was extended to 120 s. The other parameters were the same as those in the

retraining sessions.

Probe session (sessions 40-41). Rats were tested in two sessions of 15 empty trials

without infusions. The other parameters were the same as those in the shift sessions.

2.4 Dependent variables

Session-by-session analysis For each rat's session, the total number of lever-press

responses in each 3-s bin (such as 0-3 s and 1-4 s) was counted over all empty trials and

converted to the percentage of the maximum number of responses among the bins

(response rate). The response rates were plotted as a function of elapsed time (peak

function). The peak function was fitted with a Gaussian curve using KaleidaGraph® (ver.

4.5.3, Synergy Software, Reading, PA, USA): $R(t) = a + b \times \exp\{-.5 \times [(t - c)/d]^2\}$. Here,

t is the time from the start of the empty trial, and $R(t)$ is the response rate in bin t.

Parameter c is the time at which the response rate was at its maximum. Thus, c was

adopted as the peak time (an index of the subjective length of the to-be-timed duration).

The initial values of a, b, and d for the iterative estimation were 1, 100, and 10,

respectively. To determine the initial c, the first and last bins, which had response rates

of over 80%, were detected. The mean of the two class values of these bins was set at the

initial c. The data used for fitting ranged from 1.5 to 38.5 s for the baseline sessions, and

1.5 to 78.5 s for the shift and probe sessions, which was performed to cut off the noise

observed in the last part of the empty trials. The allowable error of the interactive

estimation was set to 0.1%. Curve fitting was also performed for the mean peak functions.

For the mean functions, the initial a, b, and d values were the same as for the individual

functions, whereas *c* was set at 20 and 40 for the baseline and the shift/probe sessions,

respectively. The data used to fit the mean functions ranged from 1.5 to 58.5 s for the

baseline sessions and 1.5 to 118.5 s for the shift and probe sessions. For the standardized

mean functions of the probe sessions (see Figure 2D), the values of the x (time) axis were

converted to the ratio of the estimated peak time in each group. The original time values

corresponding to the standardized x of 0.1 steps were substituted into the regression curve

formula. The Y values were then converted to the maximum Y value.

The discrimination index (*DI*) was also calculated for the individual peak function

using $DI = 100/\text{mean response rate of all bins in the peak function}$ (Meck & Church,

1987) as an index of precision. In addition, for the DI calculation, the range of averaging

in the denominator was from 1.5 s to 38.5 s for the baseline sessions and from 1.5 s. to

78.5 s for the shift and probe sessions, respectively.

Trial-by-trial analysis In an individual empty trial, rats showed a "low-high-low"

response pattern. That is, they started pressing the lever before the required time, kept

responding, and then stopped responding after the required time. According to an

algorithm proposed by a previous study (Church, Meck, & Gibbon, 1994), the timepoints

of the transition from low to high (start time) and high to low (stop time) can be detected.

The midpoint is the mean of the start and stop times. The start time and stop times are the

indices of timing accuracy, in parallel with the peak time of the session-by-session

analysis. The spread was calculated by stop time minus start time, which was the index

for the timing precision.

2.5 Histology

After completing all behavioral sessions, the rats were deeply anesthetized with sodium

pentobarbital (130 mg/kg, i.p.) and infused with dye (2% pontamine sky blue) using a

procedure similar to that of drug infusion to locate the tips of the internal canulae. They

were then perfused intracardially with saline, followed by ALTFiX® (FALMA, Tokyo,

Japan). Their brains were removed and soaked in 10% and then 20% sucrose phosphate

buffer (0.1 M) until they sunk in each solution. The brains were sectioned in the coronal

plane (40-μm thickness) using a cryostat (CM1850, Leica, Wetzlar, Germany). Sections

were stained with cresyl violet to assess the location of the cannula tip.

2.6 Statistical analysis

Two-way mixed ANOVAs followed by the test of simple main effects and Holm's

multiple   comparisons   were   performed   using   "anovakun"   (ver.   4.8.5.,

http://riseki.php.xdomain.jp/index.php) in R software (ver. 4.0.0, R Core Team, Vienna,

Austria).

3. Results

3.1 Histological results

A photomicrograph of a representative section is presented in Figure 1A. The tips of

the guide cannulae (arrows) were located above the dorsal hippocampus, while those of

the internal cannulae were extended 1 mm from the guide cannulae. Hence, all tips of the

internal canulae were presumed to be located within the dorsal hippocampus (Figure 1B).

Apparent injuries beyond the canula tracks could not be observed.

3.2 Session-by-session analysis

The data from three rats were excluded from the session-by-session analysis because

the peak time or R^2 of the fitting curve of their probe or baseline sessions were over ±2

SD from the mean of each group. As a result, the number of subjects used in the session-

by-session analysis was six for each group.

Response rate function. The mean peak function for the two sessions of the last

retraining sessions (baseline), shift sessions, and probe sessions were calculated for each

rat. The group mean is presented in Figure 2. Qualitative observation showed that the

functions in the baseline (Figure 2B) could be seen to overlap with each other. The peaks

of both groups were located at approximately 19 s. In the shift sessions (Figure 2C),

although the Mus group had a wider distribution than the aCSF group, the peaks of both

groups were located around 33 s. Interestingly, the functions from the probe sessions

(Figure 2D) showed a dramatic difference between the groups. The function from Mus

appeared to the left of the aCSF. The peak of the Mus group was around 26 s, whereas

that of the aCSF group was around 35 s. These functions, however, were superimposed

as a whole by standardizing the horizontal axis (Figure 2E).

Peak time. Mean peak times for the baseline, shift sessions, and probe sessions were

calculated for each rat. The group mean (± *SEM*) is presented in Figure 3A. There was no

difference in the mean peak time of the baseline and shift sessions, whereas it was lower

in the Mus group than in the aCSF group for the probe sessions. Mixed two-way ANOVA

revealed that the interaction of the group × block (baseline, shift, and probe) (F(1.68,

16.82) = 4.96, dfs were adjusted by Huynh-Feldt-Lecoutre's epsilon, $p = .025$, $\eta_G^2 = .267$).

The simple main effect of the group type was significant for the probe sessions (F(1, 10)

= 6.85, $p = .026$, $\eta_G^2 = .407$), but not for the baseline and shift sessions (Fs(1, 10) = .05

and .69, $ps = .834$ and .426, $\eta_G^2 s = .005$ and .065, respectively). The simple main effects

of the blocks were significant in both groups (aCSF: F(1.03, 5.16) = 15.55, $p = .010$, η_G^2

= .703. Mus: F(1.88, 9.41) = 24.06, $p < .001$, $\eta_G^2 = .767$). The post-hoc Holm's multiple

comparisons test showed that the peak times of the shift and probe sessions were

significantly higher than the baseline in each group. Moreover, in the Mus group, the

mean peak time of the probe sessions was significantly lower than that of the shift

sessions.

DI. The mean DI for the baseline, shift sessions, and probe sessions were calculated for

each rat; the mean *DI* values (± *SEM*) are presented in Figure 3B. There was no difference

in *DI* between the baseline and probe sessions, whereas it was lower in the Mus group

than in the aCSF group in the shift sessions. A mixed two-way ANOVA revealed a

significant interaction (F(2, 20) = 8.02, $p = .003$, $\eta_G^2 = .27$). The simple main effect of the

group type was significant in the shift sessions, but not in the probe sessions (Fs(1, 10) =

7.48 and 4.19, $ps = .021$ and .068, $\eta_G^2 s = .428$ and .295, respectively).

R². The mean R^2 (± *SEM*) of the fitting curves is shown in Table 1. All values were

over .820. Mixed two-way ANOVA did not detect any significant effects (interaction and

main effect blocks: Fs(1.05, 10.46) = 2.00, 4.42, $ps = .186$ and .059, $\eta_G^2 s = .095$ and .187,

main effect of group type: F(1, 10) = .819, $p = .389$, $\eta_G^2 = .038$).

3.3 Trial-by-trial analysis

Data from two rats were excluded from the trial-by-trial analysis because the values of

one of three indices (start time, midpoint, stop time, or spread) in the baseline or probe

sessions were over ±2 SD from the mean of each group. The midpoint was used only for

the excluding the outlier. As a result, the numbers of subjects in the trial-by-trial analysis

were 7 and 6 for aCSF and Mus, respectively.

Start time and stop time The medians of the start time and the stop time were calculated

for all trials (15 trials × 2 sessions, 30 trials) in the baseline, shift, and probe sessions for

each rat. The group mean of the medians (± *SEM*) is shown in Figure 4. For the start time

(Figure 4A), the means were different between the groups and the phases. However, a

two-way mixed ANOVA (group × phase) revealed that the interactions ($F(1.60, 17.58) =$

3.215, $p = .074$, $\eta_G^2 = .137$) and the main effect of the group ($F(1, 11) = .692$, $p = .423$,

$\eta_G^2 = .028$) were not significant. The main effect of the phase was significant ($F(1.60,$

$17.58) = 3.215$, $p = .074$, $\eta_G^2 = .433$). For the stop time (Figure 4B), the means of the Mus

group was higher than that of the aCSF group in the shift sessions, whereas the

relationship was reversed in the probe sessions. A two-way ANOVA revealed that the

interaction was significant ($F(1.45, 15.94) = 9.067$, $p = .004$, $\eta_G^2 = .355$). The simple main

effect of the group was significant in the shift session ($F(1, 11) = 12.292$, $p = .005$, η_G^2

$= .528$), but not in the probe session ($F(1, 11) = 4.501$, $p = .057$, $\eta_G^2 = .290$). The simple

main effect of the phase was significant in both groups (aCSF: $F(1.19, 7.15) = 27.811$, p

$< .001$, $\eta_G^2 = .769$, Mus: $F(1.29, 6.44) = 125.949$, $p < .001$, $\eta_G^2 = .929$). The post-hoc

Holm's multiple comparisons test showed that the stop time of the probe sessions was

significantly lower than the shift sessions in the Mus group but not in the aCSF group.

Spread. The group means of the medians (± *SEM*) of the spread are shown in Figure

4C. It is noticeable that the spread of the Mus group was shorter than that of the aCSF

group in the probe sessions. The interaction and the following test of the simple main

effect of the group in the probe sessions were significant ($F(1.65, 18.20) = 6.65$, $p = .009$,

$\eta_G^2 = .295$, and $F(1, 11) = 5.71$, $p = .036$, $\eta_G^2 = .342$, respectively). The simple main effect

of the phase was significant in both groups (aCSF: $F(1.61, 9.64) = 22.448$, $p < .001$, η_G^2

$= .719$, Mus: $F(1.30, 6.48) = 50.694$, $p < .001$, $\eta_G^2 = .880$). The post-hoc Holm's multiple

comparisons test showed that the spread of the probe sessions was significantly lower

than the shift sessions in the Mus group but not in the aCSF group.

4. Discussion

Two fundamental prerequisites; the homogeneity of the two groups before the

important experimental operations and the validity of the task, were suggested to be

guaranteed by the data. First, the peak functions of both groups almost entirely overlapped

with each other in the baseline sessions (Figure 2B). Moreover, the mean peak time, *DI*

(Figure 3), and R^2 (Table 1) in the session-by-session analysis, and the mean start time,

stop time, and spread in the trial-by-trial analysis (Figure 4) in the baseline sessions were

all similar between the groups. These findings suggest that the differences between

groups in the following analysis do not stem from the bias of the assignment of the

subjects to the groups. Second, the task's validity was confirmed by the transition of the

mean peak time (Figure 3A). In the aCSF group, the peak time of the shift session was

significantly higher than that of the baseline, suggesting that the memory of the new target

time was acquired by the animals. In addition, the extended peak time was maintained in

the probe sessions, as evidenced by the non-significant difference between these two

phases. These findings suggest that the memory of the new target duration was maintained

for at least 24 h.

In the shift session, dysfunction of the dorsal hippocampus was suggested to impair the

precision, but not the accuracy of the interval timing. The significant extension of the stop

time (Figure 4B) inevitably induced elevated response rates after the peak time in the

peak function (Figure 2C). This elevation could account for the significantly lowered *DI*

value seen in the Mus group (Figure 3B), suggesting impaired precision (discussed further

below). On the other hand, the dysfunction of the hippocampus does not seem to impair

the accuracy of the interval timing at the shift sessions, as evidenced by the lack of a

difference in peak time in the shift sessions (Figure 3A). This finding, however, does not

contradict the "classic effect" (Yin & Troger, 2011, see the Introduction section) of the

hippocampal lesion on the interval timing. Many studies have repeatedly reported that a

dysfunction of the hippocampus results in shortening of the peak time (Hata & Okaichi,

1998; Meck, 1988; Meck et al., 1984; Olton et al., 1987; Tam et al., 2015; Yin & Meck,

2014). In these studies, the effect has been reported in sessions in which the required time

of the PI procedure was constant and appeared after the chronic dysfunction of the

hippocampus. However, in our study, the required time was changed, and the

hippocampal function was reversibly inhibited only in two sessions. In this way, the

situation in our study was different from those suggestive of "classic" hippocampal

dysfunction effects. Therefore, our findings do not contradict those of previous studies.

The noticeable findings observed in the probe sessions can be explained by the notion

that the dorsal hippocampus is involved in the formation of long-term, but not short-term,

duration memories. The most important asset of the probe sessions was that the sessions

included only empty trials. Thus, the within-session short-term memory was not available

for feedback control of the behavior in empty trials. All of the memories that were

available to guide their behavior were limited to those acquired during the

training/retraining sessions and/or the shift sessions. It is known that when the required

time is increased during peak interval procedures, the peak times gradually increase in

normal rats (Lejeune, Ferrarra, Simons, & Wearden, 1997; Meck, Komeily-Zadeh, &

Church, 1984). These findings suggest that the shorter the peak time, the weaker the

memory of the new required time. Here, our data were as follows: the mean peak function

of the Mus group was located to the left of the aCSF group's function (Figure 2D). The

mean peak time of the Mus group was lower than that of the aCSF group in the probe

sessions (Figure 3A). The significantly extended peak time from the baseline sessions to

the shift sessions was maintained in the probe sessions in the aCSF group but not in the

Mus group (Figure 3A) . In the trial-by-trial analysis, the mean stop time of the Mus group

in the probe sessions was sufficiently (p=.057) lower than the aCSF group (Figure 4B).

Moreover, the significantly extended stop time from the baseline sessions to the shift

sessions was maintained in the probe sessions in the aCSF group but not in the Mus group

(Figure 4B). These results consistently suggest that the animals in the Mus group had a

poorer memory of the new required time (40 s) than the aCSF animals in the probe

sessions. Other important results were that the mean peak times were similar between

groups in the shift sessions (Figure 3A). This can be interpreted to show that the within-

session short-term memory from the food trials was available even though the muscimol

was infused, which guided the animals' adaptive behavior. Some previous studies support

this "short-term/long-term dissociation hypothesis" (Haettig, Stefanko, Multani,

Figueroa, McQuown, Wood, 2011; Lee & Kesner, 2003; Nagahara & McGaugh, 1992).

In a study on the DNMTS task in the eight-arm radial maze, an intra-dorsal hippocampal

infusion of muscimol impaired their intermediate-term (5 min) memory, but not their

short-term (10 s) memory (Lee & Kesner, 2003). Intra-dorsal hippocampal muscimol

infusion immediately after memory acquisition resulted in impairments in the

performance of an object memory test conducted in mice 24 h after the acquisition

(Haettig et al., 2011). Moreover, a muscimol infusion before the acquisition of inhibitory

avoidance into the septum, which has strong reciprocal connections with the

hippocampus, disrupted retrieval after 48 h, but not 15 s. After 15 min of acquisition of

the memory, the performance was halfway between 48 h and 15 s conditions (Nagahara

& McGaugh, 1992). Although the task used and/or the target region in our study were

different from those of the reference studies, it is not surprising that the within-session

short-term memory was available in our experiment. Taken together, our evidence

strongly suggests that dysfunction of the dorsal hippocampus impairs the formation of

long-term memory of a new duration.

The short-term/long-term dissociation hypothesis can also explain the discrepancy

between our findings, the findings of a previous study, and the lower *DI* of the Mus group

in the shift sessions. First, as mentioned in the Introduction, Meck (1988) reported that

the hippocampectomized rats did not show a delay in the acquisition of memories of the

new target duration. In this experiment, however, the effect of the hippocampal lesion

was examined in the shift sessions, each of which comprised both food and empty trials.

If this hypothesis is correct, it is quite reasonable that the previous study did not report

impairments in the acquisition of the new target memory. Therefore, this "discrepancy"

may be ostensible. Second, the lower *DI* of the Mus group in the shift sessions might be

explained by the incomplete short-term memory and the resulting dysfunction in feedback

control. A previous study suggested that normal animals modulate when to start

responding and when to stop in PI procedures, which were regulated according to the

information from their own performances in recent food trials (Meck, 1988). The study

also reported that this feedback control was impaired by the hippocampal lesion. Even if

the within-session short-term memory seemed to be available in the Mus group, the

availability might be lower than that of the aCSF group because, according to previous

studies (Lee & Kesner, 2003; Nagahara & McGaugh, 1992), the inferred time window in

which short-term memory was maintained may vary from several dozen seconds to

several minutes. However, in our PI procedure, the animals must maintain a memory for

at a least 160 s (40 s for the mean duration of ITI + 120 s for duration of an empty trial)

for feedback control. Indeed, the contents of the memories are different from each other

(durations, spatial information, and fear emotions), so the estimation may be too

simplified. However, it seems reasonable to infer that a weakness of the feedback control

resulting from the available but weak short-term memory produced a fluctuation of the

"low-high-low" pattern and then lowered the *DI* in the shift sessions. This speculation

should be confirmed by further studies.

The shortening of the spread and the superimposition of the peak functions can be

compatible without contradiction. In the literature on interval timing, it has been

repeatedly confirmed that two peak functions produced under different required times are

superimposed on the standardized time axis relative to the peak time, that is, the scalar

property (Church et al., 1994; Gibbon, 1977; Gibbon & Church, 1990, 1992; Malapani &

Fairhurst, 2002). This means that the ratio of the width of the function to the peak time is

a constant (i.e., Weber's law). In our data, the two peak functions were superimposed on

each other in the probe session (Figure 2E), suggesting that the Mus rats had a normal

interval timing based on their own target times, which were different from those of the

aCSF rats. Therefore, the narrower mean spread (Figure 4C) and peak function (Figure

2D) in the Mus group, which had a smaller peak time, were inevitable.

The most noticeable progress of our study was its extension of previous knowledge

from trace conditioning, electrophysiology, and computational models. First, data from

trace conditioning were believed to be evidence of the role of the hippocampus in

temporal duration memory (Lee et al., 2019). However, other explanations, such as the

disappearance of the memory trace of the CS presentation or an impairment of the

"continuation" proposed by the ICAT model of timing (Petter et al., 2016) can also

explain the data. Using a timing-specific task (peak interval task) and a filled duration

(i.e., the duration in which the to-be-timed stimulus is continuously presented) and

confirming the normal interval timing in the probe sessions in the Mus group, we could

exclude alternative interpretations of the data from the trace conditioning. Second,

inactivation by the muscimol demonstrated the causal, but not correlational, relationship

between the hippocampus and the acquisition of long-term duration memory. Third, our

data provided experimental support to the computational models arguing the memory role

of the hippocampus in interval timing (Oprisan, Aft, Buhusi, & Buhusi, 2018; Oprisan,

Buhusi, & Buhusi, 2018).

In summary, the present findings strongly suggest that the dorsal hippocampus plays

an important role in the formation of long-term, but not short-term, duration memory in

the supra-second range.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

T.Y. conceptualized the study. T.H. and T.K. supervised the study. T.Y., T.K., and T.H.

performed the experiments. T.H. and T.K. reanalyzed the data. T.H. and T.K. wrote the

manuscript.

DATA AVAILABILITY STATEMENT

The datasets are available from the open science framework (OSF).

https://osf.io/fqngb/?view_only=6d21bea2cec74fa784a9cd79f3fe3774

ORCID

T. Hata https://orcid.org/0000-0003-4614-6550

T. Kamada https://orcid.org/0000-0001-8102-2807

REFERENCES

Buhusi, C. V., & Meck, W. H. (2005). What makes us tick? Functional and neural

mechanisms of interval timing. *Nature Reviews Neuroscience*, *6*, 755–65. doi:

10.1038/nrn176

Church, R., Meck, W., & Gibbon, J. (1994). Application of scalar timing theory to

individual trials. *Journal of Experimental Psychology: Animal Behavior Processes*,

20, 135–155. doi: 10.1037/0097-7403.20.2.135

Dallérac, G., Graupner, M., Knippenberg, J., Martinez, R., Tavares, T., Tallot, L., …

Doyère, V. (2017). Updating temporal expectancy of an aversive event engages

striatal plasticity under amygdala control. *Nature Communications*, *8*, 13920. doi:

10.1038/ncomms13920

Eichenbaum, H. (2014). Time cells in the hippocampus: A new dimension for mapping

memories. *Nature Reviews Neuroscience*, *15*, 732–744. doi: 10.1038/nrn3827

Gibbon, J. (1977). Scalar expectancy theory and Weber's law in animal timing.

Psychological Review, *84*, 279–325. doi: 10.1037/0033-295X.84.3.279

Gibbon, J., & Church, R. M. (1990). Representation of time. *Cognition*, *37*, 23–54. doi:

10.1016/0010-0277(90)90017-e

Gibbon, J., & Church, R. M. (1992). Comparison of variance and covariance and

covariance patterns in parallel and serial theories of timing. *Journal of the*

Experimental Analysis of Behavior, *57*, 393–406. doi: 10.1901/jeab.1992.57-393

Green, L., Myerson, J., Holt, D. D., Slevin, J. R., & Estle, S. J. (2004). Discounting of

delayed food rewards in pigeons and rats: Is there a magnitude effect? *Journal of the*

Experimental Analysis of Behavior, 81, 39-50. doi: 10.1901/jeab.2004.81-39

Hata, T. (2011). Glutamate - a forgotten target for interval timing. *Frontiers in Integrative*

Neuroscience, 5, 27. doi:10.3389/fnint.2011.00027

Hata, T, & Okaichi, H. (1998). Effects of fimbria-fornix lesion on the temporal

discrimination revealed by peak interval procedure in rats. *Shinrigaku Kenkyu*, *69*,

304–9. doi: 10.4992/jjpsy.69.304

Jaldow, E. J., & Oakley, D. A. (1990). Performance on a differential reinforcement of

low-rate schedule in neodecorticated rats and rats with hippocampal lesions.

Psychobiology, *18*, 394–403. doi: 10.3758/bf03333086

Kamil, A. (1978). Systematic foraging by a nectar-feeding bird, the amakihi (Loxops

virens). *Journal of Comparative and Physiological Psychology*, *92*, 388–396. doi:

10.1037/h0077479

Lee, I., & Kesner, R. P. (2003). Time-dependent relationship between the dorsal

hippocampus and the prefrontal cortex in spatial memory. *The Journal of*

Neuroscience, *23*, 1517–1523. doi: 10.1523/jneurosci.23-04-01517.2003

Lee, A., Thavabalasingam, S., Alushaj, D., Cavdaroglu, B., & Ito, R. (2019). The

hippocampus contributes to temporal duration memory in the context of event

sequences: A cross-species perspective. *Neuropsychologia*, *137*, 107300. doi:

10.1016/j.neuropsychologia.2019.107300

Lin, T.-C. E., & Honey, R. C. (2011). Encoding specific associative memory: Evidence

from behavioral and neural manipulations. *Journal of Experimental Psychology:*

Animal Behavior Processes, *37*, 317–329. doi: 10.1037/a0022497

MacDonald, C. J., Carrow, S., Place, R., & Eichenbaum, H. (2013). Distinct hippocampal

time cell sequences represent odor memories in immobilized rats. *The Journal of*

Neuroscience, *33*, 14607–14616. doi: 10.1523/JNEUROSCI.1537-13.2013

Macdonald, C. J., Cheng, R.-K., & Meck, W. H. (2012). Acquisition of "start" and "stop"

response thresholds in peak-interval timing is differentially sensitive to protein

synthesis inhibition in the dorsal and ventral striatum. *Frontiers in Integrative*

Neuroscience, *6*. doi: 10.3389/fnint.2012.00010

Malapani, C., & Fairhurst, S. (2002). Scalar timing in animals and humans. *Learning and*

Motivation, *33*, 156–176. doi: 10.1006/lmot.2001.1105.

Mazur, J. E. (1987). An adjusting procedure for studying delayed reinforcement. In M. L.

Commons, J. E. Mazur, J. A. Nevin, & H. Rachlin (Eds.), *Quantitative analyses of*

behavior, Vol. 5. The effect of delay and of intervening events on reinforcement value

(p. 55–73). Lawrence Erlbaum Associates, Inc.

McEchron, M. D., Bouwmeester, H., Tseng, W., Weiss, C., & Disterhoft, J. F. (1998).

Hippocampectomy disrupts auditory trace fear conditioning and contextual fear

conditioning in the rat. *Hippocampus*, *8*, 638–646. doi: 10.1002/(sici)1098-

1063(1998)8:6<638::aid-hipo6>3.0.co;2-q

McEchron, M. D., Tseng, W., & Disterhoft, J. F. (2000). Neurotoxic lesions of the dorsal

hippocampus disrupt auditory-cued trace heart rate (fear) conditioning in rabbits.

Hippocampus, *10*, 739–751. doi: 10.1002/1098-1063(2000)10:6<739::aid-

hipo1011>3.0.co;2-i

Meck, W. (1988). Hippocampal function is required for feedback control of an internal

clock's criterion. *Behavioral Neuroscience*, *102*, 54–60. doi: 10.1037/0735-

7044.102.1.54

Meck, W.H., Church, R. M., & Olton, D. S. (1984). Hippocampus, time, and memory.

Behavioral Neuroscience, *98*, 3-22. doi: 10.1037//0735-7044.98.1.3.

Nagahara, A. H., & McGaugh, J. L. (1992). Muscimol infused into the medial septal area

impairs long-term memory but not short-term memory in inhibitory avoidance, water

maze place learning and rewarded alternation tasks. *Brain Research*, *591*, 54–61. doi:

10.1016/0006-8993(92)90977-h

Olton, D. S., Meck, W. H., & Church, R. M. (1987). Separation of hippocampal and

amygdaloid involvement in temporal memory dysfunctions. *Brain Research*. *404*,

180-8. doi: 10.1016/0006-8993(87)91369-2

Oprisan, S. A., Aft, T., Buhusi, M., & Buhusi, C. V. (2018a). Scalar timing in memory:

A temporal map in the hippocampus. *Journal of Theoretical Biology*, *438*, 133–142.

doi: 10.1016/j.jtbi.2017.11.012

Oprisan, S. A., Buhusi, M., & Buhusi, C. V. (2018b). A population-based model of the

temporal memory in the hippocampus. *Frontiers in Neuroscience*, *12*, 521. doi:

10.3389/fnins.2018.00521

Pastalkova, E., Itskov, V., Amarasingham, A., & Buzsaki, G., (2008). Internally

generated cell assembly sequences in the rat Hippocampus. *Science*, *321*, 1322–1327.

doi:10.1126/science.1159775

Paxinos, G., & Watson, C. (1998). The rat brain in stereotaxic coordinates. 4th ed. [CD-

ROM]. Academic Press, San Diego.

Petter, E., Lusk, N., Hesslow, G., & Meck, W. (2016). Interactive roles of the cerebellum

and striatum in sub-second and supra-second timing: Support for an initiation,

continuation, adjustment, and termination (ICAT) model of temporal processing.

Neuroscience & Biobehavioral Reviews, *71*, 739–755. doi:

10.1016/j.neubiorev.2016.10.015

Quinn, J. J., Loya, F., Ma, Q. D., & Fanselow, M. S. (2005). Dorsal hippocampus NMDA

receptors differentially mediate trace and contextual fear conditioning. *Hippocampus,*

15, 665–674. doi: 10.1002/hipo.20088.

Quinn, J. J., Oommen, S. S., Morrison, G. E., & Fanselow, M. S. (2002). Post-training

excitotoxic lesions of the dorsal hippocampus attenuate forward trace, backward

trace, and delay fear conditioning in a temporally specific manner. *Hippocampus*, *12*,

495–504. doi: 10.1002/hipo.10029.

Raybuck, J. D., & Lattal, M. K. (2011). Double dissociation of amygdala and

hippocampal contributions to trace and delay fear conditioning. *PLoS One*, *6*, e15982.

doi: 10.1371/journal.pone.0015982

R Core Team (2020). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/

Richards, J. B., Mitchell, S. H., & de Wit, H. (1997). Determination of discount functions

in rats with an adjusting-amount procedure. *Journal of the Experimental Analysis of*

Behavior, 67, 353-366. doi: 10.1901/jeab.1997.67-353

Salz, D. M., Tiganj, Z., Khasnabish, S., Kohley, A., Sheehan, D., Howard, M. W., &

Eichenbaum, H. (2016). Time cells in hippocampal area CA3. *The Journal of*

Neuroscience, 36, 7476–7484. doi: 10.1523/jneurosci.0087-16.2016

Sellami, A., Abed, A. S. A., Brayda-Bruno, L., Etchamendy, N., Valério, S., Oulé, M.,

… Marighetto, A. (2017). Temporal binding function of dorsal CA1 is critical for

declarative memory formation. *Proceedings of the National Academy of Sciences*,

114, 10262–10267. doi: 10.1073/pnas.1619657114

Tam, S. K., Jennings, D. J., & Bonardi, C. (2015). Effects of dorsal hippocampal damage

on conditioning and conditioned-response timing: A pooled analysis. *Hippocampus*,

25, 444-459. doi: 10.1002/hipo.22381

Yin, B., & Meck, W. H. (2014). Comparison of interval timing behaviour in mice

following dorsal or ventral hippocampal lesions with mice having δ-opioid receptor

658 gene deletion. *Philosophical Transactions of the Royal Society B*, *369*, 20120466.

659 doi: 10.1098/rstb.2012.0466

660 Yin, B., & Troger, A. (2011). Exploring the 4th dimension: Hippocampus, time, and

661 memory revisited. *Frontiers in Integrative Neuroscience*, 5, 36. doi:

662 10.3389/fnint.2011.00036

Table 1. Mean (± *SEM) R²* of the fitting curves

	Baseline	Shift	Probe
aCSF	0.959 (.009)	0.924 (.027)	0.939 (.014)
Mus	0.961 (.011)	0.820 (.079)	0.947 (.013)

Figure captions

Figure 1. **(A)** A representative photomicrograph showing the tracks of the canulae. The tips of the guide canulae (arrows) were located above the dorsal hippocampus. The tips of the internal canulae were extended 1 mm ventrally from these. Scale bar = 1 mm. **(B)** Placements of the internal cannula tips for the aCSF (open circles) and Mus rats (closed circles) on the brain. Reprinted from Paxinos, G. & Watson, C. *The Rat Brain in Stereotaxic Coordinates. 4th ed. [CD-ROM]*, Copyright (1998).

Figure 2. **(A)** Overview of the main parts of the experiment. Each types of trials were (+) or were not (-) included in the sessions. Note that the required time was changed from 20 s to 40 s for the shift sessions and the drugs were infused before starting these sessions. In addition, only empty trials were included in the probe sessions. Mean peak function in the baseline **(B)**, shift **(C)**, and probe **(D)** sessions. The vertical dotted lines show the original (20 s) and new (40 s) target times. The arrows show the time at which the response rate was at the maximum for each group. **(E)** Standardized mean peak functions. The horizontal axis was converted to the ratio to the peak time of each peak function. The open and closed circles are the estimated standardized response rates for the aCSF and Mus group respectively. The two curves were overlapping as a whole.

Figure 3. Mean (± *SEM*) peak time **(A)**, and DI **(B)**. * $p < .05$, ** $p < .01$ between the groups. # $p < .05$, ## $p < .01$, ### $p < .001$ *vs.* the baseline of the group.

686   **Figure 4**. Mean median (± *SEM*) of the start time **(A)**, the stop time **(B)**, and the

687   spread **(C)**. $p < .05$, ** $p < .01$ between the groups. # $p < .05$, ## $p < .01$, ### $p < .001$

688   *vs.* the baseline of the group.

Fig.1

A

B

Fig.2

Fig.3

Fig.4