Dietary folic acid deficiency impacts hippocampal morphology and cortical acetylcholine metabolism in adult male and female mice

Calli Bennett BS[1,2], Jacalyn Green PhD[3], Mae Ciancio PhD[4], Joanna Goral PhD[5], Lenore Pitstick[3] BS, Matt Pytynia BA[4], Alice Meyer MBS[5], Neha Kwatra BSc[1,6], Nafisa M. Jadavji PhD[1,7,8]

Affiliations
[1] Biomedical Sciences Program, College of Graduate Studies, Midwestern University, Glendale, US
[2] College of Osteopathic Medicine, Midwestern University, Glendale, US
[3] Biochemistry and Molecular Genetics, College of Graduate Studies, Midwestern University, Downers Grove, US
[4] Biomedical Sciences Program, College of Graduate Studies, Midwestern University, Downers Grove, US
[5] Anatomy, College of Graduate Studies, Midwestern University, Downers Grove, US
[6] College of Dental Medicine, Midwestern University, Glendale, US
[7] College of Veterinary Medicine, Midwestern University, Glendale, US
[8] Department of Neuroscience, Carleton University, Ottawa, Canada

Corresponding Author
Nafisa M. Jadavji, PhD
Department of Biomedical Sciences
College of Graduate Studies
Midwestern University
19555 N 59th Ave
Glendale, AZ 85308, USA
Phone: 623-572-3665
Fax : 623-572-3647
E-mail: njadav@midwestern.edu

Abstract

Objective

One-carbon metabolism is a metabolic network that integrates nutritional signals with biosynthesis, redox homeostasis, and epigenetics. There are sex differences in hepatic one-carbon metabolism. However, it is unclear whether there are sex differences in dietary deficiencies of one-carbon metabolism in the brain. The aim of the study was to examine the impact of sex differences and dietary folic acid deficiency on brain tissue in adult mice.

Methods

Male and female C57Bl/6J mice were placed on a folic acid deficient (FD) or control diet (CD) at six weeks of age. Mice were maintained on these diets for six months, after which animals were euthanized and brain tissue and serum were collected for analysis. Serum folate levels were measured. In brain tissue, hippocampal volume and morphology including Cornu Ammonis 1 and 3 (CA1; CA3), and dentate gyrus thickness were measured. Apoptosis within the hippocampus was assessed using active caspase-3 immunofluorescence staining. Additionally, cortical acetylcholine metabolism was measured in brain tissue using immunofluorescence staining of acetylcholinesterase (AChE) and choline acetyltransferase (ChAT).

Results

Male and female FD mice had reduced serum levels of folate. Both males and females maintained on a FD showed a decrease in the thickness of the hippocampal CA3 region. Interestingly, there was a sex difference in the levels of apoptosis within the CA1 region of the hippocampus. In cortical tissue, there were increased levels of neuronal ChAT and reduced levels of AChE in FD females and male mice.

Conclusions

The results indicated that FD impacts hippocampal morphology through increased apoptosis and changes acetylcholine metabolism within the cortex. The data from our study indicate a sex difference in apoptosis and differences in hippocampal morphology and choline metabolism as a result of dietary folic acid deficiency.

Keywords

One- carbon metabolism; folic acid; hippocampus; choline; apoptosis; acetylcholine

1. Introduction

One-carbon (1C) metabolism is a key metabolic network that integrates nutrition signals with biosynthesis, redox homeostasis and epigenetics. It plays an essential role in the regulation of cell proliferation, stress resistance, and embryonic development (1). Folate, also known as vitamin B9, is the main component of 1C metabolism. Folates act as 1C donors in the regeneration of methionine from homocysteine, nucleotide synthesis and repair, and DNA methylation(1,2).

While gastrointestinal bacteria produce small amounts of folate, the majority of folates are obtained in our diets. However, the bioavailability and stability from these dietary sources are much lower than synthetic, folic acid (3). Many countries fortify foods with folic acid in order to reduce deficiencies (4). However, even in countries in which fortifying foods is routine, deficiencies remain fairly common. Deficiencies are seen in individuals that have a high rate of tissue turnover (e.g., pregnant women), have an alcohol use disorder (5), or in individuals using folate antagonist drugs (e.g., methotrexate) (6). Additionally, deficiencies are common in the elderly population, which is especially vulnerable to malnutrition and folate absorption impairments (7) .

Folate's role in the regeneration of methionine from homocysteine proves especially significant when considering vascular and brain health. Elevated levels of homocysteine, that can be a result of folic acid deficiency, have been linked to several negative health outcomes, including increased risk for Alzheimer's Disease (8), and cerebrovascular diseases including stroke (9). Additionally, hyperhomocysteinemia is an important indicator of post-stroke recovery (10), mortality (11), and risk of recurrence (12). Homocysteine leads to downstream oxidative damage (13), indicating a potential mechanism behind its role in neuronal degeneration and atherosclerosis (14).

Choline, another component of 1C metabolism, is an essential nutrient. Folate and choline metabolism are tightly linked. In model systems studying the impact of reduced levels of dietary folates, choline metabolism is also affected (15–18). Choline can act as a 1C donor, especially in situations of diminished folate. In the brain, choline is also involved in the production of acetylcholine, the main neurotransmitter of the parasympathetic nervous system, and lipid membrane synthesis. Acetylcholine is synthesized by choline acetyltransferase (ChAT) and hydrolyzed by acetylcholinesterase (AChE). Reduced levels of choline have been reported to result in more apoptosis in cultured cells (19) and were observed in the hippocampus of offspring of choline deficient females (20). *In vivo* rodent studies have also shown cognitive impairment, as a result of a dietary choline deficiency. The hippocampus is particularly vulnerable to both choline and folate dietary deficiencies (15,17,21,22).

There is growing evidence that there are significant sex differences with regard to 1C metabolism (23–25), including quantifiable differences in homocysteine levels have (26). Males homozygotic for the polymorphism in methylenetetrahydrofolate reductase are more vulnerable to factors (e.g., smoking) that increase levels of homocysteine (26). Estrogen may serve as a protective factor, limiting hyperhomocysteinemia that in turn leads to an increased risk for vascular diseases(26). Furthermore, five key hepatic 1C metabolism enzymes are differentially expressed in males and females(24). Estrogen and other female hormones impact expression of these enzymes, leading to increased levels of homocysteine in males, and increased levels of choline and betaine in females(24). Further, folate intake and expression of genes in 1C metabolism regulate steatosis in a sex specific manner (27). Despite the significant evidence of hepatic 1C differences between sexes, limited evidence exists on sex related 1C metabolism differences in the brain. A recent study, including both male and female mice, demonstrates that there are significant

sex differences in behavior observed caused by a methylenetetrahydrofolate reductase deficiency (23,25). The aim of this study was to investigate if sex modulates the effects of dietary folic acid deficiency in brain tissue using a mouse model system. We hypothesize that in addition to dietary folate levels, sex will have a significant effect on hippocampal morphology and cortical 1C metabolism.

2. Methods

2.1 Experimental design

All experiments were done in accordance with IACUC animal welfare protocols and were approved by the Midwestern University Downers Grove IACUC Committee. Male and female C57Bl/6J mice were put on a folic acid deficient (FD) or control (CD) diets (Envigo) at 6 weeks of age (Table 1). Animals had access to food and water *ad libitum*. Each diet group had 5 males and 5 females; a total of 20 animals were used.

The CD and FD contained the same concentrations of macro-micro-nutrients, except for folic acid (Table 1). The CD contained 2.0 mg/kg folic acid (TD.04194), while FD contained 0.2 mg/kg folic acid (TD.95247) (28,29) . Mice were maintained on these diets for 6 months and at 8 months of age mice were euthanized with CO_2 inhalation and cervical dislocation. This corresponds to mature adults in humans (30).

Table 1. List of micro- and macro-nutrient contents in Envigo control (TD.04194) and folic acid deficient diet (TD.95247). The Envigo folic acid deficient (TD.95247) was identical to the control diet (TD.04194) except it contained 0.2 mg/kg folic acid.

Category	Concentration	Category	Concentration
Amino acids	**(g/kg)**	**Lipids**	**(g/kg)**
Lysine	14	Total fat	60.0
Methionine	4.7	Saturated fat	9.0
Cystine	3.6	Monounsaturated fat	13.8
Arginine	6.6	Polyunsaturated fat	36.6
Phenylalanine	8.8	4:0 Butyric acid	0.0
Tyrosine	9.2	6:0 Caproic Acid	0.0
Histidine	5.1	8:0 Caprylic Acid	0.0
Isoleucine	10.1	10:0 Capric acid	0.0
Leucine	16	12:0 Lauric acid	0.0
Threonine	7.6	14:0 Myristic Acid	0.0
Tryptophan	2.1	16:0 Palmitic acid	6.6
Valine	11.7	16:1 Palmitoleic acid	0.0
Aspartic Acid	12.1	18:0 Stearic acid	2.4
Glutamic acid	36.6	18:1 Oleic acid	13.8
Alanine	5.2	18:2 Linoleic acid	31.8
Glycine	3.2	18:3 Linolenic acid	4.8
Proline	18.1	Cholesterol	8.8 mg/kg
Serine	10		
Fat-soluble Vitamins	**(IU/kg)**	**Minerals**	**(mg/kg)**
Vitamin A	4000	Calcium	5100
Vitamin D	1000	Phosphorus	3000
Vitamin E	75	Potassium	3600
Vitamin K	0.8	Sodium	1000
Water-soluble Vitamins	**(mg/kg)**	Chlorine	1600
Biotin	0.20	Magnesium	515
Choline	1147.5	Copper	6.0
Folic acid	2.0	Iron	36.5
Niacin	30.0	Zinc	35.6
Pantothenate	14.7	Manganese	10.5
Riboflavin	6.0	Iodine	0.21
Thiamin	4.9	Selenium	0.15
Vitamin B_6	5.8	Molybdenum	0.15
Vitamin B_{12}	0.03	Chromium	1.00
Vitamin C	0.0		

2.2 Tissue collection

At time of euthanization body and brain weights were recorded. Tissue including brain and blood were collected for analysis. Serum was isolated from blood and stored at -80°C until analysis. Brain tissues were fixed in 4% paraformaldehyde overnight and then switched over to a 70% ethanol solution. Tissue was processed and embedded in paraffin in a coronal orientation. Brain tissue was sectioned using a microtome (Leica) at 5-µm thickness. Sections were slide mounted and stored at room temperature for staining.

2.3 Microbiological assay for serum folate

Lactobacillus rhamnosus (ATCC) was grown overnight in Folic Acid Casei Medium (Difco) supplemented with 0.025% sodium ascorbate (A7631, Sigma-Aldrich,) and 1.2 ng/ml of (6S)-5-formyltetrahydrofolate (Schirks Laboratories). An equal volume of 80% sterile glycerol was added to the overnight culture and 1 ml aliquots were frozen and stored at -80°C. The day of the assay, an aliquot of *L. rhamnosus* was thawed and 100ul was added to 50ml of Folic Acid Casei Medium supplemented with 0.025% sodium ascorbate. The approximately 20 mg/liter folic acid stock (Sigma-Aldrich) was 0.22 µm filter sterilized, verified by absorbance spectra at 282 (molar absorptivity coefficient= 27,000 M^{-1}, MW= 441.4), and stored at -20°C. Dilutions of the folic acid stock and test sera were made in freshly prepared 0.22 µm filter sterilized 0.5% sodium ascorbate. Through a series of dilutions, a folic acid standard curve with 100µl/well in duplicate was generated from 0 to 166 pg/ml final concentration. Test sera was diluted with sterile 0.5% sodium ascorbate before the assay. Several dilutions of sera, 1:600-1:2400 final concentration, at 100µl/well in duplicate were tested to ensure that an A_{600} fell within the standard curve. The inoculated media was added at 200µl/well to the 10 x10 Bioscreen honeycomb plate (Fisher

Scientific). The plate was loaded into the Bioscreen C accompanied with EZExperiment 1.26 software (OY Growth Curves Ab Ltd), incubated at 37°C with shaking for 24 hours, and A_{600} read at 24 hours (31).

2.4 Brain tissue morphological analysis

Series of brain tissue samples were stained with 1% cresyl violet (Sigma) to assess morphological changes. Images were taken using Nikkon Ni-U Compound Light microscope. Analysis of morphological changes focused on the hippocampal formation as previous studies demonstrate that this area is particularly vulnerable to 1C metabolism (15,17,21). Thickness measurements of the granular cell layer were taken using Fiji (NIH) (32,33) within the dentate gyrus, Cornu Ammonis 1 (CA1) and 3 (CA3) regions of the hippocampus. The total volume of the hippocampus was measured.

2.5 Immunofluorescence

Antigen retrieval, on paraffin embedded brain tissue sections, was performed prior to staining as previously described (34). A series of brain tissue sections were stained with the following antibodies: anti-choline acetyltransferase (ChAT) (Millipore, AB144P) at 1:100; anti-acetylcholinesterase (AChE) (Sigma, SAB2500018) at 1:100; or anti-active caspase-3 (Cell Signaling, 9662) at 1:100. All samples were co-stained with anti-neuronal nuclei antibody (NeuN, ab104224) (AbCam) at 1:200 and (4',6-diamidino-2-phenylindole) DAPI (Fisher Scientific) at 1:10,000.

Imaging of ChAT and AChE was conducted using a Zeiss Apotome microscope equipped with a camera. Two brain tissue sections with three subfields within each cortex were imaged.

Imaging of active caspase-3 was performed using an Olympus inverted microscope equipped with a camera. The hippocampus was analyzed using two images of CA1, CA3, or three dentate gyrus for each animal. Only cells demonstrating colocalization of ChAT, AChE, or active caspase-3 along with NeuN and DAPI were considered positive and counted.

2.6 Data and statistical analysis

All data analysis was conducted by 2 individuals blinded to experimental groups. Data were analyzed using GraphPad Prism 9.0. Two-way analysis of variance (ANOVA) was used to determine interactions and main effects between diet groups (folic acid deficient or control) and sex (male or female). Analyses included brain and body weight, morphological measurements, immunofluorescence of choline metabolism (AChE and ChAT), and active caspase-3. Pairwise post-hoc testing was performed using the Tukey's multiple comparison test. In all analyses, $p<0.05$ was considered statistically significant.

3. Results

3.1 Body and brain weight

Brain and body weight were collected at time of euthanization. There was a sex difference in body weight ($F_{(1, 14)} = 48.48$, $p < 0.001$), but there was no effect of diet ($F_{(1, 14)} = 1.97$, $p = 0.18$). There was no difference in brain weight observed (Sex: $F_{(1,14)} = 2.58$, $p = 0.13$, Diet: $F_{(1,14)} = 1.13$, $p = 0.31$). When analyzing brain weight as a percent of body weight, females had a higher percent brain weight ($F_{(1,14)} = 73.99$, $p < 0.0001$). Female FD mice had smaller brain weight as percent of body ($F_{(1,14)} = 6.217$, $p = 0.026$).

3.2 Serum folate measurements

Folate levels were measured in serum collected from all animals to confirm dietary deficiency. There was no sex difference ($F_{(1,14)} = 2.40$, $p = 0.14$). Compared to CD mice, animals maintained on the FD had reduced levels of folate in their serum (Figure 1A; $F_{(1,14)} = 10.52$, $p = 0.0059$).

3.3 Hippocampal volume and morphology

Previous work has shown that the hippocampus is affected by a 1C deficiency (15,20,22). The hippocampus volume was analyzed using FIJI, and thickness of the dentate gyrus, CA1, and CA3 regions. There was no sex difference ($F_{(1,12)} = 0.37$, $p = 0.55$) in the total hippocampal volume. However, there was a trend for lower hippocampal volume of mice maintained on FD (Figure 1B; $F_{(1,12)} = 4.37$, $p = 0.059$). There was no sex ($F_{(1,12)} = 0.86$, $p = 0.37$) or diet ($F_{(1,12)} = 1.15$, $p = 0.31$) differences in the thickness of the dentate gyrus (Figure 1C). Representative images of cresyl violet staining from all groups are shown in Figure 2A. For the CA1 thickness, there was no sex difference ($F_{(1,13)} = 0.07$, $p = 0.88$), but a trend for a diet differences ($F_{(1,13)} = 4.28$ $p = 0.06$) was observed. There was no sex difference in the thickness of CA3 regions ($F_{(1,13)} = 0.94$, $p = 0.35$). However, there was a decrease in the thickness of the CA3 region of mice maintained on a FD diet (Figure 2B; $F_{(1,13)} = 37.3$, $p < 0.0001$).

3.4 Hippocampal apoptosis

Hippocampal apoptosis was assessed by counting active caspase-3 positive neuronal cells, such cells were counted within the dentate gyrus, CA1, and CA3 regions of the hippocampus. There was no difference in neuronal apoptosis between groups within the dentate gyrus (sex: $F_{(1,12)} = 0.035$, $p = 0.86$, diet: $F_{(1,13)} = 0.033$, $p = 0.86$). There was a trend for a sex difference of apoptotic neurons within the CA1 region ($F_{(1,12)} = 3.502$, $p = 0.084$), but there was no difference

between dietary groups (F $_{(1,12)}$ = 0.00083, p = 0.99). Representative images of active caspase-3 staining within the CA3 are shown in Figures 3A to D. There was a difference in the number of positive apoptotic neurons between females and males (Figure 3E; (F $_{(1,12)}$ = 7.36, p = 0.019)), but no dietary difference (F $_{(1,12)}$ = 1.44, p = 0.25).

3.5 Cortical choline metabolism

Neuronal cortical choline metabolism was characterized by counting the number choline acetyltransferase (ChAT) positive cells. This enzyme is involved in the synthesis of acetylcholine at the synapse. Representative images are shown in Figure 4A to D. Quantification revealed no difference between sexes (F $_{(1,12)}$ = 1.79, p = 0.21), however, there were increases in ChAT positive neurons in the FD groups when compared to CD mice (Figure 4E; F $_{(1,12)}$ = 4.58, p = 0.050).

To confirm changes in choline metabolism in brain tissue, we also measured acetylcholine esterase (AchE) levels in neurons within the cortex. Representative images are shown in Figure 5A to D. Quantification revealed no difference between sexes (F $_{(1,12)}$ = 0.49, p = 0.49), but did reveal decreases in FD mice compared to CD animals (Figure 5E; F $_{(1,12)}$ = 7.29, p = 0.019).

4. Discussion

One-carbon metabolism involves multiple complex factors that coordinate gene expression and metabolism in response to nutritional signals that impact cellular anabolism, redox homeostasis, and epigenetics. There is growing evidence that sex plays a significant role in one-metabolism and levels of homocysteine (23–26). Despite this, there is limited evidence for sex differences in the brain tissue. Using a mouse model, our study investigated the impact of sex and dietary folic acid deficiency on hippocampal morphology and brain choline metabolism. As expected, the FD mice had lower serum folates than CD mice. Notably, our results indicate a

significant decrease in the thickness of the granular layer within the CA3 region in both males and females maintained on a FD. In cortical tissue we demonstrate changes in cholinergic metabolism between diet groups. The only sex difference we report in our study is the decreased apoptosis observed in the CA3 region of the hippocampus.

Consistent with prior research, our results demonstrate an intrinsic link between choline and folate metabolism (15–18). A folic acid deficient diet resulted in increased levels of ChAT, the enzyme that catalyzes the synthesis of acetylcholine from choline, and decreased levels of AChE, the enzyme that catalyzes the destruction of acetylcholine. Choline can act as a 1C donor in the liver and kidney via betaine homocysteine methyltransferase, especially when folate levels are low (34,35) . This compensatory function of choline shunts choline away from the production of acetylcholine, potentially leading to lower levels of the neurotransmitter (15,16,18) . Therefore, we suggest a potential mechanism for the demonstrated changes in cortical choline metabolism is a compensatory upregulation of ChAT and downregulation of AChE in response to decreased levels of acetylcholine. Low levels of acetylcholine have been associated with cognitive impairments, including mild cognitive impairment and dementia (36). An interesting follow-up to this study would be to assess cognitive function in adult females and males maintained on a folic acid deficient and control diets.

The hippocampus is impacted by a genetic or dietary deficiency in 1C metabolism, as shown by others, our results demonstrate this effect as well (15,17,22). The different areas of the hippocampus and their functions are topics of ongoing research. In humans, the CA3 region has been specifically linked to spatial representations and episodic memory. Additionally, the CA3 region is susceptible to neurodegeneration, possibly accounting for our results (37). Both *in vivo* and *in vitro* studies have demonstrated that reduced levels of folate result in an increase in apoptotic

markers (20,38–40), possibly due to increases in oxidative stress (41), DNA damage (39), or neuroexcitatory mechanisms (42). Recent *in vivo* studies have indicated similar results, with hyperhomocysteinemia, a direct consequence of 1C deficiency, leading to increased autophagy and apoptosis of cortical neurons(43).

While our results did not indicate a diet difference in apoptosis, there was a significant sex difference in apoptosis in the CA3 region of the hippocampus. The CA3 region of the hippocampus is the most interconnected region of the hippocampus and has been implicated in memory functions and susceptible to neurodegeneration (37). In order to measure levels of apoptosis, we stained for active caspase-3, which, as an executioner caspase leading to cell death, acts as an indirect measure of apoptosis (44). Previous research demonstrates a distinct sex difference in the activation of caspases, specifically following brain injury (45,46). These differences could be accounted for by sex hormones, potentially via the pro-apoptotic protein Bax (47). Model studies have indicated a significant sex difference in gross hippocampal morphology within the hippocampus including the dentate gyrus, CA1 and CA3 regions (48). A recent human study indicates that after adjusting for total hippocampal volume, regional sex differences exist, however, they are not present in the CA2/CA3 regions or the dentate gyrus (49).

In conclusion, our study demonstrates stronger diet effects than sex differences in brain tissue of adult mice maintained on a folic acid deficient diet. Further research should aim to include behavioral analysis assessing cognitive function of male and females maintained on a folic acid deficient diet.

Acknowledgments

The authors would like to the College of Graduate Studies for the generous support of this collaboration. CB was funded by the Kenneth A. Suarez Research Fellowship.

Disclosure statement

None

Notes on contribution

Calli Bennett BSc is a Doctor of Osteopathic Medicine student at Midwestern University. Calli graduated with an Honors Bachelor of Science degree in Anthropology from the University of Utah.

Jacalyn Green PhD is a Professor in the Department of Biochemistry and Molecular Genetics at Midwestern University Downers Grove campus.

Mae Ciancio PhD is an Associate Professor in the Biomedical Sciences Program at Midwestern University Downers Grove campus.

Joanna Goral PhD is an Associate Professor in the Department of Anatomy at Midwestern University Downers Grove campus.

Lenore Pitstick BS is a Department Research Coordinator in the Department of Biochemistry and Molecular Genetics at Midwestern University Downers Grove campus.

Matt Pytynia BA is a Senior Research Specialist in the Biomedical Sciences Program at Midwestern University Downers Grove campus.

Alice Meyer MBS is a Senior Research Associate in the Department of Anatomy at Midwestern University Downers Grove campus.

Neha Kwatra BSc is a Doctor of Dental Medicine student at Midwestern University. Neha graduated with a BSc in Physiology from the University of Arizona.

Nafisa M. Jadavji PhD is an Assistant Professor in Biomedical Sciences at Midwestern University (US) and Research Assistant Professor in Neuroscience at Carleton University (Canada).

Data availability statement

Data available on request from the authors.

Figure & Captions

Figure 1. The impact of dietary folic acid deficiency and sex on serum folate levels (A) and hippocampal volume (B), and thickness of dentate gyrus granular cell layer (C). Scatter plot with mean ± SEM of 3 to 5 mice per group. *p<0.05 indicate Tukey's pairwise comparison between indicated groups.

Figure 2. The impact of dietary folic acid deficiency on hippocampal Cornu Ammonis (CA) 1 and 3 region thickness. Representative images from all experimental groups with the CA1 and CA3 regions highlighted (A). Quantification of thickness of the CA3 region (B). Scatter plot with mean ± SEM of 3 to 5 mice per group. *p<0.05 and **p<0.01 indicate Tukey's pairwise comparison between indicated groups.

Figure 3. The impact of dietary folic acid deficiency and sex on active caspase-3 levels within the hippocampal Cornu Ammonis 3 (CA3). Representative images from experimental groups (A -D). Quantification of active caspase-3, neuronal nuclei (NeuN), and 4′,6-diamidino-2-phenylindole (DAPI) positive cells. Scatter plot with mean $\pm$ SEM of 3 to 5 mice per group. *$p<0.05$ indicates sex main effect between males and females.

Figure 4. The impact of dietary folic acid deficiency and sex on choline acetyltransferase (ChAT) levels within the cortex of adult mice. Representative images from all experimental groups (A to D). Quantification of ChAT, neuronal nuclei (NeuN), and 4′,6-diamidino-2-phenylindole (DAPI) positive cells. Scatter plot with mean $\pm$ SEM of 3 to 5 mice per group. * p<0.05 indicate Tukey's pairwise comparison between indicated groups.

Figure 5. The effect of dietary folic acid deficiency and sex on acetylcholine transferase (AChE) levels within the cortex of adult mice. Representative images from all experimental groups (A to D). Quantification of AChE, neuronal nuclei (NeuN), and 4′,6-diamidino-2-phenylindole (DAPI) positive cells. Scatter plot with mean $\pm$ SEM of 3 to 5 mice per group. * $p<0.05$ indicate Tukey's pairwise comparison between indicated groups.

References

1. Murray L, Emmerson J, Jadavji NM. Roles of Folate in Neurological Function. In: Folic Acid: Sources, Health Effects and Role in Disease. Nova Publishers Science Inc; 2017.

2. Burgess K, Bennett C, Mosnier H, Kwatra N, Bethel F, Jadavji NM. The antioxidant role of one-carbon metabolism on stroke. Antioxidants. 2020;9(11):1–16.

3. Caudill MA. Folate bioavailability: Implications for establishing dietary recommendations and optimizing status. Am J Clin Nutr. 2010 May;91(5):1455S-1455S.

4. Crider KS, Bailey LB, Berry RJ. Folic acid food fortification-its history, effect, concerns, and future directions. Nutrients. 2011 Mar;3(3):370–84.

5. Halsted CH, Villanueva JA, Devlin AM. Folate deficiency, methionine metabolism, and alcoholic liver disease. In: Alcohol. 2002. p. 169–72.

6. Morgan SL, Baggott JE, Vaughn WH, Austin JS, Veitch TA, Lee JY, et al. Supplementation with folic acid during methotrexate therapy for rheumatoid arthritis: A double-blind, placebo-controlled trial. Ann Intern Med. 1994 Dec;121(11):833–41.

7. Enderami A, Zarghami M, Darvishi-Khezri H. The effects and potential mechanisms of folic acid on cognitive function: a comprehensive review. Neurol Sci. 2018 Oct;39(10):1667–75.

8. Seshadri S, Beiser A, Selhub J, Jacques PF, Rosenberg IH, D'Agostino RB, et al. Plasma homocysteine as a risk factor for dementia and Alzheimer's disease. N Engl J Med. 2002;346(7):476–83.

9. Casas JP, Bautista LE, Smeeth L, Sharma P, Hingorani AD. Homocysteine and stroke: Evidence on a causal link from mendelian randomisation. Lancet. 2005;365(9455):224–32.

10. Kwon H-M, Lee Y-S, Bae H-J, Kang D-W. Homocysteine as a predictor of early neurological deterioration in acute ischemic stroke. Stroke. 2014 Mar;45(3):871–3.

11. Shi Z, Guan Y, Huo YR, Liu S, Zhang M, Lu H, et al. Elevated Total Homocysteine Levels in Acute Ischemic Stroke Are Associated With Long-Term Mortality. Stroke. 2015;46(9):2419–25.

12. Kumral E, Saruhan G, Aktert D, Orman M. Association of Hyperhomocysteinemia with Stroke Recurrence after Initial Stroke. J Stroke Cerebrovasc Dis. 2016;1–8.

13. Ho PI, Ashline D, Dhitavat S, Ortiz D, Collins SC, Shea TB, et al. Folate deprivation induces neurodegeneration: roles of oxidative stress and increased homocysteine. Neurobiol Dis. 2003;14:32–42.

14. Castro R, Rivera I, Blom HJ, Jakobs C, de Almeida IT. Homocysteine metabolism, hyperhomocysteinaemia and vascular disease: An overview. J Inherit Metab Dis. 2006 Feb;29(1):3–20.

15. Jadavji NM, Deng L, Leclerc D, Malysheva O, Bedell BJ, Caudill MA, et al. Severe methylenetetrahydrofolate reductase deficiency in mice results in behavioral anomalies with morphological and biochemical changes in hippocampus. Mol Genet Metab. 2012 Jun;106(2):149–59.

16. Chan A, Tchantchou F, Graves V, Rozen R, Shea TB. Dietary and genetic compromise in folate availability reduces acetylcholine, cognitive performance and increases aggression: Critical role of S-adenosyl methionine. J Nutr Health Aging. 2008 Apr;12(4):252–61.

17. Bahous RH, Jadavji NM, Deng L, Cosín-Tomás M, Lu J, Malysheva O, et al. High dietary folate in pregnant mice leads to pseudo-MTHFR deficiency and altered methyl metabolism, with embryonic growth delay and short-term memory impairment in offspring. Hum Mol Genet. 2017;26(5):888–900.

18. Dam K, Füchtemeier M, Farr T, Boehm-Strum P, Foddis M, Dirnagal U, et al. Deficiencies in methylenetetrahydrofolate reductase and dietary folic acid alter choline metabolism during chronic hypoperfusion. Behav Brain Res. 2017;321:201–8.

19. Yen C-LE, Mar M-H, Craciunescu CN, Edwards LJ, Zeisel SH. Deficiency in methionine, tryptophan, isoleucine, or choline induces apoptosis in cultured cells. J Nutr. 2002 Jul;132(7):1840–7.

20. Jadavji NM, Deng L, Malysheva OV, Caudill MA, Rozen R. MTHFR deficiency or reduced intake of folate or choline in pregnant mice results in impaired short-term memory and increased apoptosis in the hippocampus of wild-type offspring. Neuroscience. 2015;300:1–9.

21. Troen AM, Chao W-H, Crivello NA, D'Anci KE, Shukitt-Hale B, Smith DE, et al. Cognitive impairment in folate-deficient rats corresponds to depleted brain phosphatidylcholine and is prevented by dietary methionine without lowering plasma homocysteine. J Nutr. 2008 Dec;138(12):2502–9.

22. Chen Z, Schwahn BC, Wu Q, He X, Rozen R. Postnatal cerebellar defects in mice deficient in methylenetetrahydrofolate reductase. Int J Dev Neurosci Off J Int Soc Dev Neurosci. 2005 Aug;23(5):465–74.

23. Levav-Rabkin T, Blumkin E, Galron D, Golan HM. Sex-dependent behavioral effects of Mthfr deficiency and neonatal GABA potentiation in mice. Behav Brain Res. 2011;216(2):505–13.

24. Sadre-Marandi F, Dahdoul T, Reed MC, Nijhout HF. Sex differences in hepatic one-carbon metabolism. BMC Syst Biol [Internet]. 2018 Oct;12(1). Available from: https://pubmed.ncbi.nlm.nih.gov/30355281/

25. Blumkin E, Levav-Rabkin T, Melamed O, Galron D, Golan HM. Gender-Specific Effect of Mthfr Genotype and Neonatal Vigabatrin Interaction on Synaptic Proteins in Mouse Cortex. Neuropsychopharmacology. 2011 Jul;36(8):1714–28.

26. Stanisławska-Sachadyn A, Woodside JV, Brown KS, Young IS, Murray L, McNulty H, et al. Evidence for sex differences in the determinants of homocysteine concentrations. Mol Genet Metab. 2008 Apr;93(4):355–62.

27. Christensen KE, Wu Q, Wang X, Deng L, Caudill MA, Rozen R. Steatosis in mice is associated with gender, folate intake, and expression of genes of one-carbon metabolism. J Nutr. 2010;140(10):1736–41.

28. Kim MH, Kim WH, Kim CG, Kim DW. Synthesis and Evaluation of 99mTc-Labeled Folate-Tripeptide Conjugate as a Folate Receptor-Targeted Imaging Agent in a Tumor-Bearing Mouse Model. Nucl Med Mol Imaging. 2015 Sep;49(3):200–7.

29. Martínez-Vega R, Murillo-Cuesta S, Partearroyo T, Varela-Moreiras G, Varela-Nieto I, Pajares MA. Long-term dietary folate deficiency accelerates progressive hearing loss on CBA/Ca mice. Front Aging Neurosci. 2016;8(AUG):1–8.

30. Flurkey K, Currer JM, Harrison DE. Mouse Models in Aging Research. In: Fox J, Barthold S, Davisson M, Newcommer C, Quimby F, Smith A, editors. The-Mouse-in-Biomedical-Research, Normative Biology, Husbandry, and Models [Internet]. 2nd ed. Elsevier.; 2007. p. 637–72. Available from: https://mouseion.jax.org/stfb2000_2009/1685

31. Molloy AM, Scott JM. Microbiological assay for serum, plasma, and red cell folate using cryopreserved, microtiter plate method. Methods Enzymol. 1997;281:43–53.

32. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: An open-source platform for biological-image analysis. Nat Methods. 2012 Jul;9(7):676–82.

33. Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, et al. ImageJ2: ImageJ for the next generation of scientific image data. BMC Bioinformatics. 2017 Dec;18(1):529–529.

34. Prieur EAK, Pjetri E, Zeisel SH, Jadavji NM. Reduced brain volume and impaired memory in betaine homocysteine S-methyltransferase knockout mice. Appl Physiol Nutr Metab. 2017;42(11).

35. Niculescu MD, Zeisel SH. Diet, Methyl Donors and DNA Methylation: Interactions between Dietary Folate, Methionine and Choline. J Nutr. 2002 Aug;132(8):2333S-2335S.

36. Sabri O, Meyer PM, Gräf S, Hesse S, Wilke S, Becker GA, et al. Cognitive correlates of α4β2 nicotinic acetylcholine receptors in mild Alzheimer's dementia. Brain. 2018 Jun;141(6):1840–54.

37. Cherubini E, Miles R. The CA3 region of the hippocampus: How is it? What is it for? How does it do it? Front Cell Neurosci [Internet]. 2015 Feb;9(FEB). Available from: https://pubmed.ncbi.nlm.nih.gov/25698930/

38. Kasture VV, Sundrani DP, Joshi SR. Maternal one carbon metabolism through increased oxidative stress and disturbed angiogenesis can influence placental apoptosis in preeclampsia. Life Sci. 2018;206(May):61–9.

39. Kruman II, Culmsee C, Chan SL, Kruman Y, Guo Z, Penix L, et al. Homocysteine elicits a DNA damage response in neurons that promotes apoptosis and hypersensitivity to excitotoxicity. J Neurosci Off J Soc Neurosci. 2000 Sep 15;20(18):6920–6.

40. Kruman II, Kumaravel TS, Lohani A, Pedersen WA, Cutler RG, Kruman Y, et al. Folic acid deficiency and homocysteine impair DNA repair in hippocampal neurons and sensitize them to amyloid toxicity in experimental models of Alzheimer's disease. J Neurosci Off J Soc Neurosci. 2002 Mar 1;22(5):1752–62.

41. Hoffman M. Hypothesis: Hyperhomocysteinemia is an indicator of oxidant stress. Med Hypotheses. 2011 Dec;77(6):1088–93.

42. Lipton SA, Kim W-K, Choi Y-B, Kumar S, D'Emilia DM, Rayudu PV, et al. Neurotoxicity associated with dual actions of homocysteine at the N-methyl-D-aspartate receptor. Proc Natl Acad Sci. 1997 May 27;94(11):5923–8.

43. Zhang JW, Yan R, Tang YS, Guo YZ, Chang Y, Jing L, et al. Hyperhomocysteinemia-induced autophagy and apoptosis with downregulation of hairy enhancer of split 1/5 in cortical neurons in mice. Int J Immunopathol Pharmacol. 2017;30(4):371–82.

44. McIlwain DR, Berger T, Mak TW. Caspase functions in cell death and disease. Cold Spring Harb Perspect Biol. 2013 Apr;5(4):1–28.

45. Liu F, Li Z, Li J, Siegel C, Yuan R, Mccullough LD. Sex differences in caspase activation after stroke. Stroke. 2009 May;40(5):1842–8.

46. Siegel C, Li J, Liu F, Benashski SE, McCullough LD. miR-23a regulation of X-linked inhibitor of apoptosis (XIAP) contributes to sex differences in the response to cerebral ischemia. Proc Natl Acad Sci U S A. 2011 Jul;108(28):11662–7.

47. Forger NG, Rosen GJ, Waters EM, Jacob D, Simerly RB, De Vries GJ. Deletion of Bax eliminates sex differences in the mouse forebrain. Proc Natl Acad Sci U S A. 2004 Sep;101(37):13666–71.

48. Yagi S, Chow C, Lieblich SE, Galea LAM. Sex and strategy use matters for pattern separation, adult neurogenesis, and immediate early gene expression in the hippocampus. Hippocampus. 2016 Jan;26(1):87–101.

49. van Eijk L, Hansell NK, Strike LT, Couvy-Duchesne B, de Zubicaray GI, Thompson PM, et al. Region-specific sex differences in the hippocampus. NeuroImage. 2020 Jul;215:116781–116781.