Title   Plasticity of fibroblast transcriptional response to physical and biochemical cues revealed by dynamic network analysis

Authors/Affiliations

Pilhwa Lee[1]

[1] Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI, USA

Corresponding Author   Pilhwa Lee (jpilpo77@gmail.com)

Lead Contact   Pilhwa Lee

Summary

Data on human skin fibroblast transcriptional responses to external cues were used to reconstruct dynamic gene regulatory networks. The goal of the reconstruction was to determine dynamic network interactions (quantitative predictive relationships of mutual regulatory influences of and on transcription factor expression) from time course data on 56 transcript expression levels obtained following different external cues. The inherently under-determined nature of this problem was addressed in part by excluding putative regulatory motifs that did not appear to be functional in multiple independent experiments from different independent external perturbations. Data were obtained from a previously published experiment in which the 56 transcripts were assayed by bioluminescence in live cells cultured on substrates of varying levels of stiffness and exposed to different levels of arginylglycylaspartic acid (RGD) peptide. The inferred dynamical networks were validated via comparison of predictions to *a priori* known interactions from gene databases. We discovered that exposure to different substrate stiffnesses and to RGD stimulate responses that are mediated through *GATA4*, *SMAD3/4*, *ETS-1*, and *STAT5* and other genes, which can initiate hypertrophic, fibrotic, and inflammatory responses. The developed dynamical system identification method for discovering new mechanotransduction pathways is applicable to the identification of gene regulatory networks in numerous emerging applications where time-series data on multiple state variables and from multiple external perturbations are available.

Introduction

Fibroblasts, which are widely present in many tissue types, contribute to the regulation of the extracellular environment by secreting constituent proteins of the extracellular matrix. By influencing the mechanical properties of the microenvironment, fibroblasts play essential roles in numerous physiological, developmental, and pathophysiological disease processes. Specific developmental and disease processes in which fibroblast functions are known to make important contributions are cell lineage specification (*1-10*), epithelial-mesenchymal interaction (*11-13*), and epithelial-mesenchymal transition (*14-18*) as well as carcinogenesis and metastasis (*19-31*) and immunogenic plasticity in inflammatory diseases (*32-35*). In each of these settings, the ways in which fibroblasts respond to particular mechanical and biochemical stimuli determine how they operate. For example, with certain cardiac-specific factors as biochemical stimuli, fibroblasts may be reprogrammed to cardiac progenitors with multi-potency to cardiomyocytes, smooth muscles, and endothelial cells (*8*). Following an epithelial injury in the lung, fibroblasts respond by secretion of pro-fibrotic factors and can be transdifferentiated to myofibroblasts accompanied by ECM stiffening and fibrosis (*33*). Associated with triple-negative breast cancer, fibroblasts may also be transdifferentiated to myofibroblastic cancer-associated fibroblasts (CAF's). One sub-type of CAF induces regulatory T cells to inhibit T effector proliferation in heterogeneity (*31*) and promotes cancer invasion (*25*).

To understand the mechanisms underlying the plasticity of fibroblast responses to different stimuli, Shea et al. used the state-of-the-art live-cell imaging technique, TRAnscriptional Factor CEll ARray (TRACER) (*36*), to obtain time-series data on multiple transcripts from human skin fibroblasts following perturbations in extracellular matrix (ECM) stiffness and arginylglycylaspartic acid (RGD) peptide levels. RGD peptide is ubiquitously responsible for cell adhesion to the ECM. Thus, by perturbing RGD peptide levels and ECM stiffness, fibroblasts are exposed to a range of mechanical stimuli designed to yield a rich, dynamic response in cellular transcriptional states. To probe the system's interactions in the complex response observed by Shea et al., we used a dynamic network modeling approach to investigate the transcriptional response by attempting to reconstruct network-level regulatory pathways with multiple time-point data from the live-cell experiments of Peñalver Bernabé et al. (*36*).

Most approaches in RNA-level transcriptomic investigation are designed to analyze static data from a single time point. Approaches to network inference from time-series data include Bayesian inference, linear regression, and mutual information techniques (*37-41*). Recent advances in the single-cell analysis have generated dynamic models by incorporating a non-physical "pseudo time" to represent changes following a perturbation (*42-45*). Here, we apply a method designed to infer a "physical-time" differential equation-based model for transcription factor expression and regulation from time-series data on multiple transcripts (*46, 47*). We deconvolve gene-to-gene connectivity treating the transcriptomic ensemble as a kinetic dynamical system on randomized networks and projecting to the empirical time-series data from Shea et al. (*36*). Theoretically, this problem is a non-polynomial (NP) hard, that is, generally not possible to solve exactly. We have approximately handled this difficulty by decomposing a network into subnetworks with one-to-many relationships utilizing GPU-ODE based parallel reconstruction of subnetworks (*46, 47*). Those sub-networks are compared to kinetic data by a hybrid approach of global and gradient-based optimizations. By scoring gene-to-gene interaction and deriving *p*-values from multiple conditions, we identify commonly inferred directly associated genes (Figure 6, green lines) and differentially inferred pathways (Figure 6, black lines) from external differential cues. The selection of random networks for testing is unbiased in bootstrapping. i.e., there is no incorporation of *a priori* confirmed gene-to-gene interactions. The result is a set of likely dynamic network interactions that are informed solely by the time-course data and not by prior knowledge.

Our analysis focuses on approximately 60 transcriptional factors mainly involved in proliferation, differentiation, apoptosis, chemical and hormonal receptors. Based on the gene regulatory network (GRN) architecture inferred from the mechano-transduction data, we identify several putative differential mechanisms by which master regulatory genes *GATA4*, *SMAD3/4*, *ETS1*, and *STAT5* relay information from upstream signals to hypertrophic, fibrotic, or inflammatory phenotypes diversely. The influences of ECM stiffness and cell-ECM adhesion on the regulation of these genes are cooperative in some cases and counter-cooperative in others. Validation of these results is based on *a priori* known interactions from published databases. The validated results reveal the promise of this approach for larger-scale kinetic network reconstruction.

Methods

Transcriptional activity measurement, TRACER. The expression of transcriptional factor activities of human skin fibroblast was measured in time (3, 6, 9, 12, 27 hours) by the bioluminescent optical technique TRACER (*TR*anscriptional *A*ctivity *CE*ll a*R*rays). Six different external conditions are prescribed by three different RGD levels at one PEG hydrogel stiffness as well as three different PEG hydrogel stiffnesses with one RGD level in combination. That is, for RGD variant conditions, RGD concentrations are prescribed as 0.7, 2.0, and 3.3 mM with hydrogel stiffness 1.5 kPa. For hydrogel stiffness variant conditions, PEG-hydrogel stiffnesses are prescribed as 0.15 kPa, 1.5 kPa, and 6.5 kPa with RGD concentration of 2.0 mM. For full experimental details, see Peñalver Bernabé et al. (*36*).

Gene regulatory network reconstruction. The process of reconstructing fibroblast gene regulatory network takes five steps: (1.) Clustering, (2.) Identification of sub-networks, (3.) Development of consensus model from sub-network models, (4.) Global network construction, and (5.) Network validation by *a priori* known databases (Figure 1).

1. Clustering. For each of the six independent experiments representing different external conditions, the time courses of the 56 measured transcripts were compared to one another to identify clusters with similar behavior. For a given experiment, we denote the measured data for transcript i at time point t_k as $x_i(t_k)$. The first step in clustering is to normalize the data

$$y_i(t_k) = \frac{x_i(t_k) - \min(x_i)}{\max(x_i) - \min(x_i)},$$

where $\min(x_i)$ and $\max(x_i)$ are the minimum and maximum values of x_i over the time course.

The clustering of transcript time courses for a given experimental condition is based on several independent metrics of similarity. The first metric, $D_1(i,j)$, is the temporal correlation of transcriptional factors i and j defined:

$$D_1(i,j) = \frac{\sum_k (y_i(t_k) - \bar{y}_i)(y_j(t_k) - \bar{y}_j)}{\|y_i - \bar{y}_i\| \|y_j - \bar{y}_j\|},$$

where $\bar{y}_i$ is the mean value of $y_i(t_k)$ over a given time course. This calculation is a normalized inner-product of $y_i(t_k)$ and $y_j(t_k)$ with their mean values as baselines.

The second metric is a measurement of similarity in the local slopes of $y_i(t_k)$ and $y_j(t_k)$:

$$D_2(i,j) = \sqrt{\sum_k \frac{(\Delta y_i(t_k) - \Delta y_j(t_k))^2}{(\Delta y_i(t_k))^2}},$$

where $\Delta y_i(t_k) = y_i(t_{k+1}) - y_i(t_k)$.

The third metric of similarity is based on the numbers of convex and concave critical points estimated to exist in the time courses for $y_i(t_k)$ and $y_j(t_k)$. Convex critical points are identified at times t_k when $y_i(t_{k-1}) > y_i(t_k)$ and $y_i(t_k) < y_i(t_{k+1})$. Similarly, concave critical are identified at times t_k when $y_i(t_{k-1}) < y_i(t_k)$ and $y_i(t_k) > y_i(t_{k+1})$.

Given these three metrics, two time profiles from transcription factors i and j are clustered together when: (1.) the indicator of temporal correlation $D_1(i,j) > \alpha$; (2.) the indicator of differences in temporal gradients $D_2(i,j) < \beta$; and (3.) the number of critical points in $y_i(t_k)$ and $y_j(t_k)$ are the same. For the calculations presented here, the parameters α and β are set to 0.9 and 0.45, respectively.

The original 56 gene time courses were clustered into 47 clusters for the RGD-high and stiffness-medium experiment, 45 clusters for the RGD-middle and stiffness-medium experiment, 46 clusters for the RGD-low and stiffness-medium, 37 clusters for the RGD-middle and stiffness-hard experiment, 44 clusters for the RGD-middle and stiffness-medium experiment, and 37 clusters for the RGD-middle and stiffness-soft experiment. Clusters for the six experiments are illustrated in Supplementary Figures S1-S3.

2. Identification of sub-networks. Expression of transcripts is assumed to be governed by the following general model (46):

$$\dot{x}_j(t) = r_j^0 \frac{\Sigma_{i \in I_j^A}\left(\frac{x_i(t-\tau)}{K_{i,j}^A}\right)^n + e_j}{1 + \Sigma_{i \in I_j^A}\left(\frac{x_i(t-\tau)}{K_{i,j}^A}\right)^n + \Sigma_{i \in I_j^I}\left(\frac{x_i(t-\tau)}{K_{i,j}^I}\right)^n + e_j} - d_j x_j \,, \qquad \text{(Eq. 1)}$$

where the terms in the expression are defined as follows:

$x_j(t)$: mRNA expression of gene j at time t
r_j^0: basal transcription rate of gene j
d_j: mRNA degradation rate of gene j
I_j^A, I_j^I: a set of indices of activators and inhibitors on gene j
$K_{i,j}^A$: dissociation constant for activation of gene j by gene i
$K_{i,j}^I$: dissociation constant for inhibition of gene j by gene i
e_j: external signals on gene j
τ: time delay between transcription and translation
n: cooperative Hill-coefficient

The model assumes competitive and cooperative binding of inhibitory and activating factors. The time delay constant τ and the Hill-coefficient n are prescribed as 1 hour and 2, respectively. The other parameters, activation/inhibition indices $\{I_j^A, I_j^I\}$ and kinetics parameters $\{r_j^0, d_j, e_j, K_{i,j}^A, K_{i,j}^I\}$, are estimated by fitting with the experimental data.

The details of the model identification process are described in detail in (46). In brief, the model of Equation (1) is applied to each cluster, rather than at the individual gene level. For a given cluster for a given experiment, the governing equation is treated as a one-dimensional (one state variable) ordinary differential equation for a given variable $x_j(t)$ by using shape-preserving piecewise cubic interpolations of the measured data for other variables $x_i(t)$ on the right-hand side of Equation (1). Each of these one-

dimensional problems is highly underdetermined with multiple solutions that can equivalently match the observed data for a given variable $x_j(t)$. The algorithm works by performing a massively parallel search for model structures (defined by the activation/inhibition indices $\{I_j^A, I_j^I\}$) and parameter values $\{r_j^0, d_j, e_j, K_{i,j}^A, K_{i,j}^I\}$ associated with each putative structure. Using a GPU architecture, the algorithm returns >100 putative models at once for each cluster for each experimental condition.

Figures 2 shows time-course data associated with the clusters (red lines) along with the best five and worst five model fits from the total ensemble of model fits for each cluster. The time-course data is from the RGD-high and stiffness-medium condition. The blue curves are clustered to each red curve as a unique feature based on the clustering three metrics (Figure 2a). From each cluster profile, a subnetwork is reconstructed with randomized bootstrapping by GPU-ODE integration and 100 times fitting within the error threshold. For the time course profiles averaged among the clustered genes from experiments (red), best five and worst five fitted profiles generated by the kinetic model on random networks (black) are shown together (Figure 2b).

<u>3. Development of consensus model from sub-network models.</u> Step 2 (identification of sub-networks) yields sets of putative activation and inhibition interactions between genes in the clusters. Each cluster-cluster interaction represents $n \times m$ potential gene-gene interactions, where n and m are the numbers of genes in each cluster. To judge the likelihood of a putative gene-gene interaction, we first enumerate the $n \times m$ potential interactions for each cluster-cluster interaction in the set of sub-networks identified for all clusters from all experiments.

We assess the likelihood of a given identified gene-gene interaction representing a biologically real interaction based on the frequency with which that interaction appears in subnetworks identified from the six independent experiments. Our underlying assumption is that the greater the frequency with which a given interaction appears, the more likely that interaction represents real biological functions. For each putative interaction, we estimate the equivalent of an "h-index" citation index, h. Here, the factor h is defined as the fraction of experiments for which a given interaction appears with a frequency equal to h. For example, if for a given interaction $h = 0.25$, then this interaction appears with a frequency of 25% or greater in 25% or more subnetworks identified for the independent experiments.

In each condition, the h-index of statistical fidelity of activation from gene i to gene j is defined as (the occurrence of accepted random networks with activation from gene i to gene j) / (total occurrence of accepted random networks). The factor of h for inhibition from gene i to gene j is similarly calculated. The overall "h-index" is calculated by fitting the sorted collection of six individual "h-indices" from six conditions to $f = \frac{a}{x+b}$, and taking the shortest distance from the origin to the fitted reciprocal curve (Figure 3a). The overall "h-index" is used for the consensus model integrating "h-indices" from the six external conditions (Figure 3b-d).

For each external condition, we also define the significance of gene-to-gene interactions by their p-values. The p-values are calculated from statistical distribution of gene-to-gene binding occurrence events (Figure 4a and 4c) following the steps:

(1) Regarding cluster-to-cluster interactions, the occurrences of activation and inhibition of each cluster to a specific cluster are counted, from which a probability density distribution of occurrences of interaction is obtained (Figure 4a and 4c). There is a mapping $M(i, j)$ from each cluster i to its occurrence of interaction with the specific cluster j.
(2) From these probability density distributions of occurrences, we derive cumulative probability density distributions for activation and inhibition interactions, respectively.

(3) For each interaction from cluster i to cluster j (activation or inhibition), the occurrence of interaction $M(i,j)$ points to the corresponding cumulative probability density "$p_{\text{cummulation}}$".

(4) The cluster-to-cluster interactions are identified as significant for $1\text{-}p_{\text{cummulation}} < 0.05$ and colored as violet in cluster unit (Figure 4b and 4d). The insignificant clusters are colored as light violet.

(5) The p values of $n \times m$ gene-gene interaction are duplicated from the corresponding cluster-cluster interaction.

<u>4. Global network construction.</u> For a given assumed cutoff value for the frequency parameter h, we obtain a set of internal network interactions defining the global network for the 56 transcription factors studied here. Fig. 3 shows examples of network with 533 edges ($h = 0.24$), 248 edges ($h = 0.26$), and 103 edges ($h = 0.28$). The lower the cutoff h value, the more interaction edges accepted into the network.

Our goal is to simulate and explain whole-network behavior in terms of Equation (1) where each transcript is treated as a dependent variable, rather than in terms of one-variable sub-networks. To do this, we need to obtain global estimates of the activation and inhibition dissociation constants associated with each activation and inhibition interaction in the network, as well as the external signal factors for each gene for each of the six experimental conditions.

To simulate the global network in an integrated manner the external cues to 56 genes are modeled by a linear combination of saturating effects from two independent cues from RGD peptide and PEG-hydrogel stiffness:

$$e_{\text{RGD},i} = \frac{\max\left(e_{\max,i}^{\text{RGD}}, 0\right)}{1 + \left(\frac{k_{\text{RGD},i}}{[\text{RGD}]}\right)^{n_{\text{RGD},i}}} - \frac{\min\left(e_{\max,i}^{\text{RGD}}, 0\right)}{1 + \left(\frac{[\text{RGD}]}{k_{\text{RGD},i}}\right)^{n_{\text{RGD},i}}}, \qquad \text{(Eq. 2)}$$

$$e_{\text{STIFF},i} = \frac{\max\left(e_{\max,i}^{\text{STIFF}}, 0\right)}{1 + \left(\frac{k_{\text{STIFF},i}}{k_{\text{STIFFNESS}}}\right)^{n_{\text{STIFF},i}}} - \frac{\min\left(e_{\max,i}^{\text{STIFF}}, 0\right)}{1 + \left(\frac{k_{\text{STIFFNESS}}}{k_{\text{STIFF},i}}\right)^{n_{\text{STIFF},i}}},$$

$$e_i = e_{\text{RGD},i} + e_{\text{STIFF},i},$$

where $\{e_{\max,i}^{\text{RGD}}, k_{\text{RGD},i}, n_{\text{RGD},i}, e_{\max,i}^{\text{STIFF}}, k_{\text{STIFF},i}, n_{\text{STIFF},i}\}$ are estimated by fitting whole-network function to the data illustrated in Fig. 5.

For a given gene i the external cues $e_{\text{RGD},i}$ and $e_{\text{STIFF},i}$ can represent either activation or inhibition depending on the sign of the parameters $e_{\max,i}^{\text{RGD}}$ and $e_{\max,i}^{\text{STIF}}$.

Because the experiments have no data for $t = 0$, we obtain a long-time steady state for the isolated system ($[\text{RGD}] = 0$, $k_{\text{STIFFNESS}} = 0$) to use at the initial condition for each simulation. To estimate global kinetic parameters, the 56 gene profiles from 6 conditions are fit (Figure 5) using a combination of stochastic simulated annealing, gradient-based optimization, and the globally convergent method of moving asymptotes (GCMMA) (48). Using this hybrid approach, we obtain a number of independent realization of network parameter sets associated with eight relatively equivalent fits to the data (Figure S7-S14).

The directly associated genes from RGD and hydrogel stiffness cues are determined statistically with 8 replicates of the hybrid approach. We assume that statistically significant direct associations exist as activation or inhibition constitutively, i.e. commonly exist for all 6 different external conditions. The p-values are calculated from statistical distribution of cue-to-gene association occurrence events through the following steps:

1) For each replicate among 8, the association from each cue (RGD or hydrogel stiffness) to genes is identified if the input strengths $e_{\mathrm{max},i}^{\mathrm{RGD}}$ and $e_{\mathrm{max},i}^{\mathrm{STIFF}}$ in Eq. 2 are ranked within 5 for both activating and inhibiting roles.
2) Regarding cue-to-gene interactions, the occurrences of activation and inhibition of each cue to a specific gene are counted, from which a probability density distribution of occurrences of interaction is obtained. There is a mapping $M(i,j)$ from each cue i to its occurrence of interaction with the specific gene j.
3) From these probability density distributions of occurrences, we derive cumulative probability density distributions for activation and inhibition interactions, respectively.
4) For each interaction from cue i to gene j (activation or inhibition), the occurrence of interaction $M(i,j)$ points to the corresponding cumulative probability density "$p_{\mathrm{cummulation}}$".
5) The cue-to-gene interactions are identified as significant for 1-$p_{\mathrm{cummulation}}$ < 0.05 and colored as green lines (Figure 6).
6) The inferred cue-to-gene interactions are validated with *a priori* known gene databases (Table 1) and identified as validated (solid green lines) when there exists a prior report of the specific cue-to-gene interaction. Otherwise, they are represented by dotted green lines.

In the cue-gene interactions, the inferred activations (Figure 6, green →) are still "activation" and "up-regulation" whether the cues are strong or weak. Similarly, the inferred inhibitions (Figure 6, green --|) are still "inhibition" and "down-regulation" independent to the strengths of the cues.

Similarly, statistically significant gene-to-gene interactions are identified for each condition (Figure 4). By overlapping significant ($p<0.05$) gene-to-gene connectivities from two conditions (e.g., RGD-high and RGD-middle, RGD-middle and RGD-low), differential networks are constructed (Figure 6, black colored). The inferred four networks are validated with *a priori* known gene databases (Table 1) and identified as validated (solid lines) when there exists a prior report of the specific gene-to-gene interaction.

Data Availability

The manuscript contains all information necessary for the data (*36*) and the main codes (*47*) used for the analysis. If additional information is required, including the codes used in the model simulation, data will be readily available for editors, reviewers, and readers upon request.

Results and Discussion

Identified gene network pathways are validated from a priori known gene databases.

As detailed in Table 1, of 159 total interactions identified in our global analysis, 74 are validated based on prior reports in the literature.

Concerning hydrogel stiffness, hydrogel activates *STAT1*, *STAT5*, and *PEA3*, but inhibits RAR and PR. On the other hand, RGD peptide activates stemness-associated *SOX* and *GATA*. However, anti-hypertrophic transcriptional factor *FOXO3A* and ECM remodeling associated *ETS-1* are inhibited from the RGD peptide. We have placed these direct associations of mechano-transduction as common through all six conditions, i.e., those upstream interactions from dual cues (cell-ECM adhesion and ECM stiffness) to directly associated genes are prescribed as existing for all differential regulatory networks (Figure 6, green lines). The identified four differential networks are the following:

(Differential network-I) Overlapping significant gene-to-gene interactions from (RGD-high and hydrogel stiffness-medium) with those from (RGD-middle and hydrogel stiffness- medium): Among 26 gene-to-gene interactions, 19 are relevant (Figure 6a). There are a few interesting inferred pathways identified as relevant based on *a priori* known databases. When RGD is high, *ETS-1* is inhibited by RGD, but co-activated by *AR* and hydrogel stiffness mediated by *PEA3*. *RAR* inhibited by ECM stiffness is transmitted through *RAR* → *p53* → *RUNX1* → *SP1*. *GATA* is coactivated by RGD and *SRE*. On the other hand, *FOXO3A* is co-inhibited by *GATA4* and RGD.

(Differential network-II) Overlapping significant gene-to-gene interactions from (RGD-middle and hydrogel stiffness-medium) with those from (RGD-low and hydrogel stiffness-medium): Among 44 gene-to-gene interactions, 18 are relevant (Figure 6b). When RGD is low, RGD is inferred to activate *NFkB* through *SRF* (RGD → *SOX* --| *SRF* → *NFkB*). Also, *ETS-1* is co-inhibited by RGD, *RUNX2*, as well as *AP1* mediated by *STAT3*.

(Differential network-III) Overlapping significant gene-to-gene interactions from (RGD-middle and hydrogel stiffness-hard) with those from (RGD-middle and hydrogel stiffness-medium): Among 33 gene-to-gene interactions, 18 are relevant (Figure 6c). When hydrogel is hard, PR is inhibited by stiffness but activated by *GATA2*. *SOX* is co-activated by *TCF/LEF* and RGD. *GATA4* is activated by *YY1* and RGD mediated by *FOXO3A* but inhibited by hydrogel stiffness mediated by *RAR*.

(Differential network-IV) Overlapping significant gene-to-gene interactions from (RGD-mid and hydrogel stiffness-medium) with those from (RGD-mid and hydrogel stiffness-soft): Among 56 gene-to-gene interactions, 19 are relevant (Figure 6d). When hydrogel is soft, *GATA* is co-activated by RGD and *SRF*. Also, *RAR* inhibition is attenuated by soft hydrogel and transmitted to activate *GAS*, *GATA4*, and *SMAD3/4*.

Master regulatory transcriptional factors GATA4, SMAD3/4, ETS-1, and STAT5 and associated pathways respond adaptively with differential conditions from mechano-transduction.

(Master regulator I) *GATA4* and *SMAD3/4*: In different conditions, pathways around *GATA4* do switch. When hydrogel stiffness turns from hard to soft, the pro-fibrotic *SMAD3/4* is activated from *YY1* and *RAR*. When hydrogel stiffness is hard or soft, and the RGD level is in the middle, *GATA4* is "inhibited" by *FOXO3A*. However, when RGD is high, *GATA4* does "inhibit" *FOXO3A* together with RGD, i.e., *GATA4 - FOXO3A* has a mutual feedback structure depending on conditions (Figure 7a).

(Master regulator II) *ETS-1*: The network pathway with target gene *ETS-1* also switches depending on the RGD level. When the RGD level is low, *AP1* and *RUNX2* co-inhibit *ETS-1* together with RGD. However, when the RGD level is high, *AR* and hydrogel stiffness mediated by *PEA3* "co-activate" *ETS-1*. The stability of *ETS-1* is adaptively regulated by the RGD level and hydrogel stiffness (Figure 7b).

(Master regulator III) *STAT5*: When RGD is low, and hydrogel stiffness is medium, *STAT5* is co-activated by hydrogel stiffness and *ELK1*. When the condition is with the middle level of RGD and soft hydrogel, *STAT5* is co-activated by hydrogel stiffness and low adhesion mediated by *SOX*. *STAT5*, as a potential inflammatory signal, has an intrinsic unstable structure (Figure 7c).

Prediction of specific phenotypes

(Phenotype I) Fibrotic transdifferentiation: TGF-ß and *SMAD3/4* (*49*) are primary players in the fibrotic transdifferentiation from fibroblast to myofibroblast (*50*). Our analysis predicts that mechano-

transduction, i.e., when the hydrogel is soft, and *RAR* does mediate, shows a fibrotic potential with targeting *SMAD3/4*.

(Phenotype II) Regulation of inflammation and hypertrophy: Fibroblast deeply involves in hypertrophic and inflammatory phenotypes. Our analysis predicts those potentials by *GATA4* (*51-53*) and *STAT5* (*54*), respectively. Especially, hard hydrogel and low RGD drive anti-hypertrophic phenotype targeting to *GATA4*.

(Phenotype III) Stem cell lineage determination: Fibroblast is a significant component in the extracellular matrix and microenvironment determining differentiation fates in stem cell lineages. Among others, Stiffness --| *RAR* → *p53* → *RUNX1* axis and RGD → *SOX* → *STAT5* might be new pathways to be investigated further. *GATA* also involves in stemness. When RGD is high, *GATA* is co-activated by RGD and *SRE*. When hydrogel is soft, *GATA* is co-activated by RGD and *SRF*.

(Phenotype IV) Activation for invasive and metastatic states: In cancer metastasis, the transition to invasive migration is preconditioned by ECM remodeling and protease by MMP proteins. We have identified *ETS-1* as a potential master regulator through mechano-transduction (*55, 56*). When RGD is low, *ETS-1* is inhibited by proliferation driven by *AP1* and *STAT3* and keeps a stationary stage rather than invasive migration. On the other hand, when RGD is high, *ETS-1* is activated by hydrogel stiffness, highly good condition for a metastatic transition.

Utilization of a priori known pathways as a constraint in subnetwork reconstruction. The present reconstruction process is unbiased and does not utilize *a priori* known pathways. This kind of approach has pros and cons. Through the solely randomized inferences, we may discover a new pathway and functional architecture. This unbiased approach may predict an actual possibility of pathways for phenotypes of rare diseases. However, this does provide higher false discovery rates (FDR) for highly confirmed gene regulatory subnetwork modules. In a compromise, we plan to incorporate highly probable building blocks of pathways as given without randomized filtration depending on the features of transcriptomics and cell types.

Oscillation in long-run dynamics and identification of coupled oscillators. In the way of a broader investigation of the inferred gene regulatory networks, we have also obtained a long-run dynamics prescribed with multiple conditions. Interestingly several replicates show oscillatory and non-oscillatory responses depending on the conditions, i.e., bifurcating mechano-sensing features. Those oscillatory features imply some sort of coupled oscillators in the deconvoluted networks. Our inferences are based on a period of 27 hours, and this finite data for the reconstruction may provide limited constraints for the real system. For further validating the authenticity for the mechano-transduction induced oscillation, more data in long-run dynamics will clarify this issue. However, some groups have reported the intrinsic subcellular transports between cytoplasm and nucleus in a feedback loop and associated coupled oscillators (*57, 58*).

Global optimization techniques. We have utilized a hybrid approach combining global stochastic optimization such as simulated annealing and genetic algorithms and local gradient-based nonlinear programming. Some groups have incorporated Particle Swarm Optimization (PSO) (*59*), for which we may test in the future. As nonlinear programming, we have employed a globally convergent method of GCMMA and prescribed a few constraints in fitting the data. Those inequality constraints can be relaxed or further imposed and fine-tuned in the range of the constraints. In the outer loop, we may improve the fitting by a few more iterations of the global-local hybrid optimization.

Validation of gene network inference by knockdown and knockout. We have proposed the plasticity of fibroblast in mechano-transduction based on the differentially identified pathways depending on the

differential conditions. The experimental validation of these predictions is expected to be done by RNAi and CRISPR techniques, and the protein level western blot will confirm the phenotypic functional significance.

Acknowledgements

The author acknowledges valuable discussions and support in the early stages of the manuscript from Daniel Beard. The author also thanks Joseph Decker and Lonnie Shea for the introduction of the experimental data and helpful discussion. This research was partially supported by U01 HL118738-01A1 and NIGMS-P50GM094503.

Author contributions

The author designed research, performed research, analyzed data, and wrote the paper.

Conflict of interests

There is no conflict of interests.

References

1. D. E. Discher, P. Janmey, Y. L. Wang, Tissue cells feel and respond to the stiffness of their substrate. *Science* **310**, 1139-1143 (2005).
2. A. J. Engler, S. Sen, H. L. Sweeney, D. E. Discher, Matrix elasticity directs stem cell lineage specification. *Cell* **126**, 677-689 (2006).
3. M. Chiquet, L. Gelman, R. Lutz, S. Maier, From mechanotransduction to extracellular matrix gene expression in fibroblasts. *Biochim Biophys Acta* **1793**, 911-920 (2009).
4. L. K. Sthanam *et al.*, Biophysical regulation of mouse embryonic stem cell fate and genomic integrity by feeder derived matrices. *Biomaterials* **119**, 9-22 (2017).
5. J. T. Connelly *et al.*, Actin and serum response factor transduce physical cues from the microenvironment to regulate epidermal stem cell fate decisions. *Nat Cell Biol* **12**, 711-718 (2010).
6. A. Urciuolo *et al.*, Collagen VI regulates satellite cell self-renewal and muscle regeneration. *Nat Commun* **4**, 1964 (2013).
7. J. W. Shin, D. J. Mooney, Improving Stem Cell Therapeutics with Mechanobiology. *Cell Stem Cell* **18**, 16-19 (2016).
8. P. A. Lalit *et al.*, Lineage Reprogramming of Fibroblasts into Proliferative Induced Cardiac Progenitor Cells by Defined Factors. *Cell Stem Cell* **18**, 354-367 (2016).
9. F. M. Watt, Engineered Microenvironments to Direct Epidermal Stem Cell Behavior at Single-Cell Resolution. *Dev Cell* **38**, 601-609 (2016).
10. M. Notari *et al.*, The local microenvironment limits the regenerative potential of the mouse neonatal heart. *Sci Adv* **4**, eaao5553 (2018).
11. G. R. Cunha, Mesenchymal-epithelial interactions: past, present, and future. *Differentiation* **76**, 578-586 (2008).
12. D. Ribatti, M. Santoiemma, Epithelial-mesenchymal interactions: a fundamental Developmental Biology mechanism. *Int J Dev Biol* **58**, 303-306 (2014).
13. N. O. Enemchukwu *et al.*, Synthetic matrices reveal contributions of ECM biophysical and biochemical properties to epithelial morphogenesis. *J Cell Biol* **212**, 113-124 (2016).
14. J. L. Leight, M. A. Wozniak, S. Chen, M. L. Lynch, C. S. Chen, Matrix rigidity regulates a switch between TGF-β1-induced apoptosis and epithelial-mesenchymal transition. *Mol Biol Cell* **23**, 781-791 (2012).

15.	R. M. Carew, B. Wang, P. Kantharidis, The role of EMT in renal fibrosis. *Cell Tissue Res* **347**, 103-116 (2012).

16.	R. Du *et al.*, Hypoxia-induced Bmi1 promotes renal tubular epithelial cell-mesenchymal transition and renal fibrosis via PI3K/Akt signal. *Mol Biol Cell* **25**, 2650-2659 (2014).

17.	I. Pastushenko *et al.*, Identification of the tumour transition states occurring during EMT. *Nature* **556**, 463-468 (2018).

18.	T. Rozario, D. W. DeSimone, The extracellular matrix in development and morphogenesis: a dynamic view. *Dev Biol* **341**, 126-140 (2010).

19.	N. A. Bhowmick, E. G. Neilson, H. L. Moses, Stromal fibroblasts in cancer initiation and progression. *Nature* **432**, 332-337 (2004).

20.	A. Orimo *et al.*, Stromal fibroblasts present in invasive human breast carcinomas promote tumor growth and angiogenesis through elevated SDF-1/CXCL12 secretion. *Cell* **121**, 335-348 (2005).

21.	R. Kalluri, M. Zeisberg, Fibroblasts in cancer. *Nat Rev Cancer* **6**, 392-401 (2006).

22.	K. Pietras, A. Ostman, Hallmarks of cancer: interactions with the tumor stroma. *Exp Cell Res* **316**, 1324-1331 (2010).

23.	P. Lu, V. M. Weaver, Z. Werb, The extracellular matrix: a dynamic niche in cancer progression. *J Cell Biol* **196**, 395-406 (2012).

24.	V. M. Weaver *et al.*, Reversion of the malignant phenotype of human breast cells in three-dimensional culture and in vivo by integrin blocking antibodies. *J Cell Biol* **137**, 231-245 (1997).

25.	A. Siletz, E. Kniazeva, J. S. Jeruss, L. D. Shea, Transcription factor networks in invasion-promoting breast carcinoma-associated fibroblasts. *Cancer Microenviron* **6**, 91-107 (2013).

26.	A. De Boeck *et al.*, Differential secretome analysis of cancer-associated fibroblasts and bone marrow-derived precursors to identify microenvironmental regulators of colon cancer progression. *Proteomics* **13**, 379-388 (2013).

27.	A. Naba *et al.*, Extracellular matrix signatures of human primary metastatic colon cancers and their metastases to liver. *BMC Cancer* **14**, 518 (2014).

28.	L. E. Barney *et al.*, A cell-ECM screening method to predict breast cancer metastasis. *Integr Biol (Camb)* **7**, 198-212 (2015).

29.	L. Alba-Castellón *et al.*, Snail1-Dependent Activation of Cancer-Associated Fibroblast Controls Epithelial Tumor Cell Invasion and Metastasis. *Cancer Res* **76**, 6205-6217 (2016).

30.	B. Y. Nabet *et al.*, Exosome RNA Unshielding Couples Stromal Activation to Pattern Recognition Receptor Signaling in Cancer. *Cell* **170**, 352-366.e313 (2017).

31.	A. Costa *et al.*, Fibroblast Heterogeneity and Immunosuppressive Environment in Human Breast Cancer. *Cancer Cell* **33**, 463-479.e410 (2018).

32.	A. J. Booth *et al.*, Acellular normal and fibrotic human lung matrices as a culture system for in vitro investigation. *Am J Respir Crit Care Med* **186**, 866-876 (2012).

33.	C. E. Barkauskas, P. W. Noble, Cellular mechanisms of tissue fibrosis. 7. New insights into the cellular mechanisms of pulmonary fibrosis. *Am J Physiol Cell Physiol* **306**, C987-996 (2014).

34.	A. Gabrielli, E. V. Avvedimento, T. Krieg, Scleroderma. *N Engl J Med* **360**, 1989-2003 (2009).

35.	T. R. Katsumoto, M. L. Whitfield, M. K. Connolly, The pathogenesis of systemic sclerosis. *Annu Rev Pathol* **6**, 509-537 (2011).

36.	B. Penalver Bernabe *et al.*, Dynamic transcription factor activity networks in response to independently altered mechanical and adhesive microenvironmental cues. *Integrative biology : quantitative biosciences from nano to macro* **8**, 844-860 (2016).

37.	N. Friedman, Inferring cellular networks using probabilistic graphical models. *Science* **303**, 799-805 (2004).

38.	A. A. Margolin *et al.*, ARACNE: an algorithm for the reconstruction of gene regulatory networks in a mammalian cellular context. *BMC Bioinformatics* **7 Suppl 1**, S7 (2006).

39. V. A. Huynh-Thu, A. Irrthum, L. Wehenkel, P. Geurts, Inferring regulatory networks from expression data using tree-based methods. *PLoS One* **5**, (2010).

40. N. V. Xuan, M. Chetty, R. Coppel, P. P. Wangikar, Gene regulatory network modeling via global optimization of high-order dynamic Bayesian network. *BMC Bioinformatics* **13**, 131 (2012).

41. A. C. Haury, F. Mordelet, P. Vera-Licona, J. P. Vert, TIGRESS: Trustful Inference of Gene REgulation using Stability Selection. *BMC Syst Biol* **6**, 145 (2012).

42. C. Trapnell *et al.*, The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386 (2014).

43. M. Setty *et al.*, Wishbone identifies bifurcating developmental trajectories from single-cell data. *Nat Biotechnol* **34**, 637-645 (2016).

44. E. Marco *et al.*, Bifurcation analysis of single-cell gene expression data reveals epigenetic landscape. *Proc Natl Acad Sci U S A* **111**, E5643-5650 (2014).

45. L. Haghverdi, M. Büttner, F. A. Wolf, F. Buettner, F. J. Theis, Diffusion pseudotime robustly reconstructs lineage branching. *Nat Methods* **13**, 845-848 (2016).

46. J. N. Bazil, F. Qi, D. A. Beard, A parallel algorithm for reverse engineering of biological networks. *Integrative biology : quantitative biosciences from nano to macro* **3**, 1215-1223 (2011).

47. R. Thiagarajan, A. Alavi, J. T. Podichetty, J. N. Bazil, D. A. Beard, The feasibility of genome-scale biological network inference using Graphics Processing Units. *Algorithms Mol Biol* **12**, 8 (2017).

48. K. Svanberg, A Class of globally convergent optimization methods based on conservative convex separable approximations. *SIAM Journal of Optimization* **12**, 555-573 (2002).

49. J. Massague ́, Seoane, J., and Wotton, D., Smad transcription factors. *Genes Dev.* **19**, 2783–2810 (2005).

50. L. Zhou *et al.*, Inhibition of miR-29 by TGF-beta-Smad3 signaling through dual mechanisms promotes transdifferentiation of mouse myoblasts into myofibroblasts. *PLoS One* **7**, e33766 (2012).

51. C. Kang *et al.*, The DNA damage response induces inflammation and senescence by inhibiting autophagy of GATA4. *Science* **349**, aaa5612 (2015).

52. N. Pilon, D. Raiwet, R. S. Viger, D. W. Silversides, Novel pre- and post-gastrulation expression of Gata4 within cells of the inner cell mass and migratory neural crest cells. *Dev Dyn* **237**, 1133-1143 (2008).

53. R. Zheng, G. A. Blobel, GATA Transcription Factors and Cancer. *Genes Cancer* **1**, 1178-1188 (2010).

54. G. Ferbeyre, R. Moriggl, The role of Stat5 transcription factors as tumor suppressors or oncogenes. *Biochim Biophys Acta* **1815**, 104-114 (2011).

55. J. Dittmer, The biology of the Ets1 proto-oncogene. *Mol Cancer* **2**, 29 (2003).

56. P. Haines, G. H. Samuel, H. Cohen, M. Trojanowska, A. M. Bujor, Caveolin-1 is a negative regulator of MMP-1 gene expression in human dermal fibroblasts via inhibition of Erk1/2/Ets1 signaling pathway. *J Dermatol Sci* **64**, 210-216 (2011).

57. S. Sampattavanich *et al.*, Encoding Growth Factor Identity in the Temporal Dynamics of FOXO3 under the Combinatorial Control of ERK and AKT Kinases. *Cell Syst* **6**, 664-678.e669 (2018).

58. A. Hubaud, I. Regev, L. Mahadevan, O. Pourquié, Excitable Dynamics and Yap-Dependent Mechanical Cues Drive the Segmentation Clock. *Cell* **171**, 668-682.e611 (2017).

59. S. Iadevaia, Y. Lu, F. C. Morales, G. B. Mills, P. T. Ram, Identification of optimal drug combinations targeting cellular networks: integrating phospho-proteomics and computational network analysis. *Cancer Res* **70**, 6704-6714 (2010).

60. P. Fischer, D. Hilfiker-Kleiner, Survival pathways in hypertrophy and heart failure: the gp130-STAT3 axis. *Basic Res Cardiol* **102**, 279-297 (2007).

61. C. E. Barcus, P. J. Keely, K. W. Eliceiri, L. A. Schuler, Stiff collagen matrices increase tumorigenic prolactin signaling in breast cancer cells. *J Biol Chem* **288**, 12722-12732 (2013).

62. S. H. Yi, Y. Zhang, D. Tang, L. Zhu, Mechanical force and tensile strain activated hepatic stellate cells and inhibited retinol metabolism. *Biotechnol Lett* **37**, 1141-1152 (2015).

63. E. R. West, M. Xu, T. K. Woodruff, L. D. Shea, Physical properties of alginate hydrogels and their effects on in vitro follicle development. *Biomaterials* **28**, 4439-4448 (2007).

64. S. Kapila, Y. Xie, W. Wang, Induction of MMP-1 (collagenase-1) by relaxin in fibrocartilaginous cells requires both the AP-1 and PEA-3 promoter sites. *Orthod Craniofac Res* **12**, 178-186 (2009).

65. A. Petersen, P. Joly, C. Bergmann, G. Korus, G. N. Duda, The impact of substrate stiffness and mechanical loading on fibroblast-induced scaffold remodeling. *Tissue Eng Part A* **18**, 1804-1817 (2012).

66. O. Hazenbiller, N. A. Duncan, R. J. Krawetz, Reduction of pluripotent gene expression in murine embryonic stem cells exposed to mechanical loading or Cyclo RGD peptide. *BMC Cell Biol* **18**, 32 (2017).

67. J. C. Chang, S. H. Hsu, D. C. Chen, The promotion of chondrogenesis in adipose-derived adult stem cells by an RGD-chimeric protein in 3D alginate culture. *Biomaterials* **30**, 6265-6275 (2009).

68. Y. Katoh, M. Katoh, Identification and characterization of rat Wnt6 and Wnt10a genes in silico. *Int J Mol Med* **15**, 527-531 (2005).

69. R. S. Nho, J. Kahm, beta1-Integrin-collagen interaction suppresses FoxO3a by the coordination of Akt and PP2A. *J Biol Chem* **285**, 14195-14209 (2010).

70. J. H. Yang, W. H. Briggs, P. Libby, R. T. Lee, Small mechanical strains selectively suppress matrix metalloproteinase-1 expression by human vascular smooth muscle cells. *J Biol Chem* **273**, 6550-6555 (1998).

71. H. Kathuria, Y. X. Cao, M. I. Ramirez, M. C. Williams, Transcription of the caveolin-1 gene is differentially regulated in lung type I epithelial and endothelial cell lines. A role for ETS proteins in epithelial cell expression. *J Biol Chem* **279**, 30028-30036 (2004).

72. C. E. Massie *et al.*, New androgen receptor genomic targets show an interaction with the ETS1 transcription factor. *EMBO Rep* **8**, 871-878 (2007).

73. N. S. Belaguli *et al.*, Cardiac tissue enriched factors serum response factor and GATA-4 are mutual coregulators. *Mol Cell Biol* **20**, 7550-7558 (2000).

74. S. Huh *et al.*, Suppression of the ERK-SRF axis facilitates somatic cell reprogramming. *Exp Mol Med* **50**, e448 (2018).

75. F. Wu, K. Tyml, J. X. Wilson, iNOS expression requires NADPH oxidase-dependent redox signaling in microvascular endothelial cells. *J Cell Physiol* **217**, 207-214 (2008).

76. C. C. Applegate, M. A. Lane, Role of retinoids in the prevention and treatment of colorectal cancer. *World J Gastrointest Oncol* **7**, 184-203 (2015).

77. A. Shilkaitis, A. Green, K. Christov, Retinoids induce cellular senescence in breast cancer cells by RAR-β dependent and independent pathways: Potential clinical implications (Review). *Int J Oncol* **47**, 35-42 (2015).

78. J. W. Lee, A. van Wijnen, S. C. Bae, RUNX3 and p53: How Two Tumor Suppressors Cooperate Against Oncogenic Ras? *Adv Exp Med Biol* **962**, 321-332 (2017).

79. R. Pippa *et al.*, MYC-dependent recruitment of RUNX1 and GATA2 on the SET oncogene promoter enhances PP2A inactivation in acute myeloid leukemia. *Oncotarget* **8**, 53989-54003 (2017).

80. I. Mineva, M. Stamenova, W. Gartner, L. Wagner, Expression of the small heat shock protein alphaB-crystallin in term human placenta. *Am J Reprod Immunol* **60**, 440-448 (2008).

81. H. Zhou, M. E. Dickson, M. S. Kim, R. Bassel-Duby, E. N. Olson, Akt1/protein kinase B enhances transcriptional reprogramming of fibroblasts to functional cardiomyocytes. *Proc Natl Acad Sci U S A* **112**, 11864-11869 (2015).

82. X. Liu *et al.*, LPS-induced proinflammatory cytokine expression in human airway epithelial cells and macrophages via NF-κB, STAT3 or AP-1 activation. *Mol Med Rep* **17**, 5484-5491 (2018).

83. H. Zhang *et al.*, STAT3 restrains RANK- and TLR4-mediated signalling by suppressing expression of the E2 ubiquitin-conjugating enzyme Ubc13. *Nat Commun* **5**, 5798 (2014).

84. Y. Zhang *et al.*, Co-stimulation of the bone-related Runx2 P1 promoter in mesenchymal cells by SP1 and ETS transcription factors at polymorphic purine-rich DNA sequences (Y-repeats). *J Biol Chem* **284**, 3125-3135 (2009).

85. T. Egawa, D. R. Littman, Transcription factor AP4 modulates reversible and epigenetic silencing of the Cd4 gene. *Proc Natl Acad Sci U S A* **108**, 14873-14878 (2011).

86. H. Gu, J. C. Pratt, S. J. Burakoff, B. G. Neel, Cloning of p97/Gab2, the major SHP2-binding protein in hematopoietic cells, reveals a novel pathway for cytokine-induced gene activation. *Mol Cell* **2**, 729-740 (1998).

87. Y. Deng *et al.*, A Blockade of IGF Signaling Sensitizes Human Ovarian Cancer Cells to the Anthelmintic Niclosamide-Induced Anti-Proliferative and Anticancer Activities. *Cell Physiol Biochem* **39**, 871-888 (2016).

88. C. Dogra, D. S. Srivastava, A. Kumar, Protein-DNA array-based identification of transcription factor activities differentially regulated in skeletal muscle of normal and dystrophin-deficient mdx mice. *Mol Cell Biochem* **312**, 17-24 (2008).

89. M. Prencipe *et al.*, Relationship between serum response factor and androgen receptor in prostate cancer. *Prostate* **75**, 1704-1717 (2015).

90. K. H. Lee, S. Evans, T. Y. Ruan, A. B. Lassar, SMAD-mediated modulation of YY1 activity regulates the BMP response and cardiac-specific expression of a GATA4/5/6-dependent chick Nkx2.5 enhancer. *Development* **131**, 4709-4723 (2004).

91. S. Gregoire *et al.*, Essential and unexpected role of Yin Yang 1 to promote mesodermal cardiac differentiation. *Circ Res* **112**, 900-910 (2013).

92. D. Su, L. J. Gudas, Retinoic acid receptor gamma activates receptor tyrosine kinase Tie1 gene transcription through transcription factor GATA4 in F9 stem cells. *Exp Hematol* **36**, 624-641 (2008).

93. C. A. Rubel, H. L. Franco, J. W. Jeong, J. P. Lydon, F. J. DeMayo, GATA2 is expressed at critical times in the mouse uterus during pregnancy. *Gene Expr Patterns* **12**, 196-203 (2012).

94. A. L. Façanha, M. C. dos Reis, M. Montero-Lomeli, Structural study of the porcine Na+/H+ exchanger NHE1 gene and its 5'-flanking region. *Mol Cell Biochem* **210**, 91-99 (2000).

95. N. R. Sundaresan *et al.*, Sirt3 blocks the cardiac hypertrophic response by augmenting Foxo3a-dependent antioxidant defense mechanisms in mice. *J Clin Invest* **119**, 2758-2771 (2009).

96. J. D. Kormish, D. Sinner, A. M. Zorn, Interactions between SOX factors and Wnt/beta-catenin signaling in development and disease. *Dev Dyn* **239**, 56-68 (2010).

97. M. S. Harkins, P. L. Moseley, G. K. Iwamoto, Regulation of CD23 in the chronic inflammatory response in asthma: a role for interferon-gamma and heat shock protein 70 in the TH2 environment. *Ann Allergy Asthma Immunol* **91**, 567-574 (2003).

98. S. I. Lee *et al.*, The miR-302 cluster transcriptionally regulated by POUV, SOX and STAT5B controls the undifferentiated state through the post-transcriptional repression of PBX3 and E2F7 in early chick development. *Mol Reprod Dev* **81**, 1103-1114 (2014).

99.   M. C. Peeters *et al.*, The adhesion G protein-coupled receptor G2 (ADGRG2/GPR64) constitutively activates SRE and NFκB and is involved in cell adhesion and migration. *Cell Signal* **27**, 2579-2588 (2015).

100.   M. Stadler *et al.*, Transcriptional induction of the PML growth suppressor gene by interferons is mediated through an ISRE and a GAS element. *Oncogene* **11**, 2565-2573 (1995).

101.   J. Nolting *et al.*, Retinoic acid can enhance conversion of naive into regulatory T cells independently of secreted cytokines. *J Exp Med* **206**, 2131-2139 (2009).

102.   A. He, S. W. Kong, Q. Ma, W. T. Pu, Co-occupancy by multiple cardiac transcription factors identifies transcriptional enhancers active in heart. *Proc Natl Acad Sci U S A* **108**, 5632-5637 (2011).

103.   M. Urbini *et al.*, Identification of SRF-E2F1 fusion transcript in EWSR-negative myoepithelioma of the soft tissue. *Oncotarget* **8**, 60036-60045 (2017).

Figure Legends

Figure 1: *Workflow of gene network reconstruction*

Figure 2. *Gene profile clustering and subnetwork reconstruction by randomized connectivity.* (a) Clustering. The blue curved profiles are clustered to each red profile as a unique feature based on correlation and occurrences of convexity. Each dataset from 6 conditions of external cues has different clustering profiles. (b) Reconstruction of subnetworks. From each cluster profile, a subnetwork is reconstructed with randomized bootstrapping by GPU-ODE integration and 100 times fitting within the error threshold. For the time course profiles averaged among the clustered genes from experiments (red), best five and worst five fitted profiles generated by the kinetic model on random networks (black) are shown together.

Figure 3. *Global gene network integration by scoring from multiple cues.* (a) Gene-to-gene fidelity is estimated from gene-to-gene binding frequency-based score combining six scores as one. (b) For those gene-to-gene edges that scored more than the threshold of overall $h=0.24$, the gene-to-gene interactions of the activation (blue) and inhibition (red) are prescribed for the global network. (c) The gene regulatory network with the overall h-index threshold$=0.26$. (d) The gene regulatory network with the overall h-index threshold$=0.28$.

Figure 4. *Reconstruction of p-value based gene regulatory subnetworks.* The subnetwork reconstruction process generates unbiased random gene binding events of activation/inhibition for each cluster in each condition. The binding occurrences are collected as distribution (a and c), and the clusters are identified as significant for $p < 0.05$ (b and d) and colored as violet. The insignificant clusters are colored as light violet.

Figure 5. *Global optimization for gene regulatory network integration.* The subnetworks are integrated by a combination of stochastic global optimization and gradient-based programming (GCMMA) while estimating kinetic parameters from 6 different cues with the gene-to-gene interacting connectivity fixed (Figure 3c). However, the cue-to-gene connectivities are different for each of 8 replicates, and their "overall" connectivities are determined by their statistical significance. (a) Reconstructed gene profiles with 3 RGD peptide levels (low: blue, middle: green, high: red) and a medium level of hydrogel stiffness (b) Reconstructed gene profiles with 3 PEG levels for different hydrogel stiffnesses (soft: blue, medium: green, hard: red) and RGD level in the middle. For (a) and (b), the solid curves are from reconstructed kinetic models, and five circles are from TRACER data (3, 6, 9, 12, 27 hours).

Figure 6. *Identified gene network pathways are validated from a priori known gene databases.* By overlapping *p*-value based gene-to-gene connectivities commonly significant from two conditions, *differential* networks are constructed. The significant interactions from dual cues to targeted genes are shown by green colored lines (activation → or inhibition --|). They assume to exist constitutively through 6 different conditions. The inferred four networks are validated with *a priori* known gene databases and identified as "*relevant*" (solid) when there exist specific gene-to-gene interactions even though the cell line and conditions may not be the same as those of the TRACER experiments (Table). The interactions not validated are represented by dotted lines (This does mean that they are predicted, but they need experimental validations). (a) Overlapping (RGD-high and stiffness-medium) with (RGD-middle, stiffness-medium). We have inferred that 19 gene-to-gene interactions are relevant among 26 predicted edges. (b) Overlapping (RGD-middle and stiffness-medium) with (RGD-low, stiffness-medium). Among 44 gene-to-gene interactions, 18 are relevant. (c) Overlapping (RGD-middle and

stiffness-hard) with (RGD-middle and stiffness-medium). Among 33 gene-to-gene interactions, 18 are relevant. (d) Overlapping (RGD-mid and stiffness-medium) with (RGD-middle and stiffness-soft). Among 56 gene-to-gene interactions, 19 are relevant.

Figure 7. *Master regulatory transcriptional factors GATA4, SMAD3/4, ETS-1, and STAT5 and associated pathways respond adaptively with differential conditions from mechano-transduction.* Master regulatory genes are identified from validated pathways in Figure 6. The dimmed interactions are not significant in the specific conditions. (a) To indicate the adaptive gene regulations with *GATA4*, the interactions with soft hydrogel stiffness and RGD-middle, those with hard hydrogel stiffness and RGD-middle, those with high RGD and medium hydrogel stiffness are shown in sequence. *GATA4* is counter-stimulated by RGD and hydrogel stiffness, i.e., effectively activated by RGD and inhibited by hydrogel stiffness. Activation of *SMAD3/4* from *YY1* and *RAR* are turned on in soft hydrogel. When RGD is high, *GATA4* inhibits *FOXO3A*. (b) To indicate the adaptive gene regulations with *ETS1*, the interactions with low RGD and medium hydrogel stiffness, and those with high RGD and medium hydrogel stiffness are shown in sequence. *ETS1* is counter-stimulated by RGD and hydrogel stiffness. RGD – low versus high. *ETS1* is inhibited from *STAT3* and *AP1* mediated through *RUNX2* when RGD is low, but activated from medium stiffness mediated by *PEA3* when RGD is high. (c) To indicate the adaptive gene regulations with *STAT5*, the interactions with soft hydrogel stiffness and RGD-middle, and those with low RGD and medium hydrogel stiffness are shown in sequence. *STAT5* is co-stimulated by stiffness and RGD. (RGD-low, stiffness-medium) versus (RGD-mid, stiffness-soft). STAT5 is activated by *ELK1* when RGD is low, but this activator is switched over to *SOX* triggered by the middle level of RGD and soft hydrogel.

Figure 1

(I) CLUSTERING

(II) SUBNETWORK RECONSTRUCTION

(III) GENE-TO-GENE CONNECTIVITY

(V) NETWORK VALIDATION

(IV) SUBNETWORK INTEGRATION

Figure 2

Figure 3

Figure 4

Figure 5

(a)

(b)

Figure 6

(a) RGD – HIGH STIFFNESS – MEDIUM

(b) RGD – LOW STIFFNESS – MEDIUM

(c) RGD – MID STIFFNESS – HARD

(d) RGD – MID STIFFNESS – SOFT

Figure 7

Table 1: Gene-to-gene interaction validation

Direct association common from all six external conditions		
STIFFNESS → *STAT1*	*(60)*	Hypertrophy and heart failure
STIFFNESS → *STAT5*	*(61)*	ER alpha+ breast cancer
STIFFNESS --| *RAR*	*(62)*	Hepatic stellate cell
STIFFNESS --| *PR*	*(63)*	Follicle cells
STIFFNESS → *PEA3*	*(64, 65)*	Fibrocartilaginous cells, fibroblast
RGD → *SOX*	*(66, 67)*	Murine embryonic stem cell
RGD → *GATA*	*(68)*	Rat and human
RGD --| *FOXO3A*	*(66, 69)*	Human lung fibroblast
RGD --| *ETS1*	*(70)*	Human vascular smooth muscle
RGD – HIGH & MIDDLE, STIFFNESS – MEDIUM: Differential network I		
PEA3 → ETS1	*(71)*	Lung type 1 epithelial/endothelial
AR → ETS1	*(72)*	Prostate cancer
SRE → GATA	*(73)*	fibroblast
SOX --| SRF	*(74)*	Mouse embryonic fibroblast
AP1 → IRF1	*(75)*	Microvascular endothelium
RAR → p53	*(76, 77)*	Colorectal cancer, ER- breast cancer
P53 → RUNX1	*(78)*	Lung cancer
RUNX1 → SP1	*(79)*	Acute myeloid leukemia
GATA3 → HSTF	*(80)*	Placental fibroblast
GATA4 --| FOXO3A	*(81)*	Cardiac fibroblast
RGD – MIDDLE & LOW, STIFFNESS – MEDIUM: Differential network II		
AP1 → STAT3	*(82)*	Human airway epithelium
AP1 → IRF1	*(75)*	Microvascular endothelium
STAT3 --| ETS1	*(83)*	macrophage
RUNX2 --| ETS1	*(84)*	Mouse osteoblast precursor cells
AP4 → RUNX1	*(85)*	T-cell
RUNX1 → SP1	*(79)*	Acute myeloid leukemia
ELK1 → STAT5	*(86)*	Hematopoietic cells
SRF → NFkB	*(87)*	Ovarian cancer cells
SOX --| SRF	*(74)*	Mouse embryonic fibroblast
STIFFNESS – HARD & MED, RGD - MIDDLE: Differential network III		
STAT4 → AP4	*(88)*	Skeletal muscle
SRF → AR	*(89)*	Prostate cancer
YY1 → GATA4	*(90, 91)*	Cardiac crescent, cardiac progenitor
RAR → GATA4	*(92)*	Teratocarcinoma stem cells
GATA2 → PR	*(93)*	Mouse uterus
AP3 → PEA3	*(94)*	epithelium
FOXO3A --| GATA4	*(95)*	Cardiac myocyte
TCF/LEF → SOX	*(96)*	Sea urchin embryos
HSE → ISRE	*(97)*	Alveolar macrophage

STIFFNESS – MED & SOFT, RGD - MIDDLE: Differential network IV		
FOXO3A --| GATA4	*(95)*	Cardiac myocyte
SOX → STAT5	*(98)*	Chick blastodermal cell
SRE → NFkB	*(99)*	Breast cancer
RAR → GATA4	*(92)*	Teratocarcinoma stem cells
RAR → GAS	*(100)*	Acute promyelocytic leukemia
YY1 → SMAD3/4	*(50)*	Mouse myoblast
YY1 → GATA4	*(90, 91)*	Cardiac crescent, cardiac progenitor
RAR → SMAD3	*(101)*	Naïve and regulatory T cell
SRF → GATA	*(102)*	Cardiac muscle (HL-1)
SRF → E2F1	*(103)*	Myoepithelioma