Massively parallel transposon mutagenesis identifies temporally essential genes for biofilm

formation in *Escherichia coli*

Authors: Emma R Holden[1], Muhammad Yasir[1], A Keith Turner[1], John Wain[1,2], Ian G. Charles[1,2],

Mark A Webber[1,2] *

[1] Quadram Institute Bioscience, Norwich Research Park, Norwich, Norfolk, NR4 7UQ, U.K.

[2] Norwich Medical School, University of East Anglia, Norwich Research Park, Norwich, Norfolk,

NR4 7TJ, U.K.

* Corresponding author Mark.Webber@quadram.ac.uk

Keywords: Functional genomics, transposon mutagenesis, adhesion, TraDIS

Running title: *E. coli* TraDIS biofilm

Abstract

Biofilms complete a life cycle where cells aggregate, grow and produce a structured community before dispersing to seed biofilms in new environments. Progression through this life cycle requires controlled temporal gene expression to maximise fitness at each stage. Previous studies have focused on the essential genome for the formation of a mature biofilm, but here we present an insight into the genes involved at different stages of biofilm formation. We used TraDIS-Xpress; a massively parallel transposon mutagenesis approach using transposon-located promoters to assay expression of all genes in the genome. We determined how gene essentiality and expression affects the fitness of *E. coli* growing as a biofilm on glass beads after 12, 24 and 48 hours. A selection of genes identified as important were then validated independently by assaying biofilm biomass, aggregation, curli biosynthesis and adhesion ability of defined mutants. We identified 48 genes that affected biofilm fitness including genes with known roles and those not previously implicated in biofilm formation. Regulation of type 1 fimbriae and motility were important at all time points. Adhesion and motility were important for the early biofilm, whereas matrix production and purine biosynthesis were only important as the biofilm matured. We found strong temporal contributions to biofilm fitness for some genes including some which were both beneficial and detrimental depending on the stage at which they are expressed, including *dksA* and *dsbA*. Novel genes implicated in biofilm formation included *ychM* and *truA* involved in cell division, *crfC* and *maoP* in DNA housekeeping and *yigZ* and *ykgJ* of unknown function. This work provides new insights into the requirements for successful biofilm formation through the biofilm life cycle and demonstrates the importance of understanding expression and fitness through time.

Introduction

Bacteria rarely exist planktonically outside of the laboratory and are usually found as part of structured, aggregated communities called biofilms [1]. Clinically, approximately 80% of infections have been suggested to have a biofilm component [2]. Biofilm related infections are complicated by their intrinsic tolerance to antimicrobials, making infections difficult to treat and, often persistent [3-6]. Cells within a biofilm grow more slowly than those in planktonic culture and this reduced level of metabolic activity has been associated with tolerance to antimicrobials, allowing biofilms to be typically 10-1000-fold less sensitive to antibiotics than corresponding strains in planktonic conditions [7,8]. Aside from the clinical setting, there are many useful applications of biofilms, including wastewater treatment and bioprocessing [9]. Biofilms undergo a life cycle that commonly consists of initial attachment to a surface, growth and maturation of the biofilm over time with characteristic production of extracellular matrix components, followed by dispersal of planktonic cells to facilitate colonisation of new surfaces [10]. The switch between planktonic and biofilm lifestyles is driven by environmental stimuli promoting large scale changes in gene expression and regulation that are necessary to support the bacterial community through the life cycle, which is distinct to planktonic growth conditions. Expressing the right genes at the right time and place is critical for efficient production of a biofilm.

The main components of the biofilm extracellular matrix in *E. coli* are the amyloid protein curli, the polysaccharide cellulose and extracellular DNA [11]. Curli is transcribed by the divergent operons *csgBAC* and *csgDEFG*, with curli biosynthesis regulated by CsgD [12]. Cellulose biosynthetic machinery is encoded by *bcsRQABZC* and *bcsEFG*, and its production is regulated by c-di-GMP [13]. Several genes are known to be involved in the regulation of matrix production, including *ompR* [14,15], *cpxR* [14,16,17] and *rpoS* [18,19], amongst others [20-22]. Extracellular DNA is also an important component of the biofilm matrix, and the addition of DNase has negatively affected the biomass of biofilms formed by *Pseudomonas aeruginosa* [23], *Bacillus cereus* [24] and a range of Gram-negative pathogens, including *E. coli* [25].

Most previous studies focusing on identifying the genes and pathways required for biofilm

formation in *E. coli* have concentrated on the mature biofilm rather than dissecting events across

the life cycle. Various studies have taken a genome wide approach to identifying genes that impact

biofilm formation. One assessed biofilm formation of all the mutants in the Keio collection [26],

although analysis is limited to the effect of inactivation [27]. Another used a transcriptomic approach

to identify genes with altered expression in biofilms compared to in suspension over time [28]. DNA

microarrays have also been used to link the presence of different genes with biofilm capacity in

panels of isolates [29].

Large scale transposon mutagenesis represent another high-throughput, sensitive whole genome

approach to link phenotype to genotype methodologies [30-32]. These methods have the advantage

of assessing the impact of a very large number of random insertions within and without genes on a

phenotype. They have however been historically limited by an inability to assay essential genes

where transposon insertions are not viable. We sought to investigate biofilm formation using

TraDIS-Xpress, a method developed recently which allows essential genes to be assayed for their

role in survival in test conditions. In this method, gene expression changes caused by an outward-

facing inducible promoter incorporated into the transposon are captured as well as traditional

essentiality measurements [33].

This study identified 48 genes that were found to be important at different stages of biofilm

formation by *E. coli*. By investigating the genes important across the biofilm life cycle, we were

able to get a dynamic view of the main pathways important at different stages of biofilm

development. Genes identified by TraDIS-Xpress as important at each stage were then

phenotypically validated using defined mutants. Our findings reinforced the importance of

adhesion, motility and matrix production in the biofilm, and revealed the importance of genes not

previously implicated in biofilm formation. This included genes with roles in cell division (*ychM* and

truA), DNA housekeeping (*crfC* and *maoP*) and *yigZ* and *ykgJ*, the functions of which have not

been elucidated. We identified clear requirements for some pathways at specific points of the

biofilm life cycle, furthering our understanding of how biofilms maintain fitness over time.

Results

Analysis of the TraDIS-Xpress data found 48 genes that considerably affected biofilm formation over time in *E. coli*: 42 were identified as being needed to maintain the fitness of a biofilm and expression of 6 genes was predicted to result in reduced biofilm fitness (Figure 1 and Supplementary table 1). The main pathways that were consistently important in the biofilm through all the time points included type 1 fimbriae, curli biosynthesis and regulation of flagella (Figure 1). All other loci identified affected biofilm formation at specific points in the life cycle.

Fimbriae expression and motility are important at all stages of biofilm formation

Only 4 genes were found to be important throughout 12, 24 and 48 hours (figure 2a). These included *fimB* and *fimE*. The recombinase gene *fimB* which helps mediate both 'ON-to-OFF' and 'OFF-to-ON' switching of fimbriae expression was beneficial for biofilm formation at all time points. Fewer *fimB* mutants were observed in biofilm conditions compared to planktonic, and this number decreased over time. In contrast, inactivation of *fimE*, responsible for only 'ON-to-OFF' fimbrial regulation [34], increased biofilm fitness at all time points. Initially, there were only slightly more *fimE* mutants in biofilm conditions compared to planktonic at 12 hours, but this increased over time with a stark contrast seen between biofilm and planktonic conditions at the 24 and 48 hour time points. The predicted impacts on biofilm formation were phenotypically validated by testing both gene knockout mutants from the Keio collection (which contains two, independent deletion mutants for most genes in *E. coli* BW25113) [35]. Analysis of these mutants showed for both that there was no deficit in biofilm biomass (Figure 3a), but both were deficient in cell aggregation (Figure 3b). Together, the TraDIS-*Xpress* and phenotypic data suggest that the ability to regulate fimbriae expression in a phase-dependent manner is important for fitness of a biofilm, rather than being constrained in a fimbriae 'ON' or 'OFF' state.

Disruption of *lrhA*, a regulator of motility and chemotaxis [36], was beneficial for biofilm formation at all time points (Figure 2b). LrhA also has a role in type 1 fimbriae expression through activating expression of *fimE* [37], but in addition represses flagella-mediated motility. Analysis of the Δ*lrhA*

biofilm showed initial formation of microcolonies occurred faster than the wild-type (Figure 4) but at

later time points the biofilms formed by this mutant were less mature than seen with the wild-type.

There was no significant change in biomass formed by this mutant (Figure 3a) and mutants

appeared less aggregative than the wild-type (Figure 3b). These data suggest that inactivation of

lrhA impacts both adhesion and aggregation differently at distinct stages of the biofilm life cycle

and may result in a benefit to early surface colonisation but with a cost to later maturation.

Expression of the Hha toxin attenuator *tomB* was also found to be consistently important for biofilm

formation at 12, 24 and 48 hours (Figure 2b). Consistent with this prediction, the Δ*tomB* mutant

biofilm had reduced cell aggregation and curli biosynthesis, and reduced biofilm biomass (Figure 3

a,b,c).

<u>Regulatory genes are important in the early biofilm</u>

In the early biofilm, at 12 hours growth, only 13 genes were found to distinguish the planktonic and

biofilm conditions. TraDIS-*Xpress* data indicated that inactivation of transcriptional factor *dksA*

promoted biofilm formation at the 12- and 24-hour time points but not in the mature biofilm (Figure

2c). Supporting this, analysis of Δ*dksA* mutant biofilms under flow conditions showed an initial

benefit with increased adhesion at both the 12 and 24 hour time points, but reduced microcolony

formation at the 48 hour time point, suggesting *dksA* affects biofilm initiation (Figure 4). Inactivation

of Δ*dksA* was also seen to reduce cell aggregation, curli biosynthesis and biofilm biomass (Figure

3 a,b,c). Expression of *hdfR*, a negative regulator of motility [38], was found to be detrimental to

biofilm fitness in the early biofilm after 12 and 24 hours growth (Figure 2b), and Δ*hdfR* mutant

biofilms had significantly reduced biomass (Figure 3a). In addition, the stress response regulator

marR [39] and the 23S rRNA methyltransferase *rlmI* [40] were both found to be beneficial for biofilm

fitness at the 12 hour time point only, and reduced biofilm biomass was found in deletion mutants

(figure 3a). These genes have both previously been implicated in biofilm formation [40-42], but the

effect on early biofilm formation has not been described previously.

<u>Biofilms sampled after 24 hours demonstrate both adhesion and matrix production are important</u>

More pathways were identified as being important to biofilm formation at 24 hours that at 12 hours.

Two genes involved in DNA housekeeping were found to be involved in biofilm formation at this

time point. These included *dam*, encoding DNA methyltransferase [43], insertional activation of which

was not tolerated in the 24 hour biofilm, and Δ*dam* mutants were defective in aggregation

compared to the wild type (figure 3b). Also, inactivation of *maoP*, involved in Ori macrodomain

organisation [44], was predicted to confer a fitness advantage in the 24 hour biofilm compared to the

planktonic condition. TraDIS-*Xpress* data showed more reads mapped to *maoP* in the biofilm

conditions compared to the planktonic at 24 hours suggesting loss of this gene was beneficial.

Phenotypic analysis of the Δ*maoP* mutant biofilm did demonstrate a phenotype although in

opposition to the prediction, *maoP* mutants were significantly deficient in biofilm biomass

production, curli biosynthesis and one mutant displayed reduced aggregation (Figure 3 a,c).

In the 12 and 24 hour biofilms, *dsbA* (encoding disulphide oxidoreductase) was essential with no

insertions detected within this gene (Figure 2c). The role of *dsbA* in adhesion to abiotic surfaces

and epithelial cells has previously been suggested (Lee et al., 2008, Bringer et al., 2007).

Phenotypic validation of the Δ*dsbA* mutant showed a red, dry and rough (*rdar*) phenotype on

Congo red plates (Figure 3c), indicative of increased curli biosynthesis. Cell aggregation in the

Δ*dsbA* mutant was significantly improved compared to the wild type, implying a role of *dsbA* in

inhibiting cell-cell aggregation. Our data showed that *dsbA* is important in the early biofilm, but its

deletion appears to be beneficial to the formation of a mature biofilm, according to the Congo red

and aggregation data. Supporting this hypothesis, *dsbA* was not essential at the 48 hour time

point.

Inactivation of the RNase III modulator *ymdB* [45] was found to reduce fitness in the biofilm, with

fewer reads mapping here in biofilm conditions compared to planktonic at both the 24 and 48 hour

time points. This follows previous findings that both inactivation and overexpression of *ymdB*

negatively affects biofilm biomass [45]. In concordance with these findings, a Δ*ymdB* mutant had

significantly improved cell aggregation and reduced biofilm biomass (figure 3 a,b).

Curli biosynthesis became important by the 24 hour time point as no insertions mapped to *csgC*,

encoding a curli subunit chaperone [46] and more transposon insertions mapped upstream of the

curli biosynthesis regulator *csgD* [12], likely indicating its increased expression benefitted biofilm

formation. At the 48 hour time point, both genes were essential for biofilm formation, which was

also the case for the known *csgD* regulator, *ompR* [14], supported by significantly reduced biofilm

biomass and reduced aggregation in knockout mutants (figure 3a,b).

The mature biofilm grown for 48 hours requires purine biosynthesis, matrix production, motility and

solute transport

There were 38 genes found to be important for fitness of the mature biofilm after 48 hours growth,

and 25 of these genes were identified as essential at this time point only. The major pathway

implicated in biofilm formation at 48 hours was purine ribonucleotide biosynthesis, with four genes,

purD, *purH*, *purL* and *purE* [47], found to be essential at this time point only. TraDIS-Xpress did not

identify mutants in any of these genes in biofilms sampled at 48 hours, whereas several reads

mapped to these loci under planktonic conditions, as well as under both biofilm and planktonic

conditions earlier at 12 and 24 hours. Visualisation of a Δ*purD* mutant biofilm under flow conditions

saw poor biofilm formation and no microcolony formation at any time compared to the wild type

(Figure 4). Additionally, Δ*purD* and Δ*purE* mutants were deficient in biofilm biomass production,

curli biosynthesis, and Δ*purE* also showed reduced cell aggregation (Figure 3 a,b,c), confirming an

important role for purine biosynthesis in matrix production and curli biosynthesis in the mature

biofilm.

The flagella master regulatory system *flhDC* was identified as important in the mature biofilm.

Biofilms sampled after 48 hours saw fewer *flhC* mutants, while insertions interpreted as over-

expressing *flhD* increased in numbers both at the 24 and 48 hour time points, compared to

planktonic conditions. No mutants in *flgD* and *fliE*, encoding flagellar filament proteins, were

identified at 24 and 48 hours, respectively. It has previously been shown that motility is important

for initial biofilm formation [48,49], but this may not relate to biomass formation where no differences

were seen for Δ*flhD*, Δ*flhC*, Δ*fliE* and Δ*flgD* mutants.

Various pleiotropic transcriptional regulators were also important in the mature biofilm. This

included the H-NS antagonist *leuO* [50]. Increased insertions upstream of *leuO* under biofilm

conditions after 12 hours growth, as well as no *leuO* mutants in 48 hour biofilms, indicated it was

beneficial to biofilm formation. A Δ*leuO* mutant did not aggregate as well as the wild type, and one

Δ*leuO* mutant had reduced biofilm biomass (figure 3 a,b). The Δ*leuO* mutant biofilm under flow

conditions demonstrated an inability to form microcolonies after 48 hours growth (figure 4). The

leucine-responsive global regulator *lrp* [51] and a transcriptional regulator responsible for survival

under acid stress, *gadW* [52] were also found to have fewer mutants in the 48 hour biofilm compared

to the planktonic condition, indicating their importance in the mature biofilm. Reduced biofilm

biomass, aggregation and curli biosynthesis were observed for one copy of Δ*lrp*, but no differences

in biofilm formation or aggregation were seen for Δ*gadW* mutant biofilms (figure 3 a,b,c).

Inactivation of the outer membrane channels *mscL* [53], *tolC* [54] and *ompF* [55] was not tolerated in the

mature biofilm grown for 48 hours. This would indicate the importance of transport in the mature

biofilm, however inactivation of *tolC* and *ompF* did not result in a change in biofilm biomass (figure

3a).

Three genes involved in cell division or DNA replication, *crfC*, *ychM* and *truA*, were identified as

important in the 48 hour biofilm. Expression of *yhcM*, involved in cell division [56] was predicted to be

essential in the 48 hour biofilm, but this was not reflected in the phenotype of deletion mutants

(Figure 3 a). The pseudouridine synthase *truA* [57], also reported to be involved in cell division [58] was

found to be essential in the mature biofilm grown for 24 and 48 hours. Although a Δ*truA* mutant

showed no change in biofilm biomass or curli biosynthesis, aggregation was significantly improved

compared to the wild type albeit to a small degree (figure 3b), contrary to the prediction generated

by TraDIS-*Xpress*.

Discussion

We have characterised the essential genome of *E. coli* biofilms across the biofilm lifecycle by using the high throughput transposon mutagenesis screen TraDIS-*Xpress*. The identification of genes and pathways already described to be involved in biofilm formation validates the efficacy of this experimental model. The early biofilm established 12 hours after inoculation was characterised by genes involved in adhesion. The 24-hour biofilm required both adhesion and matrix production, transitioning into matrix production being of the upmost importance in the mature biofilm after 48 hours. In concordance with previous work identifying genes whose importance varies with time in the *E. coli* biofilm, we also reported that control of fimbriae expression and motility remained important at each stage of the biofilm life cycle rather than just being involved in initial attachment [28].

As well as identifying known pathways, TraDIS-*Xpress* was also able to identify genes not previously known to be involved in biofilm formation, including *yigZ*, *ykgJ*, *ychM*, *maoP*, *truA*, *crfC*. We found that expression of *maoP* was detrimental to the fitness of biofilms grown for 24 hours, but a Δ*maoP* mutant biofilm had reduced biofilm biomass and reduced curli biosynthesis compared to the wild type. Chromosomal organisation of the Ori macrodomain requires both *maoP* and *maoS* [44], however no signal is seen in our data for *maoS*. This warrants further investigation into how chromosomal macrodomain organisation affects biofilm formation. The importance of cell division in the mature biofilm was further supported by our findings of fewer *ychM* and *truA* mutants compared to planktonic conditions, both of which have not before been implicated in biofilm formation. However, deletion of either gene had no effect on biofilm biomass or curli production, and the Δ*truA* deletion mutant had improved aggregation compared to the wild type. Genes involved in cell division are clearly important for the fitness of the mature biofilm, but the essential roles of *ychM* and *truA* in this process are currently unclear but unlikely to relate to crude biomass production.

Expression of *dsbA* and repression of *dksA* was found in this study to benefit early biofilm fitness.

Based on previous studies and phenotypic analysis of knockout mutants in this study, we believe

the increase in biofilm fitness seen is due to increased adhesion in these mutants [59,60]. This study

has highlighted the benefit of close temporal gene regulation in the biofilm, where the expression

of certain genes can have a different effect on biofilm fitness at different stages of the biofilm life

cycle. For example, we found that *dsbA* was important for the early biofilm, but a *dsbA* mutant

biofilm had increased curli expression and increased aggregation. Expression of *dsbA* has been

previously found to result in repression of the curli regulator *csgD* and curli subunit *csgA*, essential

for optimal fitness of the mature biofilm [61]. Conversely, we found that expression of the

transcription factor *dksA* was detrimental in the early biofilm, whilst a *dksA* knockout biofilm had

reduced biofilm biomass, reduced curli biosynthesis and reduced aggregation. The relationship

between *dksA* expression and curli biosynthesis has been previously characterised in similar

studies [32,62]. Again, these data show differential expression of important genes at different stages

of the biofilm life cycle is essential for optimising biofilm fitness.

Purine biosynthesis was found to be important in the mature biofilm, through the essentiality of

purD, *purE*, *purL* and *purH* in biofilms grown for 48 hours. Similar findings have previously been

described in another transposon mutagenesis experiment in uropathogenic *E. coli* [32]. Inactivation

of purine biosynthetic genes was also found to impair biofilm formation in *Bacillus cerus*, but this

was thought to be due to reduced extracellular DNA in the biofilm matrix [24]. Extracellular DNA is

thought to aid adhesion and has been found to be important in the biofilms of a wide range of

bacterial species [23,25]. Our data suggest the importance is in the mature biofilm rather than initial

adhesion. A relationship between both purine and pyrimidine biosynthesis and curli production in

the biofilm has been reported [32,62,63]. More recently, curli biosynthesis in a *purL* mutant was

reported to be abrogated through addition of inosine, which is involved in the *de novo* purine

biosynthetic pathway for production of adenosine monophosphate (AMP) and guanine

monophosphate (GMP) [64]. This suggests that nucleotide production itself, rather than the

regulatory effects of the genes involved, affects curli biosynthesis, supporting one hypothesis that

disruption of the purine biosynthetic pathway may directly result in a reduction of cyclic-di-GMP,

known to regulate biofilm formation and affect curli biosynthesis [32]. Quantification of intracellular c-

di-GMP or further investigation of other c-di-GMP-dependent pathways in these mutants would

uncover the relationship between these pathways and biofilm formation.

TraDIS-*Xpress* data suggested that expression of *fimB* and deletion of *fimE* was necessary for

optimal biofilm fitness at all points in the biofilm, rather than just for initial attachment. Analysis of

Δ*fimB* and Δ*fimE* deletion mutants found no significant change in biofilm biomass and reduced

cell-cell aggregation. This finding supports previous work that reported increased fimbriae

expression across all stages of biofilm development [28]. Previous work has observed a positive

correlation between type 1 fimbriae and exopolysaccharide production in the mature biofilm [65], but

the increase in biofilm biomass to support this was not seen in our study. These data suggest that

for a population the ability to present cells both with and without fimbriae is beneficial for fitness

throughout the life cycle.

Analysis of biofilms under flow conditions found that Δ*lrhA* and Δ*tomB* mutant biofilms had a

similar appearance after 12 hours growth, could potentially indicate a similar role in the biofilm. The

role of *lrhA* in motility regulation has been well documented [36,37,66], and expression of *tomB* has

been seen to reduce motility through repression of *fliA* [67]. We found deletion of eiher *lrhA* or *tomB*

resulted in reduced aggregation. Although Δ*lrhA* and Δ*tomB* deletion mutants shared many similar

phenotypes, TraDIS-*Xpress* data predicted that *tomB* was beneficial and *lrhA* was detrimental to

biofilm formation at 12, 24 and 48 hours. Previous studies on Δ*lrhA* mutant biofilms have reported

increased adhesion, aggregation and biomass compared to the wild type [37]. This was not seen in

our study, but we did find aggregation of the Δ*lrhA* mutant changed over time, with increased

aggregates seen in biofilms grown in flow cells after 12 hours (figure 4). However, decreased

aggregation was observed in planktonic cultures after 24 hours (figure 3). The initial benefit to

biofilm formation resulting from the inactivation of *lrhA* predicted by the TraDIS-*Xpress* data agrees

with the initial phenotype of the Δ*lrhA* biofilm in the flow cells. This may be due to reduced

induction of *fimE* by LrhA [37], thereby allowing expression of type 1 fimbriae to facilitate adhesion.

We have already described how expression of both *fimB* and *fimE* is necessary for optimal fitness

of the mature biofilm, and the effect of *lrhA* on biofilm formation correlates with these findings, with

reduced aggregation in Δ*lrhA* biofilms after 24 hours (also seen in *fimB* and *fimE* mutants) and no

microcolony formation under flow conditions at 24 and 48 hours. Therefore, the importance of *lrhA*

to biofilm formation clearly appears to be time dependent, with the most important role in early

events.

Studies on the effect of *tomB* on biofilm formation have focused on its toxin-antitoxin relationship

with *hha*, which has been found to reduce expression of fimbrial subunit *fimA* and activate

prophage lytic genes causing cell death [68]. Deletion of *hha* was found to reduce motility through

flhDC and increase curli production through *csgD* [69]. We found no obvious benefit to biofilm fitness

with insertional inactivation of *hha*, but this may not be visible in our data due to these mutants

having a functional copy of *tomB*. The role of *tomB* in the biofilm has not previously been

suggested, but we predict *tomB* is involved in regulation of adhesion in the early biofilm and matrix

biosynthesis in the late biofilm.

The relationship between motility and biofilm formation is complex. Although it is widely understood

that motility is crucial for biofilm formation [48,49], it is also true that motility and curli production have

an inverse relationship, where *csgD* directly represses *fliE* [70] and induces c-di-GMP synthesis

through *adrA*, which reduces motility through *ycgR* [71-73]. This is often referred to as a lifestyle

'switch', where biofilm matrix production represses motility for a motile-to-sessile lifestyle transition

[74]. We found that although insertional inactivation of negative motility regulators *lrhA* and *hdfR*

improved biofilm fitness according to the TraDIS-*Xpress* data, a Δ*hdfR* deletion mutant actually

had reduced biofilm biomass, and deletion of either *lrhA* or *hdfR* impaired cell aggregation. Our

data found an important role for structural flagella components in the mature biofilm. Previous work

has suggested that flagella filaments are important for initial attachment and adhesion [75], however

we did not find this to be the case, with the expression of flagella filaments only appearing to

increase biofilm fitness in the mature biofilm grown for 48 hours. This is supported by previous

work that found expression of flagella is important at all stages of the developing biofilm [28]. It

appears that regulation of flagella and motility, rather than their fixed expression or absence, is

important for optimal biofilm fitness.

Previous genome-wide screens on *E. coli* biofilms have identified some of the same genes as this

study [26,30,32]. The TraDIS-*Xpress* technology used here makes for a more a powerful analysis of

biofilm formation by predicting roles of changes in gene expression as well as essentiality and by

analysing important genes over time. Differences between this work and previous studies may

reflect biofilms being grown under different conditions on different surfaces, as these

environmental factors greatly affect the pattern of gene expression and gene essentiality in the

biofilm [76]. Analysis of more strains and species, grown on different abiotic and biotic surfaces

under a range of environmental conditions may provide a wider list of essential genes for biofilm

formation shared amongst a majority of human pathogens, as well as substrate-, condition- and

species-specific genes and pathways for specific industrial, clinical and drug-development

applications. As well as temporal changes in gene expression, spatial changes have been shown

to affect biofilm development [77]. Integration of the spatial component into this model, to assay how

gene expression throughout the biofilm over time affects biofilm fitness, would be the next logical

step in furthering our understanding of biofilm development.

This study had revealed important time-specific roles for known and novel genes in biofilm

formation. We reveal some pathways have a more important role in the mature biofilm than

previously appreciated and identify genes with time dependent conditional essentiality within the

biofilm. We also identify potential new candidate genes essential for biofilm formation, which could

be targeted for novel anti-biofilm therapies. Further work using high-density transposon mutant

libraries across time and in different conditions is likely to further our understanding of biofilm

biology.

Methods

Transposon mutant library

373
374
375 The *E. coli* BW25113 transposon mutant library containing over 800,000 distinct mutants that was

376 used in this study has recently described [33]. The transposon used to construct this library

377 incorporates an outward-transcribing IPTG-inducible promoter.

Biofilm model conditions

The library was used to inoculate parallel cultures of 5 mL LB broth (without salt) with

approximately 10^7 cells. Cultures were grown in 6-well plates containing 40 sterile 5 mm glass

beads per well (Sigma, 18406). Two replicates were set up, with or without 1 mM IPTG. Plates

were incubated at 30 ˚C with light shaking for 48 hours. After 12, 24, and 48 hours of incubation, 2

mL of planktonic sample was collected from each culture and 70 beads were taken to constitute

the biofilm sample. Beads were washed twice in sterile PBS and vortexed in tubes containing PBS

to resuspend cells from the biofilm. Both planktonic and biofilm samples were centrifuged at 2100 x

g to form pellets for DNA extraction.

TraDIS-Xpress *Sequencing*

DNA was extracted from pellets following the protocol described in Trampari, et al. [78]. A

customised sequencing library was prepared to identify transposon insertions. DNA was

tagmented using a MuSeek DNA fragment library preparation kit (ThermoFisher) and customised

Tn5-i5 and i7 primers were used in PCR for 28 cycles [33]. DNA fragments of 300-500 bp were size

selected using AMPure beads (Beckman Coulter) and nucleotide sequences were generated using

a NextSeq 500 and a NextSeq 500/550 High Output v2 kit (75 cycles) (Illumina).

Informatics

Fastq files were aligned to the *E. coli* BW25113 reference genome (CP009273) using the

BioTraDIS (version 1.4.3) software suite [79] using SMALT (version 0.7.6). Insertion frequencies

were determined per gene using tradis_gene_insert_sites within the BioTraDIS toolkit. Insertion

frequencies per gene for each replicate were plotted against each other to determine the

experimental error between replicates as well as differences in insertion frequency between control

and test conditions (supplementary figure 1). The tradis_comparison.R command (also part of the

BioTraDIS toolkit) was also used to determine significant differences ($p < 0.05$, after correction for

false discovery) in insertion frequencies per gene between control and test conditions. For all

candidate loci, plot files generated by BioTraDIS were also examined manually in Artemis (version

17.0.1) [80] to confirm the results from these two approaches, as well as to identify regions where

inserts were under differential selection but did not fall within coding regions of the genome.

Validation experiments

Knockout mutants for genes identified by TraDIS-Xpress data were sourced from the Keio

collection [35] and tested for their biofilm-forming abilities. Crystal violet assays, used to assess

biofilm biomass production, were undertaken by inoculating 10^4 of each mutant strain into 200 µL

LB broth without salt in a 96-well plate. After 48 hours incubation at 30 ˚C, the culture was

removed, wells were rinsed with water, and the residual biofilms were stained for 10 minutes with

200 µL 0.01% crystal violet. The plate was then rinsed with water to remove the stain and 200 µL

70% ethanol was added to the wells to solubilise the stained biofilm. The optical density (OD) was

measured using a FLUOstar Omega plate reader (BMG Labtech) at 590 nM. Cell aggregation was

measured by leaving overnight cultures (normalised to an $OD_{600\,nm}$ of 0.3) on an unagitated surface

at room temperature. After 24 hours, the supernatant of each culture was removed by pipetting,

diluted in PBS and measured in a plate reader at 600 nm. Biofilm matrix composition was

investigated through spotting 10 µL of each mutant (representing 10^5 CFU) on agar supplemented

with 40 µg/mL Congo red (Sigma, C6277) to examine curli production. Plates were incubated at

30˚C for 48 hours and photographed to compare mutant biofilm composition to the wild type.

Adhesion and biofilm architecture were investigated under flow conditions for selected mutants

using the Bioflux system. Flow cells were primed with LB broth without salt at 5 dyne/cm^2 and

seeded with approximately 10^7 cells. The plate was left at room temperature for 2.5 hours to allow

attachment, and subsequently incubated at 30 °C at a flow rate of 0.3 dyne/cm^2. After 12, 24 and

48 hours, biofilms were visualised with an inverted light microscope and representative images at

10x, 20x and 40x magnification were taken at three locations of the flow cell. Experiments were

performed in duplicate.

Data availability

Sequence data supporting the analysis in this study has been deposited in ArrayExpress.

Temporary accession number for review E-MTAB-9873.

Code availability statement

All software packages used are described in the methods. No bespoke code was used in this

study.

Acknowledgements

ERH was supported by a studentship funded by the Quadram Institute Bioscience. The authors

gratefully acknowledge the support of the Biotechnology and Biological Sciences Research

Council (BBSRC); AKT, MY, JW, IGC and MAW were supported by the BBSRC Institute Strategic

Programme Microbes in the Food Chain BB/R012504/1 and its constituent project

BBS/E/F/000PR10349.

Author contributions

ERH designed and performed experiments, analysed the data and wrote the paper. AKT and IGC

helped design experiments and wrote the paper. MY helped design experiments. JW heled

analyse data and wrote the paper. MAW designed the experiments, analysed the data and wrote

the paper.

Competing interests

The authors have no competing interests.

References

456 1 Berlanga, M. & Guerrero, R. Living together in biofilms: the microbial cell factory and its
457   biotechnological implications. *Microbial Cell Factories* **15**, 165, doi:10.1186/s12934-016-
458   0569-5 (2016).
459 2 Bjarnsholt, T. *et al.* Biofilm formation – What we can learn from recent developments. *J.*
460   *Intern. Med.* **0**, doi:10.1111/joim.12782 (2018).
461 3 Gbejuade, H. O., Lovering, A. M. & Webb, J. C. The role of microbial biofilms in prosthetic
462   joint infections. *Acta Orthop.* **86**, 147-158, doi:10.3109/17453674.2014.966290 (2015).
463 4 Davis, S. C., Martinez, L. & Kirsner, R. The diabetic foot: the importance of biofilms and
464   wound bed preparation. *Curr. Diab. Rep.* **6**, 439-445 (2006).
465 5 Vestby, L. K., Gronseth, T., Simm, R. & Nesse, L. L. Bacterial Biofilm and its Role in the
466   Pathogenesis of Disease. *Antibiotics (Basel, Switzerland)* **9**, doi:10.3390/antibiotics9020059
467   (2020).
468 6 Wang, H., Tay, M., Palmer, J. & Flint, S. Biofilm formation of *Yersinia enterocolitica* and its
469   persistence following treatment with different sanitation agents. *Food Control* **73**, 433-437,
470   doi:https://doi.org/10.1016/j.foodcont.2016.08.033 (2017).
471 7 Mah, T.-F. *et al.* A genetic basis for *Pseudomonas aeruginosa* biofilm antibiotic resistance.
472   *Nature* **426**, 306, doi:10.1038/nature02122
473 https://www.nature.com/articles/nature02122#supplementary-information (2003).
474 8 Hoyle, B. D. & Costerton, J. W. Bacterial resistance to antibiotics: the role of biofilms. *Prog.*
475   *Drug Res.* **37**, 91-105 (1991).
476 9 Flemming, H.-C. *et al.* Biofilms: an emergent form of bacterial life. *Nature Reviews*
477   *Microbiology* **14**, 563, doi:10.1038/nrmicro.2016.94 (2016).
478 10 Kostakioti, M., Hadjifrangiskou, M. & Hultgren, S. J. Bacterial biofilms: development,
479   dispersal, and therapeutic strategies in the dawn of the postantibiotic era. *Cold Spring*
480   *Harb. Perspect. Med.* **3**, a010306-a010306, doi:10.1101/cshperspect.a010306 (2013).
481 11 Flemming, H.-C. & Wingender, J. The biofilm matrix. *Nature Reviews Microbiology* **8**, 623-
482   633, doi:10.1038/nrmicro2415 (2010).
483 12 Barnhart, M. M. & Chapman, M. R. Curli Biogenesis and Function. *Annu. Rev. Microbiol.* **60**,
484   131-147, doi:10.1146/annurev.micro.60.080805.142106 (2006).
485 13 Serra, D. O. & Hengge, R. in *Extracellular Sugar-Based Biopolymers Matrices* (eds Ephraim
486   Cohen & Hans Merzendorfer) 355-392 (Springer International Publishing, 2019).
487 14 Jubelin, G. *et al.* CpxR/OmpR Interplay Regulates Curli Gene Expression in Response to
488   Osmolarity in *Escherichia coli*. *J. Bacteriol.* **187**, 2038-2049, doi:10.1128/jb.187.6.2038-
489   2049.2005 (2005).
490 15 Vidal, O. *et al.* Isolation of an *Escherichia coli* K-12 Mutant Strain Able To Form Biofilms on
491   Inert Surfaces: Involvement of a New *ompR* Allele That Increases Curli Expression. *J.*
492   *Bacteriol.* **180**, 2442-2449 (1998).
493 16 Dorel, C., Vidal, O., Prigent-Combaret, C., Vallet, I. & Lejeune, P. Involvement of the Cpx
494   signal transduction pathway of *E. coli* in biofilm formation. *FEMS Microbiol. Lett.* **178**, 169-
495   175, doi:10.1111/j.1574-6968.1999.tb13774.x (1999).
496 17 Otto, K. & Silhavy, T. J. Surface sensing and adhesion of *Escherichia coli* controlled by the
497   Cpx-signaling pathway. *Proceedings of the National Academy of Sciences* **99**, 2287-2292,
498   doi:10.1073/pnas.042521699 (2002).
499 18 Adams, J. L. & McLean, R. J. Impact of *rpoS* deletion on *Escherichia coli* biofilms. *Appl.*
500   *Environ. Microbiol.* **65**, 4285-4287 (1999).

19 Corona-Izquierdo, F. P. & Membrillo-Hernandez, J. A mutation in *rpoS* enhances biofilm
formation in *Escherichia coli* during exponential phase of growth. *FEMS Microbiol. Lett.*
211, 105-110 (2002).

20 Gerstel, U., Park, C. & Römling, U. Complex regulation of *csgD* promoter activity by global
regulatory proteins. *Mol. Microbiol.* **49**, 639-654, doi:doi:10.1046/j.1365-
2958.2003.03594.x (2003).

21 Gerstel, U. & Romling, U. The *csgD* promoter, a control unit for biofilm formation in
Salmonella Typhimurium. *Res. Microbiol.* **154**, 659-667, doi:10.1016/j.resmic.2003.08.005
(2003).

22 Amores, G. R., de las Heras, A., Sanches-Medeiros, A., Elfick, A. & Silva-Rocha, R. Systematic
identification of novel regulatory interactions controlling biofilm formation in the
bacterium *Escherichia coli*. *Sci. Rep.* **7**, 16768, doi:10.1038/s41598-017-17114-6 (2017).

23 Whitchurch, C. B., Tolker-Nielsen, T., Ragas, P. C. & Mattick, J. S. Extracellular DNA required
for bacterial biofilm formation. *Science* **295**, 1487, doi:10.1126/science.295.5559.1487
(2002).

24 Vilain, S., Pretorius, J. M., Theron, J. & Brözel, V. S. DNA as an Adhesin: *Bacillus cereus*
Requires Extracellular DNA To Form Biofilms. *Appl. Environ. Microbiol.* **75**, 2861,
doi:10.1128/AEM.01317-08 (2009).

25 Tetz, G. V., Artemenko, N. K. & Tetz, V. V. Effect of DNase and Antibiotics on Biofilm
Characteristics. *Antimicrob. Agents Chemother.* **53**, 1204, doi:10.1128/AAC.00471-08
(2009).

26 Niba, E. T. E., Naka, Y., Nagase, M., Mori, H. & Kitakawa, M. A Genome-wide Approach to
Identify the Genes Involved in Biofilm Formation in *E. coli*. *DNA Res.* **14**, 237-246,
doi:10.1093/dnares/dsm024 (2008).

27 Aedo, S. J., Ma, H. R. & Brynildsen, M. P. Checks and Balances with Use of the Keio
Collection for Phenotype Testing. *Methods Mol. Biol.* **1927**, 125-138, doi:10.1007/978-1-
4939-9142-6_9 (2019).

28 Domka, J., Lee, J., Bansal, T. & Wood, T. K. Temporal gene-expression in *Escherichia coli* K-
12 biofilms. *Environ. Microbiol.* **9**, 332-346, doi:10.1111/j.1462-2920.2006.01143.x (2007).

29 Schembri, M. A., Kjærgaard, K. & Klemm, P. Global gene expression in *Escherichia coli*
biofilms. *Mol. Microbiol.* **48**, 253-267, doi:doi:10.1046/j.1365-2958.2003.03432.x (2003).

30 Puttamreddy, S., Cornick, N. A. & Minion, F. C. Genome-Wide Transposon Mutagenesis
Reveals a Role for pO157 Genes in Biofilm Development in *Escherichia coli* O157:H7
EDL933. *Infect. Immun.* **78**, 2377, doi:10.1128/IAI.00156-10 (2010).

31 Goh, K. G. K. *et al.* Genome-Wide Discovery of Genes Required for Capsule Production by
Uropathogenic *Escherichia coli*. *mBio* **8**, e01558-01517, doi:10.1128/mBio.01558-17 (2017).

32 Nhu, N. T. K. *et al.* Discovery of New Genes Involved in Curli Production by a Uropathogenic
Escherichia coli Strain from the Highly Virulent O45:K1:H7 Lineage. *mBio* **9**, e01462-01418,
doi:10.1128/mBio.01462-18 (2018).

33 Yasir, M. *et al.* TraDIS-Xpress: a high-resolution whole-genome assay identifies novel
mechanisms of triclosan action and resistance. *Genome Res*, doi:10.1101/gr.254391.119
(2020).

34 Klemm, P. Two regulatory *fim* genes, *fimB* and *fimE*, control the phase variation of type 1
fimbriae in *Escherichia coli*. *The EMBO Journal* **5**, 1389-1393, doi:10.1002/j.1460-
2075.1986.tb04372.x (1986).

35 Baba, T. *et al.* Construction of *Escherichia coli* K-12 in-frame, single-gene knockout
mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008-2006.0008,
doi:10.1038/msb4100050 (2006).

36 Lehnen, D. *et al.* LrhA as a new transcriptional key regulator of flagella, motility and
chemotaxis genes in *Escherichia coli*. *Mol. Microbiol.* **45**, 521-532, doi:10.1046/j.1365-
2958.2002.03032.x (2002).
37 Blumer, C. *et al.* Regulation of type 1 fimbriae synthesis and biofilm formation by the
transcriptional regulator LrhA of *Escherichia coli*. *Microbiology* **151**, 3287-3298,
doi:10.1099/mic.0.28098-0 (2005).
38 Ko, M. & Park, C. H-NS-Dependent regulation of flagellar synthesis is mediated by a LysR
family protein. *J. Bacteriol.* **182**, 4670-4672, doi:10.1128/jb.182.16.4670-4672.2000 (2000).
39 Alekshun, M. N. & Levy, S. B. Alteration of the Repressor Activity of MarR, the Negative
Regulator of the *Escherichia coli marRAB* Locus, by Multiple Chemicals In Vitro. *J. Bacteriol.*
181, 4669-4672 (1999).
40 Herzberg, M., Kaye, I. K., Peti, W. & Wood, T. K. YdgG (TqsA) controls biofilm formation in
Escherichia coli K-12 through autoinducer 2 transport. *J. Bacteriol.* **188**, 587-598,
doi:10.1128/jb.188.2.587-598.2006 (2006).
41 Holden, E. R. & Webber, M. A. MarA, RamA and SoxS as mediators of the Stress Response:
Survival at a Cost. *Front. Microbiol.* **11**, 828 (2020).
42 Kettles, R. A. *et al.* The *Escherichia coli* MarA protein regulates the *ycgZ-ymgABC* operon to
inhibit biofilm formation. *Mol. Microbiol.* **0**, doi:10.1111/mmi.14386 (2019).
43 Szyf, M. *et al.* DNA methylation pattern is determined by the intracellular level of the
methylase. *Proc. Natl. Acad. Sci. U. S. A.* **81**, 3278-3282, doi:10.1073/pnas.81.11.3278
(1984).
44 Valens, M., Thiel, A. & Boccard, F. The MaoP/*maoS* Site-Specific System Organizes the Ori
Region of the *E. coli* Chromosome into a Macrodomain. *PLoS Genet.* **12**, e1006309-
e1006309, doi:10.1371/journal.pgen.1006309 (2016).
45 Kim, T., Lee, J. & Kim, K.-s. *Escherichia coli* YmdB regulates biofilm formation independently
of its role as an RNase III modulator. *BMC Microbiol.* **13**, 266-266, doi:10.1186/1471-2180-
13-266 (2013).
46 Evans, Margery L. *et al.* The Bacterial Curli System Possesses a Potent and Selective
Inhibitor of Amyloid Formation. *Mol. Cell* **57**, 445-455,
doi:https://doi.org/10.1016/j.molcel.2014.12.025 (2015).
47 Zhang, Y., Morar, M. & Ealick, S. E. Structural biology of the purine biosynthetic pathway.
Cell. Mol. Life Sci. **65**, 3699-3724, doi:10.1007/s00018-008-8295-8 (2008).
48 Pratt, L. A. & Kolter, R. Genetic analysis of *Escherichia coli* biofilm formation: roles of
flagella, motility, chemotaxis and type I pili. *Mol. Microbiol.* **30**, 285-293,
doi:10.1046/j.1365-2958.1998.01061.x (1998).
49 Wang, F. *et al.* Flagellar Motility Is Critical for *Salmonella enterica* Serovar Typhimurium
Biofilm Development. *Front. Microbiol.* **11**, doi:10.3389/fmicb.2020.01695 (2020).
50 Shimada, T., Bridier, A., Briandet, R. & Ishihama, A. Novel roles of LeuO in transcription
regulation of *E. coli* genome: antagonistic interplay with the universal silencer H-NS. *Mol.*
Microbiol. **82**, 378-397, doi:10.1111/j.1365-2958.2011.07818.x (2011).
51 Kroner, G. M., Wolfe, M. B. & Freddolino, P. L. *Escherichia coli* Lrp Regulates One-Third of
the Genome via Direct, Cooperative, and Indirect Routes. *J. Bacteriol.* **201**, e00411-00418,
doi:10.1128/jb.00411-18 (2019).
52 Tramonti, A., De Canio, M. & De Biase, D. GadX/GadW-dependent regulation of the
Escherichia coli acid fitness island: transcriptional control at the *gadY–gadW* divergent
promoters and identification of four novel 42 bp GadX/GadW-specific binding sites. *Mol.*
Microbiol. **70**, 965-982, doi:10.1111/j.1365-2958.2008.06458.x (2008).

53 Sukharev, S. I., Blount, P., Martinac, B., Blattner, F. R. & Kung, C. A large-conductance
mechanosensitive channel in *E. coli* encoded by *mscL* alone. *Nature* **368**, 265-268,
doi:10.1038/368265a0 (1994).
54 Morona, R., Manning, P. A. & Reeves, P. Identification and characterization of the TolC
protein, an outer membrane protein from *Escherichia coli*. *J. Bacteriol.* **153**, 693-699
(1983).
55 Cai, S. J. & Inouye, M. EnvZ-OmpR interaction and osmoregulation in *Escherichia coli*. *J.
Biol. Chem.* **277**, 24155-24161, doi:10.1074/jbc.M110715200 (2002).
56 Marteyn, B. S. *et al.* ZapE is a novel cell division protein interacting with FtsZ and
modulating the Z-ring dynamics. *mBio* **5**, e00022-00014, doi:10.1128/mBio.00022-14
(2014).
57 Hamma, T. & Ferré-D'Amaré, A. R. Pseudouridine Synthases. *Chem. Biol.* **13**, 1125-1135,
doi:https://doi.org/10.1016/j.chembiol.2006.09.009 (2006).
58 Tsui, H. C., Arps, P. J., Connolly, D. M. & Winkler, M. E. Absence of *hisT*-mediated tRNA
pseudouridylation results in a uracil requirement that interferes with *Escherichia coli* K-12
cell division. *J. Bacteriol.* **173**, 7395-7400, doi:10.1128/jb.173.22.7395-7400.1991 (1991).
59 Bringer, M. A., Rolhion, N., Glasser, A. L. & Darfeuille-Michaud, A. The oxidoreductase DsbA
plays a key role in the ability of the Crohn's disease-associated adherent-invasive
Escherichia coli strain LF82 to resist macrophage killing. *J. Bacteriol.* **189**, 4860-4871,
doi:10.1128/jb.00233-07 (2007).
60 Magnusson, L. U., Gummesson, B., Joksimović, P., Farewell, A. & Nyström, T. Identical,
Independent, and Opposing Roles of ppGpp and DksA in *Escherichia coli*. *J. Bacteriol.* **189**,
5193-5202, doi:10.1128/jb.00330-07 (2007).
61 Anwar, N., Rouf, S. F., Römling, U. & Rhen, M. Modulation of Biofilm-Formation in
Salmonella enterica Serovar Typhimurium by the Periplasmic DsbA/DsbB Oxidoreductase
System Requires the GGDEF-EAL Domain Protein STM3615. *PLoS One* **9**, e106095,
doi:10.1371/journal.pone.0106095 (2014).
62 Smith, D. R. *et al.* The Production of Curli Amyloid Fibers Is Deeply Integrated into the
Biology of *Escherichia coli*. *Biomolecules* **7**, 75, doi:10.3390/biom7040075 (2017).
63 Garavaglia, M., Rossi, E. & Landini, P. The pyrimidine nucleotide biosynthetic pathway
modulates production of biofilm determinants in *Escherichia coli*. *PLoS One* **7**, e31252-
e31252, doi:10.1371/journal.pone.0031252 (2012).
64 Cepas, V. *et al.* Transposon Insertion in the *purL* Gene Induces Biofilm Depletion in
Escherichia coli ATCC 25922. *Pathogens (Basel, Switzerland)* **9**,
doi:10.3390/pathogens9090774 (2020).
65 Rodrigues, D. F. & Elimelech, M. Role of type 1 fimbriae and mannose in the development
of *Escherichia coli* K12 biofilm: from initial cell adhesion to biofilm formation. *Biofouling* **25**,
401-411, doi:10.1080/08927010902833443 (2009).
66 Li, S. *et al.* An Osmoregulatory Mechanism Operating through OmpR and LrhA Controls the
Motile-Sessile Switch in the Plant Growth-Promoting Bacterium *Pantoea alhagi*. *Appl.
Environ. Microbiol.* **85**, e00077-00019, doi:10.1128/aem.00077-19 (2019).
67 Barrios, A. F., Zuo, R., Ren, D. & Wood, T. K. Hha, YbaJ, and OmpA regulate *Escherichia coli*
K12 biofilm formation and conjugation plasmids abolish motility. *Biotechnol. Bioeng.* **93**,
188-200, doi:10.1002/bit.20681 (2006).
68 Garcia-Contreras, R., Zhang, X. S., Kim, Y. & Wood, T. K. Protein translation and cell death:
the role of rare tRNAs in biofilm formation and in activating dormant phage killer genes.
PLoS One **3**, e2394, doi:10.1371/journal.pone.0002394 (2008).

69 Sharma, V. K. & Bearson, B. L. Hha controls *Escherichia coli* O157:H7 biofilm formation by
differential regulation of global transcriptional regulators FlhDC and CsgD. *Appl. Environ.*
Microbiol. **79**, 2384-2396, doi:10.1128/AEM.02998-12 (2013).
70 Ogasawara, H., Yamamoto, K. & Ishihama, A. Role of the Biofilm Master Regulator CsgD in
Cross-Regulation between Biofilm Formation and Flagellar Synthesis. *J. Bacteriol.* **193**,
2587-2597, doi:10.1128/jb.01468-10 (2011).
71 Fang, X. & Gomelsky, M. A post-translational, c-di-GMP-dependent mechanism regulating
flagellar motility. *Mol. Microbiol.* **76**, 1295-1305, doi:10.1111/j.1365-2958.2010.07179.x
(2010).
72 Nieto, V. *et al.* Under Elevated c-di-GMP in *Escherichia coli*, YcgR Alters Flagellar Motor Bias
and Speed Sequentially, with Additional Negative Control of the Flagellar Regulon via the
Adaptor Protein RssB. *J. Bacteriol.* **202**, e00578-00519, doi:10.1128/JB.00578-19 (2019).
73 Paul, K., Nieto, V., Carlquist, W. C., Blair, D. F. & Harshey, R. M. The c-di-GMP binding
protein YcgR controls flagellar motor direction and speed to affect chemotaxis by a
"backstop brake" mechanism. *Mol. Cell* **38**, 128-139, doi:10.1016/j.molcel.2010.03.001
(2010).
74 Pesavento, C. *et al.* Inverse regulatory coordination of motility and curli-mediated adhesion
in *Escherichia coli*. *Genes Dev.* **22**, 2434-2446, doi:10.1101/gad.475808 (2008).
75 Wood, T. K., González Barrios, A. F., Herzberg, M. & Lee, J. Motility influences biofilm
architecture in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* **72**, 361-367,
doi:10.1007/s00253-005-0263-8 (2006).
76 Prouty, A. M. & Gunn, J. S. Comparative analysis of *Salmonella enterica* serovar
Typhimurium biofilm formation on gallstones and on glass. *Infect. Immun.* **71**, 7154-7158
(2003).
77 Samanta, P., Clark, E. R., Knutson, K., Horne, S. M. & Prüß, B. M. OmpR and RcsB abolish
temporal and spatial changes in expression of *flhD* in *Escherichia coli* Biofilm. *BMC*
Microbiol. **13**, 182, doi:10.1186/1471-2180-13-182 (2013).
78 Trampari, E. *et al.* Antibiotics select for novel pathways of resistance in biofilms. *bioRxiv*,
605212, doi:10.1101/605212 (2019).
79 Barquist, L. *et al.* The TraDIS toolkit: sequencing and analysis for dense transposon mutant
libraries. *Bioinformatics* **32**, 1109-1111, doi:10.1093/bioinformatics/btw022 (2016).
80 Carver, T., Harris, S. R., Berriman, M., Parkhill, J. & McQuillan, J. A. Artemis: an integrated
platform for visualization and analysis of high-throughput sequence-based experimental
data. *Bioinformatics* **28**, 464-469, doi:10.1093/bioinformatics/btr703 (2011).
677
678

679 **Figures**

681 **Figure 1:** Genes involved in biofilm formation over time in *E. coli*. Expression of genes in green

682 were found to be beneficial for, and those in red were found to be detrimental to, biofilm formation.

Figure 2: Mapped reads from TraDIS-Xpress data, plotted with BioTraDIS in Artemis. **a)** Insertion

sites in and around *fimB* and *fimE* in planktonic and biofilm conditions after 12 and 48 hours

growth. **b)** Insertion sites in and around *hdfR*, *lrhA* and *tomB* in planktonic and biofilm conditions

after 24 hours growth. **c)** Insertion sites in and around *dksA* and *dsbA* in planktonic and biofilm

conditions after 12 and 48 hours growth. For all plot files, one of two independent replicates is

shown and the transposon-located promoter is induced with 1mM IPTG in all conditions shown.

Figure 3: Phenotypic validation of selected genes involved in biofilm formation. **a**) Biofilm biomass of single knockout mutants relative to wild type *E. coli* BW25113, measured by crystal violet staining. Two biological and a minimum of two technical replicates were performed for each mutant. **b**) Cell aggregation of single knockout mutants relative to wild type *E. coli* BW25113, measured by $OD_{600\ nm}$ of the supernatant of unagitated cultures. Points show the ODs of three independent replicates. For both graphs, red points/bars show data from the first KEIO collection mutant of each gene, and blue points/bars show data from the second mutant. Error bars show 95% confidence intervals and the shaded area shows the 95% confidence interval of the wild type. Single asterisks (*) represent a significant difference between one Keio mutant copy and the wild type, and double asterisks (**) denote a significant difference between both Keio mutant copies and the wild type (Welch's *t*-test, $p < 0.05$). **c**) Colonies grown on agar supplemented with Congo red to compare curli biosynthesis between single knockout mutants and the wild type. Images are representative of 2 biological and 2 technical replicates.

709 **Figure 4**: Biofilm formation of single knockout mutants on glass analysed under flow conditions

710 after 12 and 48 hours growth. 10x Magnification. Images are representative of two independent

711 replicates. Scale bar indicates 5 µm.

714 **Figure 5**. Summary of genes important for biofilm formation by *E. coli* at different stages of

715 development.