Genomic surveillance framework and global population structure for *Klebsiella pneumoniae*

Margaret M. C. Lam[1]*, Ryan R. Wick[1], Stephen C. Watts[2], Louise T. Cerdeira[1], Kelly L. Wyres[1], Kathryn E. Holt[1,3]

[1]Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria 3004, Australia. [2]Department of Biochemistry and Molecular Biology, Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Victoria 3010, Australia. [3]London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK.

*Corresponding author: margaret.lam@monash.edu

ABSTRACT

K. pneumoniae is a leading cause of antimicrobial-resistant (AMR) healthcare-associated infections, neonatal sepsis and community-acquired liver abscess, and is associated with chronic intestinal diseases. Its diversity and complex population structure pose challenges for analysis and interpretation of *K. pneumoniae* genome data. Here we introduce Kleborate, a tool for analysing genomes of *K. pneumoniae* and its associated species complex, which consolidates interrogation of key features of proven clinical importance. Kleborate provides a framework to support genomic surveillance and epidemiology in research, clinical and public health settings. To demonstrate its utility we apply Kleborate to analyse publicly available *Klebsiella* genomes, including clinical isolates from a pan-European study of carbapenemase-producing *Klebsiella*, highlighting global trends in AMR and virulence as examples of what could be achieved by applying this genomic framework within more systematic genomic surveillance efforts. We also demonstrate the application of Kleborate to detect and type *K. pneumoniae* from gut metagenomes.

TEXT

Klebsiella pneumoniae bacteria commonly colonize the mammalian gut, but are also

recognized as a major public health threat due to their ability to cause severe

infections in healthcare settings and their association with antimicrobial resistance

(AMR)[1,2]. Reports of *K. pneumoniae* gut colonization frequencies vary by country

and demographics, but prevalence rates as high as 87% have been reported[3–6]. *K.*

pneumoniae colonization is implicated in chronic diseases of the gastrointestinal tract

including inflammatory bowel disease and colorectal cancer[7]. There is also a growing

body of evidence highlighting colonization as a reservoir for extraintestinal infections

(urinary tract infection, pneumonia, wound or surgical site infections, sepsis) in

vulnerable individuals such as neonates, the elderly, immunocompromized and

hospitalized patients[8]. Treatment of healthcare-associated (HA) *K. pneumoniae*

infections is often limited by multidrug resistance (MDR) resulting from the

accumulation of horizontally-acquired AMR genes and mutations in core genes[2].

Treatment is further complicated by increasing frequencies of strains producing

extended-spectrum β-lactamases (ESBL) and/or carbapenemases, prompting

increased reliance on colistin and β-lactamase inhibitor combinations[9,10]. The World

Health Organization has accordingly prioritized *K. pneumoniae* as a target for new

drugs and therapies[11].

Outside healthcare settings, *K. pneumoniae* is also recognized as a causative agent of

community-acquired infections including urinary tract infection and pneumonia, but

also invasive infections such as pyogenic liver abscess, endophthalmitis and

meningitis[12,13]. Invasive community-acquired infections are generally associated with

so-called hypervirulent *K. pneumoniae* (hvKp) and are most commonly reported in

East and Southeast Asia, or in individuals with East Asian ancestry[12]. Features

associated with hvKp include a K1, K2 or K5 polysaccharide capsule and

horizontally-acquired virulence factors encoding the siderophores aerobactin (Iuc)

and salmochelin (Iro), the genotoxin colibactin (Clb), and a hypermucoid phenotype

(conferred by the *rmpADC* locus)[14–18]. HvKp are rarely MDR and most remain

susceptible to drugs except ampicillin, to which *K. pneumoniae* are intrinsically

resistant due to the chromosomally-encoded β-lactamase SHV[19]. However there have

been increasing reports of hvKp carrying AMR plasmids and co-occurrence of AMR

and virulence determinants in non-hvKp isolates. The convergence of AMR and

virulence in *K. pneumoniae* potentiates invasive and difficult-to-treat infections, and

at least one fatal outbreak has been documented in China where carbapenemase-

producing hvKp are increasingly common[20–24].

Research conducted in the pre-genomic era characterized 77 distinct capsular (K)

serotypes[25], nine O types[26] and variable AMR profiles amongst the *K. pneumoniae*

population[27,28], indicating a diverse genetic and phenotypic landscape[15,29]. In recent

years, genomic studies have provided key insights into the population structure of *K.*

pneumoniae (recently summarized in Wyres et al[16]), revealing hundreds of deep-

branching phylogenetic lineages comprising sequence types (STs) or clonal groups

(CGs) defined by the seven-gene multi-locus sequence typing (MLST) scheme[29].

Some of these lineages correspond to lineages (i.e. STs and CGs) that have

accumulated large numbers of AMR genes that have become globally distributed (e.g.

CG258, CG15, ST307); these are dubbed MDR clones and have been linked with HA

infections and hospital outbreaks worldwide[30]. Others carry a high load of virulence

genes (e.g. CG23, CG65, CG86) and are recognized as hvKp associated with

community-acquired infections. Further distinguishing MDR from hvKp clones are

their K and O antigen profiles, with the former displaying a diverse range of K and O

biosynthesis loci as a result of homologous recombination between strains, while

hvKp rarely deviate from the K1, K2 or K5 types[16].

Importantly, genomic characterization of clinical isolates identified as *K. pneumoniae*

via biochemical tests or mass spectrometry (MALDI-TOF) has revealed the existence

of multiple related species and subspecies, which together form the *K. pneumoniae*

species complex (KpSC). These differ by 3-4% nucleotide divergence across core

chromosomal genes, but share the same pool of AMR and virulence genes[16].

Infections and outbreaks caused by other KpSC members have been reported but they

generally account for a significantly lower disease burden than *K. pneumoniae* (10-

20%)[19,31,32]. Genomics has also clarified that the two *K. pneumoniae* subspecies

originally defined by distinct and unusual disease manifestations (subsp.

rhinoscleromatis which causes a progressive and chronic granulomatous infection

known as rhinoscleroma, and subsp. *ozaenae* which causes atrophic rhinitis or ozena)

actually represent CGs of *K. pneumoniae* (CG3 and CG90)[15]. Like hvKp clones, these

strains also express specific capsule types (K3, K4 and K5) alongside aerobactin and

another acquired siderophore, yersiniabactin (Ybt)[16].

Due to its clinical importance and increasing AMR, *K. pneumoniae* is increasingly the

focus of surveillance efforts and molecular epidemiology studies. The sheer volume

of clinically-relevant molecular targets renders whole genome sequencing (WGS) the

most cost-efficient characterization approach, however extracting and interpreting

clinically important features is challenging. To address this, we have developed

Kleborate, a genotyping tool designed specifically for *K. pneumoniae* and the

associated species complex, which consolidates detection and genotyping of key

virulence and AMR loci alongside species, lineage (ST) and predicted K and O

antigen serotypes directly from genome assemblies. Here we describe Kleborate and

demonstrate its utility by application to publicly available datasets. First, we show

that Kleborate can rapidly recapitulate and augment the key findings from a recent

large-scale European genomic surveillance study[33]. Next, we apply Kleborate to a

curated collection of 13,156 publicly available WGS to further showcase its utility

and derive novel insights into the global epidemiology of *Klebsiella* AMR, virulence

and convergence. Finally, we show that Kleborate can also be applied to detect

clinically relevant genotypes from meta-genome assembled genomes (MAGs).

RESULTS

Integrated genomic framework and genotyping tool

Our goal was to develop a single tool that can rapidly extract genotype information

that is clinically relevant to *K. pneumoniae* and other members of the species complex

in order to support genomic epidemiology and surveillance. We have previously

reported genotyping schemes for the acquired *K. pneumoniae* virulence loci *ybt, clb,*

iuc and *iro*[34,35] (whose detection and typing is implemented in early versions of

Kleborate), and also K and O antigen typing implemented in the software Kaptive[36].

Here we expand the Kleborate framework to include additional features including

taxonomic assignment to species and subspecies, assignment to lineages via seven-

locus MLST, detection and genotyping of the *rmp* hypermucoidy locus and the *rmpA2*

gene, and identification of AMR determinants (mutations and horizontally acquired

genes, including assignment of SHV β-lactamase alleles as either ESBL, β-lactamase

inhibitor resistance, or intrinsic ampicillin resistance only, see **Methods** and

Supplementary text). Kleborate can optionally call Kaptive for K/O antigen

prediction.

Unlike generic AMR or virulence typing tools, we include only genetic features for

which there is strong evidence of an associated phenotype in *K. pneumoniae* that has

confirmed clinical relevance. These are reported in a manner that facilitates

interpretation, including summarizing virulence and AMR genotypes into scores that

reflect escalating clinical risk in *K. pneumoniae* infections. Kleborate features are

summarized in **Table 1** and methodological details for genotyping are provided in

Methods. For a typical 5.5 Mbp genome, a Kleborate run including AMR typing

takes <10 seconds on a laptop, while robust K and O serotype prediction using

Kaptive[36] adds an additional ~1 minute. Results are output in tab-delimited format,

making it easy to integrate Kleborate into existing workflows.

Species and subspecies assignment

The taxonomy of *Klebsiella* is rapidly evolving, with several new species and

subspecies recently identified[37–39]. As a consequence, many genomes in public

databases are incorrectly assigned. We therefore introduced a custom approach for

rapid and accurate species and subspecies identification for *Klebsiella*, based on Mash

distances[40] to a taxonomically-curated genome set (representative tree in **Figure S1A-**

B), avoiding the need for users to download large reference genome databases (see

Methods). This approach was validated using a set of n=285 diverse clinical isolates

and compared with species assignments based on the read-based taxonomic classifier

Kraken2 (details in **Supplementary Text**, **Table S1, Figure S1**).

Virulence and AMR scores

Genomes are scored according to the clinical risk associated with the AMR and virulence loci that are detected (see **Methods**). Here we take advantage of the structured distribution of AMR and virulence determinants within the *K. pneumoniae* population[14] to reduce the genotyping data to simple numerical summary scores that reflect the accumulation of loci contributing to clinically relevant AMR or hypervirulence: virulence scores range from 0 to 5, depending on the presence of key loci associated with increasing risk (yersiniabactin < colibactin < aerobactin); resistance scores range from 0 to 3, based on detection of genotypes warranting escalation of antimicrobial therapy (ESBL < carbapenemase < carbapenemase plus colistin resistance, see **Table 1**). These simple numerical scores facilitate downstream analyses including trend detection. For example, analysis of a non-redundant subset of 9,705 publicly available *K. pneumoniae* genomes (see below, **Table S2**) showed increasing AMR and virulence scores over time (barplots in **Figure 1A-B**). The virulence and resistance scores were correlated not only with the prevalence of individual components that contribute to the scores, but also with other components that are co-distributed in the population (lines in **Figure 1A-B**). For example, the frequencies of *rmpADC* and *rmpA2* loci over time were correlated with the virulence score (**Figure 1A**); and the resistance score was correlated with the mean number of acquired AMR genes and associated drug classes (excluding ESBLs, carbapenemases and colistin which contribute to the score) (**Figure 1C**). Consistent with this, genomes with resistance scores >0 (assigned based on the presence of ESBL and/or carbapenemase genes) typically carry many additional AMR genes conferring resistance to multiple drug classes (**Figure 1D-E**). Reducing the data to key axes of virulence and AMR also facilitates exploration of subpopulations associated with

AMR, virulence or convergence of both traits; such as specific *K. pneumoniae*

lineages or specimen types (see below).

Rapid genotyping of clinical isolates from a large-scale surveillance study

We applied Kleborate to analyse all *K. pneumoniae* clinical isolate genomes deposited

in RefSeq by the EuSCAPE surveillance study (927 carbapenem-non-susceptible, 697

carbapenem-susceptible; see **Table S2**)[33]. Kleborate rapidly and accurately

reproduced key findings from the original study, which were originally derived from

multi-step analyses comprising five independent tools and four independent databases

(each from a different public repository, one with additional manual curation): (i)

70.2% of carbapenem-non-susceptible genomes (n=651/927) carried carbapenemases,

mainly KPC-3, OXA-48, KPC-2 and NDM-1; (ii) these were dominated by a few

major clones, ST11, ST15, ST45, ST101, ST258, and ST512; (iii) individual countries

were associated with specific carbapenemase/clone combinations (see **Figure 2A**). A

detailed comparison of the results reported by Kleborate versus those reported in the

original study is provided in **Supplementary Text** and **Table S3**.

In addition to the detection of carbapenemase genes, Kleborate also identified porin

defects, which are known to contribute to the carbapenem-resistance phenotype[41,42],

in 36.5% of EuSCAPE genomes (including 60% of those with carbapenemase genes

and 19.9% of those without carbapenemase genes). These defects included

truncation/deletion of OmpK35 and/or OmpK36 (also considered in the original

study) as well as GD or TD insertions in the OmpK36 β-strand loop[41] (not considered

in original study, but here detected in 18.6% of genomes including 18 with no porin

deletion). **Figure 3** shows meropenem MICs stratified by porin defect-carbapenemase

combinations identified by Kleborate, highlighting the importance of porin defects –

including the OmpK36 β-strand loop insertions – for full expression of carbapenem

resistance in *K. pneumoniae*.

The rise in carbapenem-resistant *K. pneumoniae* infections in hospitals and its

associated morbidity and mortality[43] has led to increased interest in alternative control

strategies such as vaccines, phage therapy and antibody therapy, key targets for which

are the K and O surface antigens[44,45]. Kleborate confidently identified K and O

biosynthesis loci in 98.3% and 99.1% of EuSCAPE genomes, respectively, including

87 distinct K loci and 11 distinct O loci (**Figures S2** and **S3**). Amongst carbapenem-

non-susceptible isolates (meropenem MIC >2), 38 distinct K types were identified

and the most common were KL107 (n=173), KL17 (n=67), KL106 (n=41), KL24

(n=35), KL15 (n=19) and KL36 (n=13). Seven distinct O types were detected among

these genomes, and the most common were O2 (n=294), O1 (n=136) and O4 (n=52).

Overall, the data suggest an intervention would need to be effective against six K

types or two O types in order to provide coverage of 80% of carbapenem-resistant

infections in Europe (**Figure 2B-C**). However, it is important to explore the impact of

population structure on these findings, specifically the impact of local clonal

expansions. Kleborate aides this type of analysis by providing ST and other

genotyping information alongside the K and O locus types, which can be viewed in

the context of geographic information. Doing so revealed that each of the top three K

loci were dominated by a single ST (83.5% of KL107 were ST512; 93.0% of KL105

were ST11; 91.4% of KL17 were ST101). Importantly, the vast majority of ST512-

KL107 genomes (75.3%) originated from Italy where this ST is known to be locally

circulating[46,47], while 58% of ST11-KL105 originated from Poland and Slovakia, and

56% of ST101-KL17 originated from Serbia and Romania. When these putative local

expansions were excluded, the top 6 K loci were (KL24, KL15, KL2, KL112, KL107,

KL151) and accounted for just 34% of the remaining genomes.

Global population snapshot of *K. pneumoniae* AMR and virulence

We applied Kleborate to analyse n=13,156 *Klebsiella* genomes (see **Methods**, **Table**

S2). Here we provide a brief overview of the data followed by an exploration of

AMR, virulence and the phenomenon of convergence, with the aim to highlight the

rich information and types of inferences that can be derived from Kleborate output.

The genome data represented isolates collected from a range of sources in 99

countries between 1920–2020 (**Table S4**, although human isolates from the USA,

China and UK dominated the data set accounting for n=4,702 genomes, 35.7% of

total). The majority of these genomes were sourced from RefSeq, and among these

Kleborate identified 1.0% (n=103/10,747) as a species other than the taxon recorded

in NCBI; this is consistent with other studies and highlights the current confusion

around taxonomic designations in *Klebsiella*. The most common species was *K.*

pneumoniae (n=11,259, 86%); the rest comprised other KpSC species (9.4%), other

members of the *K. oxytoca* species complex (3.1%) and *K. aerogenes* (1.9%) (**Figure**

4, Table S4). AMR and virulence genes were concentrated in the KpSC and

particularly *K. pneumoniae* (**Figure 4, Table S5**).

The collection captured extensive phylogenetic diversity across the *K. pneumoniae*

species (see interactive phylogeny at

http://microreact.org/project/JDyan46yctyDh6weEUjWN), and Kleborate assigned

these genomes to ≥1,452 different STs (1,119 known STs across and at least 333

novel STs). Notably, 600 STs (41%) were represented by just a single genome each

(accounting for 5.3% of all genomes). We detected n=4 ST67 (subspecies

rhinoscleromatis) and n=3 ST90 (subspecies *ozanae*). A small number of STs were

overrepresented, reflecting the bias towards sequencing MDR and hypervirulent

isolates, as well as those causing hospital outbreaks. For example, 1,354 genomes

(12.0%) represented the KPC-associated ST258, which is known to dominate

carbapenem-resistant *K. pneumoniae* in the USA and southern Europe (where it has

been the subject of intense genomic investigations) but is comparatively rare in other

regions of the world[16]. To reduce the impact of these sampling biases in public

genome collections, we down-sampled to a non-redundant set of 9,705 *K.*

pneumoniae genomes representing unique combinations of ST, genetic subcluster

(Mash distance <0.0003), virulence genotype, AMR genotype, specimen type,

location and year of isolation (see **Methods**). However, we cannot fully correct for

the sampling biases inherent in the public genome data and even after subsampling,

the 30 most common STs accounted for 63.4% of genomes (n≥50 genomes each,

n=6,151 total; see **Figure S4**). **Figure 5** shows the distribution of AMR and virulence

scores amongst non-redundant genomes from these 30 common *K. pneumoniae* STs

(n>50 per ST), each of which displays high rates of AMR and/or virulence.

AMR determinants

SHV β-lactamases conferring intrinsic resistance to the penicillins were detected in

85.9% of the 9,705 non-redundant *K. pneumoniae* genomes (ESBL forms of SHV

were detected in 10.0%). Acquired AMR was widespread (77.1% of genomes had at

least one gene or mutation conferring acquired AMR detected) and 71.6% of genomes

were predicted to be MDR (acquired resistance to ≥3 drug classes[48]), a much higher

rate than is reported in most geographical regions[3,49–51], reflecting the bias within

public genome collections. The majority of genomes had a non-zero resistance score,

reflecting presence of ESBL and/or carbapenemase genes: 22.3%, 37.1% and 5.9%

genomes had resistance scores of 1, 2 and 3 respectively. Mean resistance scores

increased through time (**Figure 1B**). This trend could be an artefact of sampling bias

towards the selective sequencing of AMR isolates, however it is consistent with the

increasing AMR rates reported in surveillance studies globally[52–54].

Comparatively higher prevalence of acquired AMR genes was observed in some STs

(**Figure S4**). Many of these STs represent recognized MDR clones largely from

clinical samples that were also associated with high mean resistance scores (**Figures

6A-B**), driven by high frequency of ESBL and carbapenemase genes (**Figures 5,

S5A-B**). The most common ESBLs/carbapenemases were widely detected across the

population (46-299 STs each), including amongst the top 30 common STs (prevalence

range per ST, 0.1-100%; see **Figure S5A-B**), highlighting their mobile nature. The

notable exception was CTX-M-65, which appeared to be largely clone specific,

detected in only 9 STs and ST11 accounting for 96.7% of these genomes.

Colistin resistance determinants were detected in 8.7% of the non-redundant *K.

pneumoniae* genomes. These were mostly nonsense mutations in MgrB or PmrB

(83.5%) rather than acquisition of an *mcr* gene (15.8%, and an additional 6 genomes

with both acquired *mcr* and truncated MgrB/PmrB). The rate of detection ranged from

0-25.2% for the 30 most common STs, and was highest amongst ST512, ST437,

ST147, ST16 and ST258 (**Figure S5C**), each of which are also associated with high

rates of carbapenem-resistance. Porin mutations were detected in 37.9% of genomes

(34.0% OmpK35, 20.2% OmpK36, 16.3% both). High prevalence of specific porin

defects have been reported previously in some clones[41,42], and this was reflected in

our analysis of ST258 and its derivative ST512. We observed OmpK35 truncations in

99.9% of non-redundant ST258 genomes (with or without truncations or substitutions

in OmpK36), and truncations in OmpK35 and/or OmpK36 in all ST512 (99.4% with

OmpK35 truncations, 94.4% with the OmpK36GD mutation, see **Figure S5D**).

Virulence loci

The prevalence of acquired siderophores and colibactin loci amongst non-redundant

K. pneumoniae genomes was 44.4% *ybt*, 7.5% *clb*, 11.2% *iuc* and 7.0% *iro*. The loci

were found across diverse *K. pneumoniae* STs (391 STs with *ybt*, 56 with *clb*, 144

with *iuc*, 108 with *iro*) but were rarely detected in other *Klebsiella* species (with the

exception of *ybt* among the *K. oxytoca* species complex, see **Figure 4**) indicating

frequent mobilisation within *K. pneumoniae* but not between species (**Table S6**,

Figure S6). Mean virulence scores increased through time (**Figure 1A**). **Figure 5B**

shows the frequency of virulence scores in the top 30 most common STs in the non-

redundant genome set. Sixteen of these common STs had ≥40% of genomes carrying

the ICE*Kp*-associated *ybt* without the virulence plasmid-associated *iuc* locus (i.e.

virulence score=1-2), including well known MDR clones ST258, ST11, ST14, ST15,

ST101, ST147, ST152, ST395. Only the hvKp clones (ST23, ST86, ST65) and ST231

had a high frequency of *iuc* (virulence score ≥3).

In addition to detecting the presence of virulence loci, Kleborate reports on their

completeness, genetic lineages and associated MGE variants, which can provide

insights into their dissemination. Most of the virulence loci identified in the non-

redundant *K. pneumoniae* data set (98%) matched one of the genetic lineages

described previously[34,35] (**Table S6**). **Figure S7A** shows the frequency of *iuc* lineages

in *K. pneumoniae* STs with ≥20 non-redundant genomes and at least one genome

harbouring *iuc*. There were four STs for which >60% genomes harboured *iuc*, and

only a single *iuc* lineage was detected in each (*iuc1* in ST23, ST65, ST86; *iuc2A* in

ST82), consistent with long-term persistence of a specific virulence plasmid in these

well-known hypervirulent clones. In contrast, *iuc* was less frequent among other STs,

several of which were associated with multiple *iuc* lineages (e.g. ST231, ST25,

ST35), consistent with more recent and/or transient virulence plasmid acquisitions

(mostly *iuc1*, followed by *iuc3* and *iuc5*).

Frameshift mutations (i.e. truncations) and/or incomplete loci (i.e. missing at least one

gene) were detected in 10%, 28.5%, 13.6% and 17.7% of non-redundant *K.*

pneumoniae genomes with *ybt*, *clb*, *iuc* and *iro* respectively (**Table S7**). While some

of these may erroneously arise from contig breaks in draft genome assemblies, true

truncations or missing genes may reflect a lack of function. The latter is likely true for

instances where we observe conserved frameshift mutations across entire lineages,

e.g. frameshift mutations were detected in *iucA* for all *iuc3+* genomes and in *iroC* for

all *iro3+* and *iro4+* genomes.

The hypermucoidy locus *rmpADC* was detected in 8.4% of non-redundant *K.*

pneumoniae genomes (and just eight genomes of other KpSC species, **Table S6**). The

majority of these genomes (67.2%, belonging to >79 STs) carried intact copies of all

three genes, thus likely express the hypermucoid phenotype. Intact *rmpADC* was

common in *iuc*-positive genomes of the hvKp clones ST23 and ST86, as well as MDR

clones ST29 and ST101 (**Figure S7B**). Many other *iuc*-positive genomes carried

rmpADC loci with truncated or missing genes, which likely do not confer the

hypermucoid phenotype. Notably, these included hvKp clones ST65 and ST82, as

well as MDR clones ST231, ST15 and ST14. The *rmpA2* gene was detected in 7.4%

of non-redundant *K. pneumoniae* genomes, but was mostly present in truncated form

(89.0% of *rmpA2*+ genomes) due to frameshifts within a poly-G tract[55]. The latter

highlights the importance of considering not only the presence/absence of a given

gene, but also whether it encodes a full-length protein, which may have important

clinical implications.

Facilitating detection of AMR-virulence convergence

AMR and virulence determinants have until recently been segregated in non-

overlapping *K. pneumoniae* populations[14,19], as clearly indicated by the distributions

of AMR and virulence scores among STs (**Figures 5, 6A**). However, reports of

convergent AMR-virulent strains with the potential to cause difficult-to-treat

infections are increasingly common[16,56]. Kleborate facilitates rapid identification of

such strains on the basis of resistance and virulence scores (convergence defined as

virulence score ≥3 and resistance score ≥1, **Figure 6C**). Based on these scores, we

observed a total of 601 convergent *K. pneumoniae* (510 non-redundant) with the

highest proportion corresponding to a virulence score of 4 (indicative of

yersiniabactin plus aerobactin/virulence plasmid detection) and resistance score of 2

(carbapenem resistance).

The majority of convergent genomes (74.5%) were concentrated within a small

number of STs comprising the well-known hypervirulent (ST23, ST86, ST65) and

MDR lineages (ST11, ST15, ST231 and ST147) (**Figures 6C-D, 7**). We combined the

genotyping data and information from a Mash-distance-based neighbour-joining tree

(http://microreact.org/project/JDyan46yctyDh6weEUjWN) to define unique

convergence events (defined as unique combinations of ST, virulence and resistance

determinants, and phylogenetic cluster). This identified n=173 convergence events,

accounted for by either acquisition of the virulence plasmid by MDR/other clones

(n=84 events; 475 genomes), or acquisition of ESBLs/carbapenemases by

hypervirulent clones (n=89 events; 126 genomes) (**Figure 7**, **Table S8**).

The most common virulence plasmid, KpVP-1 (*iuc1* ± *iro1*), accounted for 54% of

virulence plasmid acquisition events (n=45 acquisitions), while *iuc3* plasmids, the *E.*

coli derived *iuc5* (±*iro5*) and *iuc/iro* unknown (i.e. novel or divergent *iuc/iro* loci)

accounted for 21%, 11% and 14%, respectively (**Figure 7**). AMR acquisitions by

hypervirulent clones involved the ESBL/carbapenemase genes that are most common

in the general *K. pneumoniae* population: KPC-2 (26%), OXA-232 (17%) and CTX-

M-15 (18%). The majority of convergence events (87%) were associated with just a

small number of genomes (i.e. n≤3); however, five events were associated with >20

genomes in the complete dataset, which may indicate clonal expansion and

dissemination of the corresponding convergent strains locally and/or between

countries. One such event corresponded to the ST11-KPC + KpVP-1 deletion variant

strain that was originally reported in 2017[20] and has since been recognized as widely

distributed in China[20–24]. The complete public genome set (i.e. counting redundant

genomes) included 148 genomes corresponding to this specific ST11 convergence

event mostly from China but also from France (n=2). Notably though, this was only

one of 50 convergence events that we detected in China, including 8 involving

acquisition of *iuc1* or *iuc5* by ST11 (see **Table S8**, and interactive tree at

http://microreact.org/project/JDyan46yctyDh6weEUjWN). Additional events associated

with >20 genomes included (i) ST231-MDR + virulence plasmids carrying novel *iuc*

lineages detected in India, Pakistan, Switzerland, Thailand and USA, (ii) ST15-CTX-

M-15 + KpVP-1 in Pakistan, (iii) ST15-MDR + KpVP-1 in China and Nepal, and (iv)

another distinct ST11-KPC-2 + KpVP-1 event in China. Including the above three

examples, 11 convergence events appeared to involve intercountry expansion of

which one has been previously documented[57].

Overall, convergent genomes were detected originating from most geographical

regions for which genome data was available, but some regions had many more

events than others (**Figure 7, Table S8**). This uneven distribution may stem from a

skew in the number of genomes available per region (e.g. due to variation in

accessibility or application of genome sequencing). Nevertheless, the number of

convergent genomes in the eastern, southeastern and southern parts of Asia were

noticeably high, driven by the frequency of convergence events detected in China

(n=50 events) and Thailand (n=26 events) as well as putative clonal expansions of

these strains as discussed above (**Figure 7**). Of note, AMR acquisitions by

hypervirulent lineages were particularly frequent within East and Southeast Asia

where hypervirulent infections are most frequently reported, alongside countries from

eastern and northern Europe.

Outside of *K. pneumoniae,* convergence events were rare: we detected n=2 *K.*

quasipneumoniae subsp. *similipneumoniae* (ST367 with KpVP-1 and CTX-M-15;

ST3387 with *iuc3* and CTX-M-55) and n=2 *K. variicola* subsp. *variicola* (ST595 with

KpVP-1 and KPC-2; ST1848 with *iuc5* and KPC-2).

Genotyping *K. pneumoniae* from metagenome data

There is increasing interest in detection and typing of *K. pneumoniae* direct from gut

metagenome data[58], due to the role of *K. pneumoniae* gut colonization as a source of

acute infections and as a contributor to chronic diseases[7,8]. We tested Kleborate's

performance by application to n=40 metagenomes from which at least one KpSC

isolate was cultured and sequenced, as part of the Baby Biome Study[59]. We compared

the results of running Kleborate on metagenome-assembled genomes (MAGs, i.e.

species-specific contig bins extracted from whole-metagenome assemblies) vs. KpSC

isolate whole genome sequence(s) cultured from the same fecal sample. Thirty-two

metagenomes had >1% relative abundance of KpSC, and genotyping of MAGs from

these yielded results consistent with genotyping of cultured isolates for 26/32 samples

(16 with identical genotypes reported for species, ST, K/O locus, virulence and AMR;

10 with close matches; see **Fig. S8, Tables S9-S10**). As expected, MAG-derived

genotypes were closest to those of isolates when only one KpSC strain was cultured

from the sample (see **Fig. S8, Table S10**). Kleborate analysis of whole metagenome

assemblies (as opposed to individual MAGs) is not recommended: species detection

and ST assignment matched that of the corresponding WGS isolates for only n=4/40

metagenome assemblies, which is unsurprising as the whole metagenomes include

sequences derived from dozens of different bacteria, many of which harbour

homologs of genotyping targets.

DISCUSSION

Whole genome sequencing is being increasingly implemented in research and public

health labs as a cost- and time- effective option for tracking pathogens and AMR.

However, identification of clinically-relevant features remains a key bottleneck that

hinders widespread adoption of genome surveillance. We have presented a

comprehensive framework and tool for rapid genotyping of *Klebsiella* species

genomes: Kleborate is a single unified approach for species detection, MLST and

genotyping of key virulence and AMR determinants. It focusses only on genomic

features for which there is strong evidence of a clinically relevant phenotype in KpSC

and presents the data in a readily interpretable format, with numerical summaries and

categorical scores corresponding to measures of clinical risk.

A key strength of the Kleborate framework is its species-specific approach. This is

particularly important for accurate interpretation of AMR and virulence gene screens

from WGS, wherein the use of generic databases and tools can result in confusion.

Notable examples include the intrinsic *oqxAB* and *fosA* alleles, which unlike for other

Enterobacterales, do not confer resistance to quinolones and fosfomycin when

expressed in KpSC. Kleborate does not report these intrinsic alleles, neither does it

report intrinsic virulence determinants such as the siderophore enterobactin, which is

known to play a role in KpSC pathogenicity but for which the presence alone cannot

be considered to indicate enhanced virulence of one isolate over another. Correct

taxonomic identification of *K. pneumoniae* can be difficult in itself, hence the inbuilt

speciation tool is an important feature (and here identified nearly 100 RefSeq

genomes with incorrect species/subspecies assignments).

Another strength of our approach is the rich data output by Kleborate, which

facilitates in-depth investigation of population structure, AMR and virulence

epidemiology. This allows rapid exploration and understanding of: (i) hypervirulence-

associated loci and the molecular drivers of their dissemination (**Figures S4** and **S7**);

(ii) molecular mechanisms of complex AMR phenotypes e.g. carbapenem resistance

(**Figure 3**); (iii) AMR and virulence trends (**Figures 1**, **5** and **6**); (iv) emerging

convergent AMR-virulent strains so that they can be targeted for surveillance and

infection control (**Figure 7**); (v) overrepresented STs and genotypes, which may be

indicative of transmission clusters that should be targeted for further investigation (as

demonstrated for the EuSCAPE surveillance genomes, **Figure 2A**); (vi) surface

antigen epidemiology, which can inform the design of novel vaccines and

therapeutics (**Figure 2B-C**). Notably Kleborate can also yield useful genotyping

results from metagenomics data (**Figure S8**), which is gradually being adopted for

clinical and surveillance applications relevant to *K. pneumoniae*. User interpretation

of Kleborate's extensive data output can be guided by the accompanying web-based

visualization app, Kleborate-Viz. Through this app, many of the analyses and plots

presented in this manuscript can be rapidly replicated, and further explored in an

interactive manner.

Kleborate is designed to facilitate detection and tracking of clinically relevant AMR

and virulence traits from genome data, and analysis of public data not only identified

specific clones and genes associated with one or the other of these traits (**Figures 5,**

6), but also 601 genomes in which the two converge (carrying *iuc+* virulence

plasmids and ESBL and/or carbapenemase genes; **Figure 7**). We estimated at least

173 unique AMR-hypervirulence convergence events; the majority were detected

within a single isolate (n=119 events), but many others appear to be associated with

local outbreaks or larger-scale spread and apparently across multiple countries (**Table**

S8). Some of the convergence events in China and other countries in the neighbouring

South and Southeast Asia regions have been extensively reported[16,20,49,56], but to our

knowledge a significant number had not been recognized previously. These include

ST231-MDR (most with OXA-232, remainder with ESBL only) + *iuc*, which has

been reportedly circulating in India[49], and our analysis also detected in Pakistan,

Thailand, Switzerland and USA.

Kleborate has already been widely adopted by the *Klebsiella* research community – at

least 74 studies have reported using the Kleborate software package, including larger-

scale genome surveillance studies in South and Southeast Asia, the Caribbean and the

United States[31,49,50] (full list in **Table S11**). Kleborate is freely available as a

standalone command-line tool for local high-throughput analyses or incorporation

into existing bioinformatics workflows (https://github.com/katholt/Kleborate), and

can be easily accessed through the online tool PathogenWatch

(https://pathogen.watch/). With such broad accessibility and utility, Kleborate is

poised to become a cornerstone of the *Klebsiella* genomic surveillance toolkit that can

help inform containment and control strategies targeting this priority pathogen.

METHODS

Kleborate software: implementation and genotyping logic

Kleborate (v.2) is a command-line tool written in Python and is freely available under

the GNU v3.0 license at http://github.com/katholt/Kleborate. It takes as input one or

more whole genome assemblies (FASTA format), types each one against a series of

screening databases outlined in detail below, and returns results in a tab-delimited text

file (one genome per row). On default settings, Kleborate will report assembly quality

metrics, taxonomic assignment, MLST and virulence loci genotypes. Screening for

AMR determinants, and/or K/O serotyping via Kaptive[36], is optional (**Table 1**).

Assembly quality

Assembly quality metrics, reported to help users assess the reliability of genotyping

results, are: contig count, contig N50, largest contig size, total genome size, and

number of ambiguous bases (e.g. 'N'). Low quality warnings are flagged if: (i)

ambiguous bases are detected; (ii) assembly length falls outside the expected range of

4.5-7.5 Mbp; or (iii) N50 is below 10,000 bp. Users should carefully consider the

genotyping outputs for low quality assemblies.

Taxonomic assignment

Kleborate's species prediction function provides a convenient way to confirm species,

including differentiating between the closely related members of the KpSC which are

frequently misclassified using laboratory techniques. Kleborate calculates Mash[40]

distances between the input genome/s and a curated collection of reference assemblies

from different *Klebsiella* and other Enterobacterales, and reports the species with the

smallest distance. Mash distance ≤0.02 is reported as a strong match, ≤0.04 as weak

(only when no strong matches are found, see **Supplementary Text** for further

details).

MLST

Genomes assigned to species in the KpSC are assigned sequence types using

nucleotide BLAST against the established *K. pneumoniae* chromosomal seven-locus

MLST scheme[29] described and maintained on the *K. pneumoniae* BIGSdb site hosted

at the Pasteur Institute (http://bigsdb.pasteur.fr/klebsiella/klebsiella.html).

Virulence gene detection and typing

Virulence loci (*ybt, iuc, iro, clb, rmpADC, rmpA2*) are detected using nucleotide

BLAST search against the database of known alleles. The best hit allele for each gene

(with ≥90% identity and ≥80% coverage) is reported in the main virulence columns.

If the majority of genes expected for the locus are present, then the alleles are used to

calculate STs which are reported along with their associated lineage and MGE (based

on previously defined schemes: YbST for *ybt*, CbST for *clb*, AbST for *iuc*, SmST for

iro, according to the previously defined schemes[34,35]; and a novel RmST scheme for

the *rmpADC* locus). To generate the RmST typing scheme we used the same 2,733

genomes from our original virulence plasmid study[35] to screen and extract the

sequences for *rmpADC* and define allele numbers and STs. These ST sequences

cluster into four distinct lineages associated with distinct MGEs (*rmp1* with KpVP-1,

rmp2 with KpVP-2, *rmp2A* with the *iuc2A* virulence plasmids, and *rmp3* with

ICE*Kp1*; to be described in detail elsewhere). Where the best hit for a gene is a weak

match (80-90% identity, 40-80% coverage) this is reported in the 'spurious hits'

column. Truncations are detected by translating the best-matching nucleotide

sequence for each query gene into amino acids and comparing to the reference length

(expressed as % amino acid length from the start codon, those <90% are reported).

The presence of *ybt*, *clb* and *iuc* are used to assign a virulence score as follows:

0=none present, 1=yersiniabactin only, 2=colibactin without aerobactin (regardless of

yersiniabactin, however *ybt* is almost always present when *clb* is), 3=aerobactin only,

4=aerobactin and yersiniabactin without colibactin, and 5= all three present. The

presence of *iro* (salmochelin) is not used to calculate the virulence score because its

presence is very strongly associated with aerobactin.

Detection and typing of antimicrobial resistance determinants

When AMR detection is switched on, Kleborate screens for known acquired AMR

determinants using a curated version of the CARD AMR nucleotide database (v3.0.8

downloaded February 2020; see doi.org/10.6084/m9.figshare.13256759.v1 for full

details on curation). Genes are identified using nucleotide BLAST (and amino acid

search with tBLASTx if no exact nucleotide match is found). Gene truncations and

spurious hits are identified as described above for virulence genes. Unlike the

acquired forms, the intrinsic variants of *oqxAB,* chromosomal *fosA* and *ampH* are not

associated with clinical resistance in KpSC and are therefore not reported. However,

SHV, LEN or OKP β-lactamase alleles intrinsic to KpSC species are known to confer

clinical resistance to penicillins and are reported in the Bla_chr column. Acquired

SHV variants, and individual SHV sequence mutations known to confer resistance to

extended-spectrum β-lactams or β-lactamase inhibitors, are reported separately (see

Supplementary Text, Tables S12-S13 for details).

Chromosomally encoded mutations and gene loss or truncations known to be

associated with AMR are reported for genomes identified as KpSC species. These

include fluoroquinolone resistance mutations in GyrA (codons 83 and 87) and ParC

(codons 80 and 84), and colistin resistance from truncation or loss of MgrB and PmrB

(defined as <90% amino acid sequence coverage). Mutations in the OmpK35 and

OmpK36 osmoporins reportedly associated with reduced susceptibility to β-

lactamases[41,42] are also screened and reported for KpSC genomes, and include

truncation or loss of these genes and OmpK36GD and OmpK36TD transmembrane β-

strand loop insertions[41]. SHV β-lactamase, GyrA, ParC and OmpK mutations are

identified by alignment of the translated amino acid sequences against a reference

using BioPython, followed by interrogation of the alignment positions of interest (see

Supplementary Text, Tables S12-S13 for a list of relevant positions).

AMR genes and mutations are reported by drug class, with β-lactamases further

categorized by enzyme activity (β-lactamase, ESBL or carbapenemase, with/without

resistance to β-lactamase inhibitors). Horizontally acquired AMR genes are reported

separately from mutational resistance and contribute to the AMR gene count; these

plus chromosomal mutations count towards the number of acquired resistance classes

(intrinsic SHV alleles, reported in Bla_chr column, are not included in either count).

Resistance scores are calculated as follows: 0=no ESBL or carbapenemase, 1=ESBL

without carbapenemase (regardless of colistin resistance); 2=carbapenamase without

colistin resistance (regardless of ESBL); 3=carbapenemase with colistin resistance

(regardless of ESBL).

Serotype prediction

By default, genomes are screened against the *wzi* database in the *Klebsiella* BIGSdb

(using nucleotide BLAST) which is used to predict capsule (K) type based on a

previously defined scheme[60]. This allows rapid typing however the relationship

between *wzi* allele and K type is not one-to-one[36] . If surface antigen prediction is

important to users they can obtain more robust identification of K and O antigen

(LPS) loci by switching on serotype prediction with Kaptive[36] (--kaptive), which adds

a few minutes per genome to Kleborate's runtime.

Data visualization

To facilitate interpretation of Kleborate's rich data output we provide a web-based

application (Kleborate-Viz, https://kleborate.erc.monash.edu/), implemented in R

Shiny, which takes as inputs a Kleborate results file (required), sample metadata

(CSV format, optional) and MIC data (CSV format, optional).

Genome analysis

The analyses reported here result from applying Kleborate v2.0.0 to publicly available

genome collections. A total of 13,156 *Klebsiella* WGS assemblies, encompassing

non-duplicate isolates with unique BioSample accessions identified from published

studies (some deposited as read sets only, which were assembled using Unicycler

v0.4.7[61], data sources summarized in **Table S14**) plus any additional genomes

designated as *Klebsiella* in NCBI's RefSeq repository of genome assemblies (as of

17th July 2020). In order to minimize the impact of sampling bias favouring common

MDR and/or virulent lineages and those causing outbreaks, we subsampled the

collection into a 'non-redundant' dataset of 11,277 genomes (9,705 *K. pneumoniae*)

as follows. Pairwise Mash distances were calculated using Mash v2.1, and used to

cluster genomes using single-linkage clustering with a threshold of 0.0003. These

clusters were further divided into non-redundant groups with unique combinations of

(i) Mash cluster, (ii) chromosomal ST, (iii) virulence gene profiles (i.e. presence of

ybt/clb/iro/iuc loci and lineage assignment), (iv) AMR profiles, (v) year and country

of isolation, and (vii) specimen type where available. For each resulting non-

redundant group, one genome was selected at random as the representative for

analyses. The full list of genomes, including database accessions, isolate information,

cluster/group assignment, and Kleborate results are provided in **Table S2**. The subset

of 1,624 *K. pneumoniae* assemblies deposited in RefSeq by the European EuSCAPE

surveillance study[33] (out of 1,649 reported in original study; **Table S2**) were used for

the EuSCAPE analyses reported in **Figures 2** and **3**. The Kleborate-Viz web

application is pre-loaded with the non-redundant and EuSCAPE WGS datasets

reported in this paper, and can be used to reproduce the plots shown in **Figures 1A-C,**

2B-C, 3, 6A-B and to further explore the Kleborate results.

Metagenome analysis

We downloaded metagenomic reads, and matched isolate WGS assemblies, for n=47

infant gut microbiota samples deposited by the Baby Biome Study[59]. Metagenome

reads were assembled using SPAdes version 3.13.1[62] with the --meta flag and the

resulting contigs binned using MaxBin v2.2.7[63]. Seven metagenomes failed to

assemble due to memory and compute walltime constraints, hence we report results

for 40 samples (**Table S10**). Kleborate was run separately on the full metagenome

assemblies, all contig bins (from which the *Klebsiella* bin could then be identified),

and the matched WGS assemblies. Metagenomic read sets were also analysed using

Kracken 2.0.7[64] and Bracken v2.5[65] (with a custom GDTB release 89 database[66]) to

estimate the relative abundance of KpSC reads in each metagenome.

Statistical analysis

Statistical analyses and data visualisations were conducted using R v1.1.456. Figures

were generated with ggplot v3.2.0 and pheatmap v1.0.12. Correlations between

virulence and resistance scores, and the prevalence of virulence and resistance

determinants over time, were analysed using Spearman's rank-order correlation (i.e.

non-parametric test).

Acknowledgements

We thank Prof Sylvain Brisse and the curators of the *K. pneumoniae* BIGSdb at

Institut Pasteur for hosting and maintaining the MLST schemes

(https://bigsdb.pasteur.fr/klebsiella/klebsiella.html); and Prof David Aanensen and

team at the Centre for Genomic Pathogen Surveillance for making Kleborate available

online within Pathogenwatch (http://pathogen.watch).

Funding

This work was supported by the Bill and Melinda Gates Foundation of Seattle (grant

OPP1175797 to K.E.H.) and the Viertel Charitable Foundation of Australia (Senior

Medical Research Fellowship to K.E.H.). K.L.W is supported by the National Health

and Medical Research Council of Australia (Investigator Grant APP1176192).

References

1.   World Health Organisation. *Global Priority List of Antibiotic-Resistant*

Bacteria to Guide Research, Discovery, and Development of New Antibiotics.

(2017).

2.   Wyres, K. L. & Holt, K. E. *Klebsiella pneumoniae* as a key trafficker of drug

resistance genes from environmental to clinically important bacteria. *Curr.*

Opin. Microbiol. **45**, 131–139 (2018).

3.   Gorrie, C. L. *et al.* Gastrointestinal carriage is a major reservoir of *K.*

pneumoniae infection in intensive care patients. *Clin Infect Dis* **65**, 208–215

(2017).

4.   Martin, R. M. *et al.* Molecular epidemiology of colonizing and infecting

isolates of *Klebsiella pneumoniae. mSphere* **1**, (2016).

5.   Chung, D. R. *et al.* Fecal carriage of serotype K1 *Klebsiella pneumoniae* ST23

strains closely related to liver abscess isolates in Koreans living in Korea. *Eur J*

Clin Microbiol Infect Dis **31**, 481–486 (2012).

6.   Lin, Y.-T. *et al.* Seroepidemiology of *Klebsiella pneumoniae* colonizing the

intestinal tract of healthy chinese and overseas chinese adults in Asian

countries. *BMC Microbiol.* **12**, 13 (2012).

7.   Kaur, C. P., Vadivelu, J. & Chandramathi, S. Impact of *Klebsiella pneumoniae*

in lower gastrointestinal tract diseases. *J. Dig. Dis.* **19**, 262–271 (2018).

8.   Podschun, R. & Ullmann, U. *Klebsiella* spp. as Nosocomial Pathogens:

Epidemiology, Taxonomy, Typing Methods, and Pathogenicity Factors. *Clin*

Microbiol Rev **11**, 589–603 (1998).

9.   Petrosillo, N., Taglietti, F. & Granata, G. Treatment Options for Colistin

Resistant *Klebsiella pneumoniae*: Present and Future. *J. Clin. Med.* **8**, 934

(2019).

10.   Tooke, C. L. *et al.* β-Lactamases and β-Lactamase Inhibitors in the 21st

Century. *J. Mol. Biol.* **431**, 3472–3500 (2019).

11.   Geneva: World Health Organization. *Prioritization of Pathogens To Guide*

Discovery, Research and Development of New Antibiotics for Drug-Resistant

Bacterial Infections, Including Tuberculosis. (2017).

12.   Shon, A. S., Bajwa, R. P. S. & Russo, T. A. Hypervirulent (hypermucoviscous)

Klebsiella pneumoniae: a new and dangerous breed. *Virulence* **4**, 107–118

(2013).

13.   Siu, L. K., Yeh, K., Lin, J., Fung, C. & Chang, F. *Klebsiella pneumoniae* liver

abscess: a new invasive syndrome. *Lancet Infect Dis* **12**, 881–887 (2012).

14.   Wyres, K. L. *et al.* Distinct evolutionary dynamics of horizontal gene transfer

in drug resistant and virulent clones of *Klebsiella pneumoniae*. *PLoS Genet.* **15**,

e1008114 (2019).

15.   Brisse, S. *et al.* Virulent clones of *Klebsiella pneumoniae*: Identification and

evolutionary scenario based on genomic and phenotypic characterization. *PLoS*

One **4**, (2009).

16.   Wyres, K. L., Lam, M. M. C. & Holt, K. E. Population genomics of *Klebsiella*

pneumoniae. *Nat. Rev. Microbiol.* **18**, 344–359 (2020).

17.   Walker, K. A. *et al.* A *Klebsiella pneumoniae* Regulatory Mutant Has Reduced

Capsule Expression but Retains Hypermucoviscosity. *MBio* **10**, e00089-19

(2019).

18.   Walker, K. A., Treat, L. P., Sepúlveda, V. E. & Miller, V. L. The Small Protein

RmpD Drives Hypermucoviscosity in *Klebsiella pneumoniae*. *MBio* **11**,

e01750-20 (2020).

19. Holt, K. E. *et al.* Genomic analysis of diversity, population structure, virulence,

and antimicrobial resistance in *Klebsiella pneumoniae*, an urgent threat to

public health. *Proc Natl Acad Sci USA* **112**, E3574--81 (2015).

20. Gu, D. *et al.* A fatal outbreak of ST11 carbapenem-resistant hypervirulent

Klebsiella pneumoniae in a Chinese hospital: a molecular epidemiological

study. *Lancet Infect Dis* **18**, 37–46 (2018).

21. Xu, M. *et al.* High prevalence of KPC-2-producing hypervirulent *Klebsiella*

pneumoniae causing meningitis in Eastern China. *Infect. Drug Resist.* **12**, 641–

653 (2019).

22. Dong, N. *et al.* Genome analysis of clinical multilocus sequence Type 11

Klebsiella pneumoniae from China. *Microb. genomics* **4**, e000149 (2018).

23. Wong, M. H. Y. *et al.* Emergence of carbapenem-resistant hypervirulent

Klebsiella pneumoniae. *Lancet Infect Dis* **18**, (2018).

24. Yao, H., Qin, S., Chen, S., Shen, J. & Du, X.-D. Emergence of carbapenem-

resistant hypervirulent *Klebsiella pneumoniae*. *Lancet Infect Dis* **18**, (2018).

25. Ørskov, I. D. A. & Fife-Asbury, M. A. New *Klebsiella* capsular antigen, K82,

and the deletion of five of those previously assigned. *Int J Syst Bacteriol.* **27**,

386–387 (1977).

26. Trautmann, M. *et al.* O-antigen seroepidemiology of *Klebsiella* clinical isolates

and implications for immunoprophylaxis of *Klebsiella* infections. *Clin. Diagn.*

Lab. Immunol. **4**, 550–555 (1997).

27. Elhani, D. *et al.* Molecular epidemiology of extended-spectrum beta-lactamase-

producing *Klebsiella pneumoniae* strains in a university hospital in Tunis,

Tunisia, 1999&-2005. *Clin. Microbiol. Infect.* **16**, 157–164 (2010).

28. Chen, L. *et al.* Carbapenemase-producing *Klebsiella pneumoniae*: molecular

and genetic decoding. *Trends Microbiol.* **22**, 686–696 (2014).

29.   Diancourt, L., Passet, V., Verhoef, J., Grimont, P. A. & Brisse, S. Multilocus

sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin*

Microbiol. **43**, 4178–4182 (2005).

30.   Wyres, K. L. & Holt, K. E. *Klebsiella pneumoniae* Population Genomics and

Antimicrobial-Resistant Clones. *Trend Microbiol* **24**, 944–956 (2016).

31.   Long, S. W. *et al.* Population Genomic Analysis of 1,777 Extended-Spectrum

Beta-Lactamase-Producing *Klebsiella pneumoniae* Isolates, Houston, Texas:

Unexpected Abundance of Clonal Group 307. *MBio* **8**, e00489--17 (2017).

32.   Potter, R. F. *et al.* Population Structure, Antibiotic Resistance, and

Uropathogenicity of *Klebsiella variicola*. *MBio* **9**, e02481-18 (2018).

33.   David, S. *et al.* Epidemic of carbapenem-resistant *Klebsiella pneumoniae* in

Europe is driven by nosocomial spread. *Nat. Microbiol.* (2019).

doi:10.1038/s41564-019-0492-8

34.   Lam, M. M. C. *et al.* Genetic diversity, mobilisation and spread of the

yersiniabactin-encoding mobile element ICE*Kp* in *Klebsiella pneumoniae*

populations. *Microb Genom* **Jul 9**, (2018).

35.   Lam, M. C. C. *et al.* Tracking key virulence loci encoding aerobactin and

salmochelin siderophore synthesis in *Klebsiella pneumoniae*. *Genome Med.* **10**,

77 (2018).

36.   Wick, R. R., Heinz, E., Holt, K. E. & Wyres, K. L. Kaptive Web: user-friendly

capsule and lipopolysaccharide serotype prediction for *Klebsiella* genomes. *J*

Clin Microbiol. **56**, e00197-18 (2018).

37.   Martínez-Romero, E. *et al.* Genome misclassification of *Klebsiella variicola*

and Klebsiella quasipneumoniae isolated from plants, animals and humans.

Salud Publica Mex **60**, 52–62 (2018).

38.   Rodrigues, C. *et al.* Description of *Klebsiella africanensis* sp. nov., *Klebsiella*

variicola subsp. *tropicalensis* subsp. nov. and *Klebsiella variicola* subsp.

variicola subsp. nov. *Res. Microbiol.* **S0923-2508**, 30019–1 (2019).

39.   Long, S. W. *et al.* Whole-genome sequencing of a human clinical isolate of the

novel species *Klebsiella quasivariicola* sp. nov. *Genome Announc.* **5**, e01057-

17 (2017).

40.   Ondov, B. D. *et al.* Mash: Fast genome and metagenome distance estimation

using MinHash. *Genome Biol.* **17**, 132 (2016).

41.   Fajardo-Lubia´n, A., Ben Zakour, N. L., Agyekum, A., Qi, Q. & Iredell, J. R.

Host adaptation and convergent evolution increases antibiotic resistance

without loss of virulence in a major human pathogen. *PLoS Pathog.* **15**,

e1007218 (2019).

42.   Wong, J. L. C. *et al.* OmpK36-mediated Carbapenem resistance attenuates

ST258 *Klebsiella pneumoniae in vivo*. *Nat. Commun.* **10**, 3957 (2019).

43.   Hauck, C. *et al.* Spectrum of excess mortality due to carbapenem-resistant

Klebsiella pneumoniae infections. *Clin. Microbiol. Infect.* **22**, 513–519 (2016).

44.   Opoku-Temeng, C., Kobayashi, S. D. & DeLeo, F. R. *Klebsiella pneumoniae*

capsule polysaccharide as a target for therapeutics and vaccines. *Comput.*

Struct. Biotechnol. J. **17**, 1360–1366 (2019).

45.   Venturini, C. *et al.* Fine capsule variation affects bacteriophage susceptibility

in *Klebsiella pneumoniae* ST258. *FASEB J.* **34**, 10801–10817 (2020).

46.   Arena, F. *et al.* Population structure of KPC carbapenemase-producing

Klebsiella pneumoniae in a long-term acute-care rehabilitation facility:

identification of a new lineage of clonal group 101, associated with local

hyperendemicity. *Microb. genomics* **6**, e000308 (2020).

47.   Ferrari, C. *et al.* Multiple *Klebsiella pneumoniae* KPC Clones Contribute to an

Extended Hospital Outbreak. *Front. Microbiol.* **10**, 2767 (2019).

48.   Magiorakos, A.-P. *et al.* Multidrug-resistant, extensively drug-resistant and

pandrug-resistant bacteria: an international expert proposal for interim standard

definitions for acquired resistance. *Clin. Microbiol. Infect.* **18**, 268–281 (2012).

49.   Wyres, K. L. *et al.* Genomic surveillance for hypervirulence and multi-drug

resistance in invasive *Klebsiella pneumoniae* from South and Southeast Asia.

Genome Med. **12**, 11 (2020).

50.   Heinz, E., Brindle, R., Morgan-McCalla, A., Peters, K. & Thomson, N. R.

Caribbean multi-centre study of *Klebsiella pneumoniae*: whole genome

sequencing, antimicrobial resistance and virulence factors. *bioRxiv* (2019).

doi:10.1101/541136

51.   Musicha, P. *et al.* Genomic analysis of *Klebsiella pneumoniae* isolates from

Malawi reveals acquisition of multiple ESBL determinants across diverse

lineages. *J. Antimicrob. Chemother.* **74**, 1223–1232 (2019).

52.   Alvarez-Uria, G., Gandra, S., Mandal, S. & Laxminarayan, R. Global forecast

of antimicrobial resistance in invasive isolates of Escherichia coli and

Klebsiella pneumoniae. Int. J. Infect. Dis. **68**, 50–53 (2018).

53.   Brolund, A. *et al.* Worsening epidemiological situation of carbapenemase-

producing Enterobacteriaceae in Europe, assessment by national experts from

37 countries, July 2018. *Eurosurveillance* **24**, (2019).

54.   Bell, J. *et al. Gram-negative Sepsis Outcome Program 2019 Report.* (2019).

55.   Yu, W.-L., Lee, M.-F., Tang, H.-J., Chang, M.-C. & Chuang, Y.-C. Low

prevalence of *rmpA* and high tendency of *rmpA* mutation correspond to low

virulence of extended spectrum β-lactamase-producing *Klebsiella pneumoniae*

isolates. *Virulence* **6**, 162–172 (2015).

56.   Chen, L. & Kreiswirth, B. N. Convergence of carbapenem-resistance and

hypervirulence in *Klebsiella pneumoniae. Lancet Infect Dis* **18**, 2–3 (2018).

57.   Lam, M. M. C. *et al.* Convergence of virulence and multidrug resistance in a

882   single plasmid vector in multidrug-resistant *Klebsiella pneumoniae* ST15. *J*

883   *Antimicrob Chemother.* (2019). doi:https://doi.org/10.1093/jac/dkz028

884   58.   Chen, Y. *et al.* Preterm infants harbour diverse *Klebsiella* populations,

885   including atypical species that encode and produce an array of antimicrobial

886   resistance- and virulence-associated factors. *Microb. Genomics* **6**, (2020).

887   59.   Shao, Y. *et al.* Stunted microbiota and opportunistic pathogen colonization in

888   caesarean-section birth. *Nature* **574**, 117–121 (2019).

889   60.   Brisse, S. *et al. wzi* Gene Sequencing, a Rapid Method for Determination of

890   Capsular Type for *Klebsiella* Strains. *J. Clin. Microbiol.* **51**, 4073–4078

891   (2013).

892   61.   Wick, R. R., Judd, L. M., Gorrie, C. & Holt, K. E. Unicycler: Resolving

893   bacterial genome assemblies from short and long sequencing reads. *PLoS*

894   *Comput Biol* **13**, e1005595 (2017).

895   62.   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a

896   new versatile metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

897   63.   Wu, Y.-W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated

898   binning algorithm to recover genomes from multiple metagenomic datasets.

899   *Bioinformatics* **32**, 605–607 (2016).

900   64.   Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with

901   Kraken 2. *Genome Biol.* **20**, 257 (2019).

902   65.   Lu, J., Bretwieser, F. P., Thielen, P. & Salzberg, S. L. Bracken: estimating

903   species abundance in metagenomics data. *PeerJ Comput. Sci.* **3**, (2017).

904   66.   Parks, D. H. *et al.* A complete domain-to-species taxonomy for Bacteria and

905   Archaea. *Nat. Biotechnol.* **38**, 1079–1086 (2020).

67.   Bialek-davenet, S. *et al.* Genomic Definition of Hypervirulent and Multidrug-

Resistant *Klebsiella pneumoniae* Clonal Groups. *Emerg. Infect. Dis.* **20**, 1812–

1820 (2014).

68.   Neubauer, S. *et al.* A Genotype-Phenotype Correlation Study of SHV β-

Lactamases Offers New Insight into SHV Resistance Profiles. *Antimicrob.*

Agents Chemother. **64**, e02293-19 (2020).

69.   Kidd, T. J. *et al.* Molecular mechanisms and virulence of colistin-resistant

Klebsiella pneumoniae. Eur. Respir. J. **48**, PA2625 (2016).

70.   Cannatelli, A. *et al.* In vivo evolution to colistin resistance by PmrB sensor

kinase mutation in KPC-producing *Klebsiella pneumoniae* is associated with

low-dosage colistin treatment. *Antimicrob. Agents Chemother.* **58**, 4399–4403

(2014).

71.   Cannatelli, A. *et al.* MgrB Inactivation Is a Common Mechanism of Colistin

Resistance in KPC-Producing *Klebsiella pneumoniae* of Clinical Origin.

Antimicrob. Agents Chemother. **58**, 5696 LP – 5703 (2014).

72.   Drlica, K. & Zhao, X. DNA gyrase, topoisomerase IV, and the 4-quinolones.

Microbiol. Mol. Biol. Rev. **61**, 377–392 (1997).

Author Contributions

Study design: K.E.H. Data analysis: M.M.C.L., K.L.W. and K.E.H. Code

development: R.R.W., K.E.H., S.C.W., L.T.C., and K.L.W. Manuscript writing:

M.M.C.L., K.L.W., and K.E.H. All authors contributed to manuscript editing.

Competing Interests

None declared.

Figure 1. Relationships between Kleborate virulence and resistance scores and the prevalence of key virulence and antimicrobial resistance (AMR). Data shown summarise Kleborate results for non-redundant set of 9,705 publicly available *K. pneumoniae* genomes (**Table S2**). **(A)** Mean virulence score (barplot, right y-axis) and prevalence of individual virulence loci (line plots, left y-axis) over time. Ybt, yersiniabactin; clb, colibactin; iuc, aerobactin; iro, salmochelin; rmpADC, hypermucoidy *rmp* locus; rmpA2, *rmpA2* gene. Correlations between mean virulence score and prevalence of each locus are noted. **(B)** Mean resistance score (barplot, right y-axis) and prevalence of carbapenemases (carb), acquired colistin resistance genes (col gene), mutations in MgrB/PmrB (col mut) and genes conferring resistance to extended-spectrum β-lactams (ESBL) (line plots, left y-axis).

Correlations between mean resistance score and prevalence of each resistance type are noted. **(C)** Mean number of acquired AMR genes and classes, over time. Correlations with mean resistance score are noted. **(D)** Histograms showing total number of acquired AMR classes predicted per genome, stratified by resistance score. **(E)** Histograms showing total number of acquired AMR genes detected per genome, stratified by resistance score. Correlations reported in **A-C** are Spearman rank correlations; significance levels are indicated with asterisks: *p<0.01, **p<0.001.

Figure 2. Kleborate genotyping results for European *K. pneumoniae* surveillance isolates. Data shown summarise Kleborate results for 927 carbapenem-non-susceptible and 697 carbapenem-susceptible *K. pneumoniae* genomes from the EuSCAPE study (data included in **Table S2**). **(A)** Geographical and lineage distribution of carbapenemase genes. Each circle represents a genome, colored by carbapenemase (see inset legend). Barplots summarise the number of genomes from each *K. pneumoniae* lineage (top) and country (right), colored by carbapenemase. **(B-C)** Cumulative prevalence of **(B)** capsule (K) locus

and **(C)** O antigen locus types, for carbapenem non-susceptible (meropenem MIC>2) isolates, ordered by overall prevalence. Thick line indicates curve for whole data set; others give results separately for different United Nations geographical regions (see inset legend).

Figure 3. Distribution of meropenem MIC, stratified by Kleborate-detected carbapenemase genes and OmpK35/36 porin mutations, for European *K. pneumoniae* surveillance isolates. Data shown summarise Kleborate results for 1,490 *K. pneumoniae* genomes from the EuSCAPE study (data included in **Table S2**). Each circle represents the reported meropenem MIC for an isolate, coloured by type of porin mutation/s identified by Kleborate from the corresponding genome assembly (colour key in inset legend, prevalence of each genotype across 1490 genomes is indicated in brackets). Isolates are stratified by carbapenemase gene (enzymes labelled on x-axis) and OmpK mutations[41,42] reported by Kleborate. Wt, full-length OmpK35 and OmpK36 with no GD/TD insertion in the OmpK36 β-strand loop; mut, otherwise; trunc, truncation. Dashed lines indicate EUCAST breakpoints for clinical resistance (red, MIC >8) and non-susceptibility (black, MIC >2).

Figure 4. Summary of genome collection metadata, and Kleborate-derived virulence and antimicrobial resistance (AMR) genotypes, for all publicly available *Klebsiella* genomes. Data shown summarise Kleborate results for 11,277 non-redundant *Klebsiella* genomes publicly available as at July 17, 2020 (**Table S2**). From left to right: barplots showing source information by geographical region and sample type (coloured as per inset legend); heatmaps showing prevalence of virulence loci (blue) and predicted AMR drug classes (red) (as per inset scale bars). Genomes are summarised by species, ordered by species complex: KpSC, *K. pneumoniae* species complex; KoSC, *K. oxytoca* species complex; and other *Klebsiella*. In the heatmaps, the total number of genomes in which each type of virulence/AMR determinant was detected are indicated below each column. Column names are as follows: ybt, yersiniabactin; clb, colibactin; iuc, aerobactin; iro, salmochelin; rmp, hypermucoidy Rmp; rmpA2, hypermucoidy rmpA2; AGly, aminoglycosides; Col, colistin; Fcyn, fosfomycin; Flq, fluoroquinolone; Gly, glycopeptide; MLS, macrolides; Phe, phenicols; Rif, rifampin; Sul, sulfonamides; Tet, tetracyclines; Tgc, tigecycline; Tmt, trimethoprim; Bla, β-lactamases; inhR, β-lactamase inhibitor; ESBL, extended-spectrum β-lactamases; ESBL_inhR, extended-spectrum β-lactamase with resistance to β-lactamase inhibitors; Carb, carbapenemase; Bla_chr, intrinsic chromosomal β-lactamase; SHV,

mutations in SHV; Omp, truncations/mutations in *ompK35/ompK36*; Col, truncations in *mgrB/pmrB* conferring colistin resistance; Flq, mutations in *gyrA/parC* conferring resistance to fluoroquinolones.

Figure 5. Distribution of resistance and virulence scores among genomes belonging to the 30 most common _K. pneumoniae_ lineages. Data shown summarise Kleborate results for non-redundant set of 9,705 publicly available _K. pneumoniae_ genomes (**Table S2**). Lineages were defined on the basis of multi-locus sequence types (STs) reported by Kleborate, and ordered from highest to lowest difference between mean virulence and mean resistance score. Minimum genome count per ST shown is 50. Ybt, yersiniabactin; clb, colibactin; iuc, aerobactin; VP, virulence plasmid; ESBL, extended-spectrum β-lactamase; Carb, carbapenemase; Col, colistin resistance determinant

Figure 6. Insights from resistance and virulence scores. Data shown summarise Kleborate results for non-redundant set of 9,705 publicly available *K. pneumoniae* genomes (**Table S2**). (**A-B**) Mean resistance and virulence scores grouped by (**A**) lineage and (**B**) sample type. Each circle represents a single lineage (multi-locus sequence type, ST) or sample type as labelled; size indicates the number of genomes (as per inset legend); colour indicates groups per inset legend. (**C**) Heatmap showing number of genomes with each combination of resistance and virulence scores. Convergent genomes correspond to a virulence score ≥3 (carrying *iuc*) and resistance score of ≥1 (carrying ESBL and/or carbapenemase gene/s), as indicated by the red box. (**D**) Barplots showing lineage distribution of convergent genomes,

for each combination of resistance score and virulence score. Lineages are grouped into hypervirulent (hv), multidrug resistant (MDR) and other; and coloured by ST (as per inset legend).

Figure 7. Convergence of AMR and virulence determinants in the *K. pneumoniae* population, identified by Kleborate analysis of public genomes. (A) Geographical and lineage distribution of convergence events. Each circle represents a unique convergence event (i.e. a monophyletic clade harbouring both ESBL/carbapenemase genes and *iuc*; see interactive tree at https://microreact.org/project/JDyan46yctyDh6weEUjWN, summary of events in **Table S8**, assignment of genomes to events in **Table S2**). Circles are scaled by the number of total genomes linked to the event and colored to indicate whether convergence is inferred to have occurred via acquisition of AMR gene/s (ESBL or carbapenemase/s) by a hypervirulent lineage or via acquisition of an *iuc*-encoding plasmid by an AMR lineage, as per inset legend. Marginal barplots show the number of convergence events (color blocks) and genomes (block heights) associated with each lineage (top) or geographical region (right). Lineages were defined on the basis of multi-locus sequence types (STs), number of convergence events estimated for each is labelled at the top of each bar. **(B)** Distribution of convergent genomes by location. Countries from which convergent genomes were detected are colored on the map; circles represent the number of convergent genomes detected in each UN-defined geographical region (indicated by color, as per inset legend), scaled and labelled

with the minimum estimated number of unique convergence events specific to each region (excluding inter-regional convergence events). The total number of convergence events affecting each region, including region-specific and inter-regional convergence events, are given in brackets in the inset legend.

Table 1. Genome features reported by Kleborate

Feature	Description
Assembly quality	Contig count, N50, largest contig, ambiguous bases
Identification	Species[16], MLST[29,67] (if *K. pneumoniae* species complex)
Acquired virulence determinants	Presence, genotypes, associated MGEs, truncations • yersiniabactin[34], • colibactin[34], • aerobactin[35], • salmochelin[35], • hypermucoidy loci *rmpADC* and *rmpA2*
Virulence score	0=no yersinabactin, colibactin or aerobactin; 1=yersiniabactin only; 2=yersiniabactin and colibactin (or colibactin only); 3= aerobactin without yersiniabactin or colibactin; 4= aerobactin with yersiniabactin (no colibactin); 5=yersiniabactin, colibactin and aerobactin
Serotype prediction	*wzi* allele and associated K locus[60] (default), Full K and O locus typing via *Kaptive*[36] (optional)
AMR determinants (optional)	
Acquired genes	Total count, alleles grouped by drug class, truncations
Mutations in core genes	SHV beta-lactamase (ESBL or inhibitors)[68], OmpK35/OmpK36 osmoporins[41,42] (carbapenems), MgrB/PmrB[69–71] (colistin), GyrA/ParC[72] (fluoroquinolones)

Number of drug classes	Excludes penicillins since resistance is intrinsic
Resistance score	1=ESBL; 2=Carbapenemase; 3=Carbapenemase plus colistin resistance; 0 otherwise

Table 2. Prevalence of virulence loci, ESBL and carbapenemase genes in non-redundant *Klebsiella* genomes

Species Complex	Species	Total no. genomes	Virulence prevalence	ESBL prevalence	Carbapenemase prevalence
K. pneumoniae species complex	*K. pneumoniae*	9705	Ybt: 4309, 44% Clb: 794, 8% Iuc: 1090, 11% Iro: 683, 7% Rmp: 782, 8% RmpA2: 716, 7%	4634, 48%	4173, 43%
	K. quasipneumoniae subsp. *quasipneumoniae*	119	-	31, 26%	29, 24%
	K. quasipneumoniae subsp. *similipneumoniae*	363	Ybt: 8, 2% Iuc: 6, 2% Iro: 4, 1% Rmp: 4, 1% RmpA2: 3, 0.8%	138, 38%	32, 9%
	K. quasivariicola	16	-	3, 19%	-
	K. africana	1	-	-	-
	K. variicola subsp. *variicola*	498	Ybt: 15, 3% Iuc: 4, 0.8% Iro: 5, 1% Rmp: 4, 0.8% RmpA2: 2, 0.4%	52, 10%	36, 7%
	K. variicola subsp. *tropica*	18	-	2, 11%	1, 6%
K. oxytoca species complex	*K. oxytoca*	98	Ybt: 96, 98%	9*, 9%	6, 6%
	K. grimontii	75	Ybt: 41, 55%	1*, 1%	3, 4%
	K. huaxiensis	4	-	1*, 25%	-
	K. michiganensis	144	Ybt: 102, 71% Clb: 1, 0.7%	21*, 15%	33, 23%
	K. pasteurii	21	Ybt: 21, 100%	1*, 5%	-

	K. spallanzanii	4	-	-*	-
NA	_K. aerogenes_	209	Ybt: 101, 48% Clb: 95, 45% Iro: 190, 91%	24, 11%	34, 16%
	K. indica	2	-	-	-

*excluding OXY genes that are conserved in _K. oxytoca_ species complex