Recurrent emergence and transmission of a SARS-CoV-2 Spike deletion H69/V70

Kemp SA[1,2,3*], Harvey WT[4*], Lytras S[5], The COVID-19 Genomics UK (COG-UK) consortium[6],

Carabelli AM[3], Robertson DL[5*], Gupta RK[2,3*]

[1]Division of Infection and Immunity, University College London, London, UK.

[2]Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), Cambridge,

UK.

[3]Department of Medicine, University of Cambridge, Cambridge, UK.

[4]Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,

Glasgow, UK

[5]MRC - University of Glasgow Centre for Virus Research, Glasgow, UK.

[6]https://www.cogconsortium.uk. Full list of consortium names and affiliations are in

Appendix 1.

*Authors contributed equally to this work

Address for correspondence:

Ravindra K. Gupta

Cambridge Institute for Therapeutic Immunology and Infectious Diseases

Jeffrey Cheah Biomedical Centre

Puddicombe Way

Cambridge CB2 0AW, UK

Tel: +44 1223 331491

rkg20@cam.ac.uk

Abstract

SARS-CoV-2 Spike amino acid replacements in the receptor binding domain (RBD) occur

relatively frequently and some have a consequence for immune recognition. Here we report

recurrent emergence and significant onward transmission of a six-nucleotide deletion in the

S gene, which results in loss of two amino acids: H69 and V70. Of particular note this

deletion, ΔH69/V70, often co-occurs with the receptor binding motif amino acid

replacements N501Y, N439K and Y453F. One of the ΔH69/V70+ N501Y lineages, B.1.1.7, is

comprised of over 4000 SARS-CoV-2 genome sequences from the UK and includes eight

other S gene mutations: RBD (N501Y and A570D), S1 (ΔH69/V70 and Δ144/145) and S2

(P681H, T716I, S982A and D1118H). Some of these mutations have presumably arisen as a

result of the virus evolving from immune selection pressure in infected individuals and at

least one, lineage B.1.1.7, potentially from a chronic infection. Given our recent evidence

that ΔH69/V70 enhances viral infectivity (Kemp et al. 2020), its effect on virus fitness

appears to be independent to the RBD changes. Enhanced surveillance for the ΔH69/V70

deletion with and without RBD mutations should be considered as a priority. Permissive

mutations such as ΔH69/V70 have the potential to enhance the ability of SARS-CoV-2 to

generate new variants, including vaccine escape variants, that would have otherwise

significantly reduced viral infectivity.

Background

SARS-CoV-2's Spike surface glycoprotein engagement of hACE2 is essential for virus entry

and infection[1], and the receptor is found in respiratory and gastrointestinal tracts[2]. Despite

this critical interaction and the constraints it imposes, it appears the RBD, and particularly

the receptor binding motif (RBM), is relatively tolerant to mutations[3,4], raising the real

possibility of virus escape from vaccine-induced immunity and monoclonal antibody

treatments. Spike mutants exhibiting reduced susceptibility to monoclonal antibodies have

been identified in *in vitro* screens[3,5,6], and some of these mutations have been found in

clinical isolates[7]. Due to the susceptibility of the human population to this virus, the acute

nature of infections and limited use of vaccines to date there has been limited selection

pressure placed SARS-CoV-2[8]; as a consequence few mutations that could alter antigenicity

have increased significantly in frequency.

The unprecedented scale of whole genome SARS-CoV-2 sequencing has enabled

identification and epidemiological analysis of transmission and surveillance, particularly in

the UK[9] . As of December 18[th,] there were 270,000 SARS-CoV-2 sequences available in the

GISAID Initiative (https:gisaid.org/). However, geographic coverage is very uneven with

some countries sequencing at higher rates than others. This could result in novel variants

with altered biological or antigenic properties evolving and not being detected until they are

already at high frequency.

Studying SARS-CoV-2 chronic infections can give insight into virus evolution that would

require many chains of acute transmission to generate. This is because the majority of

infections arise as a result of early transmission during pre or asymptomatic phases prior to

peak adaptive responses, and virus adaptation not observed as the virus is usually cleared

by the immune response[10,11]. We recently documented *de novo* emergence of antibody

evasion mutations mediated by S gene mutations in an individual treated with convalescent

plasma (CP)[12]. Dramatic changes in the prevalence of Spike variants ΔH69/V70 (an out of

frame six-nucleotide deletion) and D796H variant followed repeated use of CP, while *in vitro*

the mutated ΔH69/V70+D796H variant displayed reduced susceptibility to CP, at the same

time retaining infectivity comparable to wild type[12]. The ΔH69/V70 itself conferred a two-

fold increase in Spike mediated infectivity using pseudotyped lentiviruses. Worryingly, other

deletions in the N-Terminal Domain (NTD) have been reported to arise in chronic infections[7]

and provide escape from NTD-specific neutralising antibodies[13].

Here we analysed the available GISAID Initiative data for circulating SARS-CoV-2 Spike

sequences containing ΔH69/V70 and performed phylogenetic and structural modelling from

the pandemic data and across sarbecoviruses in different species. We find, while occurring

independently, the Spike ΔH69/V70 often emerges after a significant RBM amino acid

replacement that increases binding affinity to hACE2. Protein structure modelling indicates

this mutation could also contribute to antibody evasion as suggested for other NTD

deletions[13].

Results

The deletion H69/V70 is present in over 6000 sequences worldwide, 2.5% of the available data (Figure 1, Supplementary Figure 1), and largely in Europe from where most of the sequences in GISAID are derived (Figure 1C-D). Many of the sequences are from the UK and Denmark where sequencing rates are high compared to other countries. ΔH69/V70 occurs in variants observed in different global lineages, representing multiple independent acquisitions of this SARS-CoV-2 deletion (Figure 1A). While variants with deletions in this region of Spike are observed in GISAID[13], the earliest unambiguous sequence that includes the ΔH69/V70 was detected in Sweden in April 2020, an independent deletion event relative to other ΔH69/V70 variants. The prevalence of ΔH69/V70 has since increased since August 2020 (Figure 1C-D). Further analysis of sequences revealed, firstly, that single deletions of either 69 or 70 were uncommon and, secondly, some lineages of ΔH69/V70 alone were present (Figure 1A), as well as ΔH69/V70 in the context of other mutations in Spike, specifically those in the RBM (Figure 1A, E, F).

To gauge the importance of this part of Spike molecule, we examined the 69/70 region of Spike in a set of other known *Sarbecoviruses*, relatives to SARS-CoV-2 (Figure 2). We observe substantial variability in the region, specifically caused by indels, with some viruses including SARS-CoV having 6-7 amino acid deletions (Fig 2B). This is indicative of plasticity in this protein region that could allow the sarbecoviruses to alter their Spike conformation. The second closest relative to SARS-CoV-2 for this region after RaTG13 is the cluster of 5 CoVs sampled in trafficked pangolins in the Guangxi province[14]. Looking at the 69/70 region in these virus sequences raises the interesting observation that one of the five viruses in the cluster, P1E, has amino acids 69H and 70L present, while the other four have a double amino acid deletion (Figure 2B). Given that SARS-CoV-2 and RaTG13 have the homologous HV insertion at these positions, the most parsimonious explanation is that the proximal common ancestor between SARS-CoV-2 and the Guangxi pangolin cluster had the insertion, which was then lost while circulating in the pangolin population, similar to what we now see with SARS-CoV-2 in humans. Interestingly, the double amino acid deletion in the pangolin viruses in in-frame in contrast to what is seen in SARS-CoV-2 (e.g. lineage B.1.1.7, Figure 2C).

To estimate the structural impact of **ΔH69/V70**, the protein structure of the NTD possessing

the double deletion was modelled. The **ΔH69/V70** deletion was predicted to alter the

conformation of a protruding loop comprising residues 69 to 76, pulling it in towards the

NTD (Figure 3A). In the post-deletion structural model, the positions of the alpha carbons of

residues either side of the deleted residues, Ile68 and Ser71, were each predicted to occupy

positions 2.9Å from the positions of His69 and Val70 in the pre-deletion structure.

Concurrently, the positions of Ser71, Gly72, Thr73, Asn74 and Gly75 are predicted to have

changed by 6.5Å, 6.7Å, 6.0Å, 6.2Å and 8Å, respectively, with the overall effect of these

residues moving inwards, resulting in a less dramatically protruding loop. The position of

this loop in the structure prior to the occurrence of the **ΔH69/V70** is shown in the context of

the wider NTD in Figure 3B. The locations of main RBD mutations observed with **ΔH69/V70**

are shown in Figure 3C and D. Residues belonging to a similarly exposed, nearby loop that

form the epitope of a neutralising, NTD-binding epitope are also highlighted.

We next examined the lineages where S gene mutations in the RBD were identified at high

frequency, in particular co-occurring with N439K (Figure 3C,D), an amino acid replacement

reported to be defining variants increasing in numbers in Europe and other regions[3] (Figure

1E, Supplementary figure 2). N439K appears to have reduced susceptibility to a small subset

of monoclonals targeting the RBD, whilst retaining affinity for ACE2 *in vitro*[3]. The proportion

of viruses with **ΔH69/V70** only increased from August 2020 when it appeared with the

second N439K lineage, B.1.141[3] (Figure 1E). As of November 26[th,] remarkably there were

twice as many cumulative sequences with the deletion as compared to the single N439K

indicating it may be contributing to the success of this lineage (Figure 1E). Due to their high

sampling rates the country with the highest proportion of N439K+**ΔH69/V70** versus N439K

alone is England. The low levels of sequencing in most countries indicate N439K's

prevalence could be relatively high[3]. In Scotland, where early growth of N439K was high

(forming N439K lineage B.1.258 that subsequently went extinct with other lineages after the

lockdown[3]), there is now an inverse relationship with 546 versus 177 sequences for N439K

and N439K+**ΔH69/ΔV70** respectively (Figure 1E). These differences therefore likely reflect

differing epidemic growth characteristics and timings of the introductions the N439K

variants with or without the deletion.

The second significant cluster with ΔH69/V70 and RBD mutants involves Y453F, another

Spike RBD mutation that increases binding affinity to ACE2[4] (Figure 3C,D) and has been

found to be associated with mink-human infections[15]. In one SARS-CoV-2 mink-human sub-

lineage, termed 'Cluster 5', Y453F and ΔH69/V70 occurred with F486L, N501T and M1229I

and was shown to have reduced susceptibility to sera from recovered COVID-19 patients

(https://files.ssi.dk/Mink-cluster-5-short-report_AFO2). Y453F has been described as an

escape mutation for mAb REGN10933[16]. The ΔH69/V70 was first detected in the Y453F

background on August 24[th] and thus far appears limited to Danish sequences

(Supplementary Figure 3).

A third lineage containing the same deletion ΔH69/V70 has arisen with another RBD

mutation N501Y (Figure 4A, C, Supplementary Figure 4). Based on its location it might be

expected to escape antibodies similar to COV2-2499[5] (Figure 3C, D). In addition, when SARS-

CoV-2 was passaged in mice for adaptation purposes for testing vaccine efficacy, N501Y

emerged and increased pathogenicity[17]. Sequences with N501Y alone were isolated both in

the UK, Brazil, and USA in April 2020, and recently in South Africa[18]. A newly described

N501Y-derived lineage in South Africa, (B.1.351, Fig 1A) (also termed 501Y.V2) is

characterised by eight mutations in the Spike protein, including N501Y and two other

important residues in the receptor-binding domain (K417N and E484K) which are important

residues in RBM[18]. The positions of residues 417, 484, and 501 proximal to the bound hACE2

are shown in Figure 4C. The E484K substitution has been identified as antigenically

important being reported as an escape mutation for several monoclonal antibodies

including C121, C144[19], REGN10933 and REGN10934[16]. The increase in hACE2 binding

affinity caused by N501Y may have been permissive for the mutation K417N. Specifically,

the increased ACE2 affinity induced by N501Y (+0.24 $\Delta\log_{10} K_d$) will be compensated by

K417N (-0.45 $\Delta\log_{10} K_d$)[20]. Residue K417 is also identified as antigenically significant with

K417E facilitating escape from mAb REGN10933[16].

N501Y + ΔH69/V70 sequences were first detected in the UK on 20[th] September 2020, with

the cumulative number of N501Y + ΔH69/V70 mutated sequences now exceeding the single

mutant N501Y lineage (Figure 1F). On closer inspection these sequences were part of a new

lineage (B.1.1.7), termed VOC 202012/01 by Public Health England as they are associated

with relatively high numbers of infections (Figure 4A-C, Supplementary Figure 4). In addition

to RBD N501Y + NTD ΔH69/V70 this new variant had five further S mutations across the RBD

(N501Y) and S2 (P681H, T716I, S982A and D1118H), as well as NTD Δ144[21] (Figure 4B). The

variant has now been identified in a number of other countries, including Hong Kong, Japan,

Australia, France, Spain, Singapore, Israel, Switzerland, and Italy. This lineage has a relatively

long branch due to 23 unique mutations (Figure 4A and supplementary figure 4), consistent

within host evolution and spread from a chronically infected individual[21]. Notably a related

sequence can be identified that contains ΔH69/V70, N501Y and D1118H (Figure 4A, black

box). However, the available sequences did not enable us to determine whether the B1.1.7

mutations N501Y + ΔH69/V70 arose as a result of a N501Y virus acquiring ΔH69/V70 or vice

versa.

The B.1.1.7 lineage has some notable features. Firstly the Spike Δ144 mutation could lead to

loss of binding of the S1-binding neutralising antibody 4A8[13] (Figure 3B). The Y144 sidechain

is itself around 4.5Å from the nearest atoms of 4A8 complexed with Spike and the deletion

is expected to alter the positions neighbouring residues that directly interact with 4A8;

contacting residues (145, 146, 147, 150, 152, 246 and 258)[22] are shown in magenta in Figure

3B. Secondly the P681H mutation lies within the furin cleavage site. Furin cleavage is a

property of some more distantly related coronaviruses, and in particular not found in SARS-

CoV-1[23]. When SARS-CoV-2 is passaged *in vitro* it results in mutations in the furin cleavage

site, suggesting the cleavage is dispensable for *in vitro* infection[24]. The significance of furin

site mutations may be related to potential escape from the innate immune antiviral IFITM

proteins by allowing infection independent endosomes[25]. The significance of the multiple S2

mutations is unclear at present, though D614G, also in S2 was found to lead to a more open

RBD orientation to explain its higher infectivity[26]. T716I and D1118H occur at residues

located close to the base of the ectodomain (Figure 4B) that are partially exposed and

buried, respectively. The residue 982 is located centrally, in between the NTDs, at the top of

a short helix (approximately residues 976-982) that is completely shielded by the RBD when

spike is in the closed form, though becomes slightly more exposed in the open

conformation. Residue 681 is part of the loop (residues 676-689) containing the furin-

cleavage site, the structure of which is disordered in both cleaved and uncleaved forms[27],

though the surface-exposed locations of modelled residues 676 and 689 (orange in Figure

4B) indicate that the unmodelled residues 677-688 form a prominently-exposed loop; the

significant structural flexibility of which has prevented inclusion in structural models[22,28].

Given the association between ΔH69/V70 and RBD gene mutations, we hypothesised similar

to our chronic infection[12], that this deletion might enhancing virus infectivity and

compensate for potential loss of infectivity due to RBD mutation. In the absence of virus

isolates we used a lentiviral pseudotyping approach to test the impact of ΔH69/V70 on virus

Spike protein mediated infection. A D614G bearing Spike protein expressing DNA plasmid

was co-transfected in HEK 293T producer cell lines along with plasmids encoding lentiviral

capsid and genome for luciferase. Mutant Spike plasmids bearing ΔH69/V70, N501Y and

N501Y+ΔH69/V70 were constructed by site directed mutagenesis. Plasmids were expressed

in 293T virus producer cells and infectious titres measured in supernatants from cells by

infection of target cells expressing ACE2 and TMPRSS2 receptors. The infectious titres were

adjusted for virus input as determined by reverse transcriptase activity in virions (Figure 5).

As we previously showed, ΔH69/V70 Spike pseudotyped virus had higher infectivity

compared to wild type (Figure 5)[12]. N501Y showed a significantly lower infectivity compared

to wild type and the addition of ΔH69/V70 restored infectivity back to wild type levels

(Figure 5), consistent with our hypothesis.

Discussion

We have presented data demonstrating multiple, independent, and circulating lineages of

SARS-CoV-2 variants bearing a Spike ΔH69/V70. This deletion spanning six nucleotides, is

mostly due to an out of frame deletion of six nucleotides, has frequently followed receptor

binding amino acid replacements (N501Y, N439K and Y453F that have been shown to

increase binding affinity to hACE2 and reduce binding with monoclonal antibodies) and its

prevalence is rising internationally. Interestingly the presence of sequence at site 69/70

appears to be unique to SARS-CoV-2 and the closest bat sarbecovirus, RaTG13, and one of

the pangolin sequences. We speculate it may have been lost in the other pangolins as these

viruses are presumably originated in bats infecting the pangolins after importation to China.

The ΔH69/V70 deletion was also shown to increase Spike mediated infectivity by two-fold

over a single round of infection, and appeared to occur with a mutation that conferred

reduced susceptibility to neutralising antibodies[29]. Over the millions of replication rounds

per day in a SARS-CoV-2 infection even modest reductions in antibody susceptibility could

be significant. Therefore, ΔH69/V70 may be a 'permissive' mutation that enhances

infection[28], with the potential to enhance the ability of SARS-CoV-2 to tolerate mutations,

which could include immune/ vaccine escape variants that would have otherwise

significantly reduced viral fitness. We show here experimentally that the ΔH69/V70 deletion

is indeed able to rescue an infectivity defect induced by N501Y.

Interestingly, one of the three changes, K417N, in the South African lineage, B.1.351[18],

decreases binding affinity to hACE[20] so moderating the strength of binding, which would

have been increased by the N501Y mutation, i.e., a similar outcome for Spike infectivity but

through a direct mechanism.   This may explain the lack of ΔH69/V70 in the South African

B.1.35 lineage, though it will be important to know whether this lineage also acquires the

ΔH69/V70 deletion in the future.

The potential for SARS-CoV-2 to evolve and fix mutations is exemplified by D614G, an amino

acid replacement in S2 that alters linkages between S1 and S2 subunits on adjacent

protomers as well as RBD orientation, infectivity, and transmission[26,30,31]. The example of

D614G also demonstrates that mechanisms directly impacting important biological

processes can be indirect. Similarly, a number of possible mechanistic explanations may

underlie ΔH69/V70. For example, the fact that it sits on an exposed surface and is estimated

to alter the conformation of a particularly exposed loop might be suggestive of immune

interactions and escape, although allosteric interactions could alternatively lead to the

higher infectivity recently reported[31].

The finding of a lineage (B.1.1.7), termed VOC 202012/01, bearing seven S gene mutations

across the RBD (N501Y, A570D), S1 (ΔH69/V70 and Δ144) and S2 (P681H, T716I, S982A and

D1118H) in UK requires urgent experimental characterisation. The detection of a high

number of novel mutations suggests this lineage has either been introduced from a

geographic region with very poor sampling or viral evolution may have occurred in a single

individual in the context of a chronic infection[12]. We detected a sequence basal to B1.1.7
that had both mutations as well as D1118H in S2.

This variant bears some concerning features: firstly, the ΔH69/V70 deletion which increases
infectivity in vitro by two fold[12] . Secondly the Δ144 which may affect binding by antibodies
related to 4A8[13]. Thirdly VOC 202012/01 bears the N501Y mutation that may have higher
binding affinity for ACE2, and which has arisen independently in other countries, including
South Africa where it has led to establishment and explosive transmission of the lineage
B.1.351[18]. Finally, the VOC 2020/1201 lineage has a second RBD mutation A570D that could
alter Spike RBD structure and a mutation near to the furin cleavage site could represent
further adaptive change. The emergence of variants with higher numbers of mutations so
far in the UK and South Africa may herald an era of re-infection and threaten future vaccine
efficacy if left unchecked.

Given the emergence of multiple clusters of variants carrying RBD mutations and the
ΔH69/V70 deletion, limitation of transmission takes on a renewed urgency. Continued
emphasis on testing/tracing, social distancing and mask wearing are essential, with
investment in other novel methods to limit transmission[32]. In concert, comprehensive
vaccination efforts in the UK and globally should be accelerated in order to further limit
transmission and acquisition of further mutations. If geographically limited then focussed
vaccination may be warranted. Research is vitally needed into whether lateral flow devices
for antigen and antibody detection can detect emerging strains and the immune responses
to them. Finally, detection of the deletion and other key mutations by rapid diagnostics
should be a research priority as such tests could be used as a proxy for antibody escape
mutations to inform surveillance at global scale.

Acknowledgements
COG-UK is supported by funding from the Medical Research Council (MRC) part of UK
Research & Innovation (UKRI), the National Institute of Health Research (NIHR) and Genome
Research Limited, operating as the Wellcome Sanger Institute. RKG is supported by a
Wellcome Trust Senior Fellowship in Clinical Science (WT108082AIA). SAK is supported by
the Bill and Melinda Gates Foundation via PANGEA grant: OPP1175094. DLR is funded by the

MRC (MC UU 1201412). WH is funded by the MRC (MR/R024758/1). We thank Dr James

Voss for the kind gift of HeLa cells stably expressing ACE2. SL is funded by Medical Research

Council MC_UU_12014/12. This study was also partly funded by Rosetrees Trust.

Conflicts of interest

RKG has received consulting fees from UMOVIS lab, Gilead Sciences and ViiV Healthcare,

and a research grant from InvisiSmart Technologies.

Methods

Phylogenetic Analysis

All available full-genome SARS-CoV-2 sequences were downloaded from the GISAID

database (http://gisaid.org/)[33] on 26[th] November. Duplicate and low-quality sequences

(>5% N regions) were removed, leaving a dataset of 194,265 sequences with a length of

>29,000bp. All sequences were realigned to the SARS-CoV-2 reference strain MN908947.3,

using MAFFT v7.475 with automatic flavour selection and the --keeplength --addfragments

options[34]. Major SARS-CoV-2 clade memberships were assigned to all sequences using the

Nextclade server v0.10 (https://clades.nextstrain.org/).

Maximum likelihood phylogenetic trees were produced using the above curated dataset

using IQ-TREE v2.1.2[35]. Evolutionary model selection for trees were inferred using

ModelFinder[36] and trees were estimated using the GTR+F+I model with 1000 ultrafast

bootstrap replicates[37]. All trees were visualised with Figtree v.1.4.4

(http://tree.bio.ed.ac.uk/software/figtree/) and ggtree v1.14.6 rooted on the SARS-CoV-2

reference sequence and nodes arranged in descending order. Nodes with bootstraps values

of <50 were collapsed using an in-house script.

To reconstruct a phylogeny for the 69/70 Spike region of the 17 *Sarbecoviruses* examined in

Fig 2, Rdp5[38] was used on the codon Spike alignment to determine the region between

amino acids 1 and 256 as putatively non-recombinant. A tree was reconstructed using the

protein alignment of this region with FastTree (default parameters)[39]. Alignment

visualisation was done using BioEdit[40].

Structural modelling

The structure of the post-deletion NTD (residues 14-306) was modelled using I-TASSER[41], a

method involving detection of templates from the protein data bank, fragment structure

assembly using replica-exchange Monte Carlo simulation and atomic-level refinement of

structure using a fragment-guided molecular dynamics simulation. The structural model

generated was aligned with the spike structure possessing the pre-deletion conformation of

the 69-77 loop (PDB 7C2L[22]) using PyMOL (Schrödinger). Figures prepared with PyMOL

using PDBs 7C2L, 6M0J[42], 6ZGE28 and 6ZGG[28].

Pseudotype virus preparation

Viral vectors were prepared by transfection of 293T cells by using Fugene HD transfection

reagent (Promega). 293T cells were transfected with a mixture of 11ul of Fugene HD, 1µg of

pCDNAΔ19Spike-HA, 1ug of p8.91 HIV-1 gag-pol expression vector22,23, and 1.5µg of

pCSFLW (expressing the firefly luciferase reporter gene with the HIV-1 packaging signal).

Viral supernatant was collected at 48 and 72h after transfection, filtered through 0.45um

filter and stored at -80˚C as previously described24. Infectivity was measured by luciferase

detection in target 293T cells transfected with TMPRSS2 and ACE2.

Standardisation of virus input by SYBR Green-based product-enhanced PCR assay (SG-PERT)

The reverse transcriptase activity of virus preparations was determined by qPCR using a

SYBR Green-based product-enhanced PCR assay (SG-PERT) as previously described[43].

Briefly, 10-fold dilutions of virus supernatant were lysed in a 1:1 ratio in a 2x lysis solution

(made up of 40% glycerol v/v 0.25% Trition X-100 v/v 100mM KCl, RNase inhibitor 0.8 U/ml,

TrisHCL 100mM, buffered to pH7.4) for 10 minutes at room temperature.

12µl of each sample lysate was added to thirteen 13µl of a SYBR Green master mix

(containing 0.5µM of MS2-RNA Fwd and Rev primers, 3.5pmol/ml of MS2-RNA, and

0.125U/µl of Ribolock RNAse inhibitor and cycled in a QuantStudio. Relative amounts of

reverse transcriptase activity were determined as the rate of transcription of bacteriophage

MS2 RNA, with absolute RT activity calculated by comparing the relative amounts of RT to

an RT standard of known activity.

References

1   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable
bat origin. *nature* **579**, 270-273 (2020).

2   Sungnak, W. *et al.* SARS-CoV-2 entry factors are highly expressed in nasal epithelial
cells together with innate immune genes. *Nat Med* **26**, 681-687,
doi:10.1038/s41591-020-0868-6 (2020).

3   Thomson, E. C. *et al.* The circulating SARS-CoV-2 spike variant N439K maintains
fitness while evading antibody-mediated immunity. *bioRxiv* (2020).

4   Starr, T. N. *et al.* Deep mutational scanning of SARS-CoV-2 receptor binding domain
reveals constraints on folding and ACE2 binding. *Cell* **182**, 1295-1310. e1220 (2020).

5   Greaney, A. J. *et al.* Complete mapping of mutations to the SARS-CoV-2 spike
receptor-binding domain that escape antibody recognition. *Cell Host & Microbe*
(2020).

6   Starr, T. N. *et al.* Prospective mapping of viral mutations that escape antibodies used
to treat COVID-19. *bioRxiv* (2020).

7   Choi, B. *et al.* Persistence and Evolution of SARS-CoV-2 in an Immunocompromised
Host. *The New England journal of medicine* **383**, 2291-2293,
doi:10.1056/NEJMc2031364 (2020).

8   MacLean, O. A. *et al.* Natural selection in the evolution of SARS-CoV-2 in bats, not
humans, created a highly capable human pathogen. *BioRxiv* (2020).

9   consortiumcontact@cogconsortium.uk, C.-G. U. An integrated national scale SARS-
CoV-2 genomic surveillance network. *Lancet Microbe* **1**, e99-e100,
doi:10.1016/S2666-5247(20)30054-9 (2020).

10   Mlcochova, P. *et al.* Combined Point-of-Care Nucleic Acid and Antibody Testing for
SARS-CoV-2 following Emergence of D614G Spike Variant. *Cell Rep Med* **1**, 100099,
doi:10.1016/j.xcrm.2020.100099 (2020).

11   He, X. *et al.* Temporal dynamics in viral shedding and transmissibility of COVID-19.
Nat Med **26**, 672-675, doi:10.1038/s41591-020-0869-5 (2020).

12   Kemp, S. A. *et al.* Neutralising antibodies drive Spike mediated SARS-CoV-2 evasion.
medRxiv, 2020.2012.2005.20241927, doi:10.1101/2020.12.05.20241927 (2020).

13   McCarthy, K. R. *et al.* Natural deletions in the SARS-CoV-2 spike glycoprotein drive
antibody escape. *bioRxiv*, 2020.2011.2019.389916, doi:10.1101/2020.11.19.389916
(2020).

14   Lam, T. T. *et al.* Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins.
Nature **583**, 282-285, doi:10.1038/s41586-020-2169-0 (2020).

15   Munnink, B. B. O. *et al.* Transmission of SARS-CoV-2 on mink farms between humans
and mink and back to humans. *Science* (2020).

16   Baum, A. *et al.* Antibody cocktail to SARS-CoV-2 spike protein prevents rapid
mutational escape seen with individual antibodies. *Science* **369**, 1014-1018,
doi:10.1126/science.abd0831 (2020).

17   Gu, H. *et al.* Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy.
Science **369**, 1603-1607, doi:10.1126/science.abc4730 (2020).

18   Tegally, H. *et al.* Emergence and rapid spread of a new severe acute respiratory
syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations
in South Africa. *medRxiv*, 2020.2012.2021.20248640,
doi:10.1101/2020.12.21.20248640 (2020).

19   Weisblum, Y. *et al.* Escape from neutralizing antibodies by SARS-CoV-2 spike protein
variants. *Elife* **9**, e61312, doi:10.7554/eLife.61312 (2020).

20   Starr, T. N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain
Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310 e1220,
doi:10.1016/j.cell.2020.08.012 (2020).

21   Rambaut A., L. N., Pybus O, Barclay W, Carabelli A. C., Connor T., Peacock T.,
Robertson D. L., Volz E., on behalf of COVID-19 Genomics Consortium UK (CoG-UK).
Preliminary genomic characterisation of an emergent SARS-CoV-2 lineage in the UK
defined by a novel set of spike mutations, <https://virological.org/t/preliminary-
genomic-characterisation-of-an-emergent-sars-cov-2-lineage-in-the-uk-defined-by-a-
novel-set-of-spike-mutations/563> (2020).

22   Chi, X. *et al.* A neutralizing human antibody binds to the N-terminal domain of the
Spike protein of SARS-CoV-2. *Science* **369**, 650-655, doi:10.1126/science.abc6952
(2020).

23   Andersen, K. G., Rambaut, A., Lipkin, W. I., Holmes, E. C. & Garry, R. F. The proximal
origin of SARS-CoV-2. *Nat Med* **26**, 450-452, doi:10.1038/s41591-020-0820-9 (2020).

24 Davidson, A. D. *et al.* Characterisation of the transcriptome and proteome of SARS-
CoV-2 reveals a cell passage induced in-frame deletion of the furin-like cleavage site
from the spike glycoprotein. *Genome Med* **12**, 68, doi:10.1186/s13073-020-00763-0
(2020).

25 Peacock, T. P. *et al.* The furin cleavage site of SARS-CoV-2 spike protein is a key
determinant for transmission due to enhanced replication in airway cells. *bioRxiv*
(2020).

26 Yurkovetskiy, L. *et al.* Structural and Functional Analysis of the D614G SARS-CoV-2
Spike Protein Variant. *Cell* **183**, 739-751 e738, doi:10.1016/j.cell.2020.09.032 (2020).

27 Wrobel, A. G. *et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform
on virus evolution and furin-cleavage effects. *Nature Structural & Molecular Biology*
27, 763-767, doi:10.1038/s41594-020-0468-7 (2020).

28 Wrobel, A. G. *et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform
on virus evolution and furin-cleavage effects. *Nat Struct Mol Biol* **27**, 763-767,
doi:10.1038/s41594-020-0468-7 (2020).

29 Kemp, S. A. *et al.* Recurrent emergence and transmission of a SARS-CoV-2 Spike
deletion ΔH69/V70. *bioRxiv*, 2020.2012.2014.422555,
doi:10.1101/2020.12.14.422555 (2020).

30 Korber, B. *et al.* Tracking changes in SARS-CoV-2 Spike: evidence that D614G
increases infectivity of the COVID-19 virus. *Cell* **182**, 812-827. e819 (2020).

31 Hou, Y. J. *et al.* SARS-CoV-2 D614G variant exhibits efficient replication ex vivo and
transmission in vivo. *Science* **370**, 1464-1468, doi:10.1126/science.abe8499 (2020).

32 Mlcochova, P. *et al.* Extended in vitro inactivation of SARS-CoV-2 by titanium dioxide
surface coating. *bioRxiv* (2020).

33 Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data - from
vision to reality. *Euro surveillance : bulletin Europeen sur les maladies transmissibles*
= European communicable disease bulletin **22**, 30494, doi:10.2807/1560-
7917.ES.2017.22.13.30494 (2017).

34 Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780,
doi:10.1093/molbev/mst010 (2013).

35   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference   in   the   Genomic   Era.   *Mol   Biol   Evol*   **37**,   1530-1534,
doi:10.1093/molbev/msaa015 (2020).
36   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*
14, 587-589, doi:10.1038/nmeth.4285 (2017).
37   Minh, B. Q., Nguyen, M. A. & von Haeseler, A. Ultrafast approximation for
phylogenetic bootstrap. *Mol Biol Evol* **30**, 1188-1195, doi:10.1093/molbev/mst024
(2013).
38   Martin, D. P., Murrell, B., Golden, M., Khoosal, A. & Muhire, B. RDP4: Detection and
analysis of recombination patterns in virus genomes. *Virus evolution* **1** (2015).
39   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: computing large minimum evolution
trees   with   profiles   instead   of   a   distance   matrix.   *Mol   Biol   Evol*   **26**,   1641-1650,
doi:10.1093/molbev/msp077 (2009).
40   Hall, T., Biosciences, I. & Carlsbad, C. BioEdit: an important software for molecular
biology. *GERF Bull Biosci* **2**, 60-61 (2011).
41   Roy, A., Kucukural, A. & Zhang, Y. I-TASSER: a unified platform for automated protein
structure and function prediction. *Nat Protoc* **5**, 725-738, doi:10.1038/nprot.2010.5
(2010).
42   Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to
the ACE2 receptor. *Nature* **581**, 215-220, doi:10.1038/s41586-020-2180-5 (2020).
43   Vermeire, J. *et al.* Quantification of reverse transcriptase activity by real-time PCR as
a fast and accurate method for titration of HIV, lenti- and retroviral vectors. *PloS one*
7, e50859-e50859, doi:10.1371/journal.pone.0050859 (2012).
502

Figure 1. A. Sub-sampled global phylogeny of SARS-CoV-2 whole genome sequences highlighting those with specific mutations in Spike: ΔH69/V70, N439K, Y453F and N501Y. Tree tips are coloured by geographic region (see key). Grey bars on the right show the presence or absence of the deletion H69/V70 and amino acid variants N439K, Y453F, and N501Y. Lineages from Rambaut et al. 2020 are shown.

Figure 1B–D. Number of new occurrences of SARS-CoV-2 sequences with the Δ H69/V70 deletion. Frequency of the ΔH69/V70 deletion from the GISAID database (accessed 16th Dec 2020) by reporting country and sampling date: **B.** Worldwide carriage of ΔH69/V70; most sequences preferentially carry RBD mutations alongside the deletion. **C.** Several distinct ΔH69/V70 lineages carrying RBD mutations 439K, 453F and 501Y have begun to emerge, predominantly in Denmark and England. **D.** Sequences carrying the ΔH69/V70 but in the absence of other major Spike mutations are restricted mostly to Denmark. **The relative increase in frequency of sequences carrying Spike mutants 439K E. 501Y F. and ΔH69/70;** viruses co-carrying the Δ69/70 deletion became dominant between September-October 2020 in terms of cumulative cases.

Figure 2. Comparison of the H69/V70 deletion site to other sarbecoviruses. A. phylogeny for the Spike region 1-256 for **B.** protein sequences from 17 Sarbecoviruses, including SARS-CoV-2 (Wuhan-Hu-1) and SARS-CoV (HSZ-Cc), with distinct genotypes at the Spike region around amino acid positions 69 and 70 (highlighted in orange box). The 69/70 HL insertion in the P1E sequence from the Guangxi pangolin virus cluster is highlighted in a yellow box. **C.** The nucleotide alignment between SARS-CoV-2 Wuhan-Hu-1, B.1.1.7, the bat sarbecovirus (RatG13) and the Guangxi pangolin viruses shows the difference between the out of frame deletion observed in the former and the in-frame deletion in the latter.

Figure 3. Structural characteristics of viruses carrying deletion at H69/V70 and co-occurring RBD mutations A. Prediction of conformational change in the spike N-terminal domain due to deletion of residues His69 and Val70. The pre-deletion structure is shown in cyan, except for residues 69 and 70, which are shown in red. The predicted post-deletion structure is shown in green. Residues 66-77 of the pre-deletion structure are shown in stick representation and coloured by atom (nitrogen in blue, oxygen in coral). Yellow lines connect aligned residues 66-77 of the pre- and post-deletion structures and the distance of 6 Å between aligned alpha carbons of Thr73 in the pre- and post-deletion conformation is labelled. **B.** Surface representation of spike homotrimer in open conformation (PDB: 7C2L) with each monomer shown in different shades of grey. On the monomer shown positioned to the right, the exposed loop consisting of residues 69-77 is shown in cyan and the neutralising antibody (4A8) binding NTD epitope described by Chi et al. is shown in magenta **C.** Surface representation of spike in closed conformation (PDB: 6ZGE) viewed in a 'top-down' view along the trimer axis. The residues associated with RBD substitutions N439K, Y453F and N501Y are highlighted in red and labelled on a single monomer. **D.** Spike in open conformation with a single erect RBD (PDB: 6ZGG) in trimer axis vertical view with the location of Δ H69/V70 in the N-terminal domain and RBD mutations highlighted as red spheres and

Figure 4. A. Lineages bearing Spike N501Y and ΔH69/V70 mutations: A sub-sampled global phylogeny of SARS-CoV-2 sequences bearing either the ΔH69/V70 or N501Y mutations. Duplicate sequences were removed, and 200 randomly sub-sampled background sequences (without either mutation) were included to create a representative sub-sample of sequences. Distinct sub-lineages of the ΔH69/V70 mutation are currently currently circulating, predominantly in the UK. The novel UK variant VOC 202012/01 (Lineage B.1.1.7, green) is also shown and in black box a related sequence carrying N501Y, ΔH69/V70 and D1118H. **B.** Surface representation of spike homotrimer in open conformation with one upright RBD overlaid with ribbon representation (PDB: 6ZGG, Wrobel et al., 2020), with different monomers shown in shades of grey. The deleted residues H69 and V70 and the residues involved in amino acid substitutions (501, 570, 716, 982 and 1118) and the deletion at position 144 are coloured red on each

Figure 5: Spike mutant ΔH69/V70 has 2 fold higher infectivity compared to wild type (D614G background) and compensates for infectivity defect of Spike N501Y. Single round Infection of target cells A and B by luciferase expressing lentivirus pseudotyped with SARS-CoV-2 Spike protein on 293T cells co-transfected with ACE2 and TMPRSS2 plasmids. Data showing Infectivity normalized for virus input using reverse transcriptase activity in virus supernatants. RLU – relative light units; U – unit of reverse transcriptase activity (RT). Data are representative of 2 independent experiments.