Predicting recombination frequency from map distance

Mikko Kivikoski[a,1], Pasi Rastas[b], Ari Löytynoja[b], Juha Merilä[a,c]

[a] Ecological Genetics Research Unit, Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, FI-00014 University of Helsinki, Finland

[b] Institute of Biotechnology, HiLIFE, FI-00014 University of Helsinki, Finland

[c] Area of Ecology and Biodiversity, School of Biological Sciences, Kadoorie Biological Science Building, Pokfulam Road, The University of Hong Kong, Hong Kong SAR

[1] corresponding author: Mikko Kivikoski, E-mail: mikko.kivikoski@helsinki.fi

Author ORCID iDs:

M.K. **0000-0001-7105-651X**, P.R. **0000-0003-2768-1339**, A.L. **0000-0001-5389-6611**, J.M. **0000-0001-9614-0072**

Keywords: Crossover interference, *Gasterosteus aculeatus*, *Homo sapiens*, Mapping function, *Pungitius pungitius*, Stickleback

Running title: Predicting recombination frequency

Abstract

Map distance is one of the key measures in genetics and indicates the

expected number of crossovers between two loci. Map distance is estimated

from the observed recombination frequency using mapping functions, the

most widely used of those, Haldane and Kosambi, being developed at the

time when the number of markers was low and unobserved crossovers had a

substantial effect on the recombination fractions. In contemporary high-

density marker data, the probability of multiple crossovers between adjacent

loci is negligible and different mapping functions yield the same result, that

is, the recombination frequency between adjacent loci is equal to the map

distance in Morgans. However, high-density linkage maps contain an

interpretation problem: the map distance over a long interval is additive and

its association with recombination frequency is not defined. Here, we

demonstrate with high-density linkage maps from humans and stickleback

fishes that the inverse of Haldane or Kosambi mapping functions fail to

predict the recombination frequency from map distance, and show that this

is because the expected number of crossovers is not sufficient to predict

recombination frequency. We formulate a piecewise function to calculate

the probability of no crossovers between the markers that yields more

accurate predictions of recombination frequency from map distance. Our

results demonstrate that the association between map distance and

recombination frequency is context-dependent and no universal solution

exists. We anticipate that our study will motivate further research on this

subject to yield a more accurate mathematical description of map distance in

the context of modern data.

Introduction

Crossovers in meiosis break the physical linkage among loci and allow

formation of recombinant chromosomes and ensure chromosome

segregation in meiosis I (Koehler et al. 1996, Hassold et al. 2021). Although

crossovers and the resultant recombinations have been studied for more than

a hundred years (Sturtevant 1913), many related questions remain

unanswered. Due to their profound importance in sexual reproduction,

substantial research efforts have focused on better understanding the among-

organism and individual variation of crossover rate (e.g., Stapley et al. 2017,

Haenel et al. 2018), and on the other hand, recent technologies have been

utilized for detecting recombinations at the gamete level (Dréau 2019, Bell

et al. 2020, Yang et al. 2022). Crossovers also have implications for

statistical properties essential in population genetics, such as the variance of

genetic relatedness (Veller et al. 2020). However, one aspect that has gained

little attention in the era of high-throughput sequencing is the interpretation

of genetic map distances.

Recombinant gametes or offspring can be utilized to build linkage maps that

quantify the physical order and map distance (i.e., expected number of

crossovers) between loci. Map distances are estimated with mapping

functions that attempt to account for the non-additivity of the recombination

frequencies due to multiple crossovers between adjacent loci. The two most

widely used and referred to mapping functions are probably those of

Haldane (1919) and Kosambi (1944, e.g. Lynch and Walsh 1998, Hill and

Weir 2011, Otto and Payseur 2019). However, modern sequencing methods

and increasing marker density have reduced the utility of these functions in

linkage map reconstruction; the probability of multiple crossovers between

closely positioned adjacent loci is negligible and all mapping functions yield

essentially the same result, $r = d$, where r is recombination frequency and d

is map distance in Morgans (e.g., Purcell et al. 2007). As the map distances

are estimated only for adjacent loci, they are additive over longer intervals.

Consequently, recombination frequency over intermediate or long map

distances (e.g., 50 cM) does not follow any simple association.

Although the importance of mapping functions in linkage map construction

has decreased over time with increasing access to dense marker data,

inverse mapping functions have recently been utilized to predict genetic

shuffling in meiosis (Veller et al. 2019) and from that the variance in

genetic relatedness (Veller et al. 2020). It is technically trivial to translate

map distance to recombination frequency with an inverse of mapping

function, but as all mapping functions effectively yield the same map for

high-density data, it is not clear which inverse mapping function, if any,

would yield the true recombination frequencies. Veller et al. (2020)

observed that empirical variance in genetic relatedness among human

(*Homo sapiens*) subjects did not match those predicted with the inverse of

Kosambi function, indicating that the inverse of Kosambi function is invalid

for translating map distances into recombination frequencies.

Here, we show with empirical data from humans and from nine-spined

(*Pungitius pungitius*) and three-spined (*Gasterosteus aculeatus*)

sticklebacks, that the inverse of Kosambi, Haldane, or linear mapping

functions do not translate additive map distances correctly to recombination

frequencies. To that end, we propose a new approach to translate map

distances to recombination frequencies using a piecewise function based on

the probability of no crossovers between the markers. We motivate this

approach with empirical data from the nine-spined stickleback and

demonstrate its performance with empirical data from humans and nine- and

three-spined sticklebacks.

Materials and methods

Number of crossovers and recombination frequency

An odd number of gametic crossovers between two loci cause
recombination, and the recombination frequency of two loci is equal to the
probability of an odd number of gametic crossovers between them.
Assuming that there is no chromatid interference, any positive number of
crossovers in the bivalent leads to equal proportions of recombinant and
non-recombinant gametes, while the absence of crossovers between two loci
always leads to non-recombinant gametes. From this follows that
recombination frequency r, can be expressed as a function of probability of
no crossovers between two loci in the bivalent p_0, so that $r = \frac{1}{2}(1 - p_0)$
(Mather 1938, Weeks et al. 2009). To translate map distance into
recombination frequency, the association between map distance and p_0 is
necessary.

Map distance of two loci is the expected number of gametic crossovers
between them. However, the distance does not tell the variation around the
expectation or the likelihood for no bivalent crossovers (p_0) in that particular
interval. Depending on the number of crossovers in the bivalent and their
localization, the same map distance can be associated with different values

of p_0 and recombination frequency (Fig. 1). To address this ambiguity, we

formulate a piecewise function $p_0(k)$ that gives the probability of no

crossovers between the two loci in the bivalent with k crossovers.

Derivation of $p_0(k)$

The impetus for the model comes from chromosome 8 (LG8) of the nine-

spined stickleback (*Pungitius pungitius*). LG8 is a metacentric chromosome

(Kivikoski et al. 2021), where maternal map lengths of the two chromosome

arms are 52 and 60 cM, corresponding on average to 1.04 and 1.20

crossovers per bivalent, respectively (Table S1). Out of 934 gametes, 265

have two maternal crossovers and 94% of those gametes have one crossover

in both chromosome arms. Because these are gametic crossovers, the

number of crossovers in the bivalent could have been two or more.

However, the inferred likelihoods for two, three, or four crossovers in the

bivalent are 0.85, 0.12, and 0.03, respectively, and likelihoods for other

crossover counts are zero (see the section 'Inference of bivalent crossover

rate' below). Assuming no chromatid interference, the proportion of the

gametes with two crossovers descending from meiosis with the same

number of crossovers is 0.79 (see supplementary methods). Given this high

proportion, those crossovers are, for the sake of example, interpreted to

largely represent bivalent crossovers in meiosis with two crossovers. The

map positions of the two crossovers are uncorrelated (r=0.029, p=0.64) and

they are distributed approximately uniformly along the map distance (Fig.

2).

Putting this observation in a general context, we assume that when there are

k crossovers in the bivalent they occur in k distinct regions, and within those

regions the exact localization is independent of other crossovers in the

bivalent. Let d be the map length of the whole chromosome. If there are

$k(k > 0)$ crossovers in the bivalent, they are assumed to occur in k non-

overlapping regions of equal size, $\frac{d}{k}$, such that every crossover occurs in a

different chromosomal region. Within each region, their localizations are

uniformly distributed and independent across the regions. For a given

marker pair, $p_0 = \sum_{k=0}^{n} P(\text{no crossovers}|\ k)\ P(k \text{ crossovers}) =$

$\sum_{k=0}^{n} p_k\ p_0(k)$, where n is the highest possible number of crossovers in the

bivalent, p_k is the probability of k crossovers in the bivalent and $p_0(k)$ is

the probability of no crossovers between the markers when there are k

crossovers in the bivalent. For markers at map positions m_i and m_j ($m_j \geq$

m_i), $p_0(k)$ is defined as:

$$p_0(k) = \begin{cases} 1, & if\ k = 0 \\ 1 - \dfrac{|m_j - m_i|}{\frac{d}{k}}, & if\ \left\lceil \dfrac{m_j}{\frac{d}{k}} \right\rceil = \left\lceil \dfrac{m_i}{\frac{d}{k}} \right\rceil\ and\ k > 0 \\ \left(1 - \dfrac{|b_i - m_i|}{\frac{d}{k}}\right)\dfrac{|b_j - m_j|}{\frac{d}{k}}, & if\ \left\lceil \dfrac{m_j}{\frac{d}{k}} \right\rceil - \left\lceil \dfrac{m_i}{\frac{d}{k}} \right\rceil = 1\ and\ k > 0 \\ 0, & if\ \left\lceil \dfrac{m_j}{\frac{d}{k}} \right\rceil - \left\lceil \dfrac{m_i}{\frac{d}{k}} \right\rceil > 1\ and\ k > 0 \end{cases}$$ (Eq. 1),

where $\left\lceil \dfrac{m_i}{\frac{d}{k}} \right\rceil$ and $\left\lceil \dfrac{m_j}{\frac{d}{k}} \right\rceil$ are the crossover regions of the markers m_i and m_j,

respectively, and b_i and b_j are the upper boundaries for these regions,

respectively, so that $b_i = \left\lceil \dfrac{m_i}{\frac{d}{k}} \right\rceil \dfrac{d}{k}$ and $b_j = \left\lceil \dfrac{m_j}{\frac{d}{k}} \right\rceil \dfrac{d}{k}$. Notation $\lceil\ \rceil$ refers to

ceiling function, and $\left\lceil \dfrac{m_j}{\frac{d}{k}} \right\rceil$ is the least integer greater than, or equal to $\dfrac{m_j}{\frac{d}{k}}$.

This function is also applicable for multiple loci. In the case of three

markers (m_i, m_j, m_k), the recombination frequencies derived with this

function meet the criteria $r_{ij} + r_{jk} \geq r_{ik}$, where r_{ij}, r_{jk}, and r_{ik} are the

recombination frequencies between markers m_i and m_j, m_j and m_k, and m_i

and m_k, respectively (Karlin and Liberman 1978, Weeks 1994; see also

Supplementary methods).

Our approach is built on three premises: (1) Absence of chromatid

interference; (2) even spacing of crossovers due to crossover interference;

and (3) the fact that the number of bivalent crossovers in meiosis I varies

across individuals, chromosomes, sexes, and species. The second and the

third premises have been demonstrated several times in the literature and are

merely facts. Crossover interference causes even distribution of crossovers

by their physical distances (micrometers) (Zhang et al. 2014, Zickler and

Kleckner 2015) and approximately by base pairs, as the chromatin packing

ratio per nucleus is roughly constant (see Fig. S2 in Veller et al. 2019).

However, the spatial distribution per map distance cannot be concluded

directly. The number of crossovers per chromosome is not constant but

varies between chromosomes, nuclei, individuals, sexes, and species (e.g.,

Stapley et al. 2017) and hence affects the spatial distribution of crossovers

(Charles 1938, Zhang et al. 2014). While the absence of chromatid

interference is a common assumption in the literature (e.g., Weinstein 1936,

Zhao and Speed 1996, Sandor et al. 2012), this has been empirically tested

and confirmed only in a few organisms, including humans and yeast (Zhao

et al. 1995, Mancera et al. 2008, Hou et al. 2014, Wang et al. 2019).

In contrast to existing mapping functions, the proposed model is not based

on the renewal process (Zhao and Speed 1996) and it does not model

crossover interference parametrically. Instead, crossover interference is

implemented structurally by assuming that crossovers occur in close

proximity less often than would be expected by chance in the absence of

crossover interference.

We applied the function for sex-specific recombination data in the 21

chromosomes of nine- and three-spined sticklebacks and in the 22 human

autosomes. The total map lengths were derived from the linkage maps and

the likelihoods for different numbers of crossovers in the bivalent were

inferred from the observed number of crossovers (see 'Inference of bivalent

crossover frequency' below). Implementation of the $p_0(k)$ function in R is

provided in the 'Data archiving' section.

Stickleback linkage maps

The crossover frequencies and locations were estimated from the linkage

maps described in Kivikoski et al. (2021). For the nine-spined stickleback,

high-density linkage maps (22,468 markers informative to conclude

crossover) were reconstructed with Lep-MAP3 software (Rastas 2017) from

a data set of 133 parents and 938 F_1 offspring. The parental fish, 46 females

and 87 males, were wild-caught individuals from the Baltic Sea coast of

Finland (Helsinki, 60°13'N, 25°11'E) that were artificially crossed in

laboratory to produce the aforementioned F_1 offspring (Kivikoski et al.

2021, see also Rastas et al. 2016). Five females were each crossed with a

different male, forming five full-sib families, and the other 41 females were

each crossed with two different males, which formed 41 half-sib families.

Identification of the single nucleotide polymorphisms (SNPs) of the parental

and the F_1 fish were based on whole-genome sequencing of the parents (5–

10X coverage; Illumina Hiseq platforms, BGI Hong Kong) and DarTseq

(Diversity Arrays Technology, Pty Ltd) genotyping of the F_1 fish (Kivikoski

et al. 2021). The read mapping was conducted with BWA-mem (ver. 0.7.15,

Li 2013) and the variants were called with SAMtools mpileup (ver 1.9, Li et

al. 2009) following the Lep-MAP3 software pipeline (Rastas 2017). The

linkage maps were built with Lep-MAP3, and the number of paternal and

maternal crossovers were inferred from the observed changes in the

haplotype phase of the F_1 offspring. Crossovers could not be inferred for

four crosses with only single offspring, and the final dataset included 934

offspring in total.

The three-spined stickleback linkage maps were based on previously

published sequencing data (Pritchard et al. 2017). In short, ninety wild-

caught parental fish (Baltic Sea, Helsinki, Finland, 60°13'N, 25°11'E) were

crossed such that the males (n=30) were each crossed with two different

females (n=60). This yielded 60 and 30 full-sib and half-sib families,

respectively, with 517 F_1 offspring. Genotyping of the parental and the F_1

fish were based on genotype-by-sequencing, according to the Restriction-

site Associated DNA (RAD) sequencing protocol of Elshire et al. (2011).

The crossing, rearing protocols and sequencing data are explained in more

detail in Leder et al. (2015) and Pritchard et al. (2017). For this study, the

RAD reads from Pritchard et al. (2017) were mapped to three-spined

stickleback reference genome (v4, Peichel et al. 2017) and the variants were

called following the Lep-MAP3 pipeline as for the nine-spined sticklebacks.

This yielded 28,187 informative markers to build linkage maps with Lep-

MAP3.

For all linkage maps (maternal and paternal maps of the nine- and three-

spined stickleback), the genetic distances between the adjacent markers

were calculated from recombination frequency with the Haldane mapping

function. The distances are additive for non-adjacent markers. There were,

on average, 1,070 and 1,342 markers per chromosome in the nine- and

three-spined stickleback maps, respectively. Hence, the inter-marker

distances were short: on average 19,571 bp corresponding to 0.054 cM and

0.106 cM in the paternal and maternal maps of the nine-spined stickleback,

respectively, and 15,461 bp corresponding to 0.043 cM and 0.075 cM in the

paternal and maternal maps of the three-spined stickleback, respectively. As

all conventional functions yield very similar results for small recombination

frequencies, the choice of the mapping function has a minor impact on the

map distances.

Analysis of human recombination data

To evaluate the general applicability of the new function, we analyzed

human data from Halldorsson et al. (2019). This consisted of sex-specific

linkage maps (their supplementary Data S1 and S2) and the crossover data

(their supplementary Data S4). Crossover locations and counts were

obtained from the column 'medsnp' of the sex-specific linkage maps for the

41,092 probands with both paternal and maternal crossover information. All

crossovers were used irrespective of their status regarding the gene

conversions (complex, non-complex, or not assessed; see Halldorsson et al.

2019). Moreover, no probands were discarded based on the total number of

crossovers in them; the highest number of crossovers per proband per

chromosome was 17 maternal crossovers in chromosome 13. The number of

markers in the linkage maps ranged from 17,894 to 90,036 depending on the

chromosome. We used R (ver. 4.1.1 R Core Team 2018) with seed value

2021 to sample 1.5% of the markers of every chromosome, which yielded

268–1,351 markers (i.e., 35,778–911,925 marker pairs) per chromosome.

For every marker pair, we estimated the sex-specific recombination

frequency by calculating the proportion of the studied probands (n=41,092)

with an odd number of crossovers between the markers.

Inference of crossover frequency

Assuming there is no chromatid interference, the probability of observing k

crossovers in a randomly sampled meiotic product depends on the number

of crossovers in the bivalent so that $P(k) = \binom{n}{k}\left(\frac{1}{2}\right)^{n}$, where n is the number

of crossovers in the bivalent and $0 \leq k \leq n$ (Weinstein 1936). The number

of crossovers in the bivalent varies not only between the sexes and

chromosomes but also between individuals and individual meioses (see

Broman and Weber 2000 for an example). Therefore, the observed

crossovers in the gamete pool are a sample of crossovers from meioses with

a different number of crossovers in the bivalent. As the sampling function

above is known, the multinomial distribution of the number of crossovers in

the bivalent can be estimated. We used the expectation-maximization (EM)

algorithm of Yu and Feingold (2001) to estimate the multinomial

distribution for different numbers of crossovers in bivalent. The algorithm

approximates the multinomial distribution that maximizes the likelihood of

the data consisting of the numbers of meiotic products with $0 \dots N$ observed

crossovers in each chromosome. According to Yu and Feingold (2001), for

data where the highest number of observed crossovers in a single meiotic

product is N, it is sufficient to estimate the multinomial distribution between

0 and $2N - 1$. The algorithm was applied separately for maternal and

paternal crossovers and for each chromosome by pooling all meiotic

products (n=934, n=517, n=41,092 for nine- and three-spined stickleback

and human data, respectively, Tables S1–S6). The estimated multinomial

distribution includes the maximum-likelihood estimate for no crossovers in

the bivalent. We applied the bootstrapping test of Yu and Feingold (2001) to

estimate if, in case of an estimate above zero, the deviation from that is

statistically significant ($p < 0.05$). For the chromosomes with non-

significant p-values, we used a restricted multinomial distribution that

restricts the likelihood of no crossovers to zero for further analyses. This

choice was made to assume the obligate crossover as a null hypothesis.

Chromosomes with p-value below 0.05 were chr15 of the three-spined

stickleback (paternal meioses) and in humans chr21 and chr22 (both

maternal and paternal meioses) and chr3 (paternal meioses). For humans,

meioses with no crossovers in chromosomes 21 and 22 have been

previously reported in cytological studies (e.g. Wang et al. 2017, Hassold et

al. 2021), but we are not aware of such findings for the chr3 and further

verification is needed. For the sticklebacks, this is the first study testing the

obligate crossover hypothesis.

Performance assessment of functions

The performance of $p_0(k)$ and the three inverse mapping functions in

predicting recombination frequency from map distance were assessed by

calculating the mean absolute error of the predictions and the intra-

chromosomal component of genetic shuffling, $\bar{r}$ (Veller et al. 2019). The

equations for the inverse mapping functions were: $r = \frac{1}{2}(1 - e^{-2d})$

(Haldane 1919), $r = \frac{1}{2}tanh(2d)$ (Kosambi 1944), and

$\begin{cases} r = d, \text{ when } d \leq 0.5 \\ r = \frac{1}{2}, \text{when } d > 0.5 \end{cases}$ (linear). In all functions, r is the recombination

frequency and d is the map distance in Morgans.

The mean absolute error of the predicted recombination frequency was

calculated as $\frac{1}{n}\sum |r_e - r_p|$, where r_e and r_p are empirical and predicted

recombination frequencies per marker pair, respectively, and n is the total

number of marker pairs. The intra-chromosomal component of $\bar{r}$ is the part

of genetic shuffling due to crossover rate and localization. A higher number

of crossovers, and their even distribution increase $\bar{r}$, while a low crossover

rate and terminal or aggregated localization decrease it. Here, we calculated

the intra-chromosomal component of $\bar{r}$, by first predicting the

recombination frequency from map distance and then converting it to

shuffling according to Eq. 10 of Veller et al. (2019) $\sum_{i<j} r_{ij}/\binom{\Lambda}{2}$, where r_{ij}

is the rate of shuffling, i.e. recombination frequency of locus pair (i, j), Λ is

the number of loci, and $\binom{\Lambda}{2}$ is the number of locus pairs.

Results

Map distance and recombination frequency do not have a fixed

association

Mapping functions formulate a fixed association between recombination

frequency and the expected number of crossovers in a gamete, the map

distance. However, the same additive map distance can be associated with

different recombination frequencies depending on the crossover positions

and the context of the focal loci (Fig. 1). This demonstrates that inverse

mapping functions have limitations in predicting recombination frequency

from map distance. Maternal crossovers in chromosome 8 of the nine-spined

stickleback have a distribution similar to the idealized example (Fig. 2),

suggesting that a structural model of crossover localization may be

applicable in predicting recombination frequency from map distance.

Another limitation of the mapping functions (and their inverses) is that they

do not account for the variation in the number of bivalent crossovers.

Inferred probabilities for bivalent crossover counts show that the number of

crossovers varies between sexes, among chromosomes, and in meioses in all

three studied species (Fig. 3, Tables S1–S6). This further indicates that the

inverse of the Kosambi or Haldane mapping function may not be suitable

for different species, sexes, or chromosomes.

Inverse Kosambi and Haldane mapping functions underestimate

recombination frequency

For the three studied species, empirical recombination frequencies lie

between the predictions of the inverse Kosambi and linear function (Fig. 4,

Fig. S1–S3). This shows that inverse Kosambi and Haldane mapping

functions systematically underestimate the recombination frequencies and

the linear function overestimates them. The poor performance of the inverse

Haldane function is not surprising as it assumes no crossover interference.

However, the fact that the inverse Kosambi function, which does implement

crossover interference, systematically underestimates recombination

frequencies implies that its crossover interference model does not reflect the

underlying biology.

The new $p_0(k)$ function outperforms the existing functions

We evaluated the new function against existing functions by calculating the

mean absolute error of predictions and the intra-chromosomal component of

genetic shuffling for the two sexes of three different species. The

predictions made with the new $p_0(k)$ functions depict the pattern of

empirical data (Fig. 4, Fig. S4–S6), and the mean absolute error of those

predictions are clearly lowest in five out of the six cases. In nine-spined

stickleback males, most meioses have one crossover and the linear function

gives a slightly lower error, than $p_0(k)$ (Table 1). The excellent overall

performance (Table S7–S9) demonstrates that the new function works for

different species and on different types of chromosomes. Consistent with the

mean absolute error, the new function is superior when assessed on the

intra-chromosomal components of $\bar{r}$ estimated from the empirical and

predicted recombination frequencies. For each species and for both sexes,

the predictions by the $p_0(k)$ function gave results closest to the empirical

results (Table 2), demonstrating its potential for the application.

Altogether, our analyses show that the inverses of the Kosambi and the

Haldane mapping functions systematically underestimate the recombination

frequencies. Although the linear function works for chromosomes with an

overall crossover rate close to 1 (map length ca. 50 cM), the $p_0(k)$ function

does not underestimate the recombination frequency to the same extent as

the other two functions. Overall, the new $p_0(k)$ function gives qualitatively

and quantitatively the best results.

Discussion

Intrinsic limitations of mapping functions

Map distance tells the expected number of crossovers between two loci and can be used as a proxy of recombination; the longer the map distance the higher the recombination frequency. In principle, translating a map distance to a recombination frequency is trivial, one only needs an inverse of a mapping function such as Kosambi (1944) or Haldane (1919). However, the problem with this approach is that in modern high-density linkage maps, map distances of non-adjacent loci are additive and an inverse mapping function is not guaranteed to give the correct recombination frequency.

The intrinsic limitations of mapping functions, namely that they describe the interference only at a general level and do not account for the variation in the strength of crossover interference or the number of crossovers, have been previously recognized (Crow 1990, Zhao and Speed 1996, Otto and Payseur 2019). However, empirical tests on the performance of inverse mapping functions with modern data are scarce.

Inverse Kosambi and Haldane mapping functions err with high-

density data

Here, we demonstrated that additive map distance can yield an array of

different recombination frequencies (Fig. 1) and showed with empirical data

from humans and nine- and the three-spined sticklebacks that the inverse of

Kosambi and Haldane mapping functions underestimate the recombination

frequencies (Fig. 4). We also formulated a new function that outperforms

those mapping functions in this task and yields lower error (Table 1) and a

more accurate estimate of genetic shuffling (Table 2). The fact that mapping

functions fail to predict recombination frequencies implies that those models

of crossover interference does not predict crossover localization correctly.

Similar findings were reported by Veller et al. (2020), who showed that the

per-chromosome variances of genetic relatedness estimated with the inverse

of Kosambi function were higher than those from cytological data that

should approximate "true" variance. Our results regarding both sexes of all

three studied species were concordant: inverse Kosambi and Haldane

functions underestimate recombination frequencies and genetic shuffling

($\overline{r}$), causing overestimation of variance in genetic relatedness, which

decreases as a function of $\overline{r}$ (Veller et al. 2020). Underestimation of

recombination frequencies means that the Kosambi map function predicts

too many crossovers (excessively long map distance) for observed

recombination frequency. The Haldane map function yields even longer

map distance, as it assumes no crossover interference.

Stickleback males have primarily one crossover per bivalent (Fig. 3, Table

S2, S4) and in those chromosomes the linear function had the best

performance (Table S7–S9). The linear function predicts recombination

frequency equal to map distance, whereas the $p_0(k)$ gives the map distance

per the total map length of the chromosome, which yields an underestimate

in certain chromosomes. This indicates that the linear function is adequate

and should be preferred for chromosomes that mainly have one bivalent

crossover, especially in species where this is a norm, such as in Lepidoptera

(Davey et al. 2017).

Accurate estimates of recombination frequency are needed in predicting

genetic shuffling and from that the variance in genetic relatedness (Visscher

et al. 2006). On the other hand, map distance per base pair (cM/Mbp) is

used as a measure of recombination in comparative genomics, especially in

non-model organisms (Stapley et al. 2017, Martin et al. 2019). This metric

is easy to obtain from linkage maps, but as shown here, meaningful

interpretation of map distance requires knowledge of the recombination

process and interpretation with an inverse mapping function can lead to

biased estimates.

Limitations of the approach

In contrast to existing methods, the approach presented here builds on the

fact that the crossover rate varies among organisms and meioses, which

leads to differences in spatial distributions of crossover sites, and this

variation should be implemented in the model. However, this approach

requires knowledge about the probability of different numbers of crossovers

in the bivalent, which is not needed by other methods and that cannot be

concluded directly from the total map length. These can be inferred from

gametic crossovers obtained from linkage maps (as done here) or analysis of

haploid offspring (Liu et al. 2015). Alternatively, cytological methods can

be used to obtain the bivalent crossover counts directly (Froenicke et al.

2002, Wang et al. 2017). Especially with gametic crossovers that contain the

sampling variance caused by the fact that only two chromatids are involved

in one crossover event, the sample size must be sufficient to obtain

reasonably accurate probability distributions. Implicitly, gametic crossovers

were, as a null expectation, assumed to present an unbiased sample of those

in the bivalent and that meiotic drive or selection for crossover count has not

occurred. However, effects of natural selection could also be involved, for

example if the offspring are studied in later life-stages instead of direct

investigation of gametes.

In the $p_0(k)$ function itself, the most important assumption is the uniform

distribution of crossovers per map distance and the arbitrarily defined

breakpoints of the "segments". These assumptions are simplifications that

allow mathematically tractable formula. The distribution of crossovers per

map distance can be studied from the gametic crossovers only to some

extent because they present a subset of those in the bivalent. However, the

likelihood for gametes with a certain number of crossovers that show all

bivalent crossovers can be calculated and the inherent uncertainty can be

estimated (Supplementary methods). Based on the distribution of gametic

crossovers per map distance, the assumptions of the model approximate the

data in many chromosomes (Fig. S7–S9). However, the non-random

distribution of error per map distance shows that all assessed functions have

a systematic bias (Fig. S10–S12).

The motivation for the approach presented here came from wild individuals

of the nine-spined stickleback, which is not a canonical organism to study

crossovers. In contrast to humans for example, the number of crossovers in

meiosis varies very little, especially in males. Despite the overall differences

in crossover rates (map lengths) of human and sticklebacks, the presented

method was demonstrated to work for human data as well.

Conclusions

The most salient finding of this study is that the inverse of the Kosambi and the Haldane mapping functions systematically underestimate recombination frequencies. Another caveat of using inverse mapping functions to translate additive map distances to recombination frequencies is that they yield one prediction per map distance, which does not match empirical findings. These findings demonstrate that (intermediate) map distances must be interpreted with care and context-specifically. We also formulated a piecewise function that allows the association between map distance and recombination frequency to be ambiguous. The fact that this function outperforms existing mapping functions in this task indicates that its implementation of crossover interference is more concordant with the data when compared with functions devised earlier. However, the presented function does not replace mapping functions in building linkage maps; it only replaces their inverses in predicting recombination from map distance.

Acknowledgements

We thank the Biodata Analytics Unit of the University of Helsinki (in particular Jukka Siren) for advice in statistics. We wish to acknowledge

CSC – IT Center for Science, Finland, for access to computational

resources. Our research was supported by Academy of Finland (grants:

129662, 134728 and 218343 to JM; 322681 to AL; 343656 to PR), the

Doctoral Programme in Wildlife Biology Research (LUOVA; funding to

MK), the Alfred Kordelin Foundation (grant 210190 to MK), National

Natural Science Foundation of China/Research Grants Council (RGC) Joint

Research Scheme 2021/2022 ('N_HKU763/21' to JM), and support from

the Helsinki Institute for Life Sciences (HiLife; grant to JM).

549   Author contribution statement

551   MK formulated the mathematical function introduced here, executed the

552   data analyses, and led the manuscript writing. PR made the stickleback

553   linkage maps and was involved in the formulation of the mathematical

554   function. AL and JM supervised the study and participated in writing the

555   manuscript.

Conflict of Interest

558 The authors declare no competing interests.

Data archiving

The stickleback linkage maps and computer code for replication of the

analyses of this study are available in Github

https://github.com/mikkokivikoski/recombinationStudies. In the repository,

the p0kFunction.r file provides an implementation of $p_0(k)$ function in R. To

facilitate reproducibility of the analyses, analytical steps following the

linkage map reconstruction step are integrated into an Anduril2 workflow

(Cervera et al. 2019).

References

Bell, A. D., Mello, C. J., Nemesh, J., Brumbaugh, S. A., Wysoker, A., &

McCarroll, S. A. (2020). Insights into variation in meiosis from 31,228

human sperm genomes. *Nature, 583*: 259–264.

Broman, K. W., & Weber, J. L. (2000). Characterization of human

crossover interference. *American Journal of Human Genetics, 66*(6): 1911–

1926.

Cervera, A., Rantanen, V., Ovaska, K., Laakso, M., Nuñez-Fontarnau, J.,

Alkodsi, A., Casado, J., Facciotto, C., Häkkinen, A., Louhimo, R., Karinen,

S., Zhang, K., Lavikka, K., Lyly, L., Pal Singh, M., & Hautaniemi, S.

(2019). Anduril 2: upgraded large-scale data integration framework.

Bioinformatics, 35(19): 3815–3817.

Charles, D. R. (1938). The spatial distribution of cross-overs in X -

chromosome tetrads of *Drosophila melanogaster*. *Journal of Genetics,*

36(1): 103–126.

Crow, J. F. (1990). Mapping functions. *Genetics, 125*(4): 669–67.

Davey, J. W., Barker, S. L., Rastas, P. M., Pinharanda, A., Martin, S. H.,

Durbin, R., McMillan, W. O., Merrill, R. M., & Jiggins, C. D. (2017). No

evidence for maintenance of a sympatric *Heliconius* species barrier by

chromosomal inversions. *Evolution Letters*, *1*(3), 138–154.

Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K.,

Buckler, E. S., & Mitchell, S. E. (2011) A robust, simple genotyping-by-

sequencing (GBS) approach for high diversity species. *PLoS One* 6: e19379.

Froenicke, L., Anderson, L. K., Wienberg, J., & Ashley, T. (2002). Male

mouse recombination maps for each autosome identified by chromosome

painting. *American Journal of Human Genetics, 71(6)*: 1353–1368.

Haenel, Q., Laurentino, T. G., Roesti, M., & Berner, D. (2018). Meta-

analysis of chromosome-scale crossover rate variation in eukaryotes and its

significance to evolutionary genomics. *Molecular Ecology, 27*(11): 2477–

2497.

Haldane, J. B. S. (1919). The combination of linkage values and the

calculation of distances between the loci of linked factors. *Journal of*

*Genetic*s, 8(4): 299–309.

Halldorsson, B. V., Palsson, G., Stefansson, O. A., Jonsson, H., Hardarson,

M. T., Eggertsson, H. P., Gunnarsson, B., Oddsson, A., Halldorsson, G. H.,

Zink, F., Gudjonsson, S. A., Frigge, M. L., Thorleifsson, G., Sigurdsson, A.,

Stacey, S. N., Sulem, P., Masson, G., Helgason, A., Gudbjartsson, D. F.,

Thorsteinsdottir, U., & Stefansson, K. (2019). Characterizing mutagenic

effects of recombination through a sequence-level genetic map. *Science,*

363(6425): eaau1043.

Hassold, T., Maylor-Hagen, H., Wood, A., Gruhn, J., Hoffmann, E.,

Broman, K. W., & Hunt, P. (2021). Failure to recombine is a common

feature of human oogenesis. *American Journal of Human Genetics, 108*(1):

16–24.

Hill, W., & Weir, B. (2011). Variation in actual relationship as a

consequence of Mendelian sampling and linkage. *Genetics Research, 93*(1),

47–64.

Karlin, S., & Liberman, U. (1978). Classifications and comparisons of

multilocus recombination distributions. *Proceedings of the National*

Academy of Sciences U.S.A., 75(12): 6332–6336.

Kivikoski, M., Rastas, P., Löytynoja, A., & Merilä, J. (2021). Automated

improvement of stickleback reference genome assemblies with Lep-Anchor

software. *Molecular Ecology Resources, 21*(6): 2166–2176.

Koehler, K. E., Hawley, R. S., Sherman, S., & Hassold, T. (1996).

Recombination and nondisjunction in humans and flies. *Human Molecular*

Genetics 5(1): 1495–1504.

Kosambi, D. D. (1944). The estimation of map distances from

recombination values. *Annals of Eugenics, 12*(1): 172–175.

Leder, E. H., McCairns, R. J. S., Leinonen, T., Cano, J. M., Viitaniemi, H.

M., Nikinmaa, M., Primmer, C. R., & Merilä, J. (2015) The evolution and

adaptive potential of transcriptional variation in sticklebacks – signatures of

selection and widespread heritability. *Molecular Biology and Evolution 32*:

674–689.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth,

G., Abecasis, G., & Durbin, R. (2009). The sequence alignment/map format

and SAMtools. *Bioinformatics, 25*: 2078–2079.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly

contigs with BWA-MEM. Preprint at http://arxiv.org/abs/1303.3997. 1– 3.

Liu, H., Zhang, X., Huang, J., Chen, J. Q., Tian, D., Hurst, L. D., & Yang,

S. (2015). Causes and consequences of crossing-over evidenced via a high-

resolution recombinational landscape of the honey bee. *Genome Biology,*

16: 15.

Lynch, M., & Walsh, B. (1998). Genetics and Analysis of Quantitative

Traits. Sinauer Associates, Sunderland, MA, pp 393–398.

Martin, S. H., Davey, J. W., Salazar, C., & Jiggins, C. D. (2019).

Recombination rate variation shapes barriers to introgression across

butterfly genomes. *PLoS Biology, 17*(2): e2006288.

Mather, K. (1938). Crossing-over. *Biological Reviews, 13*(3): 252–292.

Otto, S. P., & Payseur, B. A. (2019) Crossover interference: Shedding light

on the evolution of recombination. *Annual Reviews of Genetics,* 53: 19–44.

Peichel, C. L., Sullivan, S. T., Liachko, I., & White, M. A. (2017)

Improvement of the threespine stickleback genome using a Hi-C-based

proximity-guided assembly. *Journal of Heredity, 108*: 693–700.

Pritchard, V, L., Viitaniemi, H. M., McCairns, R. J. S., Merilä, J., Nikinmaa,

M., Primmer, C. R., & Leder, E. H. (2017). Regulatory architecture of gene

expression variation in the threespine stickleback *Gasterosteus aculeatus.*

G3: Genes, Genomes, Genetics, 7(1): 165–178.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R.,

Bender, D., Maller, J., Sklar, P., de Bakker, P. I. W., Daly, M., & Sham, P.

C. (2007). PLINK: A tool set for whole-genome association and population-

based linkage analyses. *American Journal of Human Genetics, 81*(3): 559–

575.

Rastas, P., Calboli, F. C. F., Guo, B., Shikano, T., & Merilä, J. (2016).

Construction of ultradense linkage maps with Lep-MAP2: Stickleback F2

recombinant crosses as an example. *Genome Biology and Evolution, 8*(1):

78–93.

Rastas, P. (2017). Lep-MAP3: robust linkage mapping even for low-

coverage whole genome sequencing data. *Bioinformatics, 33*(23): 3726–

3732.

R Core Team (2018) R: A Language and Environment for Statistical

Computing (R Foundation for Statistical Computing, Vienna, Austria).

Sandor, C., Li, W., Coppieters, W., Druet, T., Charlier, C., & Georges, M.

(2012) Genetic Variants in *REC8*, *RNF212*, and *PRDM9* influence male

recombination in cattle. *PLoS Genetics, 8*(7): e1002854.

Stapley, J., Feulner, P. G. D., Johnston, S. E., Santure, A. W., & Smadja, C.

M. (2017). Variation in recombination frequency and distribution across

eukaryotes: patterns and processes. *Philosophical Transactions of the Royal*

Society B 372: 20160455.

Sturtevant, A. H. (1915). The behavior of chromosomes as studied through

linkage. *Z. Abstam. Vererbung,* 13: 234–287.

Veller, C., Kleckner, N., & Nowak, M. A. (2019). A rigorous measure of

genome-wide genetic shuffling that takes into account crossover positions

and Mendel's second law. *Proceedings of National Academy of Science*

U.S.A. 116 (5) 1659–1668.

Veller, C., Edelman, N. B., Muralidhar, P., & Nowak, M. A. (2020).

Variation in genetic relatedness is determined by the aggregate

recombination process. *Genetics, 216*(4): 985–994.

Veller, C., Wang, S., Zickler, D., Zhang, L. & Kleckner, N. (2022).

Limitations of gamete sequencing for crossover analysis. *Nature* 606: E1–

E3.

Visscher, P. M., Medland, S. E., Ferreira, M. A. R., Morley, K. I., Zhu, G.,

Cornes, B. K., Montgomery, G. W., & Martin, N. G. (2006). Assumption-

free estimation of heritability from genome-wide identity-by-descent

sharing between full siblings. *PLoS Genetics* 2(3): e41.

Wang, S., Hassold, T., Hunt, P., White, M. A., Zickler, D., Kleckner, N., &

Zhang, L. (2017). Inefficient crossover maturation underlies elevated

aneuploidy in human female meiosis. *Cell, 168*(6): 977–989.

Weeks, D. E. (1994) Invalidity of the Rao map function for three loci.

Human Heredity, 44: 178–180.

Weinstein, A. (1936). The theory of multiple-strand crossing over. *Genetics,*

21(3): 155–199.

Yang, L., Gao, Y., Li, M. Park, K.-E., Liu, S., Kang, X., Liu, M., Oswalt,

A., Fang, L., Telugu, B. P., Sattler, C. G., Li, C., Cole, J. B., Seroussi, E.,

Xu, L., Yang, L., Zhou, Y., Li, L., Zhang, H., Rosen, B. D., Van Tassell, C.

P., Ma, L., & Liu, G. E. (2022). Genome-wide recombination map

construction from single sperm sequencing in cattle. *BMC Genomics 23*:

181.

Yu, K., & Feingold, E. (2001). Estimating the frequency distribution of

crossovers during meiosis from recombination data. *Biometrics, 57*(2), 427–

434.

Zhang, L., Liang, Z., Hutchinson, J., & Kleckner, N. (2014) Crossover

patterning by the Beam-Film model: Analysis and implications. *PLoS*

Genetics, 10(1): e1004042.

Zhao, H., Speed, T. P. (1996). On genetic map functions. *Genetics, 142*(4):

1369–1377.

Zickler, D., & Kleckner, N. (2015). Recombination, pairing, and synapsis of

homologs during meiosis. *Cold Spring Harbor perspectives in biology*, *7*(6):

a016626.

Figures

A

B

	A-B, B-C, C-D, D-E	A-C, C-E	A-D, B-E	A-E	B-D
p0	0.5	0	0	0	0.25
p1	0.5	1	0.5	0	0.5
p2	0	0	0.5	1	0.25
Expected no. crossovers	0.5	1	1.5	2	1
Recombination frequency	0.25	0.5	0.5	0.5	0.375
Additive map distance (cM)	25	50	75	100	50
Haldane map distance (cM)	34.7	Inf	Inf	Inf	69.3
Kosambi map distance (cM)	27.5	Inf	Inf	Inf	48.7

C

Fig. 1 Number and location of crossovers affect the recombination frequency. (A) An example of a chromosome with always two crossovers *in the bivalent* so that one occurs in the yellow and the other one in the blue area. The locations of the crossovers within their distinct regions are independent. (B) The table shows the probability for 0, 1, or 2 crossovers between the markers (p_0, p_1, and p_2, respectively) and the expected number

of crossovers *in the bivalent*, the resulting recombination frequency of the

marker pair (*in the gamete*) and the estimated map distance for every marker

pair using different mapping functions. (C) Graph showing the relationship

between the recombination frequency and map distance with different

mapping functions. Notably, intervals A-C, A-D, and A-E have the same

recombination frequencies but different map distances, whereas marker

pairs A-C and B-D have the same map distances but different recombination

frequencies.

Fig. 2 Observed spatial distribution of maternal crossovers in gametes with

two maternal crossovers. (A) Each dot corresponds to one offspring (n=265)

and coordinates show the map positions of the two crossovers, 'A' referring

to the crossover closer to chromosome start and 'B' to the one closer to

chromosome end (r=0.029, p=0.64). (B) Density plots show the distribution

of the two crossovers. Densities are calculated in 5 cM windows. In both

panels, the dashed-line rectangles show the expected distribution of

crossovers expected under the model presented here.

Fig. 3 Observed (gametic) and inferred (bivalent) crossover frequency
distributions in the nine-spined stickleback (*P. pungitius*), three-spined
stickleback (*G. aculeatus*), and human (*H. sapiens*). Each bar shows the
proportion of offspring with a certain number of crossovers (Observed) and
the inferred proportions of meiosis with a certain number of crossovers in
the bivalent in maternal and paternal meioses (Inferred). The inference is
based on the expectation-maximization algorithm by Yu and Feingold
(2001). Chromosomes are ordered from shortest to longest by length in base
pairs. Crossover counts 6–11 and 12–21 are grouped for readability.

Fig. 4 Examples of empirical recombination frequencies (left panels) and

predictions by the new function (right) in the nine-spined stickleback (top),

three-spined stickleback (middle), and human (bottom). Solid lines show the

three inverse mapping functions. Each orange and blue dot is a marker pair

in maternal and paternal data, respectively.

Tables

Table 1. Means of a per-marker pair absolute error in the predicted
recombination frequencies for both sexes of the three study species.
815
816

Species	Sex	Haldane	Kosambi	Linear	$p_0(k)$
Human	Paternal	0.0282	0.00936	0.00853	0.00375
	Maternal	0.0257	0.00709	0.00823	0.00346
Nine-spined stickleback	Paternal	0.0104	0.00442	0.00267	0.00302
	Maternal	0.0417	0.0152	0.0154	0.00910
Three-spined stickleback	Paternal	0.00867	0.00399	0.00311	0.00232
	Maternal	0.0438	0.0167	0.0139	0.00981

818 **Table 2.** Intra-chromosomal components of genetic shuffling ($\bar{r}$) for
819 autosomes. The five columns show $\bar{r}$ calculated from the empirical
820 recombination frequencies and those predicted by the four functions.
821

Species	Sex	Haldane	Kosambi	Linear	$p_0(k)$	Empirical
Human	Paternal	0.012799	0.014872	0.016796	0.015558	0.015882
	Maternal	0.018091	0.020133	0.021721	0.020672	0.020831
Nine-spined stickleback	Paternal	0.007945	0.009261	0.01041	0.009627	0.009819
	Maternal	0.01243	0.01449	0.01646	0.01509	0.01538
Three-spined stickleback	Paternal	0.007979	0.009292	0.010406	0.009733	0.009648
	Maternal	0.011867	0.013869	0.015800	0.014449	0.014815

823
824