Promising potential of high-throughput molecular phenotyping of

freshwater fishes for environmental assessment

Benjamin MARIE[1,], Julie LELOUP[2], Aurélie BONIN[3], Alison GALLET[1], Maud MOUCHET[4]*

[1] UMR 7245, CNRS/MNHN, Molécules de Communication et Adaptation des Micro-organismes (MCAM), équipe
"Cyanobactéries, Cyanotoxines et Environnement", 12 rue Buffon - CP 39, 75231 Paris Cedex 05, France.
[2] 7618 iEES-Paris Sorbonne Université - 4 place Jussieu - 75252 Paris Cedex 05, France.
[3] UMR 5553, CNRS/Université Joseph Fourrier, Laboratoire d'Écologie Alpine (LECA), BP 53, 38041 Grenoble
Cedex-9, France.
[4] UMR 7204, CNRS, SU, MNHN Centre d'Ecologie et des Sciences de la Conservation (CESCO), 43 rue Buffon - CP
135, 75231 Paris Cedex 05, France.
13

* Corresponding author: benjamin.marie@mnhn.fr

Abstract:

The recent democratisation of high-throughput molecular phenotyping allows the rapid
expansion of promising untargeted multi-dimensional approaches (*e.g.* epigenomics,
transcriptomics, proteomics, metabolomics, as well as microbiome metabarcoding), that
now represent innovative perspectives for environmental assessments. Indeed, when
developed for ecologically relevant species, these emerging "omics" analyses may present
valuable alternatives for the development of novel generations of ecological indicators, that
in turn could provide early warnings of eco(toxico)logical impairments.
This pilot study investigates the bio-indicative potential of different multi-metric tools based
on different high-throughput molecular phenotyping approaches (*i.e.* metabarcoding of the
intestine microbiome, and liver metabolomics by nuclear magnetic resonance (NMR) and
liquid chromatography coupled to high resolution mass spectrometry (LC-HRMS) on two
sentinel fish species (*Perca fluviatilis* and *Lepomis gibbosus*) from a set of eight water bodies
of the peri-urban area of Paris (France). We show that the LC-MS metabolome dataset
allows remarkably clear separation of individuals according to the species but also according
to their respective sampling lakes. Interestingly, the similar variations of *Perca* and *Lepomis*
metabolomes occur locally indicating that local environmental constraints drive the
observed metabolome variations beyond their obvious genetic differences. Thus, the
development of such reliable molecular phenotyping for environmental monitoring
constitutes a promising and innovative bio-indicative tool for environmental assessment.

Keywords: microbiome metabarcoding, untargeted metabolomics, environmental
assessment, biomarkers, sentinel species

1. INTRODUCTION

In the past decades, the use of bio-indicator aquatic organisms has been widely adopted in order to monitor, maintain or improve the quality of water bodies worldwide.[1] In parallel, the difficulties and the rational limits of direct chemical monitoring for providing sufficient information to adequately assess the risks from anthropogenic chemicals in the environment have been increasingly recognized.[2]

Recent years have offered intense implementation of molecular techniques in a wide variety of ecology research fields. Modern bio-monitoring and bio-assessment may greatly benefit from the specific development of high-throughput methods that potentially provide reliable, high quantity and quality standardized biomolecular data in a cost- and time-efficient way.[3-4]

Fishes, in particular, present unique features that make them especially relevant for bio-indication purposes. They indeed occupy in almost any aquatic habitat, as they are capable of experiencing different and variable environmental conditions.[5] Fishes are also responsive to numerous abiotic (*e.g.* temperature, water velocity, sediment load, hypoxia) or biotic (*e.g.* famine, predation, parasitism) pressures, together with anthropogenic stressors such as contaminants, that represent additional constraints that fish may experience in disturbed ecosystems.[6] Environmental stressors are known to induce biological variations, in relation to the specific physiological, developmental and reproductive capabilities of the different fish species allowing them to colonize and occupy ecosystems.[7] Thus, fishes appear to be valuable bio-indicators and their individual or community responses to environmental pressures, therefore, might have a high ecological relevance.

In Europe, for example, according to the Water Framework Directive (WFD), local ecological quality can be assessed by comparing the community structure (of fishes, but also of diatoms, sea-grasses, macro-invertebrates, etc.) in impacted sites to those observed in reference sites, using various biological quality indices.[8] However, experts face several important challenges originating from the main limitation of different diversity indexes based on fish guild: they remain especially time-consuming, requires costly sampling procedures, and can even exhibit low responsiveness to field variation.[9-10] One other important drawback of fish diversity-based bio-indicative indexes is that they also present a high sensitivity to species migration, human transplantation or fishing activities, that critically compromise their predictive power.[11] On the other hand, fish bio-indication can be

also considered through various measurable biological traits (*e.g.* population structure,

genetic or functional diversity, physiological or biological parameters). Some of them have

been already implemented, including morphometrical features,[12] punctual biochemical

markers,[13] or genetic diversity.[7] However, these attempts have also emphasized their

relative limitations in terms of operational predication ability.[14]

The recent democratisation of high-throughput molecular phenotyping allows the rapid

expansion of promising untargeted multi-dimensional approaches (*e.g.* epigenomics,

transcriptomics, proteomics, metabolomics, as well as microbiome composition by

metabarcoding). These constitute many early responding biological traits, and represent

promising perspectives for environmental assessments.[15-16] In this context, it is very likely

that emerging "omics"-based analyses, developed for ecologically relevant fish species, may

present valuable alternative for the development of new ecological evaluation tools, that

could provide even earlier warning of eco(toxico)logical impairs.[17]

To this end, the present pilot study investigates the bio-indicative potential of different

multi-metric tools based on different high-throughput molecular phenotyping approaches

(*i.e.* metabarcoding of the intestine microbiome and liver metabolomics by NMR and LC-MS)

on two sentinel fish species (*Perca fluviatilis* and *Lepomis gibbosus*) from a set of eight water

bodies of the peri-urban area of Paris (France). We hypothesize that molecular phenotyping

approaches can be exploited for environmental diagnosis, and provide valuable multi-metric

characters indicative of ecological conditions for both evaluation or prediction purposes,

together with a prospective mechanistic understanding that allows cause-effect relationship

investigation. We compare here the potential phenotypic bio-indication of two sentinel

species, as these organisms could present different or comparable sensitivity and

responsiveness to local environmental constraints, that may reinforce the proof of concept

of their respective interest for "omics"-based bio-indication.

2. MATERIALS AND METHODS

2.1. Fish sampling

Field sampling campaigns were performed during late summer 2015 (7-10[th] September) in
eight peri-urban pounds around the Paris' area (Île-de-France region, France), chosen for
their respective eutrophication levels and the presence or the absence of recurrent
cyanobacterial blooms: Cergy-Pontoise (Cer), Champs-sur-Marne (Cha), Maurepas (Mau),
Rueil (Rue), Verneuil (Ver), Varennes-sur-Seine (Var), Fontenay-sur-Loing (Fon), and Triel
(Tri) pounds (Supplementary figure S1).[18-19] These sites were sampled by the Hydrosphère
company (www.hydrosphere.fr) with an electric fishing device (FEG 8000, EFKO, Leutkirch,
Germany) for capturing fish alive. The investigation of the fish guild indicates that only the
perch (*Perca fluviatilis*) and pumpkinseed sunfish (*Lepomis gibbosus*) were present in all or
almost all these pounds (supplementary table S1) and were thus selected as sentinel species
for further molecular phenotyping analyses by metabarcoding and metabolomics.
Briefly, alive caught fishes (n=5-10 young-of-the-year per pounds and per species) were
directly measured (12.0±4.8 cm), weighed (9.3±2.6 g), briefly euthanized by neck dislocation
and then liver and intestine of each individual was shortly sampled, flash-frozen in liquid
nitrogen and kept at -80°C until analyses, in accordance with European animal ethical
concerns and regulations.
In every lake, sub-surface chlorophyll-*a* equivalent concentrations attributed to the four
main phytoplankton groups (Chlorophyta, Diatoms, Cyanobacteria and Cryptophyta) were
measured with an *in-situ* fluorometer (Fluoroprobe II, Bbe-Moldenke, Germany). Sub-surface
water samples filtered on 20-μm mesh size were also collected for phytoplanktonic
community analysis and further metabolomics characterisation, and then kept at -80°C until
analysis.
125

2.2. Intestinal microbiome sequencing and OTU analyses
Intestinal microbiome DNA was extracted from each fish individual (4-5 replicate fishes per
pound), stored and filtered following Mouchet et al..[20] After DNA concentration and purity
checking, a 295 pb fragment of the 16S rRNA-encoding gene corresponding to the V5-V6
variable region was amplified using 789-F and 1080-R primers and sequenced on an Illumina
MiSeq platform (2x250 bp, Fasteris, Switzerland). Raw reads analysis was processed through
the FROGS pipeline (Find Rapidly OTU with Galaxy Solution) implemented on a galaxy
instance (2.3.0) (https://galaxy.migale.inra.fr/). Sequences were dereplicated before
clustering using SWARM algorithm (2.1.5) with, at first, a denoising step using an

aggregation distance equal to 3.[21] The taxonomic assignation of each cluster was performed

using the BLAST tools. OTU table was rarefied, normalized by Total Sum Scaling using

MicrobiomeAnalyst platform.[22]

2.3. Liver metabolite extraction and metabolomics analyses

The liver extraction was performed on 132 individuals (comprising 78 *Perca* and 54 *Lepomis*)

with methanol/chloroform/water (ratio 2/2/1.8 – 22 mL.g^{-1} at 4°C) and the polar fraction

was analyzed on a 600-MHz NMR spectrometer equipped with a 5-mm cryoprobe (Advance

III HD Bruker, Germany) with a noesygppr1d pulse sequence as previously described.[19] ^{1}H-

NMR spectra were treated with the Batman R package for deconvolution, peak assignment

and quantification of 222 putative metabolites.[23]

The liver extracted polar phase was additionally injected (2 µL) on C_{18} column (Polar

Advances II 2.5 pore - Thermo), then eluted at a 300 µL.min^{-1} flow rate with a linear gradient

of acetonitrile in 0.1% formic acid (5 to 90 % in 21 min) with a ultra-high performance liquid

chromatography (UHPLC) system (ELUTE, Bruker). Consecutively, the individual metabolite

contents were analyzed using an electrospray ionization hybrid quadrupole time-of-flight

(ESI-Qq-TOF) high-resolution mass spectrometer (Compact, Bruker) at 2 Hz speed on positive

MS mode on the 50–1500 *m/z* range. The feature peak list was generated from recalibrated

MS spectra (< 0.5 ppm for each sample, as an internal calibrant of Na formate was injected

at the beginning of each sample analysis) within a 1-15 min window of the LC gradient, with

a filtering of 5,000 count of minimal intensity, a minimal occurrence in at least 50% of all

samples, and combining all charge states and related isotopic forms using MetaboScape 4.0

software (Bruker).

Additionally, five pools of six different individuals randomly selected for *Perca* and *Lepomis*

(quality check samples) were similarly eluted then analyzed on positive autoMS/MS mode at

2-4 Hz on the 50-1500 *m/z* range for further metabolite annotation. Molecular networks

were performed with the GNPS (http://gnps.ucsd.edu) and/or MetGem

(http://metgem.github.io) softwares, as previously described[24] and metabolite annotation

was assayed using GNPS, Mona, HMDB and Massbank MS/MS spectral libraries.

Phytoplankton samples of each water pound, concentrated with a 20-*µ*m mesh size, were

also extracted with 75% methanol (2 min sonication, 5 min centrifugation at 15,000 g - 4°C)

and then similarly was analyzed on LC-HRMS on autoMS/MS positive mode, as earlier
described, and the metabolite list was annotated with the same pipelines.

2.4. Data matrix treatment

The resulting count table of 87 OTUs (microbiome metabarcoding), and the intensity data
tables of the 222 metabolites (^{1}H NMR) and 1252 analytes (LC-MS) were further treated for
quantile normalization and Pareto's scaling, data representation by PCA and heatmap with
hierarchical clustering and then analyzed to investigate the influences of "Species" (*Perca* or
Lepomis) "Lakes" (Che, Cha, Fon, Mau, Rue, Tri, Var or Ver) parameters on the datasets by
partial least square – differential analysis (PLS-DA) using the MixOmics R Package,[25] the
MetaboAnalyst 4 tool [26] and the MicrobiomeAnalyst platform.[22] The PLS model
performances were determined by accuracy, performance (R^2) and predictability (Q^2) values,
and classification error.

3. RESULTS AND DISCUSSION

3.1. Global informativeness of the three molecular datasets on *Perca* and *Lepomis*

Figure 1 illustrates the global relative distribution of OTUs, metabolites observed by NMR
and by LC-MS for each fish (comprising both *Perca* and *Lepomis*) on a heatmap with
hierarchical classifications and individual plot principal component analyses (PCA). Based on
the relative distribution of the 87 identified OTUs within fish individuals (44 *Perca* and 34
Lepomis), no reliable discrimination could be observed according to the species and lakes
(Fig. 1A and D). Although the 222 potential metabolites quantified by NMR on 128 fish (78
Perca and 54 *Lepomis*) present no obvious distinction for neither the "species" nor "lakes"
variables (Fig. 1B and E), the 1,252 analytes semi-quantified from the liver metabolomes by
LC-MS clearly present more discriminant patterns between species and lakes (Fig. 1C and F).
Additional investigations on those datasets were then performed using supervised models.
While unsupervised PCA were first used to evaluate the global dispersion between species
and sampling groups, a supervised model such as partial least square differential analysis
(PLS-DA) (Supplementary figure S2; supplementary tables S2, S3 and S4) allows to maximize
the separation between sample classes (*i.e.* "species" and "lakes") and to extract
information on discriminating features (variable importance on projection - VIP).

Interestingly, these differential analyses indicate that the microbiome metabarcoding
dataset allows a faint but noteworthy discrimination among the intestinal microbiomes
observed within individuals of the two species (R^2 = 0.54), but with only a limited
predictability of the models (Q^2 = 0.13 and classification error = 0.32) (Supplementary figure
S2A), while no reliable microbiome discrimination could be retrieved for the "lake" variable
according to PLS-DA investigation (Supplementary figure S2D). Concerning the ^{1}H NMR liver
metabolomics dataset, the PLS-DA model provide a comparable discrimination between
species (R^2 = 0.51 and Q^2 = 0.38) (Supplementary figure S2B), but could neither allow reliable
"lake" discrimination or prediction (Supplementary figure S2E). On the other hand, a very
strong discrimination for the "species" was achieved by PLS-DA for the LC-MS liver
metabolomics dataset (R^2 = 0.96 and Q^2 = 0.96, classification error = 0) (Supplementary
figure S2C). Only a slight distinction among the "lakes" can be observed for the LC-MS
dataset (R^2 = 0.34 and Q^2 = 0.24; classification error = 0.84) (Supplementary figure S2F),
whereas both the heatmap with hierarchical clustering (Figure 1C) and the individual plot
from the un-supervised PCA (Figure 1F) show remarkable grouping of fishes from the same
species collected in the same lakes. This surprising lack of performance of the PLS-DA
performed with grouped LC-MS data from the two fish species, *Perca* and *Lepomis*, is very
likely due to species variation among the different lakes. Indeed, this hypothesis is further
explored by investigating separately the *Perca* and *Lepomis* locality differences on their
respective intestinal microbiomes and liver metabolomes.
218
3.2. Comparison of *Perca* and *Lepomis* molecular signature variation among lakes
Considering separately the 16S rDNA metabarcoding datasets of the two fish species, the
intestine microbiome of *Perca* and *Lepomis* shows only faint specificity within the different
sampling lake groups, as investigated by both un-supervised and supervised multivariate
analyses and hierarchical classification (Supplementary figure S3). Indeed, the different
performance scoring retrieved for PLS-DA remains distinctly low, and does not support an
important structuration of the data according to the "lake" variable, whatever the species
considered. However, in the present case, the limited sample number per sampling lake
(n=4-5) could constitute a significant limitation to the informativeness of the intestinal
microbiome for local discrimination (supplementary figure S3).
The intestinal microbiome constitutes a remarkable interface between the organism and the

surrounding environment that is involved in various fish biological processes including
digestion, immunity or ecotoxicology.[27] The analysis of its microbial composition, in terms of
genotype occurrence, density and taxonomy represent remarkable biological traits for the
fish host that aim at characterizing the functionality and the ecology of this specific intestinal
microbial ecosystem.

Although the intestinal microbiome is the object of numerous investigations in microbial
ecology and related host disease/health studies,[28] this compartment has been rarely
included in environmental biomonitoring programs. However, changes in the microbial
intestinal composition may denote significant environmental variations that have potentially
altered the host homeostasis and may be influenced by the specific responsiveness of the
different fish species. However, microbial diversity estimated from 16S-rRNA sequences
does not allow taking into consideration that different genotypes can present functional
redundancy within the intestinal microbial ecosystem,[29] or alternatively, that
undistinguishable OTUs (regarding only V5-V6 16S-rRNA fragment) can mask different sub-
taxa with distinct physiological capacities, as already exemplified for *Vibrio* sp..[30] Taken
together, these technical issues can represent noticeable limitations for the use of microbial
diversity retrieved from OTU sequencing for environmental assessment, and may require
additional function-based approaches for microbial community exploration.[27] Also, one of
the most pregnant limitation for the democratisation of microbiome metabarcoding
analyses is maybe its financial cost, that still remains too important for its integration in the
development of affordable biomonitoring solutions.

Concerning the [1]H NMR metabolomics datasets of the two fish species taken separately, the
liver metabolome of *Perca* and *Lepomis* shows only a limited specificity in relation to the
"lakes". Indeed, according to the relatively low performances of the hierarchical
classifications and the discriminant analyses, the NMR liver metabolomes present restricted
discrimination on "lakes" for both species and could also support only a limited separation
predictability of the supervised models (Supplementary figure S4; supplementary table S3).
The [1]H NMR metabolomics allow the reliable quantification of main liver metabolites
(belonging mostly to amino acids, sugars, TCA metabolites, etc.), by detecting characteristic
chemical groups, with regards to their respective position within the molecules.[31] These
primary metabolites are key contributors to the cellular metabolism and are supposed to be

responsive to variable physiological conditions of fish encountering peculiar biological
outcomes and/or environmental constraints.[32] However, this approach remains focused on a
few principal metabolites involved in main cellular pathways or regulation processes, and
not being specific of subtle molecular regulation processes. Moreover, ^{1}H NMR presents only
low sensitivity and limited discrimination capability for molecules presenting similar
chemical structures, such as lipids, that exhibit an aggregated and unspecific signal on NMR
spectra.[33] For this reason, it is supposed to present lower discriminating power than less
quantitative, but more selective and more sensitive, mass spectrometry-based
metabolomics approaches.
271
In contrast, the LC-MS dataset of the two fish species considered separately clearly shows
reliable discrimination of the metabolomes of fishes originating from the different sampling
lakes (Figure 2; supplementary figure S5; supplementary figure S6). As illustrated with the
un-supervised PCA and heatmap with hierarchical classification (Figure 2), the liver
metabolomes analyzed by LC-MS present distinct local signatures that support reliable
discriminant analysis models for both *Perca* and *Leptomis*, with significant performance
scoring of the PLS-DA (R^2_{Perca} = 0.46 and Q^2_{Perca} = 0.35; $R^2_{Lepomis}$ = 0.50 and $Q^2_{Lepomis}$ = 0.38)
(supplementary figure S5; supplementary figure S6). Taken together, these observations
indicate that the locality seems to globally influence the LC-MS metabolome composition for
both species.
However, as previously observed when comparing LC-MS datasets of both fish species
together, the *Perca* and *Lepomis* liver metabolomes of the Fontenay-sur-Loing lake (Fon)
exhibited a quite species-specific trajectory, with regards to the relative composition of their
metabolomes (Figure 2A and 2D). This suggests that, although the specific "lake" signature
of the metabolome appears in good agreement between the two species for most localities,
in some environments, local specificities are variable from one species to the other one, as
represented here for the local species-specificity of the *Perca* and *Lepomis* metabolomes in
Fon lake.
Interestingly, the hierarchical classifications performed respectively on the *Perca* and
Lepomis LC-MS metabolome datasets show very similar lake relationships, grouping together
the fish from the lakes of Cer and Mau, from Cha, Fon and Rue, and from Tri, Var and Ver
(for *Perca*) or Tri and Ver for *Lepomis*, as this latter fish species was not retrieved in Var

(Figure 2B and 2E). Then, we re-explored the fish metabolome discrimination considering

together fish from the "Cer/Mau", "Cha/Fon/Rue" and "Tri/Var/Ver" (or "Tri/Ver" for

Lepomis) groups, by PCA (supplementary figure S7), and PLS-DA (Figure 2C and 2F). For both

Perca and *Lepomis* the metabolomes present an even better discrimination scoring (*Perca*:

accuracy = 0.77, R^2 = 0.79, Q^2 = 0.70; *Lepomis*: accuracy = 0.70, R^2 = 0.73, Q^2 = 0.60) and

discriminant analysis predictive error (Class. error$_{Perca}$ = 0.04; Class. error$_{Lepomis}$ = 0.10) for

these three groups than when considering the different lakes separately. Remarkably, the

observation of the metabolite semi-quantification of the best VIP presents largely similar

variations among these three groups of lakes for *Perca* and *Lepomis* (Table 1; supplementary

figure S8).

In addition, the LC-MS metabolite annotation, supported by molecular networking analysis

(supplementary figure S9; table 1), shows that fish collected from the Cer/Mar lake group

seem to present greater energetic and anti-oxidant/detoxification reserves (*e.g.* adenosine

and oxidized glutathione) than in other lake groups. In addition, the molecular networking

annotation process highlights that most, if not all, of the known metabolites (presenting

structural identity or analogy hits) are shared between the two species (*e.g.* saccharides,

nucleic acids, carnitines, glutathiones, lipids) when most of the species-specific cluster

metabolites remains uncharacterized, as corresponding molecules present no match within

public chemical databases (supplementary figure S9).[34] This representation remarkably

illustrates the portion of the specific liver molecular metabolism that remains to be

characterized for these two species.

In general, various genetic or phenotypic factors (such as development stages,

contamination levels, predator/parasite pressures, food availability, etc.) could influence and

explain local discrepancies of the metabolome of fishes collected from different

environments.[17] However, in the present case, the fact that *Perca* and *Lepomis*

metabolomes, analyzed in parallel, present similar metabolite variations clearly indicates

that local environmental constraints drive such observed phenotypic co-variations (beyond

the obvious genetic distinction between *Perca* and *Lepomis*). Then, these present results

claim for the great value of LC-MS-based metabolomic imprints for molecular phenotyping

on sentinel species and the exploration of environmental effects.[35]

Although LC-MS-based metabolomics remains a selective analytical approach (according to

both LC separation and ionisation respective properties of each molecule), it presents un-
precedent analytical capabilities in terms of sensitivity, dynamic range and number of
characterized components and it now offers unprecedented perspectives for the
investigation for molecular phenotyping applied to environmental sciences.[36] We assume
that a single LC-MS fingerprint analysis does not embed the whole metabolite picture of the
biological compartment (the fish liver, in the present case), especially because of its
selectivity performances.[37] However, it provides a specific and precise measurement of a
large number of components that can endorse high-throuput and in-depth phenotyping.
For comparison, ^{1}H NMR metabolomics allows the global quantification of the CH_2-CH_2 or
CH_2-CH=CH fatty acid bounds,[38] whereas LC-MS metabolomics can discriminate and
characterize hundreds of different lipids belonging to various sub-classes,[39] providing
undoubtedly more informative and discriminant features. In addition, multi-omics
integration tools, such as Mixomics,[40] provide innovative solutions for the comparison of
heterogeneous dataset matrices obtained from a single set of samples or individuals, and
may contribute to investigate metabolomics biomarker relevance (supplementary figures
S10-S11).

3.3. Use of LC-MS metabolomics fingerprint for environmental assessment?
Although the value of LC-MS metabolomics for investigating the impacts of environmental
stressors or contaminants, and their respective modes of action, has been well-explored in
medical sciences[41] or in ecotoxicology laboratory-based studies on aquatic models,[42-45] such
methods have been used only faintly in field research so far.[46-47] Apart from a limited
number of evidence on the utility of NMR-based metabolomics in environmental fish
studies,[17;48-49] few other examples indicate that field-based LC-MS metabolomics constitute
an emerging and powerful, but still underused, approach for increasing our understanding of
in situ biological, physiological, ecological or ecotoxicological processes.[50-52]
Our analysis constitutes one of the first attempts to push forward the potential of high-
throughput molecular phenotyping, and especially through LC-MS-based metabolomics, for
environmental assessment. Indeed, this organismal molecular phenotyping supported by
multi-variable chemometric investigation offers remarkably rich biological information that
serves at describing specific phenotypic plasticity. Moreover, this organismal
variability/responsiveness can further be confronted to local environmental factors in order

to search for correlation/causality relationships (supplementary figure S12). This effort

pushes a step further into the objective environmental omics-assisted assessment relying on

field data modelling using artificial intelligence decision-supporting tools.[17]

As described by Pompfret and co-workers,[53] environmental metabolomics exhibits very

promising perspectives for operational bio-monitoring applications, because of its reliability,

its reproducibility, and its high predictive potential. However, these authors also point out

that the responsiveness and robustness of the bio-indicative object, that is characterized

through the analytical prism of the metabolomics, remains crucial and has to be carefully

evaluated and tested with an appropriate experimental design.[53] On the other hand,

freshwater fishes are already considered as good indicators of the ecological status of

aquatic ecosystems, as various fish biodiversity indexes have already been proposed and are

still under-evaluation at local or large scales.[8]

However, for ethical concerns, fish bio-monitoring would also gain at being less invasive and

deleterious for the organisms. To this end, non-lethal mucus sampling has been investigated

by LC-MS and has demonstrated the remarkable informativeness of this approach for

environmental studies.[51] In parallel, recent attempts have also been made concerning the

interest of planktonic or benthic microbial communities characterized by metabarcoding for

water quality bio-indication,[16] but these efforts still remain explorative.

The main challenge we face remains the development of more integrative approaches

connecting chemical, biological and ecological evaluations, in the context of anthropized

natural environments experiencing multi-stressor pressures. A major caveat of the use of

fish environmental metabolomics may now be the lack of dedicated databases[54] fulfilled by

studies considering together different species, populations, development stages, seasons

and environments. This pre-requisite knowledge could be used to provide baseline reference

data that would support machine learning or artificial neural network tools for the training

of decision-making models.

ASSOCIATED CONTENT

Supporting information

Map of sampled lakes (Supplementary figure S1); individual plots of PLS-DA performed with intestine metabarcoding of 16S rDNA (A and D), Liver NMR (B and E) and LC-MS (C and F) metabolomics datasets for *Perca* and *Lepomis*, discriminated according to the "species" (A-C) and "sampling lakes" (D-F) parameters (Supplementary figure S2); individual plots of PCA, heatmap with hierarchical clustering and PLS-DA testing the difference among the different lakes according to the intestinal microbiomes of *Perca* (A-C) and *Lepomis* (D-F) presenting insufficient accuracy, predictability and quality performances for sampling lake discrimination (Supplementary figure S3); individual plots of PCA, heatmap with hierarchical clustering and PLS-DA testing the differences among lakes according to the liver NMN metabolomes of *Perca* (A-C) and *Lepomis* (D-F) presenting insufficient accuracy, predictability and quality performances for sampling lake discrimination (Supplementary figure S4); individual plots of PLS-DA, testing the differences among lakes according to the LC-MS metabolomes of *Perca*, with corresponding best VIP list and metabolite box-plots (Supplementary figure S5); individual plots of PLS-DA, testing the differences among lakes according to the LC-MS metabolomes of *Lepomis*, with corresponding best VIP list and metabolite box-plots (Supplementary figure S6); Individual plots of PCA for LC-MS metabolomics of *Perca* (A) and *Lepomis* (B) representing the different lake groups highlighted by hierachical classication as shown on Fig. 2 (Supplementary figure S7); box-plots of top-16 best VIP metabolites according to PLS-DA performed on the lake groups indicated by hierarchical classification, as shown on Figure 2, for *Perca* (A) and *Lepomis* (B) (Supplementary figure S8); molecular networking of *Perca* and *Lepomis* metabolomes characterized by GNPS and *t*-SNE algorithms (Supplementary figure S9); Integration of the NMR and LC-MS metabolomics of *Perca* using MixOmics illustrates the comparable PLS-DA patterns of the datasets (a-b) and the global correlation of respective VIPs (correlation score = 0.79; c) (Supplementary figure S10); Integration of the NMR and LC-MS metabolomics of *Lepomis* using MixOmics illustrates the comparable PLS-DA patterns of the datasets (a-b) and the global correlation of respective VIPs (correlation score = 0.83; c) (Supplementary figure S11); phytoplankton composition estimated by BBE measurment of the eight lake sub-surface water (a), corresponding cyanobacteria relative composition for Fon, Tri, Var and Ver (B), and molecular networking of metabolites extracted from the filtered biomass of the respective water of the eight lakes generated with *t*-SNE algorithm, with cyanobacteria peptide clusters indicated in bold and microcystin cluster indicated in red (C) (Supplementary figure S12) (PDF)

List and numbers of collected fishes (Supplementary table S1); list of significant VIP discriminating the intestinal microbiomes (Supplementary table S2), the liver NMR metabolomes (Supplementary table S3) and the liver LC-MS metabolomes (Supplementary table S4) of *Perca* and *Lepomis* species determined by PLS-DA; list of significant VIP (>1.5) discriminating the liver LC-MS metabolomes of *Perca* and *Lepomis* from the different lakes determined by PLS-DA (Supplementary table S5) (XLS)

AUTHOR INFORMATION

Corresponding author

Benjamin Marie – UMR 7245, CNRS/MNHN, Molécules de Communication et Adaptation des Micro-organismes (MCAM),
équipe "Cyanobactéries, Cyanotoxines et Environnement", 12 rue Buffon - CP 39, 75231 Paris Cedex 05, France.

iD https://orcid.org/0000-0001-9880-5541; Phone : +33 1 40 79 32 12 ; Email : bmarie@mnhn.fr

Authors

Maud Mouchet – UMR 7204 CNRS/SU/MNHN, Centre d'Ecologie et des Sciences de la Conservation (CESCO), 43 rue Buffon
- CP 135, 75231 Paris Cedex 05, France.

iD https://orcid.org/0000-0001-5939-6802

Julie Leloup – UMR 7618 iEES, Paris Sorbonne Université - 4 place Jussieu - 75252 Paris Cedex 05 - France

iD https://orcid.org/0000-0002-2777-284X

Aurélie Bonin - UMR 5553, CNRS/Université Joseph Fourrier, Laboratoire d'Écologie Alpine (LECA), BP 53, 38041 Grenoble
Cedex-9, France.

iD https://orcid.org/0000-0001-7800-8609

Alison Gallet - UMR 7245, CNRS/MNHN, Molécules de Communication et Adaptation des Micro-organismes (MCAM),
équipe "Cyanobactéries, Cyanotoxines et Environnement", 12 rue Buffon - CP 39, 75231 Paris Cedex 05, France.

Author contributions

B.M. and M.M. designed experiment. B.M., A.B. and M.M. performed sample analysis. B.M., A.G. and J.L. analysed data. B.M. wrote the manuscript. All authors have given approval to the final version of the manuscript.

Notes

The authors declare not conflict of interest

ACKNOWLEDGMENTS

We would like to thanks B. Sotton for his collaboration on NMR data acquisition and treatment. We thank Kenin Tambosco for his technical assitance. We are also greatful to A. Paris for its support and prolific discussions. This work was supported by grants from CNRS (Défi ENVIROMICS "Toxcyfish"

460 project) and from Sorbone Université (CARESE-SU J15R323). The NMR and the MS spectra were
461 respectively acquired at the Plateau technique de Résonance Magnétique Nucleaire and the Plateau
462 technique de spectrométrie de masse bio-organique, Muséum National d'Histoire Naturelle, Paris,
463 France. This work benefitted from the French GDR "Aquatic Ecotoxicology" framework which aims at
464 fostering stimulating scientific discussions and collaborations for more integrative approaches.

REFERENCES

(1) Lorenz, C. M. Bioindicators for ecosystem management, with special reference to
freshwater systems. In *Trace Metals and other Contaminants in the Environment*
(**2003**, Vol. 6, pp. 123-152). Elsevier.

(2) Reyjol, Y.; Argillier, C.; Bonne, W.; Borja, A.; Buijse, A. D.; Cardoso, A. C.; ... Prat, N.
Assessing the ecological status in the context of the European Water Framework
Directive: where do we go now? *Science of the Total Environment* **2014**, *497*, 332-
344.

(3) Schimel, J. Microbial ecology: Linking omics to biogeochemistry. *Nature microbiology*
2016, *1*(2), 1-2.

(4) Paul, V. J., Freeman, C. J., & Agarwal, V. Chemical ecology of marine sponges: new
opportunities through "-Omics". *Integrative and Comparative Biology* **2019**, *59*(4),
765-776.

(5) Harris, J. H. The use of fish in ecological assessments. *Australian Journal of Ecology* **1995**,
20(1), 65-80.

(6) McClanahan, T. R.; Schroeder, R. E.; Friedlander, A. M.; Vigliola, L.; Wantiez, L.; Caselle, J.
E.; ...; Oddenyo, R. M. Global baselines and benchmarks for fish biomass: comparing
remote reefs and fisheries closures. *Marine Ecology Progress Series*, **2019**, *612*, 167-
192.

(7) Hamilton, P.; Cowx, I.; et al.; Tyler, C. R. Population-level consequences for wild fish
exposed to sublethal concentrations of chemicals–a critical review. *Fish and Fisheries*
2016, *17*(3), 545-566.

(8) Poikane, S.; Ritterbusch, D.; Argillier, C.; Białokoz, W.; Blabolil, P.; Breine, J., ...; Noges, P.
Response of fish communities to multiple pressures: development of a total
anthropogenic pressure intensity index. *Science of the Total Environment* **2017**, *586*,

502-511.

(9) Birk, S.; Bonne, W.; Borja, A.; Brucet, S.; Courrat, A.; Poikane, S.; ... & Hering, D. Three

hundred ways to assess Europe's surface waters: an almost complete overview of

biological methods to implement the Water Framework Directive. *Ecological*

Indicators **2012**, *18*, 31-41.

(10) Birk, S.; Willby, N. J.; Kelly, M. G., Bonne, W., Borja, A., Poikane, S., & Van de Bund, W.

Intercalibrating classifications of ecological status: Europe's quest for common

management objectives for aquatic ecosystems. *Science of the Total Environment*

2013, *454*, 490-499.

(11) Shin, Y. J.; Houle, J. E.; Akoglu, E.; Blanchard, J. L.; Bundy, A.; Coll, M.; ...; Salihoglu, B.

The specificity of marine ecological indicators to fishing in the face of environmental

change: a multi-model evaluation. *Ecological Indicators* **2018**, *89*, 317-326.

(12) Esin, E. V.; Edosov, A. E. The effect of chronic volcanic pollution on the morphometric

characteristics of juvenile Dolly Varden (*Salvelinus malma* W.) on the Kamchatka

peninsula. *Hydrobiologia* **2016**, *783*(1), 295-307.

(13) Zhang, Z.; Wang, J.; Pan, Z.; Zhang, Y.; Zhang, X.; Tian, H.; ... ; Ru, S. Distribution of

vitellogenin in Japanese flounder (*Paralichthys olivaceus*) for biomarker analysis of

marine environmental estrogens. *Aquatic Toxicology* **2019**, *216*, 105321.

(14) Bahamonde, P. A.; Munkittrick, K. R.; Martyniuk, C. J. Intersex in teleost fish: are we

distinguishing endocrine disruption from natural phenomena?. *General and*

comparative endocrinology, **2013**, *192*, 25-35.

(15) Bahamonde, P. A.; Feswick, A.; Isaacs, M. A.; Munkittrick, K. R.; Martyniuk, C. J. Defining

the role of omics in assessing ecosystem health: Perspectives from the Canadian

environmental monitoring program. *Environmental toxicology and chemistry*, **2016**,

35(1), 20-35.

(16) Cordier, T.; Alonso-Sáez, L.; Apothéloz-Perret-Gentil, L.; Aylagas, E.; Bohan, D. A.;

Bouchez, A.; ... Keeley, N. Ecosystems monitoring powered by environmental

genomics: a review of current strategies with an implementation roadmap. *Molecular*

Ecology. **2020**, DOI: 10.1111/mec.15472

(17) Marie, B. Disentangling of the ecotoxicological signal using "omics" analyses, a lesson

from the survey of the impact of cyanobacterial proliferations on fishes. *Science of*

The Total Environment **2020**, *736,* 139701.

(18) Maloufi, S.; Catherine, A.; Mouillot, D.; Louvard, C.; Couté, A.; Bernard, C.; Troussellier,
M. (Environmental heterogeneity among lakes promotes hyper β-diversity across
phytoplankton communities. *Freshwater Biology, 61*(5), 633-645.

(19) Sotton, B.; Paris, A.; Le Manach, S.; Blond, A.; Duval, C.; Qiao, Q.; ... ; Marie, B. Specificity
of the metabolic signatures of fish from cyanobacteria rich lakes. *Chemosphere* **2019**,
226, 183-191.

(20) Mouchet, M.; Bouvier, C.; Bouvier, T.; Troussellier, M.; Escalas, A.; Mouillot, D. Genetic
difference but functional similarity among fish gut bacterial communities through
molecular and biochemical fingerprints. *FEMS Microbiology Ecology* **2012**, *79*, 568–
580, https://doi.org/10.1111/j.1574-6941.2011.01241.x

(21) Escudié, F.; Auer, L., Bernard, M.; Mariadassou, M.; Cauquil, L.; Vidal, K.; ... Pascal, G.
FROGS: find, rapidly, OTUs with galaxy solution. *Bioinformatics* **2018**, *34*(8), 1287-
1294.

(22) Chong, J.; Liu, P.; Zhou, G.; Xia, J. Using MicrobiomeAnalyst for comprehensive
statistical, functional, and meta-analysis of microbiome data. *Nature Protocols* **2020**,
15, 799-821.

(23) Hao, J.; Liebeke, M.; Astle, W.; De Iorio, M.; Bundy, J. G.; Ebbels, T. M. D. Bayesian
deconvolution and quantification of metabolites in complex ^{1}D NMR spectra using
BATMAN. *Nature protocols* **2014**, 9(6), 1416–1427.

(24) Kim Tiam, S.; Gugger, M.; Demay, J.; Le Manach, S.; Duval, C.; Bernard, C.; Marie, B.
Insights into the Diversity of Secondary Metabolites of *Planktothrix* Using a Biphasic
Approach Combining Global Genomics and Metabolomics. *Toxins* **2019**, *11*(9), 498.

(25) Rohart, F.; Gautier, B.; Singh, A.; Lê Cao, K. A. mixOmics: An R package for 'omics feature
selection and multiple data integration. *PLoS computational biology* **2017**, *13*(11),
e1005752.

(26) Chong, J.; Wishart, D. S.; Xia, J. Using metaboanalyst 4.0 for comprehensive and
integrative metabolomics data analysis. *Current protocols in bioinformatics* **2019**,
68(1):e86.

(27) Duperron, S.; Halary, S.; Gallet, A.; Marie, B. Microbiome-aware ecotoxicology of
organisms: relevance, pittfalls and challenges. *Frontiers in Public Health* **2020**, 8, 407.

(28) Selber-Hnatiw, S., Rukundo, B., Ahmadi, M., Akoubi, H., Al-Bizri, H., Aliu, A. F., et al.
(2017). Human Gut Microbiota: Toward an Ecology of Disease. *Front. Microbiol.* 8,
1265. doi:10.3389/fmicb.2017.01265.

(29) Andam, C. P. Clonal yet different: understanding the causes of genomic heterogeneity in
microbial species and impacts on public health. *mSystems.* **2019**, *4*:e00097-19.

(30) Sawabe, T.; Kita-Tsukamoto, K.; Thompson, F.L. Inferring the evolutionary history of
vibrios by means of multilocus sequence analysis. *J Bacteriol.* **2007**, 189:7932–6.

(31) Martin, J. C.; Maillot, M.; Mazerolles, G.; Verdu, A.; Lyan, B.; Migne, C.; ... ; Manach, C.
Can we trust untargeted metabolomics? Results of the metabo-ring initiative, a large-
scale, multi-instrument inter-laboratory study. *Metabolomics* **2015**, *11*(4), 807-821.

(32) Roques, S.; Deborde, C.; Richard, N., Marchand; Y., Larroquet, L.; Prigent, S.; ...;
Fauconneau, B. Proton-NMR Metabolomics of Rainbow Trout Fed a Plant-Based Diet
Supplemented with Graded Levels of a Protein-Rich Yeast Fraction Reveal Several
Metabolic Processes Involved in Growth. *J. Nutrition* **2020**, *150*(9), 2268-2277.

(33) Emwas, A. H.; Roy, R.; McKay, R. T.; Tenori, L.; Saccenti, E.; Gowda, G. A.; ... Wishart, D.
S. NMR spectroscopy for metabolomics research. *Metabolites* **2019**, *9*(7), 123.

(34) da Silva, R. R.; Dorrestein, P. C.; Quinn, R. A. Illuminating the dark matter in
metabolomics. *Proceedings of the National Academy of Sciences* **2015**, *112*(41),
12549-12550.

(35) Lohr, K. E.; Khattri, R. B.; Guingab-Cagmat, J.; Camp, E. F.; Merritt, M. E.; Garrett, T. J.,
Patterson, J. T. Metabolomic profiles differ among unique genotypes of a threatened
Caribbean coral. *Scientific reports* **2019**, *9*(1), 1-11.

(36) Beyoğlu, D.; Idle, J. R. (2020). Metabolomic and lipidomic biomarkers for premalignant
liver disease diagnosis and therapy. *Metabolites* **2020**, *10*(2), 50.

(37) Gika, H.; Virgiliou, C.; Theodoridis, G.; Plumb, R. S.; Wilson, I. D. Untargeted LC/MS-
based metabolic phenotyping (metabonomics/metabolomics): The state of the art.
Journal of Chromatography B **2019**, *1117*, 136-147.

(38) Marchand, J.; Martineau, E.; Guitton, Y.; Le Bizec, B.; Dervilly-Pinel, G.; Giraudeau, P. A
multidimensional ^{1}H NMR lipidomics workflow to address chemical food safety
issues. *Metabolomics* **2018**, *14*(5), 60.

(39) Dreier, D. A.; Bowden, J. A.; Aristizabal-Henao, J. J.; Denslow, N. D.; Martyniuk, C. J.
Ecotoxico-lipidomics: An emerging concept to understand chemical-metabolic

relationships in comparative fish models. *Comparative Biochemistry and Physiology*

Part D: Genomics and Proteomics **2020**, *36,* 100742.

(40) Singh, A.; Shannon, C. P.; Gautier, B.; Rohart, F.; Vacher, M.; Tebbutt, S. J.; Lê Cao, K. A.

DIABLO: an integrative approach for identifying key molecular drivers from multi-

omics assays. *Bioinformatics* **2019**, *35*(17), 3055-3062.

(41) Zhang, X. W.; Li, Q. H.; Dou, J. J. Mass spectrometry-based metabolomics in health and

medical science: a systematic review. *RSC Advances*, **2020**, *10*(6), 3092-3104.

(42) Le Manach, S.; Sotton, B.; Huet, H.; Duval, C.; Paris, A.; Marie, A.; ... ; Marie, B.

(Physiological effects caused by microcystin-producing and non-microcystin

producing *Microcystis aeruginosa* on medaka fish: A proteomic and metabolomic

study on liver. *Environmental pollution* **2018**, *234*, 523-537.

(43) Huang, S. S.; Benskin, J. P.; Veldhoen, N.; Chandramouli, B.; Butler, H.; Helbing, C. C.;

Cosgrove, J. R. A multi-omic approach to elucidate low-dose effects of xenobiotics in

zebrafish (*Danio rerio*) larvae. *Aquatic toxicology* **2017**, *182*, 102-112.

(44) Ekman, D. R.; Skelton, D. M.; Davis, J. M.; Villeneuve, D. L.; Cavallin, J. E.; Schroeder, A.;

... Collette, T. W. Metabolite profiling of fish skin mucus: a novel approach for

minimally-invasive environmental exposure monitoring and surveillance.

Environmental Science & Technology **2015**, *49*(5), 3091-3100.

(45) Gil-Solsona, R., Nácher-Mestre, J., Lacalle-Bergeron, L., Sancho, J. V., Calduch-Giner, J.

A., Hernández, F., & Pérez-Sánchez, J. Untargeted metabolomics approach for

unraveling robust biomarkers of nutritional status in fasted gilthead sea bream

(*Sparus aurata*). *PeerJ* **2017**, *5*, e2920.

(46) Bundy, J. G.; Davey, M. P.; Viant, M. R. Environmental metabolomics: a critical review

and future perspectives. *Metabolomics* **2009**, *5*(1), 3.

(47) Lankadurai, B. P.; Nagato, E. G.; Simpson, M. J. Environmental metabolomics: an

emerging approach to study organism responses to environmental stressors.

Environmental Reviews **2013***, 21*(3), 180-205.

(48) Cappello, T.; Brandão, F.; Guilherme, S.; Santos, M. A.; Maisano, M.; Mauceri, A.; ...

Pereira, P. Insights into the mechanisms underlying mercury-induced oxidative stress

in gills of wild fish (*Liza aurata*) combining [1]H NMR metabolomics and conventional

biochemical assays. *Science of the Total Environment* **2016**, *548*, 13-24.

(49) Wei, F.; Sakata, K.; Asakura, T.; Date, Y.; Kikuchi, J. Systemic homeostasis in

metabolome, ionome, and microbiome of wild yellowfin goby in estuarine

ecosystem. *Scientific reports* **2018**, *8*(1), 1-12.

(50) Meador, J. P.; Bettcher, L. F.; Ellenberger, M. C.; Senn, T. D. Metabolomic profiling for

juvenile Chinook salmon exposed to contaminants of emerging concern. *Science of*

The Total Environment **2020**, *747*, 141097.

(51) Reverter, M.; Sasal, P.; Banaigs, B.; Lecchini, D.; Lecellier, G.; Tapissier-Bontemps, N. Fish

mucus metabolome reveals fish life-history traits. *Coral Reefs* **2017**, *36*(2), 463-475.

(52) Goode, K. L.; Dunphy, B. J.; Parsons, D. M. Environmental metabolomics as an ecological

indicator: Metabolite profiles in juvenile fish discriminate sites with different nursery

habitat qualities. *Ecological Indicators*, **2020**, *115*, 106361.

(53) Pomfret, S. M.; Brua, R. B.; Izral, N. M.; Yates, A. G. Metabolomics for biomonitoring: an

evaluation of the metabolome as an indicator of aquatic ecosystem health.

Environmental Reviews, **2020**, *28*(1), 89-98.

(54) Viant, M. R.; Ebbels, T. M.; Beger, R. D.; Ekman, D. R.; Epps, D. J.; Kamp, H.; ... ; Rocca-

Serra, P. Use cases, best practice and reporting standards for metabolomics in

regulatory toxicology. *Nature communications* **2019**, *10*(1), 1-10.

Legends of figures

Fig. 1. Visualization of the dataset structuration for intestine metabarcoding 16S rDNA (A), liver NMR (B) and LC-MS (C) metabolomics on heatmaps with hierarchical clustering according to Euclidean distance for *Perca* and *Lepomis* fishes collected during the 7-10th of September, 2015 within eight pounds of the peri-urban Paris' area.

Fig. 2. PCA (A and D), heatmap with hierarchical classification (B and E) and PLS-DA (C and F) of LC-MS metabolome of *Perca* (A-C) and *Lepomis* (D-F) liver according to the "sampling lake" parameter.

Table 1. Annotated VIP (score >2) of respective *Perca* and *Lepomis* LC-MS metabolomics dataset according to PLD-DA performed on lake, and corresponding variation among the here lake groups highlighted by hierarchical classification (according to Figure 2B and 2E).

650 **Fig. 1.**

653   **Fig. 2.**

654
655

656 **Table 1.**

MW (Da)	Annotation	VIP comp. 1 *Perca*	VIP comp. 1 *Lepomis*
267.09601	Adenosine	8.86 (▫ > ▫ = ▫)	15.23 (▫ > ▫ = ▫)
612.1507	Glutathione (oxidized)	6.44 (▫ > ▫ > ▫)	13.32 (▫ > ▫ > ▫)
307.08355	Glutathione (reduced)	7.02 (▫ > ▫ > ▫)	2.28 (▫ > ▫ > ▫)
383.10736	*Analog*: Cyanidin arabinosine	4.25 (▫ > ▫ > ▫)	2.67 (▫ > ▫ > ▫)
217.13223	*Analog*: Carnitine	1.55 (▫ > ▫ > ▫)	4.18 (▫ > ▫ > ▫)
229.08904	Ergothioneine	0.10	4.03 (▫ > ▫ > ▫)
626.16763	*Analog*: Glutathione (oxidized)	1.58 (▫ > ▫ = ▫)	3.76 (▫ > ▫ > ▫)
663.10912	Nicotinamide adenine dinucleotide	0.4	3.61 (▫ > ▫ = ▫)
203.1173	Acetyl-carnitine	3.50 (▫ = ▫ > ▫)	1.82 (▫ = ▫ > ▫)
567.33206	Lysophosphatidylcholine 22:6	2.99 (▫ > ▫ = ▫)	0.78
250.02948	*Analog*: N-acetylglutamic acid	2.98 (▫ > ▫ = ▫)	2.30 (▫ > ▫ = ▫)
429.14971	*Analog*: Adenosine 5'-monophosphate	0.19	2.96 (▫ > ▫ = ▫)
463.07428	Adenylo succinic acid	2.87 (▫ > ▫ = ▫)	2.90 (▫ > ▫ > ▫)
248.14845	*Analog*: N-acetylglutamic acid	0.48	2.78 (▫ > ▫ > ▫)
268.0806	*Analog*: Adenosine	1.60 (▫ > ▫ = ▫)	2.72 (▫ > ▫ = ▫)
265.0952	Tryptophan	0.19	2.65 (▫ > ▫ = ▫)
347.06212	Adenosine-5' mono-phosphate	1.96 (▫ > ▫ = ▫)	2.55 (▫ > ▫ = ▫)

Environmental sciences

Environmental assessment

Molecular phenotyping

Bioindication

Sentinel species

Microbiome metabarcoding

NMR metabolomics

LC-MS metabolomics

?

Modeling