AdRoit: an accurate and robust method to infer complex transcriptome composition

Tao Yang[1], Nicole Alessandri-Haber[1], Wen Fury[1], Michael Schaner[1], Robert Breese[1], Michael LaCroix-Fralish[2],

Jinrang Kim[1], Christina Adler[1], Lynn E. Macdonald[1], Gurinder S. Atwal[1], Yu Bai[1, *]

Affiliations

1. Regeneron Pharmaceuticals, Inc., Tarrytown NY 10591

2. Cellular Longevity, Inc., San Francisco, CA 94103

*Corresponding author

Abstract

Bulk RNA sequencing technology provides the opportunity to understand biology at the whole

transcriptome level without the prohibitive cost of single cell profiling. Advances in spatial

transcriptomics enable to dissect tissue organization and function by genome-wide gene

expressions. However, the readout of both technologies is the overall gene expression across

potentially many cell types without directly providing the information of cell type constitution.

Although several in-silico approaches have been proposed to deconvolute RNA-Seq data

composed of multiple cell types, many suffer a deterioration of performance in complex

tissues. Here we present AdRoit, an accurate and robust method infer the cell composition

from transcriptome data comprised of multiple cell types. AdRoit uses gene expression profile

obtained from single cell RNA sequencing as a reference. It employs an adaptive learning

approach to correct the sequencing technique difference between the single cell data and the

bulk or spatial transcriptome data, enabling cross-platform readout comparability. Our

systematic benchmarking and applications, which include deconvoluting complex mixtures that

encompass 30 cell types, demonstrate its superior sensitivity and specificity compared to other

existing methods as well as its utilities. In addition, AdRoit is computationally efficient and runs

orders of magnitude faster than many existing methods.

Introduction

RNA sequencing is a powerful tool to understand the biology of normal and disease tissues at

the whole transcriptome level. It helps to elucidate disease mechanism and identify potential

treatment targets[1]. Due to the presence of heterogeneous cell populations, the bulk tissue

transcriptome only characterizes the overall gene expression across multiple cell types. The cell

type identifies and their prevalence remain unknown in the bulk data. However, knowledge of

the cell type composition is often critical to understand the biology. For instance, the

constitution of stromal and immune cells sculpts the tumor microenvironment, which is

essential in cancer progression or response to immune check point inhibition[2–6]. Excessive

expression of cytokines in particular leukocyte types underlines the etiology of many chronic

inflammatory diseases[7–11]. Such information cannot be directly read out from the bulk RNA-Seq.

Recent breakthroughs in spatial transcriptomics methods enable characterizing whole

transcriptome-wise gene expressions at spatially resolved locations in a tissue section[12].

However, it remains challenging to reach a single cell resolution while measuring tens of

thousands of genes transcriptome-wise. Some widely used technologies can achieve a

resolution of 50-100 µm, equivalent to 3–30 cells depending on the tissue type[12,13]. The

transcripts therein may originate from more than one cell type, resulting in another form of

RNA-Seq data of multiple cell types. Unlike the bulk RNA-Seq, the profiling data at each spot

contains substantial dropouts as merely a few cells are sequenced, imposing additional

challenges to demystify the cell type content. We refer to bulk RNA-Seq data and spatial

transcriptome data at the multi-cell resolution as compound RNA-Seq data hereafter.

The rapid development of single-cell RNA-Seq (scRNA-Seq) technologies has allowed for cell-

type specific transcriptome profiling[14]. It provides the information missing from the RNA-Seq

data of tissues comprised of multiple cell types (e.g., bulk RNA-Seq). Nevertheless, the

technologies have low sensitivity and substantial noise due to the high dropout rate and the

cell-to-cell variability. Consequently, scRNA-Seq technologies require a large number of cells

(thousands to tens of thousands) to ensure statistical significance in the results. In addition, the

cells must remain viable during the capture. These requirements render the scRNA-Seq

technologies costly, prohibiting their application in clinical studies that involve many subjects or

cannot allow real time tissue dissociation and cell capture. Furthermore, scRNA-Seq

technologies may not be well suited to characterizing cell-type proportions in solid tissues

because the dissociation and capture steps may not have the same efficiency for different cell

types [15–17].

Because sequencing at the single cell level is not always feasible and also for the purpose of

better interpreting many highly valuable existing bulk RNA-Seq or spatial transcriptomic data

sets, in silico approaches have been developed to infer cell type proportions from compound

RNA-Seq data[18–24]. The most common strategy is to conduct a statistical inference through the

maximum likelihood estimation (MLE)[25] or the maximum a posterior probability estimation

(MAP)[26] on a constrained linear regression framework, wherein the unobserved mixing

proportion of a finite number of cell types are part of the latent variables to be optimized[19,21–

24]. The deconvolution methods are often applied to dissect the immune cell compositions in

blood samples[27–31]. However, their performance in more complex tissues, such as the nervous,

ocular, respiratory or gastrointestinal organs, remains unclear. These tissues often contain

many cell types ($10\text{-}10^2$) and the difference among related cells can be subtle, rendering the

deconvolution a challenging task. For example, a recent study on the mouse nervous system

using scRNA-Seq found more than 200 cell clusters and many are highly similar neuronal

subtypes[32].

Earlier works often utilized the transcriptome profiling of the purified cell populations to

estimate the gene expressions of cell type (e.g. Cibersort)[19]. More recently, acquiring cell type

specific expression from the scRNA-Seq data was shown to be an intriguing alternative[21–24].

Although it provides higher throughput by measuring multiple cell types in one experiment,

profiling at the single cell level has a high noise floor. The accuracy of deconvolution using

scRNA-Seq data as the reference may be affected by the data noise if not treated properly.

Moreover, the platform difference between the compound data and the single cell data cannot

be ignored.

To overcome these challenges, additional information from the data may be considered. A

recent method that weighs genes according to their expression variances across samples

greatly improved the accuracy[22], highlighting the importance of gene variability in inferring cell

type composition. Some other methods and applications have pointed out the importance of

cell type specific genes[24,28,31,33]. In these works, the cell type specific expression was only used

to select the input genes (e.g., markers). Nonetheless, it measures how informative a gene is in

distinguishing cell types and thus can be incorporated as a part of the model. To address the

platform difference between the compound data and the single cell data it is usually assumed

there exists a single scaling factor or a linearly scaled bias for all genes that can be learned and

corrected accordingly[22,23]. This assumption does not hold, as the impact of changing platforms

is different for each gene. Though learning a uniform scaling factor would correct the difference

in most genes, a few genes that remain significantly biased can easily confound the estimation,

especially under a linear model framework. Thus, a gene-wise correction should be considered.

In this work, we present a new deconvolution method, AdRoit. A unified framework that jointly

models the gene-wise technology bias, as well as the cell type specificity and cross-sample

variability of genes. The method estimates the cell type constitution in the compound RNA-Seq

samples using relevant single cell data as a training source. Genes used for deconvolution were

automatically selected from the single cell data based on their information richness. Uniquely,

it uses an adaptively learning approach to estimate gene-wise scaling factors, addressing the

issue that each platform impact genes differently. AdRoit further makes use of regularization to

avoid collinearity among closely related cell subtypes that are common in complex tissues. Over

a set of comprehensive benchmarking data with a varying cell composition complexity, AdRoit

showed superior sensitivity and specificity to other existing methods. Applications to real RNA-

Seq bulk data and spatial transcriptomics data revealed strong and expected biologically

relevant information. We believe AdRoit offers an accurate and robust tool for cell type

deconvolution and will enhance the value of bulk RNA-Seq and the spatial transcriptome

profiling.

Results

Overview of the AdRoit framework

AdRoit estimates the proportions of cell types from transcriptome data of a mixed cell

population including but not limited to bulk RNA-Seq and spatial transcriptome. It directly

models the raw reads without normalization, preserving the difference in total amounts of RNA

transcript in different cell types. The method utilizes, as a reference, the relevant pre-existing

scRNA-Seq data with cell identity annotation. It selects informative genes, estimates the mean

and dispersion of the expression of selected genes per cell type, and constructs a weighted

regularized linear model to infer percent combinations (Fig. 1a). Because sequencing platform

bias may impact genes differently[15,34,35], a uniform scaling factor for all genes does not

sufficiently eliminate such bias. A key innovation of AdRoit is that it uniquely adopts an adaptive

learning approach, where the bias is first estimated for each gene, then adjusted such that

more biased genes are corrected with a larger scaling factor (Fig. 1b).

Fig. 1

AdRoit considers a comprehensive set of other relevant factors including the cross-sample

variability and the cell type specificity of genes, as well as the collinearity of expression profiles

among closely related cell types. The cross-sample variability of a gene is the variation of its

expression in the same cell type across multiple bulk samples. It shall be distinguished from the

expression change in the gene over different cell types. AdRoit decreases the weight of genes

with high cross-sample variability whilst increasing the weight of those with an expression

highly specific to certain cell types. The definition of cross-sample variability and cell type

specificity also accounts for the overdispersed nature of counts data. Lastly, AdRoit adopts a

linear model to ensure the interpretability of the coefficients. At the same time, AdRoit includes

a regularization term to minimize the impact of statistical collinearity. Each of the factors

contributes an indispensable part to AdRoit, leading to an accurate and robust deconvolution

method for inferring complex cell compositions.

To evaluate the performance, we compared AdRoit with NNLS[18,36], MuSiC[22], Bisque[37], and

SPOTlight[38] for bulk data deconvolution, and Stereoscope[23], Cell2location[39], and SPOTlight[38] for

spatial transcriptomics data deconvolution. When evaluating the algorithms, a common

practice is to pool the single cell data to synthesize a "bulk" sample with the known ground

truth of the cell composition. We measured the performance by comparing the estimated cell

proportions with true proportions using four metrics: mean absolute difference (mAD), root

mean square deviation (RMSD), and two correlation statistics (i.e., Pearson and Spearman).

While Pearson reflects linearity, Spearman measures whether the estimated results and the

ground truth are monotonically related, even if their relationship is not linear, which avoids the

artificial high linearity scores driven by outliers when the majority of estimates are small. Thus,

both correlations statistics were included. Good estimations feature low mAD and RMSD along

with high correlation statistics. When estimating cell proportions for a synthetic sample, cells

from this sample were excluded from the input single cell reference (i.e., leave-one-out) to

avoid overfitting. We further applied AdRoit to real bulk RNA-Seq data and validated the results

by available RNA fluorescence *in-situ* hybridization (RNA-FISH) data. The estimates were further

confirmed by the biological knowledge of human pancreatic islets. We also used AdRoit to map

cell types within spatial spots, and the accuracy was verified by *in-situ* hybridization (ISH)

images from Allen mouse brain atlas[40].

AdRoit excels in both simple and complex cell constitutions

We started with a simple human pancreatic islets data set that contains 1492 cells and four

distinct endocrine cell types (Alpha, Beta, Delta, and PP cells)[41] (Supplementary Fig. 1a;

Supplementary Table 1). The synthesized bulk data were constructed by mixing the single cell

data at known proportions. All methods except SPOTlight achieved satisfactory performance

according to the evaluation metrics, AdRoit performed the best as reflected by scatterplots of

estimated proportion vs. true proportion (Supplementary Fig. 1b, Supplementary Table 2). It

had the lowest mAD and RMSD, and the highest correlation values (MuSiC's correlations are

comparable) among the five methods tested (Supplementary Fig. 1c). Given that MuSiC is a

weighted NNLS and consistently showed an improved accuracy here and in the original

publication[22], we excluded NNLS from further evaluation hereafter.

We then tested the methods on a complex tissue - human trabecular meshwork (TM). We

acquired published single cell data that contains 8758 cells and 12 cell types from 8 donors[42].

The data include 3 similar types of endothelial cells, 2 types of Schwann cells, and 2 types of TM

cells (Supplementary Fig. 2; Supplementary Table 3). Cells from each donor were pooled as

individual synthetic bulk samples. The cell type proportions vary from <1% to 43%. These

proportions were the ground truth cell composition and were compared head-to-head with the

estimated proportions inferred by four methods. For each synthetic bulk sample, estimations

were performed using a reference built from cells of remaining donors (i.e., leaving-one-out). In

each of the 8 samples, the estimates made by AdRoit best approximated the true proportions.

In particular, AdRoit had the lowest mAD (0.016) and RMSD (0.025), and the highest correlation

values (Pearson = 0.97; Spearman = 0.94) among the methods (Fig. 2a). We further assessed the

deviation of the estimates from the true proportions for each cell type. AdRoit consistently had

the lowest deviations from the true proportions for all cell types, as well as the lowest variation

among 8 samples (Fig. 2b, blue dots), indicating robustness over various cell types and samples

(Supplementary Fig. 3, Supplementary Table 4).

Fig. 2

AdRoit has better sensitivity and specificity

To assess the level of granularity that AdRoit can achieve when deconvoluting similar cell types,

we used well-studied, closely related immune cells from myeloid lineage and lymphoid lineage.

We obtained B cells, naïve CD4$^+$ T cells, memory CD4$^+$ T cells, CD8$^+$ T cells, natural killer (NK)

cells, dendritic cells (DC), CD14$^+$ monocytes, FCGR3A$^+$ monocytes, and platelets from a public

release of the human peripheral blood mononuclear cell (PBMC) single cells by 10x Genomics

(see also Data Availability). We simulated bulk samples that each contained a mixture of

myeloid lineage or lymphoid lineage cell types. Such a sample was created by randomly

selecting cells from a given set of cell types and mixing them according to their predefined

percentages. We repeated this procedure 100 times for a series of mixing proportions among

myeloid or T lymphoid cells. The bulk mixtures of CD14$^+$ monocytes, FCGR3A$^+$ monocytes and

dendritic cells (DC) followed three schemes of proportions: 0.33:0.33:0.33 (mix0), 0.1:0.45:0.45

(mix1) and 0.1:0.3:0.6 (mix2). The same ratios were applied to the mixtures of naïve CD4$^+$ T,

memory CD4$^+$ T and CD8$^+$ cells. All the nine cell types in the PBMC data were used to build the

reference for the deconvolution. AdRoit accurately estimated the known percentages in all

cases (Fig. 3a, Supplementary Table 5). In addition, there were no noticeable false positive

predictions of any excluded cell types. These results highlight that AdRoit can capture the fine

difference among similar cell types and quantitatively distinguish them.

Fig. 3

We next systematically benchmarked the sensitivity and specificity of each of the algorithms. In

the context of the cell type deconvolution, a false negative occurs when the proportion of an

existing cell type is predicted to be zero (or below a given threshold). Conversely, a non-zero

prediction (or above a given threshold) of an absent cell type results in a false positive. False

negatives and false positives measure the sensitivity and specificity of a deconvolution

algorithm, respectively. Both quantities are crucial to establish the utility of the algorithm.

Particularly, in real-world applications, it is often difficult to know *a prior* what cell types exist in

a bulk sample, users may inform the algorithm to consider more possible cell types than what

actually exist in the sample. False positive predictions in this situation would make the

algorithm unusable.

We designed a simulation to test sensitivity and specificity. we selected 6 out of the 12 human

trabecular meshwork cell types, i.e., Schwann-cell like cell, TM1, smooth muscle cell,

melanocyte, macrophage, and pericyte, from each donor sample and pooled them within that

sample to synthesize 8 new bulk samples. The unselected 6 cell types were considered absent

in the bulk samples. Some cell types selected highly resembled those in absence, challenging

the programs to pinpoint the right cell type present in the bulk among similar candidates. We

provided the full list of 12 single cell types as the reference to the programs to estimate the cell

type proportions. Consistently across 8 samples, AdRoit had the most accurate estimates for

the 6 present cell types, and 0 or close to 0 estimated values for the non-existing cell types in

the simulated bulk data. MuSiC, Bisque and SPOTlight were notably less accurate on the 6

selected cell types, as well as they had many non-negligible values for the 6 cell types excluded

in the 8 synthetic samples (Fig. 3b, Supplementary Table 6). For example, smooth muscle cells

accounted for ~14% in donor 4 but were largely missed (~0.03%) by MuSiC. We noted that TM2

had false non-zero estimates from all methods though not included. This is because TM2 is

easily mistaken as TM1 due to their high similarity[42]. Nonetheless, AdRoit's estimates of TM2

were consistently low across samples (<1% for 44 out of 48 estimates), while the other

methods had significantly larger estimates of TM2 that occasionally even exceeded the TM1

estimates. For a systematic comparison, we constructed the receiver operating characteristic

(ROC) curve by varying the threshold of detection (i.e. a cutoff below which the cell type was

deemed undetected) (Fig. 3c). AdRoit had a significantly higher area under the curve (AUC) than

the other methods (AdRoit: 0.95, Bisque: 0.79, MuSiC: 0.74, SPOTlight: 0.37), implying both

better sensitivity and specificity.

As complex tissues often contain several tens of different cell types, we continued the

evaluation of the sensitivity and specificity in more complicated cell mixtures. We utilized the

published mouse brain single cell atlas by Zeisel et al that contains a comprehensive set of

neuronal and supporting cells[32]. Without loss of generality, we consolidated the original

annotation into 46 major cell types (Methods, Supplementary Table 7). To synthesize a bulk

mixture, we randomly selected 30 cell types and pooled all their associated cells. This

procedure was repeated independently 100 times for thorough coverage of different cell type

combinations. The deconvolution was performed using all 46 cell types as the reference. As

illustrated in Fig. 3d, AdRoit was able to estimate the proportions closely consistent with the

ground truth while the other three tools made less satisfactory predictions (see also

Supplementary Table 8). The ROC curve further indicates that AdRoit has the best sensitivity

and specificity given its highest AUC (Fig. 3e; AdRoit: 0.99, Bisque: 0.77, MuSiC: 0.80, SPOTlight:

0.67). These observations convince us that AdRoit offers a leading performance on highly

complex cell constitutions.

AdRoit outperforms in deconvoluting closely related cell subtypes

To further evaluate AdRoit when multiple highly homologous subtypes of cells are present in a

complex tissue, we performed scRNA-Seq experiment on mouse lumbar dorsal root ganglion

(DRG) from five mice. Following the standard analysis pipeline (Methods), we obtained 3352

single cells after quality control procedures. After clustering and annotation, we discovered 14

cell types including multiple subtypes of neuronal cells (Fig. 4a, Supplementary Table 9). The

heatmap of the top marker genes showed distinct patterns of the major cell types as well as

similar patterns of the subtypes (Supplementary Fig. 4a), with the cell type proportions varying

from 0.5% to 33.71% (Supplementary Fig. 4b). These 14 cell types include 3 subtypes of

neurofilament containing neurons (i.e., NF_Calb1, NF_Pvalb, NF_Ntrk2.Necab2), 3 subtypes of

non-peptidergic neurons (i.e., NP_Nts, NP_Mrgpra3, NP_Mrgprd), and 5 subtypes of

peptidergic neurons (i.e., PEP1_Dcn, PEP1_S100a11.Tagln2, PEP1_Slc7a3.Sstr2,

PEP2_Htr3a.Sema5a, PEP3_Trpm8). Also discovered were tyrosine hydroxylase containing

neurons (Th), satellite glia, and endothelial cells. Such complex compositions formed a

challenging testing ground for evaluating the ability to distinguish closely related cell types. We

again performed the leave-one-out deconvolution on five synthesized bulk samples.

Fig. 4

AdRoit had highly accurate estimations on all cell subtypes across samples (Fig. 4b). Worth

noting, for the rare cell types that account for less than 5%, AdRoit still had a good estimation

that is close to the true proportions and never missed a single cell type, showing that AdRoit is

very robust on rare cell types. For example, 0.51% endothelial cells were predicted to be 0.35%,

and 1.05% NF2_Ntrk2.Necab2 cells were predicted to be 0.85% (Supplementary Fig. 5,

Supplementary Table 10). Conversely, Bisque, MuSiC, and SPOTlight were notably less accurate,

especially for the cell types less than 5%, and all three missed multiple cell types including some

large clusters accounting for ~10% of the cells (PEP1_Slc7a3.Sstr2 cells of Sample5). We

continued to examine how much the variability of the estimates was in each sample. We

computed the same 4 metrics used previously to evaluate the performance on each of the 5

synthetic samples and compared them head-to-head among the algorithms. This fine

comparison showed AdRoit significantly outperformed on nearly every sample (Fig. 4c).

Moreover, the performance metrics of AdRoit were highly consistent across samples with the

lowest variability among all four methods.

AdRoit excels on spatial transcriptomics data

Given the promising performance on complex tissues, we moved forward to test AdRoit's

applicability to spatial transcriptome data. Spatial transcriptomics data differs from bulk RNA-

Seq data in that each spot only contains transcripts from a handful of cells (3-30)[12]. Some of the

spots contain multiple cells of the same type, while others may have heterogeneous cell types

at varying mixing percentages (e.g., spatial spots at the boundary of different cell types). As the

spot is a pool of only a few cells, the variations across spatial spots are expected to be greater

than the changes in bulk samples, imposing an additional challenge for deconvolution. We

simulated a large number of spatial spots (3200 in total) by using sampled cells from the DRG

single cell data above (Methods), then compared AdRoit with Stereoscope, Cell2location, and

SPOTlight over a range of simulation scenarios.

We first tested whether the methods could correctly infer a single cell type when the spots

contain cells from that same type. For each of the 14 cell types from DRG, we sampled 10 cells

and pooled them to form a spatial spot. We repeated the simulation 100 times for robust

testing, then used the full set of 14 cell types as reference to deconvolute the 1400 simulated

spots. All methods except SPOTlight were able to identify the correct cell types with high

accuracy on the target cell types (i.e., percentages close to 1) and comparably low estimated

values (i.e., percentages close to 0) for other cell types excluded from the

simulation(Supplementary Fig. 6, Supplementary Table 11).

We then proceeded to a more difficult scenario where we sampled cells from the 5 PEP neuron

subtypes and mixed them. We created three simulation schemes for a comprehensive

evaluation: 1) 5 PEP subtypes had the same percent of 0.2; 2) PEP1_Dcn was 0.1 and the other

4 were 0.225; 3) PEP1_S100a11.Tagln2 and PEPE1_Dcn were 0.1, PEP2_Htr3a.Sema5a and

PEP1_Slc7a3.Sstr2 were 0.2, and PEP3_Trpm8 was 0.4. Again, each simulation scheme was

repeated 100 times. Under each scheme, the estimates by AdRoit consistently centered around

the true proportions while the predictions of irrelevant cell types remained close to 0 (Fig. 5a,

Supplementary Table 11). In comparison, Cell2location systematically overestimated the

PEP1_Slc7a3.Sstr2 by about 2.5%. Stereoscope more notably overestimated

PEP2_Htr3a.Sema5a and underestimated the PEP3_Trpm8. SPLOTlight had high noise for some

cell types (e.g., endothelial and satellite glia) and generally deviated from the truth for multiple

cell types under all simulation schemes.

Fig. 5

We expanded the simulated spatial spots to the mixture of 3 NP cell subtypes and the mixture

of 3 NF subtypes. In addition, we blended either NP_Mrgpra3 cells, NF_Calb1 cells, or

PEP3_Trpm8 cells with three other cell types (i.e., Th, satellite glia and endothelial). For all

these simulated spatial spots, AdRoit's estimates consistently centered around the true

proportions, whereas the other methods deviated by different degrees in most of the simulated

schemes (Supplementary Fig. 7, Supplementary Table 11). We speculate one important reason

for the underperformance of some methods such as Stereoscope is that they normalize the

total UMI counts to the same value for all cells. However, in real world, different cell types

unlikely have the same amount of RNA transcripts (e.g. immune cells have about 10 fold fewer

total UMIs than the neurons). Our simulation pooled the cells by adding up the raw UMI counts

per gene, which we believe best mimics the real data. Normalizing the data by total UMI not

only eliminates this factual difference between cell types but may distort the data from the

assumed negative binomial distribution. AdRoit avoids this problem by model the raw UMI

counts.

Next, we asked how sensitive the methods are in detecting rare cell populations. We simulated

mixtures of 3 PEP subtypes (i.e., PEP1_Slc7a3.Sstr2, PEP2_Htr3a.Sema5a, and PEP3_Trpm8)

wherein the percentage of PEP3_Trpm8 cells varied from 0.01 to 0.1 by 0.01, and the other two

cell types sharing the remaining percentage equally (Methods). At each given mixing ratio, the

simulation was repeated 100 times. We then checked how accurately the concentration of

PEP3_Trpm8 cells was estimated. The medians of AdRoit's estimates were closest to the true

proportions (Fig. 5b, red dashed lines) among all 4 methods, followed by Cell2location.

Stereoscope's estimates were systematically lower than the true values and failed to detect the

PEP3_Trpm8 cells when the simulated proportion was below 0.06. On the other hand,

SPOTlight constantly overestimated the presence of PEP3_Trpm8 cells. The median of its

predictions remained around 9% even when the true population was at 1%. Such observations

are consistent with the frequent false positives made by SPOTlight when deconvoluting the

simulated bulk samples (Fig. 3b). This comparison implied AdRoit is more advantageous in

detecting low percent cells. For a complete evaluation, we replicated the comparison using 5

additional sets of cell mixtures: NF_Calb1 with NF_Pvalb and NF2_Ntrk2.Necab2 (NP subtypes);

NP_Mrgpra3 with NP_Mrgprd and NP_Nts (NP subtypes); NF_Calb1 with Th, satellite glia and

endothelial (NF_Calb1 + others); NP_Mrgpra3 with Th, satellite glia and endothelial

(NP_Mrgpra3 + others); and PEP_Trpm8 with Th, satellite glia and endothelial (PEP_Trpm8 +

others). In each set, the first cell type mentioned was considered rare with its percent varied

from 0.01 to 0.1. The rest cell types shared the remaining proportion evenly. At each given

percent of the rare cell type, we computed how many times out of 100 the rare cell component

was detected (estimation > 0.005). AdRoit had systematically high detection rates especially

when the percent is below 3%. (Fig. 5c, Supplementary Table 12). Note that the apparent high

detection rates in SPOTlight were merely the result of its high false positive estimates (Fig. 3b

and Fig. 5b). The detection sensitivity of Cell2location was comparable to that of AdRoit.

Notably, given a rare cell population of 5%, both AdRoit and Cell2location achieved a detection

rate >90% under all simulation schemes, making them powerful tools to uncover rare cells.

Application to real bulk RNA-Seq data of human pancreatic islets

Though using synthetic bulk data based on the mixing of single cells is a useful benchmarking

strategy, the bulk and single cell RNA-Seq often use distinct RNA library preparations and

sequencing protocols. The capability of a method to deconvolute real bulk samples shall be

addressed to ensure it is useful in real-world applications. We acquired 70 real human

pancreatic islets bulk samples from published studies[41,43,44] (Supplementary Table 13) and used

single cell data of the same tissue[41] as the reference to infer the percentages of 4 endocrine cell

types (i.e., Alpha, Beta, Delta, PP). The 70 bulk samples were collected from 39 distinct donors,

including 26 healthy donors, and 13 donors with type 2 diabetes (T2D). Each donor contributed

1 to 5 bulk RNA sample repeats.

Replicates from the same donor are expected to have similar compositions and thus were used

to assess the reproducibility of the estimates from AdRoit. For all cell types, AdRoit had

consistent estimates for the same donors (Fig. 6a, Supplementary Table 14). The average

standard deviations did not exceed 1% for all 4 cell types (i.e., Alpha: 0.010; Beta: 0.008; Delta:

0.004; PP: 0.002). To seek an independent validation, we obtained cell sorting results by RNA-

FISH for 4 of the 39 donors[41] (Supplementary Table 15). The estimated cell proportions in the 4

samples agreed well with the percentages measured by RNA-FISH (Fig. 6b). The consistency

held for both major cells (Alpha and Beta) and the minor cells (Delta and PP). The observed

reproducibility and the success of the independent validation showed AdRoit is reliable in

deconvoluting real bulk RNA-Seq data.

Fig. 6

We then asked if AdRoit can detect known biological differences between healthy and T2D

donors. Loss of functional insulin-producing Beta cells is a prominent characteristic of T2D[45–47],

typically reflected by an elevated level of hemoglobin A1c (HbA1c)[48,49]. Among the healthy

donors, most Beta cell proportions estimated by AdRoit ranged from 50% to 75% (Fig. 6c),

agreeing with the known percent range of Beta cells in human islets tissue[50,51]. A significant

decrease in the estimated Beta cell proportions was seen in T2D patients (p value = 4.1e-6 by

two-sided t-test). Further, a linear regression of estimated Beta cell proportions on HbA1c

levels showed a statistically significant negative association (p value = 1.8e-6), showing that

AdRoit adequately reflected the cell composition difference between healthy donors and T2D

patients.

Application to mouse brain spatial transcriptomics

We lastly demonstrated an application to the real spatial transcriptomics data. Given that the

molecular architecture of brain tissues has been well studied, for this evaluation, we chose a set

of mouse brain spatial transcriptomics data generated by 10x genomics, containing 2703 spatial

spots. The consolidated version of the aforementioned mouse brain single cell atlas [32] was used

as the reference for deconvolution (Methods).

The cell contents inferred by AdRoit per spot appear to accurately match the expected cell

types at that location (Supplementary Fig. 8, Supplementary Table 16). For example, the three

subtypes of cortex excitatory neurons each occupied a sub-area in the cerebral cortex region.

As another example, the shape of the hippocampal region was delineated by the estimated

percentages of dentate gyrus granule/excitatory neurons. As an independent validation, we

checked the consistency between the estimated cell types with the *in-situ* hybridization (ISH)

images from Allen mouse brain atlas[52]. We chose 4 genes highly expressed in 4 brain regions

respectively, i.e., Spink8 for the hippocampal field CA1; C1ql2 for the dentate gyrus; Clic6 for

the choroid plexus; and Synpo2 for the thalamus[32]. The spots enriched with the 4 indicative cell

types, hippocampal CA1 excitatory neuron type 2, dentate gyrus granule neuron type 2, choroid

plexus cell and thalamus excitatory neuron type 1, as mapped by AdRoit, precisely co-localized

with the regions that showed a strong signal of the 4 marker genes on the ISH images

respectively (Fig. 6d). This agreement confirmed that the spatial mapping of cell types by AdRoit

is reliable.

Computational efficiency

Besides the accuracy and robustness, another major advantage of AdRoit is its magnitude

higher computational efficiency. AdRoit uses a two-step procedure to do the inference. The first

step prepares the reference using the single cell data where the per-gene means and

dispersions are estimated, and cell type specificity is subsequently computed. The built

reference can be saved and reused. We tested the running time on the reference construction

using the mouse brain single cell dataset discussed earlier that contains ~15,000 cells. It took

about 4.5 minutes on a CPU that has 24 cores (23 used for parallel computing). The second step

inputs the built reference and the compound RNA-Seq data and does the estimation.

Deconvoluting ~2700 RNA-Seq samples of mixed cell populations took around 5 minutes.

Therefore, AdRoit in total took less than 10 minutes and ~3Gb memory usage on a regular CPU.

As a comparison, MuSiC and Bisque took about 1 hour and 30 minutes on the same data using

the same CPU setup. SPOTlight took about 2 hours with the default parameters (cl_n = 100, hvg

= 3000, ntop = 200, transf = "uv", method = "nsNMF", min_cont = 0). Stereoscope ran about 24

hours continuously with the published parameter setting (-scb 256 -sce 75000 -topn_genes

5000 -ste 75000 -lr 0.01 -stb 100 -scb 100) on a powerful V100 GPU with 80 cores and 16G

memory. The efficiency of Cell2location was comparable to Stereoscope, taking about 18 hours

on the V100 GPU (posterior n_sample = 1000, n_iteration = 10000, selection_specificity =

0.45). These methods can be prohibitive when seeking a quick turnaround.

Discussion

In this work we have demonstrated that AdRoit is capable of deconvoluting the cell

compositions from the compound RNA-Seq data with a leading accuracy, measured by the

consistency between the true and predicted cell proportions. Its advantage over the existing

state-of-the-art methods was verified over a wide range of use cases. In particular, AdRoit

excelled in complex tissues composed of more than ten different cell types with a wide range of

cell proportions (e.g., trabecular meshwork, mouse brain and dorsal root ganglion). In these

cases, AdRoit performed significantly better than the comparators on deconvoluting bulk RNA-

Seq data. AdRoit is also more accurate and sensitive than tools specifically designed for

deconvoluting spatial transcriptomics spots, especially in detecting low percent rare cells.

Previous benchmarking often assumed the types of cells in the synthetic bulk data are not more

or less than the cell types collected in the reference, and thus the only unknown was the

proportion of each cell type. This assumption may not hold. Missing existing cell types or false

predictions of non-existing ones can hinder the utility of an algorithm. Thus, besides the overall

accuracy, we also examined the sensitivity and specificity of the algorithms. We observed a

superior sensitivity and specificity in AdRoit, an important leverage for its usage in practice.

The reference single cell data used by AdRoit came from different platforms, such as the 10x

Genomics Chromium Instrument (the mouse dorsal root ganglion), and the Fluidigm C1 system

(the human pancreatic islets data). AdRoit consistently exhibited excellent performance across

all benchmarking datasets independent of their single cell sequencing technology platforms.

More importantly, this statement holds not only for deconvoluting the synthesized bulk data,

but also for the real bulk RNA-Seq data. The latter typically does not apply the unique molecular

barcoding and requires a significantly different cDNA amplification procedure from what is used

in the single cell RNA-Seq (Methods). Moreover, the sequencing depth, read mapping and gene

expression quantification are dissimilar as well. The fact that AdRoit accurately dissected the

cell compositions in the real bulk samples based on the single cell reference data further

supports its cross-platform applicability.

We attribute the power of AdRoit to its comprehensive modeling of relevant factors. Firstly, we

think a common rescaling factor is insufficient to correct the platform difference between

single cells and the compound data. Rather, the impact of platform difference on genes is quite

different and not linearly scaled. Correcting such differences entails rescaling factors specifically

tailored to each gene. AdRoit uses an adaptive learning approach to estimate such gene-wise

correcting factors and does the correction in a unified model. In addition, the contribution of a

gene in a cell type to the loss function is jointly weighted by its specificity and variability in a cell

type, where specificity and variability are defined in a way accounting for the overdispersion

property of counts data. Our observations over the multiple benchmarking data sets also show

that the coexistence of similar cell types may have induced a collinearity condition that

negatively impacted the regression-based methods developed by others. Being able to alleviate

this problem gives AdRoit an edge. All these factors help AdRoit to distinguish similar cell

clusters while remaining sensitive enough to separate rare cell types.

Technically, the input profiles of individual cell types to AdRoit need not come from the single

cell RNA-Seq. Bulk RNA-Seq profiles of individual isolated cell types can be used as well.

Nevertheless, using single cell RNA-Seq data as the reference has a few key advantages. It is a

high throughput approach wherein multiple cell types can be interrogated simultaneously. Prior

knowledge of the cell types in presence as well as their specific gene markers are not required,

which allows novel cell types to be identified. Although detection of lowly expressed genes has

been a challenge for the single cell RNA-Seq, significant enhancements have been

demonstrated. For example, the number of detectable genes currently can reach an order of

10,000 per cell and keeps improving[53]. As AdRoit focuses on the informative genes whose

expressions are generally high, the detection limit of the single cell RNA-Seq does not impose a

significant drawback. Indeed, given the single cell reference profiles, AdRoit successfully

deconvoluted the real bulk RNA-Seq data and spatial transcriptomics data. The results suggest

that, besides enriching our understanding of the bulk transcriptome data, AdRoit can leverage

the usage of the vast amount and continuously growing single cell data as well.

AdRoit is a reference-based deconvolution algorithm. A comprehensive collection of the

possible cell components is important. However, completeness may not always be guaranteed.

Even with the single cell acquisition that is independent of prior knowledge, rare and/or fragile

cell types may not survive through the capture procedure and hence are excluded. It is also

difficult to generate a solid reference profile for cells that are versatile from sample to sample

(e.g., tumor cells). Currently AdRoit deals implicitly with the components unknown to the

reference. If an unknown cell type reassembles one of the referenced ones, it may be

considered as part of the known cell type and their joint population is predicted. Such an

outcome is acceptable as treating two similar cell types as one is still biologically meaningful

although the resolution of the system may be compromised. If the unknown component is

dissimilar to all the known ones, it will be ignored by AdRoit because its representative markers

are unlikely among the top weighted genes associated with the known components. At the

same time, the distinct component is expected to have a unique gene expression pattern and

thus unlikely to interfere significantly with the gene expressions from the known cell types.

Therefore, AdRoit essentially deconvolutes the relative populations among the known cell

components. For example, AdRoit was able to correctly uncover the populations of 4 endocrine

cell types from the human islet bulk data despite the absence of many other cell types such as

macrophages, Schwann cells and endothelial cells in the input single cell reference[20]. Although

under such a circumstance, the absolute percentages of the cells remain obscure, we expect

their relative proportions can be studied and valuable. A future improvement is to explicitly

model the unknown cell types and estimate their percentages upon the signals in the

compound data that cannot be explained by the contribution from the known components.

AdRoit may also be coupled with other programs[54] to output the deconvoluted cell type specific

transcriptome profiles. The inferred cell type proportions by AdRoit can serve as the input to

the downstream program and benefit the outcome with the reliability and accuracy

demonstrated in this work.

Methods

Gene selection

AdRoit selects genes that provide information about cell type identity, excluding non-

informative genes that potentially introduce noise. There are two ways for selecting such

genes: 1) union of the genes whose expression is enriched in one or more cell types in the

single cell UMI count matrix. These genes are referred to as marker genes; 2) union of the

genes that vary the most across all the cells in the single cell UMI count matrix, referred to as

the highly variable genes. For marker genes, we recommend selecting the top ~200 genes,

ranked by fold change, from each cell type for resolving complex compound transcriptome

data. Considering some genes may be enriched in more than one cell type, we further require

selected markers presenting in no more than 5 cell types to ensure specificity. We also suggest

selecting a minimum of 1000 total unique marker genes for an accurate estimation. If not

satisfied, one may consider expanding the number of top genes and/or loosen the statistical

significance cutoff.

AdRoit also offers the option to use highly variable genes. To avoid the selected highly variable

genes being dominated by large cell clusters, AdRoit first balances the cell types in the single

cell UMI count matrix by finding the median size (i.e., the number of cells) of all clusters, then

samples cells from each cluster to make its size equal to the median. Next, AdRoit computes the

variance of each gene across all the cells in the balanced single cell UMI matrix. The variance-

stabilizing transformation (VST)[55] is applied to normalize the data prior to the variance

calculation. Genes with the top 2000 largest variances are then selected.

In both ways, mitochondria genes are excluded as their expression does not have information

of cell identity. The results shown in the current paper were based on the marker genes derived

as described above. We also demonstrated that using the balanced highly variable genes

yielded comparably accurate estimations (Supplementary Fig. 9).

Estimate gene-wise mean and variance per cell type

Modeling single cell RNA-Seq data is challenging due to cell heterogeneity, technical sensitivity,

and noise. Some genes may not be detected by chance, while others may be found to be highly

dispersed. These factors can lead to excessive variability even within the same cell type. AdRoit

combats high noise and computational complexity by building models with estimated mean and

dispersion per gene per cell type. This strategy reduces the data complexity while preserves the

cell type specific information.

Although typical analyses of RNA-Seq data start with normalization, AdRoit does not do

normalization before the mean estimation. Performing a normalization across all cell types

forces every cell type to have the same amount of RNA transcripts, measured by the total

unique molecular identifier (UMI) counts per cell. However, different cell types can have

dramatically different amounts of transcripts. For example, the amount of RNA transcripts in

neuronal cells is about 5 times higher than that in glial cells. Thus, normalization can falsely

alter the relative abundance of cell types, misleading the estimation of cell type percentages.

To avoid this problem, AdRoit models the means using the raw UMI counts.

Studies have shown that UMI counts follow a negative binomial distribution[56,57]. We therefore

fit negative binomial distributions to genes of each cell type and build the model based on the

estimated means and dispersions from the selected genes. More specifically, let X_{ik} be the set

of single cell UMI counts of gene $i \in 1,..,I$ for all cells in cell type $k \in 1,...,K$. I is the number of

selected genes, and K denotes the number of cell types in the single cell count matrix. The

distribution of X_{ik} follows a negative binomial distribution,

582                                $X_{ik} \sim NB(\lambda_{ik}, p_{ik}),$ (1)

583   where λ_{ik} is the dispersion parameter of the gene i in cell type k, and p_{ik} is the success

584   probability, i.e., the probability of one observed UMI belonging to gene i in cell type k. The two

585   parameters are estimated by MLE. The likelihood function is

586                        $LH(\lambda_{ik}, p_{ik} | X_{ik}) = \prod_{i=1}^{n_k} f(X_{ik} | \lambda_{ik}, p_{ik}),$ (2)

587   where n_k is the number of cells in cell type k, and f is the probability mass function of the

588   negative binomial distribution. The MLE estimates are then given by

589                        $(\widehat{\lambda_{ik}}, \widehat{p_{ik}}) = \underset{\lambda_{ik}, p_{ik}}{arg\max} LH(\lambda_{ik}, p_{ik} | X_{ik}).$ (3)

590   Once the success probability and the dispersion are estimated, the mean and the variance of

591   the negative binomial distribution can be computed numerically,

592                                $\mu_{ik} = \frac{\widehat{\lambda_{ik}} \cdot \widehat{p_{ik}}}{1 - \widehat{p_{ik}}},$ (4)

593                                $\sigma_{ik}^2 = \frac{\widehat{\lambda_{ik}} \cdot \widehat{p_{ik}}}{(1 - \widehat{p_{ik}})^2}.$ (5)

594   Model fitting using MLE has been readily coded in many R packages. We choose the 'fitdist'

595   function from the 'fitdistrplus' package[58] for its fast computation speed and flexibility in

596   selecting distributions. The mean and variance estimations are done for each selected gene in

597   each cell type, resulting in a $I \times K$ matrix each.

Cell type specificity of genes

Genes with a cell-type specific expression pattern better distinguish cell types, thus are more

important for resolving cell type composition. In line with this property, AdRoit weighs more

the genes with a higher cell type specificity. Highly specific genes usually have consistently high

expression and thus relatively low variance among cells within a cell type. To compute the cell

type specificity of a gene, we first identify the cell type in which the gene has the highest

expression (i.e., most expressed cell type), then defines the specificity of this gene as the mean-

to-variance ratio within that cell type. A high ratio renders a high weight to the gene in the

model. Here the mean and the variance refer to the estimated values from the negative

binomial model (μ_{ik} and σ_{ik}^2 in eq. 4 and 5). Let k' be the index of the cell type that has the

highest mean expression of gene i,

$$k' = arg\max_{k}\{\mu_{ik} \mid k \ \epsilon \ 1 \dots K\}, \qquad (6)$$

then the weight of the cell type specificity for gene i, denoted w_i^S, is given by,

$$w_i^S = \frac{\mu_{ik'}}{\sigma_{ik'}^2}, \qquad (7)$$

and it is computed for each gene in the set of selected genes.

Cross-sample gene variability

The variability of a gene contrasts with how stable a gene is across samples. The idea of

weighting genes based on their variability across samples is first explored by Wang et al[22],

wherein the variability was defined as the cross-sample variance. By weighting down the high

variability genes, the authors achieved a great advantage over the traditional unweighted

method. Genes with a low cross-sample variability better represent the sample population,

hence are more trustworthy for learning the cell composition. AdRoit incorporates the same

notion to weigh the importance of genes, however, it defines the variability more

comprehensively. AdRoit acknowledges the dependency between the gene expression level

and its variance and thus computes a variance-to-mean ratio (VMR) as the cross-sample

variability. Here the mean and the variance are computed across samples. The distribution of

the VMR values is less skewed than that of the variance alone. The VMR also circumvents

underweighting genes with a low expression, or overweighting genes highly dispersed. Let Y_{ij}

denote the counts for gene i in sample $j \in 1,...,J$, then

$$Y_{ij} \sim NB(\lambda_{ij}, p_{ij}), \qquad (8)$$

where λ_{ij} is the dispersion parameter of the gene i in sample j, and p_{ij} is the success

probability. Again, we use MLE to estimate $\widehat{\lambda_{ij}}$ and $\widehat{p_{ij}}$, following which the cross-sample mean

and variance can be numerically computed:

$$\mu_i^S = \frac{\widehat{\lambda_{ij}} \cdot \widehat{p_{ij}}}{1 - \widehat{p_{ij}}}, \qquad (9)$$

$$(\sigma_i^2)^S = \frac{\widehat{\lambda_{ij}} \cdot \widehat{p_{ij}}}{\left(1 - \widehat{p_{ij}}\right)^2}, \qquad (10)$$

and the cross-sample variability VMR for gene i is then defined as

$$VMR_i = \frac{(\sigma_i^2)^S}{\mu_i^S} = \frac{1}{w_i^C}, \qquad (11)$$

where w_i^C is later used in the model. The cross-sample variability weight is computed for each

gene in the set of selected genes.

Typically, the RNA-Seq data of mixed cells to be deconvoluted include multiple samples to

control for biological variability. In this case, the VMR is computed as described above using the

cross-sample mean and variance of the gene. In case the RNA-Seq data to be deconvoluted lack

multiple samples whereas the single cell reference contains replicates, AdRoit pools all cells per

replicate to synthesize multiple compound samples and subsequently estimate the VMRs with

them. If neither input has multiple repeats, AdRoit takes a bootstrapping approach to

subsample and pool cells from each cell type in the single cell reference several times to

generate multiple compound samples.

Gene-wise scaling factor to correct the platform bias

When comparing the RNA-Seq data of cell mixtures to the single cell data, a rescaling factor is

often used to account for the library size and platform difference. The existing methods adopt a

single rescaling factor for each sample, i.e., all genes in a sample are multiplied by the same

factor[22,23]. This operation is based on a strong assumption that the platform difference impacts

every gene equally and linearly scalable among different cell types, which hardly holds. In

addition, because linear models are sensitive to outliers, the estimation of cell proportions can

be steered away from the truth by genes that are significantly affected by the platform bias.

Therefore, applying a uniform scaling factor to all genes is inappropriate.

To overcome this problem, AdRoit estimates a gene-wise scaling factor per sample via an

adaptive learning strategy and rescales each gene with its respective scaling factor. To proceed,

we first input the mean gene expression from the mixture samples (μ_i^S in eq. 9) and the

estimated means of each cell type from the single cell data (μ_{ik} in eq. 4), then apply a non-

negative least square regression (NNLS) to get a coarse estimation of the proportions of each

cell type, τ_k. For each gene i, a predicted mean expression ($\sum_k^K \widehat{\tau_k} \mu_{ik}$ in eq. 13) in the sample of

mixture cells is computed as the weighted sum of the means of this gene in each cell type

wherein the weights are τ_k. The regression equation is given by,

666
$$\mu_i^S = A \cdot (\textstyle\sum_k^K \tau_k \, \mu_{ik} + \varepsilon), \qquad 0 < \tau_k, \; \textstyle\sum_k^K \tau_k = 1 \quad (12)$$

667   where A is a constant to ensure the sum of τ_k's is 1 and ε is the error term. We use the 'nnls'

668   function in the 'nnls' package[59] to estimate τ_k's. Next, we calculate the ratio between the mean

669   expression from the mixture samples and the predicted counterpart, and define the gene-wise

670   rescaling factor as the logarithm of the ratio plus 1,

671
$$r_i = \log \left(\frac{\mu_i^S}{\sum_k^K \widehat{\tau_k}\mu_{ik}} + 1 \right). \qquad\qquad (13)$$

Given the dispersed nature of count data, the logarithm of the ratio is a more appropriate

statistic as it results in relatively stable scaling factors. The addition of 1 avoids taking logarithm

on zeros. By multiplying the gene-wise rescaling factor, the genes affected more strongly by the

platform bias (i.e., "outliers") receive a larger correction (Fig. 1b).

Weighted and regularized loss function for training

AdRoit employs a non-negative least square regression model to infer the cell compositions in a

RNA-Seq sample of cell mixtures. The model (eq. 14) approximates the gene expression

observed in the mixture sample by the weighted (the weights are the mixing percentages) sum

of the cell type-specific gene expressions from the single cell reference, after correcting the

platform difference.

683
$$y_{ij} \approx r_i \cdot \textstyle\sum_k^K \beta_k \widehat{\mu_{ik}} \qquad (\beta_1, \dots, \beta_K > 0) \qquad (14)$$

where y_{ij} is the expression of gene i in the mixture transcriptome sample j; the coefficient β_k is

the mixing proportion for cell type k; $\widehat{\mu_{ik}}$ is the estimated mean expression of gene i in cell type

k computed according to eq. 4; r_i is the gene-wise scaling factor computed by eq. 13 to correct

the bias due to the technology difference. To infer the β_k coefficients, AdRoit uniquely

incorporates all the factors discussed in the previous sections. This is implemented in a

weighted sum-of-squared loss function L, where the weights consist of two components, w_i^C in

eq. 7 and w_i^S in eq. 11. Thus, a larger weight is given to the genes with a higher cell type

specificity and a lower cross-sample variability. In cases of complex tissues where many similar

cell subtypes exist, strong collinearity among the subtypes can render the model sensitive to

noise and prone to overfitting. AdRoit handles this problem by including an L2 norm of the

mixing proportions as the regularization component in the loss function. For a compound

transcriptome sample j, the loss function is given by,

$$L_j(\beta_1, \ldots, \beta_K | y_{ij}, w_i^C, w_i^S, r_i, \widehat{\mu_{\iota k}}) = \sum_i^I w_i^C \cdot w_i^S \cdot (y_{ij} - r_i \cdot \sum_k^K \beta_k \widehat{\mu_{\iota k}})^2 + \sum_k^K \beta_k^2. \quad (15)$$

where i is the index of a gene among the selected genes used for the deconvolution; $y_{ij}, \beta_k, r_i,$

and $\widehat{\mu_{\iota k}}$ are the same as defined in eq. 14; w_i^S and w_i^C are computed as described above in eq. 7

and 11, respectively. The β_k coefficients are estimated by minimizing the loss function subject

to the constraint $\beta_1, \ldots, \beta_K > 0$,

$$\widehat{\beta_1}, \ldots, \widehat{\beta_K} = \operatorname*{argmax}_{\beta_1, \ldots, \beta_K > 0} L_j. \quad (16)$$

The minimization is done by a gradient projection method proposed by Byrd et al[60]. We derive

the gradient function by taking the partial derivative of the loss function with respect to β_k,

$$G_k = \nabla_{\beta_k} L_j = -2 \sum_i^I r_i \cdot \widehat{\mu_{\iota k}} \cdot w_i^C \cdot w_i^S \cdot \left(y_{ij} - r_i \cdot \sum_k^K \beta_k \widehat{\mu_{\iota k}}\right) + 2\beta_k. \quad (17)$$

AdRoit uses the function 'optim' from the R package 'stats' to perform the minimization[61],

providing the loss function (eq. 15) and the gradient (eq. 17). To get the final estimates of cell

type proportions, we rescale the coefficients β_k's to ensure a summation of 1,

$$\theta_k = \frac{\widehat{\beta_k}}{\Sigma_k^K \widehat{\beta_k}}. \tag{18}$$

Each mixture RNA-Seq sample j is independently estimated by the model described above.

Evaluation statistics

We compared the estimated cell type proportions with the ground truth by calculating the

following four statistics. The mean absolute difference (mAD) and root mean square deviation

(RMSD) are given by,

$$mAD = \frac{\Sigma_k^K |\theta_k - \theta_k^0|}{K} \tag{19}$$

$$RMSD = \frac{\Sigma_k^K (\theta_k - \theta_k^0)^2}{K} \tag{20}$$

where θ_k and θ_k^0 are the estimated proportions and true proportions, respectively.

Pearson correlation coefficient is computed as,

$$\rho_p = \frac{\Sigma_k^K (\theta_k - \overline{\theta_k})(\theta_k^0 - \overline{\theta_k^0})}{\sqrt{\Sigma_k^K (\theta_k - \overline{\theta_k})^2} \sqrt{\Sigma_k^K (\theta_k^0 - \overline{\theta_k^0})^2}} \tag{21}$$

where $\overline{\theta_k}$ and $\overline{\theta_k^0}$ are the means of the estimated proportions and true proportions,

respectively. Spearman correlation coefficient is given by,

$$\rho_s = \frac{\Sigma_k^K (r_k - \overline{r_k})(r_k^0 - \overline{r_k^0})}{\sqrt{\Sigma_k^K (r_k - \overline{r_k})^2} \sqrt{\Sigma_k^K (r_k^0 - \overline{r_k^0})^2}} \tag{22}$$

where r_k is the rank of θ_k.

Generation of synthesized bulk RNA-Seq and spatial transcriptomics data

Bulk RNA-Seq data used for the benchmarking were synthesized by adding up the raw UMI

counts per gene from all cells within a subject. Let t_k denote a cell of cell type k, $t_k \in 1, \ldots, T_k$,

where T_k is the total number of cells in cell type k. Let Y_{ij}^B be the count of gene i in a

synthesized bulk sample associated with the subject j, and X_{ijt_k} be the UMI count of gene i in

cell t_k within the subject j, then

731
$$Y_{ij}^B = \sum_k^K \sum_{t_k}^{T_k} X_{ijt_k} \qquad (23)$$

732   The true proportion of cell type k is given by,

733
$$\theta_k^0 = \frac{T_k}{\sum_k^K T_k} \qquad (24)$$

735   To simulate the spatial transcriptomic spots, we first sampled 10 cells without replacement

736   from each cell type, then weighed them by their respective mixing proportions before adding

737   them up. For example, to simulate a spot with p_k percent of cell type k, the count Y_{ij}^S of gene i

738   in a spatial spot j is given by,

739
$$Y_{ij}^S = \sum_k^K p_k \sum_{n=1}^{10} X_{ikn} \qquad (25)$$

where X_{ikn} is the UMI count of gene i in a selected cell n of cell type k. For each mixing scheme,

the simulation was repeated 100 times.

Single cell RNA sequencing of mouse dorsal root ganglion

As described previously[62], lumbar DRGs were isolated from five 18-week old adult C57BL/6

mice (three males and two females) and transferred to a dissociation buffer (Dulbecco's

modified Eagle's medium supplemented with 10% heat-inactivated Fetal Calf Serum) (Gibco; cat

A38400-02). To generate a single cell suspension, DRGs were subjected to a 2 step-enzymatic

dissociation followed by a mechanical dissociation. In brief, DRGs were first incubated with

0.125% collagenase P from Clostridium histolyticum (Roche Applied Science; cat #

11249002001) for 90 minutes in an Eppendorf Thermomixer C (37°C; intermittent 750 rpm

shaking for about 10 sec every 2 minutes). Then, DRGs were transferred to a Hank's Balanced

Salt Solution (HBSS, Mg^{2+} and Ca^{2+} free; Invitrogen) supplemented with 0.25% Trypsin

(Worthington biochemical corp.; cat # LS003707) and 0.0025% EDTA and incubated for 10

minutes at 37°C in the Eppendorf Thermomixer C. Trypsin was neutralized by the addition of 2.5

mg/ml MgSO4 (Sigma; cat #M-3937) and DRGs were triturated with Pasteur pipettes. The

resulting cell suspension was passed through a 70 μm mesh filter to remove remaining chunks

of tissues and centrifuged for 5 minutes at 2500 rpm at room temperature. The pellet was

resuspended in HBSS (Ca^{2+}, Mg^{2+} free; Invitrogen) and the cell suspension was run on a 30%

Percoll Plus gradient (Sigma GE17-5445-02) to further remove debris. Finally, cells were

resuspended in PBS supplemented with 0.04% BSA at a concentration of 200 cells/μl and cell

viability was determined using the automated cell analyzer NucleoCounter® NC-250™. The

suspended single cells were loaded on a Chromium Single Cell Instrument (10X Genomics) with

about 6000 cells per lane to minimize the presence of doublets. 2000-3000 cells per lane were

recovered. RNA-Seq libraries were constructed using Chromium Single Cell 3' Library, Gel Beads

& Multiplex Kit (10X Genomics). Single end sequencing was performed on Illumina NextSeq500.

Read 1 starts with a 26-bp UMI and cell barcode, followed by an 8-bp i7 sample index. Read 2

contains a 55-bp transcript read. Sample de-multiplexing, alignment, filtering, and UMI

counting were conducted using Cell Ranger Single-Cell Software Suite[63] (10X Genomics, v2.0.0).

Mouse mm10 Genome assembly and UCSC gene model were used for the alignment.

Data preprocessing

DRG single cell data

The UMI data output from the Cell Ranger Single-Cell Software Suite (10X Genomics, v2.0.0)

was analyzed using Seurat package[64] to assess the cell quality and identify cell types, similar to

what was described previously[42]. Cells with the number of detected genes less than 500 or over

15000, or with a UMI ratio of mitochondria encoded genes versus all genes over 0.1 were also

removed. The UMI data was normalized by the 'NormalizeData' method in Seurat with default

settings. To avoid potential sample-to-sample variation caused by the technical variation at

various experiment steps, we employed Seurat data integration method. The top 2000 variable

genes of each of the 5 DRG samples were identified using 'FindVariableFeatures' with

selection.method='vst'. Based on the union of these variable genes, the anchor cells in each

sample were identified by 'FindIntegrationAnchors'. All the samples were then integrated by

'IntegrateData'. We subsequently scaled the integrated data ('ScaleData') and performed

dimension reduction ('RunPCA'). Cells were then clustered based on the first 15 principal

components by applying 'FindNeighbors' and 'FindClusters' (resolution=0.6, algorithm=1).

Marker genes for each cluster were identified using 'FindAllMarkers'. Parameters were used

such that these genes were expressed in at least 25% of the cells in the cluster, and on average

2-fold higher than the rest of cells with a multiple-testing adjusted Wilcoxon test p value of less

than 0.01. The specificity of the canonical cell type-specific genes or cell cluster-specific genes

791 were further examined by visualizations (Supplementary Fig. 4) and used to define the cell type

792 for each cluster. At the end, the original UMI data from 17271 genes and 3352 cells that passed

793 the quality control were organized into a matrix (genes as rows and cell identifiers as columns).

794 This matrix, together with the cell type label for each cell therein, were loaded into AdRoit as

795 the reference profiles.

797 *Mouse brain single cell atlas data*

798 The scRNA-Seq reference data of the mouse brain were obtained from Zeisel et. al[32]. Among all

799 the available data, we only retained 96,572 cells that were acquired from the brain regions, had

800 an assigned cell type by the authors and a minimal total UMI of 1000. These cells corresponded

801 to 183 clusters at the finest taxonomy level in the original study. As many of the clusters are

802 highly similar, we decided to merge some of them to simplify the reference landscape. First, the

803 top 50 cluster enriched markers were derived using Scanpy[65] via the 'rank_genes_groups'

804 function (method='wilcoxon'), following the normalization ('normalize_per_cell'), log

805 transformation ('log1p') and regressing out ('regress_out') the variances associated with the

806 total UMI and the percentage of mitochondrial chromosome encoded genes per cell. Then, the

807 pair-wise overlapping p-values among the clusters were calculated using the top 50 marker

808 genes assuming a hypergeometric null distribution. Last, clusters with overlapping p-values

809 more significant than 1e-10 were merged and new names were assigned by combinedly

810 considering the original annotation, the molecular features and the specificity to certain brain

811 regions. A total of 46 cell types were determined that cover all the 12 brain regions and their

812 important substructures[40] (Supplementary Table 7). To make the reference dataset more

813 manageable in size and more balanced in the representation of cell types, we down sampled

814 each cluster to no more than 360 cells. A final set of 14,666 cells over 46 cell types was used for

815 the deconvolution of the mouse brain spatial transcriptome data.

817 *Human Islets single cell and bulk RNA-Seq data*

818 We used 1492 high quality human islets single cells and the associated annotations from Xin et

819 al[41]. The RPKM expression table was directly downloaded and used as-is. The RNA-FISH data

820 was also from this study[41]. The real bulk RNA-Seq data of human islets was acquired from a

821 large scale study referred in multiple publications[41,43,44]. We only included the data from donors

822 with a valid HbA1C level measurement in the regression analysis of the Beta cell proportion

823 with respect to the HbA1C level (Fig. 5c, Supplementary Table 13).

825 *Trabecular Meshwork single cell data*

826 We acquired the UMI counts, tSNE coordinates and the cell type annotation of 8759 high

827 quality cells from the authors of Patel et al[42]. Details of the quality control and the cell type

828 identification were presented in the original publication.

830 *Mouse Brain Spatial transcriptomics data by 10x Visium platform*

831 The filtered count matrix, tissue image and the spatial coordinates of a coronal section of an

832 adult C57BL/6 mouse brain were download from the 10x Genomics website (see Data

833 availability). A total of 2698 in-tissue spots were provided in the data set and used as-is.

Mouse Brian ISH images

The ISH images were directly downloaded from Allen mouse Brain Atlas[40] by searching the gene

names. The images were used without further editing except for cropping.

Data availability

The mouse DRG single cell data were deposited at NCBI GEO (accession number: GSE163252).

The bulk RNA-Seq and RNA-FISH data for human pancreatic islets were initially published as

aggregated data where the data processing and experimental procedure were described

therein[41,43,44]. We acquired the individual sample data from the authors and released them

along with the current study (Supplementary Table 10 and Supplementary Table 12). The other

public data analyzed in this study were obtained using GEO accession number GSE81608

(human pancreatic islets single cell data), NCBI SRA accession number PRJNA616025 (human

trabecular meshwork single cell data), and NCBI SRA accession number SRP135960 (mouse

brain single cell data). The 10x Genomics PBMC data was downloaded from the link

(https://support.10xgenomics.com/single-cell-gene-expression/datasets/1.1.0/pbmc3k), and

the Visium mouse brain spatial transcriptomic data was downloaded from the link

(https://support.10xgenomics.com/spatial-gene-

expression/datasets/1.1.0/V1_Adult_Mouse_Brain_Coronal_Section).

Code availability

AdRoit's source code is available on Github (https://github.com/TaoYang-dev/AdRoit).

Software

The statistical analyses were done with R statistical software (v3.6.0)[61] and python (v3.7.2)[66].

The packages used include Seurat (v3.0.1)[64], scanpy (v1.6.0)[65], dplyr (v0.8.0.1)[67], doParallel

(v1.0.14)[68], data.table (v1.12.4)[69], fitdistrplus (v1.1-1)[58], nnls (v1.4)[59], MuSiC (v0.1.1)[22],

BisqueRNA (v1.0.4)[21], SPOTlight (v1.0.4)[38], Cell2location (v0.05-alpha)[39], Stereoscope(v_03)[23].

Reference

1.   Wang, Z., Gerstein, M. & Snyder, M. RNA-Seq: A revolutionary tool for transcriptomics.

Nature Reviews Genetics (2009) doi:10.1038/nrg2484.

2.   Chu, G. C., Kimmelman, A. C., Hezel, A. F. & DePinho, R. A. Stromal biology of pancreatic

cancer. *Journal of Cellular Biochemistry* (2007) doi:10.1002/jcb.21209.

3.   Bussard, K. M., Mutkus, L., Stumpf, K., Gomez-Manzano, C. & Marini, F. C. Tumor-

associated stromal cells as key contributors to the tumor microenvironment. *Breast*

Cancer Research (2016) doi:10.1186/s13058-016-0740-2.

4.   Munn, D. H. & Bronte, V. Immune suppressive mechanisms in the tumor

microenvironment. *Current Opinion in Immunology* (2016)

doi:10.1016/j.coi.2015.10.009.

5.   Gonzalez, H., Hagerling, C. & Werb, Z. Roles of the immune system in cancer: From tumor

initiation to metastatic progression. *Genes and Development* (2018)

doi:10.1101/GAD.314617.118.

6.   Garner, H. & de Visser, K. E. Immune crosstalk in cancer progression and metastatic

spread: a complex conversation. *Nature Reviews Immunology* (2020)

doi:10.1038/s41577-019-0271-z.

7.   Singh, U. P. *et al.* Chemokine and cytokine levels in inflammatory bowel disease patients.

Cytokine (2016) doi:10.1016/j.cyto.2015.10.008.

8.   Van Lint, P. & Libert, C. Chemokine and cytokine processing by matrix metalloproteinases

and its effect on leukocyte migration and inflammation. *J. Leukoc. Biol.* (2007)

doi:10.1189/jlb.0607338.

9.   Zelová, H. & Hošek, J. TNF-α signalling and inflammation: Interactions between old

acquaintances. *Inflammation Research* (2013) doi:10.1007/s00011-013-0633-0.

10.   Koelman, L., Pivovarova-Ramich, O., Pfeiffer, A. F. H., Grune, T. & Aleksandrova, K.

Cytokines for evaluation of chronic inflammatory status in ageing research: Reliability

and phenotypic characterisation. *Immun. Ageing* (2019) doi:10.1186/s12979-019-0151-1.

11.   Landskron, G., De La Fuente, M., Thuwajit, P., Thuwajit, C. & Hermoso, M. A. Chronic

inflammation and cytokines in the tumor microenvironment. *Journal of Immunology*

Research (2014) doi:10.1155/2014/149185.

12.   Ståhl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by spatial

transcriptomics. *Science* (2016) doi:10.1126/science.aaf2403.

13.   Vickovic, S. *et al.* High-definition spatial transcriptomics for in situ tissue profiling. *Nat.*

Methods (2019) doi:10.1038/s41592-019-0548-y.

14.   Tang, F. *et al.* mRNA-Seq whole-transcriptome analysis of a single cell. *Nat. Methods*

(2009) doi:10.1038/nmeth.1315.

15.   Denisenko, E. *et al.* Systematic assessment of tissue dissociation and storage biases in

single-cell and single-nucleus RNA-Seq workflows. *Genome Biol.* (2020)

doi:10.1186/s13059-020-02048-6.

16.   Nguyen, Q. H., Pervolarakis, N., Nee, K. & Kessenbrock, K. Experimental considerations

for single-cell RNA sequencing approaches. *Frontiers in Cell and Developmental Biology*

(2018) doi:10.3389/fcell.2018.00108.

17.   Tanay, A. & Regev, A. Scaling single-cell genomics from phenomenology to mechanism.

Nature (2017) doi:10.1038/nature21350.

18.   Abbas, A. R., Wolslegel, K., Seshasayee, D., Modrusan, Z. & Clark, H. F. Deconvolution of

blood microarray data identifies cellular activation patterns in systemic lupus

erythematosus. *PLoS One* (2009) doi:10.1371/journal.pone.0006098.

19.   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.

Nat. Methods (2015) doi:10.1038/nmeth.3337.

20.   Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas

Reveals Inter- and Intra-cell Population Structure. *Cell Syst.* (2016)

doi:10.1016/j.cels.2016.08.011.

21.   Tsoucas, D. *et al.* Accurate estimation of cell-type composition from gene expression

data. *Nat. Commun.* (2019) doi:10.1038/s41467-019-10802-z.

22.   Wang, X., Park, J., Susztak, K., Zhang, N. R. & Li, M. Bulk tissue cell type deconvolution

with multi-subject single-cell expression reference. *Nat. Commun.* (2019)

doi:10.1038/s41467-018-08023-x.

23.   Andersson, A. *et al.* Single-cell and spatial transcriptomics enables probabilistic inference

of cell type topography. *Commun. Biol.* **3**, 565 (2020).

24.   Newman, A. M. *et al.* Determining cell type abundance and expression from bulk tissues

with digital cytometry. *Nat. Biotechnol.* (2019) doi:10.1038/s41587-019-0114-2.

25.   Myung, I. J. Tutorial on maximum likelihood estimation. *J. Math. Psychol.* (2003)

doi:10.1016/S0022-2496(02)00028-7.

26.   Bassett, R. & Deride, J. Maximum a posteriori estimators as a limit of Bayes estimators.

Math. Program. (2019) doi:10.1007/s10107-018-1241-0.

27.   Zhao, Y. & Simon, R. Gene expression deconvolution in clinical samples. *Genome*

Medicine (2010) doi:10.1186/gm214.

28.   Chiu, Y. J., Hsieh, Y. H. & Huang, Y. H. Improved cell composition deconvolution method

of bulk gene expression profiles to quantify subsets of immune cells. *BMC Med.*

Genomics (2019) doi:10.1186/s12920-019-0613-5.

29.   Kang, K. *et al.* CDSeq: A novel complete deconvolution method for dissecting

heterogeneous samples using gene expression data. *PLoS Comput. Biol.* (2019)

doi:10.1371/journal.pcbi.1007510.

30.   Qiao, W. *et al.* PERT: A Method for Expression Deconvolution of Human Blood Samples

from Varied Microenvironmental and Developmental Conditions. *PLoS Comput. Biol.*

(2012) doi:10.1371/journal.pcbi.1002838.

31.   Zaitsev, K., Bambouskova, M., Swain, A. & Artyomov, M. N. Complete deconvolution of

cellular mixtures based on linearity of transcriptional signatures. *Nat. Commun.* (2019)

doi:10.1038/s41467-019-09990-5.

32.   Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* (2018)

doi:10.1016/j.cell.2018.06.021.

33.   Donovan, M. K. R., D'Antonio-Chronowska, A., D'Antonio, M. & Frazer, K. A. Cellular

945  deconvolution of GTEx tissues powers discovery of disease and cell-type associated

946  regulatory variants. *Nat. Commun.* (2020) doi:10.1038/s41467-020-14561-0.

947 34. Phipson, B., Zappia, L. & Oshlack, A. Gene length and detection bias in single cell RNA

948  sequencing protocols. *F1000Research* (2017) doi:10.12688/f1000research.11290.1.

949 35. Chen, G., Ning, B. & Shi, T. Single-cell RNA-Seq technologies and related computational

950  data analysis. *Frontiers in Genetics* (2019) doi:10.3389/fgene.2019.00317.

951 36. Chen, D. & Plemmons, R. J. Nonnegativity constraints in numerical analysis. in *The Birth*

952  *of Numerical Analysis* (2009). doi:10.1142/9789812836267_0008.

953 37. Jew, B. *et al.* Accurate estimation of cell composition in bulk expression through robust

954  integration of single-cell information. *Nat. Commun.* (2020) doi:10.1038/s41467-020-

955  15816-6.

956 38. Elosua-Bayes, M., Nieto, P., Mereu, E., Gut, I. & Heyn, H. SPOTlight: seeded NMF

957  regression to deconvolute spatial transcriptomics spots with single-cell transcriptomes.

958  *Nucleic Acids Res.* (2021) doi:10.1093/nar/gkab043.

959 39. Kleshchevnikov, V. *et al.* Comprehensive mapping of tissue cell architecture via

960  integrated single cell and spatial transcriptomics. *bioRxiv* (2020).

961 40. Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature*

962  (2007) doi:10.1038/nature05453.

963 41. Xin, Y. *et al.* RNA Sequencing of Single Human Islet Cells Reveals Type 2 Diabetes Genes.

964  *Cell Metab.* (2016) doi:10.1016/j.cmet.2016.08.018.

965 42. Patel, G. *et al.* Molecular taxonomy of human ocular outflow tissues defined by single-

966  cell transcriptomics. *Proc. Natl. Acad. Sci.* **117**, 12856 LP – 12867 (2020).

967   43.   Xin, Y. *et al.* Pseudotime ordering of single human B-cells reveals states of insulin

968   production and unfolded protein response. *Diabetes* (2018) doi:10.2337/db18-0365.

969   44.   Gutierrez, G. D. *et al.* Gene signature of proliferating human pancreatic a cells.

970   *Endocrinology* (2018) doi:10.1210/en.2018-00469.

971   45.   Cerf, M. E. Beta cell dysfunction and insulin resistance. *Frontiers in Endocrinology* (2013)

972   doi:10.3389/fendo.2013.00037.

973   46.   Maedler, K. & Donath, M. Y. Beta-cells in type 2 diabetes: a loss of function and mass.

974   *Hormone research* (2004).

975   47.   Donath, M. Y. *et al.* Mechanisms of β-cell death in type 2 diabetes. *Diabetes* (2005)

976   doi:10.2337/diabetes.54.suppl_2.S108.

977   48.   Calanna, S. *et al.* Alpha- and beta-cell abnormalities in haemoglobin A1c-defined

978   prediabetes and type 2 diabetes. *Acta Diabetol.* (2014) doi:10.1007/s00592-014-0555-5.

979   49.   Kanat, M. *et al.* The Relationship Between β-Cell Function and Glycated Hemoglobin.

980   *Diabetes Care* **34**, 1006 LP – 1010 (2011).

981   50.   Nepton, S. Beta-Cell Function and Failure. in *Type 1 Diabetes* (2013). doi:10.5772/52153.

982   51.   Dolenšek, J., Rupnik, M. S. & Stožer, A. Structural similarities and differences between

983   the human and the mouse pancreas. *Islets* (2015) doi:10.1080/19382014.2015.1024405.

984   52.   Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature*

985   **445**, 168–176 (2007).

986   53.   Vieth, B., Parekh, S., Ziegenhain, C., Enard, W. & Hellmann, I. A systematic evaluation of

987   single cell RNA-Seq analysis pipelines. *Nat. Commun.* (2019) doi:10.1038/s41467-019-

988   12266-7.

989   54.   Wang, J., Roeder, K. & Devlin, B. Bayesian estimation of cell type-specific gene expression

990   with prior derived from single-cell data. *Genome Res.* (2021) doi:10.1101/gr.268722.120.

991   55.   Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome*

992   *Biol.* (2010) doi:10.1186/gb-2010-11-10-r106.

993   56.   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-

994   Seq data using regularized negative binomial regression. *Genome Biol.* (2019)

995   doi:10.1186/s13059-019-1874-1.

996   57.   Svensson, V. Droplet scRNA-Seq is not zero-inflated. *Nature Biotechnology* (2020)

997   doi:10.1038/s41587-019-0379-5.

998   58.   Delignette-Muller, M. L. & Dutang, C. fitdistrplus: An R package for fitting distributions. *J.*

999   *Stat. Softw.* (2015) doi:10.18637/jss.v064.i04.

59.   Mullen, Katharine M., I. H. M. van S. nnls: The Lawson-Hanson algorithm for non-

negative least squares (NNLS). *R Packag. version 1.4* (2012).

60.   Byrd, R. H., Lu, P., Nocedal, J. & Zhu, C. A Limited Memory Algorithm for Bound

Constrained Optimization. *SIAM J. Sci. Comput.* (1995) doi:10.1137/0916069.

61.   The R Core Team. R: A Language and Environment for Statistical Computing. *R*

Foundation for Statistical Computing (2019).

62.   Alessandri-Haber, N. *et al.* Hypotonicity induces TRPV4-mediated nociception in rat.

Neuron (2003) doi:10.1016/S0896-6273(03)00462-8.

63.   Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat.*

Commun. (2017) doi:10.1038/ncomms14049.

64.   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* (2019)

doi:10.1016/j.cell.2019.05.031.

65.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: Large-scale single-cell gene expression data

analysis. *Genome Biol.* (2018) doi:10.1186/s13059-017-1382-0.

66.   van Rossum, G. & Drake, F. L. *Python 3 Reference Manual. Scotts Valley, CA* (2009).

67.   Wickham, H. & Francois, R. dplyr: A Grammar of Data Manipulation. *R Packag. version*

0.4.2. (2015).

68.   Weston, S., Calaway, R. & Tenenbaum, D. *doParallel: Foreach Parallel Adaptor for the*

Parallel Package. *Cran* (2014).

69.   Dowle, M. & Srinivasan, A. data.table: Extension of 'data.frame'. R Package Version

1.12.8. *Manual* (2019).

1021

Acknowledgements

We thank Yurong Xin for pointing us to the relevant public data resource. We also thank Gabor

Halasz and Yuan Zhu for the helpful discussions regarding the memory usage, and Shawn

Mishra for the manuscript proofreading.

1026

Author contributions

Y.B., T.Y., N.A.-H., M.L.-F., L.E.M. and G.S.A. designed the research. T.Y., Y.B., and W.F.

developed the algorithm. T.Y., Y.B. and J.K. participated in the data analysis. M.S. and R.B.

performed the DRG tissue collection. C.A. performed the single cell library preparation and

sequencing experiment. Y.B., T.Y. and N.A.-H. wrote the manuscript.

1032

Competing interests

1034 T.Y., Y.B., W.F. and G.S.A. have filed a patent application relating to the AdRoit computational

1035 framework. M.L.-F. is an employee of Cellular Longevity. All other authors are employees

1036 and/or shareholders of Regeneron Pharmaceuticals, although the manuscript's subject matter

1037 does not have any relationship to any products or services of this corporation.

1038

Figure legends

Fig. 1: Schematic representation of AdRoit computational framework. a, AdRoit inputs

compound (bulk or spatial) RNA-Seq data, single cell RNA-Seq data and cell type annotations. It

first selects informative genes and estimates their means and dispersions, then computes the

cell type specificity of genes. Depending on the availability of multiple samples, cross-sample

gene variability is derived from either the compound RNA-Seq, or the single cell data (dashed

arrow; see also Methods). Lastly the gene-wise scaling factors are computed to correct the

platform bias between the compound and the single cell RNA-Seq data. These quantities are

used in a weighted regularized model to infer the cell type composition. **b,** A mock example to

illustrate the role of the gene-wise scaling factor. Ideally, an accurate estimation of the cell

proportions should be represented by the slope of the green line, however fitting in the

presence of outlier genes would result in the red line. Outlier genes exist because the platform

bias affects genes differently. AdRoit adopts an adaptive learning approach that first learns a

coarse estimation of the slope (red line), from which the gene-wise scaling factors are derived

and applied to the outlier genes, moving them toward the green line. The more deviated the

gene, the larger the correction (i.e., longer arrows). After the adjustment, the new estimated

slope (blue line) is closer to the truth (green line), thus is a more accurate estimation.

1056

Fig.2: Benchmark on simulated bulk data generated from the trabecular meshwork (TM)

single cells. a, AdRoit has the closest estimation to the true cell proportion comparing to

Bisque, MuSiC and SPOTlight. Each dot is a cell type from a donor. **b,** For each cell type in TM,

AdRoit has the smallest differences from the true cell type proportion and the smallest variance

of estimates across the 8 donors. For each cell type, a dot on the graph denotes a donor, and

the bars represent the $1.5 \times$ interquartile ranges. The reference used for deconvoluting each

synthetic bulk sample excludes the cells used to synthesize that sample (leave-one-out).

1064

Fig. 3: AdRoit can achieve a high granularity and exhibits high sensitivity and specificity in

complex tissues. a. AdRoit is accurate in deconvoluting the simulated bulk samples that contain

a mixture of similar cell types from myeloid or lymphoid lineage. The vertical dashed lines

indicate the true mixing proportions. $CD14^+$ monocytes, $FCGR3A^+$ monocytes and dendritic cells

(DC) were mixed under three schemes of proportions: 0.33:0.33:0.33 (mix0), 0.1:0.45:0.45

(mix1) and 0.1:0.3:0.6 (mix2). The same ratios were applied to the mixtures of naïve $CD4^+$ T,

memory $CD4^+$ T and $CD8^+$ cells. **b,** AdRoit's estimates are more accurate and specific than those

from Bisque, MuSiC and SPOTlight on synthetic samples that contain only 6 out of the 12 cell

types. The deconvolution was done using all 12 cell types as the reference. A pair of size-

matched blue (true value) and red (estimated value) bubbles indicate an accurate prediction.

Red-only and blue-only bubbles mark false positives and false negatives, respectively. **d,**

Receiver operating characteristic (ROC) curve shows AdRoit has a significantly higher area

under the curve (AUC) than other methods, meaning better sensitivity and specificity. **e,**

Scatterplots between the ground truth and the deconvoluted cell proportions in the simulated

bulk samples of high complexity (mixtures of 30 cell types). **f,** ROC curves show AdRoit has the

best AUC among all methods on highly complex cell constitutions.

1081

Fig. 4: Benchmark on scRNA-Seq data from dorsal root ganglion (DRG) where many closely

related subtypes of neuronal cells exist. a, 14 cell types are identified from scRNA-Seq samples

of 5 mice, including multiple subtypes of neurofilaments (NF), peptidergic (PEP), and non-

peptidergic (NP) neurons. **b,** Benchmarking with the synthetic data shows the cell type

proportions inferred by AdRoit are highly accurate. In particular, AdRoit remains a high

accuracy when the cells are rare (e.g., < 5%; see also the zoom-in inserts). Each dot represents a

cell type from one sample. **c,** For each sample, mAD, RMSD, Pearson and Spearman correlations

are compared across four methods. AdRoit has the lowest mAD and RMSD, and the highest

Pearson and Spearman correlations. In addition, AdRoit's estimation is the most stable across

samples. Each dot on the boxplot is a sample. Same samples are chained by the dotted lines

across the metrics. The deconvolution was done by using the leave-one-out strategy.

1093

Fig. 5: AdRoit is more accurate and sensitive on spatial spots simulated from real DRG cells. a,

Estimations from AdRoit, Cell2location,Stereoscope and SPOTlight on simulated spatial spots

that contain 5 PEP neuron subtypes. True mixing proportions are denoted by the red dashed

lines. Three schemes are presented: 1) the proportions of 5 PEP cell types are the same and

equal to 0.2; 2) PEP1_Dcn is 0.1 and the other 4 are 0.225; 3) PEP1_Dcn and

PEP1_S100a11.Tagln2 are 0.1, PEP1_Slc7a3.Sstr2 and PEP2_Htr3a.Sema5a 0.2 are 0.2, and

PEP3_Trpm8 is 0.4. **b,** The performance of AdRoit, Cell2location, Stereoscope, and SPOTlight in

estimating rare cell populations in the spatial spots. The spots contain a mixture of three PEP

cell subtypes (i.e., PEP1_Slc7a3.Sstr2, PEP2_Htr3a.Sema5a and PEP3_Trpm8), with the percent

of PEP3_Trpm8 ranging from 1% to 10% and the other two cell types sharing the remaining

proportion equally. **c,** Compare the rate of detecting rare cells in simulated spots. An inferred

percent greater than 0.5% is deemed as a positive detection. Six sets of cell mixtures are

employed: NF_Calb1 with NF_Pvalb and NF2_Ntrk2.Necab2 (NP subtypes), NP_Mrgpra3 with

NP_Mrgprd and NP_Nts (NP subtypes), PEP3_Trpm8 with PEP1_Slc7a3.Sstr2 and

PEP2_Htr3a.Sema5a (PEP subtypes), NF_Calb1 with Th, satellite glia and endothelial (NF_Calb1

+ others), NP_Mrgpra3 with Th, satellite glia and endothelial (NP_Mrgpra3 + others), and

PEP_Trpm8 with Th, satellite glia and endothelial (PEP_Trpm8 + others). In each set, the first

cell type listed is the target of detection and varies its percent from 1% to 10%. The rest cell

types split the remaining proportion evenly. The red dashed lines mark the detection rate of

90%.

1114

Fig. 6: Applications to real-world bulk RNA-Seq data and mouse brain spatial transcriptome

data. a, The deconvoluted cell compositions in the real bulk RNA-Seq data of human Islets are

highly reproducible for the repeated samples from the same donor. **b,** AdRoit estimation of the

cell type proportions agrees with the RNA-FISH measurements. **c,** AdRoit-inferred Beta cell

proportions in type 2 diabetes patients are significantly lower than those in healthy subjects. In

addition, the estimated proportions have a significant negative linear association with the

donor HbA1C levels. **d,** The spatial mapping of four mouse brain cell types is consistent with the

locations of four region-specific markers shown on the ISH images obtained from Allen mouse

brain atlas[40]. The four genes, Spink8, C1ql2, Clic6 and Synpo2, were identified by Zeisel et al[32] as

markers of the hippocampal field CA1, dentate gyrus, choroid plexus and thalamus,

respectively.

1126

1127

1128

1129

1130

1131

1132

1133