Transcription of macromolecular N degrading genes provides a window into organic nitrogen

decomposition in soil

Ella T. Sieradzki[1], Erin E. Nuccio[2], Jennifer Pett-Ridge[2], Mary K. Firestone[1,3#]

[1]Department of Environmental Science, Policy and Management, University of California,

Berkeley, California, USA

[2]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore,

California, USA

[3] Climate & Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley,

California, USA

[#]Address correspondence to: mkfstone@berkeley.edu

Abstract

Background: Nitrogen (N) is commonly limiting in soil in part because most N is present in

macromolecular organic compounds and not directly available to plants. Using extracellular

enzymes, the soil microbial community present near roots (rhizosphere) is largely responsible for

transforming organic substrates, which yields nitrogen in forms that are plant-available.

Microbial genetic potential (genes) and the cycling of inorganic nitrogen pools have been

extensively documented and widely explored in rhizospere soil. But there is scarce information

on microbial gene expression for macromolecular nitrogen decomposition and how it relates in

space and time to life stages of plant roots. Here we use a suite of time-resolved

metatranscriptomes from rhizosphere and bulk soil to follow bacterial and fungal extracellular

protease and chitinase expression during rhizosphere aging. We also explore the effect of adding

plant root litter as a distinct source of macromolecular carbon and nitrogen.

Results: Expression of extracellular proteases in rhizosphere soil increased with plant growth

in the absence of litter, whereas chitinase (*chit1*) was highly upregulated in the detritusphere.

Structural groups of proteases were dominated by serineproteases, despite the higher abundance

of metalloproteases previously observed in soil and aquatic systems. Extracellular proteases of

Betaprotebacteria were more highly expressed in the rhizosphere, whereas those of

Deltaroteobacteria and Fungi responded strongly to the presence of litter. We identified distinct

functional guilds of taxa specializing in decomposition of proteins in the rhizosphere,

detritusphere and in the vicinity of aging roots. We also identify a sub-guild that appears to

specialize in protein decomposition in the presence of growing roots as well as litter that

increases its activity in aging rhizosphere soil, which may be particularly N limited. Taxonomic

membership of these guilds partially overlapped with functional guilds identified by carbon

decomposing enzymes, making up multifunctional guilds.

Conclusions: We found organic N decomposers have distinct spatiotemporal preferences that are

selected during plant growth. Rhizosphere microbes which could potentially benefit plants by

degrading macromolecular nitrogen were not comparatively active in bulk soil, even in the

presence of root litter, implying that they are enhanced near plant roots and less competitive in

bulk soils.

Keywords

Rhizosphere, microbiome, protease, chitinase, functional guilds, metatranscriptome, gene

expression, litter

Background

Plants are commonly limited by nitrogen (N) in temperate soils since their access to the

largest soil N pools is constrained by the activity of microbes responsible for mineralization of

macromolecular nitrogen and N_2 fixation [1–3]. To a large degree, plants depend on microbial

degradation of macromolecular forms of N such as proteins and chitin [4–6] and can potentially

increase N availability by releasing exudates that stimulate microbial turnover of organic N pools

(via 'priming') [7–10]. Depolymerization of high molecular weight detrital organic N is one of

the rate-limiting steps in soil N mineralization [11,12] and depends on the activity of

extracellular enzymes such as lysozyme, protease, chitinase, nuclease and urease. The resulting

N monomers can be further broken down to NH_4 or in some cases taken up directly by the plant

[13]. While the genetic potential and bulk-scale activity of extracellular N degrading enzymes is

well established, the spatial and temporal dynamics of gene expression underpinning N

mineralization are ill-defined. In addition, soil microorganisms capable of degrading

macromolecular N are highly diverse [14], and it is unclear when and where specific taxonomic

groups are active [15].

Patterns and controls of soil N depolymerization and mineralization may be highly

dependent on the habitat (e.g., rhizosphere, detritusphere), since availability of organic-N

substrates varies in both time and space, as does the prevalence of fungi, fauna and bacterial

degraders, which can enable or limit N-cycling. In the rhizosphere, organic N is available as

amino acids, nucleotides, niacin, and choline (derived from plant exudates [10]), lignoproteins

and aromatics from sloughed off cells, nucleic acids, and microbial cell wall amino sugar

polymers (N-acetylglucosamine, N-acetylmuramic acid) derived from the bloom of cells that

develops as roots grow [15]. Moreover, root exudates can drive destabilization of mineral-

associated organic matter (MAOM), creating an additional source of macromolecular N [13].

Prior work indicates that root exudates can increase degradation of soil organic matter by up to

380% [16]. Chitinase and protease enzyme activities were shown to be significantly upregulated

in the rhizosphere of wild oatgrass (*Avena spp.*), and N mineralization was highly dependent on

quorum sensing [17]. This contrasts with the detritusphere, which has a relatively higher

proportion of aromatic N and lignoproteins. Protease enzyme activity has been shown to be

enhanced by litter addition [18] and root exudates [19], as well as vary with type of plant litter

[20,21]. Obviously, these spatial habitats (rhizosphere and detritosphere) can overlap—as when a

root initially grows through bulk soil, or when a growing root begins to age and resemble

detritus.

Soil N pools and by proxy, N mineralization activity, change with time [22] in part due to

the succession of both exudate quality and microbial communities [10,23]. At the bulk soil level,

activity of extracellular proteases is thought to drive soil N cycling [12], but what controls this

activity is not always clear [24]. Subtle distinctions caused by time and soil habitat are likely

masked because most studies are conducted on whole soils, which contain a mixture of

rhizosphere, detritusphere and bulk regions. In grassland soils, the rhizosphere is a particularly

critical hotspot for microbial biomass and activity due to rapid recycling of root exudation and

debris, and functions as a quasi-digestive system decomposing molecules inaccessible to plants.

However, root exudation, which serves as a carbon source for rhizosphere bacteria, declines near

older root sections [25]. Therefore, it has been hypothesized that bacteria in a mature rhizosphere

environment may be forced to target less labile, higher C:N sources of organic N such as plant

litter [26]. Indeed, it has been shown that the mature *Avena* sp. rhizosphere has an increased organic N demand relative to the young rhizosphere, where the mature rhizosphere has higher rates of gross nitrogen mineralization and ammonia consumption [28].

Of the vast number of microorganisms that exist in a soil, at any given time, only a small proportion (e.g., 8%) are active in the rhizosphere [15,17,27,28]. Gene expression studies show that soil bacteria, in the rhizosphere and in the presence of plant litter, form functional guilds [29–31]. However, there has been little focus on guilds based on macromolecular N degradation. To maintain cellular C:N stoichiometry, microbial carbon acquisition must be accompanied by N uptake. Therefore, we postulate that an analogous functional guild organization occurs in expression of extracellular proteases as with CAZy guilds, as exoproteases are diverse enzymes that target a variety of substrates and have also been shown to be influenced by soil characteristics [32–34]. However, while a general balance of C and N degradation is expected to maintain cellular C:N, it does not follow that the same taxa should be involved in both processes. The capacity to utilize specific C and N forms is taxon-specific; e.g., some bulk soil bacteria may degrade carbohydrates as a C source and take up inorganic N, whereas some rhizosphere bacteria may take up simple root exudates as a C source but degrade proteins as an N source.

We hypothesized that community-wide expression and taxon-specific differential expression of extracellular proteases and chitinases differed in rhizosphere vs. bulk soil during root growth and that expression profiles would be altered by root litter amendments. To address that hypothesis, we used metatranscriptomes collected over time from rhizosphere microcosms, using a robust, well-developed experimental approach [17,23,26,29] with a well-characterized annual grassland soil from Northern California. Mature wild oat grass *Avena fatua* [22] plants were monitored for 3 weeks to determine the role of diverse indigenous rhizosphere bacteria in

macromolecular nitrogen degradation by extracellular proteases and chitinases throughout root

growth and aging. In addition, we compared functional guilds based on extracellular protease

expression with guilds defined by expression of carbohydrate active enzymes (CAZy) capable of

degrading root litter.

Methods

Experimental design, sample collection and sequencing

The experimental design, sample collection and sequence data processing are described

in detail in [29]. Briefly, common wild oat *Avena fatua* was grown for six weeks in rhizobox

microcosms containing soil from the Hopland Research and Extension Center (HREC) in

northern California, a Bearwallow–Hellman loam (pH 5.6, 2% total C) packed at field bulk

density. Roots were then allowed to grow into a sidecar soil region with a transparent wall,

where the root growth timeline was marked at 3 days, 6 days, 12 days and 22 days. In half of the

sidecars, the soil was amended with dried *A. fatua* root detritus (C:N = 13.4) chopped to 1 mm.

Bulk soil bags, inaccessible to roots, were placed in each sidecar; a bulk soil treatment amended

with root detritus was also included. At each timepoint, three replicate microcosms were

destructively harvested for paired rhizosphere and bulk soil, both detritus amended and

unamended, yielding a total of 48 samples representing four treatments at four timepoints:

rhizosphere, rhizosphere + litter, bulk, and bulk + litter.

Harvested soil (1 g) was placed immediately in Lifeguard Soil Preservation Reagent

(MoBio) and processed according to the company protocol. Roots and supernatant were

removed, and the soil was stored in -80°C. DNA and RNA were co-extracted using a phenol-

chloroform procedure [35,36] and separated with an AllPrep kit (Qiagen). RNA was DNase

treated (TURBO DNase, Thermo-Fisher Scientific), depleted in ribosomal RNA (Ribo-Zero

rRNA Removal Kit, Epicentre) and reverse transcribed into cDNA. Metatranscriptomes were

sequenced for 48 samples on an Illumina HiSeq 2000 2x150 (TruSeq SBS v3) at the Joint

Genome Institute (JGI), in Walnut Creek CA.

Expression of chitinase and extracellular protease genes identified in assembled

metatranscriptomes

Raw reads were quality-trimmed (Q20) and rRNA and tRNA reads were removed.

Library size, evenness, richness and Shannon diversity were comparable between experimental

groups with a mean library size of 43 M paired end reads after filtering. Quality-controlled

metatranscriptomic reads were assembled into contigs within each sample [29]. Contigs smaller

than 200bp were discarded and the remaining contigs from all samples were clustered at 99%

identity with cd-hit-est keeping the longest sequence as the cluster representative [37]. Open

reading frames (ORFs) were predicted using Prodigal [38]. Extracellular protease ORFs were

identified by reciprocal BLAST to extracellular peptidases from the MEROPS database [14,39].

ORFs were assigned a peptidase group (serine-, metallo-, cysteine-peptidase and others) by their

best reciprocal BLAST hit. Taxonomy was determined by best BLAST hit to the NCBI non-

redundant database (NR, access date July 29[th] 2019). Chitinase ORFs were identified using six

chitinase hidden Markov models (HMMs) from the Kyoto Encyclopedia of Genes and Genomes

(KEGG): K01183, K03791, K13381, K17523, K17524 and K17525. Only the first three were

detected in our dataset. Reads were then mapped back to ORFs requiring minimum identity 95%

and 75% breadth using bbmap [40]. Read counts were normalized using DESeq2 [41]. Heat

maps of normalized counts were generated in R using the heatmap2 function in gplots [42].

Normalized transcript counts per gene per time point were compared between groups

(rhizosphere, litter and litter-amended rhizosphere) using ANOVA and Tukey post-hoc test.

Boxplots were generated in ggplot2 [43].

Expression of extracellular protease genes from a curated collection of Hopland-soil genomes

184   The quality-controlled reads were mapped using BBsplit [40] requiring 80% identity

against a dereplicated reference database of 282 HREC soil genomes including isolates [10],

single amplified genomes (SAGs) [29], metagenomic assembled genomes (MAGs) (NCBI

PRJNA517182) and stable isotope probing MAGs (SIP-MAGs) [44]. On average, 12.3% (range

6.2-31.9%) of the reads per library mapped unambiguously to genomes from the reference

database. This additional approach was taken to investigate gene expression within the context of

a genome and to search for overlap in guild membership between extracellular protease defined

guilds and previously defined CAZy guilds [29]. Three MAGs that were classified as unknown

or domain bacteria in previous studies were assigned taxonomy using GTDB-Tk version 1.3.0

[45,46]. Verification of the taxonomic assignment of MAG Burkholderiales_62_29 was done

using GToTree [47] with complete reference genomes of Betaproteobacteria from RefSeq (Feb

28, 2020).

196   Open reading frames (ORFs) were predicted in all genomes using prodigal [38] and

annotated using KEGG [48] and ggKbase (http://ggkbase.berkeley.edu). Extracellular proteases,

which do not have hidden Markov models (HMMs) capable of separating them from intracellular

proteases, were identified by gene nucleotide identity of at least 90% and coverage of at least

60% to *de novo* assembled ORFs of extracellular protease from the metatranscriptomes. Gene

counts were identified using Rsubread featureCounts [49].

Differential expression of chitinases in the genome collection was already performed in a

previous publication on expression of carbohydrate active enzymes (CAZy) and was therefore

not repeated here [29].

Statistical analyses

All features and their abundance (represented by metatranscriptomic read counts

normalized to sequencing depth) were analyzed with DESeq2 [41] requiring p-value < 0.05

(adjusted for multiple comparisons). Ordination and visualization were conducted in R using

ggplot2 [43] and vegan [50]; PERMANOVA (vegan function adonis) was used to detect

significant treatment factors affecting expression of nitrogen depolymerization genes. We define

'guilds' as groups of taxa with similar gene upregulation patterns of extracellular proteases in

both time and soil habitat. Guilds were assigned by hierarchical clustering based on differential

expression of extracellular proteases compared to unamended bulk soil, generating four

functional guilds: "Rhizosphere", "Detritusphere", "Aging root", and "Low response". Effects of

the four experimental treatments (bulk, rhizosphere, litter and litter-amended rhizosphere) on

protease gene expression were assessed by ANOVA (p adjusted for multiple comparisons), N=3

per timepoint.

Results

Expression of enzymes targeting macromolecular N

Multiple genes related to degradation of macromolecular N were identified in this

experiment: extracellular nuclease (Xds), urease (UreABC), lysozyme, chitinase (chit1) and

extracellular proteases. Expression of extracellular proteases was an order of magnitude higher

than the extracellular N degrading enzymes (sup. fig. S1), and like chitinases, was affected by

the presence of litter and roots. In contrast, extracellular nuclease (Xds), urease (UreABC) and

lysozyme were not influenced by either living or decaying roots, thus, we chose to focus on

extracellular proteases and chitinases.

Normalized transcript abundance for 4948 extracellular protease genes from bacteria

(4846) and fungi (102) were quantified in the sampled soils. Overall, extracellular protease

expression was significantly affected by time (3-way PERMANOVA, $F=2.8$, $p=0.038$), litter

amendment ($F=118.9$, $p<0.001$) as well as interactions between time:treatment ($F=26.9$,

$p<0.001$), time:location ($F=7.7$, $p<0.001$) and location:treatment ($F=15.5$, $p<0.001$). Expression

in unamended soil (no litter) generally increased over time (fig. 1A), whereas litter-amended soil

had initially high expression that then decreased and leveled off over time (fig. 1B).

The 73 distinct chitinase transcripts were expressed at a substantially lower level than the

extracellular proteases. The expression of the *chit1* gene increased over time in litter-amended

soils and was higher than in the non-litter treatments (3-way PERMANOVA, $p<0.001$) (fig. 1C).

Expression of fungal chitinases was lower than bacterial chitinases, and both were generally

lower than eukaryotic chitinases (fig. 1D). Transcripts for chitinase KEGG orthologs *CHI3L1_2*,

CHI3L1_4 and *chiA* were not detected at all and the putative chitinase K03791 had low

expression and no significant effects with time or treatment (data not shown). Chitinase CHID1

was detected but was most closely related to plants and therefore disregarded. A companion

study analyzed differential expression of chitinases within our curated genome collection and

while only one MAG was found to upregulate chitinase, it did so in the presence of litter [29].

Figure 1: Aggregated expression of extracellular proteases in bulk and rhizosphere soils from

common wild oat *Avena fatua* microcosms, grown without root litter amendment (A), or with

litter amendment (B); expression of chitinase gene *chit1* (C) and mean counts of chitinases at the

domain level (D) for unamended bulk soil (B), litter-amended bulk soil (BL), unamended

rhizosphere (R) and litter-amended rhizosphere (RL). Boxplots represent 75% of the data,

whiskers denote 90% of the data and dots (e.g. panel A bulk litter 6 days) represent outliers. Note

that the scale of the Y axis varies between panels.

Structural groups of extracellular proteases

Structural groups of extracellular proteases can be soil-specific and pH dependent [32].

The main groups of extracellular proteases found in our assembled metatranscriptomes were

serine-, metallo- and cysteine-proteases (2679, 1949 and 496, respectively out of 5295 variants

clustered at 99% identity). Aggregated expression patterns of all variants in each group reflected

the same order (serine>metallo>cysteine). Expression of serine- was consistently higher than

metallo-protease across all treatments (ANOVA F=2392, p=0) (fig. 2, sup. table S4). Expression

of serine-proteases was also significantly greater in the presence of litter compared to no litter

(diff=2337, p<0.001), but root litter amendment did not affect expression of metallo- or cysteine-

proteases, and there was no significant effect of time or location on these structural groups (sup.

table S4).

Figure 2: Aggregated normalized expression of extracellular protease structural groups over time

in bulk, rhizosphere and root litter-amended soils. Normalized expression per replicate is plotted

as dots on top of the bars. Experimental groups shown per time point are (left to right):

unamended bulk soil (B), litter-amended bulk soil (BL), unamended rhizosphere (R) and litter-

amended rhizosphere (RL). Extracellular protease groups are (left to right): serine-, metallo- and

cysteine-protease.

Taxonomy of extracellular proteases

De novo assembled ORFs of extracellular proteases were taxonomically assigned by

BLASTP best hit against the NCBI non-redundant database (NR). Since extracellular proteases

are not marker genes, we considered taxonomic assignments only at the order level or higher,

with the exception of *Rhizobacter*, for which amino acid percent identity values were extremely

high (84±8.4% amino acid identity).

In the rhizosphere, Betaproteobacteria exoproteases were significantly upregulated

(ANOVA, $p<0.05$) (fig. 3A). Of 547 variants of exoproteases from this class, 442 were assigned

as Burkholderiales. In contrast, at most timepoints, proteases of Cyanobacteria and

Thaumarchaeota were significantly downregulated in the rhizosphere compared to bulk soil (fig.

3B; sup. table S1). In multiple other taxonomic groups, proteases were significantly upregulated

only in the presence of litter: Fungi (fig. 3C), class Deltaproteobacteria (fig. 3D), as well as

highly represented phyla Bacterioidetes and Verrucomicrobia (sup. fig. S2; sup. table S1) and

classes Chitinophagia and Gammaproteobacteria (sup. table S2). Interestingly, exoprotease

expression declined with time for the predatory bacteria Myxococcales (fig. 3E),

Bdellovibrionales and Cytophagia, while clades such as Chloroflexi (fig. 3F) and Actinobacteria

had increased exoprotease expression at the final sampling point. While the Actinobacteria and

Acidobacteria had a high number of variants and high expression of exoproteases, we did not

detect a significant effect of either time or soil habitat/ amendment. Normalized protease

expression data are summarized by phylum (sup. fig. S2), class (sup. fig. S3) and order (sup. fig.

S4), and ANOVA F and p values are in sup. table S1, sup. table S2 and sup. table S3

respectively.

Figure 3: Expression of extracellular proteases over time for select taxonomic groups: (A)

Betaprotebacteria (n=547) were upregulated in the rhizosphere, (B) Thaumarchaeota (n=112)

were downregulated in the rhizosphere, (C) Fungi (n=99) were upregulated in the presence of

litter, (D) Detaproteobacteria (n=228) were upregulated in the presence of litter, (E)

Myxococcales (n=147) were upregulated in the presence of litter and decreased over time and (F)

Chloroflexi (n=153) increased in the last time point. The legend in panel F applies to all panels.

Note that as expression levels varied over taxonomic groups and to emphasize patterns, the y

axes do not have the same scale.

Functional guilds

To define functional guilds with a population-centric analysis, we mapped transcriptome

reads to a collection of 282 genomes and metagenome-assembled genomes generated from the

same soil and site [29]. Read counts were used to determine differential expression compared to

unamended bulk soil. Each genome contained multiple genes coding for extracellular proteases.

Hierarchical clustering of the mean differential expression of extracellular protease genes within

each genome was used to define four functional guilds: "rhizosphere", "detritusphere", "aging

root" and "low response" (fig. 4) following definitions first described in [29]. Similarly,

hierarchical clustering based on differential expression of the most highly upregulated *de novo*

assembled ORFs compared to unamended bulk soil revealed clear rhizosphere and detritusphere

guilds (sup. fig. S5). There was some overlap between guilds identified here and guilds identified

based on expression of CAZy genes, but the degree of overlap varied by guild (sup. fig. S6).

Within the aging root guild, we noticed one member, representing an aggregated

population transcriptome, that increased differential expression of extracellular proteases at the

last time point in the presence of both roots and litter more than would have been predicted by

the sum of the treatments alone. As this pattern differed from the rest of the guild, we identified

this member as a sub-guild labeled "late synergist" due to its expression pattern. This MAG,

Burkholderiales_62_29, was identified by two independent phylogenomic analyses as

Rhizobacter (sup. fig. S7). Burkholderiales_62_29 has 16 different extracellular protease genes,

all of which had similar significant upregulation patterns. While the sub-guild here contained

only a single *Rhizobacter* MAG, the implication of mapping transcriptomic reads at 80% identity

is that each MAG represents a taxonomic "cloud" of at least genus-level diversity. A 16S-rRNA

survey of the same samples revealed 10 operational taxonomic units (OTUs) of order

Burkholderiales which, like *Rhizobacter*, are not assigned a family [29]. Additionally, this sub-

guild may include more taxa for which we have no MAGs and therefore could not be included in

this analysis.

Figure 4: Functional guilds defined by hierarchical clustering of extracellular protease

differential gene expression during a 22-day *Avena fatua* microcosm experiment. Each row

represents the mean differential expression of extracellular protease genes mapped to a reference

genome. Treatments are (left to right): litter-amended bulk soil, litter-amended rhizosphere and

unamended rhizosphere. Time points are indicated at the bottom in days. We note that a genome

may contain more than one exoprotease gene and that reads were mapped at 80% identity,

therefore each genome also represents closely related taxa. Differential expression values per

gene that were not statistically significant were converted to zero (0) before averaging.

Discussion

Nitrogen in grassland soil is present mostly in the form of organic macromolecules such

as proteins and chitin [22,51]. Rhizosphere bacteria can acquire nitrogen through degradation of

extracellular macromolecular N, and this process can be enhanced by root exudates especially in

unfertilized or N limited soil [1,9,52]. Amino acids resulting from protein degradation are taken

up by microorganisms, and roughly 30% of the amino-acid carbon is respired, leading to

excretion of excess ammonium which could benefit the plant [53,54]. While various types of

extracellular enzymes degrade macromolecules containing N, we focus here on protease and

chitinase as they were significantly affected by litter amendment and by proximity to the plant

(bulk soil vs. rhizosphere). A previous study based on the same experiment showed that there

was very little change in the bacterial or fungal community composition throughout the

experiment [29].

Chitinase expression, while two orders of magnitude lower than that of proteases,

increased over time in litter-amended samples. This effect could imply that the litter attracted

saprotrophic fungi and possibly arthropods, the cell walls of which contain chitin [55]. As most

of the chitinases were most closely related to eukaryotic enzymes, it is likely that the chitin-

containing hyphae were preyed upon by soil fauna. A previous study with *Avena fatua* in the

same soil without a preceding addition of chitin showed that chitinase specific activity was

highest near root hairs (several days old rhizosphere), medium in bulk soil and lowest near

mature roots (one week old rhizosphere) [17]. In contrast, in rhizosphere with no litter addition,

we saw little change in chitinase expression over time. The slight differences in these results may

simply reflect the fact that the preceding study assayed enzyme activity and the current study

assessed gene expression.

We found that in unamended soil, expression of extracellular proteases was higher in the

rhizosphere compared to bulk soil, potentially resulting from input of carbon from root exudates

creating a higher demand for nitrogen as well as microbial competition over inorganic nitrogen

with the plant. Indeed, DeAngelis et al. showed that protease specific activity was significantly

higher in young rhizosphere compared to bulk soil, but activity was not significantly different

based on root age [17]. In litter-amended soil, expression of extracellular protease was highest

regardless of location at 3 days, suggesting that litter-added carbon overwhelms the effect of root

exudates at this early point in time. At 12 and 22 days there is no difference between litter-

amended and unamended soil in either rhizosphere or bulk soil, implying that the effect of litter

has waned. Changes in the quantity and quality of rhizodeposits can lead to changes in gene

expression over time in the rhizosphere. An increased demand for N is consistent with the higher

rates of gross nitrogen mineralization and ammonium consumption found in the rhizosphere of

mature compared to young *Avena sp.* [26]. The trend of increase in expression of extracellular

protease over time in our unamended rhizosphere soils suggests that as the rhizosphere ages it

becomes more nitrogen limited; thus, organic nitrogen would need to be mineralized to support

the microbial community.

Like many enzymatic reactions, macromolecular N depolymerization can be regulated by

substrate availability and end-product concentration [56]. In our study, litter amendment initially

led to high expression of extracellular proteases, suggesting that this expression was substrate

dependent [57,58] and linked carbon and nitrogen cycling [59]. However, protease activity in

grassland soil has also been shown to be affected by a multitude of parameters, such as end-

product concentration, aboveground net primary productivity [34] and soil aggregation [33]. We

observed an increase over time in expression in unamended samples although no nitrogen or

carbon sources were added. We hypothesize that this increase over time may be related to

depletion of end products (inorganic nitrogen), leading soil bacteria to invest more in expressing

depolymerization enzymes in order to access macromolecular nitrogen.

Spatiotemporal patterns of differential expression of genes were used to identify

functional guilds in which extracellular proteases were upregulated at a specific location or time.

These microbial guilds, which differed in taxonomic composition, underwent distinct

successional patterns that appear to be driven by niche conditions in the rhizosphere,

detritusphere, aging root and low response. Nuccio et al. [29], analyzing the same experimental

dataset, showed that microbial community composition (by 16S-rRNA) in each experimental

group changed very little compared to gene expression over the course of this experiment [29].

Therefore, the changing expression patterns over time that we observed are likely related to

changes in environmental conditions or cues, such as macromolecular N availability and

inorganic N availability, as opposed to wholesale shifts in community composition.

Carbon and nitrogen cycling by soil bacteria are linked through substrates that contain

both as well as by the need to maintain cellular C:N. The upregulation of carbon

depolymerization may promote increased access to nitrogen through expression of extracellular

proteases under the same conditions, leading to an overlap in guild membership. Such a

causative correlation has previously been observed in loam agricultural soil using protease

enzyme assays [59]. Therefore, we compared guild membership between guilds previously

defined by CAZy expression [29] and guilds defined here by hierarchical clustering of

extracellular protease expression. We found that rhizosphere guild membership was similar

between CAZy and protease guilds, while the detritusphere guilds had little overlap. All

members of the rhizosphere guild defined by extracellular proteases were also assigned to this

guild by CAZy expression, and all were Betaproteobacteria known to be associated with

rhizosphere soil [23,60,61]. In the sub-guild of "aging root" labeled "late synergist",

Burkholderiales bin 62_29, was assigned to the rhizosphere guild by CAZy expression. The

diverse detritusphere guild defined by extracellular protease expression included facultative

predatory bacteria of the order Myxococcales (Deltaproteobacteria). Predatory bacteria produce

proteases to bind to and digest prey bacteria [62] which may be attracted to litter. The small

amount of overlap between the protease-defined and CAZy-defined detritusphere guilds could

potentially be due to protein/amino acid supply of carbon backbones which reduces the need for

degradation of carbohydrates by CAZy genes in extracellular protein degraders. Guild overlap or

lack thereof contributes to further defining these multifunctional guilds in enzyme space.

Extracellular proteases have several structural types that differ in their active site. For

example, metalloproteases require zinc in their active site, therefore their activity may by limited

by zinc availability. There is currently very little information on when and where these different

types are active in soil, but it has been suggested that their activity might depend on soil

characteristics [32]. A recent publication by Nguyen et al. [14] showed that most of the

extracellular proteases in soil and aquatic environments are Metallo-, Serine- and

Cysteinepeptidases (MP, SP and CP respectively) in decreasing order of abundance. We found

these same clusters were dominant in our data, but expression patterns from our study revealed

dominance of SP over MP. Many bacterial phyla have dozens of copies of genes for secreted

peptidase and these genes are enriched especially in Actinobacteria and Proteobacteria, and more

so in soil/aquatic environments as opposed to animal associated bacteria. Specifically,

Betaproteobacteria are enriched in SP (mean 13 SP copies out of a mean of 25 extracellular

proteases per genome), with published genomes of isolates of Burkholderiales genera such as

Burkholderia, *Janthinobacterium* and *Massilia* containing more than 50 and up to 155 copies per

genome [14]. As Betaproteobacteria are enriched in the *Avena* rhizosphere [23], we might have

expected higher expression of SP in this environment compared to bulk soil. Surprisingly, we did

not observe that pattern; this could be explained by the high expression of genes from

Actinobacteria in all groups, as they share the same enrichment pattern with Betaproteobacteria

(SP>MP). Nguyen et al. also showed that while bacteria and archaea may carry genes coding for

multiple types of these enzymes within the same genome, the enzyme profile of proteolytic

activity was soil-specific [14,32]. Given that we see an expression profile of structural protease

groups different from that expected from genomes, this may hint that soil bacteria modify

expression of these groups in response to soil chemistry.

Conclusions

We show that extracellular proteases and chitinases are significantly upregulated by the presence

of roots and/or plant litter compared to other macromolecular N degrading enzymes. Patterns of

extracellular protease expression differ with taxonomy and can be used to define functional

guilds. Three guilds, "rhizosphere", "detritusphere" and "aging root" had membership overlap

with guilds previously defined by carbohydrate degrading gene expression based on analysis of a

CAZy genes dataset from the same experiment. This suggests that some of the members of these

guilds co-express carbohydrate- and protein-degradation enzymes. These guilds also exhibited a

pattern of temporal succession throughout plant maturation, where expression of protein

degrading enzyme genes in the rhizosphere could be triggered by the host plant through control

of exudation, or alternatively, may be a response to competition with the plant for labile nitrogen.

The taxonomic resolution provided by metatranscriptomes opens a window to the identification

of microbes that could enhance N availability to plants. The identity of these microbes has been

previously obscured in bulk analyses such as enzyme assays, rate measurements and even qPCR.

It is not far-fetched to imagine augmentation of soil or seeds with bacteria that specialize in

macromolecular organic N degradation to reduce the use of fertilizers. Ideally, augmentation

should use bacteria endemic to the specific plant rhizosphere. We show here that the identity of

such bacteria depends on plant life stage and soil characteristics such as organic material load.

References

1. Moreau D, Bardgett RD, Finlay RD, Jones DL, Philippot L. A plant perspective on nitrogen

cycling in the rhizosphere. Power S, editor. Funct Ecol. 2019;33:540–52.

2. Myrold, DD. Transformations of nitrogen. Principles and applications of soil microbiology.

Prentice Hall; 1999; 550.

3. Vitousek P, Howarth R. Nitrogen limitation on land and in the sea: How can it occur?.

Biogeochemistry. 1991.

4. Harte J, Kinzig AP. Mutualism and competition between plants and decomposers:

implications for nutrient allocation in ecosystems. Am Nat. 1993;141:829–46.

5. Paul EA, Clark FE. Soil microbiology and biochemistry. Academic Press. New York, USA.

1996.

6. Kinzig AP, Harte J. Selection of micro-organisms in a spatially explicit environment and

implications for plant access to nitrogen. Journal of Ecology. 1998. p. 841–53.

7. Dijkstra FA, Carrillo Y, Pendall E, Morgan JA. Rhizosphere priming: a nutrient perspective.

Front Microbiol. 2013;4:216.

8. Zhu B, Gutknecht JLM, Herman DJ, Keck DC, Firestone MK, Cheng W. Rhizosphere priming

effects on soil carbon and nitrogen mineralization. Soil Biol Biochem. 2014;76:183–92.

9. Meier IC, Finzi AC, Phillips RP. Root exudates increase N availability by stimulating

microbial turnover of fast-cycling N pools. Soil Biol Biochem. 2017;106:119–28.

10. Zhalnina K, Louie KB, Hao Z, Mansoori N, da Rocha UN, Shi S, et al. Dynamic root exudate

chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community

assembly. Nat Microbiol. 2018;3:470–80.

11. Chapin FS, Stuart Chapin F, Matson PA, Mooney HA. Principles of Terrestrial Ecosystem

Ecology. 2002.

12. Schimel JP, Bennett J. Nitrogen mineralization: challenges of a changing paradigm. Ecology.

2004. p. 591–602.

13. Jilling A, Keiluweit M, Contosta AR, Frey S, Schimel J, Schnecker J, et al. Minerals in the

rhizosphere: overlooked mediators of soil nitrogen availability to plants and microbes.

Biogeochemistry. 2018;139:103–22.

14. Nguyen TTH, Myrold DD, Mueller RS. Distributions of Extracellular Peptidases Across

Prokaryotic Genomes Reflect Phylogeny and Habitat. Front Microbiol. 2019;10:413.

15. Pett-Ridge J, Shi S, Estera-Molina K, Nuccio E, Yuan M, Rijkers R, et al. Rhizosphere

Carbon Turnover from Cradle to Grave: The Role of Microbe–Plant Interactions. Rhizosphere

Biology: Interactions Between Microbes and Plants. 2021. p. 51–73.

16. Zhu B, Cheng W. Rhizosphere priming effect increases the temperature sensitivity of soil

organic matter decomposition. Global Change Biology. 2011. p. 2172–83.

17. DeAngelis KM, Lindow SE, Firestone MK. Bacterial quorum sensing and nitrogen cycling in

rhizosphere soil. FEMS Microbiol Ecol. 2008;66:197–207.

18. Keiblinger KM, Schneider T, Roschitzki B, Schmid E, Eberl L, Hämmerle I, et al. Effects of

stoichiometry and temperature perturbations on beech leaf litter decomposition, enzyme

activities and protein expression. Biogeosciences. Copernicus GmbH; 2012;9:4537–51.

19. Tian K, Kong X, Yuan L, Lin H, He Z, Yao B, et al. Priming effect of litter mineralization:

the role of root exudate depends on its interactions with litter quality and soil condition. Plant

Soil. Springer Science and Business Media LLC; 2019;440:457–71.

20. Mukhopadhyay S, Joy VC. Influence of leaf litter types on microbial functions and nutrient

status of soil: Ecological suitability of forest trees for afforestation in tropical laterite wastelands.

Soil Biol Biochem. Elsevier; 2010;42:2306–15.

21. Srilakshmi A, Dakshayani R, Saigopal DVR, Narasimha G. Influence of forest litter on soil

enzyme activities. Ecol Environ Conserv. researchgate.net; 2012;18:105–12.

22. Jackson LE, Strauss RB, Firestone MK, Bartolome JW. Plant and soil nitrogen dynamics in

California annual grassland. Plant Soil. Springer; 1988;110:9–17.

23. Shi S, Nuccio E, Herman DJ, Rijkers R, Estera K, Li J, et al. Successional Trajectories of

Rhizosphere Bacterial Communities over Consecutive Seasons. MBio. 2015;6:e00746.

24. Brzostek ER, Finzi AC. Substrate supply, fine roots, and temperature control proteolytic

enzyme activity in temperate forest soils. Ecology. 2011;92:892–902.

25. Jaeger CH 3rd, Lindow SE, Miller W, Clark E, Firestone MK. Mapping of sugar and amino

acid availability in soil around roots with bacterial sensors of sucrose and tryptophan. Appl

Environ Microbiol. 1999;65:2685–90.

26. Herman DJ, Johnson KK, Jaeger CH, Schwartz E, Firestone MK. Root Influence on Nitrogen

Mineralization and Nitrification in Avena barbata Rhizosphere Soil. Soil Sci Soc Am J. Madison,

WI: Soil Science Society; 2006;70:1504–11.

27. Pett-Ridge J, Firestone MK. Using stable isotopes to explore root-microbe-mineral

interactions in soil. Rhizosphere. 2017;3:244–53.

28. Vieira S, Sikorski J, Dietz S, Herz K, Schrumpf M, Bruelheide H, et al. Drivers of the

composition of active rhizosphere bacterial communities in temperate grasslands. ISME J.

2020;14:463–75.

29. Nuccio EE, Starr E, Karaoz U, Brodie EL, Zhou J, Tringe SG, et al. Niche differentiation is

spatially and temporally regulated in the rhizosphere. ISME J. 2020;14;999–1014.

30. Yergeau É, Quiza L, Tremblay J. Microbial indicators are better predictors of wheat yield

and quality than N fertilization. FEMS Microbiology Ecology. 2020;96(2):fiz205.

31. Geddes BA, Paramasivan P, Joffrin A, Thompson AL, Christensen K, Jorrin B, et al.

Engineering transkingdom signalling in plants to control gene expression in rhizosphere bacteria.

Nat Commun. 2019;10:3430.

32. Nguyen T, Kleber M, Myrold DD. Contribution of different catalytic types of peptidases to

soil proteolytic activity. Soil Biol Biochem. 2019;138:107578.

33. Fukumasu J, Shaw LJ. The role of macro-aggregation in regulating enzymatic

depolymerization of soil organic nitrogen. Soil Biol Biochem. 2017;115:100–8.

34. Greenfield LM, Hill PW, Seaton FM, Paterson E, Baggs EM, Jones DL. Is soluble protein

mineralisation and protease activity in soil regulated by supply or demand? Soil Biol Biochem.

2020;150:108007.

35. Griffiths RI, Whiteley AS, O'Donnell AG, Bailey MJ. Rapid method for coextraction of

DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based

microbial community composition. Appl Environ Microbiol. 2000;66:5488–91.

36. Barnard RL, Osborne CA, Firestone MK. Responses of soil bacterial and fungal communities

to extreme desiccation and rewetting. ISME J. 2013;7:2229–41.

37. Huang Y, Niu B, Gao Y, Fu L, Li W. CD-HIT Suite: a web server for clustering and

comparing biological sequences. Bioinformatics. 2010;26:680–2.

38. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic

gene recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:119.

39. Rawlings ND, Barrett AJ, Finn R. Twenty years of the MEROPS database of proteolytic

enzymes, their substrates and inhibitors. Nucleic Acids Res. 2016;44:D343–50.

40. Bushnell B. BBTools software package. URL http://sourceforge net/projects/bbmap. 2014.

41. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biol. 2014;15:550.

42. Warnes GR, Bolker B, Bonebakker L, Gentleman R, Liaw WHA, Lumley T, et al. gplots:

Various R programming tools for plotting data. 2016; R package version 3.0;

https://cran.microsoft.com/snapshot/2016-03-30/web/packages/gplots/gplots.pdf

43. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer; 2009; ;10:978-0.

44. Starr EP, Shi S, Blazewicz SJ, Probst AJ, Herman DJ, Firestone MK, et al. Stable isotope

informed genome-resolved metagenomics reveals that Saccharibacteria utilize microbially-

processed plant-derived carbon. Microbiome. 2018;6:122.

45. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics. 2019; 36(6):1925–7.

46. Parks DH, Chuvochina M, Chaumeil P-A, Rinke C, Mussig AJ, Hugenholtz P. A complete

domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol. 2020;38:1079–86.

47. Lee MD. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics.

2019;35:4162–4.

48. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res.

2000;28:27–30.

49. Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better for

alignment and quantification of RNA sequencing reads. Nucleic Acids Res. 2019;47:e47.

50. Oksanen J, Kindt R, Legendre P, O'Hara B, Simpson GL, Solymos P, et al. The vegan

package. Community ecology package [http://r-forge r-project org/projects/vegan/]. 2008.

51. Olk DC. Organic Forms of Soil Nitrogen. Nitrogen in Agricultural Systems. 2008;57–100.

52. Cheng W, Parton WJ, Gonzalez-Meler MA, Phillips R, Asao S, McNickle GG, et al.

Synthesis and modeling perspectives of rhizosphere priming. New Phytol. 2014;201:31–44.

53. Hill PW, Jones DL. Plant-microbe competition: does injection of isotopes of C and N into the

rhizosphere effectively characterise plant use of soil N? New Phytol. 2019;221:796–806.

54. Roberts P, Stockdale R, Khalid M, Iqbal Z, Jones DL. Carbon-to-nitrogen ratio is a poor

predictor of low molecular weight organic nitrogen mineralization in soil. Soil Biology and

Biochemistry. 2009;41(8):1750–2.

55. Bowman SM, Free SJ. The structure and synthesis of the fungal cell wall. Bioessays.

2006;28:799–808.

56. Geisseler D, Horwath WR, Joergensen RG, Ludwig B. Pathways of nitrogen utilization by

soil microorganisms – A review. Soil Biology and Biochemistry. 2010. p. 2058–67. Available

from: http://dx.doi.org/10.1016/j.soilbio.2010.08.021

57. Allison SD, Vitousek PM. Responses of extracellular enzymes to simple and complex

nutrient inputs. Soil Biology and Biochemistry. 2005;37(5):937–44.

58. Noll L, Zhang S, Zheng Q, Hu Y, Wanek W. Wide-spread limitation of soil organic nitrogen

transformations by substrate availability and not by extracellular enzyme content. Soil Biol

Biochem. 2019;133:37–49.

59. Geisseler D, Horwath WR. Relationship between carbon and nitrogen availability and

extracellular enzyme activities in soil. Pedobiologia . 2009;53:87–98.

60. Berg G, Eberl L, Hartmann A. The rhizosphere as a reservoir for opportunistic human

pathogenic bacteria. Environmental Microbiology. 2005;7(11):1673–85.

61. DeAngelis KM, Brodie EL, DeSantis TZ, Andersen GL, Lindow SE, Firestone MK.

Selective progressive response of soil microbial community to wild oat roots. ISME J.

2009;3:168–78.

62. Pasternak Z, Pietrokovski S, Rotem O, Gophna U, Lurie-Weinberger MN, Jurkevitch E. By

their genes ye shall know them: genomic signatures of predatory bacteria. ISME J. 2013;7:756–

69.

Declarations

Ethics approval and consent to participate: not applicable

Consent for publication: not applicable

Competing interests: The authors declare that they have no competing interests

Availability of data and material: All data used in this publication, including raw reads and

genome collection, has been published previously. Metagenomes assembled MAGs can be found

in NCBI PRJNA517182, stable isotope probing MAGs in http://ggkbase.berkeley.edu/, single

amplified genomes in IMG under study name Mediterranean Grassland Soil Metagenome and

single amplified genomes in IMG, see sup. table S2 in Nuccio et al., 2020 for accession numbers

[29]. Raw reads can be found in JGI IMG. For JGI ID's (accession numbers) see supplemental

table S1 in Nuccio et al. 2020 [29].

Acknowledgements

The authors would like to thank Katerina Estera-Molina, Rachel Hestrin, Rachel Neurath and

Nameer Baker for their support and helpful feedback on the analyses. We thank Shenjing Shi for

contributing to the initial conceptualization of the experiment. Finally, we thank Evan Starr, Ulas

Karaoz, Eoin Brodie, Jizhong Zhou, Susannah G. Tringe, Rex R. Malmstrom, Tanja Woyke and

Jillian F. Banfield for assisting in the creation of the genome collection.

Funding

This research was supported by the U.S. Department of Energy Office of Science, Office of

Biological and Environmental Research Genomic Science program under Awards DE-

SC0020163 and DE-SC0016247 to M.K.F at UC Berkeley and awards SCW1589 and SCW1678

to J.P-R. at Lawrence Livermore National Laboratory. Sequencing was conducted as part of

Community Sequencing Awards 1487 to JPR and 1472 to MKF. Work conducted at LLNL was

contributed under the auspices of the US Department of Energy under Contract DE-AC52-

07NA27344.

Authors' contributions: ETS performed the data analysis, data interpretation and drafted the

manuscript. EEN, JPR and MKF conceived and designed the original rhizosphere-detritusphere

study, EEN helped to interpret the data and revise the manuscript. JPR and MKF both interpreted

the data and substantially revised the manuscript. All authors approved the submitted version and

agreed both to be personally accountable for the author's own contributions and to ensure that

questions related to the accuracy or integrity of any part of the work, even ones in which the

author was not personally involved, are appropriately investigated, resolved, and the resolution

documented in the literature.