Sex-biased admixture and assortative mating shape genetic variation and influence demographic inference in admixed Cabo Verdeans

Katharine L Korunes,[1] Giordano Bruno Soares-Souza,[2] Katherine Bobrek,[3] Hua Tang,[4] Isabel Inês Araújo,[5] Amy Goldberg,*[1,6] Sandra Beleza**[2,6]

1) Evolutionary Anthropology, Duke University, Durham, North Carolina, 27705, USA
2) Department of Genetics and Genome Biology, University of Leicester, Leicester, LE1 7RH, United Kingdom
3) Department of Anthropology, Emory University, Atlanta, Georgia, 30322, USA
4) Department of Genetics, Stanford University School of Medicine, Stanford, California, 94305 USA
5) Faculdade de Ciências e Tecnologia, Universidade de Cabo Verde (Uni-CV), Praia, Ilha de Santiago, Cabo Verde
6) These authors contributed equally to this work
*Correspondence: amy.goldberg@duke.edu
**Correspondence: sdsb1@leicester.ac.uk

ORCIDs
Katharine L Korunes: 0000-0002-2648-4707
Giordano Bruno Soares-Souza: 0000-0001-6590-5589
Katherine Bobrek: 0000-0003-0936-8779
Hua Tang: 0000-0002-0177-8864
Isabel Inês Araújo: 0000-0001-7766-1059
Amy Goldberg: 0000-0001-9306-1539
Sandra Beleza: 0000-0002-7004-3431

1 Abstract

Genetic data can provide insights into population history, but first we must understand the

patterns that complex histories leave in genomes. Here, we consider the admixed human

population of Cabo Verde to understand the patterns of genetic variation left by social and

demographic processes. First settled in the late 1400s, Cabo Verdeans are admixed descendants

of Portuguese colonizers and enslaved West African people. We consider Cabo Verde's well-

studied historical record alongside genome-wide SNP data from 563 individuals from 4 regions

within the archipelago. We use genetic ancestry to test for patterns of nonrandom mating and

sex-specific gene flow, and we examine the consequences of these processes for common

demographic inference methods and for genetic patterns. Notably, multiple population genetic

tools that assume random mating underestimate the timing of admixture, but incorporating

non-random mating produces estimates more consistent with historical records. We consider

how admixture interrupts common summaries of genomic variation such as runs-of-

homozygosity (ROH). While summaries of ROH may be difficult to interpret in admixed

populations, differentiating ROH by length class shows that ROH reflect historical differences

between the islands in their contributions from the source populations and post-admixture

population dynamics. Finally, we find higher African ancestry on the X chromosome than on

the autosomes, consistent with an excess of European males and African females contributing to

the gene pool. Considering these genomic insights into population history in the context of

Cabo Verde's historical record, we can identify how assumptions in genetic models impact

inference of population history more broadly.

Introduction

Patterns of genetic variation are often used to infer human population histories, such as fluctuations in population size, the timing of gene flow, and changes in population structure[1–7]. These studies have been informative about ancient and modern populations that may lack historical records[8–12]. However, by necessity, population-genetic models make a variety of simplifying assumptions that often neglect the demographic and social dynamics of human populations. For example, many classical models assume random mating, but within empirical populations mating is often nonrandom, known as assortative mating. Positive assortative mating in human populations can keep haplotype blocks together, affecting evolutionary processes and our ability to infer population history[13–15]. Further, male and female demographic processes can differ and vary over time. Comparisons between the X chromosome and autosomes have shown that population sizes and migration rates differ between males and females of many human populations[16–20].

The challenges of inferring human demographic history are particularly apparent on short timescales of tens of generations, where changes in allele frequencies may be difficult to observe and biased by dynamics not captured by classic models. Admixed populations provide an opportunity to examine evolutionary processes on short timescales using admixture linkage-disequilibrium (LD) structure rather than potentially small changes in allele frequencies. Understanding the patterns generating the distribution of ancestry within an admixed

population is also important as genomic data has shown how widespread admixture is

throughout human evolution[8,11,21–25].

In admixed populations, nonrandom mating can be driven by factors that correlate with genetic

ancestry. For example, spouse pairs within multiple Latino populations are correlated in their

genomic ancestry[13,15]. Recent work has begun to assess the implications of these patterns,

including effects on common demographic inference strategies and on genetic/phenotypic

variation[7,14,24]. One genetic consequence of assortative mating is an increase in long runs of

homozygosity (ROH), which arise when long tracts of homozygous genotypes occur due to

inheritance of identical by descent (IBD) haplotypes from recent common ancestors. Across

modern human populations, ROH are common and typically range in length from kilobases to

megabases[26–29]. In the field of population genetics, we are just beginning to learn about ROH

dynamics and expectations in admixed populations, which do not necessarily fit expectations

built from homogenous populations[29,30]. For example, we know that distinct length classes of

ROH reflect demographic factors from different timescales, but as we explore here, it is unclear

what these distributions of ROH mean in populations that have undergone major demographic

transitions such as recent admixture. Patterns of genetic variation can also be shaped by

differences in male and female demographic processes. In many human populations,

population sizes and migration rates differ between males and females, leading to differences in

ancestry patterns on the X chromosome vs the autosomes[16–20].

Here, we use the admixed population of Cabo Verde as a case study to understand the patterns

of genetic variation left by multifaceted social and demographic processes in humans. Admixed

populations of African ancestry, such as the population of Cabo Verde, are often excluded in the

context of medical genetics and human evolutionary genetics, despite their importance and

widespread global presence[31–33]. We consider patterns of genetic variation alongside historical

records that document many aspects of Cabo Verdean history, including settlement patterns,

timing, and sociocultural dynamics. Specifically, we use distributions of genetic ancestry to test

for patterns of nonrandom mating and sex-specific gene flow. We examine the consequences of

these processes for genetic variation, such as patterns of homozygosity, and for common

demographic inference methods. In turn, by elucidating the patterns of genetic variation social

processes leave in human genomes for a population with a well-studied historical and

ethnographic record, we can better use population-genomic methods to explore the history of

populations without extensive records.

Cabo Verdean history

The recently admixed human population of Cabo Verde presents a particularly relevant

opportunity to examine how sex-biased admixture and nonrandom mating shape genetic

patterns and influence demographic inference. Cabo Verde is an archipelago off the coast of

Senegal, inhabited today by admixed descendants of Portuguese colonizers and enslaved West

African people who settled the unpopulated islands beginning in the mid-1400s[34–37]. Since the

archipelago was unoccupied prior to the 1460s, we have historical knowledge of the start of

admixture and the contributing source populations, with written records clearly documenting

arrival times by island. Additionally, the island geography imposes population structure within

Cabo Verde, and historical records document differences in population sizes, mating patterns,

and social customs by island population (see Methods for historical data sources).

The settlement of the islands was influenced by island geography and ecology, and is often

divided into three different temporal stages, which are associated with changes in the

economy[38]. The initial settlement stage (beginning in the 1460s) corresponds to the peopling of

the Southern island Santiago, followed by the nearby island of Fogo 20 years later (Fig 1).

During this time, the economy was centered around large cotton plantations and expanding

trade with the Senegambian coast. This trade included the exportation of great numbers of

enslaved Africans to Cabo Verde, the majority of which became part of the commercial trade to

the Americas[39,40]. In the 17th century, following the decline of the slavery-based economy of

Santiago and Fogo, many free farmers seeking better farming conditions migrated to the

northwestern islands of Santo Antão and São Nicolau. These migrations comprised the second

settlement stage. In contrast to Santiago and Fogo, these northwestern islands did not become

populous economic centers[38]. During the third and final settlement stage, São Vicente in the

northwestern group of islands was colonized by settlers from the neighboring islands in the

mid 19th century. With the opening of Mindelo's harbor at that time, São Vicente prospered

from the advent of commercial Atlantic shipping, becoming the second most important island

of the archipelago after Santiago in terms of population size. In the 19th century, the eastern

islands Sal, Boa Vista, and Maio were also open to English and North American ships, but these

islands never became densely populated[38,40].

Previous investigations of genetic ancestry in Cabo Verde[34,35,37] have shown extensive admixture

in the archipelago and differences in the mean African genetic ancestry across islands, which

may reflect the differences in the settlement history and suggest restricted gene flow between

islands. Using genome-wide SNP data from 563 individuals spanning the three settlement

stages of Cabo Verde, we integrate several population genomic approaches to infer the sex-

specific demographic history of the last ~20 generations since founding, and consider the

potential biases of nonrandom mating on common inference methods. Overall, we provide

insights into the population history of Cabo Verde and demonstrate how admixed populations

can provide powerful test cases for understanding evolutionary processes on short timescales.

Methods

Study population and ancestry reference panels

We used genotype data from Beleza et al. (2013)[35], which included 563 Cabo Verdeans from the

following regions of the archipelago: Santiago (n = 172), Fogo (n = 129), Boa Vista (n = 26), and

the three northwestern islands in aggregate (Northwest Cluster; n = 236) (Figure 1). In

accordance with historical records and previous work showing that Iberian and Senegambian

populations are suitable proxies for the ancestry sources of Cabo Verde[37], we leveraged data

from the 1000 Genomes Project to estimate admixture proportions and call local ancestry in the

Cabo Verde individuals, as described below. We merged the Cabo Verde genotypes with

125 genotypes from 107 GWD (Gambian in Western Division - Mandinka) and 107 IBS (Iberian

126 Population in Spain) samples called from high-coverage resequencing data released through the

127 International Genome Sample Resource[41,42]. Our final merged dataset consisted of 884,656

128 autosomal and 20,967 X chromosome SNPs shared between the Cabo Verde samples and the

129 reference samples, with average missingness rates of 0.0017 by SNP for autosomes, and 0.0024

130 for the X chromosome.

Historical records

132 Throughout our analyses, we draw comparisons between genetic results and historical records.

133 We used primary historical documents, mainly historical letters to the Portuguese Crown,

134 documenting demographic characteristics of the islands across time. Our analyses of the

135 historical data took into consideration the generalized interpretations of historian scholars such

136 as Correia e Silva (2001; 2002)[38,43], Cabral (2001; 2012)[44,45] and Baleno (2001)[39]. These sources

137 document the dates of the stages of settlement mentioned above, along with associated with

138 changes in the economy. We refer to the interpretation of genealogical data by Cabral (2012)[45]

139 and Barbosa (1997)[46] for evidence of the structure of the Fogo's society and for evidence of first-

140 cousin marriages within Fogo[46].

Characterization of ancestry

142 To produce estimates of admixture proportions, we first performed unsupervised clustering of

143 the samples using ADMIXTURE[47]. ADMIXTURE was run separately for the autosomal and X

144 chromosome datasets after pruning based on linkage disequilibrium (LD) using the indep-

pairwise option of PLINK v1.9 with a 50-SNP sliding window incremented by 10 SNPs, and an

LD threshold of r2 = 0.5[48]. This pruned dataset included 514,551 autosomal and 12,706 X

chromosome SNPs. We estimated individual ancestries by averaging over ten independent

unsupervised ADMIXTURE runs using K = 2, given the historical and genetic support for two

source populations[35,37]. Using the LD-pruned dataset, we also visualized the data using PLINK's

principal component analysis (--pca) function, confirming that the samples cluster

approximately based on their geographic memberships, and that West African vs European

ancestry clearly separates on the first principal component (Supp Fig 1). We phased all samples

using SHAPEIT2[49]. After running SHAPEIT -check to exclude sites not contained within the

reference map, we ran SHAPEIT to yield phased genotypes at 881,279 autosomal SNPs and

20,793 X chromosome SNPs. We then called local ancestry with RFMix v1.5.4[50] PopPhased

under a two-way admixture model using the West African and European reference genotypes

described above. These local ancestry calls are publicly available via Zenodo[51]. Ancestry

proportions estimated with ADMIXTURE and RFMix here are highly correlated with results

from Beleza et al. (2013)[35], using the same genotype data but with different software (*frappe*[52]

and SABER[53]) and an older reference dataset (Supp Fig 2).

Inference of admixture timing

We applied three distinct strategies for estimating the timing of the onset of admixture in Cabo

Verde: ALDER[4], MultiWaver 2.0[5], and a method based on patterns of linkage disequilibrium

between local ancestry tracts [7]. We first converted the genotypes to EIGENSTRAT format using

convertf[54,55]. We then used ALDER to date admixture timing on each island using default

parameters with mindis = 0.005 and source populations GWD and IBS. We ran MultiWaver

using the ancestry tracts inferred by RFMix, default parameters, and 100 bootstraps. Finally, we

followed the pipeline of Zaitlen et al. (2017)[7] and its accompanying scripts to use the RFMix

local ancestry calls to measure local ancestry disequilibrium (LAD) decay in 10 Mb windows,

overlapping by 1 Mb. Specifically, we started with the first SNP on each autosome, used the 10

Mb window end point to identify the SNP closest to the inside of this boundary, and then used

local ancestry calls at these positions to determine LAD. We repeated this process along each

chromosome to obtain LAD in 279 autosomal windows. Using possible values of admixture

generations ranging from 5-25, we determined the best fit using island-specific mean LAD

decay over the 279 autosomal windows, assortative mating parameters estimated with

ANCESTOR (see below), a starting autosomal admixture proportion of 0.65, and either no

migration or with migration (migration rate = 0.01).

Tests for assortative mating

We tested for assortative mating over the last generation using ANCESTOR[15,56]. ANCESTOR

uses phased local ancestry tracts in the current generation to estimate the ancestral proportions

of the two parents of each individual. We used the inferred parental ancestries to test for

assortative mating seen as a positive correlation in ancestry between inferred mating pairs, and

we used the strength of the observed positive correlation (Pearson's R for each population,

using the median value from all chromosomes) as the assortative mating parameter in our

application of the LAD-based method for inferring admixture timing. We also examined

evidence of nonrandom mating by estimating genetic relatedness within Cabo Verde using both

IBD (see below) and kinship coefficients[57].

Identification of IBD tracts and ROH

Following the analysis pipeline of S. R. Browning et al. (2018)[2], we inferred segments of IBD

using the haplotype-based IBD detection method Refined IBD[58], and we estimated ancestry-

specific population size using IBDNe[1]. After running Refined IBD, we used its gap-filling utility

to remove gaps between segments less than 0.6 cM that had at most one discordant

homozygote. We then filtered out IBD segments smaller than 5 cM, as short segments below

this threshold are difficult to accurately detect[59]. To visualize IBD sharing in Fig 1 and Supp Fig

3, the sum of IBD shared in each pairwise comparison of individuals was plotted by using

Cytoscape 3.8 to group nodes by island, position nodes within each island according to a

prefuse force-directed algorithm, and scale the color of edges based on log-transformed total

IBD length[60].

We called ROH using GARLIC v.1.1.6[61], which implements the ROH calling pipeline of

Pemberton et al. (2012)[29]. We performed this analysis separately for the autosomes and X

chromosome. Based on the pipeline described in Szpiech et al. (2017), we used a single constant

genotyping error rate of 0.001, we allowed GARLIC to automatically choose a window size for

each population (--auto-winsize), and we used the resample flag to mitigate biases in allele

frequency estimates caused by differing sample sizes. This resulted in GARLIC selecting a

window size of 50 SNPs in each of the Cabo Verde regions and in GWD, and a window size of

40 for IBS. Using a three-component Gaussian mixture model, GARLIC classified ROH into

three length groups: small/class A, medium/class B, and long/class C ROH. Across all

populations, Class A/B and class B/C size boundaries were inferred as approximately 300 kb

and 1 Mb, respectively (see Supp Table 1 for population-specific parameters, including LOD

cutoffs and size boundaries). Using only the females, we then classified ROH for the X

chromosome.

Estimation of sex bias

To examine sex bias, we applied a mechanistic model of sex-biased admixture[3] to infer

admixture parameters under a model of constant migration. We performed this analysis for all

563 Cabo Verdean individuals pooled together, and then repeated the analysis for each island

independently. We provided the model with the following parameters: grid increment size

(0.02), the observed autosomal and X admixture proportions (based on ADMIXTURE estimates

described above), the proportion of females in the sample, and the percentile cutoff for

parameters to keep (0.1%).

Results

Patterns of shared ancestry reflect the colonization history of the islands

Using genome-wide SNP data from 563 individuals, we examined four island regions of Cabo

Verde (Fig 1), which differ in their settlement histories. The partitioning of the considered island

regions was also supported by genetic patterns. Namely, these regions showed quantitatively

distinct distributions of IBD tracts, as we explore below, and were supported by clustering

patterns in PCA (Supp Fig 1), in which the three northwestern islands grouped together (and

with the eastern island Boa Vista) despite the different settlement time of São Vicente Island.

Global ancestry proportions reflect extensive European and West African admixture and concur

with previous results[35] (Supp Fig 2), with the median West African autosomal ancestry in Cabo

Verdeans varying by island from 50% in Fogo, 56% in the northwestern islands in aggregate

(the Northwest Cluster), 63% in Boa Vista, and 75% in Santiago. These inter-island differences

alongside historical knowledge provide a comparative dataset for understanding how ancestry

patterns can be used to infer demographic processes.

The process of admixture can influence measures such as IBD and ROH that are often used to

inform inference of population history, and theoretical expectations for these measures are less

clear for admixed populations compared to homogeneous populations. Thus, we use multiple

methods to examine relatedness in Cabo Verde, and we use this case study to underscore the

need for further empirical and theoretical work to understand the dynamics of IBD and ROH in

admixed populations. Using IBD to examine relatedness between and within regions of Cabo

Verde (Fig 1), we found that patterns of shared ancestry reflect the successive settlement history

of the islands. Patterns of IBD within and between populations provide opportunities to

examine common ancestry based on the number and sizes of segments of IBD. Notably,

Santiago has the lowest mean number and total length of IBD segments between individuals

(Supp Fig 4-5). In contrast, we observe the highest levels of IBD within and between the

Northwest Cluster and Boa Vista.

Focusing on pairs of individuals with relatively high levels of IBD, we observe that recent

relatives both within and across islands are common (Fig 1; see Supp Fig 3 for lower levels of

IBD and within-island networks). Notably, relatedness parameters estimated under simple

models of structure likely do not hold in many human populations, such as recently admixed

populations. Kinship estimated under a framework[57] specifically designed for arbitrary

population structures (Supp Fig 6A) is consistent with both the overall trends in IBD tracts

(Supp Fig 4-5), and supports the presence of a subset of very closely related individuals within

the data. We observed that the ordering of the global ancestry proportions according to a

kinship matrix ordered using seriation, which places low kinship values away from the

diagonal[57] (Supp Fig 6), shows that Cabo Verdeans with greater European ancestry tend to

share greater kinship compared to individuals with greater West African ancestry.

The lower levels of IBD (Fig 1, Supp Fig 3-5) and lower kinship estimates (Supp Fig 6) observed

in Santiago are consistent with expectations based on the island's history, as Santiago was the

first Cabo Verdean island to be founded and has the largest population size. Santiago is also

likely to have relatively high genetic diversity as a result of having the highest proportion of

West African ancestry, based on the expectation that kinship in human populations generally

reflects serial bottlenecks due to dispersal from Africa[62,63]. In contrast, the high levels of IBD

within and between the Northwest Cluster and Boa Vista reflect well-documented serial

founding and migrations during the settlement of the islands. In general, kinship estimates also

reflect the colonization history of the islands, with Cabo Verdeans of greater European ancestry

sharing greater kinship compared to individuals with greater West African ancestry (Supp Fig

6). Exceptions to this trend were most noticeable in individuals from Fogo and the Northwest

Cluster, where it was most apparent that some observed relatedness patterns do not result in

smooth ordering of global ancestry proportions (Supp Fig 6B). This observation led us to

hypothesize that other sociocultural processes beyond proportions of ancestry, such as

nonrandom mating, may drive patterns of relatedness in Cabo Verde.

Nonrandom mating shapes patterns of ancestry and influences

demographic inference

To test for ancestry assortative mating within Cabo Verde, we examined whether the genomic

ancestries of individuals in mating pairs correlate with each other. To this end, we applied

ANCESTOR to computationally infer the parental ancestry proportions that likely preceded the

ancestry haplotypes we observe today. We found that the inferred ancestries of mating pairs in

the previous generation are significantly positively correlated (Fig 2).

We next considered how the observed genetic evidence of assortative mating can be leveraged

in the context of inferring population history. To estimate the timing of the onset of admixture,

we first applied two strategies, ALDER[4] and MultiWaver 2.0[5], that are not intended to account

for assortative mating. ALDER uses the extent of LD decay among neighboring loci to infer

mixture proportions and dates. In contrast, MultiWaver uses the length distribution of ancestral

tracks. Assuming a generation time of 25 years, decay rates estimated with ALDER suggest

admixture timing in the mid to late 1700s (Fig 3). MultiWaver chose a multi-wave admixture

model for Santiago (Fig 3A) and the Northwest Cluster (Fig 3C), but selected an isolation model

for Fogo (Fig 3B) and Boa Vista (Fig 3D). To incorporate nonrandom mating, we applied the

method of Zaitlen et al. (2017)[7], which describes the decay of local ancestry disequilibrium

(LAD) as a function of assortative mating strength, migration rate, recombination rate, and the

number of generations since admixture began. Results from the LAD-based method described

in Zaitlen et al. (2017)[7] estimated more generations of admixture than ALDER or MultiWaver,

with older estimates of admixture timing under a model including both assortative mating and

constant migration.

The estimates of admixture timing from ALDER and MultiWaver are noticeably more recent

than the known founding of Cabo Verde in late 1400s. Results from the LAD-based method

described in Zaitlen et al. (2017)[7] align more closely with historical records (Fig 3), particularly

when we allowed for both assortative mating and constant migration. These results suggest that

explicitly accounting for nonrandom mating improves estimates of admixture timing, while the

assumption of random mating leads to underestimation of the age of admixture.

Runs-of-homozygosity (ROH) reflect the contributions of the source populations and patterns of nonrandom mating.

We identified ROH using the method of Pemberton et al. (2012) as implemented in the software GARLIC[29,61]. This method classifies ROH into three categories (short, medium, and long) based on the length distributions in each population. ROH arise when IBD haplotypes are inherited from a common ancestor. Thus, distinct length classes reflect inheritance from ancestors at different timescales in the population's history. We examined the distributions of ROH within Cabo Verde, partitioning the distributions into shorter ROH (short and medium, as called with GARLIC) and long ROH (Fig 4A-B). While ROH in homogeneous populations are often modeled as three length classes, we observe that pre- and post-admixture dynamics in Cabo Verde are best captured by focusing on long ROH vs all shorter ROH lengths. Long ROH primarily reflects post-admixture dynamics such as recent parental relatedness in the admixed population, whereas shorter ROH reflects primarily pre-admixture homozygosity of older haplotypes and background relatedness from the source populations.

Since methods for detecting and interpreting ROH have been developed primarily in homogeneous populations[29,64], we consider ROH in Cabo Verde as an example of ROH distributions in admixed populations where we can explore results in the context of ancestry and population history. For example, to test for genetic evidence that population bottlenecks contributed to observed patterns, we estimated ancestry-specific population sizes (Supp Fig 7). Ancestry-specific population sizes in Cabo Verde suggest bottlenecks of both source

populations, followed by population expansion within the past 10 generations[1,2]. If the length of

ROH in an individual's genome is driven by distance from Africa, as expected based on

previous work[28,29], we predict the total length of ROH per genome to negatively correlate with

West African admixture proportion. Consistent with this expectation, we observe a negative

correlation between the total ROH and West African ancestry proportions in Santiago, Fogo,

and the Northwest Cluster (Supp Fig 8A). Breaking total ROH into shorter and long ROH, we

see that this negative correlation is driven primarily by shorter ROH (Fig 4; Supp Fig 8B-C).

Despite recent colonization bottlenecks, we observed that some admixed individuals in Cabo

Verde present lower levels of ROH than either of the source population proxies. Comparing the

Cabo Verdean islands to each other, we observed the lowest amounts of all length classes of

ROH in Santiago (pairwise Mann-Whitney U tests $p < 1 \times 10^{-8}$). The low levels of all length

classes of ROH in Santiago are consistent with several historical attributes of the islands,

including serial founding beginning with the settlement of Santiago. Serial founding resulting

in bottlenecks is expected to increase shorter ROH by reducing the number of ancestral

lineages. Because shorter ROH reflects background relatedness from the source populations,

shorter ROH are expected to be more sensitive to the contributions of the source populations.

Our classification of ROH into shorter and long ROH (Fig 4) revealed that Santiago has

significantly lower levels of shorter ROH compared to even the West African reference

population (Mann-Whitney U $p = 5.87 \times 10^{-7}$), while long ROH are not depleted (Mann-Whitney

U $p = 0.841$). This observation that Santiago has significantly lower levels of shorter ROH

compared to even the West African reference is perhaps surprising given the canonical view of

African source populations as generically having the lowest ROH in worldwide samples. This is

consistent with the idea that shorter ROH are contributed to the population from older shared

ancestors in the source populations, and these tracts are often interrupted with local ancestry

from another source population upon admixture. In contrast, long ROH are likely driven by

post-admixture dynamics such as small population size and mating preferences, and long ROH

tracts may span multiple local ancestries.

While ROH patterns in Cabo Verde are consistent with several historical observations, we

emphasize that caution is warranted in applying methods and expectations built using

homogeneous populations. We also note that, though we used a model-based ROH detection

approach that included steps to mitigate genotyping errors and biases in allele frequency

estimates, sensitivity and specificity of ROH detection is generally lower for shorter ROH. These

sources of uncertainty necessitate caution in inferring population history from ROH

distributions. However, the observed differences in ROH among the islands suggest that ROH

are sensitive to population genetic processes even on the short timescale since Cabo Verde's

settlement. Together, these observations and caveats underscore the need for future work

testing the effects of admixture on ROH and on ROH detection, as we explore in the Discussion.

Ancestry patterns on the X chromosome vs the autosomes reflect sex-biased demographic processes

On all islands, autosomal versus X chromosome ancestry patterns suggest that male and female contributions differ significantly by source population (Fig 5A; Wilcoxon Signed-Rank Test $p < 1 \times 10^{-8}$ for Santiago, Fogo, and the Northwest Cluster; $p = 0.0013$ for Boa Vista). To quantitatively explore these differences in male vs female demographic history, we used a model-based approach to estimate differences between male and female contributions[3]. A model of constant admixture is supported over a model of instantaneous admixture by historical records and by our comparison of admixture timing estimates (above). Using the method of Goldberg and Rosenberg for a model of constant admixture, we compared observed and expected X-chromosomal and autosomal ancestry over a grid of possible parameter values. Following the implementation from Goldberg et al. (2017)[65], Fig 5B presents results from the 0.1% of parameter sets closest to observed data based on a Euclidean distance between observed population mean ancestries on the X and autosomes and expected ancestries under the model (Supp Fig 9 shows the same data represented by plotting the male contributions). These estimated sex-specific contributions from West African and European source populations under a model of constant admixture suggest greater contributions from West African women and European men.

Given our observation that source populations make distinct contributions to ROH in Cabo Verde, we next investigate differences in autosomal vs X chromosome ROH content. Differences

in autosomal vs X chromosome ROH may arise in part due to the source populations

contributing in a sex-biased manner. Notably, this effect would be seen most in shorter ROH,

since shorter ROH reflect the homozygosity of older haplotypes and background relatedness

from the source populations. However, it is challenging to disentangle processes shaping X vs

autosomal ROH, due to the smaller effective population size of the X chromosome. Our results

suggest that sex-biased admixture processes in Cabo Verde are reflected in ROH, with lower

levels of shorter ROH on the X chromosome than on autosomes (Fig 6A). European individuals

have higher levels of ROH than West African individuals, and the higher contributions of

African X chromosomes (vs European X chromosomes) may drive the lower levels of short

ROH in Cabo Verde on the X chromosome vs the autosomes (Fig 6A). Long ROH reflects

different dynamics, potentially including the reduced post-admixture population size and other

sex-specific processes.

Discussion

In this study, we leveraged patterns of genetic variation in Cabo Verde to infer the demographic

history of the past ~20 generations. We found that distinct genetic patterns of four island

regions within the archipelago reflect the colonization history of the islands, including island-

specific settlement timing, admixture dynamics, mating patterns, and sex-biased demography.

Together these results demonstrate how patterns of ancestry and genetic variation are shaped

by social and demographic forces on short timescales. By better understanding how complex

population histories generate genetic variation, we can improve interpretation of inference from

populations without historical records.

Patterns of genetic variation and admixture in Cabo Verde reflect colonization history and subsequent sociocultural dynamics

The observed island-specific genetic patterns reflect the complex ecological and social factors

that shaped the settlement of Cabo Verde. Historical evidence suggests that the islands were

founded in a stepping-stone pattern, with settlement stages that were accompanied by

important economic and sociocultural shifts[38]. Early written records of the population

(described below) suggest that admixture began early during the settlement of Cabo Verde,

despite the highly racially stratified, slavery-based system that characterized the first settlement

stage[44]. The second and third stages were carried out by mostly already-admixed individuals

who had become a significant group within Cabo Verdean society and who migrated from the

southern to the northern islands[43]. We found that the staggered settlement history and the

island-specific population dynamics shaped patterns of ancestry and genetic variation within

Cabo Verde.

Admixture in Cabo Verde is consistent with a model of continuous gene flow from Europe and

West Africa, with European males and African females contributing predominantly to the Cabo

Verdean gene pool, as seen in other admixed populations that result from the trans-Atlantic

slave trade[66,67]. PCA (Supp Fig 1), IBD (Fig 1, Supp Fig 3-5), and kinship (Supp Fig 6) are

consistent with genetic drift occurring during the archipelago's settlement history of

consecutive founder effects and subsequent relative isolation of the islands. Specifically,

individuals are genetically more related within an island than among islands, and considerably

more individuals from the northern islands and Fogo are connected to each other by high IBD

pairwise connections (Supp Fig 4-5). We observed differences in IBD that reflect the

contributions of the source populations, such as Santiago having both the highest West African

ancestry proportions and the lowest overall levels of IBD sharing. The higher African genomic

ancestry on Santiago is consistent with previous results[34,35], and indicates differences in

settlement patterns despite the fact that colonization of the other islands involved migrations

from Santiago and was based on the same slavery-based system.

Using a mechanistic model of sex-biased admixture[3], we used differences in ancestry on the X

chromosome vs the autosomes to examine sex-biased migration in Cabo Verde. We found that

admixture occurred primarily through the mating of European males and African females,

consistent with historical work that documents sex bias during the settlement of Cabo Verde.

First records of the demographic distribution in Cabo Verde are found in a letter by a judicial

official Pero de Guimarães to the Portuguese King in 1513, which stated the presence of 118

European individuals in Santiago, of which only four were (single) women[68]. The marginal

presence of European females throughout the founding of Cabo Verde contributed to the

extensive genetic admixture currently found in the archipelago. The distribution of ROH in

autosomes versus X chromosome is influenced by initial male versus female contributions to the

admixed population's gene pool. Throughout the islands, the higher contributions of African X

chromosomes (vs European X chromosomes) may drive the lower levels of short ROH in Cabo

Verde on the X chromosome vs the autosomes (Fig 6A). These results underscore the long-

lasting genetic impacts of the trans-Atlantic slave trade, as has been recently shown in admixed

populations across the Americas[67].

Despite only a few hundred years of unique population histories among the islands, we also

found that patterns of genetic variation reflect island-specific patterns in post-admixture

dynamics. Observed patterns of IBD, kinship, and long ROH are all consistent with overall high

levels of relatedness within Cabo Verde, particularly in Fogo and the northwestern islands.

These patterns of relatedness may be shaped by both founder effects and by post-admixture

dynamics such as nonrandom mating keeping similar haplotypes together rather than

distributing them randomly throughout mating pairs in the population. Indeed, we inferred a

positive correlation between mating pairs of the previous generation, suggesting assortative

mating on the short timescale since the founding of the islands. Mating patterns and sex-biased

admixture are distinct—but interacting—processes in the history of Cabo Verde. Sex-biased

migration determined the gene pool within Cabo Verde, while patterns of nonrandom mating

within the islands shaped subsequent generations of admixture. These mating patterns were

tied to social and economic structure. European enslavers sexually coerced enslaved African

women during slavery. After the end of slavery, racially unbalanced control of landownership

and higher social status of Europeans resulted in continued socioeconomic pressures for African

women to have children with European men. Such unions often resulted in improved

socioeconomic status for both the women and their children. For example, the first admixed

individual to assume an important position within the administration of Santiago was recorded

in 1546[69]. Over generations, Cabo Verde became a population of extensively admixed ancestry,

with mating patterns shaped by the interconnected effects of social position and skin color. This

type of nonrandom mating may have been more prominent in the second half of the 17[th]

century, due to the large exodus of the white population following the decay of the slave trade.

With this population shift, historical sources say that the socioeconomic elite of the islands

became predominately the "white children of the islands"[44], leading to further racial

segregation. This historical evidence of nonrandom mating patterns—together with our genetic

observations—demonstrates how quickly changes in social structure can impact population

homozygosity and differentiation.

The island of Fogo stands out as having unique social and historical processes, even though it

was settled shortly after Santiago. Fogo's society was, since its origins, a conservative rural

society whose main economic activity was to produce goods to trade in the African coast. The

socioeconomic elite was patriarchal and aristocratic[45], and composed primarily of related

families, which promoted first-cousin marriages[46]. The sustenance of this class depended on

land ownership, which allowed the slavery-based system to be perpetuated longer in Fogo than

in Santiago[38,45]. The unique attributes of Fogo are reflected in population genetic patterns. For

example, we found that African ancestry proportion align with kinship patterns overall, but

exceptions were most obvious in individuals from Fogo and the Northwest Cluster, where some

observed relatedness patterns are not as clearly explained by global ancestry proportions. In our

estimation of admixture timing using local ancestry disequilibrium, we observed that Fogo has

greater levels of local ancestry disequilibrium than the other islands. Higher local ancestry

disequilibrium may be driven by assortative mating. Additionally, it may be that Fogo has

experienced stronger founder effects, which would decrease the number of ancestral lineages.

Indeed, the higher levels of shorter ROH within Fogo (Fig 4) are consistent with founder effects

increasing background relatedness and thus increasing shorter ROH, though we emphasize the

need for further work to understand the dynamics of ROH in admixed populations.

Patterns of genetic variation and admixture in Cabo Verde influence demographic inference

The admixture dynamics, founder effects, and mating patterns within Cabo Verde shape

summaries of genetic variation that are often used to inform demographic inference, such as

local ancestry disequilibrium, IBD, and ROH. To investigate the effects of these population

dynamics on demographic inference, we estimated admixture timing using multiple population

genetic tools. Estimates of admixture timing from genetic data were most concordant with

historical records when using inference based on local ancestry disequilibrium[7], allowing for

assortative mating and migration. Under this method, we tested a range of assortative mating

strengths (0 vs values inferred using ANCESTOR) and migration rates (0 vs 1% each

generation). The migration levels are used to demonstrate the trends of timing estimates using

constant migration and assortative mating, rather than to obtain a specific estimate of the

migration parameters. Under random mating, haplotypes are distributed randomly in mating

pairs in a population, allowing recombination to shuffle haplotypes over generations. In

contrast, assortative mating can keep ancestral haplotypes more intact, leading to

underestimation of the number of generations of admixture when estimating admixture timing.

Our evidence of assortative mating in Cabo Verde underscores the importance of accounting for

nonrandom mating in understanding admixture and inferring demographic history. Assortative

mating has been documented with respect to genetic ancestry, socioeconomic factors, and

phenotypic characteristics[7,13–15,70], but remains an under-recognized force in human population

genetics and demographic inference.

Notably, assortative mating alone was not enough to obtain time of admixture estimates

concordant with historical records. Admixture timing estimates were most concordant with the

historical records when allowing both assortative mating and constant admixture. An

admixture scenario that allows multi-wave or constant admixture model is compatible with

historical work suggesting constant migration from both Europe and West African throughout

the settlement of the islands. Importantly, the authors of ALDER and MultiWaver point out that

methods for inferring admixture timing have varying sensitivities to different admixture

scenarios[4,5]. MultiWaver may not be able to accurately infer demographic histories that deviate

from its pre-defined admixture models. ALDER notes that multi-wave or continuous admixture

can lead to more recent estimates of admixture timing, as the method inherently assumes

single-point admixture. Despite this caveat, ALDER is frequently applied in cases where

admixture is not strictly instantaneous.

All of the timing methods we used placed admixture timing for the different islands closer

together than historical dates of settlement, consistent with historical expectations that the

initial admixture in the southern islands was significant, and that many individuals that

occupied the northern and eastern islands during the second and third settlement stages of

Cabo Verde were already admixed. This observation, together with IBD and kinship patterns,

support the serial founding of the groups of islands as the main model of settlement of Cabo

Verde, as opposed to their independent settling as some historical data suggest[38]. This type of

serial founding scenario is common throughout recent human migration, underscoring that

settlement patterns, in addition to settlement timing, are critical components of accurately

inferring human demographic history.

We found that Cabo Verde's island-specific demographic history and admixture dynamics have

important genomic consequences, as observed with ROH. Despite the relatively recent

colonization of the islands, some individuals presented even lower overall levels of ROH than

African reference populations. We found that low overall levels of ROH in Cabo Verde are

driven by shorter length ROH. This observation is consistent with the idea that shorter ROH

can be attributed to older shared ancestors from the source populations, and these tracts can be

interrupted with admixed ancestry upon admixture. However, many admixed individuals still

present excess long ROH, likely reflecting post-admixture processes such as serial founding and

assortative mating. The observed island-specific distributions of ROH are consistent with the

colonization history of the islands. For example, the cultural dynamics and higher rates of first-

cousin marriages in Fogo[46], possibly driving high levels of long ROH (Fig 4B). In contrast,

Santiago has both the oldest population and the largest population size, and has comparatively

low levels of both shorter and long ROH. These observations suggest that more work, both

empirical and theoretical, is needed to understand the interacting forces of local ancestry and

ROH.

In sum, we provide insights into the population history of Cabo Verde and demonstrate how

admixed populations can provide powerful test cases for understanding demographic processes

and genomic consequences in recent human history. We show that patterns of shared ancestry

between and within the islands (quantified with IBD and kinship estimates) reflect serial

founder effects as well as settlement patterns such as post-admixture nonrandom mating. We

find that accounting for nonrandom mating allows us to improve inference of admixture timing

and better contextualize genomic consequences of admixture dynamics, such as ROH. We find

that differences in ancestry on the X chromosome vs the autosomes reflect sex-biased

demographic processes. Given the ubiquity of admixture throughout modern human

population, these results provide important, generalizable considerations for the study of recent

human evolution.

Supplemental Information Description

This article has 9 associated supplementary figures and 1 supplementary table.

Acknowledgements

We thank Jorge Rocha for his contribution during project conception and sampling, and Greg Barsh for his support in later stages of data collection. We thank Noah Zaitlen for sharing scripts to assist in our application of the Zaitlen et al. (2017) local ancestry disequilibrium method. This work was supported by MRC-UK MR/M01987X/1 and FCT-Portugal PTDC/BIA-BDE/64044/2006 awarded to Sandra Beleza; NIH NIGMS grant R35 GM133481 awarded to Amy Goldberg; NIH NIGMS grant F32 GM139313 awarded to Katharine Korunes; and by CNPq (CSF/PDE 207651/2014-0) awarded to Giordano Bruno Soares-Souza. We acknowledge the source of the reference datasets: 1000G 30x data generated at the New York Genome Center with funds provided by NHGRI Grant 3UM1HG008901-03S1. Finally, we would like to thank the Cabo Verdean participants for their invaluable contributions, and the University of Cabo Verde administration for their support.

Data and Code Availability

Sampling consent forms do not allow for public release of genotype data. Inferred local ancestry information can be found at https://doi.org/10.5281/zenodo.4021277. Code generated for this study can be found at https://github.com/agoldberglab/CaboVerde_Demographic_Analyses.

References

1. Browning, S.R., and Browning, B.L. (2015). Accurate Non-parametric Estimation of Recent
Effective Population Size from Segments of Identity by Descent. Am. J. Hum. Genet. *97*, 404–
418.

2. Browning, S.R., Browning, B.L., Daviglus, M.L., Durazo-Arvizu, R.A., Schneiderman, N.,
Kaplan, R.C., and Laurie, C.C. (2018). Ancestry-specific recent effective population size in the
Americas. PLOS Genet. *14*, e1007385.

3. Goldberg, A., and Rosenberg, N.A. (2015). Beyond 2/3 and 1/3: The Complex Signatures of
Sex-Biased Admixture on the X Chromosome. Genetics *201*, 263–79.

4. Loh, P.-R., Lipson, M., Patterson, N., Moorjani, P., Pickrell, J.K., Reich, D., and Berger, B.
(2013). Inferring admixture histories of human populations using linkage disequilibrium.
Genetics *193*, 1233–54.

5. Ni, X., Yuan, K., Liu, C., Feng, Q., Tian, L., Ma, Z., and Xu, S. (2019). MultiWaver 2.0 :
modeling discrete and continuous gene flow to reconstruct complex population admixtures.
Eur. J. Hum. Genet. *27*, 133–139.

6. Schiffels, S., and Durbin, R. (2014). Inferring human population size and separation history
from multiple genome sequences. Nat. Genet. *46*, 919–925.

7. Zaitlen, N., Huntsman, S., Hu, D., Spear, M., Eng, C., Oh, S.S., White, M.J., Mak, A., Davis, A.,
Meade, K., et al. (2017). The Effects of Migration and Assortative Mating on Admixture Linkage
Disequilibrium. Genetics *205*, 375–383.

8. Busby, G.B.J., Hellenthal, G., Montinaro, F., Tofanelli, S., Bulayeva, K., Rudan, I., Zemunik, T.,
Hayward, C., Toncheva, D., Karachanak-Yankova, S., et al. (2015). The Role of Recent
Admixture in Forming the Contemporary West Eurasian Genomic Landscape. Curr. Biol. *25*,
2518–2526.

9. Fernandes, V., Brucato, N., Ferreira, J.C., Pedro, N., Cavadas, B., Ricaut, F.-X., Alshamali, F.,
and Pereira, L. (2019). Genome-Wide Characterization of Arabian Peninsula Populations:
Shedding Light on the History of a Fundamental Bridge between Continents. Mol. Biol. Evol.
36, 575–586.

10. Font-Porterias, N., Arauna, L.R., Poveda, A., Bianco, E., Rebato, E., Prata, M.J., Calafell, F.,
and Comas, D. (2019). European Roma groups show complex West Eurasian admixture
footprints and a common South Asian genetic origin. PLOS Genet. *15*, e1008417.

11. Lazaridis, I., Patterson, N., Mittnik, A., Renaud, G., Mallick, S., Kirsanow, K., Sudmant, P.H.,
Schraiber, J.G., Castellano, S., Lipson, M., et al. (2014). Ancient human genomes suggest three
ancestral populations for present-day Europeans. Nature *513*, 409–413.

12. Skoglund, P., Malmström, H., Raghavan, M., Storå, J., Hall, P., Willerslev, E., Gilbert, M.T.P.,
Götherström, A., and Jakobsson, M. (2012). Origins and genetic legacy of Neolithic farmers and
hunter-gatherers in Europe. Science *336*, 466–469.

13. Risch, N., Choudhry, S., Via, M., Basu, A., Sebro, R., Eng, C., Beckman, K., Thyne, S.,
Chapela, R., Rodriguez-Santana, J.R., et al. (2009). Ancestry-related assortative mating in Latino
populations. Genome Biol. *10*, R132.

14. Sebro, R., Peloso, G.M., Dupuis, J., and Risch, N.J. (2017). Structured mating: Patterns and
implications. PLOS Genet. *13*, e1006655.

15. Zou, J.Y., Park, D.S., Burchard, E.G., Torgerson, D.G., Pino-Yanes, M., Song, Y.S.,
Sankararaman, S., Halperin, E., and Zaitlen, N. (2015). Genetic and socioeconomic study of mate
choice in Latinos reveals novel assortment patterns. Proc. Natl. Acad. Sci. U. S. A. *112*, 13621–6.

16. Arbiza, L., Gottipati, S., Siepel, A., and Keinan, A. (2014). Contrasting X-Linked and
Autosomal Diversity across 14 Human Populations. Am. J. Hum. Genet. *94*, 827–844.

17. Bustamante, C.D., and Ramachandran, S. (2009). Evaluating signatures of sex-specific
processes in the human genome. Nat. Genet. *41*, 8–10.

18. Keinan, A., Mullikin, J.C., Patterson, N., and Reich, D. (2009). Accelerated genetic drift on
chromosome X during the human dispersal out of Africa. Nat. Genet. *41*, 66–70.

19. Ramachandran, S., Rosenberg, N.A., Feldman, M.W., and Wakeley, J. (2008). Population
differentiation and migration: Coalescence times in a two-sex island model for autosomal and
X-linked loci. Theor. Popul. Biol. *74*, 291–301.

20. Wilkins, J.F., and Marlowe, F.W. (2006). Sex-biased migration in humans: what should we
expect from genetic data? BioEssays *28*, 290–300.

21. Busby, G.B., Band, G., Si Le, Q., Jallow, M., Bougama, E., Mangano, V.D., Amenga-Etego,
L.N., Enimil, A., Apinjoh, T., Ndila, C.M., et al. (2016). Admixture into and within sub-Saharan
Africa. ELife *5*, e15266.

22. Hellenthal, G., Busby, G.B.J., Band, G., Wilson, J.F., Capelli, C., Falush, D., and Myers, S.
(2014). A Genetic Atlas of Human Admixture History. Science *343*, 747–751.

23. Laso-Jadart, R., Harmant, C., Quach, H., Zidane, N., Tyler-Smith, C., Mehdi, Q., Ayub, Q.,
Quintana-Murci, L., and Patin, E. (2017). The Genetic Legacy of the Indian Ocean Slave Trade:

Recent Admixture and Post-admixture Selection in the Makranis of Pakistan. Am. J. Hum. Genet. *101*, 977–984.

24. Ruiz-Linares, A., Adhikari, K., Acuña-Alonzo, V., Quinto-Sanchez, M., Jaramillo, C., Arias, W., Fuentes, M., Pizarro, M., Everardo, P., Avila, F. de, et al. (2014). Admixture in Latin America: Geographic Structure, Phenotypic Diversity and Self-Perception of Ancestry Based on 7,342 Individuals. PLOS Genet. *10*, e1004572.

25. Triska, P., Soares, P., Patin, E., Fernandes, V., Cerny, V., and Pereira, L. (2015). Extensive Admixture and Selective Pressure Across the Sahel Belt. Genome Biol. Evol. *7*, 3484–3495.

26. Blant, A., Kwong, M., Szpiech, Z.A., and Pemberton, T.J. (2017). Weighted likelihood inference of genomic autozygosity patterns in dense genotype data. BMC Genomics *18*, 928.

27. Kirin, M., McQuillan, R., Franklin, C.S., Campbell, H., McKeigue, P.M., and Wilson, J.F. (2010). Genomic Runs of Homozygosity Record Population History and Consanguinity. PLOS ONE *5*, e13996.

28. McQuillan, R., Leutenegger, A.-L., Abdel-Rahman, R., Franklin, C.S., Pericic, M., Barac-Lauc, L., Smolej-Narancic, N., Janicijevic, B., Polasek, O., Tenesa, A., et al. (2008). Runs of Homozygosity in European Populations. Am. J. Hum. Genet. *83*, 359–372.

29. Pemberton, T.J., Absher, D., Feldman, M.W., Myers, R.M., Rosenberg, N.A., and Li, J.Z. (2012). Genomic Patterns of Homozygosity in Worldwide Human Populations. Am. J. Hum. Genet. *91*, 275–292.

30. Mooney, J.A., Huber, C.D., Service, S., Sul, J.H., Marsden, C.D., Zhang, Z., Sabatti, C., Ruiz-Linares, A., Bedoya, G., Fears, S.C., et al. (2018). Understanding the Hidden Complexity of Latin American Population Isolates. Am. J. Hum. Genet. *103*, 707–726.

31. Popejoy, A.B., and Fullerton, S.M. (2016). Genomics is failing on diversity. Nature *538*, 161–164.

32. Popejoy, A.B., Ritter, D.I., Crooks, K., Currey, E., Fullerton, S.M., Hindorff, L.A., Koenig, B., Ramos, E.M., Sorokin, E.P., Wand, H., et al. (2018). The clinical imperative for inclusivity: Race, ethnicity, and ancestry (REA) in genomics. Hum. Mutat. *39*, 1713–1720.

33. Landry, L.G., Ali, N., Williams, D.R., Rehm, H.L., and Bonham, V.L. (2018). Lack Of Diversity In Genomic Databases Is A Barrier To Translating Precision Medicine Research Into Practice. Health Aff. Proj. Hope *37*, 780–785.

34. Beleza, S., Campos, J., Lopes, J., Araújo, I.I., Hoppfer Almada, A., e Silva, A.C., Parra, E.J., and Rocha, J. (2012). The Admixture Structure and Genetic Variation of the Archipelago of Cape Verde and Its Implications for Admixture Mapping Studies. PLoS ONE *7*, e51103.

35. Beleza, S., Johnson, N.A., Candille, S.I., Absher, D.M., Coram, M.A., Lopes, J., Campos, J.,
Araújo, I.I., Anderson, T.M., Vilhjálmsson, B.J., et al. (2013). Genetic Architecture of Skin and
Eye Color in an African-European Admixed Population. PLoS Genet. *9*, e1003372.

36. Fernandes, A.T., Velosa, R., Jesus, J., Carracedo, A., and Brehm, A. (2003). Genetic
differentiation of the Cabo Verde archipelago population analysed by STR polymorphisms.
Ann. Hum. Genet. *67*, 340–347.

37. Verdu, P., Jewett, E.M., Pemberton, T.J., Rosenberg, N.A., and Baptista, M. (2017). Parallel
Trajectories of Genetic and Linguistic Admixture in a Genetically Admixed Creole Population.
Curr. Biol. *27*, 2529–2535.e3.

38. Correia e Silva, A.L. (2002). Dinâmicas de decomposição e recomposição de espaços e
sociedades. In História Geral de Cabo Verde., Santos, MEM, ed. (Lisbon, Portugal; Praia, Cabo
Verde: Instituto de Investigação Científica Tropical, Instituto Nacional de Investigação,
Promoção e Património Culturais de Cabo Verde), pp. 1–66.

39. Baleno, I.C. (2001). Povoamento e Formação da Sociedade. In História Geral de Cabo Verde.,
Albuquerque, L and Santos, MEM, ed. (Lisbon, Portugal; Praia, Cabo Verde: Instituto de
Investigação Científica Tropical, Instituto Nacional de Investigação, Promoção e Património
Culturais de Cabo Verde), pp. 125–177.

40. Russell-Wood, A. (1998). The Portuguese Empire, 1415-1808: A World on the Move.
(Baltimore, Maryland: Johns Hopkins University Press).

41. Clarke, L., Fairley, S., Zheng-Bradley, X., Streeter, I., Perry, E., Lowy, E., Tassé, A.-M., and
Flicek, P. (2017). The international Genome sample resource (IGSR): A worldwide collection of
genome variation incorporating the 1000 Genomes Project data. Nucleic Acids Res. *45*, D854–
D859.

42. Fairley, S., Lowy-Gallego, E., Perry, E., and Flicek, P. (2020). The International Genome
Sample Resource (IGSR) collection of open human genomic variation resources. Nucleic Acids
Res. *48*, D941–D947.

43. Correia e Silva, A.L. (2001). Espaço, ecologia e economia interna. In História Geral de Cabo
Verde., de Albuquerque L, Santos, MEM, ed. (Lisbon, Portugal; Praia, Cabo Verde: Instituto de
Investigação Científica Tropical, Instituto Nacional de Investigação, Promoção e Património
Culturais de Cabo Verde), p.

44. Cabral, I. (2001). Ribeira Grande: vida urbana, gente, mercancia, estagnação. In História
Geral de Cabo Verde., Santos, MEM, ed. (Lisbon, Portugal; Praia, Cabo Verde: Instituto de
Investigação Científica Tropical, Instituto Nacional de Investigação, Promoção e Património
Culturais de Cabo Verde), pp. 225–273.

45. Cabral, I. (2012). A evolução da sociedade foguense: Através de um estudo prosopográfico
da elite da ilha (sec. XVI-XVIII).

46. Barbosa, Gilda (1997). Casamento entre primos. J. Terra Nova No 246.

47. Alexander, D.H., Novembre, J., and Lange, K. (2009). Fast model-based estimation of
ancestry in unrelated individuals. Genome Res. *19*, 1655–1664.

48. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J.,
Sklar, P., de Bakker, P.I.W., Daly, M.J., et al. (2007). PLINK: a tool set for whole-genome
association and population-based linkage analyses. Am. J. Hum. Genet. *81*, 559–75.

49. Delaneau, O., Zagury, J.-F., and Marchini, J. (2013). Improved whole-chromosome phasing
for disease and population genetic studies. Nat. Methods *10*, 5–6.

50. Maples, B.K., Gravel, S., Kenny, E.E., and Bustamante, C.D. (2013). RFMix: a discriminative
modeling approach for rapid and robust local-ancestry inference. Am. J. Hum. Genet. *93*, 278–
88.

51. Hamid, I., Korunes, K., Beleza, S., and Goldberg, A. (2020). Rapid adaptation to malaria
facilitated by admixture in the human population of Cabo Verde (Zenodo).

52. Tang, H., Peng, J., Wang, P., and Risch, N.J. (2005). Estimation of individual admixture:
analytical and study design considerations. Genet. Epidemiol. *28*, 289–301.

53. Tang, H., Coram, M., Wang, P., Zhu, X., and Risch, N. (2006). Reconstructing Genetic
Ancestry Blocks in Admixed Individuals. Am. J. Hum. Genet. *79*, 1–12.

54. Patterson, N., Price, A.L., and Reich, D. (2006). Population Structure and Eigenanalysis.
PLOS Genet. *2*, e190.

55. Price, A.L., Patterson, N.J., Plenge, R.M., Weinblatt, M.E., Shadick, N.A., and Reich, D.
(2006). Principal components analysis corrects for stratification in genome-wide association
studies. Nat. Genet. *38*, 904–909.

56. Zou, J.Y., Halperin, E., Burchard, E., and Sankararaman, S. (2015). Inferring parental
genomic ancestries using pooled semi-Markov processes. Bioinformatics *31*, i190–i196.

57. Ochoa, A., and Storey, J.D. (2019). New kinship and FST estimates reveal higher levels of
differentiation in the global human population. BioRxiv 653279.

58. Browning, B.L., and Browning, S.R. (2013). Improving the accuracy and efficiency of
identity-by-descent detection in population data. Genetics *194*, 459–471.

59. Nelson, D., Kelleher, J., Ragsdale, A.P., Moreau, C., McVean, G., and Gravel, S. (2020).
Accounting for long-range correlations in genome-wide simulations of large cohorts. PLOS
Genet. *16*, e1008619.

60. Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., and Ideker, T. (2003). Cytoscape: A Software Environment for Integrated
Models of Biomolecular Interaction Networks. Genome Res. *13*, 2498–2504.

61. Szpiech, Z.A., Blant, A., and Pemberton, T.J. (2017). GARLIC: Genomic Autozygosity
Regions Likelihood-based Inference and Classification. Bioinformatics *33*, 2059–2062.

62. Reed, F.A., and Tishkoff, S.A. (2006). African human diversity, origins and migrations. Curr.
Opin. Genet. Dev. *16*, 597–605.

63. Tishkoff, S.A., and Williams, S.M. (2002). Genetic analysis of African populations: human
evolution and complex disease. Nat. Rev. Genet. *3*, 611–621.

64. Ceballos, F.C., Joshi, P.K., Clark, D.W., Ramsay, M., and Wilson, J.F. (2018). Runs of
homozygosity: windows into population history and trait architecture. Nat. Rev. Genet. *19*, 220–
234.

65. Goldberg, A., Günther, T., Rosenberg, N.A., and Jakobsson, M. (2017). Ancient X
chromosomes reveal contrasting sex bias in Neolithic and Bronze Age Eurasian migrations.
Proc. Natl. Acad. Sci. *114*, 2657–2662.

66. Eltis, D., Richardson, D., and Blight, D.W. (2010). Atlas of the Transatlantic Slave Trade (Yale
University Press).

67. Micheletti, S.J., Bryc, K., Esselmann, S.G.A., Freyman, W.A., Moreno, M.E., Poznik, G.D.,
Shastri, A.J., Agee, M., Aslibekyan, S., Auton, A., et al. (2020). Genetic Consequences of the
Transatlantic Slave Trade in the Americas. Am. J. Hum. Genet. *107*, 265–277.

68. Pero de Guimarães (1991). Corregedor de Cabo Verde, carta de 22 de Maio de 1513. In
História Geral de Cabo Verde., de Albuquerque L, Santos, MEM, ed. (Lisbon, Portugal; Praia,
Cabo Verde: Instituto de Investigação Científica Tropical, Instituto Nacional de Investigação,
Promoção e Património Culturais de Cabo Verde), pp. 219–223.

69. Seibert, G. (2014). Crioulização em Cabo Verde e São Tomé e Príncipe: divergências
históricas e identitárias. Afro-Ásia 41–70.

70. Robinson, M.R., Kleinman, A., Graff, M., Vinkhuyzen, A.A.E., Couper, D., Miller, M.B.,
Peyrot, W.J., Abdellaoui, A., Zietsch, B.P., Nolte, I.M., et al. (2017). Genetic evidence of
assortative mating in humans. Nat. Hum. Behav. *1*, 1–13.

Fig 1 | Shared ancestry between and within the islands in the context of geography. Each island has a corresponding cluster of nodes representing all sampled individuals, with the individuals localized to be adjacent to the island where they were sampled. Node placement within islands is determined by a force-directed algorithm using pairwise shared IBD, meaning that the spread of each cluster reflects the level of relatedness in each population. Edges between the nodes represent total IBD tract length for pairs of individuals sharing more than 150 cM of total IBD, with edges colored using the log transformed total IBD length.

Fig 2 | Correlation between inferred parental ancestries. (A) For each Cabo Verdean individual, the inferred Parent 1 vs Parent 2 ancestry proportions shown for an example chromosome (Chromosome 7), colored by island. (B) shows the set of correlation coefficients between inferred parental ancestries for each autosomal chromosome.

Fig 3 | Timelines of inferred generations of admixture for each island of Cabo Verde. The island regions are listed in order of known settlement timing beginning with Santiago (A), followed by Fogo (B), and then the later migrations to the Northwest Cluster (C) and Boa Vista (D). For each island, admixture timing inferred with population genetic methods are shown in comparison to historical records. Historical estimates of when each island was first settled are shown in generations, based on a range of generation times (20-30 year generation time). LAD-based estimates of admixture timing reflect the range of estimates generated under the assumption of constant migration rate (m = 0.01; yielding older estimates) to the assumption of no migration (m = 0; yielding more recent estimates). MultiWaver results show the admixture generations inferred under the admixture model selected by MultiWaver, with 0.95 CI.

Fig 4 | ROH content by island and in the context of ancestry. (A-B) Violin plots show the population-specific distributions of the total (summed over each genome) length of autosomal ROH per individual. Solid black dots represent the within-population means. (C-D) Total length of autosomal ROH per individual plotted against West African ancestry proportions and colored by population.

Fig 5 | Sex-biased admixture in Cabo Verde. (A) The distribution of West African ancestry proportion on the autosomes and X chromosome for each of the island regions, estimated with ADMIXTURE. (B) Under a model of constant admixture over time, the fraction of the total contribution of genetic material originating from females for West African and European source populations. Here, we show the distribution of parameter sets for the smallest 0.1% of Euclidean distances between the model-predicted and observed X and autosomal ancestry from a grid of possible parameter values. The range of sex-specific contributions from West African and European source populations that produce ancestry estimates closest to those observed in Cabo Verde are shown for Cabo Verde as a whole (left), and then broken down by region, with medians (dashed lines).

Fig 6 | Autosomal (chromosome 7) vs X chromosome distributions of ROH by class. (A) shows the population-specific distributions of the total length of shorter ROH per individual on the X chromosome compared to an autosome. Chromosome 7 was chosen as the autosomal point of comparison, given that it is the autosome most similar in size to the X chromosome. Totals are shown for females only, so that the same samples are being compared across the X chromosome and chromosome 7. Totals are plotted per Mb to account for slight differences in chromosome lengths. (B) shows the same information for long ROH.