Widespread premature transcription termination of *Arabidopsis thaliana* NLR genes by the spen protein FPA

Matthew T. Parker[1*], Katarzyna Knop[1*], Vasiliki Zacharaki[1*], Anna V. Sherwood[1], Daniel Tome[2], Xuhong Yu[3], Pascal Martin[3], Jim Beynon[2], Scott Michaels[3], Geoffrey J. Barton[1] and Gordon G. Simpson[1,4^]

[1]School of Life Sciences, University of Dundee, Dow Street, Dundee DD1 5EH, UK.
[2]School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK.
[3]Department of Biology, Indiana University, Bloomington, IN 47405, USA.
[4]The James Hutton Institute, Invergowrie DD2 5DA, UK.

*Contributed equally

^Correspondence g.g.simpson@dundee.ac.uk

Keywords: NLR, Nanopore, direct RNA sequencing, spen, FPA, m[6]A, alternative polyadenylation, chimeric RNA, disease resistance, transcription termination

Abstract

Genes involved in disease resistance are some of the fastest evolving and most diverse components of genomes. Large numbers of nucleotide-binding, leucine-rich repeat receptor (NLR) genes are found in plant genomes and are required for disease resistance. However, NLRs can trigger autoimmunity, disrupt beneficial microbiota or reduce fitness. It is therefore crucial to understand how NLRs are controlled. Here we show that the RNA-binding protein FPA mediates widespread premature cleavage and polyadenylation of NLR transcripts, thereby controlling their functional expression and impacting immunity. Using long-read Nanopore direct RNA sequencing, we resolved the complexity of NLR transcript processing and gene annotation. Our results uncover a co-transcriptional layer of NLR control with implications for understanding the regulatory and evolutionary dynamics of NLRs in the immune responses of plants.

Introduction

In plants and animals, NLR (nucleotide-binding, leucine-rich repeat receptor) proteins function to detect the presence and activity of pathogens (Barragan and Weigel, 2020; Jones et al., 2016; Tamborski and Krasileva, 2020). Plant genomes can encode large numbers of NLR genes, which often occur in physical clusters (Jiao and Schneeberger, 2020; Wei et al., 2016). Powerful selective pressure drives the rapid birth and death of NLR genes, resulting in intraspecific diversity in NLR alleles and gene number. Consequently, the near-complete repertoire of Arabidopsis NLR genes was only recently revealed using long-read DNA sequencing of diverse Arabidopsis accessions (Van de Weyer et al., 2019).

In plants, NLR proteins generally comprise an N-terminal Toll/interleukin receptor (TIR), coiled-coil (CC) or RPW8 domain that facilitates signalling; a central nucleotide-binding NB-ARC domain that acts as a molecular switch; and C-terminal leucine-rich repeats (LRRs) that interact with target proteins. NLRs can recognise pathogen effectors either directly by binding to them through LRR domains or indirectly by detecting modifications to host proteins caused by effector action. In some cases, domains of host proteins targeted by pathogen effectors have been incorporated into NLRs as integrated domains (or decoys) (Le Roux et al., 2015). NLRs that interact directly with effectors are under high levels of diversifying selection to modify their recognition specificities, resulting in significant allelic polymorphism (Prigozhin and Krasileva, 2021). Genomic variation also yields diversity in NLR protein organisation, through domain swapping or truncating mutations, and NLR isoforms that lack NB-ARC or LRR domains can function in plant immune responses (Nishimura et al., 2015; Swiderski et al., 2009; Zhang and Gassmann, 2007). The consequence of this diversity is that there is no one-size-fits-all explanation of how NLR proteins function (Barragan and Weigel, 2020).

The benefit of NLRs to the host is disease resistance, but the costs of increased NLR diversity or activity can include detrimental autoimmunity (Rodriguez et al., 2016), reduced association with beneficial microbes (Yang et al., 2010) and a general reduction in fitness

(Tian et al., 2003). In some cases, autoimmunity caused by epistatic interactions involving
NLRs can cause hybrid necrosis (Li et al., 2020). Therefore, a key question is how NLRs are
regulated to enable limited expression for pathogen surveillance but enhanced expression
during defence responses. This problem is compounded by selective pressure on NLRs
because regulatory processes must keep pace with the emergence of new NLR genes and
gain or loss of function in others. Consequently, the regulation of NLRs is one of the most
important and difficult challenges faced by plants.

NLR control measures occur at different stages of gene expression (Lai and Eulgem,
2018). For example, microRNAs limit the expression of many NLRs by targeting conserved
regions encoded in NLR mRNAs and triggering cascades of phased siRNAs that broadly
suppress NLR activity (Cai et al., 2019; Canto-Pastor et al., 2019; Shivaprasad et al., 2012;
Zhai et al., 2011). Alternative splicing, which promotes the simultaneous expression of more
than one NLR isoform, is required for the functions of both the *N* gene which provides
resistance to tobacco mosaic virus (Dinesh-Kumar and Baker, 2000), and *RECOGNITION*
OF PSEUDOMONAS SYRINGAE 4 (*RPS4*), which confers resistance to *Pseudomonas*
syringae DC3000 in Arabidopsis (Zhang and Gassmann, 2007). Alternative polyadenylation
at intragenic heterochromatin controls the expression of Arabidopsis *RECOGNITION OF*
PERONOSPORA PARASITICA 7 (*RPP7*), with functional consequences for immunity against
the oomycete pathogen *Hyaloperonospora arabidopsidis* (Tsuchiya and Eulgem, 2013).
Finally, RNA surveillance pathways control NLRs. For example, null mutants defective in
nonsense mediated RNA decay (NMD) are lethal in Arabidopsis because they trigger NLR
RPS6-dependent autoimmunity (Gloggnitzer et al., 2014). Conversely, mutations in the RNA
exosome, which degrades RNAs in a 3′ to 5′ direction, suppress *RPS6*-dependent
autoimmune phenotypes (Takagi et al., 2020). Fine tuning of different levels of NLR control
may be integrated to produce quantitative patterns of disease resistance (Corwin and
Kliebenstein, 2017), but our understanding of how this occurs globally is fragmentary and
incomplete (Adachi et al., 2019).

The RNA-binding protein FPA was first identified as a factor required for the control
of Arabidopsis flowering time (Koornneef et al., 1991). Loss-of-function *fpa* mutants flower
late due to elevated levels of the floral repressor, FLC (Schomburg et al., 2001). However,
this cannot be the only function of FPA because it is much more widely conserved than *FLC*.
FPA is a member of the spen family of proteins, which are defined by three N-terminal RNA
recognition motifs and a C-terminal protein interaction SPOC domain (Ariyoshi and
Schwabe, 2003). We previously showed that FPA controls the site of cleavage and
polyadenylation in some mRNAs, including autoregulation of *FPA* pre-mRNA (Duc et al.,
2013; Hornyik et al., 2010; Lyons et al., 2013). These findings were extended to show that
FPA can affect poly(A) site choice at genes with intronic heterochromatin, including *RPP7*
(Deremetz et al., 2019). The poly(A) selection mechanism used by FPA remains unclear.
FPA might mediate poly(A) choice either directly by recruiting the RNA 3′ end processing
machinery to sensitive sites or indirectly, for example by influencing splicing, chromatin

modifications or the rate of transcription by RNA Polymerase II (Pol II). We previously used Helicos BioSciences direct RNA sequencing (Helicos DRS) to map the 3′ ends of Arabidopsis polyadenylated transcripts and identify genes affected by transcriptome-wide loss of FPA function (Duc et al., 2013; Sherstnev et al., 2012). A limitation of this approach was that it could only identify RNA 3′ end positions, and so could not resolve other potential roles of FPA in gene expression.

In this study, we used two approaches to gain a clearer understanding of how FPA functions. We first investigated which proteins FPA associates with inside living plant cells. Next, we analysed the global impact of different levels of FPA activity on gene expression. For this, we combined Helicos DRS with short-read Illumina RNA-Seq and Oxford Nanopore Technologies (Nanopore) DRS, which can reveal the authentic processing and modification of full-length mRNAs (Parker et al., 2020). Using these combined data together with new computational approaches to study RNA processing, we found that the predominant role of FPA is to promote poly(A) site choice. In addition, we uncovered an unusual degree of complexity in the processing of NLR mRNAs, which is sensitive to FPA. The finding that premature transcription termination functions as an additional layer of NLR expression control has implications for understanding the dynamics of NLR regulation and evolution.

Results

FPA co-purifies with proteins that mediate mRNA 3′ end processing

In order to understand how FPA controls the site of mRNA 3′ end formation, we used *in vivo* interaction proteomics–mass spectrometry (*IVI-MS*) to identify which proteins FPA associates with inside living plant cells. First, we fixed molecular interactions using formaldehyde infiltration of Arabidopsis seedlings expressing FPA fused to YFP (*35S::FPA:YFP*). Wild-type Columbia-0 (Col-0) seedlings treated in the same way were used as a negative control. We then purified nuclei and performed GFP-trap immunopurification followed by liquid chromatography–tandem mass spectrometry (LC-MS/MS) to identify FPA-associated proteins. By comparing the proteins detected in three biological replicates of *35S::FPA:YFP* and Col-0, we identified 203 FPA co-purifying proteins with a median $\log_2$ fold change in abundance of greater than two (Figure 1, Supplementary file 1). At least 56% (113) of the enriched proteins are poly(A)+ mRNA-binding proteins as established by orthogonal RNA-binding proteome analysis (Bach-Pages et al., 2020; Reichel et al., 2016).

Consistent with FPA control of mRNA 3′ end formation, 14 highly conserved cleavage and polyadenylation factors (CPFs) co-purified with FPA (Figure 1A, Supplementary file 1). These include members of the cleavage and polyadenylation specificity factor (CPSF) complex, cleavage stimulating factor (CstF) complex, and cleavage factor I and II (CFIm/CFIIm) complexes. The U2AF and U2 spliceosome components that interact with CFIm–CPSF to mediate terminal exon definition were also detected (Kyburz et al., 2006) (Figure 1B, Supplementary file 1). We additionally detected both subunits of Pol II. Characteristically, Serine[5] of the Pol II C-terminal domain (CTD) heptad repeat is

phosphorylated when Pol II is at the 5′ end of genes, and Ser[2] is phosphorylated when Pol II
is at the 3′ end (Komarnitsky et al., 2000). The position-specific phosphorylation of these
sites alters the RNA processing factors which are recruited to the CTD at the different stages
of transcription. We found that the kinase CDKC;2, which phosphorylates Ser[2] (Wang et al.,
2014), and the phosphatase CPL1 (homolog of yeast Fcp1), which dephosphorylates Ser[5]
(Koiwa et al., 2004), co-purified with FPA. We also detected the homolog of the human
exonuclease XRN2 (known as XRN3 in Arabidopsis), which mediates Pol II transcription
termination (Krzyszton et al., 2018).

A second major class of proteins that co-purified with FPA are components of the
autonomous flowering pathway (Andres and Coupland, 2012; Simpson, 2004)
(Figure 1C,Supplementary file 1). FPA functions in the autonomous pathway to limit
expression of the floral repressor *FLC*. FPA activity is associated with alternative
polyadenylation of long non-coding RNAs that are transcribed antisense to the *FLC* locus
(Andres and Coupland, 2012; Simpson, 2004). Consistent with this, conserved CPF proteins
such as FY (WDR33) (Hornyik et al., 2010; Liu et al., 2007), PCFS4 (Xing et al., 2008), CSTF64
and CSTF77 (Liu et al., 2010) were previously identified in late flowering mutant screens.
Other detected autonomous pathway factors are proteins with established roles in pre-
mRNA processing, including HLP1 (Zhang et al., 2015), FLK (Mockler et al., 2004) and
EMB1579/RSA1 (Zhang et al., 2020b). Notably, FLK has been found to associate with PEP,
HUA1 and HEN4 (Zhang et al., 2015), and we identified all four of these as FPA co-purifying
proteins. In addition to regulating *FLC*, the FLK–PEP complex has been shown to control
alternative polyadenylation within pre-mRNA encoding the floral homeotic transcription
factor AGAMOUS (Mockler et al., 2004). Their co-purification with FPA suggests that this
role may be more global and involve direct interactions at RNA 3′ ends.

A third group of proteins that co-purified with FPA are conserved members of the
mRNA N[6]-methyladenosine (m[6]A) writer complex (Ruzicka et al., 2017) (Figure 1D,
Supplementary file 1). The m[6]A modification mediated by this complex is predominately
targeted to the 3′ untranslated region (UTR) of Arabidopsis protein-coding mRNAs (Parker
et al., 2020). The co-purification of FPA with m[6]A writer complex components may be
explained by either a direct role for FPA in m[6]A modification or, more simply, because both
CPF and m[6]A writer proteins are found at RNA 3′ ends.

The picture that emerges from this analysis is that FPA is located in proximity to
proteins that promote cleavage, polyadenylation, transcription termination and RNA
modification at the 3′ end of Pol II-transcribed genes.

FPA co-localises with RNA Pol II Ser[2] at the 3′ end of Arabidopsis genes
We next used an orthogonal approach to investigate the association of FPA with proteins
that function at the 3′ end of Pol II-transcribed genes. We performed chromatin
immunoprecipitation sequencing (ChIP-Seq) using antibodies against FPA and Pol II
phosphorylated at either Ser[5] or Ser[2] of the CTD heptad repeat (Yu et al., 2019). Our

metagene analysis revealed that FPA is enriched at the 3′ end of genes and co-localises
with Pol II phosphorylated at Ser[2] of the CTD (Figure 1E, Figure 1–figure supplement 1). The
close relationship between FPA and Pol II Ser[2] is reinforced by changes in the distribution
of Pol II isoforms in *fpa* mutants. For example, we previously showed that FPA is required
for 3′ end processing at *PIF5* (Duc et al., 2013). Pol II Ser[2] was enriched at the 3′ end of *PIF5*
in Col-0, but depleted from this region in *fpa-7* mutants (Figure 1–figure supplement 2).
Together, these orthogonal ChIP-Seq and *IVI-MS* analyses reveal the close association of
FPA with proteins involved in 3′ end processing and transcription termination at the 3′ end
of Arabidopsis genes.
196
FPA predominantly promotes poly(A) site choice
We next asked which RNA processing events are controlled by FPA. We used a
combination of Illumina RNA-Seq and Helicos and Nanopore DRS technologies to analyse
three different genetic backgrounds expressing different levels of FPA activity: wild-type
Col-0, loss-of-function *fpa-8* and a line overexpressing FPA fused to YFP (*35S::FPA:YFP*). In
combination, these orthogonal sequencing technologies can reveal different features of
transcriptomes: Helicos DRS short reads identify the 3′ ends of mRNAs, but cannot reveal
the full properties of the corresponding transcripts (Ozsolak et al., 2009); Illumina RNA-Seq
produces short reads derived from all expressed regions, meaning that changes in RNA 3′
end processing can only be detected by differences in coverage (Xia et al., 2014); and
Nanopore DRS long reads define the 3′ ends of mRNAs in the context of reads that can
correspond to full-length transcripts (Parker et al., 2020). For each genotype, we performed
three biological replicates with Helicos DRS, six with Illumina RNA-Seq and four with
Nanopore DRS. The resultant sequencing statistics are detailed in Supplementary file 1.
We first assessed the utility of the three sequencing technologies to map changes
in mRNA processing by focusing on the *FPA* locus. FPA autoregulates its expression by
promoting premature cleavage and polyadenylation within intron 1 of *FPA* pre-mRNA (Duc
et al., 2013; Hornyik et al., 2010). Consistent with this, a proximal poly(A) site in the first
intron and distal sites in the terminal intron and exon of *FPA* could be mapped in Col-0
using Nanopore and Helicos DRS (Figure 2A). Using all three data types, we detected a
quantitative shift towards selection of distal poly(A) sites in the loss-of-function *fpa-8* mutant
and a strong shift to proximal poly(A) site selection when FPA is overexpressed
(*35S::FPA:YFP*). Nanopore DRS provided the clearest picture of alternative polyadenylation
events because full-length reads reveal poly(A) site choice in the context of other RNA
processing events.
We next asked how transcriptome-wide RNA processing is affected by FPA activity.
Since mutations in FPA cause readthrough of annotated 3′UTRs (Duc et al., 2013), we
applied the software tool StringTie2 (Pertea et al., 2015) to create a bespoke reference
annotation with Nanopore DRS reads from Col-0, *fpa-8* and *35S::FPA:YFP*. We then
measured how changes in FPA expression altered the 3′ end distribution at each locus

using the earth mover's distance (EMD; also known as the Wasserstein distance). EMD
indicates the "work" required to transform one normalised distribution into another based
on the proportion of 3′ ends that would have to be moved and by what distance. We used
an EMD permutation test, in which reads are randomly shuffled between conditions, to
estimate p-values for each locus. Loci with an EMD greater than 25 and a false discovery
rate (FDR) less than 0.05 were considered differentially polyadenylated.

Using this approach on Nanopore DRS data, we identified 285 and 293 loci with
alternative polyadenylation events in *fpa-8* and *35S::FPA:YFP*, respectively (Figure 2B). In
all, 77.9% (222) of loci with alternative polyadenylation in *fpa-8* displayed a positive change
in the mean 3′ end position, indicating a predominant shift to distal poly(A) site selection
(Figure 2B, left panel). These loci also had greater effect sizes than those with shifts towards
proximal poly(A) sites (Figure 2C, left panel). In contrast, 56.7% (166) of loci with alternative
polyadenylation in *35S::FPA:YFP* displayed a negative change in the mean 3′ end position,
indicating a shift towards proximal poly(A) sites (Figure 2B, right panel). These loci had
greater effect sizes than those with positive changes in 3′ end profile (Figure 2C, right
panel). A total of 16 loci displayed a shift to distal poly(A) site selection in *fpa-8* and to
proximal poly(A) site selection in *35S::FPA:YFP* (hypergeometric test $p=3.9\times10^{-7}$),
demonstrating that loss of function versus overexpression of FPA can result in reciprocal
patterns of poly(A) site choice.

We used the same approach to identify loci with FPA-dependent alternative
polyadenylation in Helicos DRS data. We identified 319 and 299 genes with alternative
polyadenylation events in *fpa-8* and *35S::FPA:YFP*, respectively (Figure 2D and E).
Consistent with Nanopore DRS analysis, the predominant shifts in *fpa-8* and *35S::FPA:YFP*
were towards distal (79.0% or 252 loci) and proximal (75.3% or 225 loci) poly(A) sites,
respectively. In all, 44 loci displayed a shift to distal poly(A) sites in *fpa-8* and to proximal
poly(A) sites in *35S::FPA:YFP* (hypergeometric test $p=4.8\times10^{-30}$), again demonstrating
reciprocal poly(A) site selection depending on FPA activity. Of the 222 loci identified with
shifts to distal poly(A) sites in *fpa-8* using Nanopore DRS, 39.6% (88) were also detected
using Helicos DRS (Figure 2–figure supplement 1). Likewise, 44.0% of loci (73) with proximal
polyadenylation detected in *35S::FPA:YFP* using Nanopore DRS were also detected using
Helicos DRS. Across the DRS datasets, we identified 59 loci for which reciprocal poly(A) site
regulation by FPA could be detected by Nanopore DRS and/or Helicos DRS.

In order to analyse the Illumina RNA-Seq data, we developed annotation-agnostic
software for detecting alternative RNA 3′ end processing events, using a similar approach
to the existing tools DERfinder (Collado-Torres et al., 2017), RNAprof (Tran Vdu et al., 2016)
and DEXSeq (Anders et al., 2012). We segmented Illumina RNA-Seq data by coverage and
relative expression in *fpa-8* or *35S::FPA:YFP* compared with Col-0. Segmented regions
were grouped into transcriptional loci using the annotations generated from Nanopore
DRS datasets. Differential usage of regions within each locus was then tested using
DEXSeq. Using this approach, we identified 2535 loci with differential RNA processing

events in *fpa-8*: 1792 were upregulated, 390 were downregulated, and 353 had both upregulated and downregulated regions (FDR<0.05, absolute logFC>1; Figure 2F, left panel). A total of 1747 loci with differential RNA processing events were identified in *35S::FPA:YFP*: 997 were upregulated, 532 were downregulated, and 218 had both upregulated and downregulated regions (Figure 2F, right panel). The median effect size for differentially processed regions was greater for upregulated regions than for downregulated regions in *fpa-8*. This is consistent with an increase in transcriptional readthrough events and elevated expression of intergenic regions and downstream genes. In contrast, the median effect size for differentially processed regions was similar for up- and downregulated regions in *35S::FPA:YFP*. This is consistent with an increase in the relative expression of proximal exonic and intronic regions, and loss of expression of distal exonic regions caused by preferential selection of proximal poly(A) sites. Similar results were seen for differential splice junction usage analysis (Figure 2G), suggesting that changes in splicing are the indirect effects of altered 3′ end processing in *fpa-8*, rather than direct effects of FPA on splice site choice. Evidence of this can be seen at the *PIF5* locus, where readthrough results in increased cryptic and canonical splicing of downstream *PAO3* (Figure 2–figure supplement 2).

We next asked whether FPA influences RNA modification. Our *IVI-MS* analysis had revealed that conserved members of the Arabidopsis m^{6}A writer complex co-purify with FPA (Figure 1D,Supplementary file file 1). The human proteins most closely related to FPA are RBM15/B, which co-purify with the human m^{6}A writer complex and are required for m^{6}A deposition (Patil et al., 2016). We used LC-MS/MS to analyse the m^{6}A/A (adenosine) ratio in mRNA purified from Col-0, *fpa-8*, *35S::FPA:YFP* and a mutant defective in the m^{6}A writer complex component VIR (*vir-1*). Consistent with previous reports, the level of mRNA m^{6}A in the hypomorphic *vir-1* allele was reduced to approximately 10% of wild-type levels (Parker et al., 2020; Ruzicka et al., 2017) (Figure 2–figure supplement 3). However, we detected no differences in the m^{6}A level between genotypes with altered FPA activity. Therefore, we conclude that FPA does not influence global levels of mRNA m^{6}A methylation.

Finally, we asked whether the FPA-dependent global changes in alternative polyadenylation result from an indirect effect on chromatin state. We previously showed that FPA controls the expression of histone demethylase IBM1 by promoting proximal polyadenylation within *IBM1* intron 7 (Duc et al., 2013). IBM1 functions to restrict H3K9me^2 levels, and *ibm1* mutants accumulate ectopic heterochromatic marks in gene bodies, which affects RNA processing at certain loci (Miura et al., 2009; Saze et al., 2008). When we analysed two independent ChIP-Seq datasets of H3K9me^2 in *ibm1–4* mutants (Inagaki et al., 2017; Lai et al., 2020), we found that only 10.6% of loci with altered poly(A) site choice in *35S::FPA:YFP* have altered H3K9me^2 in *ibm1* mutants compared with 14.2% of all loci tested (hypergeometric p=0.97; Figure 2–figure supplement 4). This result suggests that FPA-dependent regulation of poly(A) site choice is not an indirect consequence of FPA control of *IBM1*.

307   Overall, these analyses reveal that the primary function of FPA is to control poly(A)
308   site choice. FPA predominantly promotes poly(A) site selection; hence, *fpa* loss-of-function
309   backgrounds exhibit readthrough at sites used in the wild type, whereas FPA
310   overexpression results in increased selection of proximal poly(A) sites.

312   NLRs are major targets of FPA-dependent poly(A) site selection

We next asked which groups of genes are sensitive to FPA-dependent alternative
polyadenylation. We used InterPro annotations (Mitchell et al., 2019) to perform protein
family domain enrichment analysis of the loci affected by FPA (revealed by the Nanopore
and Helicos DRS analyses). We found that sequences encoding NB-ARC, Rx-like coiled coil
(CC), and/or LRR domains were enriched amongst the loci with increased proximal
polyadenylation in *35S::FPA:YFP* (Figure 3A and B). This combination of domains is
associated with NLR disease resistance proteins.

The Col-0 accession contains at least 206 genes encoding some combination of TIR,
CC, RPW8, NB-ARC and LRR domains, which might be classified as NLRs or partial NLRs
(Van de Weyer et al., 2019). In general, these can be grouped according to their encoded
N-terminal domain as TIR (TNLs), CC (CNLs) or RPW8 (RNLs) genes. We manually examined
these NLR genes to identify those with alternative polyadenylation. Reannotation of some
loci was required to interpret the effects of FPA regulation. For example, we found that the
TNL gene *AT5G46490,* located in the RPS6 cluster, is incorrectly annotated as two loci,
AT5G46490 and *AT5G46500* (Figure 3–figure supplement 1). Nanopore DRS evidence
indicates that this is actually a single locus with a previously unrecognised 2.7 kb intron
containing a proximal poly(A) site, the use of which is regulated by FPA. This interpretation
is supported by nanoPARE data (Schon et al., 2018), which showed no evidence of capped
5′ ends originating from the annotated downstream gene. Use of the distal poly(A) site
introduces an additional ~400 amino acids to the C-terminus of the protein. This C-terminal
region has homology to other NLRs in the *RPS6* cluster, and is predicted to introduce
additional LRR repeats (Martin et al., 2020) (Figure 3–figure supplement 2).

Notably, we could also reannotate the chromosomal region around *RPS6* itself. The
extreme autoimmunity phenotypes of NMD mutants and mitogen-activated kinase pathway
mutants require RPS6 but the mechanisms involved are not understood (Gloggnitzer et al.,
2014; Takagi et al., 2020). Nanopore DRS indicates that the 3′UTR of *RPS6* is complex, with
multiple splicing events and poly(A) sites (Figure 3C). We also detected transcripts
expressed from this region that do not appear to be contiguous with *RPS6* 3′UTR reads.
Instead, these reads correspond to an independent unannotated gene that overlaps the
RPS6 3′UTR. This interpretation is supported by capped RNA 5′ ends detected in this region
by nanoPARE (Schon et al., 2018). In addition, Nanopore DRS analysis of the RNA exosome
mutant *hen2-2* (Parker et al., 2021) revealed that this unannotated gene is expressed at
relatively high levels, but that the transcripts are subject to degradation. Consequently,
steady-state levels of RNA expressed from this locus are relatively low in Col-0. The gene

encodes a TIR domain similar to that of RPS6 (Figure 3D). Therefore, use of the distal *RPS6* poly(A) site constitutes readthrough into the downstream TIR-domain-only NLR. Based on these analyses, we conclude that long-read Nanopore DRS data have the potential to correct NLR gene annotation at complex loci that cannot be resolved by genome annotation software or short-read Illumina RNA-Seq.

Widespread premature transcription termination of NLRs includes frequent selection of poly(A) sites in protein-coding exons

Of the 206 NLR genes examined, 124 had a sufficient level of expression levels to identify alternative polyadenylation in the Nanopore DRS data; of these 124, 62 (50.0%) were found to have FPA-dependent alternative polyadenylation (Tables 1–3). Of the 74 expressed NLRs located in major clusters, 44 (59.5%) were regulated by FPA (chi^2 *p*=0.02) (Lee and Chae, 2020). The localisation of NLRs to large genomic clusters is known to facilitate diversification (Barragan and Weigel, 2020). Consistent with this, 20 (71.4%) of the 28 expressed NLRs reported to be under high levels of diversifying selection were regulated by FPA (chi^2 *p*=0.02) (Prigozhin and Krasileva, 2021). In addition, FPA-regulated NLRs tended to be located in regions with higher levels of synteny diversity (Jiao and Schneeberger, 2020), although in this case the association was not significant (*t*-test *p*=0.09; Figure 4–figure supplement 1). Overall, these findings suggest that FPA-dependent alternative polyadenylation is associated with rapidly evolving NLRs.

The effects of FPA regulation can be broadly classified into three modes of control involving (i) readthrough and chimeric RNAs, (ii) intronic poly(A) sites and (iii) poly(A) sites within protein-coding exons. At certain complex loci, FPA can affect poly(A) site choice using combinations of these different modes of regulation.

For 17 NLR genes, we found that a change in FPA activity altered the formation of readthrough or chimeric RNAs containing one or more NLR loci (Table 1). The duplicated *RPP7a/b*-like genes *AT1G58848* and *AT1G59218* (which form part of the *RPP7* cluster containing five CNL-class NLRs) displayed increased readthrough into downstream transposable elements (TEs) in *fpa-8* (Figure 4A). EMD tests could not be performed at these loci due to the multi-mapping of reads at these duplicated genes (*AT1G58848* and *AT1G59218*). Loss of FPA function can also lead to clusters of two or more NLR genes being co-transcribed as a single transcriptional unit. For example, the TNL-class gene *AT1G63730*, located in the *B4/RLM1* cluster, forms chimeric RNA with the downstream TNL-class gene *AT1G63740* in *fpa-8* (Helicos EMD=1099, FDR=0.02; Figure 4–figure supplement 2).

We identified another 17 NLR genes with intronic polyadenylation regulated by FPA (Table 2). Of these, four contained poly(A) sites in 5′UTR introns (which would result in non-coding transcripts) and three contained alternative poly(A) sites after the stop codon (which could alter potential regulatory sequences contained in 3′UTRs). The remainder contained poly(A) sites in introns between protein-coding exons. Selection of these poly(A) sites introduce premature stop codons that result in truncated open reading frames (ORFs). For

example, we identified a proximal poly(A) site within the third intron of *AT1G69550*, which
encodes a TNL-type singleton NLR (Figure 4B). Use of this poly(A) site results in mRNAs with
a premature stop codon; the encoded protein lacks most of the predicted LRR domain. In
fpa-8, readthrough at this poly(A) site is increased (Helicos EMD=1271, FDR=1.2×10^{-4}),
resulting in an increase in the relative number of full-length transcripts.
The most common form of FPA-dependent NLR regulation was premature
termination within exons (Table 3). We identified 45 NLRs controlled in this way: at 44 of
these loci, termination occurred within protein-coding exons. In most cases, this results in
stop-codonless transcripts that are predicted targets of non-stop decay (Szádeczky-Kardoss
et al., 2018). Many of these proximal exonic poly(A) sites could be identified at lower levels
in Col-0. For example, at *RPP28* (AT2G14080), which encodes a TNL-class singleton NLR,
we detected multiple exonic poly(A) sites located within the second and fourth exons, which
encode the NB-ARC and LRR domains, respectively (Figure 4C). Selection of these exonic
poly(A) sites was increased in *35S::FPA:YFP* (Helicos EMD=859, FDR=5.4×10^{-9}) and
decreased in *fpa-8* (Helicos EMD=912, FDR=7.6×10^{-9}). FPA was also found to promote
premature termination in the protein-coding sequence of single-exon, intronless NLR
genes. For example, at *RPP13* (AT3G46530), which encodes a CNL-class NLR protein, FPA
overexpression causes selection of proximal poly(A) sites located within the region
encoding the LRR domain (Helicos EMD=228, FDR=1.8×10^{-4}; Figure 4–figure
supplement 3).
Although the most frequent consequence of FPA selection of exonic poly(A) sites
was stop-codonless transcripts, we also identified examples where the protein-coding
potential was altered. For example, *AT5G40060* encodes a TNL-class NLR but has a
premature stop codon between the TIR and NB-ARC domains. Consequently, full-length
transcription results in an mRNA with an upstream ORF (uORF) encoding the TIR domain
and a larger downstream ORF encoding NB-ARC and LRR domains (Figure 4D). However,
transcripts with such large uORFs are targets of NMD in plants (Nyikó et al., 2009).
Therefore, FPA-regulated proximal polyadenylation in the region encoding the NB-ARC
domain results in a transcript containing only the uORF, which is not a predicted NMD target
and may be more efficiently translated into a TIR-only protein.
In seven of the identified genes, exonic proximal polyadenylation is associated with
retention of an upstream intron (Table 3). As a result, premature stop codons are
introduced, resulting in a truncated coding region. For example, the TNL-type NLR *RPS4*
was previously shown to be regulated by alternative splicing induced by the effector
AvrRps4 (Zhang and Gassmann, 2007). We identified an increase in *RPS4* intron 3 retention
in *35S::FPA:YFP* compared with Col-0 that was associated with proximal polyadenylation
events in exon 4 (Helicos EMD=34, not significant; Figure 4–figure supplement 4).
Therefore, inter-dependence between splicing and poly(A) site choice may explain *RPS4*
control.

FPA controlled NLR poly(A) site selection at 16 complex loci with combinations of
intronic, exonic and readthrough sites. One example is *RPP4* (AT4G16860), a TNL-class NLR
known to mediate Arabidopsis resistance to *Hpa* isolate Emoy2 (*Hpa*-Emoy2) (van der
Biezen et al., 2002). *RPP4* is part of the *RPP5* cluster, comprising seven TNL-class NLRs. In
agreement with a previous study (Wang and Warren, 2010), we found that in wild-type Col-
0, *RPP4* can be transcribed as a chimeric RNA together with the downstream *AtCOPIA4* TE
(AT4G16870) through selection of one of the two distal poly(A) sites located within the TE
(Figure 5) (Wang and Warren, 2010) or selection of a third poly(A) site in the downstream
gene *AT4G16857*. Use of the proximal poly(A) site within the TE is associated with an
approximately 8-kb cryptic splicing event between the 5′ splice site of the first exon of *RPP4*
and a 3′ splice site located within the TE. Both Nanopore DRS and Illumina RNA-Seq data
provide evidence for this cryptic splicing event, which skips all *RPP4* exons downstream of
exon 1, removing most of the *RPP4* coding sequence and introducing a stop codon
(Figure 5, Inset 1). The resulting transcript is predicted to encode a TIR-domain-only protein.
Loss of FPA function decreases chimeric RNA formation by shifting poly(A) site selection
towards a proximal poly(A) site located within the protein-coding region of the final exon
(Figure 5–figure supplement 1). This results in the production of *RPP4* transcripts lacking in-
frame stop codons (Figure 5, Inset 2). Furthermore, in *35S::FPA:YFP*, we observed increased
selection of a proximal poly(A) site located within the first intron of *RPP4*, which would also
encode a truncated RPP4 protein. We conclude that FPA-regulated alternative
polyadenylation at *RPP4* produces transcripts with unusually long 3′UTRs, alternative
protein isoforms and transcripts that cannot be efficiently translated.
448
FPA controls *RPP7* by promoting premature termination within protein-coding exon
6
To examine the functional impact of FPA on the regulation of NLRs, we focused on *RPP7*.
RPP7 encodes a CNL-class NLR protein crucial for strain-specific resistance to *Hpa* isolate
Hiks1 (*Hpa*-Hiks1) (Tsuchiya and Eulgem, 2013). The full-length expression of *RPP7* is
controlled by elongation factors that interact with H3K9me^2, which is associated with the
COPIA-type retrotransposon (*COPIA-R7*) located in *RPP7* intron 1 (Saze et al., 2013). Using
Nanopore and Helicos DRS data, we identified at least two poly(A) sites within the *COPIA-*
R7 element, both of which were selected more frequently in *fpa-8* (Figure 6A, Figure 6–
figure supplement 1). We also identified two poly(A) sites within the second intron of *RPP7*.
The use of both sites is reciprocally regulated by FPA, with a moderate decrease in *fpa-8*
and an increase in *35S::FPA:YFP*. All of these intronic proximal poly(A) sites are located
before the start of the *RPP7* ORF and generate transcripts that do not encode RPP7 protein.
At the 3′ end of *RPP7*, we found three alternative poly(A) sites located in the terminal intron,
in addition to the previously reported most distal and most commonly used poly(A) site in
the terminal exon (Figure 6A, Inset 1) (Tsuchiya and Eulgem, 2013). Selection of each of
these poly(A) sites is associated with alternative splicing events that lead to the generation

of four possible 3'UTR sequences. Termination at the 3'UTR intronic poly(A) sites is
suppressed by FPA: their usage is increased in *fpa-8* and decreased in *35S::FPA:YFP*. These
data indicate that FPA influences *RPP7* intronic polyadenylation at a larger number of
poly(A) sites than previously supposed.
The major effect of FPA on *RPP7* is within protein-coding exon 6, where we identified
three poly(A) sites (Figure 6A, Inset 2): two at the end of the region encoding the NB-ARC
domain and one within the region encoding the LRR repeats. Cleavage and polyadenylation
at these sites result in transcripts without in-frame stop codons, thereby disrupting the
coding potential of *RPP7* mRNA. These poly(A) sites were identified in both Helicos and
Nanopore DRS data, indicating that they are unlikely to be caused by alignment errors. The
relative selection of exon 6 poly(A) sites depends on FPA expression: in Col-0, 25% of *RPP7*
Nanopore DRS reads terminate at one of these exon 6 poly(A) sites; and when FPA is
overexpressed, this figure increases to 63%. Consistent with this, a relative drop in coverage
at exon 6 was also observed in *35S::FPA:YFP* Illumina RNA-Seq data. Consequently, only
23% of *RPP7* transcripts are expected to encode RPP7 protein in the FPA-overexpressing
line. In contrast, 4% of *RPP7* Nanopore DRS reads identified in *fpa-8* terminate in exon 6,
and 79% of transcripts are expected to be protein coding. In an orthogonal approach, we
used RNA gel blot analysis to visualise *RPP7* mRNAs in Col-0, *fpa-8* and *35S::FPA:YFP*
backgrounds and detected a clear decrease in signal corresponding to full-length *RPP7*
transcripts in *35S::FPA:YFP* (Figure 6B). These data support previous evidence of FPA-
dependent control of *RPP7* (Deremetz et al., 2019) but reveal that the predominant
mechanism is via exonic transcription termination.
488
FPA modulates *RPP7*-dependent, race-specific pathogen susceptibility
We next asked whether FPA-dependent premature transcription termination at *RPP7* exon
6 has a functional consequence. Since FPA reduced the level of full-length protein-coding
RPP7 transcripts, we asked whether increased FPA activity might compromise RPP7-
dependent immunity. To test this hypothesis, we carried out pathogenesis assays using the
oomycete strain *Hpa*-Hiks1. RPP7 provides race-specific immunity to *Hpa*-Hiks1 in Col-0.
The Keswick (Ksk-1) accession is susceptible to *Hpa*-Hiks1 (Lai et al., 2019) and we used it
as a control in these studies.
We inoculated Arabidopsis seedlings with *Hpa*-Hiks1 spores in three independent
experiments. Four days after inoculation, we checked susceptibility by counting the number
of sporangiophores. With the exception of Ksk-1, all of the lines we tested were in a Col-0
background. As expected, Col-0 plants were resistant to infection
(median: 0 sporangiophores per plant), and Ksk-1 plants were sensitive to infection
(median: 5 sporangiophores per plant; $p=1.7\times10^{-32}$; Figure 6C). *fpa-7* mutants were as
resistant to infection as Col-0 (median: 0 sporangiophores per plant, $p=0.19$). This is
consistent with our finding that full-length *RPP7* transcript expression is not reduced in the
absence of FPA. *fpa-8* mutants were also resistant to infection (median: 0 sporangiophores

per plant), however there was slight variability in their resistance compared to *fpa-7* ($p=2.4\times10^{-12}$). This variability was not restored by complementation with a *pFPA::FPA* transgene ($p=0.23$) indicating that it is not caused by loss of FPA function, and is likely to result from other mutations in the *fpa-8* background. In contrast, *35S::FPA:YFP* plants were significantly more sensitive to *Hpa*-Hiks1 than *pFPA::FPA* (median: 3 sporangiophores per plant; $p=3.8\times10^{-9}$), indicating that overexpression of FPA compromises immunity. We conclude that FPA-mediated transcription termination of *RPP7* pre-mRNA has a functional consequence for race-specific immunity. Therefore, FPA control of poly(A) site selection can modulate NLR function.

Discussion

We have identified a novel role for the RNA-binding protein FPA in the control of plant innate immunity. Using *IVI-MS* proteomics and ChIP-Seq, we showed that FPA is closely associated with proteins involved in RNA 3′ processing and co-localises with Ser[2] phosphorylated Pol II at the 3′ ends of genes. Integrative analysis using three RNA sequencing technologies confirmed that the major effect of modulating FPA activity is to alter poly(A) site selection. An unexpected finding was that half of expressed NLR loci were sensitive to FPA activity. In most cases, FPA promoted the use of poly(A) sites within protein-coding exons of NLR genes. At *RPP7*, an increase in exonic polyadenylation caused by FPA overexpression was shown to compromise immunity to *Hpa*-Hiks1. The widespread nature of this control mechanism suggests that transcription termination plays an important role in the regulatory and evolutionary dynamics of NLR genes.

Uncovering protein assemblies that mediate 3′ end processing in living plant cells
We used an *in vivo* formaldehyde cross-linking approach to identify proteins that co-localise with FPA inside living plant cells. These data provide in-depth knowledge of the proteins involved in Arabidopsis RNA 3′ end processing and clues to the function of the uncharacterised proteins identified here. Components of the m[6]A writer complex also co-purify with FPA. However, unlike related proteins in human and Drosophila (Knuckles et al., 2018; Patil et al., 2016), we found that FPA is not required to maintain global levels of m[6]A modification in Arabidopsis.

Two Arabidopsis PCF11 paralogs with Pol II CTD-interacting domains (CIDs), PCFS2 and PCFS4, co-purified with FPA, but two paralogs lacking CIDs, PCFS1 and PCFS5, did not. PCF11 was previously shown to have functionally separable roles in transcription termination and cleavage and polyadenylation (Sadowski et al., 2003): the N-terminal PCF11 CID is required for transcription termination, whereas the C-terminal domains are required for cleavage and polyadenylation. The specific interaction of FPA with CID-containing PCF11 paralogs suggests that FPA regulates alternative polyadenylation by altering Pol II speed and transcription termination. The human SPOC domain protein PHF3 can bind to two adjacent Ser[2] phosphorylated heptads of the CTD of Pol II via two

546 electropositive patches on the surface of its SPOC domain (Appel, Franke et al. 2020). One
547 of these patches, and the key amino acid residues within it, is conserved in the structure of
548 the FPA SPOC domain (Zhang et al., 2016). Consequently, FPA might also interact with the
549 CTD, possibly in conjunction with CID domains of PCFS2 and PCFS4. Such interactions
550 could explain the specific co-association of FPA and Pol II Ser^2 at the 3′ end of genes.
551
Widespread control of NLR transcription termination by FPA
An unanticipated finding of this study is that Arabidopsis NLR loci were the most enriched
group of FPA targets. NLRs function in the immune response and, consistent with this
crucial role, they are under powerful and dynamic selective pressure. Defining the inventory
of Arabidopsis NLRs depended on long-range DNA sequencing of diverse accessions (Van
de Weyer et al., 2019). Here we show that long-read Nanopore DRS provides insight into
the authentic complexity of NLR mRNA processing and enables the accurate annotation of
NLR genes. For example, our reannotation of the *RPS6* locus is essential to understand the
recurring role of RPS6 in autoimmunity. The autoimmune phenotypes of mutants defective
in NMD or the mitogen-activated kinase pathway are RPS6 dependent, but the mechanisms
involved are unclear (Gloggnitzer et al., 2014; Takagi et al., 2020). We found that *RPS6* is
transcribed through a previously unrecognised downstream gene that encodes an RPS6-
like TIR domain. We showed that expression of the downstream gene is dependent on the
RNA exosome component HEN2. In addition, mutations in *HEN2* were recently identified
as suppressors of RPS6-dependent autoimmune phenotypes (Takagi et al., 2020). It is clear
that accurate annotation of complex NLR loci facilitates the interpretation of basic features
of NLR function.
Of the 124 NLRs with detectable expression in Nanopore DRS data, 62 were
sensitive to FPA activity. FPA controls 3′ end formation of NLR genes in three different
transcript locations (Figure 7): (i) 3′UTRs, where it can prevent readthrough and chimeric
RNA formation; (ii) introns, where it promotes proximal polyadenylation; and (iii) protein-
coding exons, where it promotes stop-codonless transcript formation. The consequences
of such complex control of RNA 3′ end formation are wide-ranging and likely to be context
dependent (Mayr, 2019).
Where FPA controls readthrough and chimeric RNA formation, it affects 3′UTR
length, sequence composition and cryptic splice site usage. Long or intron-containing
3′UTRs are targeted by NMD, leading to RNA decay or suppressed translatability. Long,
unstructured 3′UTRs influence intermolecular RNA interactions and phase separation,
changing the subcellular localisation of mRNAs (Ma et al., 2021). The close proximity of
mRNAs in the resulting granules may enable co-translational protein complex formation.
Readthrough transcription may also disrupt the expression of downstream genes by
transcription interference (Proudfoot, 1986).
FPA-dependent premature transcription termination at intronic poly(A) sites can
introduce novel stop codons, resulting in transcripts that encode truncated NLR proteins

with altered functions. For example, some TIR domain-only proteins are known to be active
in NLR regulation, resulting in constitutive signalling activity (Zhang et al., 2004) or act as
competitive inhibitors by titrating full-length NLR protein (Williams et al., 2014). In other
cases, TIR-domain-only proteins are sufficient for pathogen recognition (Nishimura et al.,
2017). The TE-containing 3′UTR of *RPP4* appears to be required for resistance to the
pathogen *Hpa*-Emoy2, although the mechanism involved is unclear (Wang and Warren,
2010). We discovered that cryptic splicing of *RPP4* exon 1 to a novel 3′ splice site within the
TE can produce a unique transcript that encodes only the RPP4 TIR domain. It will be
interesting to examine whether the TIR-only RPP4 isoform is required for full pathogen
resistance. We also found that intron retention at *RPS4,* which is essential for RPS4-
dependent resistance against *P. syringae* DC3000 (Zhang and Gassmann, 2007), is linked
to exonic proximal polyadenylation. Intron retention without accompanying proximal
polyadenylation will result in transcripts with long 3′UTRs that are likely to be sensitive to
NMD, whereas proximally polyadenylated transcripts could be translated into truncated
protein. Therefore, a combination of alternative polyadenylation and splicing probably
underpins *RPS4* control. In future, sensitive proteomic analyses will be important to
determine the impact of alternative polyadenylation on NLR protein isoform expression.
A remarkable finding was that FPA mostly targets the protein-coding exons of NLR
genes and, even controls premature transcription termination within the ORF of single-exon
NLR genes such as *RPP13*. Premature transcription termination in protein-coding exons
results in the production of stop-codonless transcripts that cannot be efficiently translated
into protein. These truncated transcripts may be subject to decay by RNA surveillance
mechanisms (e.g. the non-stop decay pathway) or act as non-coding RNA decoys to titrate
the levels of regulatory microRNAs (Shivaprasad et al., 2012). Increased rates of NLR
transcription in plants under pathogen attack could promote elongation through such
"regulatory" poly(A) sites. In this way, the expression of NLR proteins might be restricted
during pathogen surveillance but kept poised for rapid activation during infection.
Since the evolution of *cis*-regulatory elements controlling poly(A) site choice within
introns or 3′UTRs is free from the constraints of protein-coding functionality, why should
protein-coding exons be targeted so frequently? One possibility is that this enables the
expression of newly created NLR genes to be kept under tight control, thereby facilitating
rapid evolution whilst reducing the chances of autoimmunity (Figure 7). This hypothesis is
strengthened by the finding that many NLRs with high allelic diversity (Prigozhin and
Krasileva, 2021) are regulated by FPA. Alternative polyadenylation might also function to
hide NLR genes from negative selection and contribute to cryptic genetic variation in a
similar way to the mechanism proposed for NMD- and microRNA-mediated NLR control
(Raxwal and Riha, 2016; Shivaprasad et al., 2012). Cryptically spliced chimeric RNAs, with
subsequent retrotransposition, can be a source of new genes (Akiva et al., 2006). Therefore,
the control of transcription termination could directly facilitate the neofunctionalisation of
NLRs. In the future, it will be important to compare patterns of transcription termination at

NLRs across Arabidopsis accessions. For example, analysis of transcriptomic data will
determine whether proximal polyadenylation is conserved in NLRs with high allelic
diversity, whilst an integrative analysis of transcriptomic and genomic data will establish
whether chimeric NLR transcripts identified in some accessions are found as
retrotransposed genes in others.

At least two distinct patterns of alternative polyadenylation mediate *RPP7*
regulation, one involving intronic heterochromatin (Tsuchiya and Eulgem, 2013) and
another involving FPA-dependent termination in exon 6. The latter mechanism is conserved
across all NLR genes of the Col-0 *RPP7* locus (Table 3). Alleles of these RPP7-like NLR genes
have been identified as the causes of specific cases of hybrid necrosis (Bomblies and
Weigel, 2007). In these cases, autoimmunity is explained by allele-specific physical
interactions between RPP7 protein and the RPW8-only protein HR4 (Li et al., 2020). We
found that not only are *RPP7*-like genes targeted by FPA-dependent premature transcript
termination, but so too is *HR4* (Table 2). This raises the possibility that FPA could rescue
hybrid necrosis by limiting the expression of these proteins. FPA also appears to regulate
the proximal polyadenylation of *DANGEROUS MIX 10 (DM10)*, producing transcripts that
could encode a protein with truncated LRR repeats. *DM10* alleles with LRR truncations have
been demonstrated to cause autoimmunity in specific crosses (Barragan et al., 2021),
suggesting that in other cases FPA overexpression could trigger or enhance autoimmune
phenotypes. Consequently, modulation of transcription termination may shift the balance
of costs and benefits associated with NLR gene expression. This phenomenon is not likely
to be restricted to FPA because mutations in the RNA 3′ processing factor CPSF30 can also
suppress autoimmunity (Bruggeman et al., 2014).

The impact of FPA overexpression on gene expression and autoimmunity revealed
here derives from artificial transgene expression. However, pathogens could similarly
modulate NLR activity by evolving effectors that target the expression or activity of factors
controlling NLR poly(A) site choice. Consistent with this idea, the HopU1 effector of
P. syringae targets the RNA-binding protein AtGRP7 (Fu et al., 2007), which co-purified with
FPA. In addition, the Pi4089 effector of the oomycete pathogen *Phytophthora infestan*s
targets the KH-domain RNA-binding protein StKRBP1 in potato; as a result, the abundance
of StKRBP1 increases and infection by *P. infestans* is enhanced (Wang et al., 2015). This
precedent reveals that effector-mediated increases in RNA-binding protein abundance can
transform host RNA-binding proteins into susceptibility factors. Phylogenetic analysis of
StKRBP1 suggests that a direct homolog is absent in Brassicaceae. However, the most
closely related Arabidopsis proteins are FLK and PEP (Zhang et al., 2020a), both of which
co-purify with FPA and have been shown to regulate poly(A) site choice (Rodriguez-Cazorla
et al., 2015). FPA, GRP7, FLK and PEP, along with other RNA-binding proteins, act in concert
to fine-tune the timing of flowering through the regulation of *FLC*. In a similar way, RNA-
binding protein-dependent modulation of NLR expression might explain how quantitative
disease resistance occurs (Corwin and Kliebenstein, 2017).

New ways to analyse RNA processing

An essential feature of our study was the introduction of new approaches to study RNA processing and 3' end formation. The use of long-read Nanopore DRS transformed our understanding of the complexity of NLR gene expression by providing insight that short-read Illumina RNA-Seq and Helicos DRS could not. We recently showed that Nanopore DRS mapping of RNA 3' ends closely agrees with short-read Helicos DRS, and that Nanopore DRS is not compromised by internal priming artefacts (Parker et al., 2020). Consequently, we used Nanopore DRS to quantify alternative patterns of cleavage and polyadenylation. We also introduced a new approach to analyse alternative polyadenylation by applying the EMD metric. EMD incorporates information on the both the relative abundance and the genomic distance between alternative poly(A) sites. This is valuable because large distances between poly(A) sites are more likely to impact the mRNA coding potential or trigger mRNA surveillance compared with subtle changes in 3'UTR length.

A limitation of short-read analyses of RNA processing is their dependence upon reference transcript annotations because these may be incomplete. For example, in disease or mutant conditions, RNA processing often occurs at novel sites that are not present in reference transcriptomes (as was the case here for NLR genes). For this reason, using long-read sequencing data to generate bespoke reference transcriptomes for the genotypes under analysis can increase the value of short-read sequencing data. Until the throughput of long-read sequencing matches that of short-read technologies, a combined approach is likely to be generally useful in interpreting transcriptomes.

Concluding remarks

It is difficult to identify alternative polyadenylation from conventional short-read RNA-Seq data. As a result, the impact of alternative polyadenylation is probably under-reported. Here we show that premature transcription termination of NLR genes is widespread. Using Nanopore DRS, we could improve the accuracy of NLR annotation and revealed a layer of NLR gene regulation that may also influence the dynamics of NLR evolution. The continued development of approaches that reveal full-length native RNA molecules is likely to provide new insight into other important, but previously unrecognised, aspects of biology.

Materials and methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain (*Arabidopsis thaliana*)	Columbia (Col-0)	NA	ABRC: CS22625	Country of Origin: USA
Strain (Arabidopsis thaliana)	Keswick (Ksk-1)	Lai et al. 2018	ABRC: CS1634	Country of Origin: UK
Gene (*Arabidopsis thaliana)*	*FPA*	NA	TAIR/ABRC: AT2G43410	-
Gene (Arabidopsis thaliana)	*RPP7*	NA	TAIR/ABRC: AT1G58602	-
Genetic reagent (*Arabidopsis thaliana*)	*fpa-7*	Duc et al., 2013	ABRC: SALK_021959C	T-DNA insertion mutant in Col-0 background. Gifted by R. Amasino, UW-Madison.
Genetic reagent (*Arabidopsis thaliana*)	*fpa-8*	Baurle et al., 2007	TAIR: 4515120225	EMS point mutation in Col-0 background. Gifted by C. Dean, John Innes Centre
Genetic reagent (*Arabidopsis thaliana*)	*35S::FPA:YFP fpa-8*	Baurle et al., 2007	NA	Transgenic line in *fpa-8* background*,* gifted by C. Dean, John Innes Centre
Genetic reagent (*Arabidopsis thaliana*)	*pFPA::FPA fpa-8*	Zhang et al., 2016	NA	Transgenic line in *fpa-8* background.
Genetic reagent (*Arabidopsis thaliana*)	*vir-1*	Růžička et al., 2017	TAIR: 6532672723	EMS point mutant in Col-0 background. Gifted by K. Růžička, Brno.
Commercial assay, kit	Rneasy Plant Mini kit	QIAGEN	Cat#: 74904	-
Commercial assay, kit	SuperScript™ III Reverse Transcriptase	Thermo Fisher Scientific	Cat#: 18080044	-
Commercial assay, kit	NEBNext® Ultra™ Directional RNA Library Prep Kit for Illumina®	New England Biolabs	Cat#: E7420	-
Commercial assay, kit	Dynabeads™ mRNA Purification Kit	Thermo Fisher Scientific	Cat#: 61006	-
Commercial assay, kit	Nanopore Direct RNA sequencing kit	Oxford Nanopore Technologies	Cat#: SQK-RNA001	-
Commercial assay, kit	MinION Flow cell r9.4	Oxford Nanopore Technologies	Cat#: FLO-MIN106	-
Peptide, recombinant protein	T4 DNA ligase	New England Biolabs	Cat#: M0202	-

Commercial assay, kit	Quick Ligase reaction buffer	New England Biolabs	Cat#: B6058S	-
Commercial assay, kit	Agencourt RNAclean XP magnetic beads	Beckman Coulter	Cat#: A63987	-
Commercial assay, kit	Qubit RNA BR Assay Kit	Thermo Fisher Scientific	Cat#: Q10210	-
Commercial assay or kit	RNA ScreenTape System	Agilent	Cat#: 5067-5576 - 5067-5578	-
Antibody	FPA antibody	Covance	NA	Rabbit polyclonal antibody. Raised against FPA amino acids 536-901.
Chemical compound	[γ-32P]-ATP	Perkin Elmer	Cat#: BLU012H250UC	-
Commercial assay or kit	DECAprime II DNA labelling kit	Thermo Fisher Scientific	Cat#: AM1455	-
Commercial assay or kit	Illustra MicroSpin G-50 Columns	GE Healthcare	Cat#: 27-5330-01	-
Commercial assay or kit	RiboRuler High Range RNA Ladder	Thermo Fisher Scientific	Cat#: SM1821	-
Peptide, recombinant protein	FastAP Thermosensitive Alkaline Phosphatase	Thermo Fisher Scientific	Cat#: EF0651	-
Peptide, recombinant protein	T4 Polynucleotide Kinase	Thermo Fisher Scientific	Cat#: EK0031	-
Peptide, recombinant protein	Nuclease P1	Merck	Cat#: N8630-1VL	-
Peptide, recombinant protein	Calf Intestinal Alkaline Phosphatase	New England Biolabs	Cat#: M0290S	-
Chemical compound	N6-Methyladenosine (m6A), Modified adenosine analog	Abcam	Cat#: ab145715	-
Chemical compound	Adenosine, Endogenous P1 receptor agonist	Abcam	Cat#: ab120498	-
Commercial assay or kit	GFP-Trap Agarose	Chromotek	Cat#: gta-20	-
Software, algorithm	d3pendr	10.5281/zenodo.4319112	NA	Scripts to perform differential 3' end analysis using Nanopore DRS or Helicos DRS data
Software, algorithm	Simpson_Barton_FPA_NLRs	10.5281/zenodo.4319108	NA	All pipelines, scripts and notebooks used for analyses in this manuscript.

Plants

Plant material and growth conditions

The wild-type Col-0 accession and *fpa-7* were obtained from the Nottingham Arabidopsis

Stock Centre. The *fpa-8* mutant (Col-0 background) and *35S::FPA:YFP* in *fpa-8* (Baurle et

al., 2007) were provided by C. Dean (John Innes Centre). Generation of the *pFPA::FPA* line

was previously described (Zhang et al., 2016). Surface-sterilised seeds were sown on MS10
medium plates containing 2% agar, stratified at 4°C for 2 days, germinated in a controlled
environment at 20°C under 16-h light/8-h dark conditions and harvested 14 days after
transfer to 20°C.
710
IVI-MS
Preparation of IVI-MS samples
Seedlings were harvested 14 days after germination and cross-linked with 1 % (v/v)
formaldehyde under vacuum. The cross-linking reaction was stopped after 15 min by the
addition of glycine to a final concentration of 0.125 M and returned to vacuum for a further
5 min. Nuclei were isolated from frozen ground plant tissue using Honda buffer (20 mM
Hepes-KOH pH 7.4, 10 mM $MgCl_2$, 440 mM sucrose, 1.25 % (w/v) Ficoll, 2.5 % (w/v) Dextran
T40, 0.5 % (v/v) Triton X-100, 5 mM DTT, 1 mM PMSF, 1% (v/v) Protease Inhibitor Cocktail;
(Sigma)) and collected by centrifugation at 2000*g* for 17 min at 4°C. Nuclei were washed
twice with Honda buffer (centrifugation at 1500*g* for 15 min at 4°C between washes) and
lysed in nuclear lysis buffer (50 mM Tris-HCl pH 8, 10 mM EDTA, 1 % (w/v) SDS, 1 mM PMSF,
1 % (v/v) Protease Inhibitor Cocktail) by sonication for four cycles of 30 s pulses with low
power and 60 s cooling between pulses using a Bioruptor UCD-200 (Diagenode).
Following centrifugation (16,100*g* for 10 min at 4°C), the supernatant was diluted 10-fold
with sample dilution buffer (16.7 mM Tris-HCl pH 8, 167 mM NaCl, 1.1 % (v/v) Triton X-100,
1 % (v/v) Protease Inhibitor Cocktail). Cross-linked protein complexes were isolated with
GFP-trap agarose beads (Chromotek) and incubated at 4°C with constant rotation for 5 h,
followed by centrifugation (141*g* for 3 min at 4°C). Beads were washed three times with
washing buffer (150 mM NaCl, 20 mM Tris-HCl pH 8, 2 mM EDTA pH 8, 1 % (v/v) Triton X-
100, 0.1 % (w/v) SDS, 1 mM PMSF) by centrifugations between washes (400*g* for 3 min at
4°C). Samples were incubated at 90°C for 30 min to reverse the cross-linking prior to SDS-
PAGE. Each biological replicate was separated into five fractions following SDS-PAGE,
subjected to in-gel digestion with trypsin and submitted for LC-MS/MS analysis (LTQ
Orbitrap Velos Pro mass spectrometer; Thermo Fisher Scientific). Three biological
replicates were performed for each genotype.
736
IVI-MS data analysis
Raw peptide data files from *IVI-MS* were analysed by MaxQuant software (version 1.6.10.43)
(Cox and Mann, 2008). Peptide tables were then loaded using Proteus (version 0.2.14)
(Gierlinski et al., 2018) and summarised to protein level counts using the hi-flyer method
(mean of the top three most abundant peptides). Because wild-type plants lacking GFP
were used as controls, a large number of the proteins enriched by immunoprecipitation
were below the detection threshold in the control. This group of proteins can be classified
as "missing not at random" (MNAR). In all proteomics experiments, there will also be a
number of proteins which are not detected purely by chance: these are referred to as

"missing at random" (MAR). We treated proteins that were missing from all replicates of a condition as MNAR, and proteins that were missing only from a subset of replicates as MAR. Using the imputeLCMD package (version 2.0) (Lazar, 2015), a K nearest neighbours' strategy was used to impute MAR examples, and a quantile regression imputation of left centred data (QRILC) approach was used to impute MNAR examples. Differential expression analysis was performed on imputed data using limma (version 3.40.0) (Ritchie et al., 2015). Because imputation is not deterministic (i.e. will lead to different outcomes every time), we improved the robustness of the analysis by performing 999 bootstraps of the imputation and differential expression, and summarising the results using the median $\log_2$ fold change and harmonic mean p value.

ChIP-Seq

Preparation of libraries for ChIP-Seq

ChIP against FPA and Pol II phosphorylated at either Ser^5 or Ser^2 of the CTD heptad repeat was performed as previously described (Yu et al., 2019). Polyclonal antibodies against FPA amino acids 536–901 were raised in rabbit by Covance.

ChIP-Seq data processing

FPA and Pol II ChIP-Seq data are available at ENA accession PRJNA449914. $H3K9me^2$ ChIP-Seq data were downloaded from ENA accessions PRJDB5192 (Inagaki et al., 2017) and PRJNA427432 (Lai et al., 2020). Reads were aligned to the TAIR10 reference genome using Bowtie2 (version 2.3.5.1) (Langmead and Salzberg, 2012) with the parameters –mm –very-sensitive –maxins 800 –no-mixed –no-discordant. Counts per million normalised coverage profiles were generated using deepTools (version 3.4.3) (Ramirez et al., 2014). For 3′ end centred metagene profiles, we determined the major 3′ position per gene using the Araport11 annotation and existing Col-0 Helicos DRS data (Sherstnev et al., 2012). Metagenes centred on these positions were then generated in Python 3.6 using pyBigWig (version 0.3.17) (Ramirez et al., 2014), Numpy (version 1.18.1) (Harris et al., 2020) and Matplotlib (version 3.1.3) (Hunter, 2007). For differential $H3K9me^2$ analysis, read counts per gene (including intronic regions) were generated using pysam (version 0.16.0), and differential expression analysis was performed using edgeR (version 3.22.5) (Robinson et al., 2010).

RNA

Total RNA isolation

Total RNA was isolated using RNeasy Plant Mini kit (QIAGEN) and treated with TURBO DNase (Thermo Fisher Scientific) according to the manufacturers' instructions. The total RNA concentration was measured using a Qubit 1.0 Fluorometer and Qubit RNA BR Assay Kit (Thermo Fisher Scientific), whilst RNA quality and integrity was assessed using a

785 NanoDrop™ 2000 spectrophotometer (Thermo Fisher Scientific) and Agilent 2200
786 TapeStation System (Agilent).
787
Nanopore DRS
Preparation of libraries for DRS using nanopores
Total RNA was isolated from Col-0, *fpa-8* and *35S::FPA:YFP* seedlings as described above.
mRNA was isolated and Nanopore DRS libraries prepared (using the SQK-RNA001
Nanopore DRS Kit; Oxford Nanopore Technologies) as previously described (Parker et al.,
2020). Libraries were loaded onto R9.4 flow cells (Oxford Nanopore Technologies) and
sequenced using a 48-h runtime. Four biological replicates were performed for each
genotype.
796
Nanopore DRS data processing
Nanopore DRS reads were basecalled using the Guppy (version 3.6.0) high accuracy
model. Reads were mapped to the Arabidopsis *TAIR10* genome (The Arabidopsis
Genome, 2000) using minimap2 (version 2.17) with the parameters -a -L --cs=short -x splice
-G20000 --end-seed-pen=12 --junc-bonus=12 -uf. Spliced alignment was guided using
junctions from the Araport11 annotation (Cheng et al., 2017). Nanopore DRS reads can
suffer from "oversplitting" – where the signal originating from a single RNA molecule is
incorrectly interpreted as two or more reads (Parker et al., 2020). These errors can be
systematic and result in false positive 3′ ends. To filter these errors, we identified reads that
were sequenced consecutively through the same pore and also mapped contiguously on
the genome (within 1 kb of each other). In this way, we filtered all except the most 3′ reads,
which should contain the genuine RNA 3′ end. Pipelines for processing Nanopore DRS data
were built using Snakemake (Koster and Rahmann, 2012).
810
Helicos DRS
Preparation of samples for Helicos DRS
Total RNA was isolated from the Col-0, *fpa-8* and *35S::FPA:YFP* seedlings as described
above. Samples were processed by Helicos BioSciences as previously described (Ozsolak
et al., 2009; Sherstnev et al., 2012). Three biological replicates were performed for each
genotype.
817
Helicos DRS data processing
Helicos DRS reads were mapped to the *Arabidopsis* TAIR10 genome using Heliosphere
(version 1.1.498.63) as previously described (Sherstnev et al., 2012). Reads were filtered to
remove those with insertions or deletions of >4 nt and to mask regions with low complexity,
as determined using DustMasker (Camacho et al., 2009) (from BLAST+ suite version 2.10.1)
set at DUST level 15 (Sherstnev et al., 2012).
824

Differential 3′ end analysis of Nanopore and Helicos DRS datasets

Transcriptional loci were first identified in Col-0, *fpa-8* and *35S::FPA:YFP* Nanopore DRS reads using the long-read transcript assembly tool StringTie2 version 2.1.1 (Pertea et al., 2015). Novel transcriptional loci were merged with annotated loci from the Araport11 reference (Cheng et al., 2017). To detect sites with altered 3′ end distributions in *fpa-8* and *35S::FPA:YFP* mutants, we pooled the replicates of either Nanopore or Helicos DRS data and identified reads overlapping each transcriptional locus. These reads were used to build distributions of 3′ end locations. The difference in 3′ end distributions between the treatment and control (Col-0) was measured using EMD. To identify loci with statistically significant differences in 3′ distributions, we performed an EMD permutation test using 999 bootstraps: for this, reads for each locus were randomly shuffled between the treatment and control samples to create null distributions, and the EMD recalculated. The histogram of null EMDs was fitted using a gamma distribution, and the *p*-value (probability of achieving the observed EMD or greater by chance) was calculated from the distribution. *P*-values were corrected for multiple testing using the Benjamini–Hochberg method. Genes with an EMD>25 and an FDR<0.05 were considered to be differentially alternatively polyadenylated, and the directionality of change was identified using the difference in mean 3′ position. Software developed to perform differential 3′ analysis is available on GitHub at https://github.com/bartongroup/d3pendr and 10.5281/zenodo.4319113, and can be used with Nanopore DRS, Helicos DRS, or Illumina 3′ tag-based datasets.

Illumina RNA sequencing

Preparation of libraries for Illumina RNA sequencing

Total RNA was isolated from the Col-0, *fpa-8* and *35S::FPA:YFP* seedlings as described above. mRNA was isolated and sequencing libraries prepared using the NEBNext Ultra Directional RNA Library Prep Kit for Illumina (New England Biolabs) by the Centre for Genomic Research (University of Liverpool). 150-bp paired-end sequencing was carried out on Illumina HiSeq 4000. Six biological replicates were performed for each genotype.

Illumina RNA sequencing data processing

Illumina RNA-Seq data were assessed for quality using FastQC (version 0.11.9) and MultiQC (version 1.8) (Andrews, 2010; Ewels et al., 2016). Reads were mapped to the TAIR10 genome using STAR (version 2.7.3a) (Dobin et al., 2013) with a splice junction database generated from the Araport11 reference annotation (Cheng et al., 2017). Counts per million normalised coverage tracks were created using samtools (version 1.10) and deepTools (version 3.4.3) (Ramirez et al., 2014). To identify expressed regions in each locus, the coverage profiles of each treatment and control replicate were first extracted using pyBigWig (version 0.3.17) (Ramirez et al., 2014). These were normalised such that the area under each profile was equal to the mean area under the profiles. A normalised coverage threshold of 1 was used to identify expressed regions of the loci. These regions were further

segmented when at least two-fold differences in expression within a 25-nt window were
found between control and treatment conditions (and then regions smaller than 50 nt
removed). Expression of the segmented regions was then calculated using featureCounts
(version 2.0.0) (Liao et al., 2013). Each read pair was counted as one fragment, and only
properly paired, concordant and primary read pairs were considered. Differential usage
within transcriptional loci was assessed using DEXSeq (version 1.32.0) (Reyes et al., 2013).
Loci were considered to be differentially processed if they had a locus-level FDR<0.05 and
at least one region with an absolute logFC>1 and FDR<0.05. For differential splice junction
usage analysis, counts of splice junctions annotated in the bespoke Nanopore DRS-derived
annotation, plus Araport11 annotation, were generated for each locus using pysam (version
0.16.0). Differential splice junction usage was assessed using DEXSeq (version 1.32.0)
(Reyes et al., 2013). Loci were considered to be differentially spliced if they had a locus-
level FDR<0.05 and at least one junction with an absolute logFC>1 and FDR<0.05.
878
Gene tracks
Gene track figures were generated in Python 3.6 using Matplotlib (version 3.1.3) (Hunter,
2007). For gene tracks where any condition had >200 Nanopore DRS read alignments, 200
representative alignments were selected by random sampling without replacement (except
for the *FPA* gene track figure, where 500 read alignments were sampled). NanoPARE data
(Schon et al., 2018) were processed as previously described (Parker et al., 2020). For
reannotated gene loci, domains were predicted using the InterproScan web client (Mitchell
et al., 2019) and LRRs were predicted using LRRpredictor web client (Martin et al., 2020).
Protein alignments were created and visualised in Jalview (version 2.11) (Waterhouse et al.,
2009) using T-Coffee (Notredame et al., 2000).
889
Protein domain family enrichment analysis
To conduct protein domain enrichment analysis, InterPro domain annotations of
Arabidopsis proteins were downloaded from BioMart (Smedley et al., 2009) and converted
to genomic co-ordinates using the Araport11 annotation (Cheng et al., 2017). Domain
families overlapping each locus tested for alternative polyadenylation using either
Nanopore or Helicos DRS were identified using pybedtools (version 0.8.1) (Dale et al.,
2011). To identify enriched domain families, domains were randomly shuffled between
tested loci in 19,999 bootstraps, and the number of times that each domain class
overlapped by chance with significantly alternatively polyadenylated loci was recorded.
This was compared with the observed overlap of each domain family with alternatively
polyadenylated loci to calculate *p*-values, which were corrected for multiple testing using
the Benjamini–Hochberg method.
902

Manual annotation of alternatively polyadenylated NLR genes

To identify which of the 206 previously annotated NLR genes present in Col-0 were alternatively polyadenylated in *fpa-8* and *35S::FPA*, we devised a standard operating procedure for visual inspection. Genes that had Nanopore DRS read coverage in at least two conditions were considered to be expressed. Genes were considered to be alternatively polyadenylated if they had multiple 3′ end locations with each supported by at least four Nanopore DRS reads, and if there was a clear difference in Nanopore DRS coverage in the treatment condition compared with Col-0. Helicos and Illumina corroboration of poly(A) sites and coverage changes was also taken into consideration.

Genomic organisation of alternatively polyadenylated NLR genes

To test whether expressed NLR genes with FPA-dependent alternative polyadenylation were associated with NLR gene clusters, we used previously produced cluster assignments for Col-0 NLR genes (Lee and Chae, 2020). We also tested the association of FPA-dependent alternative polyadenylation with previously produced hypervariable NLR classifications (Prigozhin and Krasileva, 2021). The association of alternatively polyadenylated genes with both major NLR gene clusters and hypervariable NLRs was assessed using a Chi squared test. To test whether FPA-regulated NLRs are found in regions with high synteny diversity, we used 5-kb sliding window estimates of synteny diversity calculated from seven diverse Arabidopsis ecotypes (Jiao and Schneeberger, 2020). For each expressed NLR gene, the window with the largest overlap was used as the estimate of synteny diversity. The association with alternatively polyadenylated genes was assessed using a *t*-test.

RNA gel blot analysis of *RPP7* mRNAs

RNA gel blot analysis was carried out as previously described (Quesada et al., 2003) with minor modifications. *RPP7* mRNA was detected using a probe annealing to the second exon of the *RPP7* (*AT1G58602*) gene (200 bp PCR product amplified with the following primers: Forward: 5′-TCGGGGACTACTACTACTCAAGA-3′ and Reverse: 5′-TCTTGATGGTGTGAAAGAATCTAGT-3′). *Beta-TUBULIN* mRNA was used as a loading control and visualised by a probe annealing to the third exon of the *Beta-TUBULIN (AT1G20010) gene* (550 bp PCR product amplified with the following primers: Forward: 5′-CTGACCTCAGGAAACTCGCG-3′ and Reverse: 5′- CATCAGCAGTAGCATCTTGG-3′). The probes were 5′ labelled using [γ-^{32}P]-ATP (Perkin Elmer) and DECAprime™ II DNA labelling kit (Thermo Fisher Scientific) and purified on illustra G-50 columns (GE Healthcare Life Sciences). mRNA isoforms were visualised and quantified using an Amersham Typhoon Gel and Blot Imaging System (GE Healthcare Bio-Sciences AB). The RiboRuler High Range RNA Ladder (Thermo Fisher Scientific), used to identify the approximate size of RNA bands, was first dephosphorylated using FastAP Thermosensitive Alkaline Phosphatase (Thermo Fisher

942 Scientific) and then labelled with [γ-^{32}P]-ATP (Perkin Elmer) using T4 Polynucleotide Kinase
943 (Thermo Fisher Scientific) before gel loading.
944
m^{6}A LC-MS/MS
Total RNA was isolated and checked as described above. mRNA was extracted twice from
approximately 75 µg of total RNA using the Dynabeads mRNA Purification Kit (Thermo
Fisher Scientific) according to the manufacturer's instructions. The quality and quantity of
mRNA was assessed using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific)
and Agilent 2200 TapeStation System (Agilent). Samples for m^{6}A LC-MS/MS were prepared
as previously described (Huang et al., 2018) with several modifications. First, 100 ng mRNA
was diluted in a total volume of 14 ml nuclease-free water (Thermo Fisher Scientific) and
digested by nuclease P1 (1 U, Merck) in 25 µl buffer containing 20 mM NH$_4$Oac (pH 5.3) at
42°C for 2 h. Next, 3 µl freshly made 1 M NH$_4$HCO$_3$ and calf intestinal alkaline phosphatase
(1 U, New England Biolabs) were added, and samples were incubated at 37°C for 2 h. The
samples were then diluted to 50 µl with nuclease-free water and filtered (0.22 µm pore size,
4 mm diameter; Millipore). LC-MS/MS was carried out by the FingerPrints Proteomics
facility at the University of Dundee. m^{6}A/A ratio quantification was performed in
comparison with the curves obtained from pure adenosine (endogenous P1 receptor
agonist, Abcam) and m^{6}A (modified adenosine analog, Abcam) nucleoside standards.
Statistical analysis was performed using a 2-way *t*-test.
962
Pathogenesis assays
Pathogenesis assays were carried out as previously described (Tome et al., 2014). The *Hpa*-
Hiks1 isolate was maintained by weekly sub-culturing on Ksk-1 plants. A solution containing
Hpa-Hiks1 spores was used to inoculate 14-day-old Col-0, Ksk-1, *fpa-7, fpa-8, pFPA::FPA*
and *35S::FPA:YFP* seedlings. Sporangiophores were counted 4 days after inoculation. The
experiment was repeated three times using up to 45 plants per genotype per each repeat.
Statistical analysis was performed with negative binomial regression using Statsmodels
(version 0.11.0) (Seabold and Perktold, 2010), plants were grouped by experimental repeat
during testing to control for variation between repeats.
972
Code availability
All pipelines, scripts and notebooks used to generate figures are available from GitHub at
https://github.com/bartongroup/Simpson_Barton_FPA_NLRs and Zenodo at
10.5281/zenodo.4319108. The software tool developed for detecting changes in poly(A)
site choice in Nanopore and Helicos DRS data are available from GitHub at
https://github.com/bartongroup/d3pendr and Zenodo at 10.5281/zenodo.4319112.
979

Data availability

IVI-MS data are available from the ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2019) partner repository, with the dataset identifier PXD022684. FPA and Pol II ChIP-Seq data are available from ENA accession PRJNA449914 (Yu et al., 2019). Col-0 Nanopore DRS data are available from ENA accession PRJEB32782 (Parker et al., 2020). *fpa-8* and *35S::FPA:YFP* Nanopore DRS data are available from ENA accession PRJEB41451. *Hen2-2* Nanopore DRS data are available from ENA accession PRJEB41381. Col-0, *fpa-8* and *35S::FPA:YFP* Helicos DRS data are available from Zenodo DOI 10.5281/zenodo.4309752. Col-0, *fpa-8* and *35S::FPA:YFP* Illumina RNA-Seq data are available from ENA accession PRJEB41455.

Acknowledgments

We thank Paul Birch and Ingo Hein for comments on the manuscript and David Baulcombe, Ian Henderson and Wenbo Ma for helpful NLR discussions. We thank Abdelmadjid Atrih (Centre for Advanced Scientific Technologies, School of Life Sciences) for the m^{6}A LC-MS/MS analysis.

Competing interests

The authors declare no competing interests.

Funding

This work was supported by awards from the BBSRC (BB/M010066/1; BB/J00247X/1; BB/M004155/1), the University of Dundee Global Challenges Research Fund, a University of Dundee PhD studentship to V.Z. and a European Union Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant agreement No. 799300 to K.K. The FingerPrints Proteomics Facility of the University of Dundee is supported by a Wellcome Trust Technology Platform Award (097945/B/11/Z).

References

Adachi, H., Derevnina, L., and Kamoun, S. (2019). NLR singletons, pairs, and networks: evolution, assembly, and regulation of the intracellular immunoreceptor circuitry of plants. Curr Opin Plant Biol *50*, 121-131.

Akiva, P., Toporik, A., Edelheit, S., Peretz, Y., Diber, A., Shemesh, R., Novik, A., and Sorek, R. (2006). Transcription-mediated gene fusion in the human genome. Genome Res *16*, 30-36.

Anders, S., Reyes, A., and Huber, W. (2012). Detecting differential usage of exons from RNA-seq data. Genome Res *22*, 2008-2017.

Andres, F., and Coupland, G. (2012). The genetic basis of flowering responses to seasonal cues. Nat Rev Genet *13*, 627-639.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

Ariyoshi, M., and Schwabe, J.W. (2003). A conserved structural motif reveals the essential
transcriptional repression function of Spen proteins and their role in developmental
signaling. Genes Dev *17*, 1909-1920.
Bach-Pages, M., Homma, F., Kourelis, J., Kaschani, F., Mohammed, S., Kaiser, M., van der
Hoorn, R., Castello, A., and Preston, G. (2020). Discovering the RNA-Binding Proteome of
Plant Leaves with an Improved RNA Interactome Capture Method. Biomolecules *10, 661*.
Barragan, A.C., Collenberg, M., Wang, J., Lee, R.R.Q., Cher, W.Y., Rabanal, F.A., Ashkenazy,
H., Weigel, D., Chae, E., and Patricia, W. (2021). A Truncated Singleton NLR Causes Hybrid
Necrosis in Arabidopsis thaliana. Molecular Biology and Evolution *38*, 557-574.
Barragan, A.C., and Weigel, D. (2020). Plant NLR Diversity: The Known Unknowns of Pan-
NLRomes. The Plant Cell, koaa002.
Baurle, I., Smith, L., Baulcombe, D.C., and Dean, C. (2007). Widespread role for the
flowering-time regulators FCA and FPA in RNA-mediated chromatin silencing. Science *318*,
109-112.
Bomblies, K., and Weigel, D. (2007). Hybrid necrosis: autoimmunity as a potential gene-flow
barrier in plant species. Nat Rev Genet *8*, 382-393.
Bruggeman, Q., Garmier, M., de Bont, L., Soubigou-Taconnat, L., Mazubert, C., Benhamed,
M., Raynaud, C., Bergounioux, C., and Delarue, M. (2014). The Polyadenylation Factor
Subunit CLEAVAGE AND POLYADENYLATION SPECIFICITY FACTOR30: A Key Factor of
Programmed Cell Death and a Regulator of Immunity in Arabidopsis. Plant Physiology *165*,
732-746.
Cai, Q., Liang, C., Wang, S., Hou, Y., Gao, L., Liu, L., He, W., Ma, W., Mo, B., and Chen, X.
(2019). Author Correction: The disease resistance protein SNC1 represses the biogenesis of
microRNAs and phased siRNAs. Nat Commun *10*, 642.
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden,
T.L. (2009). BLAST+: architecture and applications. BMC bioinformatics *10*, 421-421.
Canto-Pastor, A., Santos, B., Valli, A.A., Summers, W., Schornack, S., and Baulcombe, D.C.
(2019). Enhanced resistance to bacterial and oomycete pathogens by short tandem target
mimic RNAs in tomato. Proc Natl Acad Sci U S A *116*, 2755-2760.
Cheng, C.-Y., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S., and Town, C.D.
(2017). Araport11: a complete reannotation of the *Arabidopsis thaliana* reference genome.
The Plant Journal *89*, 789-804.
Collado-Torres, L., Nellore, A., Frazee, A.C., Wilks, C., Love, M.I., Langmead, B., Irizarry, R.A.,
Leek, J.T., and Jaffe, A.E. (2017). Flexible expressed region analysis for RNA-seq with
derfinder. Nucleic Acids Res *45*, e9.
Corwin, J.A., and Kliebenstein, D.J. (2017). Quantitative Resistance: More Than Just
Perception of a Pathogen. Plant Cell *29*, 655-665.
Cox, J., and Mann, M. (2008). MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat
Biotechnol *26*, 1367-1372.
Dale, R.K., Pedersen, B.S., and Quinlan, A.R. (2011). Pybedtools: a flexible Python library for
manipulating genomic datasets and annotations. Bioinformatics *27*, 3423-3424.
Deremetz, A., Le Roux, C., Idir, Y., Brousse, C., Agorio, A., Gy, I., Parker, J.E., and Bouche, N.
(2019). Antagonistic Actions of FPA and IBM2 Regulate Transcript Processing from Genes
Containing Heterochromatin. Plant Physiol *180*, 392-403.

Dinesh-Kumar, S.P., and Baker, B.J. (2000). Alternatively spliced N resistance gene transcripts: their possible role in tobacco mosaic virus resistance. Proc Natl Acad Sci U S A *97*, 1908-1913.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Duc, C., Sherstnev, A., Cole, C., Barton, G.J., and Simpson, G.G. (2013). Transcription termination and chimeric RNA formation controlled by *Arabidopsis thaliana* FPA. PLoS Genet *9*, e1003867.

Ewels, P., Magnusson, M., Käller, M., and Lundin, S. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics *32*, 3047-3048.

Fu, Z.Q., Guo, M., Jeong, B.R., Tian, F., Elthon, T.E., Cerny, R.L., Staiger, D., and Alfano, J.R. (2007). A type III effector ADP-ribosylates RNA-binding proteins and quells plant immunity. Nature *447*, 284-288.

Gierlinski, M., Gastaldello, F., and Barton, G.J. (2018). Proteus: an R package for downstream analysis of MaxQuant output. bioRxiv, 416511.

Gloggnitzer, J., Akimcheva, S., Srinivasan, A., Kusenda, B., Riehs, N., Stampfl, H., Bautor, J., Dekrout, B., Jonak, C., Jimenez-Gomez, J.M.*, et al.* (2014). Nonsense-mediated mRNA decay modulates immune receptor levels to regulate plant antibacterial defense. Cell Host Microbe *16*, 376-390.

Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N.J.*, et al.* (2020). Array programming with NumPy. Nature *585*, 357-362.

Hornyik, C., Terzi, L.C., and Simpson, G.G. (2010). The spen family protein FPA controls alternative cleavage and polyadenylation of RNA. Dev Cell *18*, 203-213.

Huang, H., Weng, H., Sun, W., Qin, X., Shi, H., Wu, H., Zhao, B.S., Mesquita, A., Liu, C., Yuan, C.L.*, et al.* (2018). Recognition of RNA N^6-methyladenosine by IGF2BP proteins enhances mRNA stability and translation. Nature Cell Biology *20*, 285-295.

Hunter, J.D. (2007). Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering *9*, 90-95.

Inagaki, S., Takahashi, M., Hosaka, A., Ito, T., Toyoda, A., Fujiyama, A., Tarutani, Y., and Kakutani, T. (2017). Gene-body chromatin modification dynamics mediate epigenome differentiation in Arabidopsis. EMBO J *36*, 970-980.

Jiao, W.B., and Schneeberger, K. (2020). Chromosome-level assemblies of multiple Arabidopsis genomes reveal hotspots of rearrangements with altered evolutionary dynamics. Nat Commun *11*, 989.

Jones, J.D.G., Vance, R.E., and Dangl, J.L. (2016). Intracellular innate immune surveillance devices in plants and animals. Science *354*, aaf6395-aaf6395.

Knuckles, P., Lence, T., Haussmann, I.U., Jacob, D., Kreim, N., Carl, S.H., Masiello, I., Hares, T., Villasenor, R., Hess, D.*, et al.* (2018). Zc3h13/Flacc is required for adenosine methylation by bridging the mRNA-binding factor Rbm15/Spenito to the m(6)A machinery component Wtap/Fl(2)d. Gene Dev *32*, 415-429.

Koiwa, H., Hausmann, S., Bang, W.Y., Ueda, A., Kondo, N., Hiraguri, A., Fukuhara, T., Bahk, J.D., Yun, D.J., Bressan, R.A.*, et al.* (2004). Arabidopsis C-terminal domain phosphatase-like 1 and 2 are essential Ser-5-specific C-terminal domain phosphatases. Proc Natl Acad Sci U S A *101*, 14539-14544.

Komarnitsky, P., Cho, E.J., and Buratowski, S. (2000). Different phosphorylated forms of RNA polymerase II and associated mRNA processing factors during transcription. Genes Dev *14*, 2452-2460.

Koornneef, M., Hanhart, C.J., and van der Veen, J.H. (1991). A genetic and physiological analysis of late flowering mutants in Arabidopsis thaliana. Mol Gen Genet *229*, 57-66.

Koster, J., and Rahmann, S. (2012). Snakemake--a scalable bioinformatics workflow engine. Bioinformatics *28*, 2520-2522.

Krzyszton, M., Zakrzewska-Placzek, M., Kwasnik, A., Dojer, N., Karlowski, W., and Kufel, J. (2018). Defective XRN3-mediated transcription termination in Arabidopsis affects the expression of protein-coding genes. Plant J *93*, 1017-1031.

Kyburz, A., Friedlein, A., Langen, H., and Keller, W. (2006). Direct interactions between subunits of CPSF and the U2 snRNP contribute to the coupling of pre-mRNA 3' end processing and splicing. Mol Cell *23*, 195-205.

Lai, Y., Cuzick, A., Lu, X.M., Wang, J., Katiyar, N., Tsuchiya, T., Le Roch, K., McDowell, J.M., Holub, E., and Eulgem, T. (2019). The Arabidopsis RRM domain protein EDM3 mediates race-specific disease resistance by controlling H3K9me2-dependent alternative polyadenylation of RPP7 immune receptor transcripts. Plant J *97*, 646-660.

Lai, Y., and Eulgem, T. (2018). Transcript-level expression control of plant NLR genes. Mol Plant Pathol *19*, 1267-1281.

Lai, Y., Lu, X.M., Daron, J., Pan, S., Wang, J., Wang, W., Tsuchiya, T., Holub, E., McDowell, J.M., Slotkin, R.K.*, et al.* (2020). The Arabidopsis PHD-finger protein EDM2 has multiple roles in balancing NLR immune receptor gene expression. PLoS Genet *16*, e1008993.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods *9*, 357-359.

Lazar, C. (2015). imputeLCMD: a collection of methods for left-censored missing data imputation. R package, version *2*.

Le Roux, C., Huet, G., Jauneau, A., Camborde, L., Tremousaygue, D., Kraut, A., Zhou, B., Levaillant, M., Adachi, H., Yoshioka, H.*, et al.* (2015). A receptor pair with an integrated decoy converts pathogen disabling of transcription factors to immunity. Cell *161*, 1074-1088.

Lee, R.R.Q., and Chae, E. (2020). Variation Patterns of NLR Clusters in Arabidopsis thaliana Genomes. Plant Communications *1,* 100089.

Li, L., Habring, A., Wang, K., and Weigel, D. (2020). Atypical Resistance Protein RPW8/HR Triggers Oligomerization of the NLR Immune Receptor RPP7 and Autoimmunity. Cell Host Microbe *27*, 405-417.

Liao, Y., Smyth, G.K., and Shi, W. (2013). The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. Nucleic Acids Res *41*, e108-e108.

Liu, F., Quesada, V., Crevillen, P., Baurle, I., Swiezewski, S., and Dean, C. (2007). The Arabidopsis RNA-binding protein FCA requires a lysine-specific demethylase 1 homolog to downregulate FLC. Mol Cell *28*, 398-407.

Liu, F.Q., Marquardt, S., Lister, C., Swiezewski, S., and Dean, C. (2010). Targeted 3' Processing of Antisense Transcripts Triggers Arabidopsis FLC Chromatin Silencing. Science *327*, 94-97.

Lyons, R., Iwase, A., Gansewig, T., Sherstnev, A., Duc, C., Barton, G.J., Hanada, K., Higuchi-Takeuchi, M., Matsui, M., Sugimoto, K.*, et al.* (2013). The RNA-binding protein FPA regulates flg22-triggered defense responses and transcription factor activity by alternative polyadenylation. Sci Rep *3*, 2866.

Ma, W., Zhen, G., Xie, W., and Mayr, C. (2020). Unstructured mRNAs form multivalent RNA-
RNA interactions to generate TIS granule networks. bioRxiv, 2020.2002.2014.949503.
Martin, E.C., Sukarta, O.C.A., Spiridon, L., Grigore, L.G., Constantinescu, V., Tacutu, R.,
Goverse, A., and Petrescu, A.-J. (2020). LRRpredictor—A New LRR Motif Detection Method
for Irregular Motifs of Plant NLR Proteins Using an Ensemble of Classifiers. Genes-Basel *11,*
286.
Mayr, C. (2019). What Are 3' UTRs Doing? Csh Perspect Biol *11*.a034728.
Mitchell, A.L., Attwood, T.K., Babbitt, P.C., Blum, M., Bork, P., Bridge, A., Brown, S.D., Chang,
H.Y., El-Gebali, S., Fraser, M.I.*, et al.* (2019). InterPro in 2019: improving coverage,
classification and access to protein sequence annotations. Nucleic Acids Res *47*, D351-D360.
Miura, A., Nakamura, M., Inagaki, S., Kobayashi, A., Saze, H., and Kakutani, T. (2009). An
Arabidopsis jmjC domain protein protects transcribed genes from DNA methylation at CHG
sites. EMBO J *28*, 1078-1086.
Mockler, T.C., Yu, X., Shalitin, D., Parikh, D., Michael, T.P., Liou, J., Huang, J., Smith, Z.,
Alonso, J.M., Ecker, J.R.*, et al.* (2004). Regulation of flowering time in Arabidopsis by K
homology domain proteins. Proc Natl Acad Sci U S A *101*, 12759-12764.
Nishimura, M.T., Anderson, R.G., Cherkis, K.A., Law, T.F., Liu, Q.L., Machius, M., Nimchuk,
Z.L., Yang, L., Chung, E.-H., El Kasmi, F.*, et al.* (2017). TIR-only protein RBA1 recognizes a
pathogen effector to regulate cell death inArabidopsis. Proc Natl Acad Sci U S A *114*, E2053-
E2062.
Nishimura, M.T., Monteiro, F., and Dangl, J.L. (2015). Treasure your exceptions: unusual
domains in immune receptors reveal host virulence targets. Cell *161*, 957-960.
Nyikó, T., Sonkoly, B., Mérai, Z., Benkovics, A.H., and Silhavy, D. (2009). Plant upstream ORFs
can trigger nonsense-mediated mRNA decay in a size-dependent manner. Plant Mol Biol *71*,
367-378.
Ozsolak, F., Platt, A.R., Jones, D.R., Reifenberger, J.G., Sass, L.E., McInerney, P., Thompson,
J.F., Bowers, J., Jarosz, M., and Milos, P.M. (2009). Direct RNA sequencing. Nature *461*, 814-
818.
Parker, M.T., Knop, K., Barton, G.J., and Simpson, G.G. (2021). 2passtools: two-pass
alignment using machine-learning-filtered splice junctions increases the accuracy of intron
detection in long-read RNA sequencing. Genome Biology *22*.1-24
Parker, M.T., Knop, K., Sherwood, A.V., Schurch, N.J., Mackinnon, K., Gould, P.D., Hall,
A.J.W., Barton, G.J., and Simpson, G.G. (2020). Nanopore direct RNA sequencing maps the
complexity of Arabidopsis mRNA processing and m6A modification. eLife *9*, e49658.
Patil, D.P., Chen, C.K., Pickering, B.F., Chow, A., Jackson, C., Guttman, M., and Jaffrey, S.R.
(2016). m(6)A RNA methylation promotes XIST-mediated transcriptional repression. Nature
537, 369-373.
Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D.J.,
Inuganti, A., Griss, J., Mayer, G., Eisenacher, M.*, et al.* (2019). The PRIDE database and
related tools and resources in 2019: improving support for quantification data. Nucleic Acids
Res *47*, D442-D450.
Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T., and Salzberg, S.L.
(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.
Nat Biotechnol *33*, 290-295.
Prigozhin, D.M., and Krasileva, K.V. (2021). Analysis of intraspecies diversity reveals a subset
of highly variable plant immune receptors and predicts their binding sites. The Plant Cell,
koab013.

Proudfoot, N.J. (1986). Transcriptional interference and termination between duplicated alpha-globin gene constructs suggests a novel mechanism for gene regulation. Nature *322*, 562-565.

Quesada, V., Macknight, R., Dean, C., and Simpson, G.G. (2003). Autoregulation of *FCA* pre-mRNA processing controls *Arabidopsis* flowering time. EMBO J *22*, 3142-3152.

Ramirez, F., Dundar, F., Diehl, S., Gruning, B.A., and Manke, T. (2014). deepTools: a flexible platform for exploring deep-sequencing data. Nucleic Acids Res *42*, W187-191.

Raxwal, V.K., and Riha, K. (2016). Nonsense mediated RNA decay and evolutionary capacitance. Biochim Biophys Acta *1859*, 1538-1543.

Reichel, M., Liao, Y., Rettel, M., Ragan, C., Evers, M., Alleaume, A.M., Horos, R., Hentze, M.W., Preiss, T., and Millar, A.A. (2016). *In planta* determination of the mRNA-binding proteome of Arabidopsis etiolated seedlings. Plant Cell *28*, 2435-2452.

Reyes, A., Anders, S., Weatheritt, R.J., Gibson, T.J., Steinmetz, L.M., and Huber, W. (2013). Drift and conservation of differential exon usage across tissues in primate species. Proc Natl Acad Sci U S A *110*, 15377-15382.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res *43*, e47.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139-140.

Rodriguez, E., El Ghoul, H., Mundy, J., and Petersen, M. (2016). Making sense of plant autoimmunity and 'negative regulators'. FEBS J *283*, 1385-1391.

Rodriguez-Cazorla, E., Ripoll, J.J., Andujar, A., Bailey, L.J., Martinez-Laborda, A., Yanofsky, M.F., and Vera, A. (2015). K-homology nuclear ribonucleoproteins regulate floral organ identity and determinacy in arabidopsis. PLoS Genet *11*, e1004983.

Ruzicka, K., Zhang, M., Campilho, A., Bodi, Z., Kashif, M., Saleh, M., Eeckhout, D., El-Showk, S., Li, H., Zhong, S.*, et al.* (2017). Identification of factors required for m^{6}A mRNA methylation in Arabidopsis reveals a role for the conserved E3 ubiquitin ligase HAKAI. New Phytol *215*, 157-172.

Sadowski, M., Dichtl, B., Hubner, W., and Keller, W. (2003). Independent functions of yeast Pcf11p in pre-mRNA 3' end processing and in transcription termination. EMBO J *22*, 2167-2177.

Saze, H., Kitayama, J., Takashima, K., Miura, S., Harukawa, Y., Ito, T., and Kakutani, T. (2013). Mechanism for full-length RNA processing of Arabidopsis genes containing intragenic heterochromatin. Nat Commun *4*, 2301.

Saze, H., Shiraishi, A., Miura, A., and Kakutani, T. (2008). Control of genic DNA methylation by a jmjC domain-containing protein in Arabidopsis thaliana. Science *319*, 462-465.

Schomburg, F.M., Patton, D.A., Meinke, D.W., and Amasino, R.M. (2001). FPA, a gene involved in floral induction in Arabidopsis, encodes a protein containing RNA-recognition motifs. Plant Cell *13*, 1427-1436.

Schon, M.A., Kellner, M.J., Plotnikova, A., Hofmann, F., and Nodine, M.D. (2018). NanoPARE: parallel analysis of RNA 5' ends from low-input RNA. Genome Res *28*, 1931-1942.

Seabold, S., and Perktold, J. (2010). Statsmodels: Econometric and Statistical Modeling with Python. In Proceedings of the 9th Python in Science Conference, pp. 92-96.

Sherstnev, A., Duc, C., Cole, C., Zacharaki, V., Hornyik, C., Ozsolak, F., Milos, P.M., Barton, G.J., and Simpson, G.G. (2012). Direct sequencing of *Arabidopsis thaliana* RNA reveals patterns of cleavage and polyadenylation. Nat Struct Mol Biol *19*, 845-852.

Shivaprasad, P.V., Chen, H.M., Patel, K., Bond, D.M., Santos, B.A., and Baulcombe, D.C. (2012). A microRNA superfamily regulates nucleotide binding site-leucine-rich repeats and other mRNAs. Plant Cell *24*, 859-874.

Simpson, G.G. (2004). The autonomous pathway: epigenetic and post-transcriptional gene regulation in the control of Arabidopsis flowering time. Curr Opin Plant Biol *7*, 570-574.

Smedley, D., Haider, S., Ballester, B., Holland, R., London, D., Thorisson, G., and Kasprzyk, A. (2009). BioMart--biological queries made easy. BMC Genomics *10*, 22.

Swiderski, M.R., Birker, D., and Jones, J.D.G. (2009). The TIR Domain of TIR-NB-LRR Resistance Proteins Is a Signaling Domain Involved in Cell Death Induction. Molecular Plant-Microbe Interactions *22*, 157-165.

Szádeczky-Kardoss, I., Csorba, T., Auber, A., Schamberger, A., Nyikó, T., Taller, J., Orbán, T.I., Burgyán, J., and Silhavy, D. (2018). The nonstop decay and the RNA silencing systems operate cooperatively in plants. Nucleic Acids Res *46*, 4632-4648.

Takagi, M., Iwamoto, N., Kubo, Y., Morimoto, T., Takagi, H., Takahashi, F., Nishiuchi, T., Tanaka, K., Taji, T., Kaminaka, H., *et al.* (2020). Arabidopsis SMN2/HEN2, Encoding DEAD-Box RNA Helicase, Governs Proper Expression of the Resistance Gene SMN1/RPS6 and Is Involved in Dwarf, Autoimmune Phenotypes of mekk1 and mpk4 Mutants. Plant Cell Physiol *61*, 1507-1516.

Tamborski, J., and Krasileva, K.V. (2020). Evolution of Plant NLRs: From Natural History to Precise Modifications. Annu Rev Plant Biol *71*, 355-378.

The Arabidopsis Genome, I. (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. Nature *408*, 796-815.

Tian, D., Traw, M.B., Chen, J.Q., Kreitman, M., and Bergelson, J. (2003). Fitness costs of R-gene-mediated resistance in Arabidopsis thaliana. Nature *423*, 74-77.

Tome, D.F., Steinbrenner, J., and Beynon, J.L. (2014). A growth quantification assay for Hyaloperonospora arabidopsidis isolates in Arabidopsis thaliana. Methods Mol Biol *1127*, 145-158.

Tran Vdu, T., Souiai, O., Romero-Barrios, N., Crespi, M., and Gautheret, D. (2016). Detection of generic differential RNA processing events from RNA-seq data. RNA Biol *13*, 59-67.

Tsuchiya, T., and Eulgem, T. (2013). An alternative polyadenylation mechanism coopted to the Arabidopsis RPP7 gene through intronic retrotransposon domestication. Proc Natl Acad Sci U S A *110*, E3535-3543.

Van de Weyer, A.L., Monteiro, F., Furzer, O.J., Nishimura, M.T., Cevik, V., Witek, K., Jones, J.D.G., Dangl, J.L., Weigel, D., and Bemm, F. (2019). A Species-Wide Inventory of NLR Genes and Alleles in Arabidopsis thaliana. Cell *178*, 1260-1272 e1214.

van der Biezen, E.A., Freddie, C.T., Kahn, K., Parker, J.E., and Jones, J.D. (2002). Arabidopsis RPP4 is a member of the RPP5 multigene family of TIR-NB-LRR genes and confers downy mildew resistance through multiple signalling components. Plant J *29*, 439-451.

Wang, X., Boevink, P., McLellan, H., Armstrong, M., Bukharova, T., Qin, Z., and Birch, P.R. (2015). A Host KH RNA-Binding Protein Is a Susceptibility Factor Targeted by an RXLR Effector to Promote Late Blight Disease. Mol Plant *8*, 1385-1395.

Wang, Y.H., and Warren, J.T., Jr. (2010). Mutations in retrotransposon AtCOPIA4 compromises resistance to Hyaloperonospora parasitica in Arabidopsis thaliana. Genet Mol Biol *33*, 135-140.

Wang, Z.W., Wu, Z., Raitskin, O., Sun, Q., and Dean, C. (2014). Antisense-mediated FLC transcriptional repression requires the P-TEFb transcription elongation factor. Proc Natl Acad Sci U S A *111*, 7468-7473.

Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., and Barton, G.J. (2009). Jalview
Version 2--a multiple sequence alignment editor and analysis workbench. Bioinformatics *25*,
1189-1191.
Wei, C., Chen, J., and Kuang, H. (2016). Dramatic Number Variation of R Genes in Solanaceae
Species Accounted for by a Few R Gene Subfamilies. Plos One *11*, e0148708.
Williams, S.J., Sohn, K.H., Wan, L., Bernoux, M., Sarris, P.F., Segonzac, C., Ve, T., Ma, Y.,
Saucet, S.B., Ericsson, D.J.*, et al.* (2014). Structural basis for assembly and function of a
heterodimeric plant immune receptor. Science *344*, 299-303.
Xia, Z., Donehower, L.A., Cooper, T.A., Neilson, J.R., Wheeler, D.A., Wagner, E.J., and Li, W.
(2014). Dynamic analyses of alternative polyadenylation from RNA-seq reveal a 3'-UTR
landscape across seven tumour types. Nat Commun *5*, 5274.
Xing, D.H., Zhao, H.W., Xu, R.Q., and Li, Q.S.Q. (2008). Arabidopsis PCFS4, a homologue of
yeast polyadenylation factor Pcf11p, regulates FCA alternative processing and promotes
flowering time. Plant Journal *54*, 899-910.
Yang, S., Tang, F., Gao, M., Krishnan, H.B., and Zhu, H. (2010). R gene-controlled host
specificity in the legume-rhizobia symbiosis. Proc Natl Acad Sci U S A *107*, 18735-18740.
Yu, X., Martin, P.G.P., and Michaels, S.D. (2019). BORDER proteins protect expression of
neighboring genes by promoting 3' Pol II pausing in plants. Nat Commun *10*, 4359.
Zhai, J., Jeong, D.H., De Paoli, E., Park, S., Rosen, B.D., Li, Y., Gonzalez, A.J., Yan, Z., Kitto, S.L.,
Grusak, M.A.*, et al.* (2011). MicroRNAs as master regulators of the plant NB-LRR defense
gene family via the production of phased, trans-acting siRNAs. Genes Dev *25*, 2540-2553.
Zhang, P., Berardini, T.Z., Ebert, D., Li, Q., Mi, H., Muruganujan, A., Prithvi, T., Reiser, L.,
Sawant, S., Thomas, P.D.*, et al.* (2020a). PhyloGenes: An online phylogenetics and functional
genomics resource for plant gene function inference. Plant Direct *4*.
Zhang, X.C., and Gassmann, W. (2007). Alternative splicing and mRNA levels of the disease
resistance gene RPS4 are induced during defense responses. Plant Physiol *145*, 1577-1587.
Zhang, Y., Dorey, S., Swiderski, M., and Jones, J.D. (2004). Expression of RPS4 in tobacco
induces an AvrRps4-independent HR that requires EDS1, SGT1 and HSP90. Plant J *40*, 213-
224.
Zhang, Y., Gu, L., Hou, Y., Wang, L., Deng, X., Hang, R., Chen, D., Zhang, X., Zhang, Y., Liu, C.*,
et al.* (2015). Integrative genome-wide analysis reveals HLP1, a novel RNA-binding protein,
regulates plant flowering by targeting alternative polyadenylation. Cell Res *25*, 864-876.
Zhang, Y., Li, Z., Chen, N., Huang, Y., and Huang, S. (2020b). Phase separation of Arabidopsis
EMB1579 controls transcription, mRNA splicing, and development. Plos Biol *18*, e3000782.
Zhang, Y., Rataj, K., Simpson, G.G., and Tong, L. (2016). Crystal Structure of the SPOC
Domain of the Arabidopsis Flowering Regulator FPA. Plos One *11*, e0160694.
1335
1336

Figures

Figure 1: **FPA associates with proteins that function to process the 3′ ends of Pol II-transcribed RNAs and promote transcription termination.**

(A-D) Volcano plots representing proteins co-purifying with FPA using *IVI-MS*. Only proteins detected in all three biological replicates of the *35S::FPA:YFP* line are shown (light grey). The following classes are highlighted: **(A)** CPFs in dark blue; **(B)** Pol II-associated factors in green; terminal exon definition factors in dark orange; **(C)** autonomous pathway components in yellow and factors controlling alternative polyadenylation in light orange; and **(D)** m⁶A writer complex components in light blue. **(E)** ChIP-Seq metagene profile showing the normalised occupancy of FPA (green) and Pol II phosphorylated at Ser⁵ (pink) and Ser² (brown) of the CTD (Yu et al., 2019) relative to the major 3′ position of each gene, as measured using Helicos DRS. Only long genes (> 2.5 kb) are included (*n*=10,215).

1350
Figure 2: FPA regulates poly(A) site selection.
Loss of FPA function is associated with the preferential selection of distal poly(A) sites, whereas FPA
overexpression leads to the preferential selection of proximal poly(A) sites. **(A)** Illumina RNA-Seq,
Helicos DRS and Nanopore DRS reveal FPA-dependent RNA 3′ end processing changes at the *FPA*
(*AT2G43410*) locus. The *35S::FPA:YFP* construct has alternative transgene-derived untranslated
regions, so mRNAs derived from the transgene do not align to the native FPA 5′UTR and 3′UTR. **(B)**
Histograms showing change in mean RNA 3′ end position for significantly alternatively
polyadenylated loci (EMD>25, FDR<0.05) in *fpa-8* (left panel) and *35S::FPA:YFP* (right panel)
compared with Col-0, as detected using Nanopore DRS. Orange and green shaded regions indicate
sites with negative and positive RNA 3′ end position changes, respectively. **(C)** Effect size of
significant proximal (orange) and distal (green) alternative polyadenylation events in *fpa-8* (left
panel) and *35S::FPA:YFP* (right panel) compared with Col-0, as measured using the EMD. **(D)**
Histograms showing change in mean RNA 3′ end position for significantly alternatively
polyadenylated loci (EMD>25, FDR<0.05) in *fpa-8* (left panel) and *35S::FPA:YFP* (right panel)
compared with Col-0, as detected using Nanopore DRS. Orange and green shaded regions indicate
sites with negative and positive RNA 3′ end position changes, respectively. **(E)** Effect size of
significant proximal (orange) and distal (green) alternative polyadenylation events in *fpa-8* (left
panel) and *35S::FPA:YFP* (right panel) compared with Col-0, as measured using the EMD. **(F)**
Boxplots showing the effect size (absolute log$_2$ fold change (logFC)) of alternatively processed loci
identified using Illumina RNA-Seq in *fpa-8* (left panel) and *35S::FPA:YFP* (right panel) respectively.
Down- and upregulated loci are shown in orange and green, respectively. For each locus, the region
with the largest logFC was selected to represent the locus. Loci with both up- and downregulated
regions contribute to both boxes. **(G)** Boxplots showing the effect size (absolute logFC) of loci with
alternative splice junction usage identified using Illumina RNA-Seq in *fpa-8* (left panel) and
35S::FPA:YFP (right panel) respectively. Down- and upregulated loci are shown in orange and green,
respectively. For each locus, the junction with the largest logFC was selected to represent the locus.
Loci with both up- and downregulated junctions contribute to both boxes.

Figure 3: Nanopore and Helicos DRS identify NLR genes regulated by alternative polyadenylation.

(A-B) Protein domain enrichment analysis for loci with increased proximal poly(A) site selection in *35S::FPA:YFP* line, as detected using **(A)** Nanopore DRS or **(B)** Helicos DRS. **(C)** Nanopore DRS reveals the complexity of RNA processing at *RPS6*. Protein domain locations (shown in grey) represent collapsed InterPro annotations. The novel TIR domain was annotated using InterProScan (Mitchell et al., 2019). **(D)** Protein alignment of the predicted TIR domain from the novel gene downstream of RPS6, with the sequence of the TIR domains from RPS6 and RPS4. Helix and strand secondary structures (from UniProt: RPS4, Q9XGM3) are shown in blue and yellow, respectively. Residues are shaded according to the degree of conservation.

1389
Figure 4: FPA-dependent alternative polyadenylation of NLR transcripts.
FPA regulates **(A)** readthrough and chimeric RNA formation at *AT1G58848* (unique mapping of short
Helicos DRS reads was not possible due to the high homology of *AT1G58848* to tandemly duplicated
NLR loci in the same cluster)*;* **(B)** intronic polyadenylation at *AT1G69550*, resulting in transcripts
encoding a protein with a truncated LRR domain; **(C)** exonic polyadenylation at *AT2G14080*, resulting
in stop-codonless transcripts; and **(D)** exonic polyadenylation at *AT5G40060*, resulting in transcripts
encoding a TIR-domain-only protein due to an upstream ORF.

1397
Figure 5: Complex FPA-dependent patterns of alternative polyadenylation at *RPP4*.
FPA regulates intronic, exonic and readthrough poly(A) site selection in *RPP4*. **(Inset 1)** A magnified
view of TIR-domain-only *RPP4* transcripts detected in *35S::FPA:YFP* caused by proximal
polyadenylation in intron 1, and distal polyadenylation within the TE associated with cryptic splicing.
(Inset 2) A magnified view of the stop-codonless transcripts produced within the protein-coding
RPP4 region in *fpa-8*.

1404
Figure 6: FPA promotes premature cleavage and polyadenylation within *RPP7* protein-coding
exon 6 that compromises plant immunity against *Hyaloperonospora arabidopsidis* isolate
Hiks1(*Hpa*-Hiks1).

(A) FPA-dependent RNA 3′ end formation changes at the *RPP7 (AT1G58602)* locus. **(Inset 1)**
Magnified view of the *RPP7* 3′UTR region with alternative RNA 3′ ends. **(Inset 2)** Magnified view of
the stop-codonless transcripts produced in protein-coding *RPP7* exon 6. **(B)** RNA gel blot visualising
RPP7 transcripts in Col-0, *fpa-8* and *35S::FPA:YFP*. Probe location in second exon is shown on (A)
(light brown). Beta-TUBULIN was used as an internal control. **(C)** FPA-dependent premature exonic
termination of *RPP7* compromises immunity against *Hpa*-Hiks1. Point plot showing median number
of sporangiophores per plant calculated 4 days after *Hpa*-Hiks1 inoculation. Error bars are 95%
confidence intervals. Each experimental replicate was generated from 7–45 plants per genotype.

1416
1417 **Figure 7: Functional consequences of FPA-dependent alternative polyadenylation at NLR loci.**
1418 Model diagram showing how FPA-regulated alternative polyadenylation at NLR loci might affect the
1419 regulatory and evolutionary dynamics of plant disease resistance.
1420

Tables

1421

1422

Gene ID	Gene name	NLR class	Chimeric pair (upstream-downstream)
AT1G12220	RPS5	CNL	AT1G12220-AT1G12230
AT1G58848	RPP7a/b	TNL	AT1G58848-AT1G58889
AT1G59218	RPP7a/b	TNL	AT1G59218-AT1G59265
AT1G61190	-	CNL	ncRNA-AT1G61190
AT1G63730	-	TNL	AT1G63730-AT1G63740
AT1G63740	-	TNL	AT1G63730-AT1G63740
AT3G46730	-	CNL	AT3G46740-AT3G46730
AT4G16860	RPP4	TNL	AT4G16860-AT4G16870-AT4G16857
AT4G16960	SIKIC3	TNL	AT4G16970-AT4G16960-AT4G16957
AT4G19060	-	NB only	AT4G19070-AT4G19060
AT4G19530	-	TNL	AT4G19530-AT4G19540
AT5G38850	-	TNL	AT5G38850-AT5G38860
AT5G40090	CHL1	TNL	ncRNA-AT5G40090
AT5G44510	TAO1	TNL	AT5G44520-AT5G44510
AT5G45490	-	CNL	AT5G45472-AT5G45490
AT5G46470	RPS6	TNL	AT5G46470-TIR gene
AT5G48780	-	TNL	AT5G48775-AT5G48780

1423 **Table 1: Readthrough and chimeric RNA formation events at NLR genes regulated by FPA.**

1424

Gene ID	Gene name	NLR class	Predicted function	Protein isoform
AT1G12210	RFL1	CNL	non-coding (5′UTR)	-
AT1G58602	RPP7	CNL	non-coding (5′UTR); alternative 3′UTR	-
AT1G63750	WRR9	TNL	protein coding	TIR only
AT1G63880	RLM1B	TNL	protein coding; non-stop	TIR only
AT1G69550	-	TNL	protein coding	LRR truncation
AT3G44480	RPP1	TNL	protein coding	LRR truncation
AT3G50480	HR4	RPW8	protein coding	RPW8 truncation
AT4G16860	RPP4	TNL	protein coding	TIR only
AT4G16900	-	TNL	protein coding	LRR truncation
AT4G19510	RPP2B	TNL	alternative 3′UTR	-
AT5G17890	DAR4/CHS3	TNL	protein coding	TIR only
AT5G40910	-	TNL	protein coding	TIR only
AT5G43730	RSG2	CNL	non-coding (5′UTR)	-
AT5G43740	-	CNL	non-coding (5′UTR)	-
AT5G46270	-	TNL	protein coding	TIR/NB-ARC only; LRR truncation
AT5G46470	RPS6	TNL	alternative 3′UTR	
AT5G46490	-	TNL	protein coding; non-stop	TIR/NB-ARC only; LRR truncation

1425 **Table 2: Intronic proximal polyadenylation events at FPA-regulated NLR genes**

1426

Gene ID	Gene name	NLR class	Predicted function	Protein isoform
AT1G10920	LOV1	CNL	protein coding*	CC-only*
AT1G27180	-	TNL	non-stop	-
AT1G31540	RAC1	TNL	non-stop; protein coding^	LRR truncation^
AT1G33560	ADR1	RNL	non-stop	-
AT1G53350	-	CNL	non-stop	-
AT1G56510	WRR4A	TNL	non-stop	-
AT1G56520	-	TNL	non-stop	-
AT1G58602	RPP7	CNL	non-stop	-
AT1G58807	RF45	CNL	non-stop	-
AT1G58848	RPP7a/b	CNL	non-stop	-
AT1G59124	RDL5	CNL	non-stop	-
AT1G59218	RPP7a/b	CNL	non-stop	-
AT1G61300	-	CNL	non-stop	-
AT1G62630	-	CNL	non-stop	-
AT1G63360	-	CNL	non-stop	-
AT1G63730	-	TNL	non-stop	-
AT1G63860	-	TNL	non-stop	-
AT1G63880	RLM1B	TNL	non-stop	-
AT1G72840	-	TNL	non-coding (5'UTR)	-
AT2G14080	RPP28	TNL	non-stop	-
AT3G44480	RPP1	TNL	non-stop; protein coding^	LRR truncation^
AT3G44630	-	TNL	non-stop	-
AT3G44670	-	TNL	non-stop; protein coding^	TIR only^
AT3G46530	RPP13	CNL	non-stop	-
AT4G16860	RPP4	TNL	non-stop	-
AT4G16890	SNC1	TNL	non-stop	-
AT4G16900	-	TNL	non-stop	-
AT4G19520	-	TNL	non-stop	-
AT4G19530	-	TNL	non-stop	-
AT4G36140	-	TNL	non-stop	-
AT5G17890	DAR4/CHS3	TNL	non-stop	-
AT5G35450	-	CNL	non-stop	-
AT5G38850	-	TNL	non-stop	-
AT5G40060	-	TNL	protein coding*	TIR only*
AT5G40910	-	TNL	non-stop	-
AT5G43470	RPP8	CNL	non-stop	-
AT5G43740	-	CNL	non-stop	-
AT5G44510	TAO1	TNL	non-stop; protein coding^	LRR truncation^
AT5G44870	LAZ5	TNL	non-stop	-
AT5G45050	RRS1B	TNL	non-stop	-
AT5G45250	RPS4	TNL	protein coding^	LRR truncation^
AT5G45260	RRS1	TNL	non-stop	-
AT5G46270	-	TNL	non-stop; protein coding^	LRR truncation^
AT5G48620	-	CNL	non-stop	-
AT5G58120	DM10	TNL	non-stop; protein coding^	LRR truncation^

1427 **Table 3: Exonic proximal polyadenylation events at NLR genes regulated by FPA.**

1428 * indicates loci where exonic proximal polyadenylation generates transcripts that may be protein

1429 coding due to upstream ORFs. ^ indicates loci where exonic proximal polyadenylation coupled with

1430 intron retention results in a protein-coding ORF.

1431

 Figure supplement legends

1433
1434   **Figure 1–figure supplement 1: FPA co-localises with Pol II Ser[2] at the 3′ end of genes.**
1435   ChIP-Seq metagene profile showing the normalised occupancy of FPA (green) and Pol II
1436   phosphorylated at Ser[5] and Ser[2] of the CTD relative to the major 3′ position of each gene, as
1437   measured using Helicos DRS. Only short genes (<2.5 kb) are included (n=17,440).
1438

1439
1440 **Figure 1-figure supplement 2: FPA controls Pol II occupancy and chimeric RNA formation at**
1441 ***PIF5.***
1442 ChIP-Seq occupancy in counts per million (CPM) of FPA and Pol II phosphorylated at Ser^5 or Ser^2 at
1443 the *PIF5* and *PAO3* loci. *fpa* mutants display readthrough of the canonical *PIF5* poly(A) site, with a
1444 concomitant loss of Ser^2 at the poly(A) site, and an increase in Ser^5 in downstream P*AO3*.
1445

1446

Figure 2–figure supplement 1: Nanopore and Helicos DRS reveal FPA-dependent RNA 3′ end

processing changes.

(A) Comparison of RNA 3′ ends identified in Nanopore and Helicos DRS datasets in *fpa-8* and

35S::FPA:YFP (compared with Col-0). Bar size indicates the number of alternatively polyadenylated

loci common to an intersection (highlighted using circles below). Bars indicating loci that are

identified as alternatively polyadenylated in a single condition (*fpa-8* or *35S::FPA:YFP*) using a single

technique (Nanopore or Helicos DRS) are presented in black; bars indicating loci identified as distally

polyadenylated in *fpa-8* using both Nanopore and Helicos DRS, in orange; bars indicating loci

identified as proximally polyadenylated in *35S::FPA:YFP* using both Nanopore and Helicos DRS, in

green; and bars indicating loci identified as reciprocally regulated by FPA (distal polyadenylation in

fpa-8, proximal in *35S::FPA:YFP*) using at least one technique, in yellow.

1458

1459
1460 **Figure 2–figure supplement 2: Splicing alterations in *fpa-8* can be explained by changes in**
1461 **RNA 3′ end formation.**
1462 Gene track showing chimeric RNA formation at the *PIF5* gene locus, as detected with Illumina RNA-
1463 Seq, Helicos DRS and Nanopore DRS.
1464

1465

Figure 2–figure supplement 3: FPA does not affect global mRNA m⁶A methylation.

Box plot showing the m⁶A/A ratio, as analysed using LC-MS/MS.

Inagaki et al.
ibm1 altered
H3K9me^2

35S::FPA:YFP
alternative poly(A)
site choice

Lai et al.
ibm1 altered
H2K9me^2

1469

1470 **Figure 2-figure supplement 4: FPA-dependent regulation of NLR expression is independent**
1471 **of IBM1.**

1472 Venn diagram showing genes with altered H3K9me^2 levels in *ibm1–4* mutants, in yellow (Inagaki et
1473 al., 2017) and orange (Lai et al., 2020); and genes with altered poly(A) site choice in *35S::FPA:YFP,*
1474 in green.

1475

1476
1477 **Figure 3-figure supplement 1: Nanopore DRS informs reannotation of the complex NLR locus**
1478 **encompassing the *AT5G46490* and *AT5G46500* annotations.**
1479 Gene track showing alternative polyadenylation at the *AT5G46490* gene locus, as detected with
1480 Illumina RNA-Seq, nanoPARE, Helicos DRS and Nanopore DRS.
1481

```
AT5G46500   11  CSRLKCVSLHISKLKHLEDALFPACGALNRVELSGSSS    48
AT5G46260  875  CSRLKCVSLHISKLKRLGKVDFKDCGALTIVDLCGCPI   912
AT5G46520  873  CRELKCVSLNIFKLKHLGEVSFSNCGALTRVDLSCYPS   910

AT5G46500   49  GM---KADNIDTAS----SSLP---QVELDFRECFNLD    76
AT5G46260  913  GME-MEANNIDTVS-----------KVKLDFRDCFNLD   938
AT5G46520  911  GVEMMKADNADIVSEETTSSLPDSCVLNVNFMDCVNLD   948

AT5G46500   77  PETVLHQESIIFKYMLFPGKEEVPSYFTYRTTGV----  110
AT5G46260  939  PETVLHQESIIFKYMLFPGKEEMPSYFTYRTTGS----  972
AT5G46520  949  REPVLHQQSIIFNSMILPG-EEVPSYFTYRTSDSQPFG  985

AT5G46500  111  --SSLTIPLLHLPLSQPFFRFRVGALVTNVKHGKNIKV  146
AT5G46260  973  --SSLTIPLLHLPLSQPFFRFRVGALVTNVKHGKNIKV 1008
AT5G46520  986  TSSSLPIPLLPTQLSQPFFRFRVCAVV-SASNGVYIGV 1022

AT5G46500  147  KCEFKDRFGNSFHVGSDDFYVYLLFTKSQKGSQMLTIL  184
AT5G46260 1009  KCEFKDRFGNSFHVGSDDFYVYLLFTKSQKGSQMLTIL 1046
AT5G46520 1023  YSRFKGRIGNKFD--S--FGEVHNFMEIEKG-IHLCIF 1055

AT5G46500  185  DCCIPLNEGNASLAQGNYYDHVDINIHITESLGSFGST  222
AT5G46260 1047  DCCIPLNEGNASLAQGNYYDHVDINIHISS--GGWRST 1082
AT5G46520 1056  DCRIRLYKDNVPLSQLN-YDHVDINIHITS--GDWRST 1090

AT5G46500  223  SELKEWGIRLLEEDSSSAENQLGNPNSTLPHVSEAEEG  260
AT5G46260 1083  FELKEWGIRLLEEDSSSAENGLGNPNSTLPHVSEAEEG 1120
AT5G46520 1091  VVLKEWGIRLL-ETGSSAENRLGNPNSTLPHVSQAEEG 1127

AT5G46500  261  NMGYYTPVQ-GLVNEIEHNGESGDNNVETERSTKHAA   296
AT5G46260 1121  NMGYYTPLQEGLVNEIEHSEESGDINVGTKRSKKRMR  1157
AT5G46520 1128  NMGYYTHVQ-GLVNEIENSEDSGDNNVETERSKKRMR  1163
```

Figure 3–figure supplement 2: Nanopore DRS informs reannotation of the complex NLR locus encompassing the *AT5G46490* and *AT5G46500* annotations.

Protein alignment showing similarity between the *AT5G46500* protein sequence (which forms the C-terminal portion of distally polyadenylation *AT5G46490–AT5G46500* mRNAs) and other NLR protein sequences in the *RPS6* cluster. LRR predictions, generated with LRRpredictor (Martin et al., 2020), are shown in orange.

Figure 4–figure supplement 1: NLR genes with FPA-regulated alternative polyadenylation are found in hotspots of rearrangements.

Boxplot showing the synteny diversity, calculated from seven diverse *A. thaliana* accessions (Jiao and Schneeberger, 2020), of expressed NLR genes with and without FPA-regulated alternative polyadenylation.

1497
Figure 4-figure supplement 2: Loss of FPA function causes chimeric RNA formation at

AT1G63730 and AT1G63740 NLR loci.

Gene track showing chimeric RNA formation at the *AT1G63730* gene locus, as detected with Illumina

RNA-Seq, Helicos DRS and Nanopore DRS.

Figure 4-figure supplement 3: FPA overexpression increases exonic proximal polyadenylation of *RPP13*.

Gene track showing proximal polyadenylation at the *RPP13* gene locus, as detected with Illumina RNA-Seq, Helicos DRS and Nanopore DRS.

1509
1510 **Figure 4-figure supplement 4: FPA overexpression causes intron retention and exonic**
1511 **proximal polyadenylation at intron 3 of *RPS4*.**
1512 Gene track showing proximal polyadenylation at the *RPS4* gene locus, as detected with Illumina
1513 RNA-Seq, Helicos DRS and Nanopore DRS.
1514

1515
Figure 5-figure supplement 1: Complex FPA-dependent patterns of alternative
polyadenylation at the *RPP4* locus.
Comparison of the expression of four classes of *RPP4* (AT4G16860) transcripts detected using **(A)**
Nanopore DRS or **(B)** Helicos DRS. *N-terminal truncation*, TIR-domain-only transcripts generated by
proximal intronic polyadenylation or distal polyadenylation and cryptic splicing; *Non-stop*, mRNAs
lacking in-frame stop codons; *Full-length*, full-length protein-coding mRNAs; and Chimeric, mRNAs
containing *RPP4*, COPIA-like retrotransposon (AT4G16870) and/or downstream *AT4G16857*.
1523

Figure 6–figure supplement 1: Complex FPA-dependent patterns of alternative polyadenylation at the *RPP7* locus.

Comparison of the expression of three classes of *RPP7* transcripts detected using **(A)** nanopore DRS or **(B)** Helicos DRS. *5'UTR (non-coding)*, mRNAs prematurely terminated within the 5'UTR; *exon 6 (non-stop)*, stop-codonless transcripts terminated at proximal poly(A) sites in exon 6; and *full-length*, protein-coding mRNAs terminated at distal poly(A) sites within the 3'UTR.

List of supplementary files:

Supplementary file 1: Proteins co-purifying with FPA, as identified by *IVI-MS* **[Linked to Figure 1].**

Supplementary file 2: Properties of the sequencing datasets produced using Nanopore DRS, Helicos DRS and Illumina RNA-Seq **[Linked to Figure 2]**.

List of source datasets:

Figure 2 source data 1: Nanopore StringTie assembly **[Linked to Figure 2A-B]**.

Figure 2 source data 2: Differential 3′ processing results for *fpa-8* vs Col-0, as identified by Nanopore DRS **[Linked to Figure 2B–C]**.

Figure 2 source data 3: Differential 3′ processing results for *35S::FPA:YFP* vs Col-0, as identified by Nanopore DRS **[Linked to Figure 2B–C]**.

Figure 2 source data 4: Differential 3′ processing results for *fpa-8* vs Col-0, as identified by Helicos DRS **[Linked to Figure 2D-E]**.

Figure 2 source data 5: Differential 3′ processing results for *35S::FPA:YFP* vs Col-0, as identified by Helicos DRS **[Linked to Figure 2D-E]**.

Figure 2 source data 6: Differentially expressed regions results for *fpa-8* vs Col-0, as identified by Illumina RNA-Seq **[Linked to Figure 2F]**.

Figure 2 source data 7: Differentially expressed regions results for *35S::FPA:YFP* vs Col-0, as identified by Illumina RNA-Seq **[Linked to Figure 2F]**.

Figure 2 source data 8: Differential splice junction usage results for *fpa-8* vs Col-0, as identified by Illumina RNA-Seq **[Linked to Figure 2G]**.

Figure 2 source data 9: Differential splice junction usage results for *35S::FPA:YFP* vs Col-0, as identified by Illumina RNA-Seq **[Linked to Figure 2G]**.

Figure 2 source data 10: m^6A/A ratios for Col-0, *fpa-8*, *35S::FPA:YFP* and *vir-1*, as detected by LC-MS/MS **[Linked to Figure 2-figure supplement 3]**.

Figure 2 source data 11: Differential H3K9me^2 results for *ibm1–4* vs Col-0 **[Linked to Figure 2-figure supplement 4]**.

Figure 6 source data 1: *Hpa*-Hiks1 susceptibility results for the Col-0, Ksk-1, *fpa-7, fpa-8, pFPA::FPA* and *35S::FPA:YFP* lines **[Linked to Figure 6C]**.