Utilization of the repeated squat-stand model for studying the directional sensitivity

of the cerebral pressure-flow relationship

Lawrence Labrecque[1,2], Jonathan D Smirl[3,4,5,6,7,8,9], Patrice Brassard[1,2].

[1]Department of Kinesiology, Faculty of Medicine, Université Laval, Québec, Canada

[2]Research center of the Institut universitaire de cardiologie et de pneumologie de
Québec, Québec, Canada
[3]Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary,
Calgary, Alberta, Canada
[4]Concussion Research Laboratory, Faculty of Health and Exercise Science, University of
British Columbia, Kelowna, BC, Canada
[5]Human Performance Laboratory, Faculty of Kinesiology, University of Calgary, Calgary,
Alberta, Canada
[6]Hotchkiss Brain Institute, University of Calgary, Calgary, Alberta, Canada
[7]Integrated Concussion Research Program, University of Calgary, Calgary, AB, Canada
[8]Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Alberta,
Canada
[9]Libin Cardiovascular Institute of Alberta, University of Calgary, Alberta, Canada
20

Running title: Hysteresis in the pressure-flow relationship

Address for correspondence:

Patrice Brassard, Ph.D.

Department of Kinesiology, Faculty of Medicine

PEPS - Université Laval

2300 rue de la Terrasse, room 0290-H

Québec (Qc) GIV OA6, Canada

Phone: 418 656-2131 extension 405621

Email: patrice.brassard@kin.ulaval.ca

Abstract

Hysteresis in the cerebral pressure-flow relationship describes the superior ability of the

cerebrovasculature to buffer cerebral blood flow changes when mean arterial pressure

(MAP) acutely increases compared to when MAP acutely decreases. This phenomenon

can be evaluated by comparing the change in middle cerebral artery mean blood velocity

(MCAv) per change in MAP during either acute increases or decreases in MAP induced by

repeated squat-stands (RSS). However, no real baseline can be employed for this

particular protocol as there is no true stable reference point. Herein, we characterized a

novel metric using the greatest MAP oscillations induced by RSS without using an

independent baseline value and adjusted for time intervals ($\Delta MCAv_T/\Delta MAP_T$). We also

examined whether this metric during each RSS transition were comparable between each

other over a 5-min period. $\Delta MCAv_T/\Delta MAP_T$ was calculated using the minimum to

maximum MCAv and MAP for each RSS performed at 0.05 Hz and 0.10 Hz. We compared

averaged $\Delta MCAv_T/\Delta MAP_T$ during MAP increases and decreases in 74 healthy subjects [9

women; 32 ± 13 years]. $\Delta MCAv_T/\Delta MAP_T$ was lower for MAP increases than MAP

decreases at 0.10 Hz RSS only (0.91 ± 0.34 vs. 1.01 ± 0.44 cm·s^{-1}/mmHg; $p = 0.0013$). For

both frequency and MAP direction, time during RSS had no effect on $\Delta MCAv_T/\Delta MAP_T$.

This novel analytical method supports the use of the RSS model to evaluate the directional

sensitivity of the pressure-flow relationship. These results contribute to the importance

of considering the direction of MAP changes when evaluating dynamic cerebral

autoregulation.

66
67
68 **News & Noteworthy**
69
70 Repeated squat-stand maneuvers are able to examine the directional sensitivity of the

71 cerebral pressure-flow relationship. These maneuvers induce stable physiological cyclic

72 changes where brain blood flow changes with blood pressure increases are buffered more

73 than decreases. These results highlight the importance of considering directional blood

74 pressure changes within cerebral autoregulation.

75
76 **Key words**

78 hysteresis, mean arterial pressure, middle cerebral artery blood velocity, squat-stand,

79 cerebral autoregulation

Introduction

Dynamic cerebral autoregulation (dCA) defines the ability of the cerebrovasculature to react to rapid blood pressure changes. Evidence suggests asymmetrical responses of cerebral blood flow (CBF) to transient changes in mean arterial pressure (MAP). Specifically, the augmentation in CBF is attenuated when MAP acutely increases compared to the decline in CBF when MAP acutely decreases (1, 8, 27, 36). Since the brain is located in the skull, an inflexible structure, this phenomenon is thought to serve as a protective mechanism against blood pressure surges (39). This notion has been previously highlighted as the cerebrovasculature is more efficient at buffering changes in blood pressure associated with the systolic aspect of the cardiac cycle (11, 32).

Initially, the above-mentioned asymmetrical responses of CBF to acute changes in MAP have been reported using cyclic inflation and deflation of thigh cuffs in patients with head injury (1) and pharmacological interventions in healthy participants (36). It is important to ensure the hysteresis-like pattern of the cerebral pressure-flow relationship exists during non-pharmacologically and physiologically induced acute increases and decreases in MAP in health and disease. Accordingly, the development of a valid non-pharmacological model to examine the directional sensitivity of the cerebral pressure-flow relationship is imperative.

Unfortunately, the previously used thigh cuff deflation technique may not represent the best model to investigate the directional sensitivity of the cerebral pressure-flow relationship, considering the poor reproducibility associated with dCA metrics quantified with this technique (reviewed in (29)). In the cerebrovascular research community, a repeated squat-stand protocol of 5 min is recognized as a reproducible method for inducing large blood pressure oscillations aimed at improving the linear interpretability of the transfer function metrics (29). This 5-min period of repeated squat-stands has been commonly used to examine the relationship between cerebral blood velocity (CBV) and

MAP in the frequency domain at two different repeated squat-stands frequencies (0.05 and 0.10 Hz) with the transfer function analysis, an analytical method which does not consider the direction of MAP (15). These two squatting frequencies are utilized since they are included in the frequency bands where dCA is thought to have the most important influence on the cerebral pressure-flow dynamics (historically, these frequency bands being 0.02-0.07 Hz for the very low frequency and 0.07-0.20 for the low frequency) (15). Also, these two frequencies are most prevalent in the literature (7, 10-12, 14, 18, 21, 22, 25, 28-31). Interestingly, the repeated squat-stand maneuver induces a cyclic physiological stress that can also be utilized to evaluate the CBV-MAP relationship in the time domain (6, 8, 27). When comparing the relative change in middle cerebral artery mean blood velocity (MCAv) per relative change in MAP during repeated squat-stands, our team previously demonstrated the directional sensitivity of the cerebral pressure-flow relationship (8). Using the autoregulation index (ARI), Panerai et al. demonstrated a better autoregulatory capacity during the squatting phase (transient increase in MAP) than the standing phase (transient decrease in MAP) (27). These results support the usefulness of analyzing MAP oscillations induced by the repeated squat-stand model in the time domain to evaluate the directional sensitivity of the cerebral pressure-flow relationship. Furthermore, the 5-min repeated squat-stand model allows for the average of several induced responses. Thus, providing a more reliable estimate of the physiological response than the utilization of only one transient response for each MAP direction as had previous be performed with the rate-of-regulation (2) and the autoregulatory index (34) measures. Additionally, we need to assume the metric used to characterize the hysteresis-like pattern during each squat-stand repetition is comparable over the 5 min, which is presently unknown.

In our previous analysis, we characterized the directional sensitivity of the cerebral pressure-flow relationship using the change in MCAv in response to transient increases versus decreases in MAP induced by repeated squat-stands. Specifically, we averaged the percent change in MCAv (from baseline) per percent change in MAP (from baseline)

(%ΔMCAv/%ΔMAP) calculated for each squat-stand executed over a period of 5 min at

the frequency of interest (0.05 and 0.10 Hz). These relative changes were calculated from

a seated baseline obtained prior-to the repeated squat-stand maneuver (8). However, the

utilization of a separate resting period as the baseline value has been recently criticized

(6). Barnes et al. argued the magnitude of MAP oscillations induced by repeated squat-

stands is greater when utilizing resting seated MAP as a reference value compared to a

resting standing MAP (6). In fact, they reasoned that the decrease in MAP from a resting

seated position to the minimum reached during a stand, is greater and would explain the

hysteresis pattern. These authors suggested using a reference value in the standing

position immediately before the repeated squat-stand protocol would be more valid.

While we agree calculating a baseline value from a separate control recording may be

misleading, the utilization of a reference value from a prolonged standing position

immediately before the beginning of the repeated squat-stand maneuver can also be

problematic. In this case, the increase in MAP from a resting quiet-stance position to the

maximum reached during a squat would be greater than the MAP decrease and balance

the difference between both. In addition, a prolonged period of standing, necessary to

reach a steady state of measured variables, has the potential to lead to orthostatic

hypotension in some individuals. In light of these issues, we consider there is no real

baseline we can employ for this particular protocol as there is not a truly stable

physiological reference point we can compare the changes during the squat-stand

maneuvers to.

A notion we did not consider in our previous study was the time intervals when the

transition of MCAv and MAP took place, which represents an additional issue with this

work (8). There are delays in physiological responses and regulating mechanisms vary

whether changes in MCAv and MAP are increasing or decreasing (1). Therefore, we

cannot assume repeated squat-stands induce MCAv and MAP changes of comparable

time intervals when MAP acutely decreases and increases via repeated squat-stands

induced at 0.05 Hz or 0.10 Hz. Thus, upward (from minimum to maximum) and downward

(from maximum to minimum) changes in MCAv/MAP of similar amplitudes, but occurring

across different time intervals, would indicate vessels are reacting differently according

to the direction of the MAP changes. We also believe taking into consideration time

intervals when MCAv and MAP changes are taking place would be more appropriate in

order to compare our metric between RSS frequencies (0.05 vs. 0.10 Hz).

The prevalent modeling techniques utilized for dCA quantification with spontaneous or

driven blood pressure oscillations do not take the directionality of blood pressure changes

into consideration (i.e. autoregulation index, transfer function analysis of spontaneous

and forced blood pressure oscillations). In other words, these approaches assume the

cerebrovasculature reacts similarly when MAP acutely increases or decreases.

Confirmation of asymmetrical sensitivity of the cerebral pressure-flow relationship using

the repeated squat-stand model thus has the potential to refine and optimize the current

methods of dCA assessment (15). If it becomes clear this hysteresis-like pattern of the

cerebral pressure-flow relationship is a common characteristic of the healthy

cerebrovasculature, which could be influenced by physiological and pathological

conditions, there will be a need for a global change in the assessment of dCA in order to

take MAP direction into account. However, the optimal physiological approach to study

this phenomenon remains to be elucidated.

For these reasons, the primary aim of this study was to refine a metric using the largest

concurrent MAP oscillations induced by the repeated squat-stand maneuver without

using an independent baseline value (either seated or standing). Accordingly, we

calculated this metric by using the minimum to maximum MCAv and MAP (in response to

squatting) compared to the maximum to minimum MCAv and MAP (in response to

standing) for each squat-stand transition taking time intervals when the MCAv and MAP

transition took place into account. The secondary aim of this study was to examine

whether the novel metric during each squat-stand transition at both 0.05 and 0.10 Hz

were comparable between each other over the 5-min period. We hypothesized the metric

would be lower when MAP acutely increases compared to when MAP acutely decreases,

and this 5-min repeated squat-stand model is an adequate protocol for the

characterization of the metric since it will be comparable between all repeated squat-

stand transitions.

Material and methods

Participants

Seventy-four participants were included in this analysis. Participants had previously been

involved in research studies at one of two institutions (Université Laval, Canada;

University of British Columbia, Canada). All participants provided informed written

consent before participating to these studies. The original research studies were

approved by the *Comité d'éthique de la recherche de l'Institut universitaire de cardiologie*

et de pneumologie de Québec (CER : 20869 and 21180) and the University of British

Columbia Clinical Review Ethical Board (CER: H11-02576, H11-03287) according to the

principles established by the Declaration of Helsinki (except for registration in a

database). Since the current study included a secondary analysis from previous

institutional review board approved studies and the use of de-identified data sharing, this

study did not require additional institutional review board review. All the participants

were healthy, free from medical condition and had no history of cardiorespiratory or

cerebrovascular disease. Apart from oral contraceptives in some women, participants

were not taking any blood pressure-altering medication. Women included in the study

were either taking oral contraceptive continuously for more than a year (n=2), wearing

an intrauterine device (n=1) or were tested during menses or the early follicular phase

(day 1 to 10) of their menstrual cycle (n=6).

Experimental protocol

The experimental protocol has been described previously (8). Participants arrived fasted

at the laboratory around 7AM and had a standardized snack (granola bar and juice). An

initial 5-min seated baseline recording was made for all participants for baseline

characteristics. Following a 5-min standing rest, the participants performed repeated

squat-stands for a period of at least 5 min at a frequency of 0.05 Hz (10-s squat, 10-s

standing) and 0.10 Hz (5-s squat, 5-s standing). As previously mentioned, these

frequencies were selected since they are within the range where dCA is thought to have

its greatest influence on the cerebral pressure-flow dynamics (40). Familiarization with

the squat-stand movement was done by mirroring the experimenter before the

experimentation. During recording, the experimenter ensured the transition speed and

depth was similar throughout the squat-stand repetition protocol for each participant.

241
Instrumentation

Maximal oxygen consumption (VO_{2max}) was determined during a preliminary visit in 53

participants. All these participants performed a graded ramp exercise test up to

exhaustion on an electromagnetically braked upright cycling ergometer. After 1 min of

unloaded pedaling, the workload increased progressively (20-30 W/min) until volitional

exhaustion. A gas exchange analysis system, combining fast responding oxygen and

carbon dioxide sensors and a pneumotach (Breezesuite, MedGraphics Corp., MN, USA;

Ultima™ CardiO2® Gas Exchange Analysis System, MGC Diagnostics®, MN, USA),

continuously recorded expired air to determine VO_2. VO_{2max} was determined as the

highest 30-s mean VO_2 coincident with a respiratory exchange ratio equivalent or greater

to 1.15.

253
The day of the study, MAP was measured with a finger photoplethysmograph with a

height correcting unit for correction of values (Finometer Pro, Finapres Medical Systems,

Amsterdam, The Netherlands or Nexfin, Edwards Lifesciences, Irvine, CA, USA). MCAv was

measured with a transcranial Doppler ultrasound (Spencer Technologies, Seattle, WA or

Doppler Box, Compumedics DWL USA, San Juan Capistrano, CA) on left temporal window.

The MCA was identified using a standardized protocol (38). Once the MCA was identified,

a probe was fixed in place with a headband and adhesive ultrasonic gel.

Partial pressure of end-tidal carbon dioxide ($P_{ET}CO_2$) was recorded beath-by-breath in 48
participants during all the repeated squat-stand through a gas analyzer with fast
responding oxygen and carbon dioxide sensors and a pneumotach (Breezesuite,
MedGraphics Corp., MN, USA; Ultima™ CardiO2® Gas Exchange Analysis System, MGC
Diagnostics®, MN, USA; ML206, AD Instruments, Colorado Springs, CO) calibrated to
known gas concentrations following manufacturer instructions before each evaluation.
The analyzer was adjusted for room temperature, atmospheric pressure and percentage
of humidity the day of testing.

All data were simultaneously sampled at 1000 Hz with an analog-to-digital converter
(Powerlab 16/30 ML880, AD Instruments) and saved for later analysis using commercially
available software (LabChart version 7.1, AD Instruments). There was no delay in
recording variables from either transcranial Doppler ultrasound units. In addition, we did
not apply any time shift in LabChart. However, to account for established delays in our
blood pressure monitoring devices, we applied a -250- and -1,000-ms time shift to ensure
Nexfin and Finometer, respectively, and transcranial Doppler ultrasound data were time
aligned. Any delays within the data collection units were negated when the structured
time-aligned analysis was performed. $P_{ET}CO_2$ was time-aligned with the other signals.

Data analysis
To characterize the cerebral pressure-flow relationship in response to transient increases
and decreases in MAP, we calculated a time-adjusted ratio between MCAv and MAP
changes for each squat-stand transition in each MAP direction. To do so, absolute changes
in MCAv ($\Delta MCAv_T$) and MAP (ΔMAP_T) for each increase (from minimum to maximum) and
decrease (from maximum to minimum) were calculated and adjusted for time intervals
when the MCAv and MAP transition took place. Specifically, we divided the changes by
the duration of the transition (i.e. time interval between minimum to maximum or
between maximum to minimum). Then, we averaged $\Delta MCAv_T/\Delta MAP_T$ for each

individual's repeated squat-stands over the 5-min recording period and for each
frequency.

$\Delta MCAv_T/\Delta MAP_T$ during transient decreases in MAP was calculated between a maximum
and the following minimum as follows (Figure 1):

293
$$\frac{\dfrac{(MCAv_{min} - MCAv_{max})}{(Time_{MCAvmin} - Time_{MCAvmax})}}{\dfrac{(MAP_{min} - MAP_{max})}{(Time_{MAPmin} - Time_{MAPmax})}} = \frac{\dfrac{\Delta MCAv}{\Delta Time_{MCAv}}}{\dfrac{\Delta MAP}{\Delta Time_{MAP}}} = \frac{\Delta MCAv_T}{\Delta MAP_T}$$

295    $\Delta MCAv_T/\Delta MAP_T$ during transient increases in MAP was calculated between a minimum
296   and the following maximum as follows (Figure 1):

298
$$\frac{\dfrac{(MCAv_{max} - MCAv_{min})}{(Time_{MCAvmax} - Time_{MCAvmin})}}{\dfrac{(MAP_{max} - MAP_{min})}{(Time_{MAPmax} - Time_{MAPmin})}} = \frac{\dfrac{\Delta MCAv}{\Delta Time_{MCAv}}}{\dfrac{\Delta MAP}{\Delta Time_{MAP}}} = \frac{\Delta MCAv_T}{\Delta MAP_T}$$

Statistical analysis
The normality of distributions was tested using Shapiro-Wilk tests. If distributions were
normal, a paired t-test was used. If not, the Wilcoxon matched pairs signed rank test was
used to compare variables between transient increases and decreases for each frequency.
To test the influence of time on $\Delta MCAv_T/\Delta MAP_T$ over the 5-min recording, a
nonparametric repeated-measures Friedman test was performed. Since this test does not
support missing values, all participants with missing values for one or more transitions
were excluded. Considering the impact of cardiorespiratory fitness and age on the
cerebral pressure-flow relationship (4, 5, 22, 24), correlations between $\Delta MCAv_T/\Delta MAP_T$
and age or VO_2max were determined using Spearman's rho correlations. P values < 0.05
were considered statistically significant.

Results

A total of 74 participants were included in this analysis (9 women, 65 men). For the 0.05

Hz analysis, 6 participants were excluded due to inconsistency of MAP and MCAv

oscillations (n=5) and poor recordings of the Nexfin (n=1). For the 0.10 Hz analysis, 3

participants were excluded because squats were not completed (n=1) and poor

recordings of the Nexfin (n=2). Final samples for 0.05 Hz and 0.10 Hz repeated squat-

stands were 68 and 71, respectively. VO_2max was determined in 53 participants. Baseline

characteristics as well as resting hemodynamic and cerebrovascular seated baseline

values are shown in Table 1. Seated baseline MAP and MCAv were missing for 4

participants.

Absolute changes in MAP during acute increases in MAP were $+43 \pm 15$ mmHg (0.05 Hz)

and $+37 \pm 14$ mmHg (0.10 Hz), whereas they were -43 ± 15 mmHg (0.05 Hz) and -37 ± 14

mmHg (0.10 Hz) during acute decreases in MAP. Absolute changes in MCAv during acute

increases in MAP were $+36 \pm 14$ cm·s^{-1} (0.05 Hz) and $+32 \pm 11$ cm·s^{-1} (0.10 Hz), whereas

they were -35 ± 14 cm·s^{-1} (0.05 Hz) and -32 ± 11 cm·s^{-1} (0.10 Hz) during acute decreases

in MAP. Averaged $P_{ET}CO_2$ during the 5-min squat-stand repetitions were not different

between frequency (0.05 Hz: 38.7 ± 3.8 Torr, 0.10 Hz: 38.2 ± 3.4 Torr; p = 0.0916) and

similar to baseline resting values (Table 1).

At 0.05 Hz, averaged time intervals for MAP and MCAv changes were shorter during acute

MAP increases than MAP decreases (MAP: 9.0 ± 1.0 vs. 11.0 ± 1.0 sec; p < 0.0001; MCAv:

9.3 ± 1.5 vs. 10.7 ± 1.4 cm·s^{-1}; p < 0.0001). At 0.10 Hz, averaged time intervals for MAP

and MCAv changes were shorter during acute increases than decreases for MAP ($4.9 \pm$

0.4 vs. 5.2 ± 0.4 sec; p = 0.0024), but not for MCAv (5.1 ± 0.5 vs. 4.9 ± 0.5 sec; p = 0.1028).

For 0.05 Hz repeated squat-stands, $\Delta MCAv_T/\Delta MAP_T$ was similar during acute increases

and acute decreases in MAP (0.88 ± 0.28 vs. 0.90 ± 0.23 cm·s^{-1}/mmHg; p = 0.2693; Figure

2). For 0.10 Hz repeated squat-stands, $\Delta MCAv_T/\Delta MAP_T$ was attenuated during acute increases when compared to acute decreases in MAP (0.91 ± 0.34 vs. 1.01 ± 0.44 cm·s^{-1}/mmHg; p = 0.0013; Figure 2).

For both frequency and MAP direction, time during repeated squat-stands had no effect on $\Delta MCAv_T/\Delta MAP_T$ (all p> 0.05; Figure 3).

Cardiorespiratory fitness was correlated to $\Delta MCAv_T/\Delta MAP_T$ for 0.05 Hz repeated squat-stands during acute MAP increases ($\rho = 0.33$, p = 0.0244) but not for acute MAP decreases ($\rho = 0.21$, p = 0.1566). Cardiorespiratory fitness was correlated to $\Delta MCAv_T/\Delta MAP_T$ for 0.10 Hz repeated squat-stands during both acute increases ($\rho = 0.49$, p = 0.0003) and decreases ($\rho = 0.48$, p = 0.0005) in MAP (Figure 4). At 0.05 Hz, age was not correlated to $\Delta MCAv_T/\Delta MAP_T$ during acute MAP increases ($\rho = -0.19$, p = 0.1278) nor acute decreases in MAP ($\rho = -0.17$, p = 0.1581). At 0.10 Hz, age was not correlated to $\Delta MCAv_T/\Delta MAP_T$ during acute MAP increases ($\rho = -0.21$, p = 0.0803), but was inversely correlated during acute MAP decreases ($\rho = -0.25$, p = 0.0346) (Figure 5).

Discussion

This study demonstrates when MAP is acutely increased from minimum to maximum during repeated squat-stands at 0.10 Hz, the change in MCAv is attenuated compared to when MAP acutely decreases, which is in line with our previous results (8). The novel finding of this study resides in the notion that changes were adjusted for time intervals and calculated on actually induced MAP oscillations, between maximums and minimums *during* the maneuvers, instead of being calculated from a seated baseline value obtained *prior-to* the repeated squat-stands. These results reinforce the evidence of hysteresis in the relationship between MCAv and MAP, indicating cerebral vessels are better adapted to buffer acute increases in MAP at 0.10 Hz, highlighting a probable influence of repeated squat-stand frequency. In addition, our findings convincingly demonstrate the repeated

squat-stand model can provide a reliable estimate of the physiological response by averaging several transient responses, considering the physiological stress imposed by each squat-stand repetition leads to comparable $\Delta MCAv_T/\Delta MAP_T$ (i.e. when the metric for each acute increase in MAP is compared) over the 5-min period, at least in the studied population.

In light of the questions regarding our previous analysis (6), it was important to improve our initial analysis in order to address different concerns, i.e. to avoid the utilization of a separate resting period and take time intervals of MCAv and MAP changes into consideration. To the best of our knowledge, this is the first study to report the presence of a directional sensitivity of the human MCAv-MAP relationship when changes are adjusted for time intervals as well as calculated between each maximums and minimums reached during 0.10 Hz, but not 0.05 Hz, for repeated squat-stands. The results of the present analysis demonstrate the use of actual MAP changes, instead of calculating MAP changes from a separate baseline period, support the presence of a frequency-dependent directional sensitivity of the cerebral pressure-flow relationship in healthy participants. Our study also reinforces the use of the 5-min squat-stand protocol to analyze the MCAv-MAP relationship in the time domain. In fact, as demonstrated in Figure 3, the repetition of the same stimulus over 5 minutes produces valid and stable data over time, reducing the potential influence of outliers.

Although the current study did not aim at examining the mechanisms underlying the asymmetrical sensitivity of the cerebral pressure-flow relationship, multiple hypotheses are discussed in the literature. First, it was shown in an animal model that cerebral sympathetic nervous activity (SNA) recorded in the superior cervical ganglia increases in response to mechanically or drug-induced increases in arterial pressure, but not during transient reduction in arterial pressure (13). These results would indicate cerebral SNA increases in response to an acute elevation in MAP to protect the brain from overperfusion. Interestingly, it is proposed that 0.10 Hz repeated squat-stand frequency

might reveal the influence of SNA on human cerebrovascular regulation (20, 40).

Therefore, the lower $\Delta MCAv_T/\Delta MAP_T$ during acute increases in MAP at 0.10 Hz repeated

squat-stands could be explained by SNA activation when MAP acutely increases, in order

to protect cerebral vessels. However, the impact of cerebral SNA on CBF in humans is still

controversial for many reasons (reviewed in (3, 9)). Below frequencies of 0.07 Hz,

cerebrovascular regulation is thought to have a myogenic influence (20, 40). Another

potential mechanism could therefore be a differing intrinsic myogenic activity when MAP

increases than when MAP decreases. Using oscillatory lower body negative pressure and

a non-linear approach to quantify dCA, it was demonstrated the utilization of calcium-

channel blockade impairs dCA in humans, indicating a role for myogenic mechanisms at

frequencies between 0.03 and 0.08 Hz (33). However, these results showed only a global

dCA response since the directionality of MAP changes were not considered (33). As MAP

direction did not influence $\Delta MCAv_T/\Delta MAP_T$ at 0.05 Hz repeated squat-stands in the

present study, it could be argued the myogenic influence on CBF is similar whether MAP

acutely increases or decreases, or that our linear approach masks differences. Recently,

Barnes et al. were interested in different components of the cerebrovascular response to

repeated acute increases and decreases in MAP during the repeated squat-stand

protocol. They demonstrated MAP would be the strongest determinant of CBV when

squatting down, overwhelming the myogenic and metabolic components of dCA (6).

Overall, these results also make the myogenic mechanism a controversial mechanism to

explain the presence of a hysteresis-like pattern in the cerebral pressure-flow relationship

(33). The absence of directional sensitivity at 0.05 Hz repeated squat-stands could also be

explained by the baroreflex latency (19, 36). In fact, during these slower oscillations,

baroreceptors have time to sense pressure loading or unloading and induce appropriate

cardiovascular adjustments. This could mitigate the impact of MAP on MCAv over a 10-

sec time interval. Finally, a transient increase in MAP could elevate superior vena cava's

resistance, inducing an increase in intracranial pressure and therefore, a reduction of the

passive pressure gradient for venous drainage (16, 17), which could represent evidence

of the Monro-Kellie doctrine, i.e. a mechanism limiting the amount of CBF increase that

is able to occur within the encapsulated skull (39).

The consistency of the $\Delta MCAv_T/\Delta MAP_T$ over the 5-min repeated squat-stand at both

frequencies is indicative the physiological stress induced is stable for both direction and

magnitude of acute changes in MAP. The absence of a drift in the values demonstrates

the participants performing repeated squat-stands do not get fatigued from the

maneuvers and the addition of squat-stand transitions has no influence on the metric, at

least in healthy participants.

The relationships between $\Delta MCAv_T/\Delta MAP_T$ and cardiorespiratory fitness during acute

increases and decreases in MAP at 0.10 Hz, and only during acute increases at 0.05 Hz

repeated squat-stand protocol, are also interesting observations. These findings suggest

a diminished ability to cope with higher frequency MAP oscillations with increasing

cardiorespiratory fitness. We have previously demonstrated an association between

cardiorespiratory fitness and TFA gain, normalized gain and phase during 0.10 Hz

repeated squat-stands in young and healthy fit men (22). TFA gain metrics represent

changes in MCAv for a given change in MAP, which makes logical that $\Delta MCAv_T/\Delta MAP_T$ is

also correlated to cardiorespiratory fitness in the current analysis. However, the fact that

the current analysis using the $\Delta MCAv_T/\Delta MAP_T$ metric takes the directionality of MAP

changes into account is notable. The reason for an augmented $\Delta MCAv_T/\Delta MAP_T$ during

acute increases and decreases in MAP, (which could be interpreted as a diminished ability

to buffer MAP changes in both directions, i.e. a lowered dCA), with increasing

cardiorespiratory fitness and whether this is deleterious or beneficial for the brain, is still

unknown. Further studies are needed to evaluate the impact of improving

cardiorespiratory fitness on the maintenance of the hysteresis-like pattern in the MCAv-

MAP relationship.

In the current analysis, age inversely correlates with $\Delta MCAv_T/\Delta MAP_T$ when MAP acutely

decreases during 0.10 Hz repeated squat-stands. These findings contradict results from

our previous work (8). Indeed, when calculating relative changes from a seated baseline

obtained prior to the repeated squat-stand protocol, age was positively correlated with

$\%\Delta MCAv/\%\Delta MAP$ when MAP acutely increased for 0.05-Hz repeated squat-stand only.

As in our previous study, the correlational analysis included here is exploratory in nature.

Statistical power may have been insufficient to address that question. In addition, while

we have included a group of participants with a large range in age, only a small number

of subjects were above 40 years old, which considerably reduce the strength of this

analysis. Further well-designed research aimed at specifically examining the influence of

age on the directional sensitivity of the cerebral pressure-flow relationship is thus

necessary to address that specific question.

Perspectives

Overall, our updated analysis for repeated squat-stands is adequate to study the

directional sensitivity of the cerebral pressure-flow relationship and optimize the

assessment of dCA. It supports the need of a global change in dCA assessment in order to

take the direction of MAP into account. It also reinforces the importance of using the

repeated squat-stands, at different frequencies, to provide a broader characterization of

the cerebral pressure-flow relationship. In addition, since we showed repeated squat-

stands induce a comparable change in $\Delta MCAv_T/\Delta MAP_T$ at each squat-stand transition, it

eliminates the questioning as to whether the addition of squat-stands (exercise) would

have an impact *per se* on this metric. A next important step of major interest would be to

evaluate the reproducibility of this metric over time. It would help determine whether

the $\Delta MCAv_T/\Delta MAP_T$ is reliable over various time frames and if it remains stable over time.

Still, further research is needed to examine the possibility of using less repetitions to

calculate the $\Delta MCAv_T/\Delta MAP_T$ metric. Indeed, a diminished number of squat-stand

repetitions would be an important methodological advantage to examine for the

presence of hysteresis in clinical populations for whom a 5-min period of repeated squat-

stands could potentially be difficult to complete.

The directional sensitivity in the cerebral pressure-flow relationship is highly relevant for

various physiological (rapid eye movement sleep, exercise), and pathological clinical

situations associated with MAP surges. Nonetheless, further research is needed to

describe this phenomenon in other populations and clinical conditions. We still do not

know whether this hysteresis-like pattern remains present in different physiological and

pathological states (arterial/carotid stiffness, uncontrolled systemic hypertension,

autonomic dysreflexia, etc.). For example, findings from a recent study using a mouse

model of arterial stiffness suggest mice with calcified carotid artery have a disrupted

cerebral autoregulation in response to steady-state MAP increases compared to healthy

animals (26). The next logical step would therefore be to examine whether there is

evidence for hysteresis in the cerebral pressure-flow relationship in human clinical

populations and the present study serves as a fundamental basis to further explore the

phenomenon. Importantly for the clinician, this directional sensitivity of brain vessels to

changes in MAP should also be integrated in the framework to guide its management in

the operating room. Finally, considering the link found with cardiorespiratory fitness in

the present study, it would be interesting to evaluate the impact of exercise training on

the maintenance of the hysteresis-like pattern.

Methodological considerations

There are some limitations to this study that need to be further discussed. Only healthy

men and women were included in this analysis and the results cannot be generalized to

clinical populations. Moreover, MCAv was evaluated through transcranial Doppler

ultrasound, which is known to be representative of flow only if the diameter of the artery

is assumed to remain stable. Changes in $P_{ET}CO_2$ have been associated with MCA's

diameter variations (23, 37). Although $P_{ET}CO_2$ was not available for a part of the male

participants, there was no difference between $P_{ET}CO_2$ during baseline and repeated

squat-stand for the participants in whom it was measured. In addition, it was previously

observed that there was no change in $P_{ET}CO_2$ between squat and stand phases (27).

Therefore, $P_{ET}CO_2$ had most likely a minor effect on our results. Only a small part of the

group was composed of women either taking oral contraceptives, wearing an intrauterine

device or being in the days 1-10 of their menstrual cycle. We are not able to determine

whether it had an influence on the results and further research is needed to address that

issue. Since days 0-10 represents a large time period, it should be shortened in future

studies in order to better control for menstrual cycle. Of note, all participants (n=58)

included in our previous work (8) were also included in the current analysis. From an

analytical standpoint, the two approaches are comparable, as both methods are

calculating changes in MCAv and MAP in response to repeated squat-stands. One

difference resides in the fact the previous method calculated relative changes from a

seated baseline, whereas the revised method calculated actual absolute changes adjusted

for time intervals. This being acknowledged, the main difference resides in the *baseline*

value used to calculate these changes in MCAv and MAP. In the original approach, we

have calculated a *baseline* including averaged MCAv and MAP during a steady-state rest.

However, this *baseline* with participants in a seated position was measured during a

distinct recording period prior to the start of the series of repeated squat-stands.

Accordingly, from a physiological standpoint, it makes definitely more sense to calculate

changes in MCAv and MAP on actually induced MAP oscillations, between maximums and

minimums during the repeated squat-stands, rather than being calculated from a seated

baseline value obtained during a distinct period prior to the repeated squat-stands. We

consider the updated analytical method proposed in the current work to be the most valid

option to examine the directional sensitivity of the cerebral pressure-flow relationship

with the repeated squat-stand model since our calculated metric cannot be influenced by

a separate resting steady-state baseline. However, the two analytical methods were not

directly compared, as our main goal of the current investigation was to examine whether

the hysteresis-like pattern in the cerebral pressure-flow relationship was present with the

updated analytical method. A direct comparison would not provide additional meaningful

information.

Conclusion

Our findings indicate that this novel analytical method supports the use of the repeated

squat-stand model to examine the directional sensitivity of the cerebral pressure-flow

relationship, during actual changes in MAP and cerebral blood velocity, without the need

for a separate resting baseline. Moreover, it shows this maneuver induces stable and

robust physiological cyclic changes and allows for interindividual variability analysis.

Importantly, the results contribute to the importance of considering the direction of MAP

changes when evaluating dCA and builds upon the prior rate-of-regulation and

autoregulatory index measures.

Acknowledgments

We would like to thank Myriam Paquette, Olivier Le Blanc, Kevan Rahimaly, Sarah Imhoff,

Simon Malenfant and Audrey Drapeau for their assistance in data collection.

Grants

This study has been supported by the Ministère de l'Éducation, du Loisir et du Sport du

Québec, the Foundation of the Institut universitaire de cardiologie et de pneumologie de

Québec and the Natural Sciences and Engineering Research Council. L.L. is supported by

a doctoral training scholarship from the Canadian Institutes of Health Research.

References

1. Aaslid R, Blaha M, Sviri G, Douville CM, Newell DW. Asymmetric dynamic cerebral autoregulatory response to cyclic stimuli. *Stroke* 38: 1465–1469, 2007.

2. Aaslid R, Lindegaard KF, Sorteberg W, Nornes H. Cerebral autoregulation dynamics in humans. *Stroke* 20: 45–52, 1989.

3. Ainslie PN, Brassard P. Why is the neural control of cerebral autoregulation so controversial? *F1000Prime Rep* 6: 14, 2014.

4. Ainslie PN, Cotter JD, George KP, Lucas SJE, Murrell C, Shave R, Thomas KN, Williams MJA, Atkinson G. Elevation in cerebral blood flow velocity with aerobic fitness throughout healthy human ageing. *J Physiol* 586: 4005–4010, 2008.

5. Bailey DM, Marley CJ, Brugniaux JV, Hodson D, New KJ, Ogoh S, Ainslie PN. Elevated aerobic fitness sustained throughout the adult lifespan is associated with improved cerebral hemodynamics. *Stroke* 44: 3235–3238, 2013.

6. Barnes SC, Ball N, Haunton VJ, Robinson TG, Panerai RB. The cerebrocardiovascular response to periodic squat-stand maneuvers in healthy subjects: a time-domain analysis. *AJP Heart Circ Physiol* 313: H1240–H1248, 2017.

7. Birch AA, Dirnhuber MJ, Hartley-Davies R, Iannotti F, Neil-Dwyer G. Assessment of autoregulation by means of periodic changes in blood pressure. *Stroke* 26: 834–837, 1995.

8. Brassard P, Ferland-Dutil H, Smirl JD, Paquette M, Le Blanc O, Malenfant S, Ainslie PN. Evidence for hysteresis in the cerebral pressure-flow relationship in healthy men. *AJP Heart Circ Physiol* 312: H701–H704, 2017.

9. Brassard P, Tymko MM, Ainslie PN. Sympathetic control of the brain circulation: Appreciating the complexities to better understand the controversy. *Auton Neurosci* 207: 37–47, 2017.

10. Brothers RM, Zhang R, Wingo JE, Hubing KA, Crandall CG. Effects of heat stress on dynamic cerebral autoregulation during large fluctuations in arterial blood pressure. *J Appl Physiol* 107: 1722–1729, 2009.

11. Burma JS, Copeland P, Macaulay A, Khatra O, Wright AD, Smirl JD. Dynamic cerebral autoregulation across the cardiac cycle during 8 hr of recovery from acute exercise. *Physiol Rep* 8: e14367, 2020.

12. Burma JS, Macaulay A, Copeland P, Khatra O, Bouliane KJ, Smirl JD. Comparison of cerebrovascular reactivity recovery following high-intensity interval training and moderate-intensity continuous training. *Physiol Rep* 8: e14467, 2020.

13. Cassaglia PA, Griffiths RI, Walker AM. Sympathetic nerve activity in the superior
cervical ganglia increases in response to imposed increases in arterial pressure. *AJP*
Regul Integr Comp Physiol 294: R1255–61, 2008.

14. Claassen JAHR, Levine BD, Zhang R. Dynamic cerebral autoregulation during
repeated squat-stand maneuvers. *J Appl Physiol* 106: 153–160, 2009.

15. Claassen JAHR, Meel-van den Abeelen ASS, Simpson DM, Panerai RB, international
Cerebral Autoregulation Research Network (CARNet). Transfer function analysis of
dynamic cerebral autoregulation: A white paper from the International Cerebral
Autoregulation Research Network. *J Cereb Blood Flow Metab* 36: 665–680, 2016.

16. Dawson EA, Secher NH, Dalsgaard MK, Ogoh S, Yoshiga CC, Gonzalez-Alonso J,
Steensberg A, Raven PB. Standing up to the challenge of standing: a siphon does
not support cerebral blood flow in humans. *AJP Regul Integr Comp Physiol* 287:
R911–4, 2004.

17. Donnelly J, Budohoski KP, Smielewski P, Czosnyka M. Regulation of the cerebral
circulation: bedside assessment and clinical implications. *Crit Care* 20:129, 2016.

18. Drapeau A, Labrecque L, Imhoff S, Paquette M, Le Blanc O, Malenfant S, Brassard
P. Six weeks of high-intensity interval training to exhaustion attenuates dynamic
cerebral autoregulation without influencing resting cerebral blood velocity in
young fit men. *Physiol Rep* 7: e14185, 2019.

19. Fisher JP, Kim A, Young CN, Ogoh S, Raven PB, Secher NH, Fadel PJ. Influence of
ageing on carotid baroreflex peak response latency in humans. *J Physiol* 587: 5427–
5439, 2009.

20. Hamner JW, Tan CO. Relative contributions of sympathetic, cholinergic, and
myogenic mechanisms to cerebral autoregulation. *Stroke* 45: 1771–1777, 2014.

21. Labrecque L, Rahimaly K, Imhoff S, Paquette M, Le Blanc O, Malenfant S, Drapeau
A, Smirl JD, Bailey DM, Brassard P. Dynamic cerebral autoregulation is attenuated
in young fit women. *Physiol Rep* 7: e13984, 2019.

22. Labrecque L, Rahimaly K, Imhoff S, Paquette M, Le Blanc O, Malenfant S, Lucas SJE,
Bailey DM, Smirl JD, Brassard P. Diminished dynamic cerebral autoregulatory
capacity with forced oscillations in mean arterial pressure with elevated
cardiorespiratory fitness. *Physiol Rep* 5, 2017.

23. Lewis NCS, Smith KJ, Bain AR, Wildfong KW, Numan T, Ainslie PN. Impact of
transient hypotension on regional cerebral blood flow in humans. *Clin Sci* 129: 169–
178, 2015.

24.   Lind-Holst M, Cotter JD, Helge JW, Boushel R, Augustesen H, van Lieshout JJ, Pott
FC. Cerebral autoregulation dynamics in endurance-trained individuals. *J Appl*
Physiol 110: 1327–1333, 2011.

25.   Malenfant S, Brassard P, Paquette M, Le Blanc O, Chouinard A, Nadeau V, Allan PD,
Tzeng Y-C, Simard S, Bonnet S, Provencher S. Compromised Cerebrovascular
Regulation and Cerebral Oxygenation in Pulmonary Arterial Hypertension. *J Am*
Heart Assoc 6, 2017.

26.   Muhire G, Iulita MF, Vallerand D, Youwakim J, Gratuze M, Petry FR, Planel E,
Ferland G, Girouard H. Arterial Stiffness Due to Carotid Calcification Disrupts
Cerebral Blood Flow Regulation and Leads to Cognitive Deficits. *J Am Heart Assoc*
8: e011630, 2019.

27.   Panerai RB, Barnes SC, Nath M, Ball N, Robinson TG, Haunton VJ. Directional
sensitivity of dynamic cerebral autoregulation in squat-stand maneuvers. *AJP:*
Regulatory, Integrative and Comparative Physiology 315: R730-40, 2018.

28.   Smirl JD, Haykowsky MJ, Nelson MD, Tzeng Y-C, Marsden KR, Jones H, Ainslie PN.
Relationship between cerebral blood flow and blood pressure in long-term heart
transplant recipients. *Hypertension* 64: 1314–1320, 2014.

29.   Smirl JD, Hoffman K, Tzeng Y-C, Hansen A, Ainslie PN. Methodological comparison
of active- and passive-driven oscillations in blood pressure; implications for the
assessment of cerebral pressure-flow relationships. *J Appl Physiol* 119: 487–501,
2015.

30.   Smirl JD, Lucas SJE, Lewis NCS, duManoir GR, Smith KJ, Bakker A, Basnyat AS, Ainslie
PN. Cerebral pressure-flow relationship in lowlanders and natives at high altitude.
J Cereb Blood Flow Metab 34: 248–257, 2014.

31.   Smirl JD, Tzeng Y-C, Monteleone BJ, Ainslie PN. Influence of cerebrovascular
resistance on the dynamic relationship between blood pressure and cerebral blood
flow in humans. *J Appl Physiol* 116: 1614–1622, 2014.

32.   Smirl JD, Wright AD, Ainslie PN, Tzeng Y-C, van Donkelaar P. Differential Systolic
and Diastolic Regulation of the Cerebral Pressure-Flow Relationship During Squat-
Stand Manoeuvres. *Acta Neurochir Suppl* 126: 263–268, 2018.

33.   Tan CO, Hamner JW, Taylor JA. The role of myogenic mechanisms in human
cerebrovascular regulation. *J Physiol* 591: 5095–5105, 2013.

34.   Tiecks FP, Lam AM, Aaslid R, Newell DW. Comparison of static and dynamic cerebral
autoregulation measurements. *Stroke* 26: 1014–1019, 1995.

35.   Tzeng Y-C, Ainslie PN. Blood pressure regulation IX: cerebral autoregulation under
672   blood pressure challenges. *Eur J Appl Physiol* 114: 545–559, 2014.

673   36.   Tzeng Y-C, Willie CK, Atkinson G, Lucas SJE, Wong A, Ainslie PN. Cerebrovascular
674   regulation during transient hypotension and hypertension in humans.
675   *Hypertension* 56: 268–273, 2010.

676   37.   Verbree J, Bronzwaer A-SGT, Ghariq E, Versluis MJ, Daemen MJAP, van Buchem
677   MA, Dahan A, van Lieshout JJ, van Osch MJP. Assessment of middle cerebral artery
678   diameter during hypocapnia and hypercapnia in humans using ultra-high-field MRI.
679   *J Appl Physiol* 117: 1084–1089, 2014.

680   38.   Willie CK, Colino FL, Bailey DM, Tzeng Y-C, Binsted G, Jones LW, Haykowsky MJ,
681   Bellapart J, Ogoh S, Smith KJ, Smirl JD, Day TA, Lucas SJE, Eller LK, Ainslie PN. Utility
682   of transcranial Doppler ultrasound for the integrative assessment of
683   cerebrovascular function. *J Neurosci Methods* 196: 221–237, 2011.

684   39.   Wilson MH. Monro-Kellie 2.0: The dynamic vascular and venous pathophysiological
685   components of intracranial pressure. *J Cereb Blood Flow Metab* 36: 1338–1350,
686   2016.

687   40.   Zhang R, Zuckerman JH, Giller CA, Levine BD. Transfer function analysis of dynamic
688   cerebral autoregulation in humans. *Am J Physiol* 274: H233–41, 1998.

Figures caption

Figure 1. Panel A represents raw tracings of blood pressure (BP), middle cerebral artery blood velocity (MCAv) and partial pressure of carbon dioxide (PCO_2) at 0.05 and 0.10 Hz repeated squat-stand. Panel B depicts schematic tracings of mean arterial pressure (MAP) and middle cerebral artery mean blood velocity (MCAv) during a repeated squat-stands performed at 0.05 Hz. Enlarged regions depict how the variables were used to calculate $\Delta MCAv_T/\Delta MAP_T$. Blue color indicates variables used to calculate $\Delta MCAv_T/\Delta MAP_T$ during acute decreases in MAP. Red color indicates variables used to calculate $\Delta MCAv_T/\Delta MAP_T$ during acute increases in MAP.

Figure 2. Comparison of the $\Delta MCAv_T/\Delta MAP_T$ during transient increases (INC) and decreases (DEC) in mean arterial pressure during 0.05 and 0.10 Hz repeated squat-stands. n = 68 for 0.05 Hz, n = 71 for 0.10 Hz. Differences assessed via Wilcoxon matched pairs signed rank test.

Figure 3. Individual $\Delta MCAv_T/\Delta MAP_T$ for each squat repetition during acute increases (INC) and decreased (DEC) for 0.05 and 0.10 Hz repeated squat-stands. Numbers on the x-axis represent squat-stand repetitions. For each squat-stand repetition, n = 64 during INC and n = 56 during DEC for 0.05 Hz, n = 58 during INC and n = 59 during DEC for 0.10 Hz. The effect of time was tested using a nonparametric repeated-measures Friedman test.

Figure 4. Relationship between VO_2max and $\Delta MCAv_T/\Delta MAP_T$ for acute increases (INC) and decreases (DEC) in MAP during repeated squat-stands at 0.05 Hz and 0.10 Hz. n = 47 for 0.05 Hz, n = 50 for 0.10 Hz. Relationships between $\Delta MCAv_T/\Delta MAP_T$ and cardiorespiratory fitness (VO_2max) were determined using Spearman's rho correlations.

Figure 5. Relationship between age and $\Delta MCAv_T/\Delta MAP_T$ for acute increases (INC) and decreases (DEC) in MAP during repeated squat-stands at 0.05 Hz and 0.10 Hz. n = 68 for 0.05 Hz, n = 71 for 0.10 Hz. Relationships between $\Delta MCAv_T/\Delta MAP_T$ and age were determined using Spearman's rho correlations.

Table 1. Baseline characteristics and resting seated values

Baseline characteristics	
n	74
Age (years)	$32 \pm 13 \, (20 - 74)$
Height (m)	$1.75 \pm 0.09 \, (1.55 - 1.94)$
Body weight (kg)	$75.5 \pm 12.8 \, (53.0 - 115.0)$
Body mass index (kg/m^2)	$24.5 \pm 3.2 \, (19.0 - 36.7)$
Maximal oxygen consumption (ml·kg^{-1}·min^{-1})	$46.9 \pm 9.6 \, (23.5 - 65.4)$ *
Resting values	
Mean arterial pressure (mmHg)	$90 \pm 13 \, (60 - 122)$ [§]
Middle cerebral artery mean blood velocity (cm·s^{-1})	$62 \pm 13 \, (34 - 100)$ [§]
End-tidal carbon dioxide partial pressure (Torr)	$38 \pm 4 \, (29 - 47)$ [†]

752 Data are mean $\pm$ SD (ranges).

753 * n = 53, [§] n = 70, [†] n = 48

773
774 **Figure 1.**
775
776

777
778   **Figure 2.**

779
780 **Figure 3.**

781
782 **Figure 4.**

783
784 **Figure 5.**