Increased connectivity among sensory and motor regions during visual and audiovisual speech perception

Jonathan E. Peelle[1], Brent Spehar[1], Michael S. Jones[1], Sarah McConkey[1],
Joel Myerson[2], Sandra Hale[2], Mitchell S. Sommers[2], Nancy Tye-Murray[1]

[1] Department of Otolaryngology, Washington University in St. Louis, St. Louis MO USA

[2] Department of Psychological and Brain Sciences, Washington University in St. Louis, St. Louis MO USA

Abbreviated title: Connectivity during audiovisual speech perception

Conflicts of interest: The authors declare no competing financial conflicts of interest.

Acknowledgments: This work was supported by grants R56 AG018029 and R01 DC016594 from the US National Institutes of Health. The multiband echo planar imaging sequence was provided by the University of Minnesota Center for Magnetic Resonance Research.

Please address correspondence to:

Dr. Jonathan Peelle
Department of Otolaryngology
Washington University in Saint Louis
Saint Louis, MO 63110
jpeelle@wustl.edu

45   **Abstract**

In everyday conversation, we usually process the talker's face as well as the sound of
their voice. Access to visual speech information is particularly useful when the auditory
signal is degraded. Here we used fMRI to monitor brain activity while adult humans (n =
60) were presented with visual-only, auditory-only, and audiovisual words. The
audiovisual words were presented in quiet and several signal-to-noise ratios. As
expected, audiovisual speech perception recruited both auditory and visual cortex, with
some evidence for increased recruitment of premotor cortex in some conditions
(including in substantial background noise). We then investigated neural connectivity
using psychophysiological interaction (PPI) analysis with seed regions in both primary
auditory cortex and primary visual cortex. Connectivity between auditory and visual
cortices was stronger in audiovisual conditions than in unimodal conditions, including a
wide network of regions in posterior temporal cortex and prefrontal cortex. In addition to
whole-brain analyses, we also conducted a region-of-interest analysis on the left
posterior superior temporal sulcus (pSTS), implicated in many previous studies of
audiovisual speech perception. We found evidence for both activity and effective
connectivity in pSTS for visual-only and audiovisual speech, although these were not
significant in whole-brain analyses. Taken together, our results suggest a prominent role
for cross-region synchronization in understanding both visual-only and audiovisual
speech that complements activity in "integrative" brain regions like pSTS.

65
66
67

Introduction

Understanding speech in the presence of background noise is notoriously challenging,
and when visual speech information is available, listeners make use of it—performance
on audiovisual (AV) speech in noise is better than for auditory-only speech in noise
(Sumby and Pollack, 1954). Although there is consensus that listeners make use of
visual information during speech perception, there is little agreement either on the
neural mechanisms that support visual speech processing or on the way in which visual
and auditory speech information are combined during audiovisual speech perception.
One longstanding perspective on audiovisual speech has been that auditory and
visual information are processed through separate channels, and then integrated at a
separate processing stage (Grant and Seitz, 1998; Massaro and Palmer, 1998).
Audiovisual integration is thus often considered an individual ability that some people
are better at and some people are worse at, regardless of their unimodal processing
abilities (Magnotti and Beauchamp, 2015; Mallick et al., 2015).
However, more recent data have brought this traditional view into question. For
example, Tye-Murray and colleagues (2016) showed that unimodal auditory-only and
visual-only word recognition scores accurately predicted AV performance, and factor
analyses revealed two unimodal ability factors with no evidence of a separate
integrative ability factor. These findings suggest that rather than a separate stage of
audiovisual integration, AV speech perception may depend most strongly on the
coordination of auditory and visual inputs (Sommers, 2021).
Theoretical perspectives on audiovisual integration have also informed cognitive
neuroscience approaches to AV speech perception. Prior functional neuroimaging
studies of audiovisual speech processing have largely focused on identifying brain
regions supporting integration. One possibility is that the posterior superior temporal
sulcus (pSTS) combines auditory and visual information during speech perception. The
pSTS is anatomically positioned between auditory cortex and visual cortex, and has the
functional properties of a multisensory convergence zone (Beauchamp et al., 2004).
During many audiovisual tasks, the pSTS is differentially activated by matching and mis-
matching auditory-visual information, consistent with a role in integration (Stevenson
and James, 2009). Moreover, functional connectivity between the pSTS and primary
sensory regions varies with the reliability of the information in a modality (Nath and
Beauchamp, 2011), suggesting that the role of the pSTS may be related to combining or
weighing information from different senses.
A complementary proposal is that regions of premotor cortex responsible for
representing articulatory information are engaged in processing speech (Okada and
Hickok, 2009). The contribution of motor regions to speech perception is hotly debated.
Evidence consistent with a motor contribution includes a self-advantage in both visual-
only and AV speech perception (Tye-Murray et al., 2013, 2015), and effects of visual
speech training on speech production (Fridriksson et al., 2009; Venezia et al., 2016).
However, premotor activity is not consistently observed in neuroimaging studies of
speech perception, and in some instances may also reflect non-perceptual processing
(Szenkovits et al., 2012; Nuttall et al., 2016). It is also possible that premotor regions
are only engaged in certain types of speech perception situations (for example, when
there is substantial background noise, or when lipreading); individual differences in

hearing sensitivity or lipreading ability also may affect the involvement of premotor
cortex.
In addition to looking for brain regions that support visual-only or AV speech
perception, we therefore broaden our approach to study the role played by effective
connectivity between auditory, visual, and motor regions. If a dedicated brain region is
necessary to combine auditory and visual speech information, we would expect to see it
active during audiovisual speech. If changes in effective connectivity (Friston, 1994;
Stephan and Friston, 2010)—that is, task-based synchronized activity—underlie visual-
only or audiovisual speech processing, we would expect to see greater connectivity
between speech-related regions during these conditions relative to auditory-only
speech. In service of these questions we tested auditory-only speech perception and
AV speech perception at a range of signal-to-noise ratios (SNRs) and obtained out-of-
scanner measures of lipreading ability from our participants (**Figure 1**).
126
127

Figure 1. a. Experimental conditions with auditory-only speech, visual-only speech, and audiovisual speech. **b.** Histogram of lipreading abilities measured outside the scanner. **c.** Within-scanner behavioral performance (subjective ratings of understanding); individual participants shown in dots, mean ± SE displayed in bars.

128
129
130
131   **Method**

132 Stimuli, behavioral data, analysis scripts, and results tables are available from
133 https://osf.io/qxcu8/. MRI data are available on OpenNeuro (Markiewicz et al., 2021) at
134 https://doi.org/10.18112/openneuro.ds003717.v1.0.0.

Materials

We created seven lists of 50 words. The stimuli were recordings of a female actor speaking single words. The talker sat in front of a neutral background and spoke words along with the carrier phrase "Say the word _______" into the camera. The actor was instructed to allow her mouth to relax to a slightly open and neutral position before each target word was spoken. The edited versions of the recordings used in the current experiment did not include a carrier phrase and were each 1.5 seconds long. Recordings were made using a Canon Elura 85 digital video camera and showed the talker's head and shoulders. Digital capture and editing were done using Adobe Premiere Elements. The original capture format for the video was uncompressed .avi; the final versions used in the study were compressed as high quality .wmv files. Audio was leveled using Adobe Audition to ensure that each word had the same root mean squared (RMS) amplitude. Conditions that included background noise used RMS-leveled six-talker babble that was mixed and included in the final version of the file.

The 350 recordings used in the study were selected from a corpus of 970 recordings of high frequency words (log HAL frequency 7.01–14.99) identified using the English Lexicon Project (Balota et al., 2007). The words that were selected for presentation in the lipreading (visual-only) or audiovisual (AV) conditions in varying signal-to-noise ratios (SNR) were selected from the larger corpus based on visual-only behavioral performance on each word from 149 participants (22–90 years old) who were tested using the entire corpus. The words selected ranged from 10%–93% correct in the lipreading-only behavioral tests. They were distributed among the six conditions that included visual information (AV in Quiet, AV +5 SNR, AV 0 SNR, AV -5 SNR, AV -10, and visual-only) so they would, on average, be equivalent for lipreading difficulty. The words used in the auditory-only condition were selected from the remaining words.

Participants

We collected data from 60 participants ranging in age from 18–34 years (M = 22.42, SD = 3.24, 45 female). All were right-handed native speakers of American English (no other languages other than English before age 7) who self-reported normal hearing and an absence of neurological disease. All provided informed consent under a protocol approved by the Washington University in Saint Louis Institutional Review Board.

Procedure

Before being tested in the fMRI scanner all participants were consented, completed a safety screening, and completed an out-of-scanner lipreading assessment. The behavioral lipreading assessment consisted of 50 single word clips selected in the same way and taken from the same corpus of recorded material used in the scanner. The lipreading assessment was complete by presenting each video clip to the participant using a laptop. Participants were encouraged to verbally provide their best guess for each clip. Only verbatim responses to the stimuli were considered correct.

Participants were positioned in the scanner with insert earphones and a viewing mirror placed above the eyes to see a two-sided projection screen located at the head-side of the scanner. Those that wore glasses were provided scanner-friendly lenses that fit their prescription. Participants were also given a response box that they held in a comfortable position on their torso during testing. Each of the imaging runs presented

trials with recordings of audio, visual-only, audiovisual speech stimuli, or printed text via
an image projected on the screen that was visible to the participant through the viewing
mirror. A camera positioned at the entrance to the scanner bore was used to monitor
participant movement. A well-being check and short conversation occurred before each
run and, if needed, participants were reminded to stay alert and asked to try to reduce
their movement.
Six runs were completed during the session. Each run lasted approximately 5.5
minutes. The first five runs were perception runs and contained 98 trials each. The
stimuli were presented in blocks of five experimental trials plus two null trials for each
condition. The result was 14 blocks resulting in 70 experimental trials plus 28 null trials.
All trials included 800 ms of quiet without a visual presentation before the stimuli began.
During the null trials participants were presented with a fixation cross instead of the
audiovisual presentation. The auditory-only condition did not include visual stimuli;
instead a black screen was presented. The blocks were quasi-randomized so that two
blocks from the same condition were never presented one right after the other and one
null trial never occurred right after another.
To keep attention high, half of the experimental trials required a response from
the participant. On response trials, a set of two dots appeared on the screen after the
audiovisual/audio presentation. The right-side dot was green and the left-side dot was
red. The participant was instructed to use the right-hand button on the response box to
indicate "yes" if they were confident that they had been able to identify the previous
word and to use the left-hand button if they felt they had not identified the previous word
correctly.
After the initial five runs, a final run of 60 trials was presented in which
participants saw a series of written words projected on the screen. The items were the
same 50 words used for the behavioral visual-only assessment, but which did not
appear in any of the other fMRI conditions. Each word stayed on the screen for 2.3
seconds, followed by two green dots that appeared for 2.3 seconds. Participants were
asked to say aloud the word that was presented during the period that the dots were on
the screen. Ten null trials were randomly distributed throughout the sequence. Null trials
lasted 1.5 seconds and included a fixation cross on the screen. The reading task was
always the final run.

Behavioral data analysis
The out-of-scanner lipreading assessment was scored by taking the percentage of
correct responses made by each participant, which we used as a covariate in the fMRI
analyses, allowing us to explore patterns of brain activity that related to more successful
lipreading ability. The in-scanner lipreading was scored similarly, except scores were
based on participants' own judgement of their accuracy. Because we had no way to
verify lipreading accuracy in the scanner, we used these to assess qualitative
differences in difficulty across condition rather than formal statistical analyses.

MRI data acquisition and analysis
MRI images were acquired on a Siemens Prisma 3T scanner using a 32-channel head
coil. Structural images were acquired using a T1-weighted MPRAGE sequence with a
voxel size of .8 x .8 x .8 mm. Functional images were acquired using a multiband

sequence (Feinberg et al., 2010) in axial orientation with an acceleration factor of 8 (TE
= 37 ms), providing full-brain coverage with a voxel size of 2 × 2 × 2 mm. Each volume
took 0.770 s to acquire. We used a sparse imaging paradigm (Edmister et al., 1999;
Hall et al., 1999) with a repetition time of 2.47 s, leaving 1.7 s of silence on each trial.
We presented words during this silent period, and during the repetition task, instructed
participants to speak during a silent period to minimize the influence of head motion on
the data.
Analysis of the MRI data was performed using Automatic Analysis version 5.4.0
(Cusack et al., 2014) (RRID:SCR_003560) that scripted a combination of SPM12
(Wellcome Trust Centre for Neuroimaging) version 7487 (RRID:SCR_007037) and FSL
(FMRIB Analysis Group; Jenkinson et al., 2012) version 6.0.1 (RRID:SCR_002823).
Functional images were realigned, co-registered with the structural image, and spatially
normalized to MNI space (including resampling to 2 mm voxels) using unified
segmentation (Ashburner and Friston, 2005) before smoothing with an 8 mm FWHM
Gaussian kernel. No slice-timing correction was used. First level models contained
regressors for the condition of interest (event onset times convolved with a canonical
hemodynamic response function). To reduce the effects of motion on statistical results
we calculated framewise displacement (FD) using the 6 realignment parameters
assuming the head as a sphere with radius 50 mm (Power et al., 2012). We censored
frames exceeding an FD of 0.5, which resulted in approximately 8% data loss across all
participants. Frames with FD values exceeding this threshold were modeled out by
adding in one additional column to the design matrix for each high-motion scan (cf.
Lemieux et al., 2007).
Psycho-physiological interaction (PPI) analyses are designed to estimate the
effective connectivity between brain regions (Friston et al., 1997); that is, the degree to
which task demands alter the functional connectivity (i.e., statistical dependence of time
series) between a seed region and every other voxel in the brain. PPI analyses thus
require identifying a seed region from which to extract a time course, and two (or more)
tasks between which to compare connectivity with the seed region. For auditory and
visual cortex ROIs (see below for definition), we extracted the time course of the seed
region using SPM's VOI functionality, summarizing the time course as the first
eigenvariate of the ROI after adjusting for effects of interest.
Contrast images from single subject analyses were analyzed at the second level
using permutation testing (FSL *randomise;* 5000 permutations) with a cluster-forming
threshold of p < .001 (uncorrected) and results corrected for multiple comparisons
based on cluster extent (p < .05). Anatomical localization was performed using
converging evidence from author experience (Devlin and Poldrack, 2007) viewing
statistical maps overlaid in MRIcroGL (Rorden and Brett, 2000), supplemented by atlas
labels (Tzourio-Mazoyer et al., 2002).

Regions of interest
We defined regions of interest (ROIs) for the left posterior temporal sulcus (pSTS), left
primary auditory cortex (A1), and left primary visual cortex (V1). For the pSTS, the ROI
was defined as a 10 mm radius sphere centered at MNI coordinates (x=-54, y=-42, z=4)
previously reported to be activated during audiovisual speech processing (Venezia et
al., 2017). The ROIs for AI and V1 were defined using the Anatomy Toolbox (Eickhoff et

al., 2005) (RRID:SCR_013273) as the combination of Areas TE1.0, TE 1.1, and TE 1.2
in the left hemisphere (Morosan et al., 2001) and the left half of area hOC1,
respectively. For the non-PPI ROI analysis, data were extracted by taking the mean of
all voxels in each ROI.
272
273   **Results**

Unthresholded statistical maps are available from NeuroVault (Gorgolewski et al., 2015)
at https://neurovault.org/collections/10922/.
We first examined whole brain univariate effects by condition, shown in **Figure 2**.
We observed temporal lobe activity in all conditions, including visual-only, and visual
cortex activity in all conditions except auditory only.
We next related the activity during visual-only speech with the out-of-scanner
lipreading score (**Figure 1b**). Across participants, lipreading accuracy ranged from 4–
74% (mean = 47.75, SD = 15.49), and correlated with in-scanner ratings (Spearman rho
= 0.38). We included out-of-scanner lipreading as a covariate to see whether individual
differences in out-of-scanner scores related to visual-only activity; we did not find any
significant relationship (positive or negative).
285

Figure 2. Univariate results for spoken word perception in all experimental conditions.

Following univariate analyses, we examined effective connectivity using psychophysiological interaction (PPI) models. We started by using a seed region in left visual cortex. As seen in **Figure 3**, compared to auditory-only speech, visual-only and all audiovisual conditions showed increased connectivity with the visual cortex seed, notably including bilateral superior temporal gyrus and auditory cortex. The same was true with an auditory cortex seed, shown in **Figure 4**. Here, compared to the visual-only condition, we see increased connectivity with visual cortex in all conditions except the auditory-only condition.

Figure 3. Psychophysiological interaction analysis for experimental conditions, using a seed from left visual cortex. Warm-colored voxels showed significantly more connectivity with visual cortex in an experimental condition than in the auditory-only condition.

295
296

Figure 4. Psychophysiological interaction analysis for experimental conditions, using a seed from left auditory cortex. Warm-colored showed significantly more connectivity with auditory cortex in an experimental condition than in the visual-only condition.

297
298   Finally, to complement the above whole-brain analyses, we conducted an ROI
299 analyses focusing on pSTS, shown in **Figure 5**. For the whole-brain univariate and PPI
300 analyses described above, we extracted values from left pSTS and used one-sample t-
301 tests to see whether activity was significantly different from 0. Significance (p < .05,
302 Bonferroni corrected for 19 tests giving p < .00263) is indicated above each condition.

Figure 5. Region-of-interest analyses highlighting the role of the left pSTS in speech processing. **a.** pSTS activity for univariate analyses (cf. Figure 2). **b.** PPI-based effective connectivity with V1 (cf. Figure 3). **c.** PPI-based effective connectivity with A1 (cf. Figure 4). Significant differences from 0, corrected for multiple comparisons, are indicated with an asterisk.

Discussion

We studied brain activity associated with visual-only and audiovisual speech perception.
We found that connectivity between auditory, visual, and premotor cortex was enhanced
during audiovisual speech processing relative to unimodal processing, and during
visual-only speech processing relative to auditory-only speech processing. These

findings are broadly consistent with a role for synchronized interregional neural activity
supporting visual and audiovisual speech perception.

Dedicated regions for multisensory speech processing
Although understanding audiovisual speech requires combining information from
multiple modalities, the way this happens is unclear. One possibility is that heteromodal
brain regions such as the pSTS act to integrate inputs from unisensory cortices. In
addition to combining inputs to form a unitary percept, regions such as pSTS may also
give more weight to more informative modalities (for example, to the visual signal when
the auditory signal is noisy) (Nath and Beauchamp, 2011).
Activity in pSTS for visual-only or AV speech was suggested by both our whole-
brain and ROI-based analyses. In particular, we observed pSTS activity for AV speech
in which the auditory and visual aspects were consistently congruent, consistent with a
role for pSTS in integrating or combining auditory and visual information. Of course,
pSTS activity is not always observed for AV speech (Erickson et al., 2014). One
potential explanation for the variability in pSTS activation across studies is nature of the
speech materials. Several previous studies identifying pSTS involvement in
multisensory speech perception have used incongruent stimuli (i.e., a McGurk task)
(McGurk and MacDonald, 1976), which differs substantially from most of our everyday
speech perception experience (Van Engen et al., 2019). Thus, the conditions under
which pSTS is recruited to support visual or AV speech perception remains an open
question.
In our univariate results, we observed activity in premotor cortex for both visual-
only speech in quiet, and AV speech at more challenging signal-to-noise ratios. These
findings are consistent with a flexible role for premotor cortex in speech perception, at
least under some circumstances, as reported in other studies of visual and audiovisual
speech perception (Venezia et al., 2017). Although our current data do not support
specific conclusions, the dependence of premotor activity on task demands may explain
some of the inconsistencies underlying the debates about the role of premotor cortex
that permeate the speech perception literature.

Effective connectivity and multisensory speech processing
A different perspective comes from a focus on multisensory effects in auditory and
visual cortex (Peelle and Sommers, 2015). Much of the support for this "early
integration" view comes from electrophysiology studies showing multimodal effects in
primary sensory regions (e.g., Schroeder and Foxe, 2005). For example, Lakatos and
colleagues (2007) found that somatosensory input reset the phase of ongoing neural
oscillations in auditory cortex, which was hypothesized to increase sensitivity to auditory
stimuli. In at least one human MEG study, audiovisual effects appear sooner in auditory
cortex than in pSTS (Möttönen et al., 2004), and visual speech may speed processing
in auditory cortex (van Wassenhove et al., 2005). These findings suggest that
multisensory effects are present in primary sensory regions, and that auditory and visual
information do not require a separate brain region in which to "integrate".
In the current data, we observed stronger connectivity between auditory and
visual cortex for visual-only and audiovisual speech conditions than for unimodal
auditory-only speech; and stronger connectivity in audiovisual speech conditions than in

unimodal visual-only speech. That is, using a visual cortex seed we found increases in
effective connectivity with auditory cortex, and when using an auditory cortex seed we
found increases in effective connectivity with visual cortex. These complementary
findings indicate that functionally coordinated activity between primary sensory regions
that is increased during audiovisual speech perception.
Beyond primary sensory cortices, we also observed effective connectivity
changes to premotor cortex for both visual-only speech and several audiovisual
conditions. The functional synchronization between visual cortex, auditory cortex, and
premotor cortex is consistent with a distributed network that orchestrates activity in
response to visual-only and audiovisual speech.
Finally, our ROI analysis showed increased effective connectivity between pSTS
and V1, but not A1, under most experimental conditions (**Figure 5**). These effective
connectivity changes with V1 are consistent with a role for pSTS in audiovisual speech
processing. However, they are also not easily reconcilable with studies reporting
connectivity differences between pSTS and both A1 and V1 (Nath and Beauchamp,
2011). Although no doubt the location and size of any pSTS ROI chosen is important,
we used the same ROI for the PPI analyses with both the A1 seed and V1 seed, and so
ROI definition alone does not seem to explain the qualitative difference between the
two.
It may be worth considering whether the pSTS plays different role in relation to
A1 and V1. Just because pSTS responds to both auditory and visual information does
not necessarily mean it treats them equally, or integrates them in a modality-agnostic
manner. Indeed, given that "unisensory" cortices show multisensory effects and
anatomical connections (Cappe & Barone, 2005), heteromodal or multisensory regions
can also exhibit modality preferences (Noyce et al., 2017). In many audiovisual tasks,
auditory information appears to be preferentially processed (Grondin and McAuley,
2009; Grondin and Rousseau, 1991; Grahn et al. 2011; Recanzone, 2003). Thus, pSTS
may be particularly important in integrating visual information into an existing auditory-
dominated percept. Relatedly, it could also be that multimodal information is inextricably
bound at early stages of perception (Rosenblum, 2008), a process which may rely on
pSTS.
The emerging picture is one in which coordination of large-scale brain
networks—that is, effective connectivity reflecting time-locked functional processing—is
associated with visual-only and audiovisual speech processing. What might be the
function of such distributed, coordinated activity? Visual and audiovisual speech appear
to rely on multisensory representations. For audiovisual speech, it may seem obvious
that successful perception requires combining auditory and visual information. However,
visual-only speech has been consistently associated with activity in auditory cortex
(Calvert et al., 1997; Okada et al., 2013). These activations may correspond to visual-
auditory associations, and auditory-motor associations, learned from audiovisual
speech that are automatically reactivated, even when the auditory input is absent.
Interestingly, our out-of-scanner lipreading scores did not correlate with any of
the whole brain results. It should be noted, however, that our sample size, while large
for fMRI studies of audiovisual speech processing, may still be too small to reliably
detect individual differences in brain activity patterns (Yarkoni and Braver, 2010).
Moreover, there may be multiple ways that brains can support better lipreading, and

such heterogeneity in brain patterns would not be evident in our current analyses.
Future studies with larger sample sizes may be needed to quantitatively assess the
degree to which users' activity might fall into neural strategies, and the degree to which
these are related to lipreading performance.
It is worth highlighting an intriguing aspect of our data, which is that auditory
cortex is always engaged, even in visual-only conditions, whereas the reverse is not
true for visual cortex (which is only engaged when visual information is present) (**Figure
2**). This observation may relate to deeper theoretical issues regarding the fundamental
modality of speech representation. That is, if auditory representations have primacy (at
least, for hearing people), we might expect these representations to be activated
regardless of the input modality (i.e., for both auditory and visual speech). In fact, this is
exactly what we have observed. Although these findings do not directly speak to the
level of detail contained in visual cortex speech representations (Bernstein and
Liebenthal, 2014), they are consistent with asymmetric auditory and visual speech
representations.

Different perspectives on multisensory integration during speech perception
An enduring challenge for understanding multisensory speech perception can be found
in differing uses of the word "integration". During audiovisual speech perception,
listeners use both auditory and visual information, and so from one perspective both
kinds of information are necessarily "integrated" into a listener's (unified) perceptual
experience. However, such use of both auditory and visual information does not
necessitate a separable cognitive stage for integration (Tye-Murray et al., 2016;
Sommers, 2021), nor does it necessitate a region of the brain devoted to integration.
The interregional coordination we observed here may accomplish the task of integration
in that both auditory and visual modalities are shaping perception. In this framework,
there is no need to first translate visual and auditory speech information into some kind
of common code (see also Altieri et al., 2011).
With any study it is important to consider how the specific stimuli used influenced
the results. Here, we examined processing for single words. Visual speech can inform
perception in multiple dimensions (Peelle and Sommers, 2015), including by providing
clues to the speech envelope (Chandrasekaran et al., 2009). These clues may be more
influential in connected speech (e.g., sentences) than in single words, as other neural
processes may come into play with connected speech.

Conclusion
Our findings demonstrate the scaffolding of connectivity between auditory, visual, and
premotor cortices that supports visual-only and audiovisual speech perception. These
findings suggest that the binding of multisensory information need not be restricted to
heteromodal brain regions (e.g., pSTS), but may also emerge from coordinated
unimodal activity throughout the brain.
438

References

Altieri N, Pisoni DB, Townsend JT (2011) Some behavioral and neurobiological
constraints on theories of audiovisual speech integration: a review and
suggestions for new directions. Seeing Perceiving 24:513–539.
Ashburner J and Friston KJ (2005) Unified segmentation. NeuroImage 26: 839–851.
Beauchamp MS, Argall BD, Bodurka J, Duyn JH, Martin A (2004) Unraveling
multisensory integration: patchy organization within human STS multisensory
cortex. Nature Neuroscience 7:1190–1192.
Bernstein LE, Liebenthal E (2014) Neural pathways for visual speech perception. Front
Neurosci 8:386.
Calvert GA, Bullmore ET, Brammer MJ, Campbell R, Williams SCR, McGuire PK,
Woodruff PWR, Iversen SD, David AS (1997) Activation of auditory cortex during
silent lipreading. Science 276:593–596.
Cappe C and Barone P (2005) Heteromodal connections supporting multisensory
integration at low levels of cortical processing in the monkey. The European
journal of neuroscience 22(11): 2886–2902.
Chandrasekaran C, Trubanova A, Stillittano S, Caplier A, Ghazanfar AA (2009) The
natural statistics of audiovisual speech. PLoS Comput Biol 5:e1000436.
Cusack R, Vicente-Grabovetsky A, Mitchell DJ, Wild CJ, Auer T, Linke AC, Peelle JE
(2014) Automatic analysis (aa): efficient neuroimaging workflows and parallel
processing using Matlab and XML. Front Neuroinform 8:90.
Devlin JT, Poldrack RA (2007) In praise of tedious anatomy. NeuroImage 37:1033–
1041.
Edmister WB, Talavage TM, Ledden PJ, Weisskoff RM (1999) Improved auditory cortex
imaging using clustered volume acquisitions. Hum Brain Mapp 7:89–97.
Eickhoff SB, Stephan KE, Mohlberg H, et al. (2005) A new SPM toolbox for combining
probabilistic cytoarchitectonic maps and functional imaging data. NeuroImage
25(4): 1325–1335.
Erickson LC, Heeg E, Rauschecker JP, et al. (2014) An ALE meta-analysis on the
audiovisual integration of speech signals. Human brain mapping 35(11): 5587–
5605.
Feinberg DA, Moeller S, Smith SM, Auerbach E, Ramanna S, Glasser MF, Miller KL,
Ugurbil K, Yacoub E (2010) Multiplexed echo planar imaging for sub-second
whole brain FMRI and fast diffusion imaging. PLoS One 5:e15710.
Fridriksson J, Baker JM, Whiteside J, Eoute D Jr, Moser D, Vesselinov R, Rorden C
(2009) Treating visual speech perception to improve speech production in
nonfluent aphasia. Stroke 40:853–858.
Friston KJ (1994) Functional and effective connectivity in neuroimaging: A synthesis.
Hum Brain Mapp 2:56–78.
Friston KJ, Buechel C, Fink GR, Morris J, Rolls E, Dolan RJ (1997) Psychophysiological
and modulatory interactions in neuroimaging. Neuroimage 6:218–229.
Gorgolewski KJ, Varoquaux G, Rivera G, et al. (2015) NeuroVault.org: a web-based
repository for collecting and sharing unthresholded statistical maps of the human
brain. *Frontiers in Neuroinformatics* 9: 8.

Grahn JA, Henry MJ and McAuley JD (2011) FMRI investigation of cross-modal
interactions in beat perception: Audition primes vision, but not vice versa.
NeuroImage 54: 1231–1243.
Grant KW, Seitz PF (1998) Measures of auditory-visual integration in nonsense
syllables and sentences. J Acoust Soc Am 104:2438–2450.
Hall DA, Haggard MP, Akeroyd MA, Palmer AR, Summerfield AQ, Elliott MR, Gurney
EM, Bowtell RW (1999) "Sparse" temporal sampling in auditory fMRI. Hum Brain
Mapp 7:213–223.
Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM (2012) FSL.
NeuroImage 62:782–790.
Lakatos P, Chen C-M, O'Connell MN, Mills A, Schroeder CE (2007) Neuronal
oscillations and multisensory interaction in primary auditory cortex. Neuron
53:279–292.
Lemieux L, Salek-Haddadi A, Lund TE, et al. (2007) Modelling large motion events in
fMRI studies of patients with epilepsy. Magnetic resonance imaging 25(6): 894–
901.
Magnotti JF, Beauchamp MS (2015) The noisy encoding of disparity model of the
McGurk effect. Psychon Bull Rev 22:701–709.
Mallick DB, Magnotti JF, Beauchamp MS (2015) Variability and stability in the McGurk
effect: contributions of participants, stimuli, time, and response type. Psychon
Bull Rev 22:1299–1307.
Markiewicz CJ, Gorgolewski KJ, Feingold F, Blair R, Halchenko YO, Miller E,
Hardcastle N, Wexler J, Esteban O, Goncalves M, Jwa A, Poldrack RA (2021)
OpenNeuro: An open resource for sharing of neuroimaging data.
bioRxiv:2021.06.28.450168 Available at:
https://www.biorxiv.org/content/10.1101/2021.06.28.450168v1.full.pdf+html
[Accessed July 5, 2021].
Massaro DW, Palmer SE Jr (1998) Perceiving Talking Faces: From Speech Perception
to a Behavioral Principle. MIT Press.
McGurk H, MacDonald J (1976) Hearing lips and seeing voices. Nature 264:746–748.
Morosan P, Rademacher J, Schleicher A, Amunts K, Schormann T, Zilles K (2001)
Human primary auditory cortex: Cytoarchitectonic subdivisions and mapping into
a spatial reference system. Neuroimage 13:684–701.
Möttönen R, Schürmann M, Sams M (2004) Time course of multisensory interactions
during audiovisual speech perception in humans: a magnetoencephalographic
study. Neurosci Lett 363:112–115.
Nath AR, Beauchamp MS (2011) Dynamic changes in superior temporal sulcus
connectivity during perception of noisy audiovisual speech. J Neurosci 31:1704–
1714.
Nath AR, Beauchamp MS (2012) A neural basis for interindividual differences in the
McGurk effect, a multisensory speech illusion. Neuroimage 59:781–787.
Noyce AL, Cestero N, Michalka SW, et al. (2017) Sensory-Biased and Multiple-Demand
Processing in Human Lateral Frontal Cortex. J Neurosci 37(36): 8755–8766.
Nuttall HE, Kennedy-Higgins D, Hogan J, Devlin JT, Adank P (2016) The effect of
speech distortion on the excitability of articulatory motor cortex. NeuroImage
128:218–226.

Okada K, Hickok G (2009) Two cortical mechanisms support the integration of visual
and auditory speech: A hypothesis and preliminary data. Neurosci Lett 452:219–
223.
Okada K, Venezia JH, Matchin W, Saberi K, Hickok G (2013) An fMRI study of
audiovisual speech perception reveals multisensory interactions in auditory
cortex. PLoS One 8:e68959.
Peelle JE, Sommers MS (2015) Prediction and constraint in audiovisual speech
perception. Cortex 68:169–181.
Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE (2012) Spurious but
systematic correlations in functional connectivity MRI networks arise from subject
motion. NeuroImage 59:2142–2154.
Rorden C, Brett M (2000) Stereotaxic display of brain lesions. Behav Neurol 12:191–
2000.
Rosenblum LD (2008) Speech Perception as a Multimodal Phenomenon. Current
directions in psychological science 17(6): 405–409.
Schroeder CE, Foxe J (2005) Multisensory contributions to low-level, "unisensory"
processing. Curr Opin Neurobiol 15:454–458.
Sommers MS (2021) Santa Claus, the tooth fairy, and auditory-visual integration. The
Handbook of Speech Perception. pp. 517–539 Available at:
https://onlinelibrary.wiley.com/doi/10.1002/9781119184096.ch19.
Stephan KE, Friston KJ (2010) Analyzing effective connectivity with functional magnetic
resonance imaging. Wiley Interdiscip Rev Cogn Sci 1:446–459.
Stevenson RA, James TW (2009) Audiovisual integration in human superior temporal
sulcus: Inverse effectiveness and the neural processing of speech and object
recognition. Neuroimage 44:1210–1223.
Sumby WH, Pollack I (1954) Visual contribution to speech intelligibility in noise. J
Acoust Soc Am 26:212–215.
Szenkovits G, Peelle JE, Norris D, Davis MH (2012) Individual differences in premotor
and motor recruitment during speech perception. Neuropsychologia 50:1380–
1392.
Tye-Murray N, Spehar B, Myerson J, Hale S, Sommers M (2016) Lipreading and
audiovisual speech recognition across the adult lifespan: Implications for
audiovisual integration. Psychol Aging 31:380–389.
Tye-Murray N, Spehar BP, Myerson J, Hale S, Sommers MS (2013) Reading your own
lips: Common-coding theory and visual speech perception. Psychon Bull Rev
20:115–119.
Tye-Murray N, Spehar BP, Myerson J, Hale S, Sommers MS (2015) The self-advantage
in visual speech processing enhances audiovisual speech recognition in noise.
Psychon Bull Rev 22:1048–1053.
Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N,
Mazoyer B, Joliot M (2002) Automated anatomical labeling of activations in SPM
using a macroscopic anatomical parcellation of the MNI MRI single-subject brain.
Neuroimage 15:273–289.
Van Engen KJ, Dey A, Sommers M, Peelle JE (2019) Audiovisual speech perception:
Moving beyond McGurk. Available at: psyarxiv.com/6y8qw.

van Wassenhove V, Grant KW, Poeppel D (2005) Visual speech speeds up the neural
processing of auditory speech. Proc Natl Acad Sci U S A 102:1181–1186.
Venezia JH, Fillmore P, Matchin W, Isenberg AL, Hickok G, Fridriksson J (2016)
Perception drives production across sensory modalities: A network for
sensorimotor integration of visual speech. Neuroimage 126:196–207.
Venezia JH, Vaden KI Jr, Rong F, Maddox D, Saberi K, Hickok G (2017) Auditory,
Visual and Audiovisual Speech Processing Streams in Superior Temporal
Sulcus. Front Hum Neurosci 11:174.
Yarkoni T, Braver TS (2010) Cognitive neurosciences approaches to individual
differences in working memory and executive control: Conceptual and
methodological issues. In: Handbook of Individual Differences in Cognition
(Gruszka A, Matthews G, Szymura B, eds), pp 87–107. New York: Springer.
586