DO MOTONEURONS' DISCHARGE RATE SLOW WITH AGING? A SYSTEMATIC REVIEW AND META-ANALYSIS

Lucas B. R. Orssatto [1*], David N. Borg [2], Linda Pendrith [1], Anthony J. Blazevich [3], Anthony J. Shield [1], Gabriel S. Trajano [1]

[1] School of Exercise and Nutrition Sciences, Faculty of Health, Queensland University of Technology (QUT), Brisbane, Australia.

[2] Griffith University, Menzies Health Institute Queensland, The Hopkins Centre, Brisbane, Australia.

[3] Centre for Human Performance, School of Medical and Health Sciences, Edith Cowan University, Joondalup, Australia.

*Corresponding author: Lucas B. R. Orssatto (l.betdarosaorssatto@qut.edu.au).

Abstract

Nervous system maladaptation is linked to the loss of maximal strength production and motor control with aging. Motor unit discharge rates are a critical determinant of force production; thus, lower discharge rates could be a mechanism underpinning maximal strength and motor control losses during aging. This meta-analysis summarized the findings of studies comparing motor unit discharge rates between young and older adults, and examined the effect of distinct muscles and contraction intensities on the magnitude of discharge rates difference between these two groups. Eligible studies were combined in a meta-analysis, including tested contraction intensities and muscles in different levels, to investigate whether there were differences in discharge rates between younger and older adults. Motor unit discharge rates were higher in younger adults compared to older adults, with a pooled standardized mean difference (SMD) for all studies of 0.63 (95%CI= 0.27 to 0.99). Contraction intensity had a significant effect on the pooled SMD, with a 1% increase in intensity associated with a 0.009 (95%CI= 0.003 to 0.015) change in the pooled SMD. These findings suggest that the reductions in motor unit discharge rates, especially at higher contraction intensities, may be an important mechanism underpinning age-related losses in maximal strength production.

Keywords: Ageing, elderly, motor unit, firing frequency, motor neurone.

1. Introduction

Aging is accompanied by a notable reduction in the ability to produce muscular force and power as well as a decline in the maximal rate of force development, which influences the capacity to perform physical activities of daily living (Bergland and Strand, 2019; Orssatto et al., 2020; Suetta et al., 2019; Tomás et al., 2018). These reductions are clearly observed after 50-60 years of age and the force, power and functional capacity losses accelerate during the subsequent decades of life (Larsson et al., 2019; Suetta et al., 2019; Vandervoort, 2002). Changes within the neuromuscular system have been investigated in order to understand the mechanisms underpinning the differences in force production across the lifespan (Hunter et al., 2016; Larsson et al., 2019; Manini et al., 2013; Orssatto et al., 2018). In addition to alterations within the muscles themselves, the role of neural factors needs to be clearly defined, given that muscle output will be limited by the neural input arriving from the motoneurons.

Both the number of recruited motor units and their discharge rates can be readily modulated by the central nervous system in order to regulate muscle forces. Higher forces are achieved by an increase in the number of active motor units and/or their discharge rates (Enoka and Duchateau, 2017). The relative importance of motor unit recruitment versus discharge rate to the force rise differs significantly between muscles (De Luca and Kline, 2012) and important changes have been observed at the motor unit level during aging (Hassan et al., 2021; Manini et al., 2013; Orssatto et al., 2021a). Over at least the last four decades, researchers have investigated the effects of human aging on motor unit discharge rates during sustained isometric contractions (Kamen et al., 1995; Kirk et al., 2016; Mota et al., 2020; Nelson et al., 1983; Vaillancourt et al., 2003). However, inconsistent results have been observed when comparing young and older individuals, with some studies identifying lower (Christie and Kamen, 2010; Connelly et al., 1999; Kamen and Knight, 2004; Kirk et al., 2019, 2018; Mathew Piasecki et al., 2016a) but others similar mean discharge rates for a given proportional level of force during sustained isometric contractions (Christie and Kamen, 2009; Kallio et al., 2010; Kamen and Roy, 2000; Kirk et al., 2016; Power et al., 2012, 2010). This inconsistency may be partly explained by the wide range of methods used across studies, with variations in the (i) muscles studied; (ii) contraction intensities used (2.5 to 100% of maximal voluntary force); and (iii) small sample sizes (ranging from 5 to 49 young and 5 to 36 older adults). Because of these differences, it has been difficult to draw

broad conclusions based on data from any single study. In this case, a systematic review and analysis that pools the available data presented across studies could provide evidence that is more robust in relation to the effect of aging on motor unit discharge rates with special attention to the effects of muscle and contraction intensity.

Given the above, the aim of the present systematic review and meta-analysis with meta-regression was to identify and summarize the findings, and then estimate the effects, of studies comparing motor unit discharge rates between young and older adults during maximal and submaximal isometric sustained contractions. Thereafter, we discuss the potential influence of muscle and contraction intensity on differences in motor unit discharge rates between younger and older individuals.

2. Methods

2.1. Systematic search strategy

A systematic literature search was conducted in three electronic databases (i.e., PubMed, Web of Sciences, and Scopus) in October 2021. The chosen search terms related to aging (e.g., age, elderly, older, and aging), motor units, and discharge rate (e.g., discharge rate, firing frequency, and inter-spike interval) (Supplementary Material 1). The reference lists of the selected studies were screened for additional studies. The search procedures are reported in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram (Liberati et al., 2009; Moher et al., 2009) (Figure 1). The systematic search strategy was designed by LBRO and GST and conducted independently by LBRO and LP.

2.2. Eligibility criteria

Studies were selected according to the following inclusion criteria: i) compared groups based on age (young = 18–40 years and elderly ≥60 years); ii) adopted electromyographic methods allowing the identification of motor unit discharge rate; iii) discharge rate was measured during a sustained isometric contraction; iv) force level chosen for testing was based on a percentage of each participant's maximal voluntary force; and v) peer-reviewed and published in English. Studies were excluded if they included participants

with neurological (e.g., Parkinson disease, Alzheimer's) or musculoskeletal disorders (e.g., osteoarthritis, limb injury), involved explosive or dynamic contractions, or if data analyses or the test task were not clearly reported.

2.3. Data extraction

Data extracted from the selected studies included the country of the corresponding author, participant sample size, participants' characteristics and ages, muscles tested, relative force level reached during testing, and mean discharge rates for both young and older groups. For studies reporting chronic or acute intervention results (e.g., exercise training, fatiguing protocol, or visual gain) the baseline or control group/condition data were extracted. Data were converted to discharge rates for studies reporting inter-spike intervals. When data were reported in graphs, WebPlot Digitizer software (version 4.1) was used to extract the data. LBRO and LP conducted all data extraction independently before comparing results; mean variation between researchers was <1%. The mean values from data extracted by LBRO and LP were used for the analyses.

2.4. Data analyzes

Our primary interest was to investigate whether there were differences in discharge rates between younger and older adults, with a specific focus on differences in various muscle groups. We were also interested in determining the effect of contraction intensity on differences in motor unit discharge rates between younger and older adults.

Estimates from studies were combined in the meta-analysis using a random effects model (restricted maximum likelihood) and presented as forest plots. We used the standardized mean difference (SMD; Hedges' *g*) in motor unit discharge rates between younger and older adults, as it was not always clear whether discharge rates were reported per motor unit or per participant. A two-level hierarchical model was fitted because most studies reported several estimates for the same muscle group (i.e., at different contraction intensities). This modelling approach accounts for the clustering of multiple measurements on the same group of subjects (Assink and Wibbelink, 2016). The model assumes that at the first level, individual results in each study are distributed around a study-level effect mean, and that at the second level, the study means are distributed around an overall effect mean. To address

our second aim, muscle and contraction intensity were included as moderators in the hierarchical model, to determine their effect on the pooled SMD.

The variance of (the distribution of) the *true* effect sizes (τ^2) was used as an indicator of between-study heterogeneity, with higher values denoting greater heterogeneity (Rücker et al., 2008). Publication bias was evaluated using a funnel plot (Egger et al., 1997). A sensitivity analysis was performed to determine the robustness of meta-estimates by comparing the pooled SMD and between-study heterogeneity before and after removing the largest and smallest effects, respectively. All analyses were conducted in R (version 4.0.3) using the RStudio environment (version 1.1.447) and the *meta* (Balduzzi et al., 2019), *metafor* (Viechtbauer, 2010) and *dmetar* (Harrer et al., 2019) packages. The α for all tests was set at 5%. The data and R code that support the findings in this article can be accessed at https://github.com/orssatto/SRMA_MU.

3. Results

3.1. Systematic search

The systematic search conducted in October 2021. It was retrieved 1195 studies and after duplicates removed, 1072 study titles and abstracts were read for eligibility. After removing 1007 studies, 65 studies were read in full and 35 were excluded according to the set criteria, resulting in 30 studies being included in the review. Figure 1 shows the PRISMA flow diagram for all steps of the systematic search.

3.2. Study characteristics

Of the 30 included studies, 16 were conducted in the United States of America, 11 in Canada, 1 in Finland, and 2 in the United Kingdom. Studies tested a total of 353 young and 370 older participants. Different muscles were tested, including upper and lower body, flexors and extensors, and hand muscles. Most studies used intramuscular electromyography, except for four studies that used multichannel surface electromyography (5-pin electrodes, n = 2; and 32- or 64-channel electrodes, n = 2). The contraction intensities ranged from 2.5% to 100% of the participants' maximal voluntary contraction forces.

Figure 1. PRISMA flow diagram for the systematic search.

Table 1. Studies characteristics

Authors	Country	Young participants		Elderly participants		Characteristics	Tested muscles	EMG methods	Contraction intensity (% of MVC)
		Age (years)	Sample (sex)	Age (years)	Sample (sex)				
Christie and Kamen, 2009	USA	23.9 ± 4.1	8 (F=4; M=4)	75.6 ± 6.7	8 (F=3; M=5)	Inactive to moderately active (30 min of exercise three times per week or less	Tibialis anterior	Intramuscular	50%
Christie and Kamen, 2010	USA	21.9 ± 3.1	30 (F=15; M=15)	72.9 ± 4.6	30 (F=15; M=15)	Relatively sedentary (structured physical activity less than 3 days/week).	Tibialis anterior	Intramuscular	100%
Connelly et al., 1999	Canada	20.8 ± 0.8	6 (M)	82.0 ± 1.7	6 (M)	Recreationally active with a mean of 6.7h/week (old) and 7.6 h/week (young)	Tibialis anterior	Intramuscular	10, 25 50, 75, and 100%
Dalton et al., 2009	Canada	23.5 ± 2.9	6 (M)	75.3 ± 4.1	6 (M)	Recreationally active. Young: University students. Old: in a program to maintain muscular endurance, cardiovascular fitness, and flexibility.	Soleus	Intramuscular	25, 50, 75, and 100%
Dalton et al., 2010	Canada	24 ± 2.4	6 (M)	83 ± 9.8	6 (M)	Healthy and recreationally active.	Triceps brachii and biceps brachii	Intramuscular	10, 25 50, 75, and 100%
Erim et al., 1999	USA	30.2 ± 5.7	10 (NR)	76.9 ± 6.6	10 (NR)	No activity level or hand usage was recorded.	First dorsal interosseous	Intramuscular	20 and 50%

Feeney et al., 2018	USA	25 ± 4	13 (F=8; M=5)	78 ± 5	12 (F=5; M=7)	No history of neurological disease or injury to the upper extremity	Extensor digitorum	Surface 32-channels electrode	10 and 20%
Kallio et al., 2010	Finland	27.3 ± 3	10 (M)	70.4 ± 5	13 (M)	Physically active males without any history of neuromuscular or vascular disease.	Soleus	Intramuscular	20 and 40%
Kamen et al., 1995	USA	R(21 – 33)	7 (NR)	R(67 – 85)	7 (NR)	Free of musculoskeletal and neuromuscular complaints. Varied in fitness levels, to give a representative sample of elderly adults.	First dorsal interosseous	Intramuscular	50 and 100%
Kamen and Roy, 2000	USA	27.7 ± 4.3	7 (NR)	74.9 ± 7.1	8 (NR)	Free of any known neuromuscular, musculoskeletal or cardiopulmonary disorders.	First dorsal interosseous	Intramuscular	50 and 100%
Kamen and Knight, 2004	USA	21 (R 18 - 29)	8 (NR)	77 (R 67 – 81)	7 (NR)	None of these persons had resistance training experience within the previous year.	Vastus lateralis	Intramuscular	10 and 50%
Kirk et al., 2016	Canada	27 ± 3	10 (M)	81 ± 4	10 (M)	Free from any known neuromuscular, respiratory, cardiovascular and metabolic illness and none were systematically trained or sedentary.	Gastrocnemius	Intramuscular	25, 50, 75, and 100%
Kirk et al., 2018	Canada	26 ± 4	11 (M)	80 ± 5	10 (M)	Involved in recreational exercise programs three times a week.	Biceps femoris, semitendinosus and semimembranosus	Intramuscular	25, 50, and 100%

Kirk et al., 2019	Canada	R(22 – 33)	10 (M)	R(77 – 88)	10 (M)	Young participants were normally active University students. The older participants were independently living and participated three times per week in an organized fitness and light exercise program.	Superior trapezius	Intramuscular	25, 50, and 100%
Kirk et al., (2021a)	Canada	28 ± 5	5 (M)	85 ± 6	5 (M)	Free from known neurological, metabolic, or orthopedic disease with no prior injury to the hand or upper limb.	First dorsal interosseous	Intramuscular	25, 50, and 100%
Kirk et al., (2021b)	Canada	28 ± 5	8 (M)	83 ± 5	13 (M)	Free of prior injury to the upper limb and any neurological, metabolic, or orthopedic illness.	Anconeus	Intramuscular	100%
Kwon and Christou, 2018	USA	22.6 ± 4.1	11 (F=6; M=5)	74.1 ± 5.7	12 (F=5; M=7)	Healthy without any neurological impairment and were moderately active.	Tibialis anterior	Surface 5-pins electrode	15%
Laidlaw et al., 2000	USA	26.6 ± 1.6	8 (F=4; M=4)	73.4 ± 6.1	14 (F=8; M=6)	No known neuromuscular disorders and their vision corrected to normal levels.	First dorsal interosseous	Intramuscular	3 and 5%
Mota et al., 2020	USA	25 ± 3	12 (M)	75 ± 8	12 (M)	During the 6 months prior to study participation, enrolled participants refrained from lower body resistance training (< three times monthly) or other structured exercise (e.g., jogging) more than 30 min per day, three times per week	Vastus lateralis	Surface 5-pins electrode	50%

Nelson et al., 1983	USA	29.6 ± 4.2	13 (NR)	65.5 ± 2.2	13 (NR)	No current or history of peripheral nerve dysfunction, no current usage of medication known to affect neuronal conduction, and no indication of muscle weakness	Abductor digiti minimi	Intramuscular	5, 10, and 20%
				74.8 ± 2.9	9 (NR)				
				83.5 ± 2.8	10 (NR)				
Patten and Kamen, 2000	USA	19.2 ± 1.6	6 (F=3; M=3)	70.3 ± 3.9	6 (F=2; M=4)	Regular participation in vigorous physical activity since at least the age of 40 years. Young and older adult subjects were matched for current participation in physical activity	Tibialis Anterior	Intramuscular	10, 20, 30, 40, 60, and 100%
Patten et al., 2001	USA	23.2 ± 3.5	6 (F=3; M=3)	75.8 ± 7.4	6 (F=3; M=3)	Mean weekly time spent participating in physical activity was similar between young (128 min) and older (138 min) adult groups.	Abductor digiti minimi	Intramuscular	100%
Pereira et al., 2019	USA	21.9 ± 2.8*	49 (F=25; M=24)	67.7 ± 5.8*	36 (F=19; M=17)	No statistical differences for physical activity level between young and elderly participants.	Biceps Brachii	Surface 64-channels electrode	5%
Piasecki et al., 2016	United Kingdom	25.3 ± 4.8	22 (M)	71.4 (6.2)	20 (M)	Habitually physically active (not sedentary or participating in competitive sport at regional level or above).	Vastus lateralis	Intramuscular	25%
Piasecki et al., 2016	United Kingdom	26 ± 4	18	71 ± 4	14	From the university population and the local community. All participants were healthy and recreationally active.	Tibialis Anterior	Intramuscular	25%

Power et al., 2010	Canada	27 ± 3	10	66 ± 3	10	Recreationally active young university students, and older adults from a local exercise program designed to maintain cardiorespiratory endurance and flexibility.	Tibialis Anterior	Intramuscular	10%
Power et al., 2012	Canada	27 ± 5	9	70 ± 5	9	Recreationally active young university students and older adults. Older adults participating in non-systematic light walking and calisthenic activities.	Biceps Brachii	Intramuscular	10%
Roos et al., 1999	Canada	26.2 ± 4.1	13 (M)	80 ± 5.3	12 (M)	Moderately active	Vastus medialis	Intramuscular	10, 25, 50, 75, and 100%
Rubinstein and Kamen, 2005	USA	19.2 (R 18-30)	34 (W)	73.1 (R >69)	12 (W)	Engaged in moderate levels of daily physical activity (3–5 days/week).	Tibialis Anterior	Intramuscular	100%
Vaillancourt et al., 2003	USA	22 ± 1	10 (F=5; M=5)	67 ± 2 (F=5; M=5)	10	The subjects were naive to the purpose of the experiment and none had a history of a neurological disorder	First dorsal interosseous	Intramuscular	5, 10, 20, 40%
				82 ± 5	10 (F=5; M=5)				

R = range; EMG, electromyography; NR, not reported; M, male; F, female; *pooled mean ± standard deviation from male and female groups.

3.3. Meta-estimates

3.3.1. Overall pooled effect

Twenty-nine studies were used for the meta-analysis, in which Kamen and Knight (2004) was not included in the meta-analysis, because we were unable to retrieve all necessary data. The pooled effects for each muscle subgroup of the 29 studies are shown in Figure 2 for lower-body muscles and 3 for upper-body muscles. Motor unit discharge rates were higher in younger adults compared to older adults, with an overall pooled SMD for all studies of 0.63 (95% CI = 0.27 to 0.99, $p < .001$). There was evidence of between-study heterogeneity ($\tau^2 = 0.721$, 95% CI = 0.388 to 1.393), justifying the use of a random effects model.

3.3.2. Effect of muscle and contraction intensity

The test of moderators was significant when both muscle (df = 12) and contraction intensity (df = 78) were included in the model, $F = 2.482$, $p = .008$. There was evidence of an effect of contraction intensity on the pooled SMD, with a 1% increase in contraction intensity associated with a 0.009 (95% CI = 0.003 to 0.015, $p = .005$) change in the SMD. Discharge rates were higher for younger adults in the muscles: *trapezius*, *biceps brachii,* and *triceps brachii* (Figure 3), and *hamstrings* and *tibialis anterior* (Figure 2). However, these results, along with the absences of differences in other muscles, should be interpreted with caution. Some muscle groups had as few as three estimates available, sometimes came from different intensities from a single study (e.g., *trapezius*). As such, the failure to find any effect is likely the result of very low statistical power. The *first dorsal interosseous* was the only muscle where there may be no difference between younger and older adults, as the estimates came from six different studies. However, the 95% CI is compatible with small negative and small positive effects, and greater precision on this internal is likely required to conclude that discharge rates are not different between younger and older adults in this muscle.

3.3.3. Publication bias and sensitivity analysis

There was evidence of funnel plot asymmetry (Figure 4A). Removing the study by Dalton et al., (2010) from the meta-analysis removed the asymmetry from the funnel plot (Figure 4B) and led to a marked reduction in the pooled SMD ($g = 0.47$, 95% CI = 0.23 to 0.71, $p < .001$) and a large decrease in between-study heterogeneity ($\tau^2 = 0.202$, 95% CI = 0.053 to 0.521).

Figure 2. Estimates of differences in motor unit discharge rates between younger and older adults in lower body muscles. Positive estimates denote higher motor unit discharge rates for younger adults and negative estimates denote higher discharge rates for older adults. Note that studies ID are repeated accordingly to the contraction intensity adopted by them. This has been taken into account in the model, preventing the same of the unit-of-analysis problem. SMD, standardized mean difference; 95%CI, 95% confidence interval lower and upper limits.

Figure 3. Estimates of differences in motor unit discharge rates between younger and older adults in upper body muscles. Positive estimates denote higher motor unit discharge rates for younger adults and negative estimates denote higher discharge rates for older adults. Note that studies ID are repeated accordingly to the contraction intensity adopted by them. This has been taken into account in the model, preventing the same of the unit-of-analysis problem. SMD, standardized mean difference; 95%CI, 95% confidence interval lower and upper limits.

Figure 4. Publication bias funnel plot with all studies (Panel A) and after removing the study by Dalton et al., (2010) (Panel B). Note that Dalton et al., (2010) induced funnel plot asymmetry in Panel A, which has been improved after removing it in Panel B, which led to a marked reduction in the pooled Hedges'g (i.e., SMD).

Removing the smallest effect ($g = -1.37$, *first dorsal interosseous*, 5% contraction intensity (Vaillancourt et al., 2003)) had little influence on both the pooled SMD and between-study heterogeneity (SMD = 0.64, 95% CI = 0.29–0.99; $p < .001$; $\tau^2 = 0.701$, 95% CI = 0.375–1.359). Removing the largest effect ($g = 13.65$, *triceps brachii*, 100% contraction intensity (Dalton et al., 2010)) had little influence on the pooled SMD, but resulted in a marked reduction in heterogeneity (SMD = 0.62, 95% CI = 0.28–0.95; $p < .001$; $\tau^2 = 0.627$, 95% CI = 0.328–1.232).

4. Discussion

4.1. Main findings

The present meta-analytic review interrogated the current literature to determine the effects of aging on motor unit discharge rates. We summarized the findings of studies comparing motor unit discharge rates between young and older adult participants during

sustained maximal and submaximal isometric contractions. We also attempted to determine the influence of muscle and contraction intensity on motor unit discharge rates during aging. The main findings were that motor unit discharge rates were higher in young adults (SMD = 0.63), with the 95% confidence interval compatible with small to large effects. Moreover, the magnitude of these differences is greater as contraction intensities increases. It was possible to observe clear differences between young and older adults motor unit discharge rates in *tibialis anterior* and possibly the absence of difference in the *first dorsal interosseous*. The small number of studies testing other specific muscles did not allow to further understand a potential effect of muscle on the differences between young and older adults discharge rates. These results show why inconsistent results have been observed in the extant literature as to the effects of aging on motor unit discharge rates, and they thus contribute to our understanding of the effects of aging on neural control during sustained isometric submaximal and maximal contractions.

4.2. Effect of aging and contraction intensity

The reduction in discharge rates, particularly at high contraction levels, might reflect both functional and structural changes within the nervous system during aging. Hypothetically, mechanisms contributing to a potential motor unit discharge rate reduction may include i) a reduced ionotropic input onto the motor unit; ii) the effect of neuromodulation on motoneuron excitability; and iii) the motoneuron structural integrity and motor unit remodeling.

At the motoneuron and motor unit levels, it is also observed several age-related structural and functional changes that may explain the greater reduction in motor unit discharge rate in higher-force contractions. Central to this is that motoneurons are affected by axonal demyelination, atrophy and degeneration during aging (Jang and Remmen, 2011; McKinnon et al., 2015; Misgeld, 2011; Selman et al., 2012). This motoneuron deterioration can trigger muscle fiber denervation, resulting in some fibers being unable to contribute to force production (Aare et al., 2016). Whilst a compensatory mechanism allows low-threshold motoneurons to reinnervate some nearby denervated type II muscle fibers (Hepple and Rice, 2015), it does not prevent the reduction in motor unit discharge rate since the lower-threshold reinnervating motoneuron will exhibit slower discharge rates (Deschenes, 2011; Mathew Piasecki et al., 2016b). As a result, older adults have fewer (McNeil et al., 2005; Mittal and

Logmani, 1987; Tomlinson and Irving, 1977) but larger MUs, a greater proportion of which are innervated by lower-threshold motoneurons (Dalton et al., 2008; Lexell and Taylor, 1991). Although debate exists (Piotrkiewicz and Türker, 2017), motor units innervated by higher recruitment-threshold motoneurons should discharge at higher rates during contractions close to maximal force. However, at lower contraction intensities, motor units innervated by lower threshold motoneurons typically discharge at higher rates than those innervated by higher threshold neurons, which typically start to discharge at higher rates as force level increases (Hu et al., 2014; Oya et al., 2009). As older adults typically possess a greater proportion of lower-threshold motor units, this might explain the lower rate relative to younger adults at higher intensities contractions.

The synaptic input onto the motoneurons is another determinant factor to the rate in which a group of motor units discharges. As discussed before, individual motoneurons present a non-linear input-output relation between the received synaptic input and motoneuron discharging due to the presence of persistent inward currents. However, when analyzing a group of motoneurons, it allow the observation of a more linear transformation of common synaptic input into a cumulative train of action potentials to the muscle fibers (Negro and Farina, 2011). Therefore, the discharge output from a group of motoneurons is controlled by receiving ionotropic input in the low-frequency band-width (i.e., effective neural drive) whose gain is adjusted by motoneuronal neuromodulation (Binder et al., 2020; Negro et al., 2009; Orssatto et al., 2021b). The synaptic input onto the motoneurons is decreased with aging (Chase et al., 1985; Maxwell et al., 2018; Rowe et al., 2006), which could be a consequence of reduced intracortical facilitation and increased intracortical inhibition (McGinley et al., 2010; Opie et al., 2020; Todd et al., 2003).

Motor unit discharge rates are also influenced by neuromodulatory inputs (i.e., serotonin and noradrenaline) from the monoaminergic system, which affects motoneuron intrinsic excitability (Binder et al., 2020). Monoamines exert strong facilitation onto motoneurons by triggering persistent inward currents generated by voltage-sensitive sodium and calcium channels. Neuromodulation can change the monoaminergic input onto motoneurons adjusting the amplification of synaptic input according to the motor task demand. When persistent inward current receive a strong monoaminergic drive and is strongly active, the motoneuron intrinsic excitability is increased, allowing it to discharge in higher frequencies (Gorassini et al., 2002; Orssatto et al., 2021b). Therefore, higher

intensities contractions presents a greater motoneuronal persistent inward current strength (Orssatto et al., 2021b). In fact, estimates of motoneuronal persistent inward currents have been shown to be reduced in upper and lower limbs of older adults (Hassan et al., 2021; Orssatto et al., 2021a). It has been hypothesized, based in animal and human evidence, that these reductions are mainly explained by the deteriorating age-related adaptations within the monoaminergic system, which indicate reduced noradrenaline and serotonin secretions, and hence input onto motoneurons in older adults (Ko et al., 1997; Liu et al., 2020, 2019; Míguez et al., 1999; Shibata et al., 2006) (See discussions in Orssatto et al., (2021a) and Hassan et al., (2021) for further details). Thus, lower monoaminergic input onto motoneurons would limit its ability to discharge in higher frequencies, especially in higher intensities contractions (Orssatto et al., 2021).

In summary, the lower peak discharge rates observed in older adults could be a result of a reduced synaptic input (descending drive and afferent feedback) received by motoneurons and its capacity to integrate and amplify synaptic input onto discharge rate output via persistent inward currents. This, in addition to the smaller number and proportion of higher-threshold motoneurons may collectively limit motor unit discharge rates during high-intensity, more than low-intensity, contractions.

4.3. Muscle subgroups analysis

Age-dependent differences in discharge rates are not readily apparent in the muscle subgroups analyses. Most of these results are underpowered due to the small number of studies and samples sizes for each muscle, except for *tibialis anterior* and *first dorsal interosseous*. Keeping this in mind, the following interpretations should be taken with caution.

The differences observed in *tibialis anterior* (n = 8) and preliminary evidence of differences for *hamstrings* (n = 1) but not in *soleus* (n = 2), *gastrocnemius* (n = 1) and *quadriceps* (n = 3) suggest a pattern of reduction in discharge rates in motor units from lower-limb flexors, rather than from lower-limb extensors. Differences between muscles of different anatomical location might speculatively be explained by the characteristics of the descending tracts that innervate these muscle groups (Taylor et al., 2000). For example, previous studies in the ventral spinal cord of the decerebrate cat and neonatal rat as well as estimations of motoneuron excitability in humans have shown a greater excitability of extensor than flexor motoneurons (Cotel et al., 2009; Hounsgaard et al., 1988; Wilson et al.,

2015). Lower body extensors commonly serve an anti-gravity role, meaning that they are active for longer periods and produce greater cumulative force than flexor muscles during activities of daily living, including during upright standing and locomotor propulsion (e.g., walking) (Masani et al., 2013; Soames and Atha, 1981). These functions may be associated with a lesser risk of discharge rate loss with aging. Additional evidence for this assertion is provided by the findings that, while disuse can aggravate the deleterious effects of aging on the nervous system, master-level (i.e., older adult) athletes show a substantial preservation of neural function (Aagaard et al., 2010; Hvid et al., 2018; McGregor et al., 2011; Unhjem et al., 2016). However, further studies testing *hamstrings, soleus, gastrocnemius*, and *quadriceps* should be developed to confirm this hypothesis.

For the upper body muscles, there were no differences between younger and older adults for any muscles responsible for hand movements, such as *first dorsal interosseous* (n = 6), *extensor digitorum* (n = 1), and *abductor digiti minimi* (n = 2). It could be speculated, with caution, that hand muscles are constantly used during the lifespan, which could have prevented discharge rate loss with aging, as discussed before. Conversely, the estimates from *anconeus* (n = 1), *trapezius* (n = 1), *biceps brachii* (n = 3), and *triceps brachii* (n = 1) could suggest lower discharge rates for older adults. Interestingly, the SMD for *triceps brachii* and *biceps brachii* from Dalton et al., (2010) is considerably bigger than the observed difference in other studies and muscles, which noticeably affected the subgroups, overall pooled estimates and unsurprisingly, between-study heterogeneity (See Figure 4). For example, Pereira et al., (2019) and Power et al., (2012) showed a very low and non-significant SMD on *biceps brachii*. Therefore, any specific conclusion regarding these muscles should be avoided.

4.4. Strengths and Limitations

Despite the relevance of the current findings, some limitations should be mentioned. First, the small number of studies investigating different specific muscles other than tibialis anterior and *first dorsal interosseous* reduced the statistical power for them. Also, our search only identified studies conducted in more economically developed countries (i.e., USA, Canada, Finland, and United Kingdom). It is unclear if, for example, whether the potentially longer life expectancy and better physical health of individuals from developed countries might influence the research findings. Also, studies investigating sex-related effects of aging

on motor unit discharge rates are lacking. Only one study (Pereira et al., 2019) compared older and younger men and women, while other studies only tested men, combined male and female data, or did not report the participants' sex (Table 1). As such, it was not possible to explore potential sex-related differences on motor unit discharge rate reductions during aging; future studies are clearly warranted to address this issue.

Another important limitation of the current literature is mostly adopting sustained isometric contractions in which motor unit discharge rates were measured during the contraction plateau were included in our systematic search. This methodological restriction allowed a better control for between-study comparisons because motor unit discharge behavior is contraction mode dependent (i.e., isometric vs. concentric vs. eccentric) and varies with movement velocity (i.e., fast, or slow) and the phase of isometric force production (i.e., torque rise, plateau, or decline). However, this inclusion criterion could mask some potential age-related differences that may occur in other contractions or movement tasks. For example, evidence exists that motor unit discharge rates during explosive contractions may be more sensitive to aging (Klass et al., 2008). Also, lower motor unit discharge rates were observed for older adults during non-sustained isometric contractions (Hassan et al., 2021; Lucas B.R. Orssatto et al., 2021a). Conversely, dynamic contractions have been shown to present no age-related discharge rates differences compared to isometric contractions (Kirk et al., 2021b). However, more studies are required before clear conclusions can be drawn. Therefore, our findings may only be reflective of those during isometric contractions.

5. Conclusions

The present systematic review identified 30 studies reporting contrasting results regarding the differences in motor unit discharge rates, during sustained isometric contractions between young and older adults. In overall, motor unit discharge rates were lower in older adults. Also, the magnitude of differences between young and older adults is clearly augmented as contraction intensity increases. These findings suggest the reductions in motor unit discharge rates as an important mechanism underpinning the age-related losses in maximal strength production and motor function.

The muscle subgroups analysis indicated that discharge rates were lower in older adults for *tibialis anterior, hamstrings, biceps brachii, triceps brachii, and trapezius.*

However, *tibialis anterior* was the only muscle where a clear difference was observed and estimates came from several studies, rather than only a single study. The absence of differences for *soleus, gastrocnemius, quadriceps, extensor digitorum,* and *abductor digiti minimi* should also be interpreted with caution since the estimates were obtained from one or two studies only. The *first dorsal interosseous* was the only muscle where no difference was observed, and estimates came from several studies. However, greater precision on the SMD is needed to definitively conclude there is no difference between younger and older adults in this muscle. Nonetheless, this evidences that the differences in motor unit discharge rates should be continuously explored in different muscles along with the mechanisms underpinning these reductions.

It is important to consider that our findings may only be reflective of motor unit behavior during isometric contractions. There was also considerable methodological heterogeneity among studies regarding the muscles examined, contraction intensities used, discharge rate measurement techniques, and the age-range of participants. These aspects of study design require careful consideration by future investigations.

References

Aagaard, P., Suetta, C., Caserotti, P., Magnusson, S.P., Kjær, M., 2010. Role of the nervous system in sarcopenia and muscle atrophy with aging: Strength training as a countermeasure. Scand. J. Med. Sci. Sport. 20, 49–64.

Aare, S., Spendiff, S., Vuda, M., Elkrief, D., Perez, A., Wu, Q., Mayaki, D., Hussain, S.N.A., Hettwer, S., Hepple, R.T., 2016. Failed reinnervation in aging skeletal muscle. Skelet. Muscle 6, 29. https://doi.org/10.1186/s13395-016-0101-y

Assink, M., Wibbelink, C., 2016. Fitting Three-Level Meta-Analytic Models in R □: A Step-by-Step Tutorial Mark Assink and Carlijn J . M . Wibbelink. Quant. Methods Psychol. 12, 154–174.

Balduzzi, S., Rücker, G., Schwarzer, G., 2019. How to perform a meta-analysis with R: A practical tutorial. Evid. Based. Ment. Health 22, 153–160. https://doi.org/10.1136/ebmental-2019-300117

Bergland, A., Strand, B.H., 2019. Norwegian reference values for the Short Physical

Performance Battery (SPPB): The Tromsø Study. BMC Geriatr. 19, 1–10. https://doi.org/10.1186/s12877-019-1234-8

Binder, M.D., Powers, R.K., Heckman, C.J., 2020. Nonlinear Input-Output Functions of Motoneurons. Physiology (Bethesda). 35, 31–39. https://doi.org/10.1152/physiol.00026.2019

Chase, M.H., Morales, F.R., Boxer, P.A., Fung, S.J., 1985. Aging of motoneurons and synaptic processes in the cat. Exp. Neurol. 90, 471–478. https://doi.org/10.1016/0014-4886(85)90035-4

Christie, A., Kamen, G., 2010. Short-term training adaptations in maximal motor unit firing rates and afterhyperpolarization duration. Muscle and Nerve 41, 651–660. https://doi.org/10.1002/mus.21539

Christie, A., Kamen, G., 2009. Motor unit firing behavior during prolonged 50% MVC dorsiflexion contractions in young and older adults. J. Electromyogr. Kinesiol. 19, 543–552. https://doi.org/10.1016/j.jelekin.2008.03.005

Connelly, D.M., Rice, C.L., Roos, M.R., Vandervoort, A.A., 1999. Motor unit firing rates and contractile properties in tibialis anterior of young and old men. J. Appl. Physiol. 87, 843–852. https://doi.org/10.1152/jappl.1999.87.2.843

Cotel, F., Antri, M., Barthe, J.Y., Orsal, D., 2009. Identified ankle extensor and flexor motoneurons display different firing profiles in the neonatal rat. J. Neurosci. 29, 2748–2753. https://doi.org/10.1523/JNEUROSCI.3462-08.2009

Dalton, B.H., Harwood, B., Davidson, A.W., Rice, C.L., 2009. Triceps surae contractile properties and firing rates in the soleus of young and old men. J. Appl. Physiol. 107, 1781–1788. https://doi.org/10.1152/japplphysiol.00464.2009

Dalton, B.H., Jakobi, J.M., Allman, B.L., Rice, C.L., 2010. Differential age-related changes in motor unit properties between elbow flexors and extensors. Acta Physiol. 200, 45–55. https://doi.org/10.1111/j.1748-1716.2010.02100.x

Dalton, B.H., McNeil, C.J., Doherty, T.J., Rice, C.L., 2008. Age-related reductions in the estimated numbers of motor units are minimal in the human soleus. Muscle and Nerve 38, 1108–1115. https://doi.org/10.1002/mus.20984

De Luca, C.J., Kline, J.C., 2012. Influence of proprioceptive feedback on the firing rate and recruitment of motoneurons. J. Neural Eng. 9. https://doi.org/10.1088/1741-2560/9/1/016007

Deschenes, M.R., 2011. Motor unit and neuromuscular junction remodeling with aging. Curr. Aging Sci. 4, 209–20.

Egger, M., Davey Smith, G., Schneider, M., Minder, C., 1997. Bias in meta-analysis detected by a simple, graphical test. BMJ 315, 629–34. https://doi.org/10.1136/bmj.315.7109.629

Enoka, R.M., Duchateau, J., 2017. Rate coding and the control of muscle force. Cold Spring Harb. Perspect. Med. 7. https://doi.org/10.1101/cshperspect.a029702

Erim, Z., Beg, M.F., Burke, D.T., De Luca, C.J., 1999. Effects of aging on motor-unit control properties. J. Neurophysiol. 82, 2081–2091. https://doi.org/10.1152/jn.1999.82.5.2081

Feeney, D.F., Mani, D., Enoka, R.M., 2018. Variability in common synaptic input to motor neurons modulates both force steadiness and pegboard time in young and older adults. J. Physiol. 596, 3793–3806. https://doi.org/10.1113/JP275658

Gorassini, M., Yang, J.F., Siu, M., Bennett, D.J., 2002. Intrinsic activation of human motoneurons: Reduction of motor unit recruitment thresholds by repeated contractions. J. Neurophysiol. 87, 1859–1866. https://doi.org/10.1152/jn.00025.2001

Harrer, M., Cuijpers, P., Furukawa, T., Ebert, D.D., 2019. dmetar: Companion R Package For The Guide "Doing Meta-Analysis in R". R package version 0.0.9000.

Hassan, A.S., Fajardo, M.E., Cummings, M., McPherson, L.M., Negro, F., Dewald, J.P.A., Heckman, C.J., Pearcey, G.E., 2021. Estimates of persistent inward currents are reduced in upper limb motor units of older adults. J. Physiol. https://doi.org/10.1113/jp282063

Hepple, R.T., Rice, C.L., 2015. Innervation and neuromuscular control in ageing skeletal muscle. J. Physiol. 00, 1–14. https://doi.org/10.1113/jp270561

Hounsgaard, J., Hultborn, H., Jespersen, B., Kiehn, O., 1988. Bistability of alpha☐motoneurones in the decerebrate cat and in the acute spinal cat after intravenous 5☐hydroxytryptophan. J. Physiol. 405, 345–367. https://doi.org/10.1113/jphysiol.1988.sp017336

Hu, X., Rymer, W.Z., Suresh, N.L., 2014. Motor unit firing rate patterns during voluntary

muscle force generation: A simulation study. J. Neural Eng. 11. https://doi.org/10.1088/1741-2560/11/2/026015

Hunter, S.K., Pereira, X.H.M., Keenan, K.G., 2016. The aging neuromuscular system and motor performance. J. Appl. Physiol. 121, 982–995. https://doi.org/10.1152/japplphysiol.00475.2016

Hvid, L.G., Aagaard, P., Ørtenblad, N., Kjaer, M., Suetta, C., 2018. Plasticity in central neural drive with short-term disuse and recovery - effects on muscle strength and influence of aging. Exp. Gerontol. 106, 145–153. https://doi.org/10.1016/j.exger.2018.02.019

Jang, Y.C., Remmen, H. Van, 2011. Age-associated alterations of neuromuscular junction. Exp. Gerontol. 46, 193–198. https://doi.org/10.1016/j.exger.2010.08.029.Age-associated

Kallio, J., Avela, J., Moritani, T., Kanervo, M., Selänne, H., Komi, P., Linnamo, V., 2010. Effects of ageing on motor unit activation patterns and reflex sensitivity in dynamic movements. J. Electromyogr. Kinesiol. 20, 590–598. https://doi.org/10.1016/j.jelekin.2009.12.005

Kamen, G., Knight, C.A., 2004. Training-related adaptations in motor unit discharge rate in young and older adults. J Gerontol A Biol Sci Med Sci 59, 1334–1338. https://doi.org/10.1093/gerona/59.12.1334

Kamen, G., Roy, A., 2000. Motor unit synchronization in young and elderly adults. Eur. J. Appl. Physiol. 81, 403–410. https://doi.org/10.1007/s004210050061

Kamen, G., Sison, S. V., Du, C.C.D., Patten, C., 1995. Motor unit discharge behavior in older adults during maximal-effort contractions. J. Appl. Physiol. 79, 1908–1913. https://doi.org/10.1152/jappl.1995.79.6.1908

Kirk, E.A., Christie, A.D., Knight, C.A., Rice, C.L., 2021a. Motor unit firing rates during constant isometric contraction: establishing and comparing an age-related pattern among muscles. J. Appl. Physiol. 130, 1903–1914. https://doi.org/10.1152/japplphysiol.01047.2020

Kirk, E.A., Copithorne, D.B., Dalton, B.H., Rice, C.L., 2016. Motor unit firing rates of the gastrocnemii during maximal and sub-maximal isometric contractions in young and old men. Neuroscience 330, 376–385. https://doi.org/10.1016/j.neuroscience.2016.05.059

Kirk, E.A., Gilmore, K.J., Rice, C.L., 2021b. Anconeus motor unit firing rates during isometric and muscle-shortening contractions comparing young and very old adults. J. Neurophysiol. 126, 1122–1136. https://doi.org/10.1152/jn.00219.2021

Kirk, E.A., Gilmore, K.J., Rice, C.L., 2018. Neuromuscular changes of the aged human hamstrings. J. Neurophysiol. 120, 480–488. https://doi.org/10.1152/jn.00794.2017

Kirk, E.A., Gilmore, K.J., Stashuk, D.W., Doherty, T.J., Rice, C.L., 2019. Human motor unit characteristics of the superior trapezius muscle with age-related comparisons. J. Neurophysiol. 122, 823–832. https://doi.org/10.1152/jn.00138.2019

Klass, M., Baudry, S., Duchateau, J., 2008. Age-related decline in rate of torque development is accompanied by lower maximal motor unit discharge frequency during fast contractions. J. Appl. Physiol. 104, 739–746. https://doi.org/10.1152/japplphysiol.00550.2007

Ko, M.L., King, M.A., Gordon, T.L., Crisp, T., 1997. The effects of aging on spinal neurochemistry in the rat. Brain Res. Bull. 42, 95–98. https://doi.org/10.1016/S0361-9230(96)00216-X

Kwon, M., Christou, E.A., 2018. Visual information processing in older adults: Reaction time and motor unit pool modulation. J. Neurophysiol. 120, 2630–2639. https://doi.org/10.1152/jn.00161.2018

Laidlaw, D.H., Bilodeau, M., Enoka, R.M., 2000. Steadiness is reduced and motor unit discharge is more variable in old adults. Muscle and Nerve 23, 600–612. https://doi.org/10.1002/(SICI)1097-4598(200004)23:4<600::AID-MUS20>3.0.CO;2-D

Larsson, L., Degens, H., Li, M., Salviati, L., Lee, Y. il, Thompson, W., Kirkland, J.L., Sandri, M., 2019. Sarcopenia: Aging-Related Loss of Muscle Mass and Function. Physiol. Rev. 99, 427–511. https://doi.org/10.1152/physrev.00061.2017

Lexell, J., Taylor, C.C., 1991. Variability in muscle fibre areas in whole human quadriceps muscle: effects of increasing age. J. Anat. 174, 239–49.

Liberati, A., Altman, D.G., Tetzlaff, J., Mulrow, C., Gøtzsche, P.C., Ioannidis, J.P.A., Clarke, M., Devereaux, P.J., Kleijnen, J., Moher, D., 2009. The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions□: Explanation and Elaboration. PLoS Med. 6.

https://doi.org/10.1371/journal.pmed.1000100

Liu, K.Y., Acosta-Cabronero, J., Cardenas-Blanco, A., Loane, C., Berry, A.J., Betts, M.J., Kievit, R.A., Henson, R.N., Düzel, E., Howard, R., Hämmerer, D., 2019. In vivo visualization of age-related differences in the locus coeruleus. Neurobiol. Aging 74, 101–111. https://doi.org/10.1016/j.neurobiolaging.2018.10.014

Liu, K.Y., Kievit, R.A., Tsvetanov, K.A., Betts, M.J., Düzel, E., Rowe, J.B., Tyler, L.K., Brayne, C., Bullmore, E.T., Calder, A.C., Cusack, R., Dalgleish, T., Duncan, J., Henson, R.N., Matthews, F.E., Marslen-Wilson, W.D., Shafto, M.A., Campbell, K., Cheung, T., Davis, S., Geerligs, L., McCarrey, A., Mustafa, A., Price, D., Samu, D., Taylor, J.R., Treder, M., Belle, J. van, Williams, N., Bates, L., Emery, T., Erzinçlioglu, S., Gadie, A., Gerbase, S., Georgieva, S., Hanley, C., Parkin, B., Troy, D., Auer, T., Correia, M., Gao, L., Green, E., Henriques, R., Allen, J., Amery, G., Amunts, L., Barcroft, A., Castle, A., Dias, C., Dowrick, J., Fair, M., Fisher, H., Goulding, A., Grewal, A., Hale, G., Hilton, A., Johnson, F., Johnston, P., Kavanagh-Williamson, T., Kwasniewska, M., McMinn, A., Norman, K., Penrose, J., Roby, F., Rowland, D., Sargeant, J., Squire, M., Stevens, B., Stoddart, A., Stone, C., Thompson, T., Yazlik, O., Barnes, D., Dixon, M., Hillman, J., Mitchell, J., Villis, L., Howard, R., Hämmerer, D., 2020. Noradrenergic-dependent functions are associated with age-related locus coeruleus signal intensity differences. Nat. Commun. 11, 1–9. https://doi.org/10.1038/s41467-020-15410-w

Manini, T.M., Hong, S.L., Clark, B.C., 2013. Aging and muscle: a neuron's perspective. Curr. Opin. Clin. Nutr. Metab. Care 16, 1–10. https://doi.org/10.1097/MCO.0b013e32835b5880.Aging

Masani, K., Sayenko, D.G., Vette, A.H., 2013. What triggers the continuous muscle activity during upright standing? Gait Posture 37, 72–77. https://doi.org/10.1016/j.gaitpost.2012.06.006

Maxwell, N., Castro, R.W., Sutherland, N.M., Vaughan, K.L., Szarowicz, M.D., de Cabo, R., Mattison, J.A., Valdez, G., 2018. α-Motor neurons are spared from aging while their synaptic inputs degenerate in monkeys and mice. Aging Cell 17. https://doi.org/10.1111/acel.12726

McGinley, M., Hoffman, R.L., Russ, D.W., Thomas, J.S., Clark, B.C., 2010. Older adults exhibit more intracortical inhibition and less intracortical facilitation than young adults.

Exp. Gerontol. 45, 671–678. https://doi.org/10.1016/j.exger.2010.04.005

McGregor, K.M., Zlatar, Z., Kleim, E., Sudhyadhom, A., Bauer, A., Phan, S., Seeds, L., Ford, A., Manini, T.M., White, K.D., Kleim, J., Crosson, B., 2011. Physical activity and neural correlates of aging: A combined TMS/fMRI study. Behav. Brain Res. 222, 158–168. https://doi.org/10.1016/j.bbr.2011.03.042

McKinnon, N.B., Montero-Odasso, M., Doherty, T.J., 2015. Motor unit loss is accompanied by decreased peak muscle power in the lower limb of older adults. Exp. Gerontol. 70, 111–118. https://doi.org/10.1016/j.exger.2015.07.007

McNeil, C.J., Doherty, T.J., Stashuk, D.W., Rice, C.L., 2005. Motor unit number estimates in the tibialis anterior muscle of young, old, and very old men. Muscle and Nerve 31, 461–467. https://doi.org/10.1002/mus.20276

Míguez, J.M., Aldegunde, M., Paz-Valiñas, L., Recio, J., Sánchez-Barceló, E., 1999. Selective changes in the contents of noradrenaline, dopamine and serotonin in rat brain areas during aging. J. Neural Transm. 106, 1089–1098. https://doi.org/10.1007/s007020050225

Misgeld, T., 2011. Lost in elimination: mechanisms of axonal loss. e-Neuroforum 2, 21–34. https://doi.org/10.1007/s13295-011-0017-2

Mittal, K.R., Logmani, F.H., 1987. Age-related reduction in 8th cervical ventral nerve root myelinated fiber diameters and numbers in man. Journals Gerontol. 42, 8–10. https://doi.org/10.1093/geronj/42.1.8

Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., Altman, D., Antes, G., Atkins, D., Barbour, V., Barrowman, N., Berlin, J.A., Clark, J., Clarke, M., Cook, D., D'Amico, R., Deeks, J.J., Devereaux, P.J., Dickersin, K., Egger, M., Ernst, E., Gøtzsche, P.C., Grimshaw, J., Guyatt, G., Higgins, J., Ioannidis, J.P.A., Kleijnen, J., Lang, T., Magrini, N., McNamee, D., Moja, L., Mulrow, C., Napoli, M., Oxman, A., Pham, B., Rennie, D., Sampson, M., Schulz, K.F., Shekelle, P.G., Tovey, D., Tugwell, P., 2009. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. PLoS Med. 6. https://doi.org/10.1371/journal.pmed.1000097

Mota, J.A., Kwon, D.P., Kennedy, M., Sobolewski, E.J., Kim, Y., Gonzales, J.U., Stock, M.S., 2020. Compensatory adjustments in motor unit behavior during fatigue differ for

younger versus older men. Aging Clin. Exp. Res. https://doi.org/10.1007/s40520-019-01438-6

Negro, F., Farina, D., 2011. Linear transmission of cortical oscillations to the neural drive to muscles is mediated by common projections to populations of motoneurons in humans. J. Physiol. 589, 629–637. https://doi.org/10.1113/jphysiol.2010.202473

Negro, F., Holobar, A., Farina, D., 2009. Fluctuations in isometric muscle force can be described by one linear projection of low-frequency components of motor unit discharge rates. J. Physiol. 587, 5925–5938. https://doi.org/10.1113/jphysiol.2009.178509

Nelson, R.M., Soderberg, G.L., Urbscheit, N.L., 1983. Comparison of skeletal muscle motor unit discharge characteristics in young and aged humans. Arch. Gerontol. Geriatr. 2, 255–264. https://doi.org/10.1016/0167-4943(83)90029-8

Opie, G.M., Hand, B.J., Semmler, J.G., 2020. Age-related changes in late synaptic inputs to corticospinal neurons and their functional significance: A paired-pulse TMS study. Brain Stimul. 13, 239–246. https://doi.org/10.1016/j.brs.2019.08.003

Orssatto, L.B. da R., Bezerra, E. de S., Schoenfeld, B.J., Diefenthaeler, F., 2020. Lean, fast and strong: Determinants of functional performance in the elderly. Clin. Biomech. Ahead of p, 103182. https://doi.org/10.1016/j.clinbiomech.2020.105073

Orssatto, L.B. da R., Wiest, M.J., Diefenthaeler, F., 2018. Neural and musculotendinous mechanisms underpinning age-related force reductions. Mech. Ageing Dev. 175, 17–23. https://doi.org/10.1016/j.mad.2018.06.005

Orssatto, Lucas B.R., Borg, D.N., Blazevich, A.J., Sakugawa, R.L., Shield, A.J., Trajano, G.S., 2021a. Intrinsic motoneuron excitability is reduced in soleus and tibialis anterior of older adults. GeroScience 43, 2719–2735. https://doi.org/10.1007/s11357-021-00478-z

Orssatto, Lucas B.R., Mackay, K., Shield, A.J., Sakugawa, R.L., Blazevich, A.J., Trajano, G.S., 2021b. Estimates of persistent inward currents increase with the level of voluntary drive in low-threshold motor units of plantar flexor muscles. J. Neurophysiol. 125, 1746–1754. https://doi.org/10.1152/jn.00697.2020

Orssatto, Lucas B R, Mesquita, R.N.O., Phillips, K.M., 2021. Looking at the bigger PICture: Understanding and counteracting the decline of persistent inward currents in older adults. J. Physiol. 1–6. https://doi.org/10.1113/JP282370

Oya, T., Riek, S., Cresswell, A.G., 2009. Recruitment and rate coding organisation for soleus motor units across entire range of voluntary isometric plantar flexions. J. Physiol. 587, 4737–4748. https://doi.org/10.1113/jphysiol.2009.175695

Patten, C., Kamen, G., 2000. Adaptations in motor unit discharge activity with force control training in young and older human adults. Eur. J. Appl. Physiol. 83, 128–143. https://doi.org/10.1007/s004210000271

Patten, C., Kamen, G., Rowland, D.M., 2001. Adaptations in maximal motor unit discharge rate to strength training in young and older adults. Muscle and Nerve 24, 542–550. https://doi.org/10.1002/mus.1038

Pereira, H.M., Schlinder-DeLap, B., Keenan, K.G., Negro, F., Farina, D., Hyngstrom, A.S., Nielson, K.A., Hunter, S.K., 2019. Oscillations in neural drive and age-related reductions in force steadiness with a cognitive challenge. J. Appl. Physiol. 126, 1056–1065. https://doi.org/10.1152/japplphysiol.00821.2018

Piasecki, Mathew, Ireland, A., Coulson, J., Stashuk, D.W., Hamilton-Wright, A., Swiecicka, A., Rutter, M.K., McPhee, J.S., Jones, D.A., 2016a. Motor unit number estimates and neuromuscular transmission in the tibialis anterior of master athletes: evidence that athletic older people are not spared from age-related motor unit remodeling. Physiol. Rep. 4, 1–11. https://doi.org/10.14814/phy2.12987

Piasecki, Mathew, Ireland, A., Jones, D.A., McPhee, J.S., 2016b. Age-dependent motor unit remodelling in human limb muscles. Biogerontology 17, 485–496. https://doi.org/10.1007/s10522-015-9627-3

Piasecki, M., Ireland, A., Stashuk, D., Hamilton-Wright, A., Jones, D.A., McPhee, J.S., 2016. Age-related neuromuscular changes affecting human vastus lateralis. J. Physiol. 594, 4525–4536. https://doi.org/10.1113/JP271087

Piotrkiewicz, M., Türker, K.S., 2017. Onion skin or common drive? Front. Cell. Neurosci. 11, 2015–2018. https://doi.org/10.3389/fncel.2017.00002

Power, G.A., Dalton, B.H., Behm, D.G., Doherty, T.J., Vandervoort, A.A., Rice, C.L., 2012. Motor unit survival in lifelong runners is muscle dependent. Med. Sci. Sports Exerc. 44, 1235–1242. https://doi.org/10.1249/MSS.0b013e318249953c

Power, G.A., Dalton, B.H., Behm, D.G., Vandervoort, A.A., Doherty, T.J., Rice, C.L., 2010.

Motor unit number estimates in masters runners: Use it or lose it? Med. Sci. Sports Exerc. 42, 1644–1650. https://doi.org/10.1249/MSS.0b013e3181d6f9e9

Roos, M.R., Rice, C.L., Connelly, D.M., Vandervoort, A.A., 1999. Quadriceps muscle strength, contractile properties, and motor unit firing rates in young and old men. Muscle and Nerve 22, 1094–1103. https://doi.org/10.1002/(SICI)1097-4598(199908)22:8<1094::AID-MUS14>3.0.CO;2-G

Rowe, J.B., Siebner, H., Filipovic, S.R., Cordivari, C., Gerschlager, W., Rothwell, J., Frackowiak, R., 2006. Aging is associated with contrasting changes in local and distant cortical connectivity in the human motor system. Neuroimage 32, 747–760. https://doi.org/10.1016/j.neuroimage.2006.03.061

Rubinstein, S., Kamen, G., 2005. Decreases in motor unit firing rate during sustained maximal-effort contractions in young and older adults. J. Electromyogr. Kinesiol. 15, 536–543. https://doi.org/10.1016/j.jelekin.2005.04.001

Rücker, G., Schwarzer, G., Carpenter, J.R., Schumacher, M., 2008. Undue reliance on I2 in assessing heterogeneity may mislead. BMC Med. Res. Methodol. 8, 2–9. https://doi.org/10.1186/1471-2288-8-79

Selman, C., Blount, J.D., Nussey, D.H., Speakman, J.R., 2012. Oxidative damage, ageing, and life-history evolution: Where now? Trends Ecol. Evol. 27, 570–577. https://doi.org/10.1016/j.tree.2012.06.006

Shibata, E., Sasaki, M., Tohyama, K., Kanbara, Y., Otsuka, K., Ehara, S., Sakai, A., 2006. Age-related changes in locus ceruleus on neuromelanin magnetic resonance imaging at 3 Tesla. Magn. Reson. Med. Sci. 5, 197–200. https://doi.org/10.2463/mrms.5.197

Soames, R.W., Atha, J., 1981. The role of the antigravity musculature during quiet standing in man. Eur. J. Appl. Physiol. Occup. Physiol. 47, 159–167. https://doi.org/10.1007/BF00421668

Suetta, C., Haddock, B., Alcazar, J., Noerst, T., Hansen, O.M., Ludvig, H., Kamper, R.S., Schnohr, P., Prescott, E., Andersen, L.L., Frandsen, U., Aagaard, P., Bülow, J., Hovind, P., Simonsen, L., 2019. The Copenhagen Sarcopenia Study: lean mass, strength, power, and physical function in a Danish cohort aged 20–93 years. J. Cachexia. Sarcopenia Muscle 1–13. https://doi.org/10.1002/jcsm.12477

Taylor, J.T., Petersen, N., Butler, J.E., Gandevia, S.C., 2000. Ischaemia after exercise does not reduce responses of human motoneurones to cortical or corticospinal tract stimulation. J. Physiol. 525, 793–801. https://doi.org/10.1111/j.1469-7793.2000.00793.x

Todd, G., Taylor, J.L., Gandevia, S.C., 2003. Measurement of voluntary activation of fresh and fatigued human muscles using transcranial magnetic stimulation. J. Physiol. 551, 661–671. https://doi.org/10.1113/jphysiol.2003.044099

Tomás, M.T., Galán-Mercant, A., Carnero, E.A., Fernandes, B., 2018. Functional capacity and levels of physical activity in aging: A 3-year follow-up. Front. Med. 4, 1–8. https://doi.org/10.3389/fmed.2017.00244

Tomlinson, B.E., Irving, D., 1977. The numbers of limb motor neurons in the human lumbosacral cord throughout life. J. Neurol. Sci. 34, 213–219. https://doi.org/10.1016/0022-510X(77)90069-7

Unhjem, R.J., Nygård, M., Hoven, L.T. van den, Sidhu, S.K., Hoff, J., Wang, E., 2016. Lifelong strength training mitigates the age-related decline in efferent drive. J. Appl. Physiol. 121, 415–423. https://doi.org/10.1152/japplphysiol.00117.2016

Vaillancourt, D.E., Larsson, L., Newell, K.M., 2003. Effects of aging on force variability, single motor unit discharge patterns, and the structure of 10, 20, and 40 Hz EMG activity. Neurobiol. Aging 24, 25–35. https://doi.org/10.1016/S0197-4580(02)00014-3

Vandervoort, A. a., 2002. Aging of the human neuromuscular system. Muscle and Nerve 25, 17–25. https://doi.org/10.1002/mus.1215

Viechtbauer, W., 2010. Conducting meta-analyses in R with the metafor. J. Stat. Softw. 36, 1–48. https://doi.org/10.18637/jss.v036.i03

Wilson, J.M., Thompson, C.K., Miller, L.C., Heckman, C.J., 2015. Intrinsic excitability of human motoneurons in biceps brachii versus triceps brachii. J. Neurophysiol. 113, 3692–3699. https://doi.org/10.1152/jn.00960.2014