***Plasmodium falciparum* serology: A comparison of two protein production**

methods for analysis of antibody responses by protein microarray

Tate Oulton[1], Joshua Obiero[2], Isabel Rodriguez[3], Isaac Ssewanyana[4,5], Rebecca A. Dabbs[1],
Bryan Greenhouse[3], Chris Drakeley[1], Phil L. Felgner[2], Will Stone[1], Kevin K.A. Tetteh[1]

[1] Department of Infection Biology, London school of Hygiene and Tropical Medicine, London,
UK

[2] Department of Physiology and Biophysics, University of California, Irvine, Irvine, CA, USA

[3] University of California San Francisco, California, USA

[4] Infectious Diseases Research Collaboration, Kampala, Uganda

[5] Makerere University Collage of Health Sciences, Kampala, Uganda

Corresponding author: Kevin K.A. Tetteh**; kevin.tetteh@lshtm.ac.uk**

Keywords: protein microarray, reverse phase protein array, RPPA, malaria, Plasmodium

Abstract

The evaluation of protein antigens as putative serologic biomarkers of infection has

increasingly shifted to high-throughput, multiplex approaches such as the protein microarray.

In vitro Transcription/Translation (IVTT) systems – a similarly high-throughput protein

expression method – are already widely utilised in the production of protein microarrays,

though purified recombinant proteins derived from more traditional whole cell based

expression systems also play an important role in biomarker characterisation. Here we have

performed a side-by-side comparison of antigen-matched protein targets from an IVTT and

purified recombinant system, on a protein microarray. The magnitude and range of antibody

responses to purified recombinants was found to be greater than that of IVTT proteins, and

responses between targets from opposing expression systems did not clearly correlate.

However, responses between amino acid sequence-matched targets from each expression

system were more closely correlated. Despite the lack of a clearly defined relationship

between antigen-matched targets produced in each expression system, our data indicate

that protein microarrays produced using either method can be used confidently, in a context

dependent manner, though care should be taken when comparing data derived from
contrasting approaches.

Introduction

To date, the majority of malaria serologic studies have focussed on antibody responses to a
small number of well-characterised, highly immunogenic *Plasmodium falciparum* antigens
that have proven to be reliable markers of exposure to infection [1–8]. However, *P.*
falciparum expresses more than 5000 proteins, each a potential antibody target [9,10].
Advances in technology have led to the development of new assay platforms that allow
proteome scale investigation of antibody responses, such as the protein microarray [11,12] -
boasting significantly greater experimental throughput than more classical monoplex
methods (*e.g.* ELISA) [13,14]. The ability to simultaneously interrogate large numbers of
putative targets, using low volumes of sample, significantly increases the rate at which
antibody responses to antigens can be characterised. As such, protein microarray based
approaches to biomarker identification and humoral response profiling in malaria, and other
infectious diseases, have been increasingly adopted [15–24].

One widely utilised form of the protein microarray is based on an in vitro transcription/
translation (IVTT) system [25] – where protein products are produced through a PCR, *in vivo*
recombination cloning and an *in vitro* expression pipeline, before being printed onto arrays
[15]. The benefits of this methodology are considerable. In principle, whole organism
proteome microarrays can be fabricated simply and quickly, enabling analysis of all potential
protein driven immune responses to a pathogen. In terms of limitations, although reaction
times are shorter, the rates of IVTT protein expression lag behind that of *in vivo* expression
systems [26]. Therefore, the inherent heterogeneity between products is not assessed for
every target making it difficult to normalise for reactivity between protein spots, which
represent an impure mix of *E. coli* and target protein.

In contrast to the IVTT array methodology, the printing of purified proteins is cheaper and
typically more quantifiable. Uniform amounts of product can therefore be incorporated into
arrays, increasing confidence when comparing quantitative antibody responses between
antigenic targets [27] and assessing relative immunogenicity. The process can be modified
to support the scale up of recombinant proteins, and furthermore, affinity purification of
protein targets reduces the risk of undesired background reactivity due to expression system
components. However, the time required to produce panels of purified proteins is far in

excess of the IVTT system. For both systems, although the production of complex

conformational proteins is possible it can sometimes be a challenge [28,29].

Here we present a comparison between an IVTT based and purified protein microarray.

Matched malarial protein targets from each methodology were assessed for comparative

reactivity in serum from Ugandan participant samples (n=899) [30] to determine the

suitability of each approach in the context of high-throughput profiling of serological

responses to protein antigens.

Results

Table 1 summarises the purified and IVTT protein targets for each antigen, with further detail

in **Supplementary Table 1**. In brief, we assessed IgG antibody responses to 35 protein

targets, derived from 11 well-characterised *P. falciparum* antigens (distinct gene products).

Each antigen was represented by at least one IVTT and one purified protein target.

Magnitude of responses between expression systems

The magnitude of response to all protein targets was compared by antigen to evaluate

differences in seroreactivity between IVTT derived and purified protein targets. As expected,

responses varied significantly between antigens and between the protein targets mapping to

each antigen.

Mean responses to all targets were compared by expression system (**Figure 1**) revealing a

greater range of response to purified proteins (IQR Log2MFI = 3.88 – 6.40) than IVTT

proteins (IQR Log2MFI 0.46 – 1.68), and a greater magnitude of response to purified than

IVTT targets (p = <0.001). Similarly, the range and median intensity of individual antibody

responses was found to be greater for purified proteins than their IVTT counterparts (*e.g.*

AMA1 - IVTT_1, median [IQR] Log2MFI = 1.66 [0.80 – 2.53]; Pure_1, median [IQR] Log2MFI

= 7.92 [6.16 – 8.52]) for all targets (p = <0.001) except MSP1 Pure_2, which more closely

reflected the level of reactivity to the two MSP1 IVTT targets (**Figure 2**).

Correlation of responses between antigen matched targets

Considering all sequence matched IVTT and purified protein targets (*i.e.* excluding pairwise

comparisons where purified protein sequence were non-overlapping with IVTT sequence for

the same antigen) there was no evidence for a general correlation in mean response

between expression platforms (Spearman's rho (r_s) = 0.279, p = 0.23). Antibody responses

to all protein targets for each antigen were therefore compared individually (representative

example in **Figure 3** and all antigens in **Supplementary Figure 1**). This allowed for

comparison between sequence matching IVTT and purified protein targets (*e.g.* EBA181

IVTT 2 vs. EBA181 Pure 1), non-matching IVTT and purified protein targets (*e.g.* EBA181

IVTT 1 vs. EBA181 Pure 1), and matching or non-matching targets produced in the same

system (*e.g.* EBA181 IVTT 1 vs. EBA181 IVTT 2). Correlations were highly variable (r_s =

1.00 to -0.045) though all but one (GAMA; r_s = -0.045, p = 0.17) demonstrated a degree of

positive, if not always statistically significant, association.

Multiple IVTT targets were produced for EBA181, HSP40, MSP1, MSP4 and MSP5. For all

other than MSP5, non-sequence matching IVTTs were produced; correlation co-efficients for

these targets were between 0.37 and 0.73 (**Supplementary Figure 1**). For EBA181 and

MSP1, IVTT targets overlap by 17 amino acids – equivalent to a small peptide in terminal

regions unlikely to cover immunogenic epitopes. As such, these targets are considered non-

overlapping. For MSP4 and MSP5, duplicate IVTT products (identical but for 1 terminal

amino acid) resulted in near perfect correlation of antibody responses (MSP4 r_s = 1.00, p =

<0.001; MSP5 r_s = 0.94, p = <0.001). Multiple purified protein targets were produced for

ACS5, ETRAMP4, ETRAMP5, HSP40 and MSP1 - none of which overlap. Correlation

between these purified protein targets in each antigen varied between 0.31 and 0.59

(**Supplementary Figure 1**).

For the 8 antigens with >1 IVTT or purified protein target, the greatest level of correlation

was found between an IVTT and purified target in 4/8 instances; between two IVTT targets

(IVTT-IVTT) in 3/8 instances; and between two purified targets (purified-purified) in 1/8

instances (**Supplementary Table 2**). Comparing correlations between antigen-matched

IVTT and purified proteins only, overlapping targets correlate more highly than non-

overlapping targets. Sample sizes were too low to test the significance of this trend within

antigens (**Figure 4**).

Discussion

Protein microarrays are a practical approach to the serological screening of large numbers of

putative malaria antigen biomarkers. The throughput and flexibility of the microarray platform

presents an opportunity to interrogate malarial antibody responses at a scale far exceeding

traditional mono- or multiplex approaches, agnostic of predicted immunological targets. Here

we have evaluated matched antigenic targets produced using two *E. coli* based expression

techniques – In Vitro Transcription/Translation (IVTT), and purified, whole-cell recombinants

– in the context of a protein microarray. We found that the magnitude of antibody responses

to purified protein targets was generally higher than for their IVTT counterparts, and that

correlation between protein target pairs at the individual serum sample level was variable

and related to degree of sequence homogeneity between targets. Our findings warn against

direct comparisons of microarray data from proteins produced in different expression

platforms without careful cross-validation of sequences and allelic types, but provide support

for the use of both IVTT and purified protein microarray platforms in the context of antigen

biomarker discovery and development.

In building this study, we hypothesised that the magnitude of responses to IVTT products -

which tended to be longer, often representing single exon sequences and therefore

potentially containing more epitopes - would be greater than purified targets truncated based

on species-specificity or domain boundaries which potentially represented fewer epitopes.

Contrary to this prediction, we found that purified proteins captured a greater range and

magnitude of responses (Purified, IQR Log2MFI = 3.88 – 6.40; IVTT, IQR Log2MFI 0.46 –

1.68; $p = <0.001$). The greater level of reactivity to purified targets may relate to differences

in the amount of protein deposited on the array, where consistent and defined amounts of

purified protein are spotted in contrast to the unquantified, and likely variable IVTT products.

These findings recommend a degree of caution in interpretation of array data from two

different platforms, for example: MSP5 showed the second highest mean MFI for any

purified protein, but showed among the lowest mean MFI of any IVTT protein.

In addition to differences in the magnitude of mean responses to targets stratified by

expression system, we observed a greater range of individual sample responses, stratified

by antigen, to purified proteins than in sequence matched IVTT-expressed targets (*e.g.*

AMA1 - IVTT_1, median [IQR] Log2MFI = 1.66 [0.80 – 2.53]; *Pichia pastoris* produced

Pure_1, median [IQR] Log2MFI = 7.92 [6.16 – 8.52]; $p = <0.001$). The *P. pastoris* AMA1

included as a control for the evaluation of the production of a conformational protein. AMA1

is a complex structure comprised of three domains defined by three disulphide bonds.

Production of AMA1 in *P. pastoris* has been fully characterised in terms of correct folding of

the purified protein [31,32] and this observation is likely a reflection of antibody reactivity to

correctly folded (*P. pastoris*) and incorrectly folded AMA1 (IVTT).

Considering all antigenic targets together, we found no evidence of correlation in mean

reactivity to sequence matched targets between expression systems ($r_s = 0.28$, $p = 0.23$).

This was not unexpected taking into account the differences in magnitude of response

between IVTT and purified proteins, and that the length of native protein sequence coverage
between IVTT and purified targets was highly variable. In relation, targets grouped by
antigen displayed highly variable correlations of response, but sequence matched proteins
generally displayed stronger correlations of response than non-sequence matched targets,
indicating the importance of capturing specific epitopes within expression sequences when
producing antigens by either expression method.

Despite the lack of a clearly defined relationship between antigen-matched targets from
opposing expression systems, we remain confident that microarrays utilising IVTT or purified
recombinant proteins are able to produce compelling and biologically relevant data. Indeed,
our data show age-dependent trends in antibody responses (typical of highly endemic
populations [1–3]) irrespective of expression system (**Supplementary figure 2**), lending
weight to the applicability of either methodology in serological assays [33–46].

The IVTT system lends itself to microarray applications, as vast numbers of proteins, or even
entire proteomes, may be produced at scale relatively quickly. However, for application to
serology there is concern that expressed proteins are not quantified before printing, and that
expression levels of product may vary considerably; product yield in bacterial-based IVTT
systems is generally considered to be lower (typically ~1 mg mL^{-1} or less) though higher
protein yields have been reported [47,48]. This may lead to artificial discrepancies in
reactivity between targets. Including expression tags on the protein products, such as
glutathione-S-transferase (GST) or poly-histidine (His), allows detection of specific protein to
confirm expression but not direct quantification. Similarly, it is important to acknowledge that
the un-purified nature of printed reaction mixtures may mask, or otherwise adversely affect,
the detection of antibody reactivity in a sample; Davies et al. report IVTT reaction
compositions of 99% *E. coli* lysate to 1% target protein [15], though this will vary
considerably, at scale, in practice.

In contrast to IVTT-based microarrays, printing purified protein allows a highly quantifiable
approach to be taken. Affinity purification and dialysis of expression products substantially
reduces the risk of background reactivity to bacterial components, and the simple
determination of target protein concentrations allows defined quantities of product to be
spotted, providing much greater confidence when comparing reactivity between targets.
However, these advantages come at a substantial cost; the need for *in silico* analysis to
design vectors, transfection procedures, expression and purification drastically slows the
rate at which putative targets can be produced and screened. As such, proteins chosen for
expression are most often down-selected from larger lists of potential markers of interest,

and important targets may be missed that would otherwise be revealed by proteome level

screening. Furthermore, longer proteins and those including transmembrane regions or

signal peptides, often fail to express satisfactorily; the precipitation of product or the

formation of inclusion bodies commonly occurs in these instances. Purified protein antigens

are therefore often much smaller fragments of full-length, native protein, selected based on

the outputs of predictive tools aiming to identify the most immunoreactive and expressible

epitopes. Ultimately, mapping precise antibody binding sequences in this way provides

greater resolution in the dissection of specific epitope reactivity in screened sera, which may

be of immunological relevance. Shorter, epitope specific sequences may in theory be

transposed to IVTT systems with a view to generating more granular serological screens,

though we accept that truncated protein targets will in some cases favour linear B cell

epitopes, while missing conformational epitopes. However, for measuring exposure to

infection there is less importance on the targeting of confirmation epitopes than would be

required for protective epitopes [33].

For a number of samples in all antigens, data points (Log2 MFI) are negative – a result of

background subtraction and normalisation processes. Protein spots showing lower than

background signal most often occur due to technical assay effects or artefacts in imaging,

and hence are not considered to be of biological relevance. An additional yet important

consideration is that data for IVTT and purified protein spots were not identically corrected

and normalised, owing to differences in the composition of respective background control

spots and the GST background reactivity subtracted from tagged purified proteins. This

makes direct comparisons between the two less certain, though the normalisation processes

employed go some way to overcoming this by transforming data to a ratio above expression

system-specific background reactivity. Furthermore, in consideration of these differences,

correlations were calculated for both methods after normalisation to Log2 ratios, and are

based on rank to allow for the identification of non-linear relationships. Nonetheless, these

methods are recognised as imperfect and a degree of uncertainty remains around

comparisons between IVTT and purified proteins.

The primary benefit of the microarray platform is the ability to screen orders of magnitude

more targets simultaneously than more standard serological assays. Our analysis shows

that both IVTT and purified proteins can be successfully used to capture malarial protein-

antigen specific antibody responses on a reverse phase protein microarray. Although

correlations of response between expression systems are not as strong as may have been

expected, a number of acknowledged technical differences in the methods of protein

production may account for this finding. Considering the data more broadly, observed trends

lend support to the utilisation of both IVTT and purified arrays depending on the objectives

and context of hypotheses to be investigated. The strengths and weaknesses of each

expression system should dictate the chosen approach on a case-by-case basis. For

example, very high density proteome level screening to identify 'shortlists' of candidate

markers based on binary categorisation of seropositivity may be best achieved using IVTT

systems. In contrast, smaller numbers of 'shortlisted' targets expressed as purified proteins

may allow for more nuanced characterisation of antibody responses on a more continuous

scale. As already described, the key limitation in the production of purified recombinants in

our current expression pipeline is throughput. The adaption of our methods to increase the

capacity of protein production would improve our ability to more widely mine the biomarker

information derived from the IVTT platform. As such, we are currently exploring a number of

existing approaches to address this methodological bottleneck [49,50].

In summary, the IVTT protein microarray approach has proven to be a powerful, high-

throughput, biomarker discovery platform with applicability across a range of infectious

diseases. When combined with a cheap, scalable and flexible protein expression platform

such as the *E.coli in vivo* expression platform we have the ability to mine potential diagnostic

and vaccine related targets.

Methods

Ethics

All serum samples were collected after written informed consent from the participant or their

parent/guardian. The protocol for sample collection was reviewed and approved by the

Makerere University School of Medicine Research and Ethics Committee (#2011−149 and

#2011−167), the London School of Hygiene and Tropical Medicine Ethics Committee (#5943

and #5944), the Durham University School of Biological and Biomedical Sciences Ethics

Committee, the University of California, San Francisco, Committee on Human Research

(#11−05539 and #11−05995) and the Uganda National Council for Science and Technology

(#HS-978 and #HS-1019).

Samples

Sera were originally collected as part of a comprehensive longitudinal surveillance study

conducted in three sub-counties in Uganda (Walukuba, Jinja District; Kihihi, Kanungu

District, and Nagongera, Tororo). The study design and methods have been previously

reported and are described in detail elsewhere [30]. A sub-selection of samples (n= was

made from individuals with a breadth of recorded clinical episodes of malaria to ensure a

range of sero-reactivity.

Protein targets

Purified protein expression

Recombinant proteins were generated and expressed in *Escherichia coli* as glutathione S-

transferase (GST)-tagged fusion proteins using previously described methods [51–53]. The

exception to this was PfAMA1, which was expressed as a histidine tagged protein in *Pichia*

pastoris [31]. Purification of the expressed proteins was performed using affinity

chromatography (Glutathione Sepharose 4B (GE Healthcare Life Sciences) or HisPur Ni-

NTA (Invitrogen) resins for GST and His tagged proteins, respectively). Protein

concentration was assessed using the Bradford protein assay, with quality, and purity

assessed by 4-20% SDS-PAGE.

IVTT protein expression

An *in vitro* transcription and translation (IVTT) system was used to express proteins of

interest as previously described [15]. Briefly, *Plasmodium falciparum* DNA (3D7 isolate)

coding sequences were PCR-amplified and cloned into T7 expression vectors via

homologous recombination. Target sequences were expressed at 21°C for 16h in *E. coli*-

based, cell-free transcription/translation reactions, and products were printed onto arrays as

un-purified, whole reaction mixtures.

Overview of compared IVTT and purified protein antigens

We assessed antibody responses to protein targets mapping to eleven antigens (*i.e.* distinct

gene products), each represented on the array by at least one IVTT and one purified protein

target. Full details are in **Table 1** and **Supplementary Table 1**. The number of purified

protein targets varied according to availability, while the number of IVTT targets was

dependent on the exon composition of each the gene sequence; multiple exon sequences

were expressed as multiple protein targets based on exon delineation. Similarly, single exon

gene sequences were generally expressed as a single protein. As a result, of the 11

antigens investigated, 8 were represented by >1 IVTT or purified protein target; 5 had >1

IVTT protein target (EBA181, HSP40, MSP1, MSP4 and MSP5) and 5 had >1 purified

protein target (ACS5, ETRAMP4, ETRAMP5, HSP40 and MSP1). Near identical IVTT

proteins (1 aa difference) were produced independently and printed in parallel for two

antigens: MSP4 and MSP5. Purified protein targets were smaller than their IVTT

counterparts, which generally cover the entire reference sequence; purified protein targets

are produced as fragments of the full-length protein, with the aim of capturing antibodies

specific to predicted epitopes based on *in silico* analysis.

Protein microarray

Prior to printing, Tween 20 was added to purified proteins to yield a final concentration of

0.001% Tween 20. Arrays were printed onto nitrocellulose-coated slides (AVID, Grace Bio-

Labs, Inc., Bend, OR, USA) using an Omni Grid Accent microarray printer (Digilabs, Inc.,

Marlborough, MA, USA). Alongside proteins of interest, buffer (PBS) and no-DNA (empty T7

vector reactions) were included as controls to allow for background normalisation of purified

and IVTT proteins respectively.

Sample probing

For analysis of antibody reactivity on the protein microarray, serum samples were diluted

1:200 in a 3 mg mL^{-1} *E. coli* lysate solution in protein arraying buffer (Maine Manufacturing,

Sanford, ME, USA) and incubated at room temperature for 30 min. Arrays were rehydrated

in blocking buffer for 30 min. Blocking buffer was removed, and arrays were probed with pre-

incubated serum samples using sealed, fitted slide chambers to ensure no cross-

contamination of sample between pads. Chips were incubated overnight at 4°C with

agitation. Arrays were washed five times with TBS-0.05% Tween 20, followed by incubation

with biotin-conjugated goat anti-human IgG (Jackson ImmunoResearch, West Grove, PA,

USA) diluted 1:200 in blocking buffer at room temperature. Arrays were washed three times

with TBS-0.05% Tween 20, followed by incubation with streptavidin-conjugated SureLight P-

3 (Columbia Biosciences, Frederick, MD, USA) at room temperature protected from light.

Arrays were washed three times with TBS-0.05% Tween 20, three times with TBS, and once

with water. Arrays were air dried by centrifugation at 500 x *g* for 5 min and scanned on a

GenePix 4300A High-Resolution Microarray Scanner (Molecular Devices, Sunnyvale, CA,

USA). Target and background intensities were measured using an annotated grid file (.GAL).

Data normalisation

Microarray spot foreground and local background fluorescence data were imported into R

(Foundation for Statistical Computing, Vienna, Austria) for correction, normalisation and

analysis. Local background intensities were subtracted from foreground using the

backgroundCorrect function of the limma package [54]. The backgroundCorrect function was

then further applied to GST-tagged purified proteins, whereby background-corrected GST

fluorescence was subtracted from background-corrected target fluorescence to account for

any GST-specific reactivity in samples. All data were then Log2 transformed and the mean

signal intensity of buffer and no-DNA control spots were subtracted from purified and IVTT

proteins respectively to give a relative measure of reactivity to targets over background

(**Supplementary Figure 3**) [20].

Acknowledgements
Many thanks to Tony Holder for the kind provision of purified recombinant proteins (GAMA
and MSP7).

337
338
References

1.   Corran P, Coleman P, Riley E, Drakeley C. Serology: a robust indicator of malaria transmission
intensity? Trends in Parasitology. 2007;23: 575–582.
doi:https://doi.org/10.1016/j.pt.2007.08.023

2.   Drakeley CJ, Corran PH, Coleman PG, Tongren JE, McDonald SLR, Carneiro I, et al. Estimating
medium- and long-term trends in malaria transmission by using serological markers of malaria
exposure. Proceedings of the National Academy of Sciences of the United States of America.
2005;102: 5108–5113. doi:10.1073/pnas.0408725102

3.   Stewart L, Gosling R, Griffin J, Gesase S, Campo J, Hashim R, et al. Rapid assessment of malaria
transmission using age-specific sero-conversion rates. PloS one. 2009;4: e6083–e6083.
doi:10.1371/journal.pone.0006083

4.   Bousema T, Youssef RM, Cook J, Cox J, Alegana VA, Amran J, et al. Serologic markers for
detecting malaria in areas of low endemicity, Somalia, 2008. Emerging infectious diseases.
2010;16: 392–399. doi:10.3201/eid1603.090732

5.   Theisen M, Vuust J, Gottschau A, Jepsen S, Høgh B. Antigenicity and immunogenicity of
recombinant glutamate-rich protein of Plasmodium falciparum expressed in Escherichia coli.
Clin Diagn Lab Immunol. 1995;2: 30–34.

6.   Carvalho LH, Fontes CJ, Fernandes AA, Marinuzzi HC, Krettli AU. Cross-reactive cellular immune
response to circumsporozoite proteins of Plasmodium vivax and P. falciparum in malaria-
exposed individuals. Parasite Immunol. 1997;19: 47–59. doi:10.1046/j.1365-3024.1997.d01-
182.x

7.   Milek RL, DeVries AA, Roeffen WF, Stunnenberg H, Rottier PJ, Konings RN. Plasmodium
falciparum: heterologous synthesis of the transmission-blocking vaccine candidate Pfs48/45 in
recombinant vaccinia virus-infected cells. Exp Parasitol. 1998;90: 165–174.
doi:10.1006/expr.1998.4315

8.   Kanunfre KA, Leoratti FMS, Hoffmann EHE, Durlacher RR, Ferreira AW, Moraes-Avila SL, et al.
Differential recognition of Plasmodium falciparum merozoite surface protein 2 variants by
antibodies from malaria patients in Brazil. Clin Diagn Lab Immunol. 2003;10: 973–976.
doi:10.1128/cdli.10.5.973-976.2003

9.   Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, et al. Genome sequence of the
369   human malaria parasite Plasmodium falciparum. Nature. 2002;419: 498–511.
370   doi:10.1038/nature01097

371   10.   Florens L, Washburn MP, Raine JD, Anthony RM, Grainger M, Haynes JD, et al. A proteomic
372   view of the Plasmodium falciparum life cycle. Nature. 2002;419: 520–526.
373   doi:10.1038/nature01107

374   11.   Burbelo PD, Ching KH, Bush ER, Han BL, Iadarola MJ. Antibody-profiling technologies for
375   studying humoral responses to infectious agents. Expert Rev Vaccines. 2010;9: 567–578.
376   doi:10.1586/erv.10.50

377   12.   Manzano-Román R, Dasilva N, Díez P, Díaz-Martín V, Pérez-Sánchez R, Orfao A, et al. Protein
378   arrays as tool for studies at the host-pathogen interface. J Proteomics. 2013;94: 387–400.
379   doi:10.1016/j.jprot.2013.10.010

380   13.   Adu B, Issahaque Q-A, Sarkodie-Addo T, Kumordjie S, Kyei-Baafour E, Sinclear CK, et al.
381   Microscopic and Submicroscopic Asymptomatic Plasmodium falciparum Infections in Ghanaian
382   Children and Protection against Febrile Malaria. Infect Immun. 2020;88.
383   doi:10.1128/IAI.00125-20

384   14.   Ondigo BN, Hamre KES, Frosch AEP, Ayodo G, White MT, John CC. Antibody Profiles to P.
385   falciparum Antigens Over Time Characterize Acute and Long-Term Malaria Exposure in an Area
386   of Low and Unstable Transmission. Am J Trop Med Hyg. 2020. doi:10.4269/ajtmh.19-0480

387   15.   Davies DH, Liang XW, Hernandez JE, Randall A, Hirst S, Mu YX, et al. Profiling the humoral
388   immune response to infection by using proteome microarrays: High-throughput vaccine and
389   diagnostic antigen discovery. Proceedings of the National Academy of Sciences of the United
390   States of America. 2005;102: 547–552. doi:DOI 10.1073/pnas.0408782102

391   16.   Arevalo-Herrera M, Lopez-Perez M, Dotsey E, Jain A, Rubiano K, Felgner PL, et al. Antibody
392   Profiling in Naive and Semi-immune Individuals Experimentally Challenged with Plasmodium
393   vivax Sporozoites. PLoS Negl Trop Dis. 2016;10: e0004563. doi:10.1371/journal.pntd.0004563

394   17.   Crompton PD, Kayala MA, Traore B, Kayentao K, Ongoiba A, Weiss GE, et al. A prospective
395   analysis of the Ab response to Plasmodium falciparum before and after a malaria season by
396   protein microarray. Proceedings of the National Academy of Sciences of the United States of
397   America. 2010;107: 6958–6963. doi:10.1073/pnas.1001323107

398   18.   Sundaresh S, Randall A, Unal B, Petersen JM, Belisle JT, Hartley MG, et al. From protein
399   microarrays to diagnostic antigen discovery: a study of the pathogen Francisella tularensis.
400   Bioinformatics. 2007;23: I508–I518. doi:10.1093/bioinformatics/btm207

401   19.   Helb DA, Tetteh KKA, Felgner PL, Skinner J, Hubbard A, Arinaitwe E, et al. Novel serologic
402   biomarkers provide accurate estimates of recent Plasmodium falciparum exposure for
403   individuals and communities. Proceedings of the National Academy of Sciences. 2015;112:
404   E4438–E4447. doi:10.1073/pnas.1501705112

405   20.   Stone WJR, Campo JJ, Ouédraogo AL, Meerstein-Kessel L, Morlais I, Da D, et al. Unravelling the
406   immune signature of Plasmodium falciparum transmission-reducing immunity. Nature
407   Communications. 2018;9: 558. doi:10.1038/s41467-017-02646-2

21. van den Hoogen LL, Walk J, Oulton T, Reuling IJ, Reiling L, Beeson JG, et al. Antibody Responses
to Antigenic Targets of Recent Exposure Are Associated With Low-Density Parasitemia in
Controlled Human Plasmodium falciparum Infections. Frontiers in Microbiology. 2019;9.
doi:10.3389/fmicb.2018.03300

22. Sen S, Sanders EC, Gabriel KN, Miller BM, Isoda HM, Salcedo GS, et al. Antibody Response to a
Nucleocapsid Epitope as a Marker for COVID-19 Disease Severity. bioRxiv. 2020.
doi:10.1101/2020.10.15.341743

23. Khan S, Nakajima R, Jain A, de Assis RR, Jasinskas A, Obiero JM, et al. Analysis of Serologic
Cross-Reactivity Between Common Human Coronaviruses and SARS-CoV-2 Using Coronavirus
Antigen Microarray. bioRxiv. 2020. doi:10.1101/2020.03.24.006544

24. Chakraborty S, Randall A, Vickers TJ, Molina D, Harro CD, DeNearing B, et al. Interrogation of a
live-attenuated enterotoxigenic Escherichia coli vaccine highlights features unique to wild-type
infection. NPJ Vaccines. 2019;4: 37. doi:10.1038/s41541-019-0131-7

25. Murray CJ, Baliga R. Cell-free translation of peptides and proteins: from high throughput
screening to clinical production. Curr Opin Chem Biol. 2013;17: 420–426.
doi:10.1016/j.cbpa.2013.02.014

26. Failmezger J. Understanding limitations to increased potential of cell-free protein synthesis.
2018 [cited 2 Dec 2020]. doi:10.18419/OPUS-9701

27. Kamuyu G, Tuju J, Kimathi R, Mwai K, Mburu J, Kibinge N, et al. KILchip v1.0: A Novel
Plasmodium falciparum Merozoite Protein Microarray to Facilitate Malaria Vaccine Candidate
Prioritization. Front Immunol. 2018;9: 2866. doi:10.3389/fimmu.2018.02866

28. Baneyx F, Mujacic M. Recombinant protein folding and misfolding in Escherichia coli. Nat
Biotechnol. 2004;22: 1399–1408. doi:10.1038/nbt1029

29. Ban B, Sharma M, Shetty J. Optimization of Methods for the Production and Refolding of
Biologically Active Disulfide Bond-Rich Antibody Fragments in Microbial Hosts. Antibodies
(Basel). 2020;9. doi:10.3390/antib9030039

30. Kamya MR, Arinaitwe E, Wanzira H, Katureebe A, Barusya C, Kigozi SP, et al. Malaria
transmission, infection, and disease at three sites with varied transmission intensity in Uganda:
implications for malaria control. Am J Trop Med Hyg. 2015;92: 903–912. doi:10.4269/ajtmh.14-
0312

31. Collins CR, Withers-Martinez C, Bentley GA, Batchelor AH, Thomas AW, Blackman MJ. Fine
mapping of an epitope recognized by an invasion-inhibitory monoclonal antibody on the
malaria vaccine candidate apical membrane antigen 1. J Biol Chem. 2007;282: 7431–7441.
doi:10.1074/jbc.M610562200

32. Collins CR, Withers-Martinez C, Hackett F, Blackman MJ. An inhibitory antibody blocks
interactions between components of the malarial invasion machinery. PLoS Pathog. 2009;5:
e1000273. doi:10.1371/journal.ppat.1000273

33. Bailey JA, Berry AA, Travassos MA, Ouattara A, Boudova S, Dotsey EY, et al. Microarray analyses
reveal strain-specific antibody responses to Plasmodium falciparum apical membrane antigen 1
variants following natural infection and vaccination. Sci Rep. 2020;10: 3952.
doi:10.1038/s41598-020-60551-z

34. Tessema SK, Nakajima R, Jasinskas A, Monk SL, Lekieffre L, Lin E, et al. Protective Immunity
against Severe Malaria in Children Is Associated with a Limited Repertoire of Antibodies to
Conserved PfEMP1 Variants. Cell Host Microbe. 2019;26: 579-590.e5.
doi:10.1016/j.chom.2019.10.012

35. Proietti C, Krause L, Trieu A, Dodoo D, Gyan B, Koram KA, et al. Immune Signature Against
Plasmodium falciparum Antigens Predicts Clinical Immunity in Distinct Malaria Endemic
Communities. Mol Cell Proteomics. 2020;19: 101–113. doi:10.1074/mcp.RA118.001256

36. Stucke EM, Niangaly A, Berry AA, Bailey JA, Coulibaly D, Ouattara A, et al. Serologic responses
to the PfEMP1 DBL-CIDR head structure may be a better indicator of malaria exposure than
those to the DBL-α tag. Malar J. 2019;18: 273. doi:10.1186/s12936-019-2905-9

37. Venkatesh A, Jain A, Davies H, Periera L, Maki JN, Gomes E, et al. Hospital-derived antibody
profiles of malaria patients in Southwest India. Malar J. 2019;18: 138. doi:10.1186/s12936-019-
2771-5

38. Friedman-Klabanoff DJ, Travassos MA, Ifeonu OO, Agrawal S, Ouattara A, Pike A, et al. Epitope-
specific antibody responses to a Plasmodium falciparum subunit vaccine target in a malaria-
endemic population. J Infect Dis. 2020. doi:10.1093/infdis/jiaa611

39. Wu L, Mwesigwa J, Affara M, Bah M, Correa S, Hall T, et al. Sero-epidemiological evaluation of
malaria transmission in The Gambia before and after mass drug administration. BMC Med.
2020;18: 331. doi:10.1186/s12916-020-01785-6

40. Labadie-Bracho MY, van Genderen FT, Adhin MR. Malaria serology data from the Guiana
shield: first insight in IgG antibody responses to Plasmodium falciparum, Plasmodium vivax and
Plasmodium malariae antigens in Suriname. Malar J. 2020;19: 360. doi:10.1186/s12936-020-
03434-y

41. Wu L, Mwesigwa J, Affara M, Bah M, Correa S, Hall T, et al. Antibody responses to a suite of
novel serological markers for malaria surveillance demonstrate strong correlation with clinical
and parasitological infection across seasons and transmission settings in The Gambia. BMC
Med. 2020;18: 304. doi:10.1186/s12916-020-01724-5

42. Steinhardt L, Ravaoarisoa E, Wiegand R, Harimanana A, Hedje J, Cotte AH, et al. School-based
sero-surveys to assess the validity of using routine health facility data to target malaria
interventions in the Central Highlands of Madagascar. J Infect Dis. 2020.
doi:10.1093/infdis/jiaa476

43. Lu A, Cote O, Dimitrova SD, Cooley G, Alamgir A, Uzzaman MS, et al. Screening for malaria
antigen and anti-malarial IgG antibody in forcibly-displaced Myanmar nationals: Cox's Bazar
district, Bangladesh, 2018. Malar J. 2020;19: 130. doi:10.1186/s12936-020-03199-4

44. Surendra H, Supargiyono, Ahmad RA, Kusumasari RA, Rahayujati TB, Damayanti SY, et al. Using
health facility-based serological surveillance to predict receptive areas at risk of malaria
outbreaks in elimination areas. BMC Med. 2020;18: 9. doi:10.1186/s12916-019-1482-7

45. Müller-Sienerth N, Shilts J, Kadir KA, Yman V, Homann MV, Asghar M, et al. A panel of
recombinant proteins from human-infective Plasmodium species for serological surveillance.
Malar J. 2020;19: 31. doi:10.1186/s12936-020-3111-5

46.   Assefa A, Ali Ahmed A, Deressa W, Sime H, Mohammed H, Kebede A, et al. Multiplex serology
490   demonstrate cumulative prevalence and spatial distribution of malaria in Ethiopia. Malar J.
491   2019;18: 246. doi:10.1186/s12936-019-2874-z

492   47.   Iskakova MB, Szaflarski W, Dreyfus M, Remme J, Nierhaus KH. Troubleshooting coupled in vitro
493   transcription–translation system derived from Escherichia coli cells: synthesis of high-yield fully
494   active proteins. Nucleic Acids Research. 2006;34: e135–e135. doi:10.1093/nar/gkl462

495   48.   Caschera F, Noireaux V. Synthesis of 2.3 mg/ml of protein with an all Escherichia coli cell-free
496   transcription–translation system. Biochimie. 2014;99: 162–168.
497   doi:10.1016/j.biochi.2013.11.025

498   49.   Konczal J, Gray CH. Streamlining workflow and automation to accelerate laboratory scale
499   protein production. Protein Expr Purif. 2017;133: 160–169. doi:10.1016/j.pep.2017.03.016

500   50.   Nettleship JE, Rada H, Owens RJ. Overview of a High-Throughput Pipeline for Streamlining the
501   Production of Recombinant Proteins. Methods Mol Biol. 2019;2025: 33–49. doi:10.1007/978-1-
502   4939-9624-7_2

503   51.   Herman LS, Fornace K, Phelan J, Grigg MJ, Anstey NM, William T, et al. Identification and
504   validation of a novel panel of Plasmodium knowlesi biomarkers of serological exposure. PLoS
505   Negl Trop Dis. 2018;12: e0006457. doi:10.1371/journal.pntd.0006457

506   52.   Tetteh KKA, Osier FHA, Salanti A, Kamuyu G, Drought L, Failly M, et al. Analysis of antibodies to
507   newly described Plasmodium falciparum merozoite antigens supports MSPDBL2 as a predicted
508   target of naturally acquired immunity. Infect Immun. 2013;81: 3835–3842.
509   doi:10.1128/IAI.00301-13

510   53.   Polley SD, Tetteh KKA, Cavanagh DR, Pearce RJ, Lloyd JM, Bojang KA, et al. Repeat sequences in
511   block 2 of Plasmodium falciparum merozoite surface protein 1 are targets of antibodies
512   associated with protection from malaria. Infect Immun. 2003;71: 1833–1842.
513   doi:10.1128/iai.71.4.1833-1842.2003

514   54.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression
515   analyses for RNA-sequencing and microarray studies. Nucleic Acids Research. 2015;43: e47–
516   e47. doi:10.1093/nar/gkv007

Figure legends

Figure 1. Mean magnitude of antibody responses to targets. The mean magnitude of

response of each protein target stratified by expression system, presented with median and

interquartile range of all mean responses.

Figure 2. Magnitude and range of response to IVTT and purified proteins. All sample

responses (n = 899) to all protein targets grouped by antigen, presented with median and

interquartile range.

Figure 3. Correlation of antibody responses and sequence mapping. A representative

example correlogram of multiple antigen-matched targets (left). Spearman's rank correlation

reported (r_s) and increasing blue colour scale indicates relative strength of correlation based

on calculated correlations for all proteins included in this analysis. Protein schematic (right)

represents amino-acid aligned representation of IVTT (green) and purified (orange) proteins

to the full-length native protein (grey). Proteins in the correlogram and schematic are

correspondingly aligned.

Figure 4. Spearman's rho correlation coefficients between antigen-matched IVTT and

purified proteins. Targets with overlapping amino acid sequences are indicated by closed

circles, compared to non-overlapping sequences indicated by open circles

Tables

Table 1. Description of antigens and their corresponding IVTT and purified protein targets.

Protein	Description	Full length (AA)	Protein target/expression system	Size (Start AA - End AA)
ACS5	Acyl CoA synthase	811	IVTT_1	811 (1–811)
			Pure_1	117 (294–410)
			Pure_2	160 (414–573)
			Pure_3	150 (578–727)
AMA1	Apical membrane antigen 1	622	IVTT_1	622 (1–622)
			Pure_1	450 (97–546)
EBA181	Erythrocyte binding antigen 181	1567	IVTT_1	754 (1–754)
			IVTT_2	752 (737–1488)
			Pure_1	585 (755–1339)
ETRAMP4	Early transcribed membrane antigen 4	136	IVTT_1	136 (1–136)
			Pure_1	25 (28–52)
			Pure_2	61 (76–136)
ETRAMP5	Early transcribed membrane antigen 5	181	IVTT_1	181 (1–181)
			Pure_1	86 (26–111)
			Pure_2	47 (135–181)
GAMA	GPI-anchored membrane antigen	738	IVTT_1	738 (1–738)
			Pure_1	99 (68–166)
HSP40	Heat shock protein 40 type II	402	IVTT_1	134 (80-213)
			IVTT_2	171 (213-401)
			Pure_1	83 (71-153)
			Pure_2	189 (214-402)
MSP1	Merozoite surface protein 1	1720	IVTT_1	870 (1-870)
			IVTT_2	868 (853-1720)
			Pure_1	45 (64-108)
			Pure_2	35 (54-63;109-133)
			Pure_3	116 (1605-1720)
MSP4	Merozite surface protein 4	272	IVTT_1	162 (1-162)
			IVTT_2	161 (1-161)
			IVTT_3	110 (163-272)
			Pure_1	65 (43-107)
MSP5	Merozoite surface protein 5	272	IVTT_1	172 (1-172)
			IVTT_2	171 (1-171)
			Pure_1	61 (147-207)
MSP7	Merozoite surface protein 7	351	IVTT_1	351 (1-351)
			Pure_1	175 (177-351)

Figure 1

Figure 2

Figure 3

Figure 4