1

Individualised Proteogenomics Applied to Analysis of Amino Acid Variants in Malignant Melanoma

Marisa Schmitt[1], Tobias Sinnberg[2], Katrin Bratl[1], Claus Garbe[2], Boris Macek[1], Nicolas C. Nalpas[1]

[1] Quantitative Proteomics, University of Tuebingen, Tuebingen, Germany

[2] Division of Dermatooncology, University of Tuebingen, Tuebingen, Germany

Keywords: Proteogenomics, Cancer, Melanoma, BRAFi Resistance, Mass Spectrometry, Nucleotide Sequencing

Correspondence to:

Prof. Dr. Boris Macek
Chair, Quantitative Proteomics
Director, Proteome Center Tuebingen
Interfaculty Institute for Cell Biology
University of Tuebingen
Auf der Morgenstelle 15
72076 Tuebingen
Germany
Phone: +49/(0)7071/29-70556
E-Mail: boris.macek@uni-tuebingen.de

Dr. Nicolas Claude Nalpas
Quantitative Proteomics
Proteome Center Tuebingen
Interfaculty Institute for Cell Biology
University of Tuebingen
Auf der Morgenstelle 15
72076 Tuebingen
Germany
Phone: +49/(0)7071/29-70552
E-Mail: nicolas.nalpas@ifiz.uni-tuebingen.de

Abstract

Analysis of patient-specific nucleotide variants is a cornerstone of personalised medicine.

Although only 2% of the genomic sequence is protein-coding, mutations occurring in these regions

have the potential to influence protein structure and may have severe impact on disease aetiology.

Of special importance are variants that affect modifiable amino acid residues, as protein

modifications involved in signal transduction networks cannot be analysed by genomics.

Proteogenomics enables analysis of proteomes in context of patient- or tissue-specific non-

synonymous nucleotide variants. Here we developed an individualised proteogenomics workflow

and applied it to study resistance to BRAF inhibitor vemurafenib in malignant melanoma cell line

A375. This approach resulted in high identification and quantification of non-synonymous

nucleotide variants, transcripts and (phospho)proteins. We integrated multi-omic datasets to

reconstruct the perturbed signalling networks associated with BRAFi resistance, prioritise key

actionable nodes and predict drug therapies with potential to disrupt BRAFi resistance mechanism

in A375 cells. Notably, we showed that AURKA inhibition is effective and specific against BRAFi

resistant A375 cells. Furthermore, we investigated nucleotide variants that interfere with protein

modification status and potentially influence signal transduction networks. Mass spectrometry

(MS) measurements confirmed variant-driven modification changes in approximately 50 proteins;

among them was the transcription factor RUNX1 displaying a variant on a known phosphorylation

site S(Ph)276L. We confirmed the loss of phosphorylation site by MS and demonstrated the impact

of this variant on RUNX1 interactome. Our study paves the way for large-scale application of

proteogenomics in melanoma.

Introduction

Accumulation of mutations is one of the hallmarks of cancer cells and malignant melanoma is a type of cancer with the highest frequency of somatic mutations (1). Recent investigations showed that mutations of signalling targets in malignant melanoma are associated with poor clinical outcome, specifically in the mitogen-activated protein kinase (MAPK)/extracellular signal-regulated kinase (ERK) pathway that affects abnormal cellular growth (2). The RAS/BRAF/MEK/ERK pathway is mutated to an oncogenic form in 30% of all cancers, with non-synonymous BRAF mutations in up to 50% of cutaneous melanomas (3). The predominant BRAF mutation is within the kinase domain with a single nucleotide substitution of valine to glutamic acid at amino acid 600 (4). This mutation can result in a 500-fold increased, dimerization-independent activation of BRAF, and thus leads to a constitutive activation of downstream signalling in cancer cells (3,5). Targeted inhibition of the mutated BRAF kinase with selective inhibitors like vemurafenib, dabrafenib or encorafenib (BRAFi) results in a reduction of MAPK pathway signalling (5). However, almost all patients rapidly develop resistance to BRAFi monotherapy after a period of approximately five months (2). The considerable majority of BRAF resistance development is caused by molecular or genetic alterations that lead to MAPK pathway reactivation. The identification of multiple cellular mechanisms of resistance has greatly improved the understanding of malignancy and clinical outcomes of BRAF[V600E] metastatic melanoma e.g. by the introduction of combined BRAF and MEK inhibition. However, variants that alter the corresponding protein modification status and therefore influence resistance, remain largely elusive. In addition, the precise effect of nsSNVs, insertions and deletions (InDels) and frameshift variants at the proteome and PTM level are still largely unknown.

The past decade has seen a revolution in high-throughput sequencing technologies, which provide

information on DNA/RNA sequence, gene structure and expression (6). Mass spectrometry (MS)-

based proteomics is experiencing a technological revolution similar to that of the high-throughput

sequencing. The current state-of-the-art "shotgun" proteomics workflows are capable of routine,

comprehensive analysis of proteomes (7,8) and posttranslational modifications (PTMs) such as

phosphorylation (9,10). However, most of the standard proteomics approaches identify peptides

and proteins by matching MS/MS spectra against protein databases derived from public

repositories (e.g. UniProt) that are not "individualised", i.e. do not contain sequence information

specific for an individual patient, tissue or cell line. Commonly used protein databases therefore

inherently prevent identification of individual non-synonymous variants. Proteogenomics

addresses this issue by combining nucleotide and protein sequencing information, thus enabling

simultaneous study and integration of DNA sequence, RNA expression and splicing, protein

isoform abundance, as well as localisation of protein PTMs in personalised fashion (11,12).

Here we applied an individualised proteogenomics approach to a single immortalised human

melanoma cell line, A375, in its parental as well as in BRAFi resistant state, to analyse non-

synonymous variants and their impact on signal transduction networks in context of acquired

resistance to kinase inhibitors. Integration of matching genomics, transcriptomics and

(phospho)proteomics datasets allowed the reconstruction of signal transduction networks specific

to individual phenotypes. Subsequently, we were able to prioritise a number of drugs based on

their disruptive potential on the signal transduction networks associated with resistance to BRAFi.

We further investigated the impact of non-synonymous amino acid variants on phosphorylation

sites and their putative functional effects on cell signalling network.

Results

To study the impact of amino acids variants on signal transduction networks, we selected a widely exploited melanoma cell line harbouring the BRAF^{V600E} mutation, A375. The cell line was established with two different phenotypes, drug-sensitive ("S" phenotype) and -resistant ("R" phenotype) against the BRAF inhibitor vemurafenib, as described previously (13). Both phenotypes were subjected to whole exome sequencing, RNA-sequencing, as well as proteomics and phosphoproteomics evaluation (**Figure S1A**). These omics measurements were then integrated to reconstruct signalling networks that are disturbed in BRAFi-resistant A375 cells and to investigate the impact of variants altering protein modification status.

BRAFi-sensitive and -resistant A375 cells were measured with high coverage across different "omic" platforms

We investigated the mutational landscape, transcript expression, as well as abundance of proteins and phosphorylation sites in context of BRAFi resistance in A375 cells. Number and type of non-synonymous nucleotide variants detected in the whole exome sequencing (WES) analysis was similar across phenotypes (**Figure 1A** and **Table S1.1**). We observed a high overlap (93.82%) of nucleotide variants between A375 R and S (**Figure S1B**), most of which have already been reported in Cosmic and/or dbSNP databases. Variants were characterised based on the reference to alternate nucleotide variant change, which revealed a higher exchange frequency of adenine to guanine (and vice versa), as well as cytosine to thymine (and vice versa). These variants represented approximately 64% of the total nucleotide changes and were consistent across phenotypes (**Figure S1C**). These non-synonymous variants were incorporated into their corresponding protein sequences, generating several thousand additional protein sequence

isoforms contained within the individualised alternate protein sequence database. Despite this

large increase in the number of protein isoform sequences, the database search space increased by

only 2% (**Figure 1B**). Over 11,000 expressed genes and 42,000 transcripts were identified by

RNA-sequencing using Oxford Nanopore Technology (**Figure 1C**), of which the vast majority

was shared between phenotypes. The individualised alternate protein sequence database was

subsequently used, together with the reference protein sequence database, for the processing of

deep proteomics and phosphoproteomics data from A375 S and A375 R. High resolution MS led

to the quantification of more than 9,000 protein groups (**Figure 1D**), most of which were shared

between phenotypes. While the number of transcripts and protein groups was similar between

phenotypes, over twice as many phosphopeptides were exclusively identified in A375 R compared

to A375 S. This points to the importance of protein phosphorylation in resistance to BRAF

inhibition in A375 cells.

Key signalling pathways are perturbed in BRAFi-sensitive and -resistant melanoma A375 cells

We next performed a quantitative comparison of BRAFi-sensitive versus -resistant phenotypes for

A375 using the allelic frequency of nucleotide variants, transcript expression, as well as protein-

and phosphorylation site abundance (Significance A with s0, see **Supplementary Information**).

Investigation of the allelic frequency in A375 R versus A375 S revealed 271 upregulated and 281

downregulated non-synonymous nucleotide variants (**Figure 2A** and **Table S1.1**). To narrow

down the driver mutations in context of resistance to BRAFi, we overlapped the somatic non-

synonymous nucleotide variants identified in A375 R against the study from Long and colleagues

(14), who analysed somatic mutations in 10 malignant melanoma patients (**Table S2.1**). While we

did not observe any shared variants between A375 R and patients' data, we did find three shared

genes harbouring different somatic variants (**Figure 2B**). Of note, the overlap of somatic mutations

between patients in the Long *et al.* study was also very limited, which highlights the need for

personalised approach to investigate cancer.

RNA-seq analysis identified 1,064 significantly changing transcripts between A375 R and A375

S phenotypes (613 upregulated and 451 downregulated) (**Figure 2C** and **Table S1.2**). We

conducted an overrepresentation analysis of Reactome pathways using transcripts found

significantly upregulated in A375 R or A375 S (**Figure 2D** and **Table S1.3**). This analysis

established the involvement of CREB signalling, RSK activation and ERK/MAPK targets within

A375 S, whereas signalling by PDGF, glycoprotein hormones and collagen degradation are found

in A375 R.

MS analysis identified 302 significantly regulated proteins between phenotypes **(Figure 2E** and

Table S1.4), including key signalling proteins such as BRAF, proteins from the RAS family and

several mitogen-activated protein kinases. A similar overrepresentation analysis based on

significantly changing proteins showed that nucleosome assembly is overrepresented in A375 R,

while pathways connected to plasma lipoproteins and immune system are overrepresented in A375

S (**Figure 2F** and **Table S1.5**). Only 22 phosphorylation sites were found differentially abundant

between phenotypes, including HMGA2 and RUNX1 (**Figure S2A** and **Table S1.6**).

Interestingly, the comparison between transcriptome and proteome revealed a low correlation in

abundance ratios of A375 R versus A375 S (**Figure S2B**). Only seven transcript/protein pairs were

found significantly changing in the same direction, including LIMA1, SMTN and SDCBP. Taken

together, our data show that transcriptomics and proteomics provided distinct functional

information in context of BRAFi-resistance, thus emphasising the need for proteogenomics to

characterise the underlying signalling pathways.

Amino acid variants are detectable within key signalling pathways and proteins

The aforementioned individualised protein database, used during the processing of the MS data,

allowed us to detect a number of variant peptides, i.e. peptides harbouring reference or alternate

amino acid. More amino acid variants were detected in drug-resistant cells compared to drug-

sensitive cells on the proteome and phosphoproteome level (**Figure S2C**). To address whether

mutational hotspots occur within specific pathways, we performed pathway overrepresentation

based on identified protein isoforms harbouring amino acid variants (**Figure S2D**). Within drug-

resistant cells, VEGF and ERbB signalling pathways were exclusively overrepresented for variant

protein isoforms, whereas HIPPO signalling pathway, Rho GTPase cycle and lysosome were

specific for A375 S cells.

Next, we investigated the most commonly mutated genes in melanoma or in context of BRAFi

resistance (**Figure S2E**). Half of these genes were found to harbour at least one nucleotide variant.

However, only the BRAF V600E variant could be identified at both the WES and proteome levels.

Most of the commonly mutated genes were quantified at the transcriptome and proteome levels.

Protein abundance changes based on label-free quantification revealed non-significant trends

between phenotypes. For example, KMTD2 had a higher intensity in drug-sensitive A375 S cells,

whereas ERBB2 showed a higher abundance in A375 R cells. Among the resistance marker genes

found within the Long et al. study, only MAP2K3 (MEK3) was found mutated in our dataset

(R26T). Among these most commonly mutated proteins in melanoma, several were

phosphorylated on known as well as new modification sites (**Table S1.6**). These results highlight

the capability of proteogenomic approach to detect amino acid variants on expressed proteins,

which in turn reveal an overrepresentation of key cell signalling pathways.

The perturbed signalling network in BRAFi resistant A375 cells can be targeted by several drugs

We next integrated the significantly changing non-synonymous nucleotide variants, transcripts,

proteins and phosphorylation sites into a protein-protein interaction network (**Figure 3A, S3A** and

Table S1.7). Only the top 200 entries were retained on the basis of their number of connections

obtained from the BioGRID database. A functional annotation analysis of these 200 entries

revealed an overrepresentation in proteins involved in several immune system pathways, several

general cancer-related pathways, as well as MAPK, Ras and PI3K/Akt signalling pathways

(**Figure S3B** and **Table S1.8**). Interestingly, less than 50% of the entries in this signalling network

harbour one or more amino acid variants (**Figure 3A**). The size of the entries scales up with the

impact factor obtained through our custom bioinformatic workflow and represents their

importance in context of cancer, melanoma and melanoma resistance (see **Supplementary**

Information). A list of drugs was retrieved from DrugBank database and their targets highlighted

within this network. Several entries were highlighted through this approach, such as AURKA,

BRAF and SULT1A1; all of them are known in context of cancer, are mutated in A375 cells, can

be targeted by drugs and are upregulated in A375 R (protein level).

Potential drugs were prioritised based on their target specificity, as well as on the number of

degrees (connections) their targets have with the rest of the signalling network (**Figure 3B**).

Among the drugs that have the potential to disrupt this resistance-specific signalling network were

Alisertib and AT9283 to inhibit AURKA, FG-2216 and PX-478 to target HIF1A, and

Tanespimycin to suppress HSP90. We experimentally validated the action of the compound

Alisertib on BRAFi-resistant and -sensitive A375 cell lines (**Figure 3C** and **S3C**). BRAFi-resistant

A375 cells were sensitive to AURKA inhibition (AURKAi) with Alisertib, regardless of the

absence/presence of BRAFi vemurafenib. Conversely, A375 S tolerated Alisertib (in absence of

BRAFi) and were able to proliferate compared to A375 R cells. Our results highlight the

effectiveness and specificity of AURKAi against BRAFi-resistant A375 cells and demonstrates

that the integration of proteogenomic datasets has potential to predict effective and individualised

drug therapy in context of BRAFi resistance.

Multiple amino acid variants directly affect protein phosphorylation status

We subsequently focused on MS-identified amino acid variants based on their reported

involvement in cancer and melanoma, as well as their effect on protein phosphorylation status

(**Figure S3D** and **Table S1.9**). We identified 44 variant protein isoforms that harboured a

phosphorylation event on (or in instant vicinity of) the variant site in A375 cells (**Figure 3D and**

S3E). For the variant directly affecting the phosphorylation site, these comprised both, reference

variant peptides that lost the phosphorylation site, and alternate variant peptides that gained a

phosphorylation site. Notably, among the phosphorylated alternate peptides, more than half

displayed a tendency for higher abundance in the resistant phenotype. Several of these variants

were among the top impact ranked variants due to their involvement in melanoma and effect on

phosphorylation status (**Figure S3E**). While, these phosphorylated variant peptides were not

significantly changing in abundance between A375 R and S, we speculate that they may still have

a functional effect, which might not necessarily be connected to BRAFi resistance.

Loss of a phosphorylation site on RUNX1 alters its interactome and transcriptional activity

We investigated further one variant with high impact in context of cancer, melanoma and

phosphorylation status that was found on the RUNX1 protein, a key transcription factor involved

in cell proliferation, differentiation and apoptosis (15). This amino acid variant led to the loss of a

known phosphorylation site due to change from serine to leucine at position 276 (**Figure 4A**). The

reference and alternate variant peptides were identified with high resolution MS in both A375 S

and R cells (**Figure S4A** to **C**). The phosphorylation site is located in a highly modified region in

close proximity to the transcription activation domain of RUNX1, previously reported to be

involved in binding of key regulatory proteins, such as P300 (16). We hypothesised that this variant

is likely to influence the interactome of RUNX1. Therefore, we generated a RUNX1 gene

knockout in A375 S cells using the CRISPR/Cas9 system. Single cell clones were selected for

further analysis based on their effective RUNX1 knockout (KO). As a control we used a non-

targeting (NonTar) control guide sequence. A375 RUNX1 KO cells showed an insertion of 215

bp in the Exon 1 of the gene compared to reference DNA of A375 S cells (**Figure S4E**). The lack

of expression of RUNX1 protein was confirmed by western blot and MS analysis (**Figure S4D**).

To study the impact of the loss of a modifiable amino acid, we performed immunoprecipitation of

Flag-tagged RUNX1_wt and RUNX1_S276L in RUNX1 KO SILAC labelled cells in three

independent replicates (**Figure 4B** and **Table S3.1**). The interactome analysis by LC-MS/MS

revealed that RUNX1 and its core binding factor CBFB were significantly enriched in both

pulldowns compared to Flag-empty vector (**Figure S4F and G**). Interestingly, the known

interaction partner histone deacetyltransferase HDAC1 was enriched in RUNX1_wt interactome

and depleted in the RUNX1_S276L interactome (**Figure 4B**).

To confirm these findings, we performed pulldown assays with synthetic peptides harbouring the

amino acid sequence for reference and alternate variant peptides of RUNX1 in A375 cells (**Figure**

4C and **Table S3.2**). As in the interactome study, HDAC1 was significantly depleted in the

pulldown of alternate versus reference variant peptide indicating that the interaction between

HDAC1 and RUNX1 is disturbed due to the variant. We also identified transcriptional repressor

SIN3A to be significantly depleted in alternate variant peptide pulldown compared to reference

pulldown similar to HDAC1. RIN1 and PTPN23 showed the same trend as HDAC1 and SIN3A;

and both proteins are known to act as regulator of RAS-mediated mitogenic activity (17,18). The

proteins enriched in alternate pulldown compared to reference variant peptide pulldown were

overrepresented in TGFβ signalling, melanogenesis and insulin signalling pathways (**Figure S4H**

and **Table S3.3**). Taken together, we demonstrate that the loss of this known phosphorylation site

has an impact on the interactome of RUNX1 (**Figure S4I**) and postulate that it leads to altered

transcriptional activity (**Figure 4D**).

Discussion

In this study, we applied an individualised onco-proteogenomics approach to a widely used cell

line model of melanoma (A375), a cancer that is well known for its high mutation load (1) and the

potential for rewiring cellular networks (12,19). Two consortia, namely the Clinical Proteomic

Tumor Analysis Consortium (CPTAC) and The Cancer Genome Atlas (TCGA), have greatly

contributed to the development of onco-proteogenomics (11,20,21). However, proteogenomics

studies are still relatively rare and, due to their complexity, out of reach of most proteomics (or

genomics) laboratories. Here, we are going beyond state-of-the-art proteogenomics workflow,

through the analysis of a single melanoma cell line in order to generate truly individualised multi-

omics abundance measurement and to predict tumour-specific drug therapies.

Proteogenomic reconstructs the signalling network linked to BRAFi resistance in A375 cells

In this study, the nucleotide variant incorporation revealed very similar numbers (~10,000

variants) across A375 phenotypes, the large majority being SNVs, which is consistent with a

previous study (22). We also observed characteristic nucleotide substitutions, whereby two thirds

of substitutions are comprised of transitions and the rest being transversions. The C to T transition

was highly represented and is known to result from sun-light exposure, which is highly relevant

for skin cancer (23). The proteome coverage we obtained for A375 cells (~9,000 protein groups)

is similar to other state-of-the-art MS-based study of cancer cell lines (24). The differentially

changing entries identified through our multi-omics approach were overrepresented in pathways

of critical importance for melanoma resistance to BRAFi, including collagen degradation, CREB

signalling, RSK activation and MAPK, Ras and PI3K/Akt signalling pathways (25). Another

aspect of pivotal importance was the distinct functional information provided by the transcriptome

and proteome datasets. This agrees with previous study (24) and greatly emphasises the importance

of proteogenomic approaches as opposed to single-omic investigations.

We then reconstructed the signalling network associated with BRAFi-resistance in A375; i.e. using

the significantly changing non-synonymous nucleotide variants, transcripts, proteins and

phosphorylation sites. This approach highlighted several hubs and high impact entries, such as

CUL3, XPO1, AURKA, HSPA5, H2AFX and HIF1A, of which several lead to reduced melanoma

patient survival or are known for their involvement in melanoma development (26-28). Several

drugs were identified and ranked based on their potential to disrupt this signalling network; notably

Alisertib, a highly specific inhibitor of AURKA, which has been previously reported for its

beneficial effect in combination with BRAF and MEK inhibitors in melanoma treatment (27). We

experimentally validated the use of Alisertib on A375 S and A375 R cells and could show that

A375 R cells are sensitive to AURKAi regardless of the absence/presence of BRAFi. Our data

confirm that AURKA has a critical role in the context of resistance and may be suitable for the

treatment of melanoma as reported previously (29). In addition, the prediction and subsequent

validation of AURKAi as specific drug therapy to treat BRAFi resistant A375 cells verify the use

of our individualised proteogenomics approach.

306 ***Proteogenomic pinpoints several peptides that are phosphorylated on the variant site***

307 Because of the higher number of phosphorylation sites within A375 R, we investigated further the

308 amino acid variants that affect phosphorylation status and are important in melanoma

309 development. Around 14.8% of all amino acids in the human proteome are serine, threonine or

310 tyrosine (30), which are predominantly modified by phosphorylation. Several studies have

311 reported that these three amino acids are disproportionally affected by missense mutations (31,32).

312 While these may not all be relevant in tumour cells, since not all genes are expressed at any one-

313 time, previous studies have shown the deleterious effect of such variants (11,12). Among the

314 variant protein isoforms with impact score medium to high, two thirds were known oncogenes or

315 tumour suppressors genes; and more than 80% had a loss of known phosphorylation sites.

316 We identified hundreds of amino acid variants by high resolution mass spectrometry, including

317 some that led to a change in the modification status of the protein. Our identification results are in

318 the same range (or higher) as other studies investigating amino acid variants using custom protein

319 sequence databases (33,34). Overrepresentation of variant protein isoforms revealed their

320 accumulation in signalling by VEGF and ERbB signalling pathways in drug-resistant cells,

321 whereas HIPPO signalling pathway and Rho GTPase cycle were overrepresented in sensitive cells.

322 The overrepresented pathways common to both phenotypes, such as MAPK signalling pathway,

323 are known to be highly activated in melanoma cells with acquired resistance (35). Interestingly,

324 most of the MS-identified amino acid variants affecting the phosphorylation status were displaying

325 a trend towards higher abundance in A375 R. Such discrepancy may be connected to BRAFi

326 resistance in A375 cells as reported in other studies (36,37). Among interesting variant, RUNX1

327 was one of the top impact scoring entries in A375 cells due to its known role in cancer and loss of

328 phosphorylation site at position S276.

Rewiring of signal transduction network due to loss of a known phosphorylation site on RUNX1

We experimentally validated this striking example of a loss of a known phosphorylation sites on

RUNX1 and showed that this variant has an impact on the interactome of RUNX1. The

transcription factor RUNX1 is mutated in 3.03% of melanoma patients and so far, 43 mutations

are described in the literature for cutaneous melanoma. The variant site S276L of RUNX1 is

located in a highly modified region of the protein and may influence the nearby transcriptional

activation domain. Wee *et al.* showed *in vitro* that the triple phosphorylation at the sites S249,

T273 and S276 are important for the interaction with the histone acetyltransferase p300 and thus

lead to the regulation of gene transcription via chromatin remodelling (38). Here, we could not

identify p300 in the interactome studies of RUNX1 by immunoprecipitation of overexpressed

RUNX1 or synthetic peptide pulldowns. However, we identified the transcriptional activator

WWTR1 (TAZ) and KAT7 and the corresponding transcriptional repressors HDAC1 and SIN3A

to be changing between reference and alternate pulldown of RUNX1. The loss of the interaction

to HDAC1 by mutating RUNX1 at S48, S303 and S424 to aspartic acid *in vitro* has been described

previously (39). Here, we hypothesised that the interaction is associated with the modification

status of the protein. The crosstalk between acetylation/deacetylation mediated and

phosphorylation/dephosphorylation may alter the transcriptional activity by RUNX1**.** It is well

known in the literature that RUNX1_wt switches between active and repressive state due to

modifications like acetylation and phosphorylation and binding of interaction partners like

HDAC1, PRMT1 and P300 (40). We postulate that RUNX1_S276L remains in the active state due

the loss of binding to transcriptional repressors like HDAC1 and SIN3A, which could lead to the

accumulation of acetylation on the protein itself as well as histones. This may result in stronger

transcriptional activity, which should be tested in further experiments. Taken together, we

postulate that this variant, which influence the modification status of the protein, changed the

interactome of RUNX1 and altered the transcriptional activity of RUNX1.

Conclusions

Proteogenomics is a powerful tool to study the mode of action of disease-associated mutations at

the genome, transcriptome, proteome and PTM level. Here, we applied a proteogenomics

workflow to study the melanoma cell line A375 sensitive and resistant to BRAF inhibition. The

investigation and integration of multi-omic datasets allowed us to reconstruct the perturbed

signalling networks associated with BRAFi resistance. This resulted in the prioritisation of key

actionable nodes and the prediction of drug therapies with the potential to disrupt BRAFi resistance

mechanism in A375 cells. Notably, we demonstrated the use of AURKA inhibitor as an effective

and specific drug against BRAFi resistant A375 cells. We also detected the loss or gain of several

phosphorylation events due to variants. We could confirm the loss of Ser276 phosphorylation site

by MS as a direct consequence of variant S276L on the transcription factor RUNX1. Our results

suggest that this mutation has an impact on the interactome of RUNX1 and may be responsible for

change in its transcriptional activity. We believe that such proteogenomics workflow is readily

applicable to other types of cancer and within cell lines as well as patient-derived samples.

Materials and Methods

Only star methods are presented below; the rest of the methods is described fully in **Supplementary Information.**

Cell culture

The human metastatic BRAF^{V600E}-mutated melanoma cell line A375 (CRL-1619, ATCC) was used in this study and authenticated by Microsynth AG. The generation of the cell line with acquired resistance to vemurafenib analogue PLX4720 (Selleckchem) (for simplicity referred to as "vemurafenib" in the Results section) was conducted as described previously (13). A375 S and R cells were grown in RPMI medium (Sigma-Aldrich) supplemented with FBS (10%, PAN Biotech) and penicillin/streptavidin (100 U/ml, PAN Biotech) at 37°C and 5% $CO2$. For immunoprecipitation assays, SILAC-labelling of cells was performed as described previously (41) and detailed description of labelling of cells, CRISPR/Cas9-mediated knockout of RUNX1 and interaction assays can be found in the **Supplementary Information**.

Incorporation of non-synonymous variants into protein databases

To integrate the proteogenomic datasets, we used an in-house bioinformatics pipeline, which is coded entirely in the R programming language (42). The transcript nucleotide sequences were extracted from GRCh38 *H. sapiens* genome assembly and Ensembl transcript annotation (via BSgenome and GenomicFeatures packages). These sequences were then *in silico* translated (from start to first stop codon) into a reference protein sequences database (Biostrings package). The called variants, within Variant Call Format (VCF) files from A375 R and A375 S, were injected into each overlapping reference transcript nucleotide sequences and then *in silico* translated. The

resulting protein sequences were written into two FASTA files containing reference variant protein

sequences and sample-specific alternate variant protein sequences.

Annotation of the biological impact of detected variants

In the current study, we prioritised amino acid variants based on their impact in context of BRAFi

resistance in melanoma. For this purpose, known variant sites in melanoma, as well as known

variant sites in cancer, were obtained from CGDS (43). These were overlapped with A375

identified variants and classified as loss/gain of sites. A list of oncogenes and tumour suppressor

genes was compiled from Cosmic, ONGene, Bushman lab and Uniprot (30,44,45), whereas a list

of genes harbouring somatic mutation in BRAFi resistant tumour was retrieved after reanalysis of

published study (14). A375 variants found on these genes were annotated as relevant in cancer

and/or BRAFi resistance.

A second impact scoring strategy was also performed to investigate protein phosphorylation-based

signal transduction networks in A375 melanoma cells. Each reference/alternate variant protein

sequence was annotated based on whether phosphorylation sites (S/T/Y) were lost and/or gained

(IRanges package). A list of known kinase motifs was retrieved from PhosphoNetworks (46) and

these motifs were searched along the reference/alternate variant protein sequences. Located kinase

motifs were overlapped with variants position to determine loss/gain of the motifs. Known human

phosphorylation sites were retrieved from PhosphoSitePlus and Phospho.ELM databases (47,48).

This study identified variants, which affected these sites, were annotated as loss/gain of

phosphorylation. In a similar fashion, known variant sites in melanoma were obtained from CGDS

(43) and overlapped with the variants from A375 R and S. A list of oncogenes and tumour

suppressor genes was compiled from Cosmic, ONGene, Bushman lab and Uniprot (30,44,45).

Variants on these genes that were identified in A375 R and S were annotated as cancer-relevant.

A Levenshtein similarity score was calculated between reference and alternate variant protein

sequences, whereby alternate sequences with less than 90% similarity to their reference were

flagged.

Each amino acid within variant protein sequences were attributed a "+1" score for every overlap

with an impact annotation. A summed score was then calculated for each amino acid within

alternate variant sequence, and the maximum summed score was reported for that variant protein

isoform. Because the score depends on the number of impacts used during the annotation, we also

computed a scaled maximum score (between 0 and 1), to allow comparison between processings.

Following the computation of all impacts, each variant protein isoform is ranked to allow

prioritisation for follow up studies.

Extraction and digestion of proteins

Cells were harvested at 80% confluence with lysis buffer (6 M urea, 2 M thiourea, 60 mM Tris pH

8.0) complemented with protease (complete Mini EDTA-free tablets, Roche) and phosphatase

inhibitors (5 mM glycerol-2-phosphate, 5 mM sodium fluoride, and 1 mM sodium orthovanadate)

and 1% N-ocetylglucoside (NOG, Sigma-Aldrich) for 10 min on ice. DNA and RNA were

removed from the cell lysate using benzonase (1 U/ml, Merck Millipore) for 10 min on room

temperature (RT). Cell debris was cleared by centrifugation (2,800 xg, 10°C, 20 min). Proteins

were precipitated from cell lysates using eight volumes of acetone (-20°C) and one volume of

methanol and incubated overnight at -20°C. The resulting solution was centrifuged (2,800 xg,

10°C, 20 min) to form a cell pellet. The pellet was washed two times with 80% acetone (-21°C)

and resuspended in lysis buffer without NOG. Protein concentration was measured using Bradford

assay. 3 mg of extracted proteins were reduced with 10 mM of dithiothreitol (DTT) for 1 h,

alkylated with 55 mM iodoacetamide for an additional hour and digested with Lys-C (Lysyl

Endopeptidase, Wako Chemicals) for 3 h at RT. After adding four volumes of 10 mM ammonium

bicarbonate, proteins were digested with trypsin (Promega Corporation) overnight. To stop the

digestion, 1% TFA was added.

Detailed description of high pH reverse phase chromatography can be found in the

Supplementary Information.

Phosphopeptide enrichment

Phosphopeptides were enriched using TiO_2 beads (Titansphere, 10 µm, GL Sciences). 1 mg of

beads (in 80%, 1% TFA) were added to acidified high pH fractions and incubated for 30 min in a

rotation wheel. Phosphopeptide-bound TiO_2 beads were sequentially washed with 30% ACN, 1%

TFA, followed by 50% ACN, 1% TFA and 80% ACN, 1% TFA. Peptides were eluted with 5%

NH_4OH into 20% TFA followed by 80% ACN in 1% FA. The eluate was reduced by vacuum

centrifugation, pH was adjusted to < 2.7 with TFA and peptides were desalted on C18 StageTips

prior LC-MS/MS measurements.

Liquid chromatography - mass spectrometry

Peptides were measured on an EASY-nLC 1200 ultra-high-pressure system (Thermo Fisher

Scientific) coupled to a quadrupole Orbitrap mass spectrometer (Q Exactive HF and HFX, Thermo

Fisher Scientific, USA) via a nanoelectrospray ion source. About 1 µg of peptides was loaded on

a 20-cm analytical HPLC-column (75 µm ID PicoTip fused silica emitter (New Objective); in-

house packed using ReproSil-Pur C18-AQ 1.9-µm silica beads (Dr Maisch GmbH)). Peptides were

eluted using a 90 min gradient for proteomic, phosphoproteomic and synthetic peptide pulldown

studies and 60 min gradients for RUNX1 interaction studies. Gradient was generated by solvent A

(0.1% FA) and solvent B (80% ACN in 0.1% FA) and 200 nl/min. Column temperature was kept

at 40°C. The mass spectrometer was operated in data-dependent mode, collecting MS spectra in

the Orbitrap mass analyzer (60,000 resolution, 300–1650 *m/z* range) with an automatic gain control

(AGC) target of 3E6 and a maximum ion injection time of 25 ms. For higher-energy collisional

dissociation (HCD), the 12 most intensive peptides were selected and MS/MS spectra were

recorded with a resolution of 30,000 (fill time 45 ms). For phosphoproteomic studies, top7 method

was used with a resolution of 60,000 for HCD scans and maximum fill time of 220 ms. For the

analysis of RUNX1 interactome (measured on Q Exactive HFX), full MS were acquired in the

range of 300 - 1750 *m/z* at a resolution of 60,000 (fill time 20 ms). Twelve most abundant precursor

ions from a survey scan were selected for HCD fragmentation (fill time 110 ms) and MS/MS

spectra were acquired at a resolution of 30,000 on the Orbitrap analyzer. Precursor dynamic

exclusion was enabled with a duration of 20 s. Synthetic peptide pulldowns were analysed with a

top7 method with a resolution of 60,000 and a fill time of 110 ms.

Mass spectrometry data processing

The raw data files were processed with the MaxQuant software suite (version 1.6.8.0 and 1.5.2.8)

(49). The Andromeda search engine searched MS/MS data against *H. sapiens* reference (99,354

entries) and cell line-specific alternate databases (A375 = 29,104 entries), as well as UniProt *H.*

sapiens (release 2019/02/13; 95,943 entries) database and commonly observed contaminants.

Carbamidomethylation of cysteine (C) was set as fixed modification and oxidation of methionine,

phosphorylation at serine, threonine or tyrosine were defined as variable modifications. Trypsin/P

was selected as a protease. No more than two missed cleavages were allowed. The MS tolerance

was set at 4.5 ppm and MS/MS tolerance at 20 ppm for the analysis using HCD fragmentation

method. The false discovery rate (FDR) for peptides and proteins was set to 1%. For label-free

quantification, a minimum of one peptide was required. For quantification of proteins in the

immunoprecipitation experiments, the amino acids (Lys4)/(Arg6) and (Lys8)/(Arg10) were

defined as 'medium' and 'heavy' labels for the comparison of RUNX1 overexpressed cell lines.

For all other parameters, the default settings were used.

Proteogenomics integration

Our in-house proteogenomic bioinformatic pipeline was used to integrate WES and MS datasets,

specifically to check which mutations were identified across omic datasets. Initially, the reference

and alternate variant protein sequences were *in silico* digested according to laboratory condition;

i.e. digestion with trypsin and up to two missed cleavages (cleaver package). The overlap of MS-

identified peptides with *in silico* digested peptides led to classification into reference (non-mutated

peptide that overlap the mutation position on reference protein), alternate (mutated peptide that

overlap the mutation position on alternate protein) or unspecific (non-mutated peptide that does

not overlap any mutated positions on reference protein) variant peptides. On the basis of this

peptide classification, we summarised the peptides identification per variant protein isoforms,

allowing coverage characterisation into reference only, alternate only, reference and alternate or

unspecific. We finally focused on PTM (as implemented in the MaxQuant processing), which here

consists in phosphorylation sites. Reference and/or alternate variant peptides found

phosphorylated were flagged as such, as well as those were the phosphorylation occurred directly

on the variant sites (either on reference or alternate variant sequences). This coverage information

is exported within MaxQuant style processing results (tab-separated file as output).

Subsequently, we reconstructed the network of protein-protein (using BioGRID database) and

drug-target (using DrugBank database) interactions; i.e. signalling network of BRAFi resistant

A375 and RUNX1 interactome network (50,51). The specificities of the drugs, interacting with

nodes from the generated network, were calculated based on all possible target reported in

DrugBank database. Drugs were prioritised further by summing the number of interactions their

targets have within the network. The generated networks were exported (using igraph and RCy3

packages) into Cytoscape for further customisation (52).

Conflict of interest

509   The authors declare no conflict of interest.

Author contributions

511   M.S., T.S., B.M. and N.C.N. designed the study. M.S. performed the proteomics experiments,

512   while K.B. helped with the interaction proteomic screen. M.S. and N.C.N. analysed the data and

513   performed statistical analysis. M.S. and N.C.N. wrote the manuscript with the input from all

514   authors.

Acknowledgements

The authors acknowledge Prof. Dr. Yulia Skokowa for fruitful discussions, Prof. Dr. Stefan

Stefanovic for synthetic peptides, Prof. Dr. Stefano Stifani for the pCMV_RUNX1 expression

plasmid, Dr. Karsten Krug and Lena Thiess for their help in the initial stages of the project, as well

as c.ATG Core Facility in Tuebingen for the WES library preparation and sequencing. This work

was supported by the High Performance and Cloud Computing Group at the Center for Data

Processing of the University of Tuebingen, the state of Baden-Wuerttemberg (bwHPC), Deutsches

Konsortium für Translationale Krebsforschung (DKTK), German Research Foundation (DFG)

grants No. INST 37/935-1 and INST 37/741-1 FUGG (to B.M.), and by intramural funding from

the University of Tuebingen for the promotion of junior researchers (to N.C.N).

Data availability

526 The high throughput nucleotide sequencing data have been deposited in the NCBI Sequence Read

527 Archive (53) with the bioproject accession number PRJNA616103. The mass spectrometry

528 proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (54)

529 partner repository with the dataset identifier PXD018305. The WES bioinformatics pipeline is

530 available online (55).

References

1. Rajasagi M, Shukla SA, Fritsch EF, Keskin DB, DeLuca D, Carmona E*, et al.* Systematic identification of personal tumor-specific neoantigens in chronic lymphocytic leukemia. Blood **2014**;124:453-62

2. Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J*, et al.* Improved survival with vemurafenib in melanoma with BRAF V600E mutation. N Eng J Med **2011**;364:2507-16

3. Flaherty KT, Robert C, Hersey P, Nathan P, Garbe C, Milhem M*, et al.* Improved Survival with MEK Inhibition in BRAF-Mutated Melanoma. New England Journal of Medicine **2012**;367:107-14

4. Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S*, et al.* Mutations of the BRAF gene in human cancer. Nature **2002**;417:949-54

5. Allen EMV, Wagle N, Sucker A, Treacy D, Goetz EM, Place CS*, et al.* The genetic landscape of clinical resistance to RAF inhibition in metastatic melnaoma. Cancer Discov **2014**;4:94-109

6. Mardis ER. A decade's perspective on DNA sequencing technology. Nature **2011**;470:198-203

7. Kim MS, Pinto SM, Getnet D, Nirujogi RS, Manda SS, Chaerkady R*, et al.* A draft map of the human proteome. Nature **2014**;509:575-81

8. Wilhelm M, Schlegl J, Hahne H, Gholami AM, Lieberenz M, Savitski MM*, et al.* Mass-spectrometry-based draft of the human proteome. Nature **2014**;509:582-7

9. Macek B, Mann M, Olsen JV. Global and site-specific quantitative phosphoproteomics: principles and applications. Annual review of pharmacology and toxicology **2009**;49:199-221

10. Olsen JV, Mann M. Status of large-scale analysis of post-translational modifications by mass spectrometry. Molecular & cellular proteomics : MCP **2013**;12:3444-52

11. Mertins P, Mani DR, Ruggles KV, Gillette MA, Clauser KR, Wang P*, et al.* Proteogenomics connects somatic mutations to signalling in breast cancer. Nature **2016**;534:55-62

12. Creixell P, Schoof EM, Simpson CD, Longden J, Miller CJ, Lou HJ*, et al.* Kinome-wide Decoding of Network-Attacking Mutations Rewiring Cancer Signaling. Cell **2015**;163:202-17

13. Sinnberg T, Makino E, Krueger MA, Velic A, Macek B, Rothbauer U*, et al.* A Nexus Consisting of Beta-Catenin and Stat3 Attenuates BRAF Inhibitor Efficacy and Mediates Acquired Resistance to Vemurafenib. EBioMedicine **2016**;8:132-49

14. Long GV, Fung C, Menzies AM, Pupo GM, Carlino MS, Hyman J*, et al.* Increased MAPK reactivation in early resistance to dabrafenib/trametinib combination therapy of BRAF-mutant metastatic melanoma. Nat Commun **2014**;5:5694

15. Otto F, Lübbert M, Stock M. Upstream and downstream targets of RUNX proteins. Journal of cellular biochemistry **2003**;89:9-18

16. Kitabayashi I, Yokoyama A, Shimizu K, Ohki M. Interaction and functional cooperation of the leukemia-associated factors AML1 and p300 in myeloid cell differentiation. EMBO J **1998**;17:2994-3004

17. Hu H, Bliss JM, Wang Y, Colicelli J. RIN1 is an ABL tyrosine kinase activator and a regulator of epithelial-cell adhesion and migration. Curr Biol **2005**;15:815-23

18. Doyotte A, Mironov A, McKenzie E, Woodman P. The Bro1-related protein HD-PTP/PTPN23 is required for endosomal cargo sorting and multivesicular body morphogenesis. Proc Natl Acad Sci U S A **2008**;105:6308-13

19. Sheynkman GM, Shortreed MR, Cesnik AJ, Smith LM. Proteogenomics: Integrating Next-Generation Sequencing and Mass Spectrometry to Characterize Human Proteomic Variation. Annu Rev Anal Chem (Palo Alto Calif) **2016**;9:521-45

20. Zhang B, Wang J, Wang X, Zhu J, Liu Q, Shi Z*, et al.* Proteogenomic characterization of human colon and rectal cancer. Nature **2014**;513:382-7

21. Krug K, Jaehnig EJ, Satpathy S, Blumenberg L, Karpova A, Anurag M*, et al.* Proteogenomic
Landscape of Breast Cancer Tumorigenesis and Targeted Therapy. Cell **2020**

22. Genomes Project C, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM*, et al.* An
integrated map of genetic variation from 1,092 human genomes. Nature **2012**;491:56-65

23. Pfeifer GP, You YH, Besaratinia A. Mutations induced by ultraviolet light. Mutat Res **2005**;571:19-
31

24. Nusinow DP, Szpyt J, Ghandi M, Rose CM, McDonald ER, 3rd, Kalocsay M*, et al.* Quantitative
Proteomics of the Cancer Cell Line Encyclopedia. Cell **2020**;180:387-402 e16

25. Tian Y, Guo W. A Review of the Molecular Pathways Involved in Resistance to BRAF Inhibitors in
Patients with Advanced-Stage Melanoma. Med Sci Monit **2020**;26:e920957-e

26. Kim H, Frederick DT, Levesque MP, Cooper ZA, Feng Y, Krepler C*, et al.* Downregulation of the
Ubiquitin Ligase RNF125 Underlies Resistance of Melanoma Cells to BRAF Inhibitors via JAK1
Deregulation. Cell Rep **2015**;11:1458-73

27. Caputo E, Miceli R, Motti ML, Taté R, Fratangelo F, Botti G*, et al.* AurkA inhibitors enhance the
effects of B-RAF and MEK inhibitors in melanoma treatment. Journal of Translational Medicine
2014;12:216

28. Cerezo M, Rocchi S. New anti-cancer molecules targeting HSPA5/BIP to induce endoplasmic
reticulum stress, autophagy and apoptosis. Autophagy **2017**;13:216-7

29. Shang Y-Y, Yao M, Zhou Z-W, Jian C, Li X, Hu R-Y*, et al.* Alisertib promotes apoptosis and autophagy
in melanoma through p38 MAPK-mediated aurora a signaling. Oncotarget **2017**;8:107076-88

30. UniProt Consortium T. UniProt: the universal protein knowledgebase. Nucleic Acids Res
2018;46:2699

31. Gentile S, Martin N, Scappini E, Williams J, Erxleben C, Armstrong DL. The human ERG1 channel
polymorphism, K897T, creates a phosphorylation site that inhibits channel activity. Proc Natl Acad
Sci U S A **2008**;105:14704-8

32. Reimand J, Bader GD. Systematic analysis of somatic mutations in phosphorylation signaling
predicts novel cancer drivers. Mol Syst Biol **2013**;9:637

33. Wen B, Xu S, Zhou R, Zhang B, Wang X, Liu X*, et al.* PGA: an R/Bioconductor package for
identification of novel peptides using a customized database derived from RNA-Seq. BMC
Bioinformatics **2016**;17:244

34. Wang D, Eraslan B, Wieland T, Hallstrom B, Hopf T, Zolg DP*, et al.* A deep proteome and
transcriptome abundance atlas of 29 healthy human tissues. Mol Syst Biol **2019**;15:e8503

35. Poulikakos PI, Persaud Y, Janakiraman M, Kong X, Ng C, Moriceau G*, et al.* RAF inhibitor resistance
is mediated by dimerization of aberrantly spliced BRAF(V600E). Nature **2011**;480:387-90

36. Parker R, Vella LJ, Xavier D, Amirkhani A, Parker J, Cebon J*, et al.* Phosphoproteomic Analysis of
Cell-Based Resistance to BRAF Inhibitor Therapy in Melanoma. Front Oncol **2015**;5:95-

37. Corazao-Rozas P, Guerreschi P, André F, Gabert P-E, Lancel S, Dekiouk S*, et al.* Mitochondrial
oxidative phosphorylation controls cancer cell's life and death decisions upon exposure to MAPK
inhibitors. Oncotarget **2016**;7:39473-85

38. Wee HJ, Voon DC, Bae SC, Ito Y. PEBP2-beta/CBF-beta-dependent phosphorylation of RUNX1 and
p300 by HIPK2: implications for leukemogenesis. Blood **2008**;112:3777-87

39. Guo H, Friedman AD. Phosphorylation of RUNX1 by cyclin-dependent kinase reduces direct
interaction with HDAC1 and HDAC3. J Biol Chem **2011**;286:208-15

40. Brettingham-Moore KH, Taberlay PC, Holloway AF. Interplay between Transcription Factors and
the Epigenome: Insight from the Role of RUNX1 in Leukemia. Front Immunol **2015**;6:499

41. Schmitt M, Sinnberg T, Nalpas NC, Maass A, Schittek B, Macek B. Quantitative proteomics links
the intermediate filament nestin to resistance to targeted BRAF inhibition in melanoma cells.
Molecular & cellular proteomics : MCP **2019**

42.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
626   Foundation for Statistical Computing; 2018.

627   43.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA*, et al.* The cBio cancer genomics portal:
628   an open platform for exploring multidimensional cancer genomics data. Cancer Discov
629   **2012**;2:401-4

630   44.   Bamford S, Dawson E, Forbes S, Clements J, Pettett R, Dogan A*, et al.* The COSMIC (Catalogue of
631   Somatic Mutations in Cancer) database and website. Br J Cancer **2004**;91:355-8

632   45.   Liu Y, Sun J, Zhao M. ONGene: A literature-based database for human oncogenes. J Genet
633   Genomics **2017**;44:119-21

634   46.   Hu J, Rho HS, Newman RH, Zhang J, Zhu H, Qian J. PhosphoNetworks: a database for human
635   phosphorylation networks. Bioinformatics **2014**;30:141-2

636   47.   Hornbeck PV, Zhang B, Murray B, Kornhauser JM, Latham V, Skrzypek E. PhosphoSitePlus, 2014:
637   mutations, PTMs and recalibrations. Nucleic Acids Res **2015**;43:D512-20

638   48.   Diella F, Cameron S, Gemund C, Linding R, Via A, Kuster B*, et al.* Phospho.ELM: a database of
639   experimentally verified phosphorylation sites in eukaryotic proteins. BMC Bioinformatics
640   **2004**;5:79

641   49.   Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range
642   mass accuracies and proteome-wide protein quantification. Nature Biotechnology **2008**;26:1367-
643   72

644   50.   Oughtred R, Stark C, Breitkreutz BJ, Rust J, Boucher L, Chang C*, et al.* The BioGRID interaction
645   database: 2019 update. Nucleic Acids Res **2019**;47:D529-D41

646   51.   Wishart DS, Knox C, Guo AC, Shrivastava S, Hassanali M, Stothard P*, et al.* DrugBank: a
647   comprehensive resource for in silico drug discovery and exploration. Nucleic Acids Res
648   **2006**;34:D668-72

649   52.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D*, et al.* Cytoscape: a software
650   environment for integrated models of biomolecular interaction networks. Genome Res
651   **2003**;13:2498-504

652   53.   Coordinators NR. Database resources of the National Center for Biotechnology Information.
653   Nucleic Acids Res **2018**;46:D8-D13

654   54.   Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ*, et al.* The PRIDE
655   database and related tools and resources in 2019: improving support for quantification data.
656   Nucleic Acids Res **2019**;47:D442-D50

657   55.   Nalpas N, Schmitt M, Macek B. Whole-exome sequencing (and nanopore RNA-seq) pipeline. 1.1:
658   Zenodo; 2020.

Figures

Figure 1: State-of-the-art proteogenomic workflow to maximise multi-omics coverage.

[A] Number of non-synonymous nucleotide variants per variant type identified by WES for A375

S and R. [B] The number of protein sequences per reference or alternate databases, as well as the

overlap in search space between databases (up to two missed cleavages). [C] Gene and transcript

counts identified by RNA-sequencing for A375 S and R. [D] Protein group and phosphopeptide

site counts quantified by LC-MS/MS for A375 S and R.

Figure 2: Comparison of drug-sensitive and resistant A375 cells at the genome,

transcriptome and proteome level. [A, C and E] Comparison of A375 R vs. A375 S based on

non-synonymous nucleotide variants [A], transcripts [C] and protein groups [E]. Summed

abundance is the $\log_{10}$ sum of A375 R and A375 S for allelic depth [A], transcripts per million

transcripts [C] and label-free quantification intensities [E]. Significantly changing entries are

coloured in red and blue when up-regulated in A375 R and A375 S, respectively (Sig. A with s0,

see **Supplementary Information**). [B] Venn diagram showing the overlap in genes with non-

synonymous somatic mutations between A375 R (identified in this study) and the patients from

Long and colleagues' study (Long et al., 2014). Within the study from Long and colleagues, the

genes with somatic non-synonymous nucleotide variants are represented as shared across one or

more patients. [D and F] The Reactome pathways that are overrepresented based on significantly

changing transcripts [D] and protein groups [F] between A375 R (red) and A375 S (blue). Up to

the top 20 Reactome pathways are displayed (ranked on FDR).

Figure 3: The disturbed signalling network in BRAFi resistant cells can be targeted by a

number of drugs. [A] The interaction signalling network is generated based on significantly

changing non-synonymous nucleotide variants (circle), transcripts (triangle), proteins (diamond)

and phosphorylation sites (square). Only the top 200 entries are displayed (ranked based on their

interaction degree). Entries that have a non-synonymous nucleotide variant only in A375 R are

shown in red, only in A375 S are in blue, shared in both phenotypes are in grey and not mutated

are in green. Entries that can be targeted by a drug are displayed with a red stroke. Entries were

prioritised based on their importance in context of melanoma and BRAFi resistance (impact factor)

and their node size increased accordingly. [B] The drugs (approved by FDA), interacting with

entries from the interaction signalling network, are displayed based on their specificity to their

target and how many connections their targets have. Colour-coding corresponds to whether one of

the drug targets contains a non-synonymous nucleotide variant found only in A375 R (red), only

in A375 S (blue), shared in both phenotypes (grey) and not mutated (green). [C] Cell viability

assay of A375 S and A375 R cells treated with AURKA inhibitor Alisertib at the indicated

concentrations or in combination with the BRAF inhibitor vemurafenib (2 µM). Cell viability was

determined with MTS assay 72h after treatment start (n=6). Error bars represent standard

deviations of replicates. [D] Scatter plot of the non-synonymous amino acid variants that have an

impact on protein phosphorylation status either as a loss of a S/T/Y amino acids (green), gain of

S/T/Y (purple) or loss/gain of S/T/Y (blue). The intensity ratio of A375 R vs. A375 S corresponds

to the identified alternate peptide, while the scaled impact factor allows ranking of non-

synonymous amino acid variants due to their importance in context of melanoma and

phosphorylation status disruption.

Figure 4: Loss of a known phosphorylation site leads to a changed interactome and altered

transcriptional activity of RUNX1. [A] Schematic overview of the transcription factors RUNX1

protein. Numbers indicate the positions of amino acids residues within the protein. Identified

phosphorylation sites are highlighted in blue and identified amino acid variants are highlighted in

red. Identified peptides by LC-MS/MS are shown in the second panel. Phosphorylated peptides

are indicated with a blue border, while reference and alternate variant peptides are highlighted in

green and red, respectively. [B] Interaction proteomics screen in A375 RUNX1_KO cells stably

overexpressing Flag-tagged RUNX1_wt or Flag-tagged RUNX1_S276L. SILAC protein

expression (log_2) of RUNX1_wt or Flag-tagged RUNX1_S276L relative to the corresponding

control cell line (Flag tag only). RUNX1 and its core binding factor CBFB are marked in black.

Significantly up and downregulated proteins are highlighted in red. Results represent three

replicates per experiment group. [C] Volcano plot of synthetic alternate peptide (Syn_Leu) versus

synthetic reference peptide (Syn_Ser) pulldowns of A375 cells. The log_2 fold change in abundance

between Syn_Leu and Syn_Ser are plotted against $-log_{10}$ p-value (n=3). Black lines indicate the

significance threshold based on student t-test (FDR < 0.01; S0=1.2). Significantly up and

downregulated proteins are highlighted in red. [D] Schematic overview of proposed interaction of

RUNX1_wt and RUNX1_S276L with main transcriptional regulators.