TITLE PAGE

Title: Modeling intrahippocampal effects of anterior hippocampal hyperactivity relevant to schizophrenia using chemogenetic excitation of long axis-projecting mossy cells in the mouse dentate gyrus

Authors: James P. Bauer,[1] Sarah L. Rader,[1] Max E. Joffe,[2,3] Wooseok Kwon,[1] Juliana Quay,[4] Leann Seanez,[1] Chengwen Zhou,[5,6] P. Jeffrey Conn,[2,3,6,7] Alan S. Lewis[1,5,6,7,*]

Affiliations:

[1]Department of Psychiatry and Behavioral Sciences, Vanderbilt University Medical Center, Nashville, TN, USA

[2]Warren Center for Neuroscience Drug Discovery, Vanderbilt University, Nashville, TN USA

[3]Department of Pharmacology, Vanderbilt University, Nashville, TN USA

[4]Quantitative and Chemical Biology Program, Vanderbilt University, Nashville, TN USA

[5]Department of Neurology, Vanderbilt University Medical Center, Nashville, TN, USA

[6]Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN USA

[7]Vanderbilt Kennedy Center, Vanderbilt University Medical Center, Nashville, TN, USA

*Corresponding author: Alan S. Lewis, MD, PhD

Department of Psychiatry and Behavioral Sciences

Vanderbilt University Medical Center

465 21st Ave S., MRBIII Room 6140B

Nashville, TN 37240

Phone: 615-875-4027

Fax: 615-343-4622

Email: alan.s.lewis@vumc.org

Running title: Modeling intrahippocampal long-axis hyperactivity

Keywords: schizophrenia; psychosis; mossy cells; hippocampus; hyperactivity; learning and memory

ABSTRACT

Background: The anterior hippocampus of individuals with early psychosis or schizophrenia is hyperactive, as is the ventral hippocampus in rodent models for schizophrenia risk. Hyperactive ventral hippocampal projections to extrahippocampal brain regions contribute to schizophrenia symptoms, but less is known about the functional effects of hyperactive projections within the hippocampal formation long axis. We approached this question by testing whether hyperactivation of ventral dentate gyrus (DG) mossy cells (MCs), which densely project intrahippocampally to the dorsal DG, influences spatial memory, a cognition dependent on intact dorsal DG function.

Methods: In CD-1 mice, we targeted dorsal DG-projecting ventral DG MCs using an adeno-associated virus intersectional strategy. In vivo fiber photometry recording of ventral DG MCs was performed during exploratory behaviors. We targeted excitatory chemogenetic constructs to ventral DG MCs and tested whether their hyperactivation impaired encoding in a spatial memory task.

Results: Ventral DG MCs were activated during behavior related to environmental information gathering (rearing) but not during non-exploratory motor behaviors. Ventral DG MCs made functional monosynaptic inputs to dorsal DG granule cells, with chemogenetic activation of ventral DG MCs leading to increased activity of dorsal DG granule cells. Finally, chemogenetic activation of ventral DG MCs during the encoding phase of an object location memory task impaired retrieval 24 hours later, without effects on locomotion or other exploratory behaviors.

Conclusions: These data suggest that localized hippocampal hyperactivity may have longitudinal intrahippocampal functional consequences, supporting study of longitudinal circuits as targets to mitigate cognitive deficits associated with schizophrenia.

INTRODUCTION

Cognitive impairment in schizophrenia (CIIS) independently predicts functional deficits (1-3). However, there are currently no FDA-approved pharmacotherapies for CIIS, suggesting underlying mechanisms are incompletely understood. Here we aim to harness the power of molecular tools to manipulate neural circuits in rodents to identify underlying mechanisms accounting for robust neuroimaging findings in patients with schizophrenia or early psychosis. Several independent groups have reported that the anterior hippocampus in such patients is reduced in volume (4-8) and hyperactive at baseline (4, 9-12). It was recently shown in early psychosis patients that greater baseline hyperactivity was associated with reduced anterior hippocampal recruitment during cognitive tasks (9). However, the exact relationship between aberrant anterior hippocampal activity and deficits in learning and memory remains unclear (13).

Many rodent models with genetic or developmental insults for modeling schizophrenia etiology also exhibit hyperactivity in the ventral hippocampus (vHPC), the rodent analogue of the human anterior hippocampus (14, 15). Previous work in these models has shown that vHPC hyperactivity drives aberrant dopamine neurotransmission through extrahippocampal projections to nucleus accumbens (reviewed in (16)) and disrupts distinct cognitive domains, such as spatial working memory, via projections to medial prefrontal cortex (17, 18). Much less is known about how hyperactivity in the anterior portion of the HPC might directly interact with the posterior HPC (analogous to the rodent dorsal HPC (dHPC)) via HPC long axis circuitry. This question is important because certain cognitive deficits present in individuals with schizophrenia are dependent on posterior HPC function, such as visuospatial contextual memory (19-22). Using functional techniques in mice may identify the cognitive consequences of hyperactivity of vHPC intrahippocampal circuits, similar to studies identifying extrahippocampal ramifications.

To model hyperactive long-distance projections originating in the ventral half of the HPC, here we focus on ventral mossy cells (vMCs), which are glutamatergic neurons with cell bodies in the dentate gyrus

(DG) hilus (23, 24). Ventral MCs send long-distance excitatory projections to both ipsi- and contralateral

DG, resulting in dense connectivity of the entire longitudinal hippocampal axis (25-28). Inhibition of

vMCs impairs spatial memory encoding (29), a surprising finding given the reliance on the dHPC for

spatial memory (30-34). Mossy cells are sensitive to even low levels of activity throughout the HPC (35),

and thus are plausible candidates to be functionally responsive to HPC hyperactivity. In this study, we

used vMCs as a longitudinal HPC model circuit by examining their in vivo activity during exploratory

behavior, testing their connectivity with target dDG granule cells, and determining the consequences of

chemogenetic vMC hyperactivation in a spatial memory task that is dependent on dHPC function.

MATERIALS AND METHODS

Detailed methods are reported in the Supplemental Information.

Animals

Wildtype CD-1 male and female mice (Charles River Laboratories, Wilmington, MA; age at delivery: six-eight weeks) were acclimated to the vivarium for at least one week. All mice were group housed except for cannula-implanted mice, which were single housed. Wildtype C57BL/6J male mice (Jackson Laboratory, Bar Harbor, ME; age at delivery: six-seven weeks) were used as intruders during resident-intruder social interactions. Mice were housed under standard conditions with a 12 h light–dark cycle (lights on at 6:00 am) with food and water available ad libitum. The Vanderbilt Institutional Animal Care and Use Committee approved all procedures.

Drugs

Clozapine N-oxide (CNO) for systemic administration was provided by the NIMH Chemical Synthesis and Drug Supply Program, dissolved in 0.5% dimethyl sulfoxide in saline, and injected intraperitoneally (i.p.) 30 mins prior to start of behavioral assay. For slice electrophysiology, CNO was purchased from Hello Bio (Princeton, NJ).

Adeno-associated viruses

Adeno-associated virus particles (AAVs) used, purchased from Addgene (Watertown, MA), were AAV.pgk.Cre, pAAV.Syn.Flex.GCaMP6f.WPRE.SV40 (36), pAAV-hSyn-DIO-mCherry, pAAV-hSyn-DIO-hM3D(Gq) (37), pENN.AAV.hSyn.Cre.WPRE.hGH, and pAAV-EF1a-double floxed-hChR2(H134R)-mCherry-WPRE-HGHpA.

Stereotaxic surgery and viral infusion

Under inhaled isoflurane anesthesia, mice were mounted in a stereotaxic frame, and AAV (0.2 – 0.4 µL depending on brain region) was infused at a rate of 50 nL/min using a 2 µL syringe (Hamilton, Reno, NV). Dorsal DG: anterior/posterior (AP): -1.94 mm, medial/lateral (ML): 1.20 mm left; dorsal/ventral (DV): -2.50 mm. Ventral DG hilus: AP: -3.40 mm, ML: ±3.00 mm, DV: -3.50 mm. Fiber photometry cannulas were targeted to the same coordinates as ventral DG hilus and cemented in place with C&B-Metabond (Parkell, Edgewood, NY). Mice were housed for at least three weeks prior to experimentation to allow for viral expression. Targeting was confirmed by fluorescence microscopy. Mice with either unilateral or bilateral vMC expression were included in behavioral analyses due to extensive bilateral projections.

Fluorescent immunostaining, microscopy, and cell counting

Primary antibodies for 40 µm free-floating sections were mouse anti-calretinin (MAB1568, Millipore, Burlington, MA (38)) and rabbit anti-cFos (226 003, Synaptic Systems, Goettingen, Germany (39)), both diluted 1:1000. Widefield fluorescent images were acquired with an AF6000 LX fluorescent microscopy system (Leica, Buffalo Grove, IL) equipped with an HCX PL FLUOTAR 10x objective (NA = 0.30). Confocal images were acquired with an LSM 710 META Inverted microscope (Zeiss, White Plains, NY) equipped with a 20x Plan-Apochromat objective (NA = 0.8). Cell counting was performed blinded from confocal images.

Electrophysiology

To validate DREADD activation of vMCs, mice were rapidly decapitated under deep isoflurane anesthesia and 300-µm horizontal sections containing the vDG were prepared in *N*-methyl-D-glucamine cutting solution and recorded in artificial cerebrospinal fluid. Hilar MCs were identified by mCherry expression, patched to obtain whole cell access, and voltage-clamped at -75 mV while 10 µM CNO was

perfused in the bath. The mean of the holding current was quantified before and after CNO perfusion using pClamp 10.4 software (Axon Instruments, Union City, CA). For optogenetic activation of vMC terminals in dDG, brain slices were prepared as previously described (40, 41). Whole-cell patch-clamp recordings were made from dDG granule cells and slices were continuously superfused with ACSF. Current-clamp mode was used to record action potentials. 473 nm blue laser light (5 ms pulse duration, 20 Hz, 1 s duration) was delivered through a fiberoptic cable to stimulate channelrhodopsin-expressing vMC axon terminals. In both sets of experiments, recordings were acquired with a Multiclamp 700B amplifier (Molecular Devices, Sunnyvale, CA), filtered at 2 kHz and digitized at 10 kHz.

Behavioral studies

General

Mice were habituated to the dimly lit testing room for at least one hr before testing. Testing occurred between 0900 and 1700. Experimental and control groups were tested on the same days and all tests were videotaped. Vehicle or CNO was injected i.p. 30 mins prior to the start of behavioral assays.

Open field test

Mice were placed into the center of a 61 cm by 61 cm square arena for 10 mins. Distance traveled, time spent in arena center, and number of center entries were quantified using ANY-maze version 6.1 (Stoelting, Wood Dale, IL). Rearing, defined as standing upright on hind legs, was quantified manually by an observer blind to treatment group.

Object location memory test

Performed largely as described previously (42). Mice were handled for about two mins daily for five days, followed by five mins of habituation daily in an empty 61 cm by 61 cm open field containing spatial cues on the wall for six days (final two days of handling were combined with habituation). The next day,

mice underwent an encoding session during which they were placed in the same arena now containing two identical 500 mL glass bottles spaced ~30 cm apart and allowed to explore the arena for 10 mins. Bottles were cleaned with dilute alcohol solution between mice. 24 hrs later, mice were returned to the same arena for a 5 min retrieval session where one bottle remained in the same location and the other bottle was moved ~40 cm to a novel location. The side from which the bottle was moved was counterbalanced across mice. Time spent exploring each bottle was quantified from video by blinded observers. Only time during which mice directly explored the object (~0-2 cm away) and were not rearing on hindlegs was counted (42). Discrimination index (DI) was quantified as (time spent exploring novel object – familiar object)/(total time spent exploring both objects). Mice demonstrating an initial side preference during the encoding session (DI > 0.20) (29) were removed from the analysis (n = 2 mCherry, 3 hM3Dq).

cFos expression

Mice were habituated to the testing room for >2 hours, administered CNO or vehicle i.p., and transcardially perfused 90 minutes later.

Resident-intruder test for fiber photometry

Performed similarly to that previously described (43). To start the assay, a C57BL/6J mouse was introduced to the home cage of the photometry cannula-implanted single housed CD-1 male and allowed to interact for >10 mins.

Fiber photometry

Photometry was performed from male CD-1 mice implanted with photometry cannulas (Doric Lenses, Quebec, Canada) using Doric Lenses system controlled by Doric Neuroscience Studio (DNS) version 5.3. Video recording was time-locked with photometry signal. Signal processing was performed by DNS photometry analyzer. $\Delta F/F_0$ was calculated for the 405 nm and 465 nm channels independently using a least mean square fit, and then $\Delta F/F_0$ (405 nm) subtracted from $\Delta F/F_0$ (465 nm) to yield a corrected trace that was subsequently lowpass filtered at 2 Hz (44) to yield the final bulk calcium signal. This signal was Z-normalized across the entire social interaction. Behavior in videos was manually annotated for exploratory rearing (centered around apogee) and offensive attack onset. Data analysis was performed using custom written scripts in Matlab R2019a (MathWorks, Natick, MA).

Statistical analysis

To compare two groups, unpaired t tests with Welch's correction or paired t tests were used, as appropriate. To compare three or more groups, one- or two-way ANOVA with Sidak's multiple comparisons test was used. One-sample t tests were used to compare experimental to hypothetical means. $p < 0.05$ was considered significant and all tests were two-tailed. Statistical analyses were performed in Prism 9 (Graphpad, San Diego, CA). Error bars depict standard error of the mean (s.e.m.).

RESULTS

Targeting vMCs projecting to the dDG

We exploited the well-organized distant bilateral axonal projection of vMCs to the inner molecular layer

of the dDG (**Figure 1A**) (25, 27, 45-47) to express proteins of interest in longitudinal-projecting vMCs in

wildtype mice. Retrograde-AAV-pgk-Cre was unilaterally infused into dDG and Cre-dependent

GCaMP6f was infused into the contralateral vDG hilus (**Figure 1B**). This resulted in the expression of

GCaMP6f in large ventral hilar cells, with a band of GCaMP6f+ projections to the contralateral vDG and

bilateral dDG inner molecular layers (**Figure 1C**). Our stereotaxic targeting of vMCs was selected to be

consistent with previous functional studies of both vMCs (29) and vDG granule cells (48), and conforms

with a recent detailed anatomical study of MC projections which defined the dDG as the rostral half of the

DG (AP -1.0 to -2.5 mm) and the vDG as the caudal half of the DG (AP -2.7 to -4.0 mm) (49). Dorsal and

vMCs differ in their expression of calretinin, which in mice is only expressed in vMCs (**Supplementary

Figure 1**) (27, 28, 50). Confirming that our targeting coordinates were consistent with previous

definitions of vMCs, in the hilus we found ~87% of GCaMP6f+ neurons were also calretinin+, while

~68% of calretinin+ neurons were GCaMP6f+. We did identify low levels of GCaMP6f off-target

expression in neighboring vDG granule cells and vCA3 pyramidal neurons (n = 4 mice; **Figure 1D, E**).

Thus, these data show that this strategy is selective but not perfectly specific for vMCs, similar to other

methods for MC targeting using D2-Cre (51) or Crlr-Cre (52) transgenic lines (52, 53).

**Activation of longitudinally-projecting vMCs during behaviors related to environmental

information gathering**

The dDG is a critical brain region for spatial information processing in the rodent (54). While recent in

vivo studies have shown dMCs play key roles in spatial coding (55-57), less is known about the in vivo

activity of vMCs. Because of the longitudinal connectivity between vMCs and the dDG, we wondered

whether vMCs might also contribute environment-based input to the dDG. We used fiber photometry (58,

59) to record bulk GCaMP6f calcium activity in vMCs (**Figure 2A**) as a proxy for neuronal firing during

exploratory rearing events. Rearing on hind legs rapidly changes the mouse's vantage point and

multisensory inputs from local cues to more global environmental landmarks thereby representing a

period of environmental information gathering (60, 61) and is associated with marked changes in DG

activity, including in the hilus (62). We found that vMC activity increased concurrent with rearing apogee

(**Figure 2B**), which was consistent within (**Figure 2C**, 46 rearing events in one mouse) and across

(**Figure 2D**, 158 rearing events from 5 mice) mice. This activity increase might simply result from the

abrupt onset of movement inherent in rearing events. To test whether the abrupt onset of movement in

another behavior that is not exploratory in nature yielded the same activity increase, we examined vMC

activity during the onset of offensive attack during social interaction. Despite offensive attack involving

rigorous onset of movement, we found no consistent change in vMC activity associated with attack

behavior in the same mice during the same behavioral session (**Figure 2E**, 50 attacks from 5 mice).

Ventral MC activity was significantly greater during rearing compared to attack (t(4) = 4.02, p = 0.016)

(**Figure 2F**). These experiments suggest that vMCs projecting to the dDG are activated during

environmental information gathering, but not non-specifically activated by any locomotor behavior.

Ventral MCs functionally target dDG granule cells

Electron microscopy has revealed that most vMC axonal synaptic targets in the dDG are on spines of

granule cell dendrites in the inner molecular layer (45, 63). To test the functionality of this longitudinal

circuit, we used AAV1 as an anterograde transsynaptic tracer of functional synapses, which has been

demonstrated in a variety of neuronal circuits (64, 65). We infused AAV1-hSyn-Cre unilaterally into the

ventral hilus to serve as the seed region and infused Cre-dependent mCherry (AAV-DIO-mCherry) into

either ipsilateral or contralateral dDG to report Cre recombination (**Figure 3A**). AAV-DIO-mCherry

infusion did not express mCherry in the absence of ventral hilar AAV1-Cre, suggesting that detectable

fluorophore is due to Cre-mediated recombination and not leak expression (**Figure 3B, left**). The addition

of AAV1-Cre in either the contralateral (**Figure 3B, middle**) or ipsilateral (**Figure 3B, right**) ventral

hilus resulted in mCherry+ granule cells in the dDG, along with some labeling of dorsal hilar neurons.

Because this method likely also captured longitudinal projecting vCA3c neurons (66), we used the

intersectional targeting technique shown in Figure 1 to express channelrhodopsin selectively in vMCs and

performed whole-cell patch-clamp recordings from dDG granule cells while stimulating

channelrhodopsin+ terminals in the inner molecular layer using 473 nm blue light at 20 Hz

(**Supplementary Figure 2**) (49). Action potentials were readily detected upon depolarization of granule

cells membrane potential to approximately -60 mV. These data support functional monosynaptic

connectivity between vMCs and dDG granule cells, and are consistent with a recent report from Houser et

al. showing functional connectivity in the opposite direction, i.e. from dorsal MCs projecting to ventral

granule cells (49).

Chemogenetic excitation of vMCs activates dDG granule cells and impairs object location memory encoding

We next aimed to model persistent activation of vMCs and test whether this vMC hyperactivation

influences a behavior dependent on the dHPC formation and dDG: spatial learning and memory (30-34,

54). We expressed the G_q-coupled Designer Receptor Exclusively Activated by Designer Drugs

(DREADD) construct hM3Dq-mCherry (37) or control mCherry in bilateral vMCs (**Figure 4A,B**). Whole

cell recordings from vMCs expressing hM3Dq (n = 4 cells from 3 mice) or mCherry controls (n = 3 cells

from 2 mice) showed a significantly greater inward current in hM3Dq-expressing vMCs after bath

application of the DREADD-ligand 10 µM CNO compared to control (t(5) = 4.24, p = 0.0068) (**Figure

4C**). To determine an appropriate dose of CNO for *in vivo* behavioral experiments, we systemically

injected CNO and stained for cFos expression 90 mins later. We observed low levels of cFos in vMCs

expressing mCherry (n = 3 mice) or hM3Dq (n = 4 mice) after vehicle treatment or in mCherry-

expressing vMCs (n = 4 mice) after 10 mg/kg CNO treatment (**Figure 4D**). cFos expression was robustly

increased in hM3Dq-expressing vMCs (n = 5 mice) by 10 mg/kg CNO treatment, as revealed by a

significant treatment x virus interaction in a two-way ANOVA (F(1,12) = 33.38, p < 10^{-4}; hM3Dq + CNO

vs. all other groups: p < 10^{-4}, Sidak post-test). However, hM3Dq + CNO-treated mice also demonstrated

dense cFos expression in neighboring granule cells, CA3, and CA1 neurons despite their not expressing

hM3Dq, suggestive of overly high levels of activation and possible seizure activity. Indeed, careful

review of recorded behavior revealed myoclonic jerks in a subset of mice (data not shown). Reducing

CNO to 1 mg/kg still significantly activated vMCs in hM3Dq-expressing MCs (n = 3 mice) compared to

mCherry-expressing vMCs (n = 3 mice) (t(2.98) = 5.58, p = 0.012) (**Figure 4E**) but did not hyperactivate

neighboring granule cells. Consistent with excitatory connectivity *in vivo* between vMCs and dDG

granule cells, 1 mg/kg CNO also significantly increased cFos expression in dDG granule cells

(t(3.77)=3.21, p = 0.036) (**Figure 4F**). Based on these studies, we selected 1 mg/kg CNO to examine the

effects of vMC hyperactivation in behavioral assays.

To test how vMC chemogenetic activation affects spatial learning and memory, we used an object

location memory task, which permits temporal separation of encoding and retrieval (42). We focused on

encoding because vMC inhibition impaired encoding but not retrieval in previous studies (29). After

several days of habituation, mice investigated two identical objects within an arena containing spatial

cues (encoding phase), and 24 hrs later returned to the arena where one object was moved to a novel

location and the other remained in place. Greater time spent investigating the novel-located object is

interpreted as intact spatial memory. Given concerns regarding behavioral tests in outbred mice, we first

confirmed that CD-1 mice showed robust object location memory in this paradigm (**Supplementary

Figure 3**). We then treated mice expressing vMC hM3Dq or mCherry with 1 mg/kg CNO 30 mins prior

to the encoding session. Mice received no treatment prior to the retrieval session 24 hrs later, thus

differences during the retrieval session are likely to be due to differences in spatial encoding (**Figure 5A**).

Two-way ANOVA for discrimination index (DI) between the object locations revealed a significant

session x virus interaction (n = 10 mCherry mice, n = 9 hM3Dq mice, $F(1,17) = 7.64$, $p = 0.013$). Sidak post-hoc tests comparing DI between hM3Dq- and mCherry-expressing mice revealed no significant difference during the encoding session ($t(34) = 0.74$, $p = 0.71$) but a significant difference during the retrieval session ($t(34) = 2.76$, $p = 0.018$) whereby control mCherry mice spent more time investigating the moved object (**Figure 5B**). The total amount of time spent investigating the objects did not significantly differ between virus groups during encoding ($t(15.65) = 0.37$, $p = 0.72$) or retrieval ($t(12.65) = 1.40$, $p = 0.18$) sessions (**Figure 5C**). In a 10 min open field test (n = 12 mCherry mice, n = 11 hM3Dq mice) (**Figure 5D**) there were no significant differences in total distance traveled ($t(19.74) = 0.27$, $p = 0.79$) or time in the arena center ($t(15.13) = 0.68$, $p = 0.51$) between virus groups (**Figure 5E, F**). The total number of rearing events also did not differ between groups ($t(20.31) = 0.66$, $p = 0.52$) (**Figure 5g**). Taken together, these data suggest that hyperactivation of vMCs impairs spatial location encoding but does not significantly change exploration of the environment or gross motor output.

DISCUSSION

Here we used an intersectional viral technique to target, record from, and activate vMCs that densely project to the dDG. Our goal was to interrogate this specific intrahippocampal projection that is anatomically well-suited to distribute localized vHPC (anterior HPC in humans) hyperactivity associated with early psychosis or schizophrenia to disrupt cognitive functions reliant on the dHPC formation and dDG. We found that 1) vMCs in vivo are activated during environmental information gathering, 2) vMCs make functional excitatory synapses with dDG granule cells, 3) chemogenetic activation of vMCs activates dDG granule cells in vivo, and 4) chemogenetic activation of vMCs impairs spatial memory encoding without significantly altering exploratory behaviors. These data suggest that vMC activity associated with rearing is likely related to a role in processing environmental cues but does not directly drive rearing or other locomotor behavior. Next, given the relative importance of the dHPC formation in spatial learning and memory compared to the vHPC (31), our data support the idea that sustained

activation of vMCs impairs spatial encoding by aberrantly activating dDG granule cells. Importantly, these results do not preclude the possibility of spatial encoding deficits via non-dDG mechanisms, and future studies will be necessary to target only the vMC to dDG projections

Since MCs target both granule cells and inhibitory interneurons, considerable work has addressed whether the net effect of MC activation is to excite or inhibit DG granule cells (23, 24, 29, 52, 67, 68), primarily in the context of seizure propagation. Less attention has been devoted to the role of MCs, especially vMCs, in behavior relevant to neuropsychiatric disease including learning, memory, novelty, and anxiety, however there is growing research interest in these topics. Distinctions between vMCs and dMCs are supported by physiological (69-72), biochemical (27, 28), and anatomical (49, 72, 73) evidence. It thus may not be sufficient to extrapolate findings from dMCs to all MCs across the HPC longitudinal axis. Of note, an elegant recent study using in vivo microscopy showed increased vMC activity in novel versus familiar environments that was not due to locomotion differences (72), in line with our fiber photometry findings.

Our results, taken together with studies showing inhibition of vMCs also impairs encoding in a similar object location memory task (29), supports an overall model that bidirectional perturbation of vMC activity, either too much or too little, degrades spatial encoding (**Figure 5h**), without influencing exploratory or locomotor behavior itself. The point of convergence is likely dDG granule cells, as previous work showed that activation *or* inhibition of dDG but not vDG granule cells impaired contextual encoding (48). Two recent studies illustrate that the effects of vMCs on learning and memory are likely task specific. Fredes *et al.* showed that vMC activation prior to shock-context pairing *enhanced* fear conditioning when the conditioning occurred in a familiar context (72), an important finding because contextual fear conditioning is markedly stronger in novel versus familiar contexts. Another study reported that vMC chemogenetic activation did not significantly impair encoding in an object location

memory task in which the delay between encoding and retrieval sessions was only one hr (74) as opposed to the 24 hr delay used in the present study and that of Bui *et al.* (29).

The present results, along with post-mortem evidence for cellular and biochemical changes in human hippocampal area CA4 (75-77) support future investigation into vMC function in rodent models for ventral hippocampal hyperactivation and cognitive deficits seen in schizophrenia. A key goal is to determine why vMC inhibition or excitation impairs spatial encoding. One plausible explanation based on the neuroanatomy of vMC projections to the dDG is through interactions with the medial and/or lateral perforant path from the entorhinal cortex, which terminates in the middle or outer molecular layer on granule cell dendrites, respectively, distal to vMC projection terminals (78, 79) (49). Hypo- or hyperactivation of vMC projections to dorsal granule cells may impair the dDG's critical function of transforming inputs into sparse ensembles of activated granule cells for pattern separation (80-85). Future studies using mouse models with genetic lesions conferring high risk for schizophrenia in humans that show vHPC hyperactivity and object location memory deficits, such as 22q11.2 deletion models (86, 87), are well-suited to test whether modulating vMC longitudinal projections may rescue spatial memory deficits. Finally, chemogenetic manipulations in this study were acute, and it has been shown that learning and memory changes following viral MC ablations are surprisingly transient (52). Given aberrant neurodevelopmental processes in schizophrenia, it will be important to test whether chronic vMC inhibition or excitation impairs spatial memory in a similar manner to acute manipulations.

ACKNOWLEDGEMENTS

This work was supported by National Institutes of Health grants MH116339 (A.S.L.) and NS107424 (C.Z.), and the Vanderbilt Department of Psychiatry and Behavioral Sciences. Experiments and data analysis were performed in part using the Vanderbilt University Medical Center Cell Imaging Shared Resource (supported by NIH grants CA68485, DK20593, DK58404, DK59637 and EY08126). We thank Marina Picciotto, Ph.D. and Ye Han, Ph.D. for helpful comments on this work. This work has been posted as a preprint to bioRxiv.

DISCLOSURES

REFERENCES

1.   Fett AK, Viechtbauer W, Dominguez MD, Penn DL, van Os J, Krabbendam L (2011): The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: a meta-analysis. *Neurosci Biobehav Rev*. 35:573-588.

2.   Green MF (1996): What are the functional consequences of neurocognitive deficits in schizophrenia? *Am J Psychiatry*. 153:321-330.

3.   Mohamed S, Rosenheck R, Swartz M, Stroup S, Lieberman JA, Keefe RSE (2008): Relationship of Cognition and Psychopathology to Functional Impairment in Schizophrenia. *American Journal of Psychiatry*. 165:978-987.

4.   Schobel SA, Chaudhury NH, Khan UA, Paniagua B, Styner MA, Asllani I, et al. (2013): Imaging patients with psychosis and a mouse model establishes a spreading pattern of hippocampal dysfunction and implicates glutamate as a driver. *Neuron*. 78:81-93.

5.   Schobel SA, Kelly MA, Corcoran CM, Van Heertum K, Seckinger R, Goetz R, et al. (2009): Anterior hippocampal and orbitofrontal cortical structural brain abnormalities in association with cognitive deficits in schizophrenia. *Schizophrenia Research*. 114:110-118.

6.   McHugo M, Talati P, Woodward ND, Armstrong K, Blackford JU, Heckers S (2018): Regionally specific volume deficits along the hippocampal long axis in early and chronic psychosis. *Neuroimage Clin*. 20:1106-1114.

7.   Szeszko PR, Goldberg E, Gunduz-Bruce H, Ashtari M, Robinson D, Malhotra AK, et al. (2003): Smaller anterior hippocampal formation volume in antipsychotic-naive patients with first-episode schizophrenia. *Am J Psychiatry*. 160:2190-2197.

8.   McHugo M, Armstrong K, Roeske MJ, Woodward ND, Blackford JU, Heckers S (2020): Hippocampal volume in early psychosis: a 2-year longitudinal study. *Translational Psychiatry*. 10:306.

9.   McHugo M, Talati P, Armstrong K, Vandekar SN, Blackford JU, Woodward ND, et al. (2019): Hyperactivity and Reduced Activation of Anterior Hippocampus in Early Psychosis. *Am J Psychiatry*. 176:1030-1038.

10.   Talati P, Rane S, Kose S, Blackford JU, Gore J, Donahue MJ, et al. (2014): Increased hippocampal CA1 cerebral blood volume in schizophrenia. *Neuroimage Clin*. 5:359-364.

11.   Schobel SA, Lewandowski NM, Corcoran CM, Moore H, Brown T, Malaspina D, et al. (2009): Differential targeting of the CA1 subfield of the hippocampal formation by schizophrenia and related psychotic disorders. *Arch Gen Psychiatry*. 66:938-946.

12.   Modinos G, Şimşek F, Azis M, Bossong M, Bonoldi I, Samson C, et al. (2018): Prefrontal GABA levels, hippocampal resting perfusion and the risk of psychosis. *Neuropsychopharmacology*. 43:2652-2659.

13.   Ragland JD (2019): Relating Basal and Phasic Hippocampal Activity in People With Psychosis: A Translational Bridge to Understanding Memory Deficits? *American Journal of Psychiatry*. 176:979-981.

14.   Grimm CM, Aksamaz S, Schulz S, Teutsch J, Sicinski P, Liss B, et al. (2018): Schizophrenia-related cognitive dysfunction in the Cyclin-D2 knockout mouse model of ventral hippocampal hyperactivity. *Transl Psychiatry*. 8:212.

15.   Lodge DJ, Grace AA (2007): Aberrant hippocampal activity underlies the dopamine dysregulation in an animal model of schizophrenia. *J Neurosci*. 27:11424-11430.

16.   Grace AA (2012): Dopamine system dysregulation by the hippocampus: implications for the pathophysiology and treatment of schizophrenia. *Neuropharmacology*. 62:1342-1348.

17.   Sigurdsson T, Duvarci S (2016): Hippocampal-Prefrontal Interactions in Cognition, Behavior and Psychiatric Disease. *Frontiers in Systems Neuroscience*. 9.

18.   Spellman T, Rigotti M, Ahmari SE, Fusi S, Gogos JA, Gordon JA (2015): Hippocampal–prefrontal input supports spatial encoding in working memory. *Nature*. 522:309-314.

19.   Mesholam-Gately RI, Giuliano AJ, Goff KP, Faraone SV, Seidman LJ (2009): Neurocognition in first-episode schizophrenia: a meta-analytic review. *Neuropsychology*. 23:315-336.

20.   Ragland JD, Layher E, Hannula DE, Niendam TA, Lesh TA, Solomon M, et al. (2017): Impact of schizophrenia on anterior and posterior hippocampus during memory for complex scenes. *Neuroimage Clin*. 13:82-88.

21.   Barnett JH, Sahakian BJ, Werners U, Hill KE, Brazil R, Gallagher O, et al. (2005): Visuospatial learning and executive function are independently impaired in first-episode psychosis. *Psychol Med*. 35:1031-1041.

22.   McCleery A, Ventura J, Kern RS, Subotnik KL, Gretchen-Doorly D, Green MF, et al. (2014): Cognitive functioning in first-episode schizophrenia: MATRICS Consensus Cognitive Battery (MCCB) Profile of Impairment. *Schizophr Res*. 157:33-39.

23.   Scharfman HE (2016): The enigmatic mossy cell of the dentate gyrus. *Nat Rev Neurosci*. 17:562-575.

24.   Scharfman HE, Myers CE (2012): Hilar mossy cells of the dentate gyrus: a historical perspective. *Front Neural Circuits*. 6:106.

25.   Ribak CE, Seress L, Amaral DG (1985): The development, ultrastructure and synaptic connections of the mossy cells of the dentate gyrus. *J Neurocytol*. 14:835-857.

26.   Frotscher M, Seress L, Schwerdtfeger WK, Buhl E (1991): The mossy cells of the fascia dentata: a comparative study of their fine structure and synaptic connections in rodents and primates. *J Comp Neurol*. 312:145-163.

27.   Blasco-Ibanez JM, Freund TF (1997): Distribution, ultrastructure, and connectivity of calretinin-immunoreactive mossy cells of the mouse dentate gyrus. *Hippocampus*. 7:307-320.

28.   Fujise N, Liu Y, Hori N, Kosaka T (1998): Distribution of calretinin immunoreactivity in the mouse dentate gyrus: II. Mossy cells, with special reference to their dorsoventral difference in calretinin immunoreactivity. *Neuroscience*. 82:181-200.

29.   Bui AD, Nguyen TM, Limouse C, Kim HK, Szabo GG, Felong S, et al. (2018): Dentate gyrus mossy cells control spontaneous convulsive seizures and spatial memory. *Science*. 359:787-790.

30.   Moser MB, Moser EI, Forrest E, Andersen P, Morris RG (1995): Spatial learning with a minislab in the dorsal hippocampus. *Proceedings of the National Academy of Sciences*. 92:9697.

31.   Moser E, Moser MB, Andersen P (1993): Spatial learning impairment parallels the magnitude of dorsal hippocampal lesions, but is hardly present following ventral lesions. *J Neurosci*. 13:3916-3925.

32.   Hannula DE, Libby LA, Yonelinas AP, Ranganath C (2013): Medial temporal lobe contributions to cued retrieval of items and contexts. *Neuropsychologia*. 51:2322-2332.

33.   Evensmoen HR, Ladstein J, Hansen TI, Møller JA, Witter MP, Nadel L, et al. (2015): From details to large scale: The representation of environmental positions follows a granularity gradient along the human hippocampal and entorhinal anterior–posterior axis. *Hippocampus*. 25:119-135.

34.   Duarte IC, Ferreira C, Marques J, Castelo-Branco M (2014): Anterior/posterior competitive deactivation/activation dichotomy in the human hippocampus as revealed by a 3D navigation task. *PLoS One*. 9:e86213.

35.   Scharfman HE, Schwartzkroin PA (1988): Electrophysiology of morphologically identified mossy cells of the dentate hilus recorded in guinea pig hippocampal slices. *J Neurosci*. 8:3812-3821.

36.   Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, et al. (2013): Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*. 499:295-300.

37.   Krashes MJ, Koda S, Ye C, Rogan SC, Adams AC, Cusher DS, et al. (2011): Rapid, reversible activation of AgRP neurons drives feeding behavior in mice. *J Clin Invest*. 121:1424-1428.

38.   Liu W, Davis RL (2014): Calretinin and calbindin distribution patterns specify subpopulations of type I and type II spiral ganglion neurons in postnatal murine cochlea. *J Comp Neurol*. 522:2299-2318.

39.   Zhou QG, Nemes AD, Lee D, Ro EJ, Zhang J, Nowacki AS, et al. (2019): Chemogenetic silencing of hippocampal neurons suppresses epileptic neural circuits. *J Clin Invest*. 129:310-323.

40.   Zhou C, Huang Z, Ding L, Deel ME, Arain FM, Murray CR, et al. (2013): Altered cortical GABAA receptor composition, physiology, and endocytosis in a mouse model of a human genetic absence epilepsy syndrome. *J Biol Chem*. 288:21458-21472.

41.   Zhou C, Lippman JJ, Sun H, Jensen FE (2012): Hypoxia-induced neonatal seizures diminish silent synapses and long-term potentiation in hippocampal CA1 neurons. *J Neurosci*. 31:18211-18222.

42.   Vogel-Ciernia A, Wood MA (2014): Examining object location and object recognition memory in mice. *Curr Protoc Neurosci*. 69:8 31 31-17.

43.   Lewis AS, Pittenger ST, Mineur YS, Stout D, Smith PH, Picciotto MR (2018): Bidirectional Regulation of Aggression in Mice by Hippocampal Alpha-7 Nicotinic Acetylcholine Receptors. *Neuropsychopharmacology*. 43:1267-1275.

44.   Zalocusky KA, Ramakrishnan C, Lerner TN, Davidson TJ, Knutson B, Deisseroth K (2016): Nucleus accumbens D2R cells signal prior outcomes and control risky decision-making. *Nature*. 531:642-646.

45.   Buckmaster PS, Wenzel HJ, Kunkel DD, Schwartzkroin PA (1996): Axon arbors and synaptic connections of hippocampal mossy cells in the rat in vivo. *J Comp Neurol*. 366:271-292.

46.   Buzsaki G, Eidelberg E (1981): Commissural projection to the dentate gyrus of the rat: evidence for feed-forward inhibition. *Brain Res*. 230:346-350.

47.   Laurberg S, Sorensen KE (1981): Associational and commissural collaterals of neurons in the hippocampal formation (hilus fasciae dentatae and subfield CA3). *Brain Res*. 212:287-300.

48.   Kheirbek MA, Drew LJ, Burghardt NS, Costantini DO, Tannenholz L, Ahmari SE, et al. (2013): Differential control of learning and anxiety along the dorsoventral axis of the dentate gyrus. *Neuron*. 77:955-968.

49.   Houser CR, Peng Z, Wei X, Huang CS, Mody I (2020): Mossy Cells in the Dorsal and Ventral Dentate Gyrus Differ in their Patterns of Axonal Projections. *The Journal of Neuroscience*.JN-RM-2455-2420.

50.   Liu Y, Fujise N, Kosaka T (1996): Distribution of calretinin immunoreactivity in the mouse

dentate gyrus. I. General description. *Exp Brain Res*. 108:389-403.

51.   Gangarossa G, Longueville S, De Bundel D, Perroy J, Herve D, Girault JA, et al. (2012):

Characterization of dopamine D1 and D2 receptor-expressing neurons in the mouse hippocampus.

Hippocampus. 22:2199-2207.

52.   Jinde S, Zsiros V, Jiang Z, Nakao K, Pickel J, Kohno K, et al. (2012): Hilar mossy cell

degeneration causes transient dentate granule cell hyperexcitability and impaired pattern separation.

Neuron. 76:1189-1200.

53.   Puighermanal E, Biever A, Espallergues J, Gangarossa G, De Bundel D, Valjent E (2015): drd2-

cre:ribotag mouse line unravels the possible diversity of dopamine d2 receptor-expressing cells of the

dorsal mouse hippocampus. *Hippocampus*. 25:858-875.

54.   Goodrich-Hunsaker NJ, Hunsaker MR, Kesner RP (2008): The interactions and dissociations of

the dorsal hippocampus subregions: how the dentate gyrus, CA3, and CA1 process spatial information.

Behav Neurosci. 122:16-26.

55.   Danielson NB, Turi GF, Ladow M, Chavlis S, Petrantonakis PC, Poirazi P, et al. (2017): In Vivo

Imaging of Dentate Gyrus Mossy Cells in Behaving Mice. *Neuron*. 93:552-559 e554.

56.   GoodSmith D, Chen X, Wang C, Kim SH, Song H, Burgalossi A, et al. (2017): Spatial

Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. *Neuron*. 93:677-690 e675.

57.   Senzai Y, Buzsaki G (2017): Physiological Properties and Behavioral Correlates of Hippocampal

Granule Cells and Mossy Cells. *Neuron*. 93:691-704 e695.

58.   Gunaydin LA, Grosenick L, Finkelstein JC, Kauvar IV, Fenno LE, Adhikari A, et al. (2014):

Natural neural projection dynamics underlying social behavior. *Cell*. 157:1535-1551.

59.   Cui G, Jun SB, Jin X, Luo G, Pham MD, Lovinger DM, et al. (2014): Deep brain optical

measurements of cell type-specific neural activity in behaving mice. *Nat Protoc*. 9:1213-1228.

60.   Knierim JJ, Hamilton DA (2011): Framing spatial cognition: neural representations of proximal and distal frames of reference and their roles in navigation. *Physiol Rev*. 91:1245-1279.

61.   Lever C, Burton S, O'Keefe J (2006): Rearing on hind legs, environmental novelty, and the hippocampal formation. *Rev Neurosci*. 17:111-133.

62.   Barth AM, Domonkos A, Fernandez-Ruiz A, Freund TF, Varga V (2018): Hippocampal Network Dynamics during Rearing Episodes. *Cell reports*. 23:1706-1715.

63.   Wenzel HJ, Buckmaster PS, Anderson NL, Wenzel ME, Schwartzkroin PA (1997): Ultrastructural localization of neurotransmitter immunoreactivity in mossy cell axons and their synaptic targets in the rat dentate gyrus. *Hippocampus*. 7:559-570.

64.   Zingg B, Chou XL, Zhang ZG, Mesik L, Liang F, Tao HW, et al. (2017): AAV-Mediated Anterograde Transsynaptic Tagging: Mapping Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. *Neuron*. 93:33-47.

65.   Zingg B, Peng B, Huang J, Tao HW, Zhang LI (2020): Synaptic Specificity and Application of Anterograde Transsynaptic AAV for Probing Neural Circuitry. *J Neurosci*. 40:3250-3267.

66.   Li XG, Somogyi P, Ylinen A, Buzsaki G (1994): The hippocampal CA3 network: an in vivo intracellular labeling study. *J Comp Neurol*. 339:181-208.

67.   Botterill JJ, Lu YL, LaFrancois JJ, Bernstein HL, Alcantara-Gonzalez D, Jain S, et al. (2019): An Excitatory and Epileptogenic Effect of Dentate Gyrus Mossy Cells in a Mouse Model of Epilepsy. *Cell reports*. 29:2875-2889 e2876.

68.   Scharfman HE (1995): Electrophysiological evidence that dentate hilar mossy cells are excitatory and innervate both granule cells and interneurons. *J Neurophysiol*. 74:179-194.

69.   Jinno S, Ishizuka S, Kosaka T (2003): Ionic currents underlying rhythmic bursting of ventral mossy cells in the developing mouse dentate gyrus. *Eur J Neurosci*. 17:1338-1354.

70.   Moretto JN, Duffy AM, Scharfman HE (2017): Acute restraint stress decreases c-fos immunoreactivity in hilar mossy cells of the adult dentate gyrus. *Brain Struct Funct*. 222:2405-2419.

71.   Duffy AM, Schaner MJ, Chin J, Scharfman HE (2013): Expression of c-fos in hilar mossy cells of the dentate gyrus in vivo. *Hippocampus*. 23:649-655.

72.   Fredes F, Silva MA, Koppensteiner P, Kobayashi K, Joesch M, Shigemoto R (2020): Ventro-dorsal Hippocampal Pathway Gates Novelty-Induced Contextual Memory Formation. *Curr Biol*.

73.   Botterill JJ, Gerencer KJ, Vinod KY, Alcantara-Gonzalez D, Scharfman HE (2020): Dorsal and ventral mossy cells differ in their long-range axonal projections throughout the dentate gyrus of the mouse hippocampus. *bioRxiv*.2020.2009.2027.315416.

74.   Botterill JJ, Vinod KY, Gerencer KJ, Teixeira CM, LaFrancois JJ, Scharfman HE (2021): Bidirectional regulation of cognitive and anxiety-like behaviors by dentate gyrus mossy cells in male and female mice. *The Journal of Neuroscience*.JN-RM-1724-1720.

75.   Roeske MJ, Konradi C, Heckers S, Lewis AS (2020): Hippocampal volume and hippocampal neuron density, number and size in schizophrenia: a systematic review and meta-analysis of postmortem studies. *Mol Psychiatry*.

76.   Harrison PJ, Law AJ, Eastwood SL (2003): Glutamate receptors and transporters in the hippocampus in schizophrenia. *Ann N Y Acad Sci*. 1003:94-101.

77.   Harrison PJ (2004): The hippocampus in schizophrenia: a review of the neuropathological evidence and its pathophysiological implications. *Psychopharmacology*. 174:151-162.

78.   Hetherington PA, Austin KB, Shapiro ML (1994): Ipsilateral associational pathway in the dentate gyrus: an excitatory feedback system that supports N-methyl-D-aspartate-dependent long-term potentiation. *Hippocampus*. 4:422-438.

79.   Kleschevnikov AM, Routtenberg A (2003): Long-term potentiation recruits a trisynaptic excitatory associative network within the mouse dentate gyrus. *Eur J Neurosci*. 17:2690-2702.

80.   Rolls ET (2010): A computational theory of episodic memory formation in the hippocampus. *Behav Brain Res*. 215:180-196.

81.   Treves A, Tashiro A, Witter MP, Moser EI (2008): What is the mammalian dentate gyrus good for? *Neuroscience*. 154:1155-1172.

82.   Myers CE, Scharfman HE (2009): A role for hilar cells in pattern separation in the dentate gyrus: a computational approach. *Hippocampus*. 19:321-337.

83.   Leutgeb JK, Leutgeb S, Moser MB, Moser EI (2007): Pattern separation in the dentate gyrus and CA3 of the hippocampus. *Science*. 315:961-966.

84.   Chawla MK, Guzowski JF, Ramirez-Amaya V, Lipa P, Hoffman KL, Marriott LK, et al. (2005): Sparse, environmentally selective expression of Arc RNA in the upper blade of the rodent fascia dentata by brief spatial experience. *Hippocampus*. 15:579-586.

85.   Neunuebel JP, Knierim JJ (2014): CA3 retrieves coherent representations from degraded input: direct evidence for CA3 pattern completion and dentate gyrus pattern separation. *Neuron*. 81:416-427.

86.   Didriksen M, Fejgin K, Nilsson SR, Birknow MR, Grayton HM, Larsen PH, et al. (2017): Persistent gating deficit and increased sensitivity to NMDA receptor antagonism after puberty in a new mouse model of the human 22q11.2 microdeletion syndrome: a study in male mice. *Journal of psychiatry & neuroscience : JPN*. 42:48-58.

87.   Kahn JB, Port RG, Anderson SA, Coulter DA (2020): Modular, Circuit-Based Interventions Rescue Hippocampal-Dependent Social and Spatial Memory in a 22q11.2 Deletion Syndrome Mouse Model. *Biol Psychiatry*.

88.   Oh SW, Harris JA, Ng L, Winslow B, Cain N, Mihalas S, et al. (2014): A mesoscale connectome of the mouse brain. *Nature*. 508:207-214.

FIGURE LEGENDS

Figure 1. Intersectional targeting of longitudinal ventral mossy cell projections. A Ventral mossy cell (vMC) to dorsal dentate gyrus (dDG) circuit of interest as illustrated by the Allen Mouse Brain Connectivity Atlas (88). AAV expressing EGFP was infused into ventral hilus of wildtype C57BL/6J mouse (top). Serial two-photon tomography shows axonal projections throughout longitudinal extent of hippocampus, with strong projection to dDG inner molecular layer (https://connectivity.brain-map.org/projection/experiment/112672268). **B** To target vMCs, retrograde-AAV-pgk-Cre was infused into dDG inner molecular layer, and AAV-hSyn-Flex-GCaMP6f was infused into contralateral vDG hilus, enabling recombination only in vMCs that project to dDG. **C** Fluorescence microscopy from CD-1 mouse following targeting strategy shown in (B) reveals GCaMP6f is expressed in vMC somata with organized projections to contralateral vDG and bilateral dDG. **D** GCaMP6f from targeting strategy in (B) is highly colocalized with calretinin, a marker for ventral but not dorsal MCs. **E** Quantification of viral targeting strategy in (B) from n = 4 mice reveals most GCaMP6f+ hilar neurons are calretinin+, with limited off-target expression in nearby ventral granule cells and CA3 pyramidal neurons.

Figure 2. In vivo fiber photometry reveals ventral mossy cells are selectively activated by behavior related to environmental information gathering. A Fluorescence microscopy of GCaMP6f expression in ventral mossy cells (vMCs) using the intersectional targeting strategy and fiber optic cannula placement. **B** Individual photometry trace surrounding an example exploratory rearing event. **C** $\Delta F/F_0$ Z-scores of rearing events aligned to time of rearing apogee in an individual mouse. Events are ordered by Z-score and consistently show increased photometry signal during rearing. **D** $\Delta F/F_0$ Z-scores for N = 5 mice were aligned to time of rearing apogee, averaged within mouse, then averaged across mice. Top, black line depicts mean $\Delta F/F_0$ Z-score, pink depicts s.e.m. Bottom, heatmap showing signal for individual mice. **E** As a control to test whether vMCs were activated non-specifically by abrupt motor behavior

onset, $\Delta F/F_0$ Z-scores for the same N = 5 mice from (D) during the same behavioral session were aligned to time of offensive attack on an interacting conspecific, averaged within mouse, then averaged across mice. Top, black line depicts mean $\Delta F/F_0$ Z-score, green depicts s.e.m. Bottom, heatmap showing signal for individual mice. **F** Area under the curve from -2 s to +2 s aligned to either rearing (pink) or attack (green) for mice in (D-E) shows significantly greater vMC activity associated with rearing than attack during the same behavioral social interaction session, suggesting vMCs are not non-selectively activated by any locomotor onset.

Figure 3. AAV1-mediated anterograde transsynaptic tracing reveals functional synapses between ventral mossy cells and dorsal granule cells. A AAV1-Cre was infused unilaterally into ventral dentate gyrus (DG) hilus, and Cre-dependent mCherry (AAV-DIO-mCherry) was infused either contralaterally (box 1) or ipsilaterally (box 2) to dorsal DG to detect anterograde transsynaptic trafficking. **B** AAV-DIO-mCherry does not express mCherry when infused alone (left), however additional infusion of AAV1-Cre in either contralateral (middle) or ipsilateral (right) DG results in dorsal DG granule cell expression of mCherry, supporting functional synaptic connectivity between the ventral hilus and dorsal DG most consistent with longitudinal mossy cell projections.

Figure 4. Chemogenetic activation of ventral mossy cells activates dorsal granule cells. A To express hM3Dq-mCherry or mCherry in ventral mossy cells (vMCs), retrograde-AAV-pgk-Cre was unilaterally infused into dorsal dentate gyrus (DG) and AAV-hSyn-DIO-hM3D(Gq)-mCherry or AAV-hSyn-DIO-mCherry was bilaterally infused into ventral DG hilus. **B** Fluorescence microscopy showing representative image of bilateral vMC targeting. **C** Whole cell recordings from vMCs expressing either mCherry (n = 3 cells|2 mice) or hM3Dq-mCherry (n = 4 cells|3 mice) revealed hM3Dq+ neurons showed significantly greater mean inward current after bath application of 10 micromolar clozapine N-oxide (CNO) than control mCherry+ neurons. **p = 0.0068. **D** Mice with vMCs expressing mCherry (n = 3-4)

or hM3Dq-mCherry (n = 4-5) were administered vehicle or 10 mg/kg CNO i.p., then perfused 90 mins later and immunostained for cFos. Only vMCs expressing hM3Dq-mCherry strongly expressed cFos, but also strongly expressed cFos in neighboring ventral DG granule cells. ****$p < 10^{-4}$. **E** Same method as in (D) but with reduced dose of CNO (1 mg/kg) showed increase in cFos in vMCs expressing hM3Dq, but no hyperactivation of neighboring granule cells. *$p = 0.012$. **F** Dorsal DG granule cell cFos was also significantly upregulated in mice with vMC expression of hM3Dq-mCherry treated with 1 mg/kg CNO as compared to mice with vMC expression of mCherry. *$p = 0.036$.

Figure 5. Ventral mossy cell hyperactivation impairs object location memory encoding. A Schematic of the object location memory task. After three days of handling, mice undergo six days of habituation to an arena with spatial cues on the wall. The following day they undergo a 10 min encoding session during which they are exposed to two identical objects. 24 hrs later they are returned to the arena for a 5 min retrieval session where one object has been moved to a novel location and the other remains in the familiar location. 1 mg/kg CNO was administered 30 mins before the encoding session only. **B** CNO administration during encoding reduced object location retrieval in mice with vMC hM3Dq expression (n = 10 mCherry, n = 9 hM3Dq). **C** Total time spent investigating both objects did not significantly differ between the mCherry or hM3Dq-expressing mice during the encoding (left) or retrieval session (right). **D** Mice were later tested in a 10 min open field test in an arena with a novel floor surface 30 mins after 1 mg/kg CNO (n = 12 mCherry, n = 11 hM3Dq). There was no significant difference between mCherry and hM3Dq-expressing mice in total distance travelled (**E**), time spent in arena center (**F**), or number of rearing events (**G**). **H** Schematic suggesting that a specific setpoint of vMC activity exists to promote optimal encoding performance, which is supported by previous work showing that inhibition of vMCs impairs spatial encoding (29) and the current study suggesting that hyperactivation of vMCs also impairs spatial encoding.

A

Injection site

Allen Mouse Brain Connectivity Atlas
Experiment 112672268
C57BL/6J, anterograde EGFP tracer

B

rg-AAV-pgk-Cre

Bregma -1.94

AAV-hSyn-Flex-GCaMP6f

Bregma -3.40

— ipsilateral projection

--- contralateral projection

C

Dorsal dentate gyrus

Ventral dentate gyrus

GCaM6f/DAPI

150 µm

D

GCaMP6f

Calretinin

DAPI/Merge

50 µm

E

GFP+Calretinin+/GFP+ (%)

GFP+Calretinin+/Calretinin+ (%)

GFP+ granule cells per 10^5 µm^2

GFP+ CA3 pyramidal neurons per 10^5 µm^2

A

AAV-DIO-mCherry

1   2

Anterograde transport

AAV1-Cre

B

mCherry
DAPI

Contralateral

Ipsilateral

100 µm

Dorsal DG: DIO-mCherry   DIO-mCherry   DIO-mCherry
Ventral DG: No virus   AAV1-Cre   1   AAV1-Cre   2

A rg-AAV-pgk-Cre

AAV-hSyn-DIO-hM3Dq-mCherry

B hM3Dq-mCherry
DAPI
150 μm

C
10 μM CNO

mCherry
hM3Dq

100 pA
30 sec

I_{CNO} (pA)
**
50
0
-50
-100
-150

D
AAV-hSyn-DIO-mCherry
Vehicle | 10 mg/kg CNO

AAV-hSyn-DIO-hM3Dq-mCherry
Vehicle | 10 mg/kg CNO

mCherry

cFos

mCherry
cFos
DAPI

100 μm

cFos+mCherry+/mCherry+ in hilus (%)
○ mCherry
● hM3Dq
****
Vehicle   10 mg/kg CNO

cFos+ granule cells per mm²
○ mCherry
● hM3Dq
150
100
50
0
Vehicle   10 mg/kg CNO
hM3Dq: too dense to quantify

E
AAV-hSyn-DIO-mCherry + 1 mg/kg CNO
mCherry | cFos | mCherry cFos DAPI
100 μm

AAV-hSyn-DIO-hM3Dq-mCherry + 1 mg/kg CNO

cFos+mCherry+/mCherry+ in hilus (%)
*
mCherry   hM3Dq

F Dorsal dentate gyrus
mCherry + 1 mg/kg CNO
mCherry
cFos
DAPI
100 μm

hM3Dq + 1 mg/kg CNO

cFos+ dorsal granule cells per mm²
300
200
100
0
*
mCherry   hM3Dq