Combining an alarmin HMGN1 peptide with PD-L1 blockade facilitates stem-like CD8+ T cell expansion and results in robust antitumor effects

Chang-Yu Chen[1,2], Satoshi Ueha[1,2], Yoshiro Ishiwata[1,2], Shigeyuki Shichino[1,2], Shoji Yokochi[1,2], De Yang[3], Joost J. Oppenheim[3], Haru Ogiwara[1,2], Shungo Deshimaru[1,2], Yuzuka Kanno[1,4], Tatsuro Ogawa[1], Shiro Shibayama[5], Kouji Matsushima[1,2,*]

[1]Division of Molecular Regulation of Inflammatory and Immune Diseases, Research Institute for Biomedical Sciences, Tokyo University of Science, Chiba, Japan

[2]Department of Molecular Preventive Medicine, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

[3]Cancer and Inflammation Program, Center for Cancer Research, National Cancer Institute at Frederick, Frederick, MD, USA

[4]Department of Medicinal and Life Sciences, Faculty of Pharmaceutical Sciences, Tokyo University of Science, Chiba, Japan

[5]Research Center of Immunology, Tsukuba Institute, ONO Pharmaceutical Co., Ltd., Tsukuba, Japan

[*]Corresponding Author and Lead Contact: Kouji Matsushima, koujim@rs.tus.ac.jp; Tokyo University of Science, 2669 Yamazaki, Noda, Chiba 278-0022, Japan

Abstract

The expansion of intratumoral stem-like CD8$^+$ T (Tstem) cells provides a potential approach to improving the therapeutic efficacy of immune checkpoint blockade (ICB). Thus, here we demonstrate a strategy to facilitate Tstem cell expansion by combining an alarmin high-mobility group nucleosome binding domain 1 (HMGN1) peptide with programmed death-ligand 1 (PD-L1) blockade. The HMGN1 peptide synergizes with PD-L1 blockade in augmenting the number of mature DCs enriched in immunoregulatory molecules (mregDCs) in tumors, and enhancing their MHC class I antigen-presenting program, which is correlated with the expansion of intratumoral Tstem cells, specifically promoting the Tstem cells but restricting terminally exhausted CD8$^+$ T (Tex) cells, owing to the regulatory molecules (*Cd28*, *Cd274*, *Nectin1*, and *Pvr*) expressed on mregDCs. Taken together, our results indicate that HMGN1 peptide serves as an immunoadjuvant to promote effective anti-PD-L1 immunotherapy and implicate that mregDCs play a role beyond facilitating Tstem cell expansion.

Keywords

alarmin, high-mobility group nucleosome binding domain 1 (HMGN1), programmed death-ligand 1 (PD-L1), mature dendritic cell enriched in immunoregulatory molecules (mregDC), stem-like CD8$^+$ T (Tstem) cell, terminally exhausted CD8$^+$ T (Tex) cell

Introduction

Blocking programmed cell death 1 and its ligand programmed death ligand 1 (PD-1/PD-L1) can increase survival in cancer patients by reinvigorating the functions of tumor-reactive CD8[+] T cells [1, 2, 3, 4, 5]; however, not all patients respond equally well to PD-1/PD-L1 blockade. The overall response rate to PD-1/PD-L1 blockade alone is below 30% in patients with melanoma, non-small-cell lung carcinoma, ovarian cancer, and renal-cell cancer patients [6, 7]. Low response rates to ICB therapies are associated with inadequate antigen presentation and insufficient generation of tumor-reactive CD8[+] T cells [8, 9]. To address these issues, researchers have attempted to combine various types of immune stimulators with PD-1/PD-L1 blockade to strengthen the antigen-presenting function of intratumoral dendritic cells (DCs) and accelerate the generation of tumor-reactive CD8[+] T cells, such as R848 (a TLR-7 agonist) [10], CpG oligodeoxynucleotide (ODN; a TLR-9 agonist) [11] or SD-101 (a novel class of CpG-ODN TLR9 agonist) [12]. Here, we explore the possibility of combining the immune stimulator called high-mobility nucleosome binding domain 1 (HMGN1) with PD-L1 blockade.

HMGN1 contains two functional domains, a nucleosome binding domain (NBD) and a chromatin unfolding domain (CHUD) [13, 14], and functions as both an intracellular and extracellular regulatory protein [15]. Intracellular HMGN1 is a transcriptional factor that supports higher-order chromatin folding and gene regulation [13, 14]. Extracellular HMGN1 acts as an alarmin, which is released from apoptotic or necrotic cells, to promote DC activation via MYD88 innate immune signal transduction adaptor (MYD88)- and toll-like receptor adaptor molecule 1 (TRIF)-dependent signaling pathways [16, 17, 18, 19, 20]. Currently, recombinant HMGN1 is considered to act as a DC adjuvant for T cell-mediated cancer therapy due to its benefits of promoting CD8[+] T cell expansion [17, 18, 19, 21]. As reported previously, an *in vitro* assay demonstrated that recombinant HMGN1 can promote DC-dependent CD8[+] T cell

expansion in co-culturing Pmel-1 CD8[+] T cells with gp100-pulsed BMDCs [21]. Moreover, an *in vivo* study also demonstrated that HMGN1 synergizes anti-CD4 depleting antibody treatment in enhancing the expansion of intratumoral CD8[+] T cells in mice [21]. Thus, we speculate that combining HMGN1 with PD-L1 blockade may facilitate intratumoral CD8[+] T cell expansion.

Intratumoral CD8[+] T cell expansion is a promising strategy to improve the therapeutic efficacy of ICB, particularly if it results in the expansion of a newly defined intratumoral CD8[+] T cell subset with stem-like properties, which acts as resource cells to mediate tumor control in response to ICB therapy [22, 23, 24]. Stem-like CD8[+] T (Tstem) cells [23, 25, 26], also known as progenitor-exhausted CD8[+] T (progenitor-Tex) cells [22, 27], precursor-exhausted CD8[+] T cells (TPEX) [24], and memory-like CD8[+] T cells [26, 28], which are characterized by high expression of SLAM family member 6 (SLAMF6 or also known as Ly108), and transcriptional factor TCF7 (also known as TCF1); the intermediate expression of PD-1; low expression of T-cell immunoglobulin and mucin domain 3 (TIM-3; also known as hepatitis A virus cellular receptor 2, HAVCR2) and lymphocyte-activation gene 3 (LAG-3); and high proliferative capacity [22, 23, 27, 29, 30]. Patients with melanoma who had increased numbers of Tstem cells displayed durable tumor regression and longer progression-free survival after ICB therapy [22, 31]. Conversely, a lack of adequate Tstem cells or an abundance of exhausted CD8[+] T (Tex) cells restricts the therapeutic efficacy of ICB therapy, and limits the survival benefits to both tumor-bearing mice and patients with cancer [31].

Here we demonstrate a new strategy to facilitate Tstem cell expansion by combining a HMGN1-derived immunostimulatory peptide with PD-L1 blockade. The HMGN1 peptide synergizes with PD-L1 blockade to increase the number of and the antigen-presenting ability of tumor-infiltrating mregDCs, which is correlated with the expansion of Tstem cells in tumors. Using single-cell RNA sequencing, we predict the cell-cell interaction among mregDC, Tstem cell, and Tex cell populations by CellPhoneDB. The regulatory molecules on mregDCs may

support Tstem cell but not Tex cell activation and expansion. Here our findings demonstrate that HMGN1 peptide serves as an immunoadjuvant to promote effective anti-PD-L1 immunotherapy and implicate that mregDCs play a role beyond facilitating Tstem cell expansion.

Results

Combined treatment with HMGN1 and PD-L1 blockade induces durable tumor regression

To determine the optimal dosage of murine HMGN1 (mH) in combination with an anti-PD-L1 antibody (αPDL1; clone 10F.9G2), we first performed a dose-response assessment of combined treatment in B16F10 or Colon26 tumor-bearing mice model. After observing the best antitumor effect at 0.08 µg mH in the B16F10 model, and at a dose range of 0.08-0.4 µg mH in the Colon26 model, followed by 200 µg αPDL1 treatment (Fig. 1A, B), we therefore, used the optimized dose of 0.08 µg mH and 200 µg αPDL1 in the following experiments. In this setting, we next evaluated the anti-tumor effects of mH/αPDL1 treatment in Colon26, Lewis lung carcinoma (LLC), EO771, and B16F10 models. Compared with αPDL1 treatment alone, combined treatment of mH/αPDL1 showed significant improvement in tumor growth inhibition in all four different tumor models: Colon26 (on day 17, $p < 0.001$; day 24, $p < 0.01$), LLC (on day 18, $p < 0.01$; day 24, $p < 0.01$), EO771 (on day 14, $p < 0.01$) and B16F10 (on day 14, $p < 0.05$) (Fig. 1A, C).

Moreover, we monitored the long-term outcomes and patterns of tumor recurrence after treatments. 70% of Colon26 (seven out of ten), 30% of LLC (three out of ten), and 70% of EO771 (seven out of ten) tumor-bearing mice curing from primary tumors after mH/αPDL1 treatment were also protected from tumor re-challenge (Fig. 1D). These results demonstrated that combining HMGN1 with PD-L1 blockade induces durable tumor regression.

minP1: a minimized immunostimulatory peptide derived from the HMGN1 NBD retains its anti-tumor effects

As clinical agents, synthetic peptides have advantages over recombinant proteins, as they are quickly and easily synthesized without the risk of endotoxin contamination, and are relatively

inexpensive; therefore, we tried to identify the immunostimulatory domain on HMGN1 and used the synthesized HMGN1-derived immunostimulatory domain as a therapeutic peptide in combination with αPDL1 treatment.

In order to determine the immunostimulatory domain on HMGN1, we first synthesized the murine HMGN1 NBD peptide (P1) and CHUD peptide (P2) and evaluated the anti-tumor effects of P1/αPDL1 and P2/αPDL1 treatments in Colon26 model, respectively (Fig.2A). According to the tumor growth assay, P1/αPDL1 treatment, but not P2/αPDL1 treatment, showed synergistic anti-tumor effects equivalent to mH/αPDL1 treatment (Fig. 2B), suggesting that HMGN1 immunostimulatory domain remains within NBD.

Next, in order to minimize the region of immunostimulatory domain within HMGN1 NBD, we further selected human HMGN1 (which has equal anti-tumor efficacy to mH [21]) and synthesized a human HMGN1 NBD peptide (hP1) and six hP1-derived N-terminally or C-terminally truncated peptides (ΔN1, ΔN2, ΔN3, ΔC1, ΔC2, ΔC3; Fig. 2C). According to the tumor growth assay, the ΔC2, ΔC3, and ΔN3 peptides failed to display synergistic anti-tumor effects in combination with αPDL1 treatment, indicating that the C-terminal region from K_{31} to K_{38} and the N-terminal region from E_{16} to R_{20} are required for function. Thus, we hypothesized that this minimal peptide (minP1, spanning 23-amino acid residues from E_{16} to K_{38}, EPKRR SARLS AKPPA KVEAK PKK) retains HMGN1-induced synergistic anti-tumor effects (Fig. 2D).

As expected, minP1 showed equal anti-tumor efficacy as to mH. Combined treatment with minP1 and αPDL1 showed significant improvement in tumor growth inhibition compared with αPDL1 treatment in both Colon26 and LLC models (Fig. 2E). After monitoring the long-term treatment outcomes, we observed 80% of Colon26 (eight out of ten) and 60% of LLC (six out of ten) tumor-bearing mice achieved complete cure from tumors, which showed similar anti-tumor efficacy as to mH. Furthermore, in tumor re-challenging assay, we found that the cured

mice resisted primary tumor re-challenge and did not develop secondary tumors (Fig. 2F),

consistent with the tumor re-challenge results of mH/αPDL1 treatment (Fig. 1D). Taken

together, these data demonstrated that minP1 retains the HMGN1 immunostimulatory function

and shows the equal anti-tumor effects to full-length HMGN1 protein.

minP1 preferentially binds on intratumoral DC populations

In order to further understand how minP1 regulates anti-tumor immunity in tumor-bearing mice,

we first tried to identify the primary target population of minP1 therapy in the tumor, secondary

lymphoid organ, and primary hematopoietic organ. To address our question, we prepared a

fluorescein isothiocyanate (FITC)-conjugated minP1 (FITC-minP1) and took advantage of

FITC-minP1 to label the cell populations that would be attached by minP1. 9 days after tumor

inoculation, total cell suspensions from the tumor (Tu), draining lymph node (dLN), spleen

(SP), and bone marrow (BM) of LLC tumor-bearing mice were pre-stained with an antibody–

fluorophore combination and incubated with FITC-minP1. Distinct immune cell populations

from Tu, dLN, SP, BM were characterized first (SFig. 1A), and then clustered using t-

distributed stochastic neighbor embedding (t-SNE) analysis (SFig. 1B). By visualizing the

result of t-SNE analysis, higher FITC-minP1 intensities were detected on monocyte,

macrophage, and DC clusters in the Tu, but not detected on any cell clusters in either dLN, SP,

or BM (SFig. 1B), which suggests that minP1 may preferentially bind to intratumoral immune

cell populations, including monocyte, macrophage, and DC populations.

To confirm the result of t-SNE analysis, we evaluated the FITC-minP1 binding affinity to

intratumoral immune cell populations by competitive protein binding assay. Using an

unlabeled-minP1 as competitive protein, we found the FITC-minP1 intensities were only

decreased on monocyte, macrophage, and DC clusters, but not CD4$^+$ T, CD8$^+$ T, or NK cell

clusters (SFig. 1C). Further in dose-dependent competitive protein binding assay, as the

concentration of unlabeled minP1(competitive protein) increased, the amounts of FITC-minP1

bound to macrophage and DC clusters decreased, but the decreasing amounts of FITC-minP1

was not observed on monocyte cluster (SFig. 1D). Moreover, the half-maximal inhibitory

concentration (IC_{50}) values of DC (69.96 μg/mL) cluster was lower than those for macrophages

(192.98 μg/mL), monocyte (>200 μg/mL), and other immune cell populations (>200 μg/mL)

in the tumor (SFig. 1D). Taken together, these results demonstrated that minP1 preferentially

binds on intratumoral DC populations and may further regulates their biological functions.

minP1 transcriptionally up-regulates MHC class I antigen presentation program on

intratumoral mregDC population

To examine the role in which how minP1 shapes anti-tumor immunity in tumor

microenvironment and how minP1 regulates the biological functions of intratumoral DC

populations, we analyzed the transcriptomic changes after minP1 treatment by using bulk RNA

sequencing (bulk RNA-seq) on whole tumor tissue and single-cell RNA sequencing (scRNA-

seq) on CD45[+] immune cell populations, respectively (Fig. 3A).

In our bulk RNA-seq data, we found a total of 1,023 differentially expressed genes (DEGs,

with adjusted *p* values < 0.05 and a fold-changes of ≥ 2 among groups) identified between

αPDL1 and minP1/αPDL1 treatment groups. Gene ontology analysis yielded a large cluster

(cluster #1) containing 546 upregulated genes involved in the positive regulation of innate

immune responses and immune effector processes, which included genes involved in antigen

processing and presentation, phagocytosis, the C-type lectin receptor signaling pathway, the

TNF signaling pathway, T cell activation, and the toll-like signaling pathway (Fig. 3B).

Interestingly, many upregulated genes in the minP1/αPDL1 treatment group, such as

histocompatibility 2, class II, locus Mb1 (*H2-DMb1*), histocompatibility 2, T region locus 22

(*H2-T22*), histocompatibility 2, K1, K region (*H2-K1*), beta-2 microglobulin (*B2m*), transporter

1, ATP-binding cassette, sub-family B (MDR/TAP; *Tap1*), calreticulin (*Calr*), cathepsin L

(*Ctsl*), legumain (*Lgmn*), proteasome activator subunit 2 (PA28 beta; *Psme2*), and *Psme2b*, are

involved in antigen processing and presentation. Moreover, upregulated genes after

combination treatment including tumor necrosis factor (*Tnf*), lysosomal-associated membrane

protein 1 (*Lamp1*), *Cd44* molecule, *Cd69* molecule, TNF receptor superfamily, member 9

(*Tnfrsf9*), and IFN gamma receptor 1 (*Ifngr1*), are involved in regulation of T cell activation

(Fig. 3C). Conversely, some downregulated genes after combination treatment including

cyclin-dependent kinase 4 (*Cdk4*), proliferation-associated 2G4 (*Pa2g4*), and ribosomal

proteins 16, 21, and S24 (*Rpl16*, *Rpl21*, *Rps24*), are involved in immunosuppression and T cell

exclusion [32] (Fig. 3C). Our bulk RNA sequencing data demonstrated that minP1 shapes anti-

tumor immunity by enhancing antigen presentation and T cell activation programs after αPDL1

treatment in tumor microenvironment.

Furthermore, in our scRNA-seq data, a total of 27 unsupervised cell clusters from tumors

were profiled and characterized by unique gene signatures of each cell populations (SFig. 2).

In order to inquire deep insight about how minP1 regulates the biological functions on

intratumoral DC populations, we selected DC clusters for the following analysis. A total of six

DC clusters were profiled and characterized by their unique gene signatures, including

conventional DC type1 (cDC1; *Xcr1*, *Clec9a*, and *Naaa*), conventional DC type 2 and its

precursor (cDC2 and pre-cDC2; *Cd209a*, *Clec10a*, and *Itgam*), plasmacytoid dendritic cells

(pDC_1 and pDC_2; *Ccr9*, *Il12a*, and *Havcr1*), and mregDC (*Fscn1*, *Il12b*, and *Ccl22*) (Fig.

3E, F). Among those DC clusters, minP1 treatment resulted in a significant increase proportion

of mregDC cluster (relative to the untreated group; Fig. 3G), and also enhanced the expression

level of MHC class I antigen presentation-related genes: *H2-K1*, *Calr,* and *Tap1* on mregDC

cluster (Fig. 3H, I). Unexpectedly, we did not observe the percentage change or expression

change in cDC1 or cDC2 cluster after minP1 treatment (Fig. 3H, I). Taken together, our

scRNA-seq data further suggested that minP1 treatment results in an increasing proportion of

mregDCs in tumor microenvironment, and also transcriptionally up-regulates MHC class I

antigen presentation program on these mregDCs.

minP1 increases the number of intratumoral mregDCs and enhances their MHC class I

expression

To confirm our scRNA-seq result and assess whether the proportion and number of mregDCs

increased after minP1 treatments, we prepared total cell suspensions from the tumor tissues of

Colon26 tumor-bearing mice on day 12 (after two rounds of minP1 treatments) and 16 (after

three times of minP1 treatments) after tumor inoculation. Using flow cytometry, we identified

a $CCR7^+$ $CD11b^+$ $CD11c^+$ $CD14^-$ $Ly6C^-$ MHC class II^+ cell population as $CCR7^+$ mregDC

from a lineage-negative cell population ($CD3^-$, $CD19^-$, $Ly6G^-$; Fig. 4A). Increasing numbers

of intratumoral $CCR7^+$ mregDCs were observed in Colon26 tumor-bearing mice treated with

minP1 or minP1/αPDL1 on days 12 and 16, compared to untreated or αPDL1 treatment

respectively ($p < 0.05$; Fig. 4B, C). In minP1/αPDL1 treatment group, increased $CCR7^+$

mregDCs exhibited higher MHC class I and lower PD-L1 expression compared to those from

the untreated or αPDL1 treatment group respectively ($p < 0.05$); however, their MHC class II

expression did not exhibit any difference (Fig. 4D), which is consistent with our scRNA-seq

results and suggests that minP1 might enhance the MHC class I antigen-presenting ability of

intratumoral $CCR7^+$ mregDCs. Moreover, consistent with the results in Colon26 tumor-bearing

mice, the increasing proportion and the number of intratumoral $IL-12b^+$ $CCR7^+$ mregDCs [33]

were also observed in LLC tumor-bearing IL-12b-yellow fluorescent protein (YFP) mice

treated with minP1 or minP1/αPDL1, compared to untreated or αPDL1 treatment respectively

($p < 0.05$; Fig. 4E-G).

To figure out where these increased intratumoral IL-12b$^+$ CCR7$^+$ mregDCs locate, we

investigated the invasive margins and the central tumors of LLC tumor-bearing IL12-YFP mice.

Using immunofluorescence staining, we identified the IL-12b-YFP (green)$^+$ and MHC class II

(red)$^+$ cell population as mregDC in the tumor. Increasing numbers of IL12b$^+$ CCR7$^+$ mregDCs

were observed in both the invasive margins and the central tumors of minP1- and

minP1/αPDL1-treated mice (Fig. 4H). Taken together, these results indicated that minP1

enhances the MHC class I antigen-presenting ability of mregDCs and increases the number of

mregDCs in both invasive margin and central tumor.

Expansion of intratumoral stem-like CD8$^+$ T cells by minP1/αPD-L1 treatment is

associated with the increased number of mregDCs in tumors

The results described earlier with our bulk RNA-seq suggested that minP1/αPDL1 treatment

synergistically shapes anti-tumor immunity by promoting regulation of T cell activation

program in tumors (Fig. 3B). To confirm our bulk RNA-seq results, we prepared total cell

suspensions from the tumor tissues of Colon26 or LLC tumor-bearing mice on days 12 (after

twice minP1 treatments) and analyzed intratumoral CD8$^+$ T cells in two different type of tumors

by flow cytometry (Fig. 5A). The increasing numbers of intratumoral CD8$^+$ T cells were

observed in minP1/αPDL1 treatment group of both Colon26 and LLC models, compared with

αPDL1 treatment group (p < 0.0; Fig. 5B). We next asked whether this synergistic effect on

enhancing the expansion of intratumoral CD8$^+$ T cells by minP1/αPDL1 treatment was

associated with the increased number of mregDCs in tumors. We measured the numbers of

mregDCs and CD8$^+$ T cells in tumors and found a high correlation between the increasing

numbers of mregDCs and CD8$^+$ T cells in tumors of both Colon26 (R = 0.78, P = 0.03) and

LLC (R = 0.83, P = 0.03) models after minP1/αPDL1 treatment; however, we did not observe

this correlation in minP1 (Colon26: R = 0.62, P = 0.13; LLC: R = 0.42, P = 0.39) and αPDL1

(Colon26: R = -0.32, P = 0.69; LLC: R = -0.70, P = 0.11) treatment groups (Fig. 5C). These

results suggested that minP1/αPD-L1 treatment brings synergistic effect on enhancing the

expansion of intratumoral CD8$^+$ T cells, which is associated with the increased number of

mregDCs in tumors.

We next asked whether we could identify which CD8$^+$ T cell subsets benefited from this

expansion. First, we sought to profile Tstem and Tex cell subsets from intratumoral CD44^{high}

PD-1$^+$ CD3$^+$ CD8$^+$ T cells by their expression level of Ly108 and TIM-3 (Fig. 5D). Tstem and

Tex cells were characterized as Ly108$^+$ TIM-3$^-$ and Ly108$^-$ TIM-3$^+$, respectively. Tstem cells

exhibited high IL-2, intermediate TNF-α, and low IFN-γ compared with Tex cells (Fig. 5E),

consistent with the previous reports [22, 23, 27, 29, 30]. In minP1/αPDL1 treatment group, increasing

proportion and number of Tstem cells were observed in both Colon26 and LLC models

compared to αPDL1 treatment group (p < 0.05, Fig. 5F, G). Conversely, the propotion and

number of Tex cells decreased in both models after minP1/αPDL1 treatment (p < 0.05, relative

to αPDL1 treatment, Fig. 5F, H). Furthermore, we also checked the correlation between the

increasing numbers of Tstem cells and mregDCs in tumors. Surprisingly, there was a high

correlation between the increasing number of mregDCs and Tstem in tumors of Colon26 (R =

0.78, P = 0.03) and LLC (R = 0.86, P = 0.02) models after minP1/αPDL1 treatment, but no

correlation was observed between the increasing number of mregDCs and Tex (Colon26: R =

-0.71, P = 0.07; LLC: R = -0.74, P = 0.09) after minP1/αPDL1 treatment (Fig. 5I). Taken

together, these results suggest that an expansion of intratumoral Tstem cells by minP1/αPD-L1

treatment may require the support from the increased number of mregDCs in tumors.

The multiple regulatory molecules on mregDC that may restrict Tex cell but facilitate

Tstem cell activation and expansion

As previously reported, an increasing number of IL12[+]CCR7[+] mregDCs correlates with the activation and expansion of intratumoral CD8[+] T cells, owing to (i) the chemokine receptor CCR7 which enables mregDC to transport antigens from tumor to lymph nodes and then support T cell priming and expansion [34, 35, 36, 37], (ii) the cytokine IL-12 which enables mregDC to augment CD8[+] T cell activation via T cell-DC crosstalk beyond the cytokines IFN-γ and IL-12 [33], and (iii) immunoregulatory molecules (PD-L1, PD-L2, CD80, and so on) which enable mregDC control T cell activation [37].

However, the mechanisms how mregDCs modulate the activation and expansion of Tstem or Tex cell remain unclear. In order to understand the mechanisms between mregDC (Fig. 3E, F) and Tstem/Tex cells (SFig. 3), we here use CellPhoneDB[38] to predict their relevant interacting ligand-receptor partners from our scRNA-seq data (Fig. 6A, B). First of all, mregDCs within the tumors express high level of chemokines CCL5, CXCL9, and CXCL16 that may recruit different CD8[+] T cell subsets into tumor tissue and locally present tumor antigens to re-stimulate those recruited CD8[+] T cells. The expression pattern of the *CCL5-CCR5* and *CXCL9-CXCR3* ligand-receptor complex between mregDCs and Tstem cells, suggests mregDCs could mediate Tstem cell recruitment. Moreover, mregDCs might also have the ability to recruit Tex cells by other ligand-receptor complexes including *CCL5-CCR5* and *CXCL16-CXCR6* (Fig. 6C, D).

After recruiting Tstem and/or Tex cells, the anti-tumor activity of those T cells might further be regulated by mregDCs within tumors. mregDCs expressed higher immunoregulatory molecules such as *Cd80*, *Cd274*, *Nectin1*, *Pvr*, and *Tnfsf9* (known as CD137 ligand) compared to cDC1 and cDC2 (Fig. 6C). The corresponding receptors or ligands to those immunoregulatory molecules such as *Pdcd1*, *Cd96*, and *Tigit* were highly expressed on Tex cells but not Tstem cells. By contrast, Tstem cells expressed slightly higher *Cd28* and *Tnfrsf9* (known as CD137) (Fig. 6C). The *CD274-PDCD1*, *NECTIN1-CD96*, and *PVR-TIGIT* co-

inhibition between mregDC and Tex may restrict the activation and expansion of Tex cells (Fig.

6D), conversely, the *CD80-CD28* and *TNFSF9- TNFRSF9* co-stimulation between mregDC

and Tstem cells may facilitate the activation and expansion of Tstem (Fig. 6D). Taken together,

these results suggest that the multiple regulatory molecules on mregDC that may restrict Tex

cell but facilitate Tstem cell activation and expansion.

On account of mregDCs' possible ability to recruit and regulate both Tstem and Tex cell

functions, we further asked whether minP1/αPDL1 treatment could further augment these

relevant interacting ligand-receptor partners between mregDC and Tstem/Tex cells which are

shown by our CellPhoneDB analysis (Fig. 6B, C). We examined the expression of each

immunostimulatory molecules and its corresponding receptor or ligand on mregDC and

Tstem/Tex cell populations in different treatment groups. In comparison of untreated control

group, *Pvr* and *Cxcl9* expression were slightly upregulated on mregDCs, *Ccr5* expression were

upregulated on Tstem cells, and no clearly changed gene expressions could be found on Tex

cells in minP1/αPDL1 treatment group (SFig. 4). Those results showed that combination

treatment synergistically augmented *CCL5-CCR5*, *CXCL9-CXCR3*, and *PVR-TIGIT* signaling

pathways between mregDC and Tstem/Tex cells, suggests those three signaling pathways may

play critical roles in regulating the recruitment and activation of Tstem/Tex cell populations

after minP1/αPDL1 treatment.

Discussion

A newly defined Tstem cells act as a resource cell population mediating tumor control in response to ICB therapy [22, 31]. It is reported that Tstem cell mainly reside nearby high-dense antigen presenting cell (APC) niches within tumors, and require the support from APCs to maintain their function and differentiation [39]. For example, adequate number of DCs reinvigorates intratumoral CD8[+]T cells from exhaustion and support CD8[+]T cell expansion in tumors by *CD28-CD80* co-stimulation [29, 40]. In this study, we found that the minP1 synergizes with PD-L1 blockade in increasing the number of mregDCs and enhancing their antigen-presenting function. As recently reported, mregDCs play an immunoregulatory role inside tumors[37]. Our results indicate that mregDCs may restricts Tex but facilitates Tstem cell activation and expansion by multiple immunoregulatory molecules on mregDCs. In particular, Tstem cells with higher levels of *CD28* expression than Tex cells could be more sensitive to *CD28-CD80* co-stimulation [23]. By contrast, Tex cells with higher levels of *PDCD1*, *CD96*, and *TIGIT* could be restricted their functions by mregDCs, owing to *PDCD1-CD274*, *CD96-NECTIN1*, and *TIGIT-PVR* co-inhibition (Fig. 6C, D), which is consistent with the previous reports [37, 41]. Although we suggested the cell-cell interactions among mregDC, Tstem, and Tex cell populations by analyzing single-cell level, those cell-cell interactions are still unclear and are currently under investigation.

　　Our work highlights three important implications for both basic and clinical research. First, we have designed an effective strategy for combining the HMGN1 peptide (minP1) with PD-L1 blockade to improve the efficacy of ICB therapy. The use of peptides as therapeutics has advantages of standardized synthesis protocols, low toxicity, and good efficacy. Compared with the HMGN1, minP1 shows a lower risk of endotoxin contamination during synthesis, and could produce in larger quantities within a short time period, and has increasing potential to use as an adjuvant for cancer immunotherapies in the clinic.

Second, the transcriptome signature of tumor tissues after minP1/αPDL1 treatment could be used as an indicator to evaluate clinical response and help determine prognosis of cancer patients. The gene cluster including *H2-DMb1, H2-T22, H2-K1, B2m, Tap1, Calr, Ctsl, Lgmn, Psme2,* and *Psme2b* is involved in antigen processing and presentation, while the cluster including *Tnf*, *Lamp1*, *Cd44*, *Cd69*, *Tnfrsf9*, *Ifngr1*, is involved in regulation of T cell activation; and these genes were all upregulated after combination treatment. Conversely, a gene cluster including *Cdk4*, *Rpl16*, *Rpl21*, *Rps24*, and *Pa2g4* was involved in immunosuppression and T cell exclusion, and was down-regulated after combination treatment. These gene clusters could be used as indicators to predict and monitor the efficacy of combination treatment.

Third, we have identified the domain responsible for the immunostimulatory functions of HMGN1. Although previous studies identified the HMGN1 NBD and CHUD based on their distinct intracellular functions [13, 14], our previous work has further demonstrated that HMGN1 also has an extracellular function, inducing an immunostimulatory response that results in synergistic antitumor effects [16, 17, 18, 19, 20, 21]. Nevertheless, the area of the protein mediating the immunostimulatory function was unclear. Here we have found a 23-amino acid peptide within the HMGN1 NBD that retains the immunostimulatory responses and synergistic antitumor effects of HMGN1. This finding will enable in-depth examination of this immunostimulatory domain to understand how HMGN1 binds and interacts with its candidate receptors, such as lymphocyte antigen 96 (LY96; also known as MD2) [16] or Gαi protein coupled receptor (GiPCR) [20].

Overall, our study provides the rationale for combining an HMGN1 immunostimulatory peptide with PD-L1 blockade for cancer immunotherapy.

Figure legends

Figure 1. Combined treatment with HMGN1 and PD-L1 blockade induces durable tumor

regression.

(A) The optimized protocol for HMGN1/αPDL1 treatment in Colon26, LLC, EO771, and

B16F10 tumor-bearing mice. **(B)** Dose-response for assessing the tumor growth rate of B16F10

or Colon26 tumor-bearing mice treated with combination treatment, in the dose range of 0.016

to 2 μg murine HMGN1. **(C)** Tumor growth analysis during mH/αPDL1 treatment, and the

tumor volume of Colon26 (on days 17 and 24), LLC (on days 18 and 24), EO771 (on days 14

and 24), and B16F10 (on days 14 and 17). **(D)** Tumor re-challenge assay. One week after tumor

clearance, cured Colon26 tumor-bearing mice were re-challenged with 4T1 and Colon26 tumor

cells, cured LLC tumor-bearing mice were re-challenged with EO771 and LLC tumor cells,

and cured EO771 tumor-bearing mice were re-challenged with LLC and EO771 tumor cells,

respectively. Tumor growth is representative of three independent experiments with at least

eight mice per group. Data are presented as mean ± SEM. *, P < 0.05, **, P < 0.01, ***, P <

0.001 for a Dunnett's post hoc test (compared with control); #, P < 0.05, ##, P < 0.01; ###, P

< 0.001 in the figure indicate Student's *t*-test (comparing between mH/αPDL1-treated and

αPDL1-treated groups). Abbreviation: mH (murine HMGN1).

Figure 2. minP1: a minimized immunostimulatory peptide derived from the HMGN1

NBD retains its anti-tumor effects.

(**A**) The amino acid sequence of murine HMGN1 (mH) and its nucleosome binding domain

(NBD, peptide 1, P1) and chromatin unfolding domain (CHUD, peptide 2, P2). **(B)** The tumor

growth analysis on the P1 or P2 in combination with αPD-L1 treatment and the tumor volume

in each mouse from day 0 to day 24. **(C)** The amino acid sequence of the human HMGN1 (hH)

nucleosome binding domain (NBD) and derived peptides with various lengths. **(D)** Tumor

growth analysis after treatment with HMGN1 NBD-derived peptides in combination with

αPDL1 treatment and the tumor volume in each mouse. **(E)** Tumor growth analysis after

treatment with the minP1(human HMGN1 NBD-derived immunostimulatory peptide) in

combination with αPDL1 treatment in both Colon26 and LLC models, and the tumor volume

of each mouse from day 0 to day 24. **(F)** Tumor re-challenge assay. One week after tumor

clearance, cured Colon26 tumor-bearing mice were re-challenged with 4T1 and Colon26 tumor

cells and cured LLC tumor-bearing mice were re-challenged with EO771 and LLC tumor cells,

respectively. Tumor growth is representative of three independent experiments with at least

eight mice per group. Data are presented as mean ± SEM. *, $P < 0.05$, **, $P < 0.01$, ***, $P <$

0.001 for a Dunnett's post hoc test (compared with control); #, $P < 0.05$, ##, $P < 0.01$; ###, P

< 0.001 in the figure indicate Student's *t*-test (comparing between minP1/αPDL1-treated and

αPDL1-treated groups).

Figure 3. minP1 transcriptionally up-regulates MHC class I antigen presentation

program on intratumoral mregDC population.

(A) Schema on the preparation of tumor cell suspensions from Colon26 tumor-bearing mice

for bulk RNA-seq and scRNA-seq on day 10. **(B)** Gene ontology analysis of the differentially

expressed genes (DEGs). Each column represents a group, while each row represents an

individual gene. The Z-scores of module groups are shown at the bottom-right corner of the

heatmap. **(C)** Heatmap of the 1,023 DEGs. **(D)** Summary diagram of antigen processing and

presentation pathway components based on the Kyoto Encyclopedia of Genes and Genomes

(KEGG) pathway: map04612 (https://www.genome.jp/dbget-bin/www_bget?path:map04612).

(E, F) A t-distributed stochastic neighbor embedding (t-SNE) projection of scRNA-seq

profiling from 439 DCs in tumors of Colon26 tumor-bearing mice. DC clusters are distinct

colors. cDC1 (*Xcr1*, *Clec9a*, *Naaa*), cDC2 (*Cd209a*, *Clec10a*, *Itgam*), pDC (*Ccr9*, *Il12ra*,

Havcr1) and mregDC (*Fscn1*, *Il12b*, *Ccl22*) were profiled. Violin plots showing the expression

distribution of selected genes in different DC clusters. The y-axis represents log-normalized

expression value. (**G**) Proportion of DCs for cDC1, cDC2, and mregDC clusters in different

treatment groups. (**H**) Expression of antigen presentation (*H2-K1*, *Tap1*, *Calr*) and DC

immunostimulatory (*Cd80*, *Cd86*, *Cd274*) and cytokine and chemokine (*Ccr7*, *Il12b*, *Ccl5*,

Ccl22) genes among DC clusters in different treatment groups. Node size is proportional to the

expression frequency in a cell cluster. Node color max to min is proportional to the gene

enrichment score in each cluster; black represents control (Ct) group. Yellow represents minP1

treatment group. Blue represents PDL1 treatment group. Red represents PDL1+minP1

treatment group. (**I**) Violin plots showing the expression distribution of selected genes (*H2-K1*,

Calr, *Tap1*) in different treatment groups. The y-axis represents log-normalized expression

value.

Figure 4. minP1 increases the number of intratumoral mregDCs in both invasive margin

and central tumor and enhances their MHC class I expression.

(**A**) Schema on the preparation of tumor immune cells from Colon26 tumor-bearing BALB/c

mice. Flow cytometry gating of CCR7$^+$ mregDCs. (**B, C**) The frequency and number of CCR7$^+$

mregDCs in the tumors on days 12 and 16 after tumor inoculation. (**D**) The expression of MHC

class I, II, and PD-L1 on CCR7$^+$ mregDCs on day16. (**E**) Schema on the preparation of tumor

immune cells from LLC tumor-bearing IL12b-YFP reporter C57BL/6 mice. Flow cytometry

gating of IL-12b$^+$ CCR7$^+$ mregDCs. (**F, G**) Frequency and number of the IL-12b$^+$ CCR7$^+$

mregDCs in the tumor of LLC tumor-bearing IL12b-YFP reporter C57BL/6 mice treated with

combination treatment on day 14 after tumor inoculation. (**H**) The number and location of IL-

12b$^+$ CCR7$^+$ mregDCs in LLC tumor sections were identified by co-expression of IL-12b-YFP

(green) and MHC class II (red). Each result is representative of three independent experiments

with at least four mice per group. Data are presented as mean ± SEM. *, P < 0.05, **, P < 0.01,

***, P < 0.001 for a Dunnett's post hoc test (compared with control); #, P < 0.05, ##, P < 0.01;

###, P < 0.001 in the figure indicate Student's t-test (comparing with minP1/αPDL1-treated

and αPDL1-treated groups). Abbreviation: GMFI (geometric mean fluorescence), 7-AAD (7-

Amino-Actinomycin D).

Figure 5. Expansion of intratumoral stem-like CD8$^+$ T cells by minP1/αPD-L1 treatment

is associated with the increased number of mregDCs in tumors.

(**A**) Schema on the preparation of tumor immune cells from Colon26 or LLC tumor-bearing

mice. (**B**) The number of CD8$^+$ T cells in the tumors of Colon26 or LLC tumor-bearing mice

on day 12 after tumor inoculation. (**C**) Pearson correlation coefficients were calculated between

numbers of mregDCs and CD8$^+$ T cells in tumors. Two-sided P values are shown from a

Pearson correlation coefficient r. (**D**) Flow cytometry gating of intratumoral Tstem (Ly108$^+$

TIM-3$^-$ CD8$^+$) and Tex (Ly108$^-$ TIM-3$^+$ CD8$^+$) cells. The tSNE defines the Tstem and Tex

populations among CD8$^+$ T cells and displays the florescence intensity of CD44, PD-1, Ly108,

and TIM-3 in this population. (**E**) Expression of the intracellular cytokines IFN-γ, IL-2, and

TNF-α. (**F, G**) The compartments, frequencies, and numbers of Tstem (Ly108$^+$ TIM-3$^-$ CD8$^+$)

and Tex (Ly108$^-$ TIM-3$^+$ CD8$^+$) cells in the tumors of LLC or Colon26 tumor-bearing mice on

day 12 after tumor inoculation. (**H**) Pearson correlation coefficients were calculated between

numbers of mregDCs and either Tstem or Tex cells in tumors. Two-sided P values are shown

from a Pearson correlation coefficient r. Each result is representative of three independent

experiments with at least four mice per group. Data are presented as mean ± SEM. *, P < 0.05,

**, P < 0.01, ***, P < 0.001 for a Dunnett's post hoc test (compared with control); #, P < 0.05,

##, P < 0.01; ###, P < 0.001 in the figure indicate Student's t-test (comparing between

minP1/αPDL1-treated and αPDL1-treated groups). Abbreviation: CD45IVS: (intravenous

CD45 staining)

Figure 6. The multiple regulatory molecules on mregDC that may restrict Tex cell but

facilitate Tstem cell activation and expansion.

(A) A t-SNE projection of scRNA-seq profiling from 762 T cells and DCs in tumors of Colon26

tumor-bearing mice. T cell and DC clusters were represented as distinct colors. **(B)** Overview

of selected ligand–receptor interactions; P values were indicated by circle size, scale on right

(permutation test). The means of the average expression level of interacting molecule 1 in cell

cluster 1 and interacting molecule 2 in cell cluster 2 were indicated by color. Assays were

carried out at the mRNA level, but were extrapolated to protein interactions. **(C)** Violin plots

exhibited the expression distribution of selected genes in T cell and DC clusters. The y-axis

represented log-normalized expression value. **(D)** Summary diagram of the main receptors and

ligands expressed on the mregDC, Tstem, and Tex cell clusters that were involved in cellular

recruitment and co-stimulation/co-inhibition. Created with BioRender.com.

Methods

Mice

Seven-week-old female BALB/c, C57BL/6 were purchased from Japan SLC, Inc. The IL-12-

YFP reporter C57BL/6 mice (B6.129-Il12b$^{tm1.1Lky}$/J) were purchased from the Jackson

Laboratory (ME, USA). For tumor growth experiment, each group contained 8 mice except

where otherwise specified. All animal experiments were conducted in accordance with

institutional guidelines with the approval of the Animal Care and Use Committee of The

University of Tokyo and Tokyo University of Science.

Cell lines and tumor models

Murine tumor cell line Colon26 was obtained from the Cell Resource Center for Biomedical

Research (RRID: CVCL_0240; Cell Banker, RIKEN BRC, Japan). Lewis lung carcinoma

(LLC) was provided from Dr. F. Abe (RRID: CVCL_4358; Nipponkayaku, Tokyo, Japan).

B16F10 was obtained from the American Type Culture Collection (ATCC; RRID:

CVCL_0159; ATCC, USA). EO771 was obtained from CH3 BioSystems (CVCL_GR23).

Colon26 (2 x 10^5 cells), LLC (2 x 10^5 cells) and B16F10 (5 x 10^5) were subcutaneously

inoculated into the right flank of mice. EO771 (2 x 10^5 cells) were implanted into the inguinal

mammary fat pad. Tumor diameter was measured by twice weekly and used to calculate tumor

volume (V, mm^3) by using the formula V= L × W × W/ 2 (where L is tumor length and W is

tumor width).

In vivo treatment

Intraperitoneal injections of αPDL1 (200 μg/injection on days 4, 8, 14, and 18; clone 10F.9G2;

BioXcell), and HMGN1 and its-derived peptdies (0.08 μg on days 9, 14, 17, and 20) were given

to tumor-bearing mice based on the experiment design. Recombinant HMGN1 proteins were

produced in *E. coli* and purified using sequential fractionation by heparin affinity column, ion

exchange column and reverse-phase column as previously described [21]. The purity of HMGN1

(> 99%) was confirmed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-

PAGE). The endotoxin level of HMGN1 (< 1 endotoxin unit per mg) was assessed by

Endospecy ES-50M Kit (Seikagaku Corporation, Japan). The protein sequence of HMGN1

was identified by Applied Biosystems Procise 492 HT (Thermo Fisher Scientific, USA). The

HMGN1 peptides were synthesized by Eurofins Genomics, Japan.

Flow cytometry

Three minutes before collecting tissues, intravascular leukocytes were stained by intravenous

injection of fluorescein isothiocyanate (FITC)-conjugated antibody (3 μg/mouse) against

CD45 [42]. Total cell suspensions were prepared by enzymatic and mechanical dissociation of

tissues, as described previously [43]. Flow-Count Fluorospheres (Beckman Coulter, USA) were

used to determine cell numbers. Cells were pretreated with Fc blocking reagent (anti-mouse

CD16/CD32 monoclonal antibody, clone 2.4G2; BioXcell), then stained with a mixture of

fluorophore-conjugated anti-mouse antibodies. Data were acquired on a CytoFlex flow

cytometer (Beckman Coulter, USA) and analyzed by using FlowJo 10.5 software (FlowJo,

LLC, USA). Nonviable cells were excluded from the analysis based on forward and side scatter

profiles, and dead cells were excluded by propidium iodide (PI) staining. For intracellular

cytokine detection, enriched tumor-infiltrating $CD8^+$ T cells were re-stimulated with 1 μg/ml

ionomycin (IM) and 25 ng/ml phorbol myristate acetate (PMA) in the presence of GolgiPlug

(BD Biosciences, USA) for 4 hours at 37 ℃. The re-stimulated $CD8^+$ T cells were stained with

surface antigens, and these cells were stained for intracellular cytokines using a

Cytofix/Cytoperm kit (BD Biosciences, USA), according to the manufacturer's instructions.

For the transcriptome analysis, CD8$^+$ T cells from the tumor were sorted on FACSAria III Cell

Sorter (BD Biosciences, USA).

Classification of immune cell population by t-SNE

Multicolor flow cytometric data was visualized by T-distributed stochastic neighbor

embedding (t-SNE) analysis [44, 45, 46], which was used to classify different immune cell

populations in various organs in tumor-bearing mice. On 9 day after tumor inoculation, the

total cell suspensions from tumor (Tu), draining lymph node (dLN), spleen (SP), and bone

marrow (BM) were first stained with a mixture of 12 fluorophores, then incubated with FITC-

conjugated minP1 (FITC-minP1). Using FlowJo software, live cells were gated on 7-AAD

negative population and minimized to 3×10^4 cells per sample. The triplicate samples were

concatenated into one group and each group were underwent by t-SNE analysis. To identify

immune cell populations in the tumor, dLN, and SP, ten parameters (B220, CD3, CD4, CD8,

CD11b, CD11c, CD19, I-A/I-E, NK1.1, and Ly6C) were used to separate B cell, CD4$^+$ T cell,

CD8$^+$ T cell, NK cell, NKT cell, monocyte, macrophage/monocyte (monocyte-to-macrophage

transition), and DC populations. To identify immune system precursor cells in the BM, eleven

parameters (B220, CD11b, CD24, CD115, CD117, CD135, Ly6C, Ly6G, CX3CR1, Sca-1)

were used to separate cell populations, including B220$^+$ B cell progenitor, Ly6G$^+$ granulocyte

progenitor, monocyte progenitor (MP), monocyte-dendritic cell progenitor (MDP), common

DC progenitor (CDP), Ly6C$^+$ monocyte, and Ly6C$^-$ monocyte.

Bulk-RNA sequencing library preparation and analysis

Total RNA was extracted from 30,000 total cells of each tumor tissue, and stored in

lysis/binding buffer (Thermo Fisher Scientific, USA). The messenger RNA (mRNA) was

isolated from total RNA by Dynabeads M-270 streptavidin with biotin-oligo (dT)25 (Thermo

Fisher Scientific, USA). First- and second-strand cDNA synthesis was performed by Superscript reverse transcriptase IV (Thermo Fisher Scientific, USA) and Kapa HiFi DNA polymerase (Roche, Switzerland). While capturing on the beads, cDNA was digested with fragmentase (NEW ENGLAND BioLabs, USA). Each cDNA fragment was ligated to an adapter carrying Ion-Barcode-common sequence 1 (CS1). Finally, an 8-cycle PCR step was performed to enrich for the desired cDNA library molecules, and those cDNA library products were purified by size selection using AMPure XP beads (Beckman Coulter, USA), and were confirmed by Agilent High Sensitivity DNA kit (Agilent Technologies, USA).

The whole transcripts were amplified from total cells of tumor tissue from Colon26-bearing mice and were used to generate the 5'end Serial Analysis of Gene Expression (SAGE)-sequencing libraries. The sequencing was performed by using an Ion 540 Chef kit, an Ion 540 Chip kit, and an Ion S5 Sequencer (Thermo Fisher Scientific) according to the manufacturer's instructions except the input library concentration was 65 pM. Adapter trimming and quality filtering of sequencing data were performed by using Cutadpat-v2.4 [47], Trimommatic-v0.36 [48]. The filtered reads were mapped on Refseq mm10 using Bowtie2-2.2.5 (parameters: -t -p 11 -N 1 -D 200 -R 20 -L 20 -i S,1,0.50 --nofw). The mapped reads per gene (raw tag counts) were be quantified as gene expression. Between-sample normalization of gene expression was performed against raw count data by using R 3.6.1 (https://cran.r-project.org/) with DESeq2 [49] and pheatmap [50] packages. Genes with adjusted p value less than 0.05 and a fold-change of $\geq$ 2 between at least three samples were identified as statistically significant differentially expressed genes (DEGs). Raw data from the experiment have been deposited in the NCBI Gene Expression Omnibus (GEO, http://www.ncbi.nlm.nih.gov/geo); accession GSE139291.

Functional analysis of DEGs was performed by using Cytoscape 3.7.1 with ClueGO plugin (v2.5.4) [51, 52]. Significantly enriched Gene Ontology (GO) terms [53] (GO-immune system process, GO levels: 3–8, version: 2019/02/27) and Kyoto Encyclopedia of Genes and Genomes

(KEGG, version: 2019/02/27) pathway terms [54] in DEGs were explored and grouped, and a

term network was constructed based on the overlap of their elements (kappa score = 0.4).

Leading terms within each group were defined as the most significantly enriched term in each

group. Terms not connected with any other term were excluded.

Single-cell RNA sequencing library preparation and analysis

For single-cell RNA sequencing (scRNA-seq), we sorted 30,000 CD45$^+$ cells from tumors by

AutoMACS (Miltenyi Biotec, USA). Sorted cells were stained with TotalSeq anti-CD45

antibodies (BioLegend, USA) that had been conjugated to oligonucleotide barcodes following

an optimized protocol similar to the Cellular Indexing of Transcriptomes and Epitopes by

Sequencing (CITE-seq) workflow. Then, labeled cells were encapsulated using the BD

Rhapsody™ (BD, USA) according to the manufacturer's instructions. For scRNA-seq, cDNA

libraries were prepared by an optimized process similar to Smart-seq2[55] using NEBNext Single

Cell RNA Library Prep Kit for Illumina (NEW ENGLAND BioLabs, USA), according to the

manufacturer's instructions. QC of cDNA and final libraries was performed by qPCR library

quantification assay (KAPA). Samples were sequenced on an Illumina Novaseq 6000 S2

(Illumina, USA) to a depth of 100 million reads per library. For mouse data, after library

demultiplexing, gene-expression libraries were aligned to the mm10 reference transcriptome

and count matrices were generated using the BD Rhapsody workflow, using the 'raw' matrix

output. Doublets were removed based on co-staining of distant sample-barcoding ("Hashtag")

antibodies. Data were log-transformed [log(TPM + 1)] for all downstream analyses, most of

which were performed using the R software package Seurat[56] (http://satijalab.org/seurat) to

analyze immune cell populations.

For each cell, three quality control metrics were calculated: (1) the total detected number of

genes; (2) the proportion of ribosome encoded transcripts; and (3) the proportion of

mitochondrially encoded transcripts. Cells were excluded from downstream analysis if fewer

than 200 genes were detected. There was an expression matrix of 7,613 cells by 22,833 genes.

Each gene expression measurement was normalized by total expression within the

corresponding cell and multiplied by a scaling factor of 1,000,000. Mean and dispersion values

were calculated were used for principal components analysis. Principal components were

determined to be significant ($P < 0.05$) using the jackstraw method and tSNE was performed

on these key principal components. Unsupervised clustering was performed using a shared

nearest neighbor modularity optimization-based algorithm, as described previously 57. In our

initial analysis of DC and T cell subsets (SFigure. 2), we observed contaminating

subpopulations and removed them from the dataset by proceeding with the re-clustering

analysis. Differential gene expression and signature enrichment analyses were performed by a

Wilcoxon rank-sum test. To perform an analysis of cell-cell interaction among DC and T cell

subsets, we analyzed our scRNA-seq data by CellPhoneDB [38] (www.CellPhoneDB.org). As

previously reported [58] , CellPhoneDB could calculate the means of the average expression

level of each receptor–ligand pair in each pairwise comparison between two cell types, and

provide a *P* value for each receptor–ligand pair. We then determined the order of interactions

that are highly enriched between cell types based on the number of significant pairs, and

manually selected biologically relevant ones. The means of the average expression level of

interacting molecule 1 in cell cluster 1 and interacting molecule 2 in cell cluster 2 were

indicated by color and *P* values were indicated by circle size, scale. Assays were carried out at

the mRNA level, but are extrapolated to protein interactions.

Immunofluorescent staining

Acetone-fixed, 8-μm tumor sections of LLC tumor-bearing IL12-YFP reporter C57BL/6 mice

on day 14 after tumor inoculation, were prestained with anti-mouse MHC class II (clone

M5/114.15.2, BioLegend) and incubated with PI. Sections were mounted with ProLong Gold

Antifade Mountant (Thermo Fisher Scientific, USA) and examined under a TCS SP5 confocal

microscope (Leica Microsystems, Wetzlar, Germany).

Quantification and statistical analysis

Data were analyzed using Prism 8.0 software (GraphPad Software, USA). For comparisons among groups in the *in vivo* study, we used one-way ANOVA with the Dunnett's test. For comparisons between the means of two variables, we used two-sided unpaired Student's *t*-test. For correlation, we used a two-sided Pearson correlation with coefficient r. All statistical analyses were presented as mean with SEM, and conducted with a significance level of $\alpha = 0.05$ ($P< 0.05$).

Acknowledgements

The authors would like to thank Shin Aoki, Shunichi Fujita, and Ai Yamashita (The University of Tokyo, Tokyo) for help with maintenance of laboratory and animal facility, Rina Matsukiyo and Kazushige Shiraishi (Tokyo University of Science, Chiba) for help with cell sorting, Junko Yasuda (Tokyo University of Science, Chiba) for help with RNA sequencing and analysis. C.Y.C. was supported by the University of Tokyo Fellowship. Research work in the laboratory of K.M. is funded by Grant-in-Aid for Scientific Research on Innovative Areas (Grant Number: 17929397).

Author contributions

Conceptualization, C.Y.Chen, S.Ueha, J.J.Openheim, S.Shibayama, and K.Matsushima; Methodology, C.Y.Chen, S.Ueha, S. Shichino, Y.Ishiwata, S.Yokochi, H.Ogiwara, S.Deshimaru, Y. Kanno, and T. Ogawa; Investigation, C.Y.Chen, S.Ueha, S. Shichino, Y.Ishiwata, De Yang, H.Ogiwara, S.Deshimaru, and K.Matsushima; Writing – Original Draft, C.Y.Chen; Review & Editing, C.Y.Chen, S.Ueha, J.J.Openheim, and K.Matsushima; Funding Acquisition, S.Ueha. S.Shibayama, and K.Matsushima; Supervision, S.Ueha, and K.Matsushima.

Declaration of interests

S.S. is a senior researcher at ONO Pharmaceutical Co., Ltd. K.M received research grant from ONO Pharmaceutical Co. The remain authors declare no competing financial interests

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary liked to this article.

1 Nature Research Reporting Summary

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Hamster Anti-Mouse CD3e, PE-CF594 conjugated, clone 145-2C11	BD biosciences	Cat#562286; RRID: AB_11153307
Rat Anti-Mouse CD4, Pacific Blue conjugated, clone RM4-4	BioLegend	Cat#116008; RRID: AB_11149680
Rat Anti-Mouse CD4, PerCP conjugated, clone RM4-5	BioLegend	Cat#100538; RRID: AB_ 893325
Rat Anti-Mouse CD8, Alexa Fluor 700 conjugated, clone 53-6.7	BioLegend	Cat#100730; RRID: AB_ 493703
Rat Anti-Mouse CD11b, BV605 conjugated, clone M1/70	BD biosciences	Cat#563015; RRID: AB_ 2737951
Hamster Anti-Mouse CD11c, APC conjugated, clone N418	BioLegend	Cat#117310; RRID: AB_ 313779
Rat Anti-Mouse CD19, PerCP conjugated, clone 6D5	BioLegend	Cat#115532; RRID: AB_ 2072926
Rat Anti-Mouse/Human CD44, APC/Cy7 conjugated, clone IM7	BioLegend	Cat#103028; RRID: AB_ 830785
Rat Anti-Mouse CD45, FITC conjugated, clone 30-F11	Tonbo Biosciences	Cat#35-0451; RRID: AB_2621689
Rat Anti-Mouse/Human CD45R (B220), PE/Cy7 conjugated, clone RA3-6B2	BioLegend	Cat#103222; RRID: AB_ 313005
Rat Anti-Mouse CD49b (PanNK), FITC conjugated, clone DX5	BD biosciences	Cat#553857; RRID: AB_ 395093
Mouse Anti-Rat CD90/mouse CD90.1 (Thy1.1), APC conjugated, clone OX-7	BioLegend	Cat#202526; RRID: AB_1595470
Rat Anti-Mouse CD115 (CSF-1R), PE conjugated, clone AFS98	BioLegend	Cat#135506; RRID: AB_1937253

Rat Anti-Mouse CD117 (c-kit), APC/Cy7 conjugated, clone 2B8	BioLegend	Cat#105826; RRID: AB_1626278
Rat Anti-Mouse CD135, biotin conjugated, clone A2F10	BioLegend	Cat#135308; RRID: AB_1953267
Rat Anti-Mouse CD197 (CCR7), biotin conjugated, clone 4B12	BioLegend	Cat#120104; RRID: AB_389232
Rat Anti-Mouse CD274 (PD-L1), PE/Cy7 conjugated, clone 10F.9G2	BioLegend	Cat#124314; RRID: AB_10643573
Rat Anti-Mouse CD279 (PD-1), PE/Cy7 conjugated, clone RMP1-30	BioLegend	Cat#109110; RRID: AB_572017
Rat Anti-Mouse CD326 (TIM-3), BV421 conjugated, clone RMT3-23	BioLegend	Cat#119723; RRID: AB_2616908
Rat Anti-Mouse Ly6A/E (Sca-1), Pacific Blue conjugated, clone D7	BioLegend	Cat#108120; RRID: AB_493273
Rat Anti-Mouse Ly6C, Alexa Fluor 700 conjugated, clone HK1.4	BioLegend	Cat#128024; RRID: AB_10643270
Rat Anti-Mouse Ly6C, APC/Cy7 conjugated, clone HK1.4	BioLegend	Cat#128026; RRID: AB_10640120
Rat Anti-Mouse Ly6G, APC conjugated, clone 1A8	BioLegend	Cat#127614; RRID: AB_2227348
Rat Anti-Mouse NK1.1, PE conjugated, clone PK136	BioLegend	Cat#108708; RRID: AB_313395
Rat Anti-Mouse I-A/I-E, BV786 conjugated, clone M5/114.15.2	BioLegend	Cat#107645; RRID: AB_2565977
Rat Anti-Mouse H-2Kd, APC conjugated, clone SF1-1.1	BioLegend	Cat#116620; RRID: AB_10645328
Rat Anti-Mouse Ter-119, PerCP conjugated, clone Ter119	BioLegend	Cat#116226; RRID: AB_893635

Rat Anti-Mouse CX3CR1, BV605 conjugated, clone SA011F11	BioLegend	Cat#149027; RRID: AB_2565937
Rat Anti-Mouse Ly108, APC conjugated, clone 330-AJ	BioLegend	Cat#134610; RRID: AB_2728155
Rat Anti-Mouse IL-2, PE conjugated, clone JES6-5H4	BioLegend	Cat#503808; RRID: AB_315302
Rat Anti-Mouse IFN-γ, APC/Cy7 conjugated, clone XMG1.2	BioLegend	Cat#505850; RRID: AB_2616698
Rat Anti-Mouse TNF-α, FITC conjugated, clone MP6-XT22	BioLegend	Cat#506304; RRID: AB_315425
Chemicals, Peptides, and Recombinant Proteins		
RPMI 1640	Nacalai tesque, Japan	#30264-85
DMEM	Nacalai tesque, Japan	#09891-25
CELLBANKER	Takara, Japan	#CB011
BD GolgiPlug™ Protein Transport Inhibitor (containing Brefeldin A)	BD Bioscience	#555029
Ionomycin	Sigma-Aldrich	#I-0634
Phorbol 12-myristate 13-acetate (PMA), PKC activator	Sigma-Aldrich	#P-8139
Mouse HMGN1 recombinant protein (mH; 1-96 a.a.): 1- MPKRKVSADGAAKAEPKRRS 21- ARLSAKPAPAKVDAKPKKAA 41- GKDKASDKKVQIKGKRGAKG 61- KQADVADQQTTELPAENGET 81- ENQSPASEEEKEAKSD	Chen et al. 2019	https://www.ncbi.nlm.nih.gov/pubmed/30696484

Mouse HMGN1 peptide 1 (P1; 7-43 a.a.): 7- SADGAAKAEPKRRS 21- ARLSAKPAPAKVDAKPKKAA 41- GKD	This paper; synthesized by Eurofins Genomics, Japan	
Mouse HMGN1 peptide 2 (P2; 50-89 a.a.): 50- VQIKGKRGAKG 61- KQADVADQQTTELPAENGET 81- ENQSPASEE	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 recombinant protein (hH; 1-100 a.a.): 1- MPKRKVSSAEGAAKEEPKRR 21- SARLSAKPPAKVEAKPKKAA 41- AKDKSSDKKVQTKGKRGAKG 61- KQAEVANQETKEDLPAENGE 81- TKTEESPASDEAGEKEAKSD	Chen et al. 2019	https://www.ncbi.nl m.nih.gov/pubmed/ 30696484
Human HMGN1 peptide 1 (hP1, 7-43 a.a.): 7- SSAEGAAKEEPKRR 21- SARLSAKPPAKVEAKPKKAA 41- AKD	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ∆C1 peptide (7-38 a.a.): 7- SSAEGAAKEEPKRR 21- SARLSAKPPAKVEAKPKK	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ∆C2 peptide (7-34 a.a.): 7- SSAEGAAKEEPKRR 21- SARLSAKPPAKVEA	This paper; synthesized by Eurofins Genomics, Japan	

Human HMGN1 ΔC3 peptide (7-30 a.a.): 7- SSAEGAAKEEPKRR 21- SARLSAKPPA	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ΔN1 peptide (12-43 a.a.): 12- AAKEEPKRR 21- SARLSAKPPAKVEAKPKKAA 41- AKD	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ΔN2 peptide (16-43 a.a.): 16- EPKRR 21- SARLSAKPPAKVEAKPKKAA 41- AKD	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ΔN3 peptide (21-43 a.a.): 21- SARLSAKPPAKVEAKPKKAA 41- AKD	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ΔN1ΔC3 peptide (12-30 a.a.): 12- AAKEEPKRR 21- SARLSAKPPA	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ΔN2ΔC3 peptide (16-30 a.a.): 16- EPKRR 21- SARLSAKPPA	This paper; synthesized by Eurofins Genomics, Japan	
Human HMGN1 ΔN3ΔC3 peptide (21-30 a.a.): 21- SARLSAKPPA	This paper; synthesized by Eurofins Genomics, Japan	

Human HMGN1 minimized peptide (minP1, 16-38 a.a.): 16- EPKRR 21- SARLSAKPPAKVEAKPKK	This paper; synthesized by Eurofins Genomics, Japan	
Deposited Data		
Transcriptome Bulk-RNA sequencing data	This paper	GEO: GSE139291
Single-cell RNA sequencing data	This paper	GEO:
Experimental Models: Cell Lines		
Mouse: Colon26	Cell Banker, RIKEN BRC, Japan	RRID: CVCL_0240
Mouse: Lewis lung carcinoma (LLC)	Dr. F. Abe, Nipponkayaku, Tokyo, Japan	RRID: CVCL_4358
Mouse: B16F10	ATCC	RRID: CVCL_0159
Mouse: EO771	CH3 BioSystems	RRID: CVCL_GR23
Experimental Models: Organisms/Strains		
Mouse: BALB/cCrSlc	SLC, Japan	RRID: MGI: 2161077
Mouse: C57BL/6NCr	SLC, Japan	RRID: MGI: 5657970
Mouse: B6.Cg-Thy1[a]/Cy Tg(TcraTcrb)8Rest/J	The Jackson Laboratory	RRID: IMSR_JAX: 005023
Mouse: B6.SJL-Ptprc[a] Pepc[b]/BoyJ	The Jackson Laboratory	RRID: IMSR_JAX: 002014
Mouse: B6.129-Il12b[tm1.1Lky]/J	The Jackson Laboratory	RRID: IMSR_JAX: 006412
Software and Algorithms		

FlowJo	FlowJo, LLC	https://www.flowjo.com/solutions/flowjo
Cutadpat-v2.4	Martin, 2011	https://cutadapt.readthedocs.io/en/stable/
Trimommatic-v0.36	Bolger et al., 2014	http://www.usadellab.org/cms/?page=trimmomatic
Bowtie2	Langmead and Salzberg, 2012	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
R 3.6.1	The R Project for Statistical Computing	https://www.r-project.org
DESeq2	Love et al., 2014	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
pheatmap	Kolde, 2015	https://www.rdocumentation.org/packages/pheatmap/versions/1.0.12/topics/pheatmap
Cytoscape v3.7.1 with ClueGO plugin (v2.5.4)	Bindea et al., 2009; Saito et al., 2012	https://cytoscape.org/index.html
GraphPad Prism version 6	GraphPad Software, San Diego California USA	https://www.graphpad.com
CellPhoneDB v2.0	Efremova M et al., 2020	https://www.cellphonedb.org

Seurat 2.3.4	Stuart T et al., 2019	https://satijalab.org/seurat/
Others		
Sequencer Ion S5	Thermo Fisher Scientific	#A27212, https://www.thermofisher.com/order/catalog/product/A27212?SID=srch-srp-A27212
Ion Chef	Thermo Fisher Scientific	#4484177, https://www.thermofisher.com/order/catalog/product/4484177?SID=srch-srp-4484177
Ion 540 Kit-Chef	Thermo Fisher Scientific	#A30011, https://www.thermofisher.com/order/catalog/product/A30011?SID=srch-srp-A30011
Ion 540 Chip Kit	Thermo Fisher Scientific	#A27766, https://www.thermofisher.com/order/catalog/product/A27766?SID=srch-srp-A27766

References

1. Iwai, Y. *et al.* Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 12293-12297 (2002).

2. Strome, S.E. *et al.* B7-H1 blockade augments adoptive T-cell immunotherapy for squamous cell carcinoma. *Cancer research* **63**, 6501-6505 (2003).

3. Blank, C., Gajewski, T.F. & Mackensen, A. Interaction of PD-L1 on tumor cells with PD-1 on tumor-specific T cells as a mechanism of immune evasion: implications for tumor immunotherapy. *Cancer Immunol Immunother* **54**, 307-314 (2005).

4. Sakuishi, K. *et al.* Targeting Tim-3 and PD-1 pathways to reverse T cell exhaustion and restore anti-tumor immunity. *J Exp Med* **207**, 2187-2194 (2010).

5. Zou, W., Wolchok, J.D. & Chen, L. PD-L1 (B7-H1) and PD-1 pathway blockade for cancer therapy: Mechanisms, response biomarkers, and combinations. *Sci Transl Med* **8**, 328rv324 (2016).

6. Brahmer, J.R. *et al.* Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *The New England journal of medicine* **366**, 2455-2465 (2012).

7. Topalian, S.L. *et al.* Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *The New England journal of medicine* **366**, 2443-2454 (2012).

8. Schumacher, T.N. & Schreiber, R.D. Neoantigens in cancer immunotherapy. *Science* **348**, 69-74 (2015).

9. Jenkins, R.W., Barbie, D.A. & Flaherty, K.T. Mechanisms of resistance to immune checkpoint inhibitors. *Br J Cancer* **118**, 9-16 (2018).

10. Nie, Y. *et al.* Development of a Curative Therapeutic Vaccine (TheraVac) for the Treatment of Large Established Tumors. *Sci Rep* **7**, 14186 (2017).

11. Salmon, H. *et al.* Expansion and Activation of CD103(+) Dendritic Cell Progenitors at the Tumor Site Enhances Tumor Responses to Therapeutic PD-L1 and BRAF Inhibition. *Immunity* **44**, 924-938 (2016).

12. Ribas, A. *et al.* SD-101 in Combination with Pembrolizumab in Advanced Melanoma: Results of a Phase Ib, Multicenter Study. *Cancer Discov* **8**, 1250-1257 (2018).

13. Bustin, M. Chromatin unfolding and activation by HMGN(*) chromosomal proteins. *Trends Biochem Sci* **26**, 431-437 (2001).

14. Bianchi, M.E. & Agresti, A. HMG proteins: dynamic players in gene regulation and differentiation. *Curr Opin Genet Dev* **15**, 496-506 (2005).

15. Hock, R., Furusawa, T., Ueda, T. & Bustin, M. HMG chromosomal proteins in development and disease. *Trends Cell Biol* **17**, 72-79 (2007).

16. Yang, D. *et al.* High-mobility group nucleosome-binding protein 1 acts as an alarmin and is critical for lipopolysaccharide-induced immune responses. *J Exp Med* **209**, 157-171 (2012).

17. Wei, F. *et al.* The Alarmin HMGN1 contributes to antitumor immunity and is a potent immunoadjuvant. *Cancer research* **74**, 5989-5998 (2014).

18. Yang, D., Bustin, M. & Oppenheim, J.J. Harnessing the alarmin HMGN1 for anticancer therapy. *Immunotherapy* **7**, 1129-1131 (2015).

19. Nie, Y., Yang, D. & Oppenheim, J.J. Alarmins and Antitumor Immunity. *Clin Ther* **38**, 1042-1053 (2016).

20. Yang, D., Han, Z., Alam, M.M. & Oppenheim, J.J. High-mobility group nucleosome binding domain 1 (HMGN1) functions as a Th1-polarizing alarmin. *Semin Immunol* **38**, 49-53 (2018).

21. Chen, C.Y. *et al.* Combined treatment with HMGN1 and anti-CD4 depleting antibody reverses T cell exhaustion and exerts robust anti-tumor effects in mice. *J Immunother Cancer* **7**, 21 (2019).

22. Miller, B.C. *et al.* Subsets of exhausted CD8(+) T cells differentially mediate tumor control and respond to checkpoint blockade. *Nat Immunol* **20**, 326-336 (2019).

23. Im, S.J. *et al.* Defining CD8+ T cells that provide the proliferative burst after PD-1 therapy. *Nature* **537**, 417-421 (2016).

24. Kallies, A., Zehn, D. & Utzschneider, D.T. Precursor exhausted T cells: key to successful immunotherapy? *Nature reviews. Immunology* (2019).

25. Vodnala, S.K. *et al.* T cell stemness and dysfunction in tumors are triggered by a common mechanism. *Science* **363** (2019).

26. Wu, T. *et al.* The TCF1-Bcl6 axis counteracts type I interferon to repress exhaustion and maintain T cell stemness. *Sci Immunol* **1** (2016).

27. Paley, M.A. *et al.* Progenitor and terminal subsets of CD8+ T cells cooperate to contain chronic viral infection. *Science* **338**, 1220-1225 (2012).

28. Snell, L.M. *et al.* CD8(+) T Cell Priming in Established Chronic Viral Infection Preferentially Directs Differentiation of Memory-like Cells for Sustained Immunity. *Immunity* **49**, 678-694 e675 (2018).

29. Hashimoto, M. *et al.* CD8 T Cell Exhaustion in Chronic Infection and Cancer: Opportunities for Interventions. *Annu Rev Med* **69**, 301-318 (2018).

30. McLane, L.M., Abdel-Hakeem, M.S. & Wherry, E.J. CD8 T Cell Exhaustion During Chronic Viral Infection and Cancer. *Annual review of immunology* **37**, 457-495 (2019).

31. Siddiqui, I. *et al.* Intratumoral Tcf1(+)PD-1(+)CD8(+) T Cells with Stem-like Properties Promote Tumor Control in Response to Vaccination and Checkpoint Blockade Immunotherapy. *Immunity* **50**, 195-211 e110 (2019).

32. Jerby-Arnon, L. *et al.* A Cancer Cell Program Promotes T Cell Exclusion and Resistance to Checkpoint Blockade. *Cell* **175**, 984-997 e924 (2018).

33. Garris, C.S. *et al.* Successful Anti-PD-1 Cancer Immunotherapy Requires T Cell-Dendritic Cell Crosstalk Involving the Cytokines IFN-gamma and IL-12. *Immunity* **49**, 1148-1161 e1147 (2018).

34. Hirao, M. *et al.* CC chemokine receptor-7 on dendritic cells is induced after interaction with apoptotic tumor cells: Critical role in migration from the tumor site to draining lymph nodes. *Cancer research* **60**, 2209-2217 (2000).

35. Ohl, L. *et al.* CCR7 governs skin dendritic cell migration under inflammatory and steady-state conditions. *Immunity* **21**, 279-288 (2004).

36. Roberts, E.W. *et al.* Critical Role for CD103(+)/CD141(+) Dendritic Cells Bearing CCR7 for Tumor Antigen Trafficking and Priming of T Cell Immunity in Melanoma. *Cancer Cell* **30**, 324-336 (2016).

37. Maier, B. *et al.* A conserved dendritic-cell regulatory program limits antitumour immunity. *Nature* **580**, 257-262 (2020).

38. Efremova, M., Vento-Tormo, M., Teichmann, S.A. & Vento-Tormo, R. CellPhoneDB: inferring cell-cell communication from combined expression of multi-subunit ligand-receptor complexes. *Nat Protoc* **15**, 1484-1506 (2020).

39. Jansen, C.S. *et al.* An intra-tumoral niche maintains and differentiates stem-like CD8 T cells. *Nature* **576**, 465-470 (2019).

40. Kamphorst, A.O. *et al.* Rescue of exhausted CD8 T cells by PD-1-targeted therapies is CD28-dependent. *Science* **355**, 1423-1427 (2017).

41. Oh, S.A. *et al.* PD-L1 expression by dendritic cells is a key regulator of T-cell immunity in cancer. *Nature Cancer* **1**, 681-691 (2020).

42. Anderson, K.G. *et al.* Intravascular staining for discrimination of vascular and tissue leukocytes. *Nat Protoc* **9**, 209-222 (2014).

43. Ueha, S. *et al.* Robust Antitumor Effects of Combined Anti-CD4-Depleting Antibody and Anti-PD-1/PD-L1 Immune Checkpoint Antibody Treatment in Mice. *Cancer Immunol Res* **3**, 631-640 (2015).

44. van der Maaten, L. Accelerating t-SNE using Tree-Based Algorithms. *J Mach Learn Res* **15**, 3221-3245 (2014).

45. van der Maaten, L. & Hinton, G. Visualizing Data using t-SNE. *J Mach Learn Res* **9**, 2579-2605 (2008).

46. Simoni, Y. *et al.* Bystander CD8(+) T cells are abundant and phenotypically distinct in human tumour infiltrates. *Nature* **557**, 575-579 (2018).

47. Martin, M. Cutadapt Removes Adapter Sequences From High-Throughput Sequencing Reads. *EMBnet.journal* **Vol. 17**, 10-12 (2011).

48. Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

49. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* **15**, 550 (2014).

50. Kolde, R. R: pheatmap: Pretty Heatmaps (2015).

51. Bindea, G. *et al.* ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* **25**, 1091-1093 (2009).

52. Saito, R. *et al.* A travel guide to Cytoscape plugins. *Nat Methods* **9**, 1069-1076 (2012).

53. Mi, H., Muruganujan, A., Casagrande, J.T. & Thomas, P.D. Large-scale gene function analysis with the PANTHER classification system. *Nat Protoc* **8**, 1551-1566 (2013).

54. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**, D353-D361 (2017).

55. Picelli, S. *et al.* Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nat Methods* **10**, 1096-1098 (2013).

56. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019).

57. Macosko, E.Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).

58. Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal-fetal interface in humans. *Nature* **563**, 347-353 (2018).

1
2

Figure 1. Combined treatment with HMGN1 and PD-L1 blockade induces durable tumor regression.

Figure 2. minP1: a minimized immunostimulatory peptide derived from the HMGN1 NBD retains its anti-tumor effects.

Figure 3. minP1 transcriptionally up-regulates MHC class I antigen presentation program on intratumoral mregDC population.

Figure 4. minP1 increases the number of intratumoral mregDCs and enhances their MHC class I expression.

Figure 5. Expansion of intratumoral stem-like CD8⁺ T cells by minP1/αPD-L1 treatment is associated with the increased number of mregDCs in tumors.

Figure 6. The multiple regulatory molecules on mregDC that may suppress Tex cell but facilitate Tstem cell activation and expansion.