Universality in biodiversity patterns: causes of the variation in species-temperature and species-productivity relationships

Eliška Bohdalková[1,2*], Anna Toszogyova[1], Irena Šímová[1,2] & David Storch[1,2]

[1] Center for Theoretical Study, Charles University and the Academy of Sciences of the Czech Republic, Prague, Czech Republic

[2] Department of Ecology, Faculty of Science, Charles University, Prague, Czech Republic

[*] eliska.bohdalkova@natur.cuni.cz

Abstract

Temperature and productivity appear as universal positive correlates of species richness. However, the strength and the shape of species-temperature (STR) and species-productivity (SPR) relationships vary widely, and the causes of this variation are poorly known. We analysed (1) published species richness data for multiple taxa sampled in various regions and (2) different clades within vertebrate classes globally, to test for the effects of spatial scale and characteristics of examined taxa and regions on the strength and direction of STRs and SPRs. Up to 55% of the variation in the strength and direction of STRs and SPRs can be explained by the variables tested, although there are striking differences in the variation of the relationships among types of data, between ectotherms and endotherms and also between STRs and SPRs. When focusing on multiple taxa sampled in different regions, only the variables concerning examined region affect the strength of the relationships: STRs are stronger in colder regions, while SPRs are stronger in warmer regions, and both relationships are strong within regions in which temperature and productivity are positively correlated. Additionally, STRs are stronger in regions with wider range of temperature. When focusing on vertebrate clades, there is additional effect of clade size: the relationships are stronger for larger clades. Moreover, clade size is the only variable consistently affecting the strength of SPRs for endotherms. The finding that both the relationships are weak in regions where

temperature and productivity are not correlated indicates that both temperature and productivity simultaneously affect spatial variation in species richness. However, species richness appears to be more universally driven by productivity than temperature, as SPRs are stronger, do not vary as much as STRs, and this variation is not tightly linked to the examined factors.

Keywords: diversity-productivity relationship, diversity-temperature relationship, macroecology, phylogenetic scale, spatial scale, species richness

Introduction

Deciphering large-scale diversity patterns is one of the main goals of ecology (Gaston 2000, Pontarp et al. 2019). It has been shown that species richness is tightly related to climate, namely to temperature, water availability and resulting ecosystem productivity (Wright 1983, Currie 1991, Hawkins et al. 2003a, Field et al. 2009), so that temperature and productivity appear as the main variables positively affecting species richness (Lennon et al. 2000, Kaspari et al. 2003, Currie et al. 2004, Evans et al. 2005, Šímová et al. 2011). However, the documented species-temperature (STR) and species-productivity (SPR) relationships vary in terms of their strength and shape. Weak or variable STRs or SPRs are sometimes interpreted as a counterevidence against the hypotheses explaining diversity patterns by the processes driven by temperature or productivity (Hawkins et al. 2007, Buckley et al. 2010). Recovering the causes of the variation of STRs and SPRs is thus a necessary precondition to understand the role of temperature and productivity in diversity patterns.

Three major classes of hypotheses attempt to explain large-scale diversity patterns (Pontarp et al. 2019): (1) hypotheses based on ecological limits, (2) hypotheses concerning diversification dynamics, and (3) hypotheses based on the history of different regions and time available for the accumulation of species. Productivity and/or temperature are comprised in most of these hypotheses. Productivity may determine the total amount of resources that can be divided among species to allow the persistence of their viable populations (the more-individuals hypothesis; Storch

et al. 2018) and higher productivity may also allow higher species' specialization promoting coexistence (Abrams 1995). Alternatively, productivity may promote larger population sizes, positively affecting mutation accumulation and easier speciation (Stevens et al. 2007). Temperature may affect mutation and consequently speciation rates as well (Rohde 1978, Allen et al. 2002) or it may constrain the number of species which can persist in a given environment due to their physiological temperature tolerances (Šímová et al. 2011). Alternatively, the effect of temperature may be indirect via affecting productivity (Šímová and Storch 2017). Higher species richness in warmer and more productive areas may also be a consequence of historically larger and more stable tropical regions (Wiens and Donoughue 2004). Importantly, synthetic macroecological theories that attempt to explain large-scale diversity patterns via a combination of several fundamental processes typically assume that productivity and temperature represent key drivers of these processes. Specifically, the neutral theory (Hubbell 2001) assumes that species richness within a region is determined by the product of total number of individuals, J, and speciation rate, v. It is reasonable to assume that J is a function of productivity (together with area) of the region, while v may be linked to temperature. These assumptions provide the basis of the theories which combine the neutral theory with the metabolic theory of ecology (Allen et al. 2007, Worm and Tittensor 2018), and are also in accord with the equilibrium theory of biodiversity dynamics (Storch et al. 2018, Storch and Okie 2019). Positive effects of productivity and temperature on species richness are thus essential features of major macroecological integrative theories.

On the other hand, there are good theoretical reasons to expect that neither STR nor SPR are universal, and that considerable variation exists around canonical, monotonically increasing relationships. Some of these reasons are statistical – smaller groups of species or small regions which do not comprise enough variation of temperature or productivity may reveal relatively weak and/or insignificant patterns even if there is an overall positive trend of increasing species richness with temperature or productivity. Other reasons may relate to biological specificities of the explored taxa – many taxa prefer particular environmental conditions (e.g. particular habitat types) which

may not be simply related to temperature or productivity, and many clades (especially young, and consequently smaller ones) would reveal diversity patterns which reflect the spreading from an area of origin rather than environmental effects (Gehrke and Linder 2011). Also, species richness patterns of some taxa may be affected by biotic interactions, namely competition with other taxa (McPeek 1996).

Even more importantly, the variation of the STR and the SPR may reflect the exact way how these variables affect species richness. For instance, if temperature affects species richness only indirectly, through its effect on productivity, then the SPR should be more universal (i.e. more regular and context-independent) than the STR, and the effect of temperature should diminish when accounting for productivity. Also, if temperature affects diversity via affecting mutation and speciation rates, the metabolic theory (Brown et al. 2004) predicts exponential STR with a predictable slope. However, this prediction holds only if total community abundance is more or less constant (Allen et al. 2007), while the STR is expected to vary if this condition is not fulfilled (Storch 2012). Since the total number of individuals is constrained by productivity (Storch et al. 2018), productivity variation may, according to the metabolic theory, also affect the slope and strength of the STR. More generally, if both productivity and temperature simultaneously affect species richness, both the STR and the SPR may depend on the variation of the other variable.

The variation of STRs and SPRs has been only rarely studied in a comprehensive way. Several studies addresses the effect of spatial scale and its two components – grain and extent – on biodiversity patterns including the STR and the SPR. Generally, with increasing *grain* size, the relationships are stronger (Rahbek and Graves 2000, 2001, González-Taboada et al. 2007, Belmaker and Jetz 2011), their slope is steeper (Hillebrand 2004), and the relative importance of temperature and productivity changes (Field et al. 2009, Belmaker and Jetz 2011). The effect of geographical *extent* is mostly studied using meta-analyses – and again, with increasing the extent of the analysis, the relationships appear stronger and steeper (Hillebrand 2004) and the importance of climate and productivity increases (Field et al. 2009). Some other studies investigated the relationships across

phylogenetic scales (*sensu* Graham et al. 2018), mostly with the aim to decompose biodiversity patterns for higher taxa (Marquet et al. 2004). They typically show that SPRs for different subclades vary and deviate from the pattern observed in the whole large clade (Currie 1991, Buckley et al. 2010, Weiser et al. 2018). Buckley et al. (2010) argue that this finding is inconsistent with the hypotheses explaining richness patterns by productivity. In contrast, Hurlbert and Stegen (2014) argue that weak and varying SPRs of smaller clades are expected if energy availability (productivity) constrains species richness of the whole large clade. Further, Hawkins et al. (2007) show that for many taxa and regions the slope of the STR significantly differs from the slope predicted by the metabolic theory of ecology. However, there are mathematical reasons why the patterns for subgroups often cannot be the same as those for the whole group (Storch and Šizling 2008). Variation of STRs and SPRs across phylogenetic scales is thus expectable.

Here, our primary goal is not to test or advocate any particular mechanism hypothesized to generate species richness patterns. Instead, using a comprehensive data sets on STRs and SPRs, we simultaneously investigate the effects of multiple factors on their strength and direction. We test several hypotheses, based on above considerations. We expect the strength of both relationships to increase with increasing (1) variance of respective environmental variables, (2) spatial grain and extent and (3) taxon size (total number of species). Assuming that both temperature and productivity simultaneously affect species richness, we additionally expect that (4) both STR and SPR are stronger in regions where temperature correlates positively with productivity, and that (5) STRs and SPRs are similarly strong and are approximately equally affected by all the abovementioned factors. In contrast, if only one of these variables directly drives species richness patterns, while the other only correlates with it or has an indirect effect (e.g. if temperature affect species richness only by its effect on productivity), then we expect (6) that one of the relationships (either the STR or the SPR) would be more universal, i.e. less variable and less dependent on the examined factors.

Material and methods

Species data

We used two different types of data for the analysis: (1) a combination of heterogeneous datasets of species richness for different ectothermic taxa spanning different geographical regions (hereafter *metadata*), (2) global species richness data for amphibians, birds and mammals (hereafter *phylodata*). *Metadata* comprise 46 sets of species richness data for a wide range of plant, invertebrate and ectothermic vertebrate taxa used in Hawkins et al. (2007) (Supporting information Table S1). The taxonomic breadth of the datasets ranges from genus (bumblebees) to divisions (vascular plants). The datasets cover all continents (except Antarctica) and their geographical extent ranges from Catalonia (ca. 30,000 km^2) to global. The datasets also differ in their spatial resolution, most of them using equal-area grid cells (grids from 10 km^2 to 611,000 km^2), but in five datasets the units are political regions (Californian plants and butterflies) or nature reserves (Chinese amphibians, reptiles and angiosperms). The datasets were kindly provided by B. Hawkins and more information about the data and their original sources can be found in Hawkins et al. (2007).

In the case of *phylodata*, we examined the STR and the SPR for every amphibian, avian and mammalian clade (defined by every node) larger than 50 species within its whole distributional range on the 1° grid. We used the data on the global distribution of mammalian and amphibian species provided by the IUCN Global Species Programme (<http://www.iucnredlist.org>), and avian species provided by BirdLife International (<http://www.datazone.birdlife.org>). These two databases contain distributional information on species ranges in georeferenced polygons for 5,298 terrestrial mammal, 6,493 amphibian, and 10,961 bird species (extinct species were excluded), which we transformed into a 1° (~100.2 km) spatial grid using the Mollweide coordinate system (equal-area projection). Using only clades larger than 50 species, our data contain 361 clades of amphibians, 502 clades of birds and 249 clades of mammals. We used amphibian phylogeny from <https://doi.org/10.5061/dryad.cc3n6j5> (Jetz and Pyron 2019), avian phylogeny from

<http://birdtree.org/> (Jetz et al. 2012) and mammalian phylogeny from

<https://doi.org/10.5061/dryad.rc416> (Rosauer et al. 2017).

Environmental data

For *metadata* we used original temperature data (mean annual temperature) from different sources,

assembled by Hawkins et al. (2007). For *phylodata* we used mean annual temperature data from

CHELSA model (Karger et al. 2017, <http://chelsa-climate.org/>, downloaded in April 2017),

which we rescaled from the original 30 arcsec resolution to the scale of species data. We used net

primary productivity (NPP) data averaged from years 2000-2012 compiled from the MODIS NPP

layers (MOD17A3, Running et al. 2011, <https://lpdaac.usgs.gov/>) for both *metadata* and

phylodata. We rescaled NPP data from the original resolution 1 km to the resolution of each dataset

(*metadata*) and to the 1° resolution for *phylodata*.

Explanatory variables

We defined several variables describing each dataset (*metadata*) or clade (*phylodata*), which we

then used as the explanatory variables in models explaining variation in the strength of the STR and

the SPR. The description of these variables is in Table 1. For the purpose of one of the analyses, we

divided the explanatory variables into three categories - phylogenetic scale (*size*), spatial scale (*area*

and *grain*) and region characteristics (*meanT, meanNPP, rangeT, rangeNPP* and *T_NPP*).

Statistical analysis

As a measure of the STR and the SPR strength we used correlation (Spearman's coefficient)

between the number of species and temperature (*S_Temp*) and the number of species and NPP

(*S_NPP*), respectively. We performed the analyses also using R-square (coefficient of

determination) of the OLS regression model (linear or quadratic, based on AIC comparison) instead

of Spearman's correlation, but it led to similar results. We thus decided to present only the analyses

using the Spearman's correlation coefficient because it reflects (unlike R-square) both the strength and the direction of the relationships, regardless whether the relationship is linear or curvilinear. Almost all the relationships are monotonous, so the rank-order correlation coefficient well characterizes their strength.

For *metadata*, we used weighted least square (WLS) regression as a basis of all analyses, since the frequency distribution of *S_Temp* and *S_NPP* was close to normal. We accounted for different sample size of the datasets by using weighting by square root of sample size. First, we performed individual regression models for each single explanatory variable alone. We tested for the curvilinearity in these univariate models by comparing the AIC of models containing the dataset characteristics in (1) linear term and in (2) both linear and quadratic term. If the AIC difference was lower than 2, we considered the relationship linear, otherwise we used the model with lower AIC (Burnham and Anderson 2002). Second, we explored how much variation in *S_Temp* and *S_NPP* was explained by the full model using all explanatory variables (in linear term) together. We are aware of the fact that these full models are not entirely statistically accurate due to collinearity in data. Still, since we only wanted to find out the total variation explained by all variables (i.e. in this case we did not explore model parameters comprising the effects of individual variables), since we present the adjusted R^2 values (which adjust for the number of predictors), and since we deal with the collinearity in further analysis, we consider this approach appropriate. Third, we used variation partitioning analysis to distinguish independent effects of the explanatory variables. Following Peres-Neto et al. (2006), we performed the variation partitioning using adjusted R^2 to assess the variation explained by the explanatory variables and their combinations, because it is the only unbiased method. To keep the results clear, we did not use all the individual explanatory variables in the partitioning. Instead, we first performed variation partitioning analysis for three groups of variables - (i) phylogenetic scale (variable *size*), (ii) spatial scale (variables *grain* and *area*) and (iii) region characteristics (the three region characteristics which had the strongest effect in the univariate models). Then, we further decomposed the effects of these three region characteristics.

Each variable entered the variation partitioning analysis in linear or quadratic form depending on the results of the univariate models. Variables *size, area* and *grain* were log-transformed in order to obtain normal distribution of the variables in all described models.

For *phylodata*, to account for the phylogenetic correlation and nestedness among our data points (clades), we used phylogenetic generalized least squares (PGLS, Freckleton et al. 2002) as a basis of all analyses. Specifically, we first estimated phylogenetic dependence among the clades by searching for the maximum likelihood value of the parameter λ, which varies from 0 (phylogenetic independence) to 1 (phylogenetic signal consistent with the Brownian evolutionary model), and then we fitted a GLS model while controlling for the phylogenetic dependence, using package 'caper' (https://CRAN.R-project.org/package=caper, Orme et al. 2013, Machac et al. 2018). This way we performed all the analyses described above - univariate models, full model and the variation partitioning analysis. We performed all the analyses for 100 randomly selected phylogenetic trees and then averaged the results from these 100 models. Variables *size* (for all three vertebrate groups) and *area* (only for amphibians) were log-transformed in order to obtain normal distribution of the variables.

PGLS accounts for the dependence among clades given by their phylogenetic distance, but not necessarily for their nestedness, causing that patterns observed within a clade may be affected by patterns concerning the subclades and vice versa (Graham et al. 2018). There is no simple solution of this problem. One possibility would be to use only a set of non-nested clades (Machac et al. 2018). Nevertheless, we preferred not to use this approach, as the selection of exclusively non-nested clades would be highly non-random, would lead to small sample sizes, and would required repeated analyses on different non-nested clades sets, which would complicate the results interpretation. Using all possible clades circumvents the problem of clade selection, and also involves a range of clades, from lower to higher taxa, providing more complex results. Results from other studies (Machac et al. 2018) indicate that the results for nested and non-nested clades are

similar, and although nestedness may bias statistical significance of the tests, it does not affect the patterns themselves. All the analyses were performed in R (R Development Core Team 2017).

Results

Strength of STRs and SPRs

NPP is in average a stronger correlate of species richness than temperature for all tested data (Fig. 1a and 1b). STRs are negative (and concurrently strong) for many datasets and vertebrate clades, while SPRs are in the majority of cases positive (e.g. for *metadata*, STRs are negative for 16 datasets, while SPRs for 5 datasets and they are weak for most of these negative values). Since negative STR is not an evidence for temperature as a driving factor of species richness patterns, NPP can be considered to be a much better predictor of species richness than temperature. Specifically, *S_NPP* is in average by 0.21 higher than *S_Temp* for *metadata* (t_{45}=-2.48, p=0.02), by 0.30 higher for amphibians (t_{360}=-14.49, p<10^{-16}), by 0.19 higher for birds (t_{501}=-14.53, p<10^{-16}) and by 0.20 higher for mammals (t_{248}=-9.24, p<10^{-16}) (paired t-tests have been used). The highest proportion of negative STRs and related highest average difference between *S_Temp* and *S_NPP* found for amphibians likely results from their dependency on water availability. Still, for many datasets and clades, temperature is a better predictor of species richness than NPP, although sometimes both variables and sometimes neither variable predicts species richness well (Fig. 1c). The notable exception comprises birds, where *S_Temp* and *S_NPP* are correlated to each other, and temperature correlates with species richness with a similar strength as NPP.

Factors affecting strength of STRs and SPRs in *metadata*

In *metadata*, we are able to explain 55% of variation in the strength of STRs (*S_Temp*) and 13% of variation in the strength of SPRs (*S_NPP*) by our explanatory variables (adjusted R^2 of the WLS full model), indicating that the NPP-richness relationship is less dependent on the examined factors. *S_Temp* strongly depends on the mean temperature (*meanT*), on the correlation between temperature

and NPP (*T_NPP*) and on the temperature range (*rangeT*) (Table 2a). Specifically, STRs are generally stronger in colder regions, in regions where temperature and NPP are positively correlated and in regions with wider range of temperature (Table 2a, Fig. 2, Supporting information Fig. S1). On the other hand, the strength of the SPR is not as well predictable by our explanatory variables (Table 2a). *S_NPP* depends on the mean temperature (*meanT*), but in the opposite direction than the *S_Temp*. It also depends on the correlation between temperature and NPP (*T_NPP*) (Table 2a). SPRs are generally stronger in warmer regions and in regions where temperature and NPP are correlated (both positively and, curiously, negatively) (Table 2a, Fig. 2, Supporting information Fig. S1).

Interestingly, the strength of STRs and SPRs for *metadata* depends neither on taxon size nor on spatial scale (in terms of both *grain* and *area*; Table 2a). The absence of a univariate effect of the geographical extent (*area*) suggests that environmental gradients (namely *rangeT*) are more important determinants of the strength of the STR than geographical extent per se. Indeed, the variation partitioning indicates that when we compare STRs and SPRs for different taxa in different (arbitrarily chosen) regions of the world, the strength of the STR and the SPR depends almost entirely on environmental characteristics of the selected region (Fig. 3 and 4), although *meanT* and *T_NPP* are correlated (Fig. 2a) and we cannot completely separate their effects. In the case of STRs (*S_Temp*), the largest portion of variation is explained by the joined effects of *meanT* and *T_NPP*, but there is also a separate effect of *T_NPP* (unlike *meanT*, Fig. 3). In contrast, in the case of SPRs (*S_NPP*), the separate effect of *meanT* is larger than the separate effect of *T_NPP* (Fig. 4). Thus, the strength of SPRs depends mostly on mean temperature within the region (SPRs being stronger in warm, low latitude regions) but the strength of STRs depends on how temperature correlates with NPP within the region, suggesting that the NPP is a more proximate driver of species richness than temperature.

Factors affecting strength of STRs and SPRs in *phylodata*

For *phylodata,* the results are different in several respects. First, our variables explain lower amount of variation in the strength of STRs than in *metadata,* especially for avian and mammalian clades. The full models explain 44% of the variation in STRs strength and 32% of the variation in SPRs strength for amphibians, 21% of the variation in STRs strength and 20% of the variation in SPRs strength for birds, and 17% of the variation in STRs strength and 23% of the variation in SPRs strength for mammals (adjusted R^2 of the PGLS full models). Second, in contrast to *metadata,* there is a significant effect of the clade size on the strength of the relationships (Table 2b), indicating that the expected effect of the taxon size is important only when comparing different subclades within one taxonomical group. Moreover, taxon size is the only important factor explaining the strength of SPRs in the univariate models in avian and mammalian clades (Table 2b). However, even the univariate effect of taxon size is not very strong (Table 2b), as STRs and SPRs are more idiosyncratic for smaller clades, while for large clades they are strong in vast majority of cases (Supporting information Fig. S2 and S3). Third, we can see the difference between amphibian (ectothermic) clades on one side and avian and mammalian (endothermic) clades on another side when exploring the results of univariate models. The strength of STRs is affected by the same variables as in *metadata* only for amphibian clades, for which the strength of STRs decreases with mean temperature (*meanT*) and increases with temperature-NPP correlation (*T_NPP*) and the range of temperature (*rangeT*) (Table 2b, Supporting information Fig. S2). In summary, for *phylodata*, (1) the most ubiquitous factor affecting the strength of STRs and SPRs is clade size (the relationships are strong for large clades), (2) clade size is the only factor consistently affecting the strength of SPRs in avian and mammalian clades and (3) only in amphibians, there are univariate effects of environmental characteristics of the clade range (mean temperature, temperature-NPP correlation and temperature range), which are similar as in the case of *metadata*.

The effect of phylogenetic scale (clade size) is shown also by the variation partitioning analysis, where partial effects of clade size are ubiquitous (Fig. 3 and 4, note that 'phylogenetic

scale' represents only one variable, i.e. *size*, while 'region characteristics' represents three variables). Variation partitioning analysis also reveals that partial effect of the region characteristics is still important for the STR while it is weak for the SPR.

Discussion

The effects of taxon size and spatial scale on the the STR and the SPR

We provide the first analysis that compares the effects of multiple taxon- and region-related variables on the strength of the STR and the SPR. We investigated this using two different types of data, representing two common types of data structure in macroecological research: (1) various taxa sampled using different sampling techniques within different regions of the world (Hawkins et al. 2007) and (2) grid-cell data of amphibian, avian and mammalian clades within their global ranges. There was a striking difference in the drivers of variation in the explored species richness patterns between the two types of data. Taxon size affected the strength of STRs and SPRs only when comparing subclades within vertebrate classes, i.e. when exploring these relationships across different phylogenetic scales (sensu Graham et al. 2018). Weiser et al. (2018) in their study analysing latitudinal gradient of species richness (LGSR) for different nested plant clades concluded that this non-universality and variability of LGSR suggests that climate or energy fail to explain LGSR and that this result instead supports the 'tropical conservatism' hypothesis. However, in accord with Hurlbert and Stegen (2014), we argue that this inconsistency in smaller taxa does not exclude temperature or productivity as important factors shaping species richness patterns. There are several possible reasons why smaller taxa may exhibit weak and variable relationships: (1) simple statistical reasons (sampling effects), (2) species richness of small clades may be affected by a specific environmental factor related to specifities of their biology, or interaction with other taxa, (3) smaller taxa may be young and their richness patterns thus may be driven by their recent history of spreading from an area of origin (Gehrke and Linder 2011).

In this respect, it is interesting that we did not find any univariate effects of clade size when comparing different taxa within different regions (i.e. the meta-analytic approach; as in e.g. Mittelbach et al. 2001, Hawkins et al. 2007, Field et al. 2009). We admit that 46 datasets is perhaps too low number for revealing such an effect. Still, considering that we were able to show other strong effects using the same data, we propose that when comparing different taxa in different regions (meta-analytic approach), our choice of the region is what really matters, relatively independently of the taxon examined. The fact that such regions almost never represent real ranges of the whole taxa would strengthen this effect - when examining richness patterns for particular, arbitrarily delimited region (e.g. California or Catalonia), the effect of its environment (mean temperature, temperature-productivity relationship or the range of temperature within the region) is always essential. The same can probably explain the absence of the univariate effect of spatial scale (both grain and extent). Therefore, in the case of the meta-analytic approach comprising very different datasets, it only matters which region is selected from the global map of temperature-productivity conditions, relatively independently of the taxon studied.

The drivers of the variation of STRs and SPRs

We have shown that the strength of both STR sand SPR is affected mostly by mean temperature and temperature-productivity correlation within examined region or within clade range, even though this does not apply generally across all taxa and the effect is weaker in the case of SPRs. To be specific, STRs are generally stronger in colder regions (within temperate zone), while SPRs are generally stronger in warmer regions (within the tropics). This is consistent with earlier studies showing that in higher latitudes, species richness is best predicted by temperature or potential evapotranspiration (PET, which is driven mostly by temperature), while in lower latitudes it is best explained by productivity or water availability (Hawkins et al. 2003a). Similarly, avian species diversity was shown to be best predicted by temperature or PET within Palearctic and Nearctic regions, and by productivity or rainfall in the Afrotropics, Neotropics and Australia (Hawkins et al. 2003b). In

colder regions, temperature likely becomes the limiting factor, while in hot regions, water availability and related ecosystem productivity is likely to be limiting. The strength of STRs in ectotherms was also affected by temperature range, which is probably driven by the fact that regions with the narrow temperature range do not form sufficient temperature gradient.

The temperature-productivity correlation within studied region (which is quite evenly distributed from strong negative to strong positive in our data, Supporting information Fig. S1, S2 and S3) appears as an important source of the STR and the SPR variation. Both these relationships are weak where the temperature-productivity correlation within the region is weak (Fig. 2), suggesting that both environmental variables simultaneously affect species richness - if only one variable causally affected species richness, the other being just a surrogate, the strength of either STR or SPR should not depend on its correlation with the other variable. The interpretation that both temperature and productivity directly drive diversity patterns seems to be consistent with synthetic theories of large-scale biodiversity patterns (Allen et al. 2007, Storch et al. 2018, Worm and Tittensor 2018, Storch and Okie 2019). These theories assume that while productivity constrains total number of individuals, temperature may affect rates of processes generating diversity. However, the temperature-productivity correlation had in fact a strong independent effect only on STRs, while in the case of SPRs the effect was considerably weaker and mostly shared with the effect of mean temperature (Fig. 4). Moreover, the regions in which temperature and productivity were negatively correlated revealed negative STRs but positive SPRs. This may indicate that temperature is not as crucial factor determining species richness, especially in the tropics, which is the region typically characterized by the negative temperature-productivity correlation (Fig. 2). In contrast, in the temperate regions, temperature may limit species richness not only via limiting biological rates, but also by limiting pools of available species that can tolerate lower temperatures (Šímová et al. 2011), in accord with the niche conservatism hypothesis (Wiens and Donoughue 2004).

The role of temperature in generating diversity patterns may be even weaker in endotherms. We have shown that while variation of STRs and SPRs strength in amphibian clades reveals similar causes as the variation in the datasets of various ectothermic taxa (*metadata*) (with the exception of differences caused by the different nature of the data discussed above), the SPRs for avian and mammalian clades are largely independent of our explanatory variables, with the exception of the taxon size discussed above. Given that SPRs are generally strong (Fig. 1) and independent of most of the examined variables in endotherms, productivity indeed appears as the key factor shaping species richness patterns for these taxa. Our results indicate that although temperature may be as good predictor of species richness as NPP for some clades and datasets (Fig. 1c), the finding that *in general* SPRs are positive, strong, and largely independent of most of the explored variables (compared to STRs, Fig. 1) indicates that productivity is more important species richness driver than temperature - and this is especially pronounced in endotherms.

Conclusion

We provide the first comprehensive test of how different variables comprising spatial scale and the characteristics of examined taxa and regions affect the strength of species-temperature and species-productivity relationships. Both relationships are stronger for larger clades, indicating that phylogenetic scale matters (Weiser et al. 2018), but only when comparing different subclades within one large taxon. When comparing different taxa in different regions (meta-analytical approach), only the role of region characteristics (mean temperature and temperature-productivity correlation within the region) has been revealed. We provide new evidence for the previously documented patterns that temperature plays a more important role in colder regions, while productivity is more important in warmer regions (Hawkins et al. 2003a, 2003b). Additionally, both relationships are weak in regions where temperature and productivity are not correlated. These findings indicate that both productivity and temperature simultaneously affect species richness, at least in ectotherms. However, the relationships between species richness and productivity are less variable and at the

same time more independent of all the studied factors, indicating that the productivity-richness relationship is more universal and context-independent than the temperature-richness relationship. This may stem from the universal productivity effects on the limits of species richness (sensu Hurlbert and Stegen 2014), i.e. species richness carrying capacity (Storch and Okie 2019). We suggest that every study aiming to test a mechanism hypothesised to form spatial patterns in species richness should pay careful attention to the effects potentially causing the variation of the strength of species richness patterns, especially when data only for arbitrarily chosen regions and/or taxa are used.

Funding

This research was supported by the Czech Science Foundation (grant no. 20-29554X).

Data availability statement

Most of the data used in the study (with the exception specified below) are freely available from the cited and publicly accessible databases. 46 datasets containing data on species richness and mean annual temperature for different taxa and regions (referred as *metadata* in the text) were assembled by Hawkins et al. (2007) and kindly provided by Bradford Hawkins for the purpose of the study. We provide the list of the datasets with the values of response and explanatory variables used in our analysis in the Supporting information.

References

Abrams, P. A. 1995 Monotonic or unimodal diversity productivity gradients - what does competition theory predict. – Ecology 76: 2019–2027.

Allen, A. P. et al. 2002. Global biodiversity, biochemical kinetics, and the energetic-equivalence rule. – Science 297: 1545–1548.

Allen, A. P. et al. 2007. Recasting the species-energy hypothesis: the different roles of kinetic and potential energy in regulating biodiversity. – In: Storch, D. et al. (eds.), Scaling biodiversity. Cambridge Univ. Press, pp. 283–299.

Belmaker, J. and Jetz, W. 2011. Cross-scale variation in species richness–environment associations. – Glob. Ecol. Biogeogr. 20: 464–474.

BirdLife International and Handbook of the Birds of the World 2017. Bird species distribution maps of the world. Version 6.0. Retrieved from <http://datazone.birdlife.org>.

Brown, J. H. et al. 2004. Toward a metabolic theory of ecology. – Ecology 85: 1771–1789.

Buckley, L. B. et al. 2010. Phylogeny, niche conservatism and the latitudinal diversity gradient in mammals. – Proc. Royal Soc. B 277: 2131–2138.

Burnham, K. and Anderson, D. 2002. Model selection and multi-model inference. – Springer.

Currie, D. J. 1991. Energy and large-scale patterns of animal- and plant-species richness. – Am. Nat. 137: 27–49.

Currie, D. J. et al. 2004. Predictions and tests of climate-based hypotheses of broad-scale variation in taxonomic richness. – Ecol. Lett. 7: 1121–1134.

Evans, K. L. et al. 2005. Species–energy relationships at the macroecological scale: a review of the mechanisms. – Biol. Rev. 80: 1–25.

Field, R. et al. 2009. Spatial species richness gradients across scales: a meta-analysis. – J. Biogeogr. 36: 132–147.

Freckleton, R. P. et al. 2002. Phylogenetic analysis and comparative data: A test and review of evidence. Am. Nat. 160: 712–726.

Gaston, K. J. 2000. Global patterns in biodiversity. – Nature 405: 220–227.

Gehrke, B. and Linder, P. 2011. Time, space and ecology: why some clades have more species than others. – J. Biogeogr. 38: 1948–1962.

González-Taboada, F. et al. 2007. Breeding bird species richness in Spain: assessing diversity hypothesis at various scales. – Ecography 30: 241–250.

Graham C. H. et al. 2018. Phylogenetic scale in ecology and evolution. – Glob. Ecol. Biogeogr. 27: 175–187.

Hawkins, B. A. et al. 2003a. Energy, water, and broad-scale geographic patterns of species richness. – Ecology 84: 3105–3117.

Hawkins, B. A. 2003b. Productivity and history as predictors of the latitudinal diversity gradient of terrestrial birds. – Ecology 84: 1608–1623.

Hawkins, B. A. et al. 2007. A global evaluation of metabolic theory as an explanation for terrestrial species richness gradients. – Ecology 88: 1877–1888.

Hillebrand, H. 2004. On the generality of the latitudinal diversity gradient. – Am. Nat. 163: 192–211.

Hubbell, S. P. 2001. The unified neutral theory of biodiversity and biogeography. – Princeton Univ. Press.

Hurlbert, A. H. and Stegen, J. C. 2014. When should species richness be energy limited, and how would we know? – Ecol. Lett. 17: 401–413.

IUCN 2017. The IUCN red list of threatened species. Version 2017–1. Retrieved from <http://www.iucnredlist.org>.

Jetz, W. and Pyron, R. A. 2019. Data from: The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. – Dryad Digital Repository, <https://doi.org/10.5061/dryad.cc3n6j5>.

Jetz, W. et al. 2012. The global diversity of birds in space and time. – Nature 491: 444–448.

Karger, D. N. et al. 2017. Climatologies at high resolution for the earth's land surface areas. – Sci. data 4: 170122.

Kaspari, M. et al. 2003. Spatial grain and the causes of regional diversity gradients in ants. – Am. Nat. 161: 459–477.

Lennon, J. J. et al. 2000. Bird diversity and environmental gradients in Britain: a test of the species–energy hypothesis. – J. Anim. Ecol. 69: 581–598.

Machac A. et al. 2018. Ecological controls of mammalian diversification vary with phylogenetic scale. – Glob. Ecol. Biogeogr. 27: 32–46.

Marquet, P. A. et al. 2004. Diversity emerging: toward a deconstruction of biodiversity patterns. – In: Lomolino, M. V. and Heaney, L. R. (eds.), Frontiers of biogeography: new directions in the geography of nature. Sinauer, pp 191–210.

McPeek, M. A. 1996. Linking local species interactions to rates of speciation in communities. – Ecology 77: 1355–1366.

Mittelbach, G. G. et al. 2001. What is the observed relationship between species richness and productivity? – Ecology 82: 2381–2396.

Orme, D. et al. 2013. CAPER: comparative analyses of phylogenetics and evolution in R. – Methods Ecol. Evol. 3:145–151.

Peres-Neto, P. R. et al. 2006. Variation partitioning of species data matrices: Estimation and comparison of fractions. Ecology 87: 2614–2625.

Pontarp, M. et al. 2019. The latitudinal diversity gradient: novel understanding through mechanistic eco-evolutionary models. – Trends Ecol. Evol. 34: 211–223.

R Core Team 2018. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

Rahbek, C. and Graves, G. R. 2000. Detection of macro-ecological patterns in South American hummingbirds is affected by spatial scale. – Proc. Royal Soc. B 267: 2259–2265.

Rahbek, C. and Graves, G. R. 2001. Multiscale assessment of patterns of avian species richness. – Proc. Natl. Acad. Sci. USA 98: 4534–453.

Rohde, K. 1978. Latitudinal gradients in species diversity and their causes II. Marine parasitological evidence for a time hypothesis. – Biol. Zentralbl. 97: 405–418.

Rosauer, D. F. et al. 2017. Data from: Phylogenetically informed spatial planning is required to conserve the mammalian tree of life. Dryad Digital Repository, <https://doi.org/10.5061/dryad.rc416>.

Running, S. et al. 2011. MOD17A3 MODIS/Terra Net Primary Production Yearly L4 Global 1km SIN Grid V055 [Data set]. NASA EOSDIS Land Processes DAAC.

Šímová, I. and Storch, D. 2017. The enigma of terrestrial primary productivity: measurements, models, scales and the diversity–productivity relationship. – Ecography 40: 239–252.

Šímová, I. et al. 2011. Global species energy relationship in forest plots: role of abundance, temperature and species climatic tolerances. – Glob. Ecol. Biogeogr. 20: 842–856.

Stevens, M. H. H. et al. 2007 Diversification rates increase with population size and resource concentration in an unstructured habitat. – Genetics 177: 2243–2250.

Storch, D. 2012. Biodiversity and its energetic and thermal controls. – In Sibly, R. M. et al. (eds.), Metabolic ecology: A scaling approach. Wiley-Blackwell, pp 120–131.

Storch, D. and Šizling, A. L. 2008. The concept of taxon invariance in ecology: Do diversity patterns vary with changes in taxonomic resolution? – Folia Geobot. 43: 329–344.

Storch, D. and Okie, J. 2019. The carrying capacity for species richness. – Glob. Ecol. Biogeogr. 28: 1519–1532.

Storch, D. et al. 2018. The more-individuals hypothesis revisited: the role of community abundance in species richness regulation and the productivity-diversity relationship. – Ecol. Lett. 21: 920–937.

Weiser, M. D. et al. 2018. Taxonomic decomposition of the latitudinal gradient in species diversity of North American floras. – J. Biogeogr. 45: 418–428.

Wiens, J. J. and Donoghue, M. J. 2004. Historical biogeography, ecology and species richness. – Trends Ecol. Evol. 19: 639–644.

Worm, B. and Tittensor, D. P. 2018. A theory of global biodiversity. – Princeton Univ. Press.

Wright, D. H. 1983. Species-energy theory: an extension of species-area theory. – Oikos, 41: 496–506.

Table 1. Explanatory variables for both types of data (*metadata* and *phylodata*).

VARIABLE	EXPLANATION	NOTE
size	total number of species in the dataset (*metadata*) or species richness of the clade (*phylodata*)	unknown for Chinese amphibians, reptiles and angiosperms, Australian tiger beetles, and Iberian/French dung beetles
area	geographical extent of the dataset in km^2 (*metadata*) or clade range size measured as the number of grid cells occupied by given clade (*phylodata*)	not defined in New World ants, North American grasshoppers, Chinese amphibians, reptiles and angiosperms (since species richness data did not cover continuously any clearly defined region)
meanT	mean annual temperature in °C within the dataset (*metadata*) or within the clade range (*phylodata*)	
meanNPP	mean NPP within the dataset (*metadata*) or within the clade range (*phylodata*)	
rangeT	range of mean annual temperature within the dataset (*metadata*) or within the clade range (*phylodata*) (the difference between the lowest and the highest value)	
rangeNPP	range of NPP within the dataset (*metadata*) or within the clade range (*phylodata*) (the difference between the lowest and the highest value)	
T_NPP	Spearman's correlation coefficient between temperature and NPP within the dataset (*metadata*) or within the clade range (*phylodata*)	
grain	resolution of the dataset - size of the grid cell for raster data, size of the sampling unit for sample data (New World Ants, North American grasshoppers) and mean area of nature reserves (considering the reserve areas did not vary by more than an order of magnitude) for Chinese amphibians, reptiles and angiosperms in km^2	defined only for *metadata* (since it is identical for all *phylodata*); not defined for Californian plants and butterflies because there it represented political units whose area varied by orders of magnitude.

Table 2. Results of the univariate models explaining the variation in the strength of the STR (left column) and in the strength of the SPR (right column) for (a) datasets of various ectothermic taxa – *metadata*, and for (b) vertebrate clades – *phylodata*. The effects significant on α=0.05 are in bold, for the explanation of the variables see Table 1. AIC corrected for small sample size (AICc), adjusted R^2 (adj. R^2) and coefficients (coefs.) are displayed.

	SPECIES-TEMPERATURE RELATIONSHIP				SPECIES-PRODUCTIVITY RELATIONSHIP			
(a)	**Model**	**AICc**	**adj. R^2**	**coefs.**	**Model**	**AICc**	**adj. R^2**	**coefs.**
M E T A D A T A	***meanT***	**54.6**	**0.50**	**-**	***meanT***	**12.6**	**0.16**	**+**
	T_NPP	**55.5**	**0.49**	**+**	***T_NPP+T_NPP²***	**17.4**	**0.10**	**-;+**
	rangeT + rangeT²	**72.1**	**0.29**	**-;+**	log(*size*)	18.6	0.01	+
	log(*area*)	78.6	0.00	+	log(*area*)	19.6	0.00	+
	log(*size*)	79.5	0.00	+	*rangeT*	20.4	0.01	-
	rangeNPP	80.8	0.12	-	log(*grain*)	20.5	0.00	-
	log(*grain*)	82.7	0.00	-	*rangeNPP*	20.6	0.00	+
	meanNPP	86.1	0.01	-	*meanNPP*	21.1	0.00	-
(b)								
A M P H I B I A N S	***T_NPP+T_NPP²***	**-210.9**	**0.33**	**+;-**	**log(*area*)+log(*area*)²**	**-277.1**	**0.22**	**-;+**
	log(*area*)+log(*area*)²	**-210.8**	**0.33**	**-;+**	**log(*size*)**	**-267.4**	**0.19**	**+**
	meanT+meanT²	**-197.5**	**0.30**	**+;-**	***meanNPP+meanNPP²***	**-259.7**	**0.17**	**-;+**
	rangeT+rangeT²	**-189.9**	**0.29**	**-;+**	***rangeT+rangeT²***	**-246.8**	**0.15**	**+;+**
	meanNPP+meanNPP²	**-185.5**	**0.25**	**-;-**	***meanT***	**-242.9**	**0.14**	**-**
	log(*size*)+log(*size*)²	**-117.4**	**0.12**	**-;+**	***T_NPP+T_NPP²***	**-222.7**	**0.10**	**+;+**
	rangeNPP	-82.0	0.02	+	*rangeNPP*	-203.6	0.03	+
B I R D S	***meanT+meanT²***	**-277.0**	**0.16**	**+;-**	**log(*size*)**	**-471.7**	**0.15**	**+**
	T_NPP	**-276.5**	**0.15**	**+**	***meanNPP+meanNPP²***	**-407.6**	**0.03**	**+;-**
	log(*size*)	**-237.8**	**0.09**	**+**	***area***	**-400.0**	**0.01**	**+**
	rangeT+rangeT²	**-237.5**	**0.09**	**+;-**	***rangeNPP***	**-398.3**	**0.01**	**+**
	meanNPP+meanNPP²	**-225.9**	**0.05**	**+;-**	***rangeT***	**-394.1**	**0.00**	**+**
	area	**-202.9**	**0.02**	**+**	***T_NPP***	**-393.0**	**0.00**	**-**
	rangeNPP	**-198.7**	**0.01**	**+**	***meanT***	**-392.4**	**0.00**	**-**
M A M M A L S	***T_NPP+T_NPP²***	**-163.1**	**0.17**	**+;-**	**log(*size*)**	**-252.4**	**0.15**	**+**
	meanT+meanT²	**-149.4**	**0.12**	**+;-**	***area***	**-223.2**	**0.04**	**+**
	log(*size*)	**-139.4**	**0.08**	**+**	***meanT+meanT²***	**-222.6**	**0.05**	**+;-**
	area	**-126.0**	**0.03**	**+**	***rangeT***	**-221.2**	**0.04**	**+**
	rangeT	**-124.6**	**0.02**	**+**	***rangeNPP***	**-214.0**	**0.01**	**+**
	meanNPP	-123.2	0.02	-	*meanNPP*	-212.6	0.00	-
	rangeNPP	-121.3	0.01	+	*T_NPP*	-212.2	0.00	-

Figure 1. Histograms showing the distribution of the frequencies of (a) the STR strength and (b) the SPR strength for each data set (the first row refers to *metadata,* the following three rows to *phylodata*). The colours are scaled from yellow (weak relationship) to red (negative relationship) and to green (positive relationship) and correspond to the colour scale in Fig. 2. Sample size (n) is presented inside each plot. (c) The relationship between the strength of the STR (*S_Temp*) and the strength of the SPR (*S_NPP*) for each data set (rows). The Pearson's correlation coefficient (r) is presented inside each plot.

Figure 2. Plots showing the strength of the STR (a) and the SPR (b) of *metadata* in relation to region characteristics. Mean values and ranges of temperature and NPP are reflected by the position of the ellipse within the plot, and the temperature-NPP correlation is reflected by the diagonal elongation of the ellipse. Each ellipse represents one dataset and it encompasses 80% of data points. Colours of the ellipses correspond to the strength of the STR (*S_Temp)* and the SPR (*S_NPP)*, respectively, according to the colour bar. Additionally, we split the datasets so that only the strong positive, strong negative and weak relationships are presented (bottom). Note that there are no strong negative SPRs. (a) STRs are strong and positive in colder regions, regions with wider temperature range and regions with positive temperature-NPP relationship. On the other hand, STRs are strong and negative in warmer (tropical) regions, in regions with narrower temperature range and those revealing negative temperature-NPP relationships. They are generally weak for regions where temperature and NPP are not correlated. (b) SPRs do not show as clear patterns as STRs, although the effect of the mean temperature is still visible – SPRs are generally stronger in warmer regions and they are mostly weak in regions where temperature and NPP are not correlated.

Figure 3. Venn diagrams describing variation partitioning for the strength of the STR (*S_Temp*) for different data (rows). In the left column, explanatory variables are divided into 3 categories - phylogenetic scale (variable *size*), spatial scale (variables *area* and *grain* - only for *metadata*) and region characteristics (three region characteristics with the strongest univariate effect on *S_Temp*, see Table 2 and Table 1). In the right column, the variation partitioning is further performed for these three region characteristics. The numbers represents adjusted R^2 (in %), zero and negative values are not displayed. Every section is coloured according to its adjusted R^2 value on the white-black scale (white corresponds to 0, black corresponds to 50).

Figure 4. Venn diagrams describing variation partitioning for the strength of the SPR (*S_NPP*) for different data (rows). In the left column, explanatory variables are divided into 3 categories - phylogenetic scale (variable *size*), spatial scale (variables *area* and *grain* - only for metadata) and region characteristics (three region characteristics with the strongest univariate effect on *S_NPP*, see Table 2 and Table 1). In the right column, the variation partitioning is further performed for these three region characteristics. The numbers represents adjusted R^2 (in %), zero and negative values are not displayed. Every part is coloured according to its adjusted R^2 value on the white-black scale (white corresponds to 0, black corresponds to 50).