Rate-of-change analysis in palaeoecology revisited: a new approach

Ondřej Mottl[1*], John-Arvid Grytnes[1], Alistair W.R. Seddon[2], Manuel J. Steinbauer[1,3], Kuber P. Bhatta[1],

Vivian A. Felde[2], Suzette G.A. Flantua[1], H. John B. Birks[2,4]

1. Department of Biological Sciences, University of Bergen, PO Box 7803, N-5020 Bergen, Norway

2. Department of Biological Sciences and Bjerknes Centre for Climate Research, University of

Bergen, PO Box 7803, N-5020 Bergen, Norway

3. Bayreuth Center of Ecology and Environmental Research (BayCEER) and Department of Sport

Science, University of Bayreuth, 95547 Bayreuth, Germany

4. Environmental Change Research Centre, University College London, Gower Street, London,

WC1 6BT, UK.

Correspondence author:

*Ondřej Mottl (ORCID number 0000-0002-9796-5081)

Email: ondrej.mottl@gmail.com

Address: Department of Biological Sciences, University of Bergen, PO Box 7803, N-5020 Bergen,

Norway

ABSTRACT

Dynamics in the rate of compositional change beyond the time of human observation are uniquely

preserved in palaeoecological sequences of terrestrial and marine origin. Periods with a high

rate-of-change (RoC) have been assigned to reflect changes due to external factors such as climate

and human forcing. However, changes in sedimentation rates and sampling strategies can result in an

uneven distribution of time intervals within a stratigraphic sequence, which are known to affect RoC

estimates and thus derived conclusions. Several approaches have been used to account for these

issues (e.g. binning of levels, data smoothing, etc.) but despite the importance and frequency of these,

there has been relatively little exploration of the implications of these challenges into quantifying RoC

in palaeoecology.

Here, we introduce R-Ratepol – an easy-to-use R package – that provides a new, robust numerical

technique for detecting and summarising RoC patterns in complex multivariate time-ordered

stratigraphical sequences. First, we compare the performance of common methods of estimating RoC

and detection of periods of high RoC (peak-point) using simulated pollen-stratigraphical data with

known patterns of compositional change and temporal resolution. Then, we apply our new

methodology to four representative European pollen sequences to detect potential drivers of
compositional change.

We show that the successful detection of known shifts in compositional changes identified by RoC
depends heavily on the smoothing methods and dissimilarity coefficients used, and the density of
stratigraphical levels. Identifying the need for an improved numerical approach, we propose a new
method of binning with a moving window in combination with a generalised additive model for peak-
point detection. Our method shows a 68% increase in the correct detection of peak-points compared
to the more traditional way of peak selection by individual levels, as well as achieving a reasonable
compromise between Type I and Type II errors. Likewise, by using the four pollen sequences from
Europe, we show that our approach also performs well in detecting periods of significant
compositional change during known onsets of human activity, early land-use transformation, and
changes in fire frequency.

Expanding the approach using R-Ratepol to the increasing number of open access paleoecological
datasets in global databases, such as Neotoma, will allow future palaeoecological and macroecological
studies to quantify major changes in biotic composition or in sets of abiotic variables across broad
spatio-temporal scale.

KEYWORDS: Compositional change; peak-point detection; stratigraphic sequences; pollen analysis;
rate-of-change; simulated data

1. INTRODUCTION
Quantifying spatio-temporal changes in biological composition or diversity is essential for monitoring
the current biodiversity crisis and for disentangling underlying drivers such as climate, land-use
change, pollution, or introduction of invasive species, as well as intrinsic population dynamics driven
by biotic interactions (e.g. Dodd et al., 1994; Dornelas et al., 2013; Gotelli et al., 2010; Magurran et al.,
2010; Monchamp et al., 2018; Morecroft et al., 2009; Richardson et al., 2006; Silvertown et al., 2006;
Steinbauer et al., 2018; Wolfe et al., 1987). Therefore, the importance of such ecological long-term
observational studies (50-100 years) is increasingly recognised (Dornelas et al., 2013; Hillebrand et al.,
2018; Magurran et al., 2019). Substantial compositional change has been observed during recent
decades (e.g. Feeley et al., 2020; Steinbauer et al., 2018), the last centuries ("The Great Acceleration";
Steffen et al., 2015), and during the Holocene as a result of human impact (Stephens et al., 2019) or
in response to regional climate change (Seddon et al., 2015; Shuman et al., 2005). However, similar
rates of substantial change in composition have also been detected on longer time scales (Kemp et

al., 2015). To understand the impacts of humans, for instance, on ecosystems it is essential to compare
temporal changes in species composition and diversity through human history and to investigate
whether such changes are unique to the epoch of human impact ("Anthropocene") or whether they
precede human-dominated systems (Birks, 1997; Birks et al., 2016).

Assemblage data of palaeoecological sequences of terrestrial and marine proxies (pollen, diatoms,
chironomids, cladocerans, corals, molluscs, sediment chemical variables, etc.) are a unique resource
for quantifying spatio-temporal changes in biotic or abiotic composition and diversity beyond the time
period of human observations. Rate-of-change (RoC) analysis was introduced into palaeoecology by
Jacobson and Grimm (1986) to quantify the rate and the magnitude of compositional change within a
Holocene pollen sequence. It was extended by Jacobson et al. (1987) to quantify and compare rates
of change within and between sequences (see also Grimm and Jacobson, 1992) in an attempt to
identify regional-scale and local-scale patterns in rates of change. RoC analysis estimates
compositional change or temporal beta-diversity between adjacent stratigraphical levels and hence
between times. Unlike other estimates of beta-diversity, RoC analysis specifically estimates the
magnitude of compositional change per *unit time*. Therefore, an essential requirement in RoC analysis
is a robust age-depth model for the stratigraphical sequence of interest to estimate the precise age of
levels within the sequence.

Despite the wide use of RoC analysis in palaeoecology (e.g. Birks and Ammann, 2000; Birks, 1997; Birks
and Birks, 2008; Correa-Metrio et al., 2012; Grindean et al., 2019; Laird et al., 1998; Solovieva et al.,
2008; Urrego et al., 2009), there has been relatively little exploration of the underlying methodology
of RoC analysis and how the various methodologies and sequence properties influence RoC estimates
(but see Bennett and Humphry, 1995; Birks, 2012; Lotter et al., 1992). Different approaches to
estimate RoC (Bennett and Humphry, 1995; Birks, 2012) involve choices in terms of approach and
technique, such as transforming the stratigraphical levels and associated assemblage data (using
binning or interpolation to constant time intervals), smoothing the data, and the metric used to
estimate the amount of compositional dissimilarity between adjacent levels (Birks, 2012).

The variety of approaches available to RoC analysis can, however, create problems. First, Anderson et
al. (2020) stress the sensitivity of RoC to temporal sampling variation within and between records,
consequently hampering conclusive comparisons. Second, there has been a lack of standardisation
between studies (i.e. incorporating different dissimilarity metrics, different temporal procedures)
which means RoC outcomes cannot be directly compared between sequences or studies (Birks, 2012).
Third, intrinsic properties of an individual sequence, such as its taxonomic richness and the density
and distribution of levels within the sequence, can influence the estimated RoC. The end result is that,

since the expected patterns of RoC in a sequence are unknown, the most appropriate choices or
methods to undertake RoC analysis are also unknown (Bennett and Humphry, 1995; Birks, 2012).

Here, we present and evaluate the performance of R-Ratepol, a new R package (R Core Team, 2018),
for RoC analysis introducing a standardised and robust method, which incorporates age uncertainties
from age-depth modelling as well as standardisation of taxa richness, and hence current best practice,
for estimating and comparing RoC estimates within and between stratigraphical sequences. First, we
present the R-Ratepol and its capacities for data commonly used in palaeoecological studies. Second,
we compare the performance of various methods of estimating RoC using simulated pollen-
stratigraphical data (Blaauw et al., 2010) with known patterns of compositional change and resolution.
In addition, we introduce a method for detecting 'peak-points', defined as a rapid change in
composition or relative abundances of variables within the sequence, which provides an intuitive,
much-needed means to compare directly RoC between sequences and thus facilitates the
interpretation of potential drivers of assemblage change on a regional scale. We compare the
effectiveness of our new method for the successful detection of peak-points with commonly used
methodologies which are also provided in R-Ratepol. Finally, we illustrate our findings by estimating
RoC values for four pollen palynological sequences with different densities of stratigraphical levels and
pollen richness and link the observed patterns to known anthropogenic activities. Our approach can
provide important information for future palaeoecological and macroecological studies attempting to
quantify, and then attribute, major changes in biotic or abiotic composition across broad spatial areas,
and to compare the observed changes in recent decades with changes that occurred within the
Holocene (e.g. Gibson-Reinemer et al., 2015).

The term 'assemblage' is used through the text to refer to sets of biotic (pollen, macrofossils, etc.) or
abiotic (sediment chemistry, isotope ratios, etc.) variables studied in levels within a stratigraphical
sediment sequence.

2. MATERIALS AND METHODS

2.1. R-Ratepol

R-Ratepol is written as an R package and includes a range of possible settings including a novel method
to evaluate RoC in a single stratigraphical sequence using assemblage data and age uncertainties for
each level. There are multiple built-in dissimilarity coefficients (DC) for different types of data, and
various levels of data smoothing that can be applied depending on the type and variance of the data.
In addition, R-Ratepol can use randomisation, accompanied by use of age uncertainties of each level

and taxon standardisation to detect RoC patterns in datasets with high data noise or variability (i.e.
numerous rapid changes in composition or sedimentation rates).

The computation of RoC in R-Ratepol is performed using the following steps (Fig. 1) and detailed
descriptions of the underlying methods and formulae are given in the Supplementary Material:

1. Assemblage and age-model data are extracted from the original source and should be
compiled together, i.e. depth, age, taxon 1, taxon 2, etc.
2. (optional) Smoothing of assemblage data: Each variable within the assemblage data is
smoothed using one of five in-built smoothing methods: none, Shepard's 5-term filter (Davis,
1986; Wilkinson, 2005), moving average, age-weighted average, Grimm's smoothing (Grimm
and Jacobson, 1992).
3. Creation of time bins: A template for all time bins in all window movements is created.
4. A single run (an individual loop) is computed:
a. (optional) Selection of one time series from age uncertainties (see section 2.1.1.2. on
randomisation)
b. Subsetting levels in each bin: Here the working units are created (WUs; see section
2.1.1.1.)
c. (optional) Standardisation of assemblage data in each WU (see section 2.1.1.2. on
randomisation)
d. Calculation of RoC between adjacent WUs: RoC is calculated as the dissimilarity
coefficient (DC) standardised by age differences between WUs. Five in-built
dissimilarity coefficients are available: *Euclidean distance*, *standardised Euclidean*
distance, *Chord distance*, *Chi-squared coefficient* (Prentice, 1980), *Gower's distance*
(Gower, 1971). The choice of DC depends on the type of data (see Supplementary
Material).
e. Summary of a single run is produced based on all moving windows
5. Step 4 is repeated multiple times (e.g. 10,000 times).
6. Validation and summary of results from all runs of RoC calculation are produced.
7. (Optional) Data beyond a certain age can be excluded (e.g. 8000 cal yr BP for the pollen
sequences in this study)
8. Detection and validation of significant peak-points. There are five in-built methods to detect
significant peak-points: *Threshold*, *Linear trend*, *Non-linear trend*, *first derivative* of a
generalised additive model (f-deriv GAM; Simpson, 2018), and *Signal-to-Noise Index* (SNI; Kelly
et al., 2011).

2.1.1. Specific methodological considerations

2.1.1.1. Selection of working units (WU; Step 3)

RoC is calculated between consecutive Working Units (WU). Traditionally, these WUs represent individual stratigraphical levels. However, changes in sedimentation rates and sampling strategies can result in an uneven temporal distribution of levels within a time sequence, which in turn makes the comparison of RoC between sequences problematic. There are various methods that attempt to minimise such problems. The first is *interpolation of levels* to evenly spaced time intervals, and the use of the interpolated data as WUs. This can lead to a loss of information when the density of levels is high. Second is *binning of levels*: assemblage data are pooled into age brackets of various size (bins) and these serve as WUs. Here, the issue is a lower resolution of WUs and their uneven size in terms of total assemblage count (bins with more levels have higher assemblage counts). Third is *selective binning*: like classical binning, bins of selected size are created, but instead of pooling assemblage data together, only one level is selected as representative of each bin. This results in an even number of WUs in bins with a similar count size in the assemblage. However, the issue of low resolution remains.

Therefore, we propose a new method of *binning with a moving window*, which is a compromise between using individual levels and selective binning. This method follows a simple sequence: bins are created, levels are selected as in selective binning, and RoC between bins is calculated. However, the brackets of the time bin (window) are then moved forward by a selected amount of time (Z), levels are selected again, and RoC calculated for the new set of WUs. This is repeated X times (where X is the bin size divided by Z) while retaining all the results. R-Ratepol currently provides several options for selecting WU, namely as individual levels, binning of levels, selective binning, and our new method of binning with a moving window.

2.1.1.2. Randomisations

Due to the inherent statistical errors in uncertainties in the age estimates from age-depth and the assemblage datasets (e.g. pollen counts in each level; Birks and Gordon, 1985), R-Ratepol can be run several times and the results summarised (Steps 5-6). Therefore, two optional settings are available by using age uncertainties and assemblage data standardisation.

Age uncertainties (Step 4a)

For each run, a single age sequence from the age uncertainties is randomly selected. The calculation between two consecutive WUs (i.e. one working-unit combination) results in a *RoC score* and a *time position* (which is calculated as the mean age position of the two WUs). However, due to random sampling of the age sequence, each WU combination will result in multiple RoC values and age positions. R-Ratepol assigns the age position of each WU combination as the median time position from all calculations (Step 6).

The final RoC value for a single WU combination is calculated as the median of the scores from all

randomisations. In addition, the 95[th] quantile from all randomisations is calculated as an error

estimate.

Data standardisation (Step 4b)

Taxa in the assemblage dataset can be standardised to a certain count (e.g. number of pollen grains

in each WU) by rarefaction. Random sampling without replacement is used to draw a selected number

of individuals from each WU (e.g. 150 pollen grains).

2.1.2. Detection of peak-points in RoC sequence (Step 8)

A rapid change in composition or relative abundances of taxa within the sequence can provide a means

of comparing RoC between sequences and interpreting the potential drivers of assemblage change.

To detect such significant peak-points of RoC scores in each sequence, each point is tested to see if it

represents a significant increase in RoC values. There are various ways to detect peak-points in a time

series and R-Ratepol is able detect peak-points using five methods:

1)   Threshold: Each point in the RoC sequence is compared to a median of all RoC scores from the

whole sequence (i.e. *threshold value*). The point is considered significant if the 95[th] quantile

of the RoC scores from all calculations is higher than the *threshold value*.

2)   Linear trend: linear model is fitted between the RoC values and their ages. Differences

between the model and each point are calculated (residuals). The standard deviation (SD) is

calculated from all the residuals. A peak is considered significant if it is 1.5 SD higher than the

model.

3)   Non-linear trend: A conservative generalised additive model (GAM) is fitted through the RoC

scores and their ages ($\mathrm{GAM} = \mathrm{RoC} \sim s(age, k = 3)$) using the *mgcv* package (Wood, 2011).

The distance between each point and the fitted value is calculated (residuals). The standard

deviation (SD) is calculated from all the residuals. A peak is considered significant if it is 1.5 SD

higher than the model.

4)   F-deriv GAM: smooth GAM model is fitted to the RoC scores and their ages ($\mathrm{GAM} =$

$\mathrm{RoC} \sim s(age)$. The first derivative as well as continuous confidence intervals are calculated

from the model using the *gratia* package (Simpson, 2019). A peak is considered significant if

the confidence intervals of the first derivative differ from 0 (for more information see

Simpson, 2018).

5)   Signal-to-noise (SNI) method: We adapted SNI from Kelly et al. (2011), which was developed

to detect changes in charcoal stratigraphical records. SNI is calculated for the whole RoC

sequence and a peak-point is considered significant if it has an SNI value higher than 3.

2.2. Testing the successful detection of peak-points using simulated data

In order to compare the performance of various methods of estimating RoC, we used simulated pollen-stratigraphical data (Blaauw et al., 2010) with known patterns of compositional change and temporal resolution to test the success of peak-point detection in the expected time period of compositional change. In addition, we used a generalised linear mixed modelling approach to evaluate the effects of the different R-Ratepol parameters.

2.2.1. Data simulation

We simulated pollen datasets following Blaauw *et al.* (2010) based on generating pollen-assemblage data in response to known changes in environmental conditions and compositional properties (Fig. S1). We used the following process of data generation: (i) we used the density of levels from a European pollen sequence, Glendalough (Sequence A) as a template for the number of levels and the corresponding ages for each level. (ii) Pollen data were created as abundance data with a total of 300 pollen grains in each level. (iii) We then adjusted the total values of each simulated taxon so that the pollen assemblage resembled a log-scale rank distribution. (iv) We added jitter to each of the pollen taxa (function *jitter*, factor = 1.5). All changes were made in order to make the simulated data more similar to data from a real-life study.

Two additional settings were altered during the simulation of datasets: (i) richness – *low richness* (LR) datasets contain 5 pollen taxa, *high richness* (HR) datasets contain 50 pollen taxa; and (ii) position of change in external environmental properties – *recent* (R) has a sudden increase of environmental properties at 2000 calibrated years before present (1950 CE: cal yr BP) and a decrease at 3000 cal yr BP, *late* (L) has a sudden increase at 5500 and a decrease at 6500 cal yr BP (Fig. S1). This results in two abrupt changes in pollen composition in each dataset. Note that the different timings of the change in environmental properties (hereafter called 'focal area') were selected to illustrate the effects of level density (datasets with the *later* change have a low density of levels within the focal area and datasets with a *recent* change have a high density). The combination of these two settings results in four types of simulated datasets: *low richness – recent change* (LR-R), *low richness – late change* (LR-L), *high richness – recent change* (HR-R), and *high richness – late change* (HR-L).

2.2.2. Using the simulated datasets to test analytical performance of RoC methods with different settings

To investigate the differences between the various settings in RoC methods, we simulated 100 datasets for each dataset type (LR-R, LR-L, HR-R, HR-L; see previous section), and compared the success rate of detecting peak-points in the expected time period for all combinations of (i) WU selection, (ii) smoothing method, (iii) dissimilarity coefficient, and (iii) peak-point detection.

We tested three types of WU selection (individual stratigraphical levels, selective binning, and binning

with moving window), five types of smoothing methods (none, Shepard's 5-term filter, moving

average, age-weighted average, Grimm's smoothing), two types of dissimilarity coefficients

appropriate for compositional data (Chord distance, Chi-squared coefficient), and five types of peak-

point detection (Threshold, Linear trend, Non-linear trend, f-deriv GAM, SNI).

For each simulated dataset we calculated RoC using combinations of all WU selections, data

smoothing, dissimilarity coefficient, and peak-point detection method (150 different settings in total).

R-Ratepol has been set up with standardisation of pollen assemblage data to 150 pollen grains in each

WU and 100 randomisations per calculation. The number of randomisations is not higher due to high

computational demands (60,000 RoC calculations, each with 100 randomisations). Randomisations

are in place only for taxa standardisation as no age uncertainties have been used in this example. We

then extracted (i) the number of WUs where R-Ratepol detected a significant peak-point during the

focal area (± 500 yr), and (ii) the number of samples, where R-Ratepol detected a peak-point during a

time of no expected change ('false positives'). We transformed these numbers of peak-points into

ratios of points detected to total points in the area (inside or outside focal area; see Fig. S2 for example

of peak-point detection).

Statistical tests to evaluate RoC options
First, we tested if the various WU selection methods affect the successful detection of peak-points

within the focal area. We pooled all RoC calculations from all dataset types and created generalised

linear mixed models using the Template Model Builder (Brooks et al., 2017) glmmTMB $_{success-method}$

with the ratio of WU marked as peak-points to all WUs in the focal area ($R_{success-method}$) as the

dependent variable with a beta error distribution. Independent variables are: WU (3 level factor),

peak-point detection method (5 level factor), dataset type (4 level factor), and all their interactions.

Individual dataset ID and RoC setting (factor combining smoothing method and DC; 10 levels) were

selected as the random factors. We then used the *dredge* function from the *MuMIn* package (Barton,

2020) to fit models with all possible combinations of predictors (with a constraint that dataset type

must be present), and ordered the models by parsimony (assessed by corrected Akaike information

criterion; AIC_C). We only selected the best model if ΔAIC_C (i.e. *delta* in the *MuMIn* package) < 2. In

the case where multiple models had similar parsimony, we selected the best model using the

compare_performance function from the *performance* package (Ludecke et al., 2020). We then used

the *emmens* package (Russell, 2020) to obtain estimated marginal means and the 95[th] quantile of the

independent variables from the final model. Similarly, we built glmmTMB $_{FalsePositive-method}$ using

the ratio of WUs marked as peak-points to all WUs outside the focal area ($R_{FalsePositives-method}$).

Next, for a more detailed exploration of different settings in R-Ratepol, we divided the data to only

include the best performing WU selection and peak-point detection combination (i.e. binning with

moving window and non-linear trend peak-point detection, see Results) and created

glmmTMB $_{success-detail}$ and glmmTMB $_{FalsePositive-detail}$ with $R_{success-detail}$ and

$R_{FalsePositives-detail}$ as the dependent variables, respectively. Both models were fitted with a beta

error distribution and the independent variables are data-smoothing type (5 level factor), dissimilarity

coefficient (2 level factor), position of environmental change (i.e. density of levels; 2 level factor),

richness of dataset (2 level factor), and all their interactions. Individual dataset ID was selected as a

random factor. As before, the *dredge* and *compare_performance* functions were used to select the

best model and reduce unnecessary predictors.

2.3. Examples of rate-of-change results using empirical palynological data

After assessing the differences between the individual settings of R-Ratepol based on the results from

the simulated datasets, we examined the RoC estimates for real palynological data representing

different pollen richness and composition, and different density of levels. We obtained pollen data

from the Neotoma database (Williams et al., 2018) using the *Neotoma R package* (Goring et al., 2015).

We chose four European sequences (A–D; Fig 2).

In each sequence, taxa were harmonised to the taxonomically highest pollen morphotype (Level =

MHVar2) using the pollen harmonisation table in Giesecke et al. (2019). To develop age-depth models,

we used the pre-selected radiometric control points provided in Giesecke et al. (2014) and calibrated

the radiocarbon dates using the IntCal13 Northern Hemisphere calibration curve (Reimer et al., 2013).

For each sequence, we constructed an age-depth model using the *Bchron* R package (Haslett and

Parnell, 2008) to generate 1000 possible age estimates for all sample levels at the sampling resolution

of the original pollen sequences. We used these 1000 draws to build posterior estimates of age

uncertainty. We calculated the median age estimate for each sample level to obtain the default age

used in subsequent analyses.

In each sequence, we excluded all levels that contained less than 150 pollen counts of terrestrial taxa,

and all levels beyond a 3000-year extrapolation of the oldest chronological control point. In addition,

we excluded all levels with an age older than 8500 cal yr BP to ensure a focus on the period with

detectable human impact.

Sequence A is from Glendalough, Ireland (53°00'10.0"N 6°22'09.0"W; Neotoma dataset id = 17334;

Haslett et al., 2006). It has a very uneven distribution of levels (N = 102) with the highest density

around 2000 cal yr BP (Fig. 2A). The sequence contains 80 pollen taxa with abundant *Corylus*, *Quercus,*

Alnus, Betula, and Poaceae. Cyperaceae, Poaceae, and Ericales all increase in the last 1000 cal yr BP,

preceded by a period with high values of *Quercus*, *Betula*, and *Alnus* until ~ 5000 cal yr BP. Before 5000

cal yr BP the sequence has large variations in *Pinus*, an increase of *Alnus*, and a large decrease in

Corylus. **Sequence B** is from Dallican Water, Scotland (60°23'14.5"N 1°05'47.3"W; Neotoma dataset

id = 4012; Bennett et al., 1992). It has a relatively even distribution of levels (N = 63; Fig. 2B) and

contains 50 pollen taxa with Ericales, *Betula*, and Poaceae being the most abundant. The pollen record

shows sudden increases of Ericales after 4000 cal yr BP. **Sequence C** is from Steerenmoos, Germany

(47°48'20.0"N 8°12'01.6"E; Neotoma dataset id = 40951; Rösch, 2000). It is an example of a very

detailed sequence, but with an uneven distribution of levels (N = 273; Fig. 2C). The sequence contains

103 pollen taxa with *Abies*, *Fagus*, *Corylus*, *Betula*, and *Quercus* being the most abundant. As with

Sequence A, the pollen stratigraphy can be separated into three major parts: i) a recent period until

1000 cal yr BP typified by high values of *Pinus*; ii) 2000–6000 cal yr BP typified by *Abies* and *Fagus*; and

iii) 6000 cal yr BP to the base with abundant *Betula*, *Corylus*, and *Quercus*. **Sequence D** is from Alanen

Laanijärvi, a boreal-forest lake in Swedish Lapland (67°58'N 20°29'W; Neotoma dataset id = 45314;

Heinrichs et al., 2005). It is an example of a sequence with 54 levels across the time of interest,

containing 44 pollen taxa with high abundance of *Pinus* and *Betula* throughout the whole sequence

(Fig. 2D).

We calculated RoC scores for the four selected European sequences (A-D) using all three methods of

WU selection with Shepard's 5-term filter smoothing and Chi-squared coefficient as the dissimilarity

coefficient. A non-linear trend peak-point detection was used in each sequence, the number of

randomisations was set to 10,000, and the size of time bin was selected as 500 yr (see Fig. S3 for an

example of changes of RoC values with size of time bin). In addition, to provide examples of the

differences between different settings of R-Ratepol, we explored data from sequence A using binning

with the moving window method of WU selection and calculated RoC scores for all combinations of

five smoothing methods and two dissimilarity coefficients (Chi-squared, Chord distance).

3. RESULTS

3.1. Comparison of success rates in peak-point detection for simulated datasets

Successful detection of peak-points ($R_{success-method}$) and the number of incorrectly (false positives)

detected peak-points ($R_{FalsePositives-method}$) are both significantly affected by WU selection and the

method of peak-point detection (Fig. 3; see Table S1 and Table S2 for the AIC_C of all models). For

detecting peak-points successfully within the focal area, our approach of binning with a moving

window results in an overall better performance in successful peak-point detection than the use of

individual levels or binning (effect of successful detection increases by 68% and 20%, respectively; Fig.

3A). Simulations show that the non-linear trend method performs best in detecting peak-points within

the focal area (14% higher than the second-best, the linear trend method; Fig. 3B), and detects a

similar amount of false positives compared to the other peak-point detection methods (with the

exception of SNI, which detects 10-fold less false positives).

While using only the binning with the moving-window approach and the non-linear trend method for

peak-point detection, the successful detection of peak-points ($R_{success-detail}$) and the number of

false positives ($R_{FalsePositives-detail}$) are both influenced by the density of levels, type of dissimilarity

coefficient, and data-smoothing type (Fig. 4; see Tables S3, S4, S5, and S6 for the best selected

models). Using Chi-squared coefficient as the dissimilarity measure results in a 10% increase in the

estimated correctly detected peak-points than Chord distance on average (Fig. 4) but the coefficient

intervals of their estimates overlap (Fig. S4A). The level of smoothing has, on average, a decreasing

effect on the successful detection of peak-points, with data without smoothing being the most

successful (10% higher effect than the second-best Shepard's 5-term filter smoothing; Fig. S4B). The

number of false positives is affected in a similar way, with no smoothing and Shepard's 5-term filter

having the lowest number of false positives. The datasets with an assemblage change which occurred

during a time period characterised by a high density of levels (i.e. recent environmental change) show,

on average, a 14-fold higher success of peak-point detection and a 7-fold higher false positive

detection than datasets with change at the time of a low density of levels (i.e. environmental change

later in the sequence; Fig. S4C). Richness of pollen does not show a significant effect on the detection

of peak-points (note that the assemblage data standardisation was used for all computation).

3.2. Examples of rate-of-change results using real palynological data

The selection of WUs affects not only the overall shape of the RoC curve in all sequences but also the

timing of the period of significant increase in RoC (Fig. 2). Binning with a moving window detects

significant periods of increased RoC scores at sequence A (Glendalough) from 4900–4200 cal yr BP;

sequence B (Dallican Water) from 3500–3000 cal yr BP; sequence C (Steerenmoos) from 6400–5800

cal yr BP; and sequence D (Alanen Laanijärvi) from 550–350 cal yr BP (Fig. 2). The binning method

shows relatively similar patterns and position of peak-points (except for sequence D). However, the

error intervals (95[th] quantile of RoC scores) are much wider. Using individual levels as WUs (Fig. 2.

'RoC-levels') results in very different positions of peak-points and a different shape of the RoC curve

with wide error intervals.

4. DISCUSSION

Based on the results of our simulations, the choice of methods and parameters is critical in the analysis of rate-of-change. Comparison of methods using simulated datasets shows that selection of WUs, peak-point detection method, dissimilarity coefficient, smoothing of the data, and the density of the levels in the period of environmental change significantly influence the detection of sudden changes in stratigraphical sequences. Our proposed method of binning with a moving window in combination with a non-linear trend peak-point detection method improves such detection of an increased RoC. The method shows a 68% increase in the effect of correct detection of peak-points compared to the more traditional approach of selection by individual levels. Peak-point detection using a non-linear trend provides a reasonable compromise between statistical error Type I (linear trend and threshold method) and Type II (SNI method).

There is little difference between using Chi-squared coefficient or Chord's distance as the dissimilarity coefficient for pollen compositional data, and we therefore recommend using both for palaeoecological data sequences that are expressed as percentages or proportions and that contain many zero values. However, R-Ratepol is created as modular tool and can be used for many different paleo-proxies present in sedimentary sequences. Euclidean distance (see Supplementary Material) is appropriate for absolute assemblage data such as pollen concentrations or pollen accumulation rates (e.g. Peglar, 1993; Peglar and Birks, 1993). Such data may require a log-transformation prior to calculating the Euclidean distance between levels or WUs. Standardised Euclidean distance (see Supplementary Material) is appropriate when the variables in an assemblage are expressed in different units, such as geochemical, isotope, and sediment variables (e.g. Bakke et al., 2009; Birks and Birks, 2006; Nesje et al., 2014)). Gower's (1971) distance is appropriate for assemblage data consisting of different numerical types (continuous quantitative, qualitive, binary) such as plant macrofossil data (e.g. Birks, 2014).

Recommendation of a specific smoothing technique of data is less straightforward. There is a trade-off between the potential loss of information by smoothing the data and the incorrect attribution of noise (for example from sampling error) as signal. In our simple simulation experiment, data with a relatively low amount of smoothing (Shepard's 5-term filter and data without smoothing) perform relatively well, independently of the density of samples (Fig. 3). All the other smoothing methods perform worse, especially if environmental change occurred during a period characterised by a low density of levels. However, detailed exploration of the RoC pattern in a real pollen sequence (sequence A; Fig. 5) shows that all smoothing methods result in a similar RoC curve and position. Therefore, there is a need for critical evaluation in future studies to assess the risks of using data smoothing, depending on the density of samples and research questions of interest.

A multitude of appropriate data-smoothing approaches (Wilkinson, 2005), dissimilarity coefficients

(e.g. Legendre and Legendre, 2012; Prentice, 1980), techniques for creating WUs, and methods of

peak-point detection (e.g. Simpson, 2018) are potentially available. Of the two selected dissimilarity

methods and five smoothing algorithms, all commonly used in palaeoecology, we conclude that

datasets with a high density of levels tend to have a higher chance of successfully detecting peak-

points. Anderson et al. (2020) reached similar conclusions in a regional-scale synthesis of fossil pollen

data from California. Here the authors suggest maintaining a consistent temporal spacing within

records, and the use of probabilistic models explicitly incorporating age-model uncertainties to

increase the precision of age estimates of each level. We show that our method of binning with a

moving window yields even better results than the traditional use of individual levels or binning, and

has an advantage here, as age-model uncertainties can be incorporated.

RoC analysis has several applications in palaeoecological research. It is a useful numerical tool to

detect patterns in stratigraphical data that cannot readily be seen by visual inspection (see Jacobson

et al., 1987 for an example at Gould Pond, Maine). It is also useful to compare the rates of change in

different proxies (pollen, diatoms, chironomids, etc.) studied in the same stratigraphical sequence

(e.g. Birks and Ammann, 2000; Szabó et al., 2020). Moreover, it is most useful when RoC results are

compared from several sequences (e.g. Grimm and Jacobson, 1992; Jacobson et al., 1987). Consistent

patterns in RoC peaks in several sequences may potentially indicate responses to extrinsic drivers such

as regional climate change, pathogenic attacks, or widespread human activity. Ecologists are

recognising the importance of quantifying RoC of extrinsic environmental drivers such as temperature,

toxins, and salinity on ecosystems (Pinek et al., 2020) and are developing new and powerful numerical

tools for space–time analysis of community compositional data over time intervals of decades (e.g.

Legendre and Gauthier, 2014). RoC peaks unique to individual sequences may, in contrast, reflect local

factors such as sequence hiatuses, sediment reworking, or spatially restricted human impact or natural

disturbance.

A connection between RoC and local factors can be seen in the sequences used in this study (Fig. 2).

Sequence A (Glendalough) shows significant peak-points at the time of the onset of human activity

and the expansion of grassland and heath (Haslett et al., 2006). Sequence B (Dallican Water) has

significant peak-points at times when major changes in land-use occurred as the island was abandoned

by humans and then reinhabited 800 years later (Bennett et al., 1992). Sequence C (Steerenmoos) has

several significant peak-points at the time of the expansion of *Abies* pollen in the late Neolithic period

and with frequent fires (Rösch, 2000). Sequence D (Alanen Laanijärvi) has significant peak-points in

the very recent period, possibly associated with changes in organic deposition, fertilisation, and timber

harvesting (Heinrichs et al., 2005). Here we can assume that our approach of binning with a moving

window is closest to the expected pattern, further supported by the relatively high, successful
detection of peak-points in the simulated datasets (Fig. 3). In contrast, the traditional use of individual
levels fails to detect the periods of significant change, or falsely assigns peak-points to other, non-
relevant periods, or a combination of both (Fig 2). Some of these errors occur, as expected, with a low
density of sample levels. These sequences and the RoC outcomes exemplify the usefulness of
quantifying rates of compositional change estimated through time.

In summary, we have developed a framework for the robust estimation of rate-of-change in
stratigraphical time-ordered sequences of terrestrial and marine palaeoecological datasets. Our
overall framework consists of four major parts:

1. Establishing a robust age-depth model for the stratigraphical sequence of interest with age
uncertainties for all individual levels.
2. Selecting consecutive working units prior to RoC estimation to allow for uneven temporal
distribution of the analysed levels within the sequence using selective binning with a moving
window.
3. Estimating compositional dissimilarity between working units.
4. Detecting statistically significant peak-points in the RoC estimates within the sequence.

Given the various choices of coefficients, methods, and evaluations in RoC analysis, it is essential when
presenting RoC results for stratigraphical assemblage data to document what choices were made.
There are, as we have shown, many decisions in RoC analysis. Publishing plots of 'Rate of Change' with
no explanation about what choices were made in the RoC analysis is to be strongly discouraged (cf.
Abrook et al., 2019).

Our RoC analysis approach as implemented in R-Ratepol is a significant improvement over existing
methods for RoC analysis that do not incorporate methods to detect statistically significant peak-
points (Birks, 2012). R-Ratepol is, we believe, a powerful and much needed addition to the toolkit of
robust numerical techniques available to palaeoecologists and palaeolimnologists for detecting and
summarising patterns in complex multivariate time-ordered stratigraphical sequences (Birks, 2010,
1997).

5. ACKNOWLEDGEMENTS
All the authors are supported by the European Research Council under the EU Horizon 2020 Research
and Innovation Programme (grant 741413 HOPE) Humans on Planet Earth – Long-term impacts on
biosphere dynamics. HJBB is indebted to John Line (University of Cambridge) for collaboration in 1990

in the development of the Fortran77 RATEPOL program, which was a starting point for R-Ratepol, to

Cathy Jenks for indispensable editorial help, and to Hilary Birks for many helpful discussions. We are

indebted to Gavin Simpson for valuable comments on an earlier manuscript. Data were obtained from

the Neotoma Paleoecology Database (www.neotoma.org) and its constituent European Pollen

Database (EPD). The work of the data contributors, data stewards, and the Neotoma and EPD

community is gratefully acknowledged.

6. AUTHORS' CONTRIBUTION

501   OM and HJBB conceived the ideas and designed the study. OM led the manuscript writing, data

502   analyses, and interpretation of the results. OM, AWRS, KPB, VAF, and SGAF designed the methodology

503   for extracting and processing palaeoecological sequences from Neotoma. OM, JAG, AWRS, MJS, and

504   HJBB contributed to methodology development. HJBB acquired the funding. All authors contributed

505   to the writing of the text and interpretation of the results and gave final approval for publication.

7. DATA AVAILABILITY

R-Ratepol is available as an R package at GITHUB (https://github.com/HOPE-UIB-BIO/R-Ratepol-

package), including all the data used in this study. Scripts for all analyses used in this study can be

found at GITHUB repository (https://github.com/HOPE-UIB-BIO/RateOfChange).

8. REFERENCES

Abrook, A.M., Matthews, I.P., Milner, A.M., Candy, I., Palmer, A.P., Timms, R.G.O., 2019.
Environmental variability in response to abrupt climatic change during the Last Glacial-
Interglacial Transition (16-8 cal ka BP): Evidence from Mainland, Orkney. Scottish J. Geol. 56,
30–46. https://doi.org/10.1144/sjg2019-006

Anderson, L., Wahl, D.B., Bhattacharya, T., 2020. Understanding rates of change: A case study using
fossil pollen records from California to assess the potential for and challenges to a regional data
synthesis. Quat. Int. https://doi.org/10.1016/j.quaint.2020.04.044

Bakke, J., Lie, O., Heegaard, E., Dokken, T., Haug, G.H., Birks, H.H., Dulski, P., Nilsen, T., 2009. Rapid
oceanic and atmospheric changes during the Younger Dryas cold period. Nat. Geosci. 2, 202–
205. https://doi.org/10.1038/ngeo439

Barton, K., 2020. MuMIn: Multi-Model Inference. R package version 1.43.17.

Bennett, K.D., Boreham, S., Sharp, M.J., Switsur, V.R., 1992. Holocene History of Environment,
Vegetation and Human Settlement on Catta Ness, Lunnasting, Shetland. J. Ecol. 80, 241.
https://doi.org/10.2307/2261010

Bennett, K.D., Humphry, R.W., 1995. Analysis of late-glacial and Holocene rates of vegetational change
at two sites in the British Isles. Rev. Palaeobot. Palynol. 85, 263–287.
https://doi.org/10.1016/0034-6667(94)00132-4

Birks, H.H., Ammann, B., 2000. Two terrestrial records of rapid climatic change during the Glacial-
Holocene transition (14,000-9,000 calendar years B.P.) from Europe. Proc. Natl. Acad. Sci. U. S.
A. 97, 1390–1394.

Birks, H.H., Birks, H.J.B., 2006. Multi-proxy studies in palaeolimnology. Veg. Hist. Archaeobot. 15, 235–
251. https://doi.org/10.1007/s00334-006-0066-6

Birks, H.J.B., 2014. Challenges in the presentation and analysis of plant-macrofossil stratigraphical
data. Veg. Hist. Archaeobot. 23, 309–330. https://doi.org/10.1007/s00334-013-0430-2

Birks, H.J.B., 2012. Analysis of Stratigraphical Data, in: Birks, H.J.B., Lotter, A.F., Juggins, S., Smol, J.P.
(Eds.), Tracking Environmental Change Using Lake Sediments Volume 5: Data Handling and
Numerical Techniques. Springer Science & Business Media, Dordrecht, pp. 355–378.

Birks, H.J.B., 2010. Numerical methods for the analysis of diatom assemblage data, in: Smol, J.P.,
Stoermer, E.F. (Eds.), The Diatoms: Applications for the Environmental and Earth Sciences.
Cambridge University Press, Cambridge, pp. 23–54.

Birks, H.J.B., 1997. Environmental change in Britain- a long-term paleaoecological perspective, in:
Mackay, A.W., Murlis, J. (Eds.), Britain's Natural Environment: A State of the Nation Review.
ENSIS Publications, London, UK, pp. 23–28.

Birks, H.J.B., Birks, H.H., 2008. Biological responses to rapid climate change at the Younger Dryas-
Holocene   transition   at   Kråkenes,   western   Norway.   The   Holocene   18,   19–30.
https://doi.org/10.1177/0959683607085572

Birks, H.J.B., Felde, V.A., Seddon, A.W.R., 2016. Biodiversity trends within the Holocene. The Holocene
26, 994–1001. https://doi.org/10.1177/0959683615622568

Birks, H.J.B., Gordon, A.D., 1985. Numerical Methods in Quaternary Pollen Analysis. Academic Press,
London.

Blaauw, M., Bennett, K.D., Christen, J.A., 2010. Random walk simulations of fossil proxy data. The
Holocene 20, 645–649. https://doi.org/10.1177/0959683609355180

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., Skaug, H.J.,
2017. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized
Linear Mixed Modeling. R J. 9, 378–400. https://doi.org/10.3929/ethz-b-000240890

Correa-Metrio, A., Bush, M.B., Cabrera, K.R., Sully, S., Brenner, M., Hodell, D.A., Escobar, J., Guilderson,
T., 2012. Rapid climate change and no-analog vegetation in lowland Central America during the
last 86,000 years. Quat. Sci. Rev. 38, 63–75. https://doi.org/10.1016/j.quascirev.2012.01.025

Davis, J.C., 1986. Statistics and Data Analysis in Geology, 2nd edn. ed. J. Wiley & Sons, New York.

Dodd, M.E., Silvertown, J., McConway, K., Potts, J., Crawley, M., 1994. Stability in the plant
communities of the Park Grass Experiment: The relationships between species richness, soil pH
and   biomass   variability.   Philos.   Trans.   R.   Soc.   B   Biol.   Sci.   346,   185–193.
https://doi.org/10.1098/rstb.1994.0140

Dornelas, M., Magurran, A.E., Buckland, S.T., Chao, A., Chazdon, R.L., Colwell, R.K., Curtis, T., Gaston,
K.J., Gotelli, N.J., Kosnik, M.A., McGill, B., McCune, J.L., Morlon, H., Mumby, P.J., Øvreås, L.,
Studeny, A., Vellend, M., 2013. Quantifying temporal change in biodiversity: Challenges and
opportunities. Proc. R. Soc. B Biol. Sci. 280, 20121931. https://doi.org/10.1098/rspb.2012.1931

Feeley, K.J., Bravo-Avila, C., Fadrique, B., Perez, T.M., Zuleta, D., 2020. Climate-driven changes in the
composition   of   New   World   plant   communities.   Nat.   Clim.   Chang.   10,   965–970.
https://doi.org/10.1038/s41558-020-0873-2

Gibson-Reinemer, D.K., Sheldon, K.S., Rahel, F.J., 2015. Climate change creates rapid species turnover
in montane communities. Ecol. Evol. 5, 2340–2347. https://doi.org/10.1002/ece3.1518

Giesecke, T., Davis, B., Brewer, S., Finsinger, W., Wolters, S., Blaauw, M., de Beaulieu, J.L., Binney, H.,
Fyfe, R.M., Gaillard, M.J., Gil-Romera, G., van der Knaap, W.O., Kuneš, P., Kühl, N., van Leeuwen,
J.F.N., Leydet, M., Lotter, A.F., Ortu, E., Semmler, M., Bradshaw, R.H.W., 2013. Towards
mapping the late Quaternary vegetation change of Europe. Veg. Hist. Archaeobot. 23, 75–86.
https://doi.org/10.1007/s00334-012-0390-y

Giesecke, T., Wolters, S., van Leeuwen, J.F.N., van der Knaap, P.W.O., Leydet, M., Brewer, S., 2019.
Postglacial change of the floristic diversity gradient in Europe. Nat. Commun. 10, 1–7.
https://doi.org/10.1038/s41467-019-13233-y

Goring, S., Dawson, A., Simpson, G.L., Ram, K., Graham, R.W., Grimm, E.C., Williams, J.W., 2015.
Neotoma: A programmatic interface to the Neotoma paleoecological database. Open Quat. 1,
1–17. https://doi.org/10.5334/oq.ab

Gotelli, N.J., Dorazio, R.M., Ellison, A.M., Grossman, G.D., 2010. Detecting temporal trends in species
assemblages with bootstrapping procedures and hierarchical models. Philos. Trans. R. Soc. B
Biol. Sci. 365, 3621–3631. https://doi.org/10.1098/rstb.2010.0262

Gower, J.C., 1971. A general coefficient of similarity and some of its properties. Biometrics 27, 857–
871.

Grimm, E.C., Jacobson, G.L., 1992. Fossil-pollen evidence for abrupt climate changes during the past
18000 years in eastern North America. Clim. Dyn. 6, 179–184.

Grindean, R., Tanţău, I., Feurdean, A., 2019. Linking vegetation dynamics and stability in the old-
growth forests of Central Eastern Europe: Implications for forest conservation and
management. Biol. Conserv. 229, 160–169. https://doi.org/10.1016/j.biocon.2018.11.019

Haslett, J., Parnell, A., 2008. A simple monotone process with application to radiocarbon-dated depth
chronologies. J. R. Stat. Soc. Ser. C Appl. Stat. 57, 399–418. https://doi.org/10.1111/j.1467-
9876.2008.00623.x

Haslett, J., Whiley, M., Bhattacharya, S., Salter-Townshend, M., Wilson, S.P., Allen, J.R.M., Huntley, B.,
Mitchell, F.J.G., 2006. Bayesian palaeoclimate reconstruction. J. R. Stat. Soc. Ser. A Stat. Soc.
169, 395–438. https://doi.org/10.1111/j.1467-985X.2006.00429.x

Heinrichs, M., Peglar, S.M., Bigler, C., Birks, H.J.B., 2005. A multi-proxy palaeoecological study of
602     Alanen Laanijärvi, a boreal-forest lake in Swedish Lapland. Boreas 34, 192–206.
603     https://doi.org/10.1080/03009480510012881

604 Hillebrand, H., Blasius, B., Borer, E.T., Chase, J.M., Downing, J.A., Eriksson, B.K., Filstrup, C.T., Harpole,
605     W.S., Hodapp, D., Larsen, S., Lewandowska, A.M., Seabloom, E.W., Van de Waal, D.B., Ryabov,
606     A.B., 2018. Biodiversity change is uncoupled from species richness trends: Consequences for
607     conservation and monitoring. J. Appl. Ecol. 55, 169–184. https://doi.org/10.1111/1365-
608     2664.12959

609 Jacobson, G.L., Grimm, E.C., 1986. A numerical analysis of Holocene forest and prairie vegetation in
610     central Minnesota. Ecology 67, 958–966. https://doi.org/10.2307/1939818

611 Jacobson, G.L., Webb, T., Grimm, E.C., 1987. Patterns and rates of vegetation change during the
612     deglaciation of eastern North America, in: Ruddiman, W.F., Wright, H.E. (Eds.), North America
613     and Adjacent Oceans During the Last Deglaciation. Geological Society of America, Boulder,
614     Colorado, pp. 277–288. https://doi.org/10.1130/dnag-gna-k3.277

615 Kelly, R.F., Higuera, P.E., Barrett, C.M., Feng Sheng, H., 2011. A signal-to-noise index to quantify the
616     potential for peak detection in sediment-charcoal records. Quat. Res. 75, 11–17.
617     https://doi.org/10.1016/j.yqres.2010.07.011

618 Kemp, D.B., Eichenseer, K., Kiessling, W., 2015. Maximum rates of climate change are systematically
619     underestimated in the geological record. Nat. Commun. 6, 1–6.
620     https://doi.org/10.1038/ncomms9890

621 Laird, K.R., Fritz, S.C., Cumming, B.F., 1998. A diatom-based reconstruction of drought intensity,
622     duration, and frequency from Moon Lake, North Dakota: A sub-decadal record of the last 2300
623     years. J. Paleolimnol. 19, 161–179. https://doi.org/10.1023/A:1007929006001

624 Legendre, P., Gauthier, O., 2014. Statistical methods for temporal and space-time analysis of
625     community composition data. Proc. R. Soc. B Biol. Sci. 281, 20132728.
626     https://doi.org/10.1098/rspb.2013.2728

627 Legendre, P., Legendre, L., 2012. Numerical Ecology, 3rd Englis. ed. Elsevier, Amsterdam.
628     https://doi.org/10.1016/B978-0-12-409548-9.10595-0

629 Lotter, A.F., Ammann, B., Sturm, M., 1992. Rates of change and chronological problems during the
630     late-glacial period. Clim. Dyn. 6, 233–239. https://doi.org/10.1007/BF00193536

Ludecke, D., Makowski, D., Waggoner, P., Patil, I., Ben-Shachar, M.S., 2020. Assessment of Regression
Models Performance. CRAN.R-Project.org.

Magurran, A.E., Baillie, S.R., Buckland, S.T., Dick, J.M.P., Elston, D.A., Scott, E.M., Smith, R.I.,
Somerfield, P.J., Watt, A.D., 2010. Long-term datasets in biodiversity research and monitoring:
Assessing change in ecological communities through time. Trends Ecol. Evol. 25, 574–582.
https://doi.org/10.1016/j.tree.2010.06.016

Magurran, A.E., Dornelas, M., Moyes, F., Henderson, P.A., 2019. Temporal β diversity—A
macroecological   perspective.   Glob.   Ecol.   Biogeogr.   28,   1949–1960.
https://doi.org/10.1111/geb.13026

Monchamp, M.E., Spaak, P., Domaizon, I., Dubois, N., Bouffard, D., Pomati, F., 2018. Homogenization
of lake cyanobacterial communities over a century of climate change and eutrophication. Nat.
Ecol. Evol. 2, 317–324. https://doi.org/10.1038/s41559-017-0407-0

Morecroft, M.D., Bealey, C.E., Beaumont, D.A., Benham, S., Brooks, D.R., Burt, T.P., Critchley, C.N.R.,
Dick, J., Littlewood, N.A., Monteith, D.T., Scott, W.A., Smith, R.I., Walmsley, C., Watson, H., 2009.
The UK Environmental Change Network: Emerging trends in the composition of plant and
animal communities and the physical environment. Biol. Conserv. 142, 2814–2832.
https://doi.org/10.1016/j.biocon.2009.07.004

Nesje, A., Bakke, J., Brooks, S.J., Kaufman, D.S., Kihlberg, E., Trachsel, M., D'Andrea, W.J., Matthews,
J.A., 2014. Late glacial and Holocene environmental changes inferred from sediments in Lake
Myklevatnet, Nordfjord, western Norway. Veg. Hist. Archaeobot. 23, 229–248.
https://doi.org/10.1007/s00334-013-0426-y

Peglar, S.M., 1993. The Mid-Holocene Ulmus decline at Diss Mere, Norfolk, UK: a year-by- year pollen
stratigraphy   from   annual   laminations.   The   Holocene   3,   1–13.
https://doi.org/10.1177/095968369300300101

Peglar, S.M., Birks, H.J.B., 1993. The mid-Holocene Ulmus fall at Diss Mere, South-East England -
disease   and   human   impact?   Veg.   Hist.   Archaeobot.   2,   61–68.
https://doi.org/10.1007/BF00202183

Pinek, L., Mansour, I., Lakovic, M., Ryo, M., Rillig, M.C., 2020. Rate of environmental change across
scales in ecology. Biol. Rev. 95, 1798-1811. https://doi.org/10.1111/brv.12639

Prentice, I.C., 1980. Multidimensional scaling as a research tool in Quaternary palynology: A review of

theory and methods. Rev. Palaeobot. Palynol. 31, 71–104. https://doi.org/10.1016/0034-

6667(80)90023-8

R Core Team, 2018. R: A language and environment for statistical computing.

Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H.,

Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas, I., Hatté, C.,

Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W.,

Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van

der Plicht, J., 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 years

cal BP. Radiocarbon 55, 1869–1887. https://doi.org/10.2458/azu_js_rc.55.16947

Richardson, A.J., Walne, A.W., John, A.W.G., Jonas, T.D., Lindley, J.A., Sims, D.W., Stevens, D., Witt,

M., 2006. Using continuous plankton recorder data. Prog. Oceanogr. 68, 27–74.

https://doi.org/10.1016/j.pocean.2005.09.011

Rösch, M., 2000. Long-term human impact as registered in an upland pollen profile from the southern

Black Forest, south-western Germany. Veg. Hist. Archaeobot. 9, 205–218.

https://doi.org/10.1007/BF01294635

Russell, L., 2020. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version

1.4.8.

Seddon, A.W.R., Macias-Fauria, M., Willis, K.J., 2015. Climate and abrupt vegetation change in

Northern Europe since the last deglaciation. The Holocene 25, 25–36.

https://doi.org/10.1177/0959683614556383

Shuman, B., Bartlein, P.J., Webb, T., 2005. The magnitudes of millennial- and orbital-scale climatic

change in eastern North America during the Late Quaternary. Quat. Sci. Rev. 24, 2194–2206.

https://doi.org/10.1016/j.quascirev.2005.03.018

Silvertown, J., Poulton, P., Johnston, E., Edwards, G., Heard, M., Biss, P.M., 2006. The Park Grass

Experiment 1856-2006: Its contribution to ecology. J. Ecol. 94, 801–814.

https://doi.org/10.1111/j.1365-2745.2006.01145.x

Simpson, G.L., 2019. gratia: graceful'ggplot'-based graphics and other functions for GAMs fitted using

"mgcv." R Packag. version 0.2-1.

Simpson, G.L., 2018. Modelling palaeoecological time series using generalised additive models. Front.

Ecol. Evol. 6, 1–21. https://doi.org/10.3389/fevo.2018.00149

Solovieva, N., Jones, V., Birks, H.J.B., Appleby, P.G., Nazarova, L., 2008. Diatom responses to 20th

century climate warming in lakes from the northern Urals, Russia. Palaeogeogr. Palaeoclimatol.

Palaeoecol. 259, 96–106. https://doi.org/10.1016/j.palaeo.2007.10.001

Steffen, W., Broadgate, W., Deutsch, L., Gaffney, O., Ludwig, C., 2015. The trajectory of the

Anthropocene: The Great Acceleration. Anthr. Rev. 2, 81–98.

https://doi.org/10.1177/2053019614564785

Steinbauer, M.J., Grytnes, J.A., Jurasinski, G., Kulonen, A., Lenoir, J., Pauli, H., Rixen, C., Winkler, M.,

Bardy-Durchhalter, M., Barni, E., Bjorkman, A.D., Breiner, F.T., Burg, S., Czortek, P., Dawes, M.A.,

Delimat, A., Dullinger, S., Erschbamer, B., Felde, V.A., Fernández-Arberas, O., Fossheim, K.F.,

Gómez-García, D., Georges, D., Grindrud, E.T., Haider, S., Haugum, S. V., Henriksen, H., Herreros,

M.J., Jaroszewicz, B., Jaroszynska, F., Kanka, R., Kapfer, J., Klanderud, K., Kühn, I., Lamprecht, A.,

Matteodo, M., Di Cella, U.M., Normand, S., Odland, A., Olsen, S.L., Palacio, S., Petey, M., Piscová,

V., Sedlakova, B., Steinbauer, K., Stöckli, V., Svenning, J.C., Teppa, G., Theurillat, J.P., Vittoz, P.,

Woodin, S.J., Zimmermann, N.E., Wipf, S., 2018. Accelerated increase in plant species richness

on mountain summits is linked to warming. Nature 556, 231–234.

https://doi.org/10.1038/s41586-018-0005-6

Stephens, L., Fuller, D., Boivin, N., Rick, T., Gauthier, N., Kay, A., Marwick, B., Geralda, C., Armstrong,

D., Barton, C.M., Denham, T., Douglass, K., Driver, J., Janz, L., Roberts, P., Rogers, J.D., Thakar,

H., Altaweel, M., Johnson, A.L., Vattuone, M.M.S., Aldenderfer, M., Archila, S., Artioli, G., Bale,

M.T., Beach, T., Borrell, F., Braje, T., Buckland, P.I., Jiménez Cano, N.G., Capriles, J.M., Castillo,

A.D., Çilingiroglu, Ç., Cleary, M.N., Conolly, J., Coutros, P.R., Covey, R.A., Cremaschi, M.,

Crowther, A., Der, L., Di Lernia, S., Doershuk, J.F., Doolittle, W.E., Edwards, K.J., Erlandson, J.M.,

Evans, D., Fairbairn, A., Faulkner, P., Feinman, G., Fernandes, R., Fitzpatrick, S.M., Fyfe, R.,

Garcea, E., Goldstein, S., Goodman, R.C., Guedes, J.D., Herrmann, J., Hiscock, P., Hommel, P.,

Horsburgh, K.A., Hritz, C., Ives, J.W., Junno, A., Kahn, J.G., Kaufman, B., Kearns, C., Kidder, T.R.,

Lanoë, F., Lawrence, D., Lee, G.A., Levin, M.J., Lindskoug, H.B., López-Sáez, J.A., Macrae, S.,

Marchant, R., Marston, J.M., McClure, S., McCoy, M.D., Miller, A.V., Morrison, M.,

Matuzeviciute, G.M., Müller, J., Nayak, A., Noerwidi, S., Peres, T.M., Peterson, C.E., Proctor, L.,

Randall, A.R., Renette, S., Schug, G.R., Ryzewski, K., Saini, R., Scheinsohn, V., Schmidt, P.,

Sebillaud, P., Seitsonen, O., Simpson, I.A., Soltysiak, A., Speakman, R.J., Spengler, R.N., Steffen,

M.L., Storozum, M.J., Strickland, K.M., Thompson, J., Thurston, T.L., Ulm, S., Ustunkaya, M.C.,

Welker, M.H., West, C., Williams, P.R., Wright, D.K., Wright, N., Zahir, M., Zerboni, A., Beaudoin,

E., Garcia, S.M., Powell, J., Thornton, A., Kaplan, J.O., Gaillard, M.J., Goldewijk, K.K., Ellis, E.,

2019. Archaeological assessment reveals Earth's early transformation through land use.

Science. 365, 897–902. https://doi.org/10.1126/science.aax1192

Szabó, Z., Buczkó, K., Haliuc, A., Pál, I., L. Korponai, J., Begy, R.C., Veres, D., Luoto, T.P., Zsigmond, A.R.,

Magyari, E.K., 2020. Ecosystem shift of a mountain lake under climate and human pressure: A

move out from the safe operating space. Sci. Total Environ. 743, 1–17.

https://doi.org/10.1016/j.scitotenv.2020.140584

Urrego, D.H., Bush, M.B., Silman, M.R., Correa-Metrio, A.Y., Ledru, M.-P., Mayle, F.E., Paduano, G.,

Valencia, B.G., 2009. Millennial-scale ecological changes in tropical South America since the Last

Glacial Maximum, in: Vimeaux, F., Sylvestre, F., Khodri, M. (Eds.), Past Climate Variability in

South America and Surrounding Regions. Springer, Dordrecht, pp. 283–300.

https://doi.org/10.1007/978-90-481-2672-9_12

Wilkinson, L., 2005. The Grammar of Graphics. Springer-Verlag, New York, USA 37.

https://doi.org/10.2307/2669493

Williams, J.W., Grimm, E.C., Blois, J.L., Charles, D.F., Davis, E.B., Goring, S.J., Graham, R.W., Smith, A.J.,

Anderson, M., Arroyo-Cabrales, J., Ashworth, A.C., Betancourt, J.L., Bills, B.W., Booth, R.K.,

Buckland, P.I., Curry, B.B., Giesecke, T., Jackson, S.T., Latorre, C., Nichols, J., Purdum, T., Roth,

R.E., Stryker, M., Takahara, H., 2018. The Neotoma Paleoecology Database, a multiproxy,

international, community-curated data resource. Quat. Res. 89, 156–177.

https://doi.org/10.1017/qua.2017.105

Wolfe, D.A., Champ, M.A., Flemer, D.A., Mearns, A.J., 1987. Long-term biological data sets: Their role

in research, monitoring, and management of estuarine and coastal marine systems. Estuaries

10, 181–193. https://doi.org/10.2307/1351847

Wood, S.N., 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation of

semiparametric generalized linear models. J. R. Stat. Soc. Ser. B Stat. Methodol. 73, 3–36.

https://doi.org/10.1111/j.1467-9868.2010.00749.x

Figure 1. Schematic visualisation of RoC calculation in R-Ratepol with binning with a moving
window. Methodological retails are provided in section 2.1 and Supplementary Material.

Figure 2. Summary of four sequences used in this study. Glendalough (A), Dallican Water (B),

Steerenmoos (C), and Alanen Laanijärvi (D). For each sequence, three different plots are shown.

First plot (left) shows the density of levels within each sequence for the last 8000 cal yr BP. Second

plot (middle) shows pollen stratigraphies for the most common taxa in the sequences. Values are presented as proportions of pollen grains in each level. Last plot (right) shows the rate-of-change score calculated with R-Ratepol for three different methods for WU selection: RoC-levels = use of individual levels, RoC-binning = binning, RoC-MW = binning with moving window. Shepard's 5-term filter smoothing and Chi-squared coefficient were selected for smoothing and dissimilarity measure, respectively. Green points indicate significant peak-points detected using the non-linear trend method. For a detailed explanation of the methods, see Materials and Methods, and Supplementary Material.

Figure 3. Comparison of the correct detection of peak-points in simulated datasets. A) Comparison of estimated marginal means of peak-point detection rates between different WU selections. Working unit selection: levels = use of subsequent levels, bins = selective binning, MW = binning with moving window. B) Comparison of estimated marginal means of peak-point detection rates between different peak-point detection methods. GAM first deriv = first derivative of a GAM; SNI = signal-to-noise index.

Figure 4. Comparison of peak-point detection rates using binning with the moving window and non-linear peak detection method. Smoothing methods: None = data without smoothing, Shep = Shepard's 5-term filter, M_avg = moving average, Age_w = age-weighted average, Grimm = Grimm's smoothing. Dissimilarity coefficients: Chord = Chord distance, Chisq = Chi-squared coefficient. For a detailed explanation of the methods, see Supplementary Material.

Figure 5. Rate-of-change (RoC) and peak-points detected in Sequence A (Glendalough) for five categories of data smoothing (columns) on two dissimilarity coefficients (rows). The binning with moving window was selected as Working Units, with 500 yr bin and 5 window shifts. Green points indicate significant peak-points detected using the non-linear trend method. Smoothing methods: None = data without smoothing, Shep = Shepard's 5-term filter, M_avg = moving average, Age_w = age-weighted average, Grimm = Grimm's smoothing. Dissimilarity coefficients: Chord = Chord distance, Chisq = Chi-squared coefficient. For a detailed explanation of the methods, see Supplementary Material.