Title:

Length–length and length–weight relationships for four small pelagic fishes in the Kuroshio–

Oyashio current system

Running title:

Morphometric relationships of small pelagic fishes

Authors:

Sho Furuichi[1]*, Yasuhiro Kamimura[1], Ryuji Yukami[1]

Affiliations:

[1]Fisheries Resources Institute, Japan Fisheries Research and Education Agency, Yokohama,

Japan

*Correspondence:

Sho Furuichi, Fisheries Resources Institute, Japan Fisheries Research and Education Agency,

Yokohama, Japan

Email: sfuru@affrc.go.jp

Summary

Length–length relationships and length–weight relationships were estimated for four small

pelagic fishes (Japanese sardine *Sardinops melanostictus*, Japanese anchovy *Engraulis*

japonicus, chub mackerel *Scomber japonicus*, and spotted mackerel *Scomber australasicus*)

in the Kuroshio–Oyashio current system. Fish samples were collected from surface–midwater

trawl surveys and commercial purse-seine fisheries between September and October of 2020

in the western North Pacific. Total length (TL), fork length (FL), and standard length (SL)

were measured to 0.01 cm, and whole body weight (W) was measured to the nearest 0.01 g

for each individual. All length–length relationships (TL–FL, TL–SL, and FL–SL) were highly

significant ($p < 0.0001$), with $r^2 > 0.98$ in all species. Length–weight relationships (W–TL,

W–FL, and W–SL) were also highly significant ($p < 0.0001$), with $r^2 > 0.98$ in all species.

This study provides a useful reference for biological studies and stock assessments of these

small pelagic fishes.

1. INTRODUCTION

Length–length relationships (LLRs) and length–weight relationships (LWRs) of fishes are key

tools in fish biology and ecology, fisheries science, and fisheries management (Froese, 2006;

Froese et al., 2011). Length–length relationships are needed to convert length measurements

from one length type into another and are also important in comparative growth studies.

Length–weight relationships are used to predict weight from length and to calculate

production and biomass of a fish population in stock assessment.

In the Kuroshio–Oyashio current system (around the Japanese Archipelago), the small

pelagic fish community is dominated by Japanese sardine *Sardinops melanostictus* Temminck

& Schlegel, 1844, Japanese anchovy *Engraulis japonicus* Temminck & Schlegel, 1846, chub

mackerel *Scomber japonicus* Houttuyn, 1782, and spotted mackerel *Scomber australasicus*

Cuvier, 1832. The ecological and commercial importance and remarkable fluctuations in the

abundances of these small pelagic fishes have inspired studies ranging in scope from the

dynamics of single stocks to the dynamics of small pelagic fish communities at ecosystem and

global scales (Chavez, 2003; Oozeki et al., 2019). Although numerous researchers from

different institutions and disciplines have studied the same ecosystem, and even the same fish

stocks, these researchers have often used different standards for fish length measurement (i.e.,

standard length [SL], fork length [FL], or total length [TL]) (e.g., Jung et al., 2008; Kang et

al., 2009; Yukami et al., 2008). Therefore, conversion equations (that is, LLRs) for these

various length measurement standards are required for comparative or integrative research

among different geographic regions and species. However, there is currently little information

on LLRs for the four small pelagic fishes in the Kuroshio–Oyashio current system. In

addition, although LWRs for these fishes have been reported, studies have been limited to

specific geographic regions and length-measurement standards.

In this study, we provide LLRs and LWRs among TL, FL, and SL for the four small

pelagic fish species (*S. melanostictus*, *E. japonicus*, *S. japonicus*, and *S. australasicus*) that

dominate the Kuroshio–Oyashio current system.

2. MATERIALS AND METHODS

Specimens were obtained from a trawl survey and from commercial purse-seine catches. In

the trawl survey, 35 surface–midwater trawls were conducted in the western North Pacific off

northern Japan and the Kuril Islands (38°20′N–46°58′N, 142°8′E–174°50′E) during

September–October 2020. A surface–midwater trawl net (opening dimensions: 25 m × 25 m;

cod-end mesh size: 17.5 mm) was towed once at each station (15–60 min duration at 3.5–4.0

kn [1.8–2.1 m/s], depths 0–30 m). At each station, fish species were identified on board, and

up to 380 individuals of each species were randomly selected and stored at –25 °C. Samples

from purse-seine catches were collected from commercial landings at the Hachinohe Port (an

important fishing port in northern Japan) in October 2020. All sampled fish were measured to

the nearest 0.01 cm in standard length (SL), fork length (FL), and total length (TL) in the

laboratory. Total weight was measured to the nearest 0.01 g.

To estimate LLRs, linear relationships between the various measurement types (TL, FL,

and SL in cm) were derived by using major-axis (MA) regression. MA regression is used to

find the axis that minimizes errors in both measured variables, and is particularly useful for

comparative and integrative study as it allows bi-directional conversions.

LWRs were estimated by using the equation: $W = a \times L^b$, where W is total body

weight (g), L is length (TL, FL, or SL in cm), a is the coefficient of proportionality, and b is

the coefficient of allometry. Parameters a and b were estimated by non-linear least-squares

regression. Additionally, 95% confidence intervals of a and b, and the coefficient of

determination (r^2) were calculated. All statistical analyses were performed in R version 3.6.3

(R Development Core Team, 2020).

3. RESULTS

Sample sizes, body-length and body-weight ranges, and estimated LWR and LLR parameters for each fish species are presented in Tables 1, 2, and 3. All LLRs were highly significant ($p <$ 0.0001), with r^2 values >0.98. All LWRs were also highly significant ($p < 0.0001$), with r^2 values >0.94.

4. DISCUSSION

In the present study, we estimated LLRs for *S. melanostictus*, *E. japonicus*, *S. japonicus*, and *S. australasicus*. This is the first comprehensive report of LLRs for these four small pelagic fish species in the Kuroshio–Oyashio current system. Although these species have been studied in various parts of the world, previous studies employed a variety of standards for fish length measurements, making comparisons between and among studies difficult. Our results will be helpful for comparative or integrative research among different geographic regions and species.

In addition to LLRs, we also estimated LWRs for the four species. The allometric coefficients (*b*) of the LWRs were within the expected range of 2.5–3.5 (Froese, 2006), although there was some variability among species. This indicates that our estimates are plausible. However, we were not able to sample specimens from across a broad temporal and spatial range. LWRs can be affected by various factors such as season and location (Jellyman et al., 2013; Valle et al., 2020). Therefore, a broader sampling scheme in future studies could provide more robust LWRs.

110   In conclusion, our results provide new information on LLRs and LWRs for four small

111   pelagic fish species in the Kuroshio–Oyashio current system. These data will be useful for

112   further biological study and management of the four species.

ACKNOWLEDGEMENTS

We are very grateful to Michiyo Ichikawa, Aki Narisawa, Akiko Sugawara, Yuko Saiki, and

Tomiko Nagai, who greatly contributed to the data collection. We thank all those who helped

in the collection of samples. We also thank English-speaking professional editors from ELSS,

Inc., for English proofreading. Funding for this study was provided by the Japan Fisheries

Research and Education Agency and the Fisheries Agency of Japan.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

DATA AVALABILITY STATEMENT

The data used to support the findings of this study are available from the corresponding

author upon request.

REFERENCES

Chavez, F. P. (2003). From Anchovies to Sardines and Back: Multidecadal Change in the

Pacific Ocean. *Science*, *299*(5604), 217–221. https://doi.org/10.1126/science.1075880

Froese, R. (2006). Cube law, condition factor and weight-length relationships: History, meta-

analysis and recommendations. *Journal of Applied Ichthyology*, *22*(4), 241–253.

https://ñoi.org/10.1111/j.1439-0426.2006.00805.x

Froese, R., Tsikliras, A. C., & Stergiou, K. I. (2011). Editorial note on weight-length relations

of fishes. *Acta Ichthyologica et Piscatoria*, *41*(4), 261–263.

https://doi.org/10.3750/AIP2011.41.4.01

Jellyman, P. G., Booker, D. J., Crow, S. K., Bonnett, M. L., & Jellyman, D. J. (2013). Does

one size fit all? An evaluation of length-weight relationships for New Zealand's

freshwater fish species. *New Zealand Journal of Marine and Freshwater Research*,

47(4), 450–468. https://doi.org/10.1080/00288330.2013.781510

Jung, S., Hwang, S. Do, & Kim, J. (2008). Fecundity and growth-dependent mortality of

Pacific anchovy (*Engraulis japonicus*) in Korean coastal waters. *Fisheries Research*,

93(1–2), 39–46. https://doi.org/10.1016/j.fishres.2008.02.004

Kang, D., Cho, S., Lee, C., Myoung, J. G., & Na, J. (2009). Ex situ target-strength

measurements of Japanese anchovy (*Engraulis japonicus*) in the coastal Northwest

Pacific. *ICES Journal of Marine Science*, *66*(6), 1219–1224.

https://doi.org/10.1093/icesjms/fsp042

Oozeki, Y., Ñiquen Carranza, M., Takasuka, A., Ayón Dejo, P., Kuroda, H., Tam Malagas, J.,

Okunishi, T., Vásquez Espinoza, L., Gutiérrez Aguilar, D., Okamura, H., & Guevara

Carrasco, R. (2019). Synchronous multi-species alternations between the northern

Humboldt and Kuroshio Current systems. *Deep-Sea Research Part II: Topical Studies in*

156  *Oceanography*, *159*, 11–21. https://doi.org/10.1016/j.dsr2.2018.11.018

157 R Development Core Team. (2020). *R: A language and environment for statistical computing*.

158  R Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/

159 Valle, M., Rosas-Puchuri, U., & Velez-Zuazo, X. (2020). Improving data deficiencies in

160  length-weight relationships for fish species around an artificial breakwater and adjacent

161  soft-bottom at the central coast of Peru. *Journal of Applied Ichthyology*, Early View.

162  https://doi.org/10.1111/jai.14089

163 Yukami, R., Aoki, I., & Mitani, I. (2008). Daily age of adult Japanese anchovy *Engraulis*

164  *japonicus* off eastern Honshu, Japan by otolith daily increment. *Fisheries Science*, *74*(6),

165  1348–1350. https://doi.org/10.1111/j.1444-2906.2008.01665.x

168 Table 1. Sample sizes and ranges of body length and weight of four small pelagic fish species

169 caught in the western North Pacific.

Species	*n*	TL (cm)		FL (cm)		SL (cm)		W (g)	
		Min	Max	Min	Max	Min	Max	Min	Max
Sardinops melanostictus (Temminck & Schlegel, 1844)	1809	12.5	25.2	11.4	22.6	10.9	21.5	12.9	142.4
Engraulis japonicus (Temminck & Schlegel, 1846)	664	4.7	16.2	4.3	15.4	4.0	14.4	0.43	28.6
Scomber japonicus (Houttuyn, 1782)	3682	6.3	46.1	5.9	43.0	5.3	39.8	1.6	874.9
Scomber australasicus (Cuvier, 1832)	312	15.7	34.4	14.6	32.2	13.6	29.4	31.2	369.5

170 Abbreviations: *n*, sample size; TL, total length; FL, fork length; SL, standard length; W, total

171 body weight.

174 Table 2. Estimated length–length relationship parameters for four small pelagic fish species in

175 the western North Pacific.

Species	Equation	a (95% CI)	b (95% CI)	r^2
Sardinops melanostictus (Temminck & Schlegel, 1844)	TL = a + b (FL)	–0.241 (–0.327 to –0.156)	1.09 (1.09 to 1.10)	0.99
	TL = a + b (SL)	0.371 (0.295 to 0.447)	1.13 (1.12 to 1.13)	0.99
	FL = a + b (SL)	0.559 (0.485 to 0.633)	1.03 (1.03 to 1.04)	0.99
Engraulis japonicus (Temminck & Schlegel, 1846)	TL = a + b (FL)	0.121 (0.064 to 0.178)	1.07 (1.06 to 1.08)	0.99
	TL = a + b (SL)	0.358 (0.307 to 0.409)	1.12 (1.11 to 1.13)	0.99
	FL = a + b (SL)	0.221 (0.167 to 0.276)	1.05 (1.04 to 1.05)	0.99
Scomber japonicus (Houttuyn, 1782)	TL = a + b (FL)	–0.169 (–0.218 to –0.121)	1.08 (1.07 to 1.08)	0.99
	TL = a + b (SL)	–0.072 (–0.123 to –0.021)	1.16 (1.16 to 1.17)	0.99
	FL = a + b (SL)	0.092 (0.054 to 0.131)	1.08 (1.08 to 1.08)	0.99
Scomber australasicus (Cuvier, 1832)	TL = a + b (FL)	0.009 (–0.300 to 0.312)	1.07 (1.05 to 1.08)	0.98
	TL = a + b (SL)	0.320 (0.026 to 0.609)	1.13 (1.12 to 1.15)	0.98
	FL = a + b (SL)	0.294 (0.079 to 0.507)	1.06 (1.05 to 1.07)	0.99

176 Abbreviations: TL, total length; FL, fork length; SL, standard length; CI, confidence interval;

177 r^2, coefficient of determination.

181 Table 3. Estimated length–weight relationship parameters for four small pelagic fish species

182 in the western North Pacific.

Species	Equation	a (95% CI)	b (95% CI)	r^2
Sardinops melanostictus (Temminck & Schlegel, 1844)	$W = a \times TL^b$	0.002 (0.002 to 0.003)	3.45 (3.41 to 3.48)	0.94
	$W = a \times FL^b$	0.002 (0.002 to 0.003)	3.54 (3.50 to 3.58)	0.94
	$W = a \times SL^b$	0.004 (0.004 to 0.005)	3.4 (3.37 to 3.44)	0.95
Engraulis japonicus (Temminck & Schlegel, 1846)	$W = a \times TL^b$	0.004 (0.003 to 0.004)	3.2 (3.16 to 3.24)	0.98
	$W = a \times FL^b$	0.006 (0.006 to 0.007)	3.08 (3.04 to 3.12)	0.97
	$W = a \times SL^b$	0.008 (0.007 to 0.008)	3.10 (3.06 to 3.13)	0.98
Scomber japonicus (Houttuyn, 1782)	$W = a \times TL^b$	0.006 (0.006 to 0.006)	3.11 (3.10 to 3.12)	0.99
	$W = a \times FL^b$	0.007 (0.007 to 0.007)	3.13 (3.12 to 3.14)	0.99
	$W = a \times SL^b$	0.009 (0.009 to 0.010)	3.12 (3.12 to 3.13)	0.99
Scomber australasicus (Cuvier, 1832)	$W = a \times TL^b$	0.005 (0.004 to 0.006)	3.20 (3.14 to 3.27)	0.96
	$W = a \times FL^b$	0.006 (0.005 to 0.007)	3.19 (3.13 to 3.25)	0.96
	$W = a \times SL^b$	0.006 (0.005 to 0.007)	3.27 (3.22 to 3.32)	0.97

183 Abbreviations: W, total body weight; TL, total length; FL, fork length; SL, standard length;

184 CI, confidence interval; r^2, coefficient of determination.