Population dynamics and ranging behaviours of provisioned silvered langur (*Trachypithecus cristatus*) in Peninsular Malaysia

Norlinda Mohd-Daut[1], Ikki Matsuda[2,3,4,5], Kamaruddin Zainul Abidin[6], and Badrul Munir Md-Zain[1*]

[1]*Department of Biological Sciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia*

[2]*Chubu University Academy of Emerging Sciences, 1200, Matsumoto-cho, Kasugai-shi, Aichi 487-8501, Japan*

[3]*Wildlife Research Center of Kyoto University, 2-24 Tanaka-Sekiden-cho, Sakyo, Kyoto 606-8203, Japan*

[4]*Japan Monkey Centre, 26 Kanrin, Inuyama, Aichi 484-0081, Japan*

[5]*Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Locked Bag 2073, 88999 Kota Kinabalu, Sabah, Malaysia*

[6]*Faculty of Applied Science, Universiti Teknologi Mara Pahang, 26400, Bandar Tun Abdul Razak, Pahang, Malaysia*

*Correspondence: E-mail: abgbadd1966@yahoo.com

Running head: Silvered langur population dynamics and ranging behaviours

Abstract

Tourists are attracted to the Bukit Melawati Kuala Selangor (BMKS) of Peninsular Malaysia,

a small hill park, both for its status as a historical site and the free-ranging silvered langurs

(*Trachypithecus cristatus*) that come for provisioning. We assessed the population trends and

group sizes of *T. cristatus* over 10 years in the BMKS and examined their ranging patterns.

Comparisons of observed populations between 2005 (190 individuals) and 2017 (193

individuals) revealed the stable demography and group sizes of the six *T. cristatus* groups in

the BMKS. Based on a total of 185 location points of the six groups in 2017, their mean

ranging area was 3.6 ha with a range of 0.86 to 6.93 ha with extensive spatial overlap. We

also found a significant positive relationship between the six groups' ranges and group sizes

in 2017. Additionally, qualitative ecological comparisons with a previous study on *T. cristatus*

in 1965 (before provisioning) suggest that the artificial food supply in the study area could

modify the population dynamics and socioecology of *T. cristatus*. The modifications might

alter their range size and territoriality in the BMKS. Overall, we found that provisioning had

negative effects on the ecology of *T. cristatus* in the BMKS. Therefore, modifying

management policies, such as banning feeding and implementing educational programs, may

contribute to their proper conservation.

Keywords: Colobinae; lutung; group size; population size, home range; Bukit Melawati

Kuala Selangor

Introduction

The monitoring of animal populations is essential for developing effective management plans for long-term conservation (Wich and Marshall 2016). Primates are long-lived mammals with slow life cycles (Charnov and Berrigan 2005; Ross 1998), so they generally respond slowly to dramatic environmental changes and human interference (Chapman et al. 2006). Therefore, long-term population monitoring data is indispensable for understanding their population dynamics (e.g. Chapman et al. 2007; Matsuda et al. 2020). Knowledge of a species' ranging behaviour could also contribute to a better understanding of its behavioural ecology and habitat requirements, leading to effective conservation strategy designs (Nathan et al. 2008). Among various environmental factors, food distribution and abundance are significant influences on population characteristics, such as demography and density (e.g. Bernard et al. 2019; Chapman et al. 2017; Hanya et al. 2004; Marshall 2010), and ranging in many primate species (e.g. Ampeng and Md-Zain 2012; Clutton-Brock 1975; Green et al. 2020; Hanya et al. 2020; Matsuda et al. 2009; Olupot et al. 1994; Raemaekers 1980; Zhou et al. 2007).

Both direct (e.g. being fed by tourists) and indirect (e.g. garbage eating) human or tourist provisioning affect primate demography (Altmann and Alberts 2003; Borries et al. 2001; Mori 1979; Sugiyama and Ohsawa 1982; Takahata 1980) and ranging behaviours (Altmann and Muruthi 1988; Asquith 1989; Kurita et al. 2008; Md-Zain et al. 2010; Ruslin et al. 2019; Sha and Hanya 2013). Initially, when primates are provisioned, the daily travel distance, and their total ranging area, generally decrease (Altmann and Muruthi 1988; José-Domínguez et al. 2015; Sengupta et al. 2015) since the high nutrient foods provided by humans are available to individuals/groups in specific areas. On the other hand, it has been reported that larger groups of free-ranging primates travel greater distances and have larger

home ranges so that their foraging activity meets the nutritional requirements of all group members (e.g. Makwana 1978; Takasaki 1981); thus, this would theoretically be the case in this population if provisioning is limited. Under any circumstances (with or without provisioning), access to energy-rich foods (artificial or natural) generally leads to higher birth rates or shorter interbirth intervals, potentially inducing larger group sizes and reducing per capita foraging success, which subsequently increases the distances travelled daily (Chapman and Chapman 2000; Janson 1988).

As a result of provisioning, animals are often over-habituated (Russon and Wallis 2014b). Therefore, provisioning can trigger human-animal conflict, which is of great concern to conservationists and wildlife managers. For example, several reported conflicts have involved situations such as monkeys attacking and injuring tourists, possibly infecting tourists and vice versa, and damage to the natural habitat (e.g. Brennan et al. 1985; Fuentes et al. 2007; Kurita 2014; Md-Zain et al. 2011; Palacios et al. 2011; Sha et al. 2009). Consequently, ecotourism that includes free-ranging primates, especially concerning provisioning, should be implemented with caution.

Silvered langurs (*Trachypithecus cristatus*), classed as vulnerable by the IUCN and widely distributed in Southeast Asia (Meijaard and Nijman 2015), are foregut-fermenting colobines (Chivers 1994; Matsuda et al. 2019). The ecological knowledge of *T. cristatus* is limited; however, according to a preliminary study (Hock and Sasekumar 1979), they are predominantly folivorous, with leaves comprising over 90% of their diet.

Tourists are attracted to the Bukit Melawati Kuala Selangor (BMKS), a small hill park in Peninsular Malaysia, due to its status as a historical site, its beautiful offshore scenery and the presence of free-ranging habituated *T. cristatus* as a result of tourist provisioning (e.g.

feeding the animals bread or cookies) (Mohd-Daut 2020). Before provisioning, the *T. cristatus* ranging area was strictly territorial, and the ranges of adjacent groups overlapped by only a few trees in the BMKS (Bernstein 1968). Their home range has been estimated to be 20 ha in size, with 350 m as the mean daily path length (Yeager and Kool 2000). Although it is unclear when tourist provisioning started, the *T. cristatus* in the site appear to be over-habituated as early as the 2000s. They have been observed approaching visitors to beg for food, staring at and touching tourists and stealing food from tourists (Mohd-Daut and Md-Zain 2021); however, the langurs do not initiate aggressive contact. Only one study (Hambali et al. 2016) reported the non-territorial ranging behaviour of one *T. cristatus* group in the site (home range: 1.0 ha; daily path length: 10–200 m), though the information on the ranging patterns in other adjacent groups was not available. Additionally, we have almost no empirical information about their ecology and behaviours in relation to the effects of provisioning in the BMKS based on long-term monitoring.

In the present study, to gain a complete picture of the demography and ranging patterns of the provisioned groups of *T. cristatus* in the BMKS, which possibly contribute to the development of their effective management plans, we sought to (1) evaluate population changes between 2005 and 2017 and (2) estimate the range size of all groups at the site. The effects of provisioning on their demography and ranging behaviour are discussed by qualitatively comparing records from a previous study, conducted before tourist provisioning (Bernstein 1968).

Materials and Methods

We observed six groups of silvered langurs *T. cristatus* from March to April 2005 and from October to December 2017 in a part of the BMKS (3°20' 26.36 N, 101°14'42.46 E), located approximately 62 km from Kuala Lumpur, Malaysia (Hambali et al. 2016). The study area covers ca. 22.5 ha, surrounded by the Kuala Selangor Nature Park, a mangrove forest along the coastline that includes a residential area. Historical structures and monkey habitats in this open area attract visitors and make the BMKS a renowned tourism area in Selangor. Although fragmented, the study area still harbours plants that are important resources in the *T. cristatus* diet, including both natural and ornamental plants (Mohd-Daut and Md-Zain 2021). The mean minimum and maximum temperatures were approximately 26.5°C and 29.8°C, respectively, and total precipitation at the site was 1,505.5 mm (Metereology Department of Malaysia).

We observed that groups with one adult male, multiple adult females and immatures were well-habituated to humans due to the tourist site's popularity. We estimated each individual's approximate age using age/sex categories based on the criteria of Aggimarangsee (2002), Bernstein (1968) and Harding (2010), with some modifications. In 2005, we identified all *T. cristatus* adult, subadult and juvenile individuals in the study groups based on 1) the shape of the face, especially that distinguished by the curve of the cheek fur; 2) the shape of the nose and eyes, which usually had characteristics resembling those of the original parent in the group and 3) any visible evidence of birth defects or injuries, such as scars or limb defects. Although all individuals identified in 2005 were not recognised in 2017, some group members with distinctive features, such as scars and limb defects, were recognisable even in 2017. Notably, all adult males were replaced from 2005 to 2017. Thus, in 2017, the recognisable individuals were mostly females, and the groups including such females were

found in similar areas to those used by the groups they were member of in 2005. In addition to the fact that *T. cristatus* appears to be a species characterised by female philopatry and male-biased dispersal (Wolf and Fleagle 1977), we considered this as evidence that these were the same groups.

At 0800 h each day, we arrived at the study site and followed the groups until they selected evening sleeping trees and started sleeping, usually at 1900 h. We completed 520 and 1,023 hr observations in 2005 and 2017, respectively. Once-daily, we checked the numbers, age class, and sex of group members. On the first day of each observation week, we attempted to record the location points for all study groups at the site using Global Positioning Systems (GPS). The location data was typically recorded once a week. Of the six groups, five were constantly observed in the study area, although one group (group F) was infrequently found. We recorded 185 location points for the six *T. cristatus* groups, with 14 to 34 points per group. We transferred GPS coordinates in decimal degrees and projected them into meters (WGS 1984 UTM zone 48N) using the projection tool in ArcMap version 10.5. Biotas 2.0 Alpha was used to calculate and investigate the ranging area. The most common method, namely Minimum Convex Polygon (MCP) 100% (Boitani and Fuller 2000), was used to analyse each group's home range. Based on the home range areas for each group (calculated by MCP), we also calculated the amount of overlap between the group areas. Google Earth was used to map and estimate the size of the land area used in the study. Areas used by the study groups were divided into three categories: vegetated areas, including all areas covered by trees; grassland and open areas and building/construction areas.

We compared the group size of six one-male-multi-female groups between 2005 and 2017, using a Wilcoxon rank-sum test (Wilcoxon 1945). We examined the relationship

between home range size and the six groups' sizes in 2017, using linear regression analysis. The relationship between the total area size of vegetated patches within each group's ranging areas and their group size was also tested using linear regression analysis. We used the Bonferroni correction to correct for multiple comparisons of those linear regression analyses (P = 0.05/N: N is the number of comparisons made, i.e. 0.025). The significance level was set to α = 0.05. We report means with standard deviation for all results. All statistical analyses were performed in the R statistical computing environment version 4.02 (R Core Development-Team 2019).

Results Population comparison between 2005 and 2017

In 2005, we counted 190 *T. cristatus* in six groups with one male and multiple females at the study site. The group size was almost the same in 2017 (193), though we found an additional all-male group (Table 1). The mean size of groups with one male and multiple females was 31.7 ± 16.5 and 31.5 ± 12.4 individuals in 2005 and 2017, respectively, and the size was not significantly different between the years (W = 17; *p*-value: 0.916). We found that, out of the six one-male-multi-female groups, two groups had more members (i.e. A and F), and two groups maintained their group size (i.e. B and E). In contrast, the other two groups had fewer members (i.e. C and D). Groups A and F experienced an increase of 10 and 2 members, respectively. On the other hand, Group C experienced a decrease of 3 members in the total group size of 24 individuals. Meanwhile, Group D, which consisted of 46 individuals, decreased by 10 members.

Ranging area in 2017

Based on 185 location points of six one-male-multi-female groups, their mean ranging size was 3.6 ± 2.08 ha with a range of 0.86 to 6.93 ha (Table 2 and Fig. 1). The size of the area calculated by MCP using the cumulative location points in each group suggests that we have not fully recorded the total area occupied by the study groups as the area size did not fully reach a plateau for each group (Fig. 2). Nonetheless, within their ranging areas, the most dominant habitat type was the vegetated area (54.8 ± 7.34 ha), followed by the grass/open area (32.7 ± 5.35) and the building/construction area (12.6 ± 3.99), though the percentage of each habitat type was different across the six groups (Table 3).

There was range area overlap among the six one-male-multi-female groups; the mean overlapped area based on all dyadic combinations for six groups was 1.28 ± 1.08 ha (21.9% ± 16.2%) with a range of 0.12–4.39 ha (1.44%–62.7%) (Table 4). The core area, which was used by all six groups, was 0.33 ha (Fig. 1).

The six groups' per capita ranging size differed across the groups with a range of 0.07 to 0.15 ha (Table 2). There was a significant positive relationship between their ranging area and group sizes (Fig. 3; R^2: 0.81; corrected p-value: 0.030). There was also a significant positive relationship between the vegetated area within each group's range and group sizes (Fig. 3; R^2: 0.82; corrected p-value: 0.025).

Discussion

The population size of *T. cristatus* in the BMKS study area was almost the same in 2005 and 2017 (190 and 193 individuals, respectively), despite the addition of an all-male group,

indicating that their numbers have remained stable over the decade. Group sizes in 2017 (27.6) were smaller than in 2005 (31.7), but it was not a significant difference. Although provisioning impacts primate demography in general (Rothman and Bryer 2019), we could not detect significant changes in population or group sizes in the study site. This result suggests that a constant supply of high nutrient artificial foods provided by tourists in the 2000s (Md-Zain et al. 2009) may have had a minimal effect on the demography of the *T. cristatus*. However, it should be noted that the current information is too limited to comprehensively evaluate the provisioning effects, and a further study focusing on the nutritional intake (in terms of calories) of *T. cristatus* in the BMKS is necessary. On the other hand, the density of *T. cristatus* using the area in 2017 was unusually higher (ca. $858/km^2$) than those of *Trachypithecus* spp. in other study sites (Newton and Dunbar 1994): mean value = $101/km^2$; range = $2.1–345/km^2$. Although the density of primates is generally influenced by provisioning and/or habitat quality (e.g. Bernard et al. 2019; Chapman et al. 2017; Hanya et al. 2004; Marshall 2010), the fact that the density at our study site is extremely high does imply that there is an effect of provisioning on population density in our study area.

Although the demography was stable even under the provisioning condition over the decade, it is reasonable to expect the *T. cristatus* population size to increase due to the effects of artificial foods provided by tourists in the 2000s but not in the 1960s in the BMKS (Bernstein 1968). In 1965, the six *T. cristatus* groups, including at least 160 individuals, were reported (Bernstein 1968). Based on these figures, their population size would have increased by ca. 30 individuals (19%) from the 1960s to the 2000s, suggesting that there has been an impact of provisioning on the population size of *T. cristatus*. However, since we only conducted the population survey in a part of the BMKS that tourists can access, additional

groups might inhabit the forested area. A population survey of *T. cristatus* in the entire

BMKS area would allow us to better conclude how tourist provisioning has influenced the

demography over the last five decades.

The artificial food supply in the study area may modify the socioecology of *T.*

cristatus, altering their range size and territoriality in the BMKS. The study group ranging

areas in 2017 (mean of the six groups = 3.6 ha) were much smaller than the 1965 ranging

areas (20 ha: Bernstein 1968; Yeager and Kool 2000), although we cannot deny the

possibility that this effect may have been an artefact of the small sample sizes, in terms of the

number of location data points for each study group. Nonetheless, it should be noted that

MCP, used in this study to calculate the home range sizes, is a more suitable measure of home

range area, especially for a data set with fewer location points (e.g. Grueter et al. 2009;

Robbins 2003). Furthermore, as the study groups sometimes moved out of the area and

briefly inhabited the forested area (NMD personal observations), the real sizes of the study

group home ranges might be underestimated. However, we observed that the *T. cristatus*

groups mostly spent their time in the BMKS study area and even slept there at night, which

would minimize errors in estimating their home range values. It may not be necessary for *T.*

cristatus to forage larger areas to satisfy their nutrient requirements as they could easily

obtain energy-rich artificial foods in the study area. Such behavioural changes reportedly led

to decreases in ranging areas or daily path lengths in other provisioned primate species, e.g.

pig-tailed macaques (José-Domínguez et al. 2015), rhesus macaque (Sengupta et al. 2015),

and baboons (Altmann and Muruthi 1988). Thus, because no artificial or feeding maintenance

occurred in 1965 (Bernstein 1968), the observed change of the *T. cristatus* range sizes in our

study area may be influenced by recent tourist provisioning.

Additionally, the six *T. cristatus* group ranging areas were more concentrated to our study area where human food was accessible and abundant in 2017 than those in 1965 (Bernstein 1968). Thus, the population density in 2017 would be more biased to such a specific tourist area of the BMKS. This bias may explain why the six *T. cristatus* groups in 1965 ranged larger areas and were more territorial, with less spatial overlap, among the groups (Bernstein 1968) compared to those in 2017. In the 1960s, the population might have been near or above carrying capacity, and groups would be territorial at high densities in the BMKS. However, when people started provisioning with high nutrient foods, the population might have dipped below carrying capacity, and the groups would be non-territorial. This explanation also fits the hypothesized theoretical relationship among home range overlap, individual density, and carrying capacity, according to Yeager and Kool's (2000) description of primate territorial behaviours.

Although the ranging behaviours of the study groups would be influenced by the provisioning (i.e. decreased ranging area), there was a significant positive relationship between ranging areas and group sizes of the six *T. cristatus* groups. Theoretically, increases in group size should reduce per capita foraging success (Chapman and Chapman 2000; Janson 1988), i.e. larger groups generally travel greater distances with a larger home range area (e.g. Makwana 1978; Takasaki 1981). We also found a significant positive relationship between vegetated areas within the ranging area of each group and group size, suggesting that provisioned foods may not be sufficient to meet their nutrient requirements. Thus, the larger *T. cristatus* groups would still need to forage natural foods within the larger vegetated areas in their ranging area. In non-human primates, the increase of day range, possibly with increasing group size, has been used as an index to evaluate the degree of food competition

within groups. This hypothesis on weak/absent food competition is supported by the fact that a flat relationship exists between group size and day range across various colobines (Yeager and Kool 2000). Therefore, a significant positive relationship between the group size and ranging area of the six *T. cristatus* groups in this study indicates their high intraspecific competition for food. However, it may be difficult to conclude if the highly competitive foods in this study site are artificial or natural. We suggest that future studies on *T. cristatus* examine their food habits and nutrients as these will be key to understanding the demography in the BMKS and their socioecology in relation to intraspecific competition in folivorous primates.

Overall, we found the negative effects of provisioning on the ecology of the *T. cristatus* in the BMKS, suggesting that the study groups need effective management. Although ecotourism does not always negatively impact primate conservation (Lhota et al. 2019a; Lhota et al. 2019b), it has often failed to deliver conservation benefits for free-ranging primates, especially provisioning (Russon and Wallis 2014a). Therefore, we recommend that local authorities take appropriate actions to make visitors aware that they should stop feeding animals through educational methods explaining the negative impacts of provisioning on the ecology of *T. cristatus* in the BMKS.

Acknowledgments

We appreciate the Department of Wildlife and National Parks Peninsular Malaysia and Kuala Selangor District Council for giving us a permit (JPHL and TN(P): 100-34/1.24 Jld 7(25)) to conduct the study at BMKS. We also appreciate the Metereology Department for providing our climate data, the botanist of the Department of Biological Sciences and Biotechnology,

Faculty Science and Technology Universiti Kebangsaan Malaysia, Ahmad Fitri Zohari, for his help in identifying the types of trees around the study area. This study was partially funded by the Society for the Promotion of Science (KAKENHI Grant Nos. #19KK0191 and # 19H03308 to IM), JSPS Core-to-Core Program, Advanced Research Networks (#JPJSCCA20170005 to S. Kohshima), and the Cooperative Research Program by KUPRI (2016-B-97).

References

Aggimarangsee N (2002) Behavior and ecology of isolated dusky langur (*Trachypithecus obscurus*) population at Khao Lommuak, Prachuap Khiri Khan Province, Thailand. Ph.D., Chiang Mai University, Thailand

Altmann J, Alberts SC (2003) Intraspecific Variability in Fertility and Offspring Survival in a Nonhuman Primate: Behavioral Control of Ecological and Social Sources. In: Wachter KW, Bulatao RA (eds) Offspring: Human Fertility Behavior in Biodemographic Perspective. National Academies Press, Washington (DC), pp 140-169

Altmann J, Muruthi P (1988) Differences in daily life between semiprovisioned and wild-feeding baboons. Am J Primatol 15:213-221. doi: 10.1002/ajp.1350150304

Ampeng A, Md-Zain BM (2012) Ranging patterns of critically endangered colobine, *Presbytis chrysomelas chrysomelas*. ScientificWorldJournal 2012:594382. doi: 10.1100/2012/594382

Asquith PJ (1989) Provisioning and the study of free-ranging primates: History, effects, and prospects. American Journal of Physical Anthropology 32:129-158. doi: 10.1002/ajpa.1330320507

Bernard H, Matsuda I, Hanya G, Phua M-H, Oram F, Ahmad AH (2019) Feeding ecology of the

proboscis monkey in Sabah, Malaysia, with special reference to plant species-poor forests.

In: Nowak K, Barnett AA, Matsuda I (eds) Primates in Flooded Habitats: Ecology and

Conservation. Cambridge University Press, Cambridge, pp 89-98

Bernstein IS (1968) The lutong of Kuala Selangor. Behaviour 32:1-16. doi: 10.1163/156853968x00054

Boitani L, Fuller TK (2000) Research Techniques in Animal Ecology: Controversies and Consequences.

Columbia University Press, New York

Borries C, Koenig A, Winkler P (2001) Variation of life history traits and mating patterns in female

langur monkeys (*Semnopithecus entellus*). Behavioral Ecology and Sociobiology 50:391-402.

doi: 10.1007/s002650100391

Brennan EJ, Else JG, Altmann J (1985) Ecology and behaviour of a pest primate: vervet monkeys in a

tourist-lodge habitat. African Journal of Ecology 23:35-44. doi: 10.1111/j.1365-

2028.1985.tb00710.x

Chapman CA et al. (2017) Primate population dynamics: variation in abundance over space and time.

Biodiversity and Conservation 27:1221-1238. doi: 10.1007/s10531-017-1489-3

Chapman CA, Chapman LJ (2000) Determinants of group size in primates: the importance of travel

costs. In: Boinski S, Garber PA (eds) On the move: how and why animals travel in groups.

University of Chicago Press, Chicago, pp 24-42

Chapman CA, Naughton-Treves L, Lawes MJ, Wasserman MD, Gillespie TR (2007) Population Declines

of Colobus in Western Uganda and Conservation Value of Forest Fragments. International

Journal of Primatology 28:513-528. doi: 10.1007/s10764-007-9142-8

Chapman CA et al. (2006) Do food availability, parasitism, and stress have synergistic effects on red colobus populations living in forest fragments? Am J Phys Anthropol 131:525-534. doi: 10.1002/ajpa.20477

Charnov EL, Berrigan D (2005) Why do female primates have such long lifespans and so few babies? or Life in the slow lane. Evolutionary Anthropology: Issues, News, and Reviews 1:191-194. doi: 10.1002/evan.1360010604

Chivers D (1994) Functional anatomy of the gastrointestinal tract. In: Davies A, Oates J (eds) Colobine monkeys: their ecology, behaviour and evolution. Cambridge University Press, Cambridge, pp 205–227

Clutton-Brock TH (1975) Ranging behaviour of red colobus (*Colobus badius tephrosceles*) in the Gombe National Park. Animal Behaviour 23:706-722. doi: 10.1016/0003-3472(75)90147-5

Fuentes A, Shaw E, Cortes J (2007) Qualitative assessment of macaque tourist sites in Padangtegal, Bali, Indonesia, and the Upper Rock Nature Reserve, Gibraltar. International Journal of Primatology 28:1143-1158. doi: 10.1007/s10764-007-9184-y

Green SJ, Boruff BJ, Niyigaba P, Ndikubwimana I, Grueter CC (2020) Chimpanzee ranging responses to fruit availability in a high-elevation environment. Am J Primatol:e23119. doi: 10.1002/ajp.23119

Grueter CC, Li D, Ren B, Wei F (2009) Choice of analytical method can have dramatic effects on primate home range estimates. Primates 50:81-84. doi: 10.1007/s10329-008-0113-2

Hambali K, Md-Zain BM, Amir A (2016) Daily movement, sleeping sites and canopy level use of habituated silvered-leaf monkeys (*Trachypithecus cristatus*)in Bukit Malawati, Kuala Selangor, Malaysia. Journal of Sustainability Science and Management 11:21-30

Hanya G et al. (2004) Environmental determinants of the altitudinal variations in relative group

densities of Japanese macaques on Yakushima. Ecological Research 19:485-493. doi:

10.1111/j.1440-1703.2004.00662.x

Hanya G, Yoshihiro SI, Hayaishi S, Takahata Y (2020) Ranging patterns of Japanese macaques in the

coniferous forest of Yakushima: Home range shift and travel rate. Am J Primatol 82:e23185.

doi: 10.1002/ajp.23185

Harding LE (2010) *Trachypithecus cristatus* (Primates: Cercopithecidae). Mammalian Species 42:149-

165. doi: 10.1644/862.1

Hock LB, Sasekumar A (1979) A preliminary study on the feeding biology of mangrove forest

primates, Kuala Selangor. Malayan Nature Journal 33.:105–112

Janson CH (1988) Food competition in brown capuchin monkeys (*Cebus apella*): quantitative effects

of group size and tree productivity. Behaviour 105:53-76. doi: 10.1163/156853988x00449

José-Domínguez JM, Huynen M-C, García CJ, Albert-Daviaud A, Savini T, Asensio N (2015) Non-

territorial Macaques Can Range Like Territorial Gibbons When Partially Provisioned With

Food. Biotropica 47:733-744. doi: 10.1111/btp.12256

Kurita H (2014) Provisioning and tourism in free-ranging Japanese macaques. In: Russon AE, Wallis J

(eds) Primate Tourism: A Tool for Conservation? Cambridge University Press, Cambridge, pp

44-55

Kurita H, Sugiyama Y, Ohsawa H, Hamada Y, Watanabe T (2008) Changes in demographic parameters

of *Macaca fuscata* at Takasakiyama in relation to decrease of provisioned foods.

International Journal of Primatology 29:1189-1202. doi: 10.1007/s10764-008-9296-z

Lhota S, Scott KSS, Sha JCM (2019a) Primates in flooded forests of Borneo: opportunities and

challenges for ecotourism as a conservation strategy. In: Barnett AA, Nowak K, Matsuda I

(eds) Primates in Flooded Habitats: Ecology and Conservation. Cambridge University Press, Cambridge, pp 331-340

Lhota S, Sha JCM, Bernard H, Matsuda I (2019b) Proboscis monkey conservation: beyond the science. In: Behie AM, Teichroeb JA, Malone NM (eds) Primate Research and Conservation in the Anthropocene. Cambridge University Press, Cambridge pp 182-196

Makwana SC (1978) Field ecology and behaviour of the rhesus macaque (*Macaca mulatta*): I. Group composition, home range, roosting sites, and foraging routes in the Asarori Forest. Primates 19:483-492. doi: 10.1007/bf02373310

Marshall AJ (2010) Effect of habitat quality on primate populations in Kalimantan: gibbons and leaf monkeys as case studies. In: Gursky S, Supriatna J (eds) Indonesian Primates. Springer New York, New York, pp 157-177

Matsuda I et al. (2020) Population dynamics of the proboscis monkey *Nasalis larvatus* in the Lower Kinabatangan, Sabah, Borneo, Malaysia. Oryx 54:583-590. doi: 10.1017/s0030605318000467

Matsuda I, Chapman CA, Clauss M (2019) Colobine forestomach anatomy and diet. J Morphol 280:1608-1616. doi: 10.1002/jmor.21052

Matsuda I, Tuuga A, Higashi S (2009) Ranging behavior of proboscis monkeys in a riverine forest with special reference to ranging in inland forest. International Journal of Primatology 30:313-325. doi: 10.1007/s10764-009-9344-3

Md-Zain BM, Mohd-Daut N, Md-Nor S (2009) Characterizing silvered leaf monkey-visitor interactions at Bukit Melawati, Kuala Selangor, Malaysia. Journal Of Wildlife And Parks 26:83-94

Md-Zain BM et al. (2010) A Comprehensive Population Survey and Daily Activity Budget on Long-tailed Macaques of Universiti Kebangsaan Malaysia. Journal of Biological Sciences 10:608-615

Md-Zain BM, Tarmizi MR, Mohd-Zaki M (2011) Campus monkeys of Universiti Kebangsaan Malaysia:

nuisance problems and st udents perception. In: Gumert MD, Fuentes A (eds) Monkey on

the Edges. Cambridge University Press, Cambrige, pp 110-117

Meijaard E, Nijman V (2015) *Trachypithecus cristatus*. The IUCN Red List of Threatened Species

2020:e.T22035A17959977. doi: https://dx.doi.org/10.2305/IUCN.UK.2020-

2.RLTS.T22035A17959977.en

Mohd-Daut N (2020) Population size and positional behaviour of silvered leaf monkey

(*Trachypithecus cristatus*) at Bukit Melawati Kuala Selangor. PhD thesis, Universiti

Kebangsaan Malaysia, Selangor, Malaysia

Mohd-Daut N, Md-Zain BM (2021) Notes on diet and anthropogenic food consumption of habituated

silvered langurs (*Trachypithecus cristatus*) at Bukit Melawati, Kuala Selangor, Selangor,

Malaysia. Malayan Nature Journal 73:1-5

Mori A (1979) Analysis of population changes by measurement of body weight in the Koshima troop

of Japanese monkeys. Primates 20:371-397. doi: 10.1007/bf02373390

Nathan R et al. (2008) A movement ecology paradigm for unifying organismal movement research.

Proc Natl Acad Sci U S A 105:19052-19059. doi: 10.1073/pnas.0800375105

Newton PN, Dunbar RIM (1994) Colobine monkey society. In: Davies AG, Oates JF (eds) Colobine

monkeys: Their ecology, behaviour and evolution. Cambridge University Press, Cambridge,

pp 311-346

Olupot W, Chapman CA, Brown CH, Waser PM (1994) Mangabey (*Cercocebus albigena*) population

density, group size, and ranging: A twenty-year comparison. Am J Primatol 32:197-205. doi:

10.1002/ajp.1350320306

Palacios G et al. (2011) Human metapneumovirus infection in wild mountain gorillas, Rwanda. Emerg

Infect Dis 17:711-713. doi: 10.3201/eid1704.100883

R-Core-Development-Team (2019) R: a language and environment for statistical computing.

Foundation for Statistical Computing, Vienna

Raemaekers J (1980) Causes of variation between months in the distance traveled daily of gibbons.

Folia Primatol (Basel) 34:46-60. doi: 10.1159/000155947

Robbins MM (2003) Home range and frugivory patterns of mountain gorillas in Bwindi Impenetrable

National Park, Uganda. International Journal of Primatology 24:467-491. doi:

10.1023/a:1023741328338

Ross C (1998) Primate life histories. Evolutionary Anthropology: Issues, News, and Reviews 6:54-63.

doi: 10.1002/(sici)1520-6505(1998)6:2<54::aid-evan3>3.0.co;2-w

Rothman JM, Bryer MAH (2019) The effects of humans on the primate nutritional landscape. In:

Behie A, Teichroeb JA, Malone N (eds) Primate Research and Conservation in the

Anthropocene. Cambridge University Press, Cambridge pp 199-215

Ruslin F, Matsuda I, Md-Zain BM (2019) The feeding ecology and dietary overlap in two sympatric

primate species, the long-tailed macaque (*Macaca fascicularis*) and dusky langur

(*Trachypithecus obscurus obscurus*), in Malaysia. Primates 60:41-50. doi: 10.1007/s10329-

018-00705-w

Russon AE, Wallis J (2014a) Primate tourism as a conservation tool: a review of the evidence,

implications, and recommendations. In: Russon AE, Wallis J (eds) Primate Tourism: A Tool for

Conservation? Cambridge University Press, Cambridge, pp 313-332

Russon AE, Wallis J (2014b) Reconsidering primate tourism as a conservation tool: an introduction to

the issues. In: Russon AE, Wallis J (eds) Primate Tourism: A Tool for Conservation?

Cambridge University Press, Cambridge, pp 3-18

Sengupta A, McConkey KR, Radhakrishna S (2015) Primates, Provisioning and Plants: Impacts of

Human Cultural Behaviours on Primate Ecological Functions. PLoS One 10:e0140961. doi:

10.1371/journal.pone.0140961

Sha JC, Gumert MD, Lee BP, Jones-Engel L, Chan S, Fuentes A (2009) Macaque-human interactions

and the societal perceptions of macaques in Singapore. Am J Primatol 71:825-839. doi:

10.1002/ajp.20710

Sha JCM, Hanya G (2013) Temporal Food Resource Correlates to the Behavior and Ecology of Food-

Enhanced Long-Tailed Macaques (*Macaca fascicularis*). Mammal Study 38:163–175. doi:

10.3106/041.038.0305

Sugiyama Y, Ohsawa H (1982) Population dynamics of Japanese monkeys with special reference to

the effect of artificial feeding. Folia Primatol (Basel) 39:238-263. doi: 10.1159/000156080

Takahata Y (1980) The reproductive biology of a free-ranging troop of Japanese monkeys. Primates

21:303-329. doi: 10.1007/bf02390462

Takasaki H (1981) Troop size, habitat quality, and home range area in Japanese macaques.

Behavioral Ecology and Sociobiology 9:277-281. doi: 10.1007/bf00299883

Wich SA, Marshall AJ (2016) An Introduction to Primate Conservation. Oxford University Press, New

York

Wilcoxon F (1945) Individual comparisons by ranking methods. Biometrics Bulletin 1:80. doi:

10.2307/3001968

Wolf KE, Fleagle JG (1977) Adult male replacement in a group of silvered leaf-monkeys (*Presbytis cristata*) at Kuala Selangor, Malaysia. Primates 18:949-955. doi: 10.1007/bf02382945

Yeager CP, Kool KM (2000) The behavioral ecology of Asian colobines. In: Whitehead PF, Jolly CJ (eds) Old World Monkeys. Cambridge Cambridge University Press, pp 496-521

Zhou Q, Huang C, Li Y, Cai X (2007) Ranging behavior of the Francois' langur (*Trachypithecus francoisi*) in the Fusui Nature Reserve, China. Primates 48:320-323. doi: 10.1007/s10329-006-0027-9

Table 1 Composition of groups of *Trachypithecus cristatus*, with several one-male-multifemale groups (OMG) and an all-male group (AMG) in 2005 and 2017.

Group ID	Group type	Year	Age and sex categories						
			Adult male	Adult female	Subadult male	Subadult female	Juvenile	Infant	Total
A	OMG	2005	1	12	0	1	3	2	19
		2017	1	14	1	4	7	2	29
B	OMG	2005	1	22	3	2	2	3	33
		2017	1	21	2	2	4	3	33
C	OMG	2005	1	13	4	3	3	3	27
		2017	1	18	1	1	1	2	24
D	OMG	2005	1	8	7	11	24	5	56
		2017	1	32	0	7	2	4	46
E	OMG	2005	1	28	1	1	7	6	44
		2017	1	33	2	4	1	3	44
F	OMG	2005	1	7	0	3	0	0	11
		2017	1	5	1	2	2	2	13
G	AMG	2005	-	-	-	-	-	-	-
		2017	4	0	0	0	0	0	4

Table 2 Range area and per capita ranging area of the study *T. cristatus* groups

Group ID	Total group members	Ranging area (ha)	Per capita ranging area (ha)
A	29	2.15	0.07
B	33	3.93	0.11
C	24	3.26	0.14
D	46	6.93	0.15
E	44	4.45	0.10
F	13	0.86	0.07
G	4	Unknown	--

Table 3 Percentage of the land used for each *T. cristatus* group.

Group	A	B	C	D	E	F
Vegetated area	47.04	61.10	47.42	61.54	61.55	49.87
Grass and open area	37.72	29.10	41.01	28.95	28.51	30.64
Building/construction area	15.24	9.89	11.56	9.51	9.94	19.49

Table 4 The area size (upper) and percent (bottom) of spatial overlap between the six one-male-multifemale groups.

Group ID	A	B	C	D	E	F
A		1.31 ha	1.73 ha	0.93 ha	0.41 ha	0.43 ha
B	27.6%		1.24 ha	1.90 ha	0.12 ha	0.21 ha
C	46.8%	20.8%		2.32 ha	1.83 ha	0.80 ha
D	11.5%	21.2%	25.6%		4.39 ha	0.86 ha
E	6.61%	1.44%	31.1%	62.7%		0.69 ha
F	16.5%	4.64%	24.2%	12.4%	15.0%	

Figure 1. Ranging area of each *T. cristatus* group (Group A–E) with compiled ranging areas of all six groups at the study site.

Figure 2 Area size calculated by MCP based on the cumulative location points in each study group.

Figure 3. Relationships between the ranging area in each group (left) and the total vegetated area within each group's ranging area with group size (the number of group members in each group). The blue dotted lines and shaded areas represent the linear regression lines for the observed samples and their 95% confidence interval ranges, respectively.

Appendix 1. Chart of age categories with associated morphological and physical characteristics according to Aggimarangsee (2002) and Hamada and Yamamoto (2010).

Age categories	Morphology and physical characteristics
Adult	Large in body size and larger than the other age categories. The body growth stops as stature and reproductive organs reach maturity. In females, genital swelling is confined to the vulva and the surrounding perineum and is expected to occur most frequently during the female cycle. In males, the testes increase in size, and musculoskeletal development occurs.
Subadult	Smaller than adults in body size but bigger than juveniles; i.e. individuals that are more than three-fourth of the body size of full-sized adults. Reproductive organs are immature.
Juvenile	Pelage of the head and body has changed to adult-coloured grey, but they are less than three-fourth of the full adult body size. Some juveniles are occasionally breast-fed.
Infants	Newborns and individuals at weaning age are categorized as infants. The pelage of the head and body is orange in color and is starting to change to gray. The face skin is slightly pinkish and slowly changes to dark gray and black as the infant reaches the juvenile stage. Total coloration changes take about three months.