Asymmetric chromatin capture and nuclear envelopes separate endogenous and ectopic chromosomes after induced cell fusion

Bharath Sunchu [1], Nicole Lee [1,2], Roberto Carlos Segura[1] & Clemens Cabernard [1]

[1] Department of Biology, University of Washington
Life Science Building
Seattle, Washington State, USA

Current address:

[2] Cancer Science Institute, National University of Singapore
Singapore

* Lead and corresponding author: ccabern@uw.edu

Metazoan cells accurately attach to, congress and segregate chromosomes during mitosis

Additionally, hybrid cells derived through fertilization or somatic cell fusion also employ

mechanisms to recognize and separate chromosomes of different origin. The underlying

mechanisms are mostly unknown but could prevent aneuploidy and tumor formation. Here,

we acutely induce fusion between *Drosophila* neural stem cells (neuroblasts) and

differentiating ganglion mother cells (GMCs) *in vivo* to define how epigenetically distinct

chromatin is recognized and segregated. We find that Nb-GMC hybrid cells align both

endogenous (neuroblast-origin) and ectopic (GMC-origin) chromosomes at the metaphase

plate through centrosome derived dual spindles. Mixing of endogenous and ectopic

chromatin is prevented through an asymmetric, microtubule-dependent chromatin capture

mechanism during interphase and physical boundaries imposed by nuclear envelopes.

Although hybrid cells fail to accurately segregate ectopic chromatin, manifested in lagging

chromosomes and chromosome bridges, transplanted brain tissue containing hybrid cells

neither reduce the lifespan nor form visible tumors in host flies. We conclude that fly neural

stem cells utilize asymmetric centrosome activity in interphase to capture and physically

separate epigenetically distinct chromatin in a microtubule-dependent manner. We propose

a novel chromosome recognition and separation mechanism that could also inform biased

chromatid segregation observed in flies and vertebrates.

Dividing cells distribute replicated chromosomes equally between the two forming sibling cells

through microtubule-dependent attachment and segregation mechanisms (McIntosh, 2016;

Maiato et al., 2017). Chromosomes are connected with microtubules via kinetochore proteins,

which are localized on centromeric DNA (Thomas et al., 2017; Yu et al., 2019). The occurrence of

biased segregation of sister chromatids or homologous chromosomes during mitosis and meiosis

respectively, as well as the spatial separation of the maternal and paternal genomes during early

vertebrate development, implies the presence of accurate chromatin recognition mechanisms

(Akera et al., 2017; Ranjan et al., 2019; Sunchu and Cabernard, 2020; Reichmann et al., 2018).

Chromosome separation has also been observed in fused human cells (Heasley et al., 2017; Rieder

et al., 1997). The molecular nature of these separation mechanisms is unclear but could entail

asymmetries in centromere binding proteins, kinetochore size or kinetochore composition (Arco et

al., 2018; Ranjan et al., 2019; Akera et al., 2019; Drpic et al., 2018).

Over a century ago, cell – cell fusion has also been proposed to initiate tumor formation (Aichel,

1911), possibly through chromosome missegregation and aneuploidy (Boveri, 2008; Molina et al.,

2020). Here, we ask how hybrid cells accurately recognize, separate and segregate epigenetically

distinct chromosomes. To this end, we acutely fused *Drosophila* neural stem cells (neuroblasts

(NBs), hereafter) with differentiating ganglion mother cells (GMCs) in the intact larval fly brain

to create hybrid cells, containing both neuroblast and GMC chromosomes. Unperturbed *Drosophila*

neuroblasts divide asymmetrically, self-renewing the neural stem cell while forming a

differentiating GMC. Neuroblasts are twice the size of GMCs and neurons and express the

transcription factor Deadpan (Dpn[+]), whereas GMCs and neurons can be identified based on

Prospero (Pros[+]) expression (Gallaud et al., 2017) (Fig. S1A). We used a 532nm pulsed laser to

induce a small lesion at the Nb – GMC interface between a dividing neuroblast and GMC, causing

the GMC chromatin to enter the neuroblast cytoplasm. Targeted mitotic neuroblasts often retain

the GMC chromosomes, creating a large apical hybrid cell containing a Dpn[+] and Pros[+] nucleus

(Nb – GMC hybrid). Most Nb – GMC hybrid cells normally localize the contractile ring marker non-

muscle Myosin to the cleavage furrow and complete cytokinesis (Fig. S1B-E). Acute cell fusion can

also result in the expulsion of the GMC chromatin, forming GMC – GMC hybrids (see

Supplementary Fig. 3c in (Roubinet et al., 2017)) but the focus here is on Nb – GMC hybrids

(hybrid cells, hereafter) only.

To better characterize the dynamics of neuroblast (endogenous) and GMC (ectopic)

chromosomes during mitosis, we induced cell fusion at different cell cycle stages in wild type

neuroblasts, expressing the canonical chromosome marker His2A::GFP. Hybrid cells derived from

Nb-GMC fusions early in the cell cycle could (1) align only the neuroblast chromosomes at the

metaphase plate, (2) congress a mix of neuroblast and GMC chromosomes or (3) separate and

align the two chromosome pools at the metaphase plate (Fig. 1A). We found that the endogenous

and ectopic chromatin was separated and distinguishable when fusions were induced in early

mitosis. Both the ectopic and endogenous chromatin aligned at the metaphase plate (Fig. 1B; video

1&2). Nb-GMC fusions can be induced at all cell cycle stages but GMC chromosomes align at the

metaphase plate more accurately in hybrid cells derived from interphase or early prophase fusions

(Fig. 1C, D). We conclude that hybrid cells derived from fusions between interphase Nbs and non-

mitotic GMCs accurately align ectopic and endogenous chromatin at the metaphase plate.

Nb-GMC hybrid cells independently align Nb and GMC chromatin at the metaphase plate

We next asked whether GMC chromatin congresses independently of neuroblast chromatin. To this

end, we measured the time between nuclear envelope breakdown (NEB) and chromosome

alignment at the metaphase plate for Nb and ectopic GMC chromatin in hybrid cells derived from

interphase and early prophase fusions (Fig. 1E). In most hybrid cells, ectopic and endogenous

chromatin was distinguishable based on differences in location and intensity (see Fig. 1B; video

1&2). In untargeted control neuroblasts (unfused), Nb chromosomes aligned at the metaphase

plate within 6.5 minutes after NEB (SD = 1.55; n = 6), which is insignificantly faster than the

neuroblast chromosomes of hybrid cells (t = 6.9 mins; SD = 2.84; n = 11). GMC chromosomes

aligned within 7.63 mins (SD = 3.69; n = 11), statistically not significantly different from

unperturbed wild type chromosomes (Fig. 1E). In most Nb-GMC hybrids, the endogenous

neuroblast and the ectopic GMC chromosomes aligned at the metaphase plate with no significant

time difference. However, in a few cases, ectopic chromatin aligned before or after the neuroblast

chromatin (Fig. 1F). These results suggest that the neuroblast and GMC chromatin can move

independently to the metaphase plate in hybrid cells.

Ectopic spindles distinguish between Nb and GMC chromatin

We next investigated the mechanisms underlying independent Nb/GMC chromosome alignment.

Ectopic chromosomes could be aligned together with the endogenous chromosomes via a single

bipolar spindle. Alternatively, hybrid cells could form multiple bipolar spindles, which attach to

either the neuroblast's, GMC's, or chromosomes from both cell types (Fig. 2Aa). Live cell imaging

showed that hybrid cells derived from interphase Nb-GMC fusions contained double spindles in

almost all cases, whereas the vast majority of cell fusions induced in metaphase only formed one

mitotic spindle (Fig. 2B, C & video 3). Most hybrid cells contained two mitotic spindles, which

predominantly formed at the same time (Fig. 2D, E). We quantified spindle alignment and

positioning dynamics by measuring the angle and distance between the two spindles during

metaphase (Fig. 2F and methods). The two spindles were initially misaligned and separated but

decreased their inter-spindle distance and angle during metaphase (Fig. 2G-I). We conclude that

Nb-GMC hybrid cells align neuroblast and GMC chromosomes separately at the metaphase plate

through the formation of independent mitotic spindles.

Ectopic spindles are nucleated from GMC centrosomes

Mitotic spindles can be nucleated through the centrosome-dependent, chromatin or microtubule

pathway but when centrosomes are present, the centrosome pathway prevails (Prosser and

Pelletier, 2017). To elucidate the mechanisms underlying ectopic spindle formation, we induced

Nb-GMC fusions in interphase wild type neuroblasts expressing a live centriole (Asterless;

Asl::GFP) and spindle marker (cherry::Jupiter), and assayed centrosome dynamics and spindle

formation throughout mitosis. Normal wild type neuroblasts usually contain two Asl::GFP positive

centrioles in mitosis, forming a single bipolar spindle. However, in Nb-GMC hybrids, we

predominantly found four Asl::GFP positive centrioles, two of which seemed to be introduced from

the GMC (Fig. 2J,K). The GMC centrosomes nucleated an ectopic bipolar spindle that subsequently

aligned with the main neuroblast spindle. Although multipolar spindle formation can be prevented

through centrosome clustering (Quintyne et al., 2005), we observed that GMC centrosomes

approached neuroblast centrosomes but still formed two parallel bipolar spindles. We conclude

that ectopic spindles are formed through the centrosome pathway, using centrioles originating

from GMCs.

Microtubule-dependent, asymmetric chromatin-centrosome attachments retain

chromosomes close to the apical neuroblast cortex during interphase

Our data suggest that Nb and GMC chromatin are being separated through an endogenous, Nb-

derived and an ectopic, GMC-derived mitotic spindle. We next investigated how these spindles

distinguish between Nb and GMC chromosomes. During mitosis, microtubules emanate from

centrosomes and attach to sister chromatids through kinetochore proteins, localizing to the

centromeric region (Fukagawa and Earnshaw, 2014). *Drosophila* male germline stem cells contain

asymmetric levels of the centromere-specific H3 variant (Centromere identifier (Cid) in flies

(Henikoff et al., 2000)) (Ranjan et al., 2019), prompting us to investigate whether Cid intensity

differs between endogenous and ectopic chromatin in hybrid cells. We induced Nb-GMC fusions

of wild type cells expressing Cid::EGFP (Ranjan et al., 2019) and measured Cid intensity on both

GMC and Nb chromatin. These measurements did not reveal a significant intensity difference

between Nb and GMC Cid (Fig. S2A). However, we noticed that endogenous Cid::EGFP was

localized in very close proximity to the apical centrosome in unperturbed interphase and prophase

wild type neuroblasts (Fig. S2B & video 4). Cid::EGFP remained associated with chromatin

throughout the neuroblast cell cycle, excluding the possibility that early Cid clusters are not

connected with chromatin (Fig. S2F & video 5).

Interphase wild type neuroblasts contain only one active apical microtubule organizing center

(MTOC), retaining the daughter-centriole containing centrosome close to the apical neuroblast

cortex. The mother-centriole-containing centrosome is inactive in interphase but matures from

prophase onward, positioning itself on the basal cell cortex (Januschke et al., 2013, 2011;

Januschke and Gonzalez, 2010; Gallaud et al., 2020). We measured the distance of individual

Cid::EGFP clusters to the apical and basal centrosome in unperturbed wild type neuroblasts (Fig.

S 2C) and found that interphase Cid was always in close proximity to the apical centrosome. After

nuclear envelope breakdown (NEB), Cid moved progressively towards the metaphase plate (Fig. S

2D). Once the basal centrosome appeared (0 mins), Cid was still closer to the apical than the basal

centrosome and this distance asymmetry was still intact 6 minutes after the appearance of the

basal centrosome (Fig. S2E).

The proximity of Cid::EGFP clusters to the active interphase MTOC suggests a microtubule-

dependent chromosome attachment mechanism. Indeed, wild type neuroblasts treated with the

microtubule-depolymerizing drug colcemid showed a strong correlation between apical MTOC

activity and Cid localization; as MTs depolymerize after colcemid addition, Cid progressively

moved away from the apical cortex towards the cell center (Fig. S2G, H & video 6). To test whether

MTOCs are connected to Cid clusters during interphase, we removed the centriolar protein

Centrobin (Cnb; CNTROB in humans). Neuroblasts lacking Cnb fail to maintain an active apical

interphase MTOC but regain normal MTOC activity during mitosis (Januschke et al., 2013).

Neuroblasts expressing *cnb* RNAi lost apical Cid localization after the apical centrosome

downregulated its MTOC activity. However, maturing centrosomes reconnected with Cid in

prophase (Fig. S2I, J & video 7). Cid's proximity to the apical MTOC ('apical' refers to the

centrosome destined to move to the apical cortex) in *cnb* RNAi expressing neuroblasts was much

more varied compared to wild type (Fig. S2L). At 6 minutes after centrosome maturation onset,

Cid – apical MTOC distance was comparable to wild type, as were Cid – basal centrosome distance

relationships (Fig. S2K-M). We conclude that Cid-containing chromatin is already attached to the

apical centrosome prior to entry into mitosis.

Asymmetric centrosome-chromatin attachments contribute to the separation of

endogenous and ectopic chromosomes

Based on these observations, we hypothesized that the separation between endogenous and

ectopic chromosomes could be due to a pre-attachment mechanism, preventing mixing of Nb and

GMC chromosomes. To test this hypothesis, we first analyzed Cid localization in relation to the

endogenous and ectopic centrosomes in wild type hybrid cells. Similar to unperturbed wild type

neuroblasts, endogenous Cid is also localized in close proximity to the endogenous apical

centrosome in wild type hybrid cells (Fig. 3A-C, G, H & video 8; 'apical' refers to the centrosome

destined to segregate into the large apical sibling cell) and appeared further away from ectopic

Cid clusters compared to ectopic centrosomes (Fig. 3A-C, I & video 8; 0 mins refers to the

appearance of the endogenous basal centrosome; '0' refers to the appearance of the ectopic

centrosomes).

We next attempted to randomize Cid – centrosome distance relationships by inducing acute fusions

in *cnb* RNAi expressing neuroblasts, since loss of interphase MTOC activity released endogenous

Cid from the apical centrosome (see above; Fig. S2G-M). In contrast to wild type hybrid cells,

endogenous Cid is roughly equidistant to the endogenous and ectopic centrosomes in *cnb* RNAi

expressing hybrid cells at 0 mins and '0' mins respectively (Fig. 3D-F, J, K & video 9). However,

ectopic Cid was still closer to the ectopic centrosomes in *cnb* RNAi expressing hybrid cells compared

to the endogenous apical centrosome (Fig. 3D, F, L & video 9). In both wild type and *cnb* RNAi

expressing hybrid cells, ectopic and basal centrosomes were about equidistant to endogenous Cid,

but ectopic centrosomes were closer to ectopic Cid (Fig. S3A-D). Taken together, we conclude that

in hybrid cells the apical MTOC forms an asymmetric attachment to endogenous Cid-containing

chromosomes prior to entry into mitosis.

Nuclear envelopes separate ectopic and endogenous chromatin in hybrid cells

We next asked whether early MTOC – Cid attachments are sufficient to prevent endogenous and

ectopic chromosome separation, tracking endogenous and ectopic Cid::EGFP after induced cell

fusion. In wild type hybrid cells, endogenous and ectopic CID clusters started to congress at the

metaphase plate and became difficult to clearly separate 8.8 mins (SD = 5.63; n = 5; Fig. 3M)

after NEB. However, 50% of *cnb* RNAi hybrid cells showed endogenous and ectopic Cid mixing

prior to NEB (Fig. 3D, N), although the time difference was not significantly different to wild type

hybrid cells (Fig. 3M).

Neuroblasts undergo semi-closed mitosis, mostly retaining a matrix composed of nuclear

envelope proteins around the mitotic spindle (Katsani et al., 2008). We imaged hybrid cells with

the nuclear envelope marker Klaroid, using the protein-trap line koi::EGFP (Buszczak et al., 2007).

We confirmed that unperturbed wild type neuroblasts contain a nuclear envelope matrix

surrounding the mitotic spindle during mitosis (Fig. S3E). Similarly, wild type hybrid cells contain

two nuclear envelopes during mitosis (Fig. 3O and Fig. S3F). Taken together, these data suggest

that asymmetric MTOC-Cid attachments in interphase and nuclear envelopes establish and

maintain the physical separation between endogenous and ectopic chromosomes in hybrid cells.

Loss of biased interphase MTOC activity removes asymmetric MTOC-Cid attachments and allows

for cross-connections between endogenous centrosomes and ectopic Cid (Fig. 3P).

Hybrid cells segregate endogenous and ectopic chromosomes independently

We next investigated whether both bipolar spindles are functional in Nb-GMC hybrid cells.

Erroneous or incomplete microtubule-kinetochore attachments trigger the spindle assembly

checkpoint (SAC), preventing or delaying anaphase entry (Musacchio, 2015). Since the

kinetochore-derived 'wait anaphase' signal is diffusible (Heasley et al., 2017), ectopic spindles

should thus also obey the SAC in Nb-GMC hybrid cells. We tested whether hybrid cells contain

functional microtubule-kinetochore attachments by measuring the time between finished

chromosome alignment at the metaphase plate and chromosome separation in Nb-GMC hybrids

expressing His2A::GFP and cherry::Jupiter (Fig. 4A, B). Unperturbed control neuroblasts usually

initiate anaphase onset within 2.63 minutes (SD=1.84; n=12) after chromosomes are aligned at

the metaphase plate. In hybrid cells derived from interphase fusions, endogenous and ectopic

chromatin entered anaphase 4.14 mins (SD= 2.23; n=14) and 5.12 minutes (SD= 1.85; n=13)

after metaphase alignment. Only ectopic chromatin for prophase induced hybrid cells showed a

significantly delayed anaphase onset (Average: 10.25 mins; SD= 4.12; n=6) (Fig. 4C). Ectopic

chromatin never separated before endogenous chromosomes but entered anaphase with a few

minutes delay (Fig. 4D). We conclude that in Nb-GMC hybrids, endogenous and ectopic

chromosomes establish correct MT-kinetochore attachments, thereby fulfilling the spindle

assembly checkpoint necessary to enter anaphase. However, given the delays in ectopic

chromosome separation, we further conclude that ectopic spindles can initiate chromatid

separation independently from the endogenous neuroblast spindle.

Hybrid cells are insufficient to induce tumors in wild type host flies

Finally, we assessed the accuracy of chromosome segregation in wild type hybrid cells. Using the

canonical chromosome marker His2A::GFP we detected chromosome missegregation - ranging

from lagging chromosomes to chromosome bridges - in all wild type hybrid cells (Fig. 4E,F & video

10). Chromosome segregation defects can result in aneuploidy and micronuclei formation (Molina

et al., 2020). We found a small percentage of hybrid cells containing micronuclei, but more

frequently discovered heterokaryons (hybrid cells containing two nuclei of different origin). In

most cases, hybrid cells fused both nuclei into one, forming synkaryons (Fig. 4G).

Aneuploidy has been proposed to be a hallmark of cancer (Molina et al., 2020) but appears

to be context dependent (Ben-David and Amon, 2020). To test whether neuroblast – GMC derived

hybrid cells are sufficient to induce tumor formation, we grafted His2A::GFP expressing larval fly

brains after successful induction of cell fusion into the abdomen of wild type hosts (Rossi and

Gonzalez, 2015) and monitored the host flies for tumor formation and life span changes. As

previously reported (Caussinus and Gonzalez, 2005), brat RNAi expressing brains formed visible

tumors in host flies by day 30 and caused a reduction in lifespan of the host. However, larval brains

without attempted fusions (wild type transplants), attempted but unsuccessful fusions (controlling

for the effect of laser ablation), or induced fusion, showed neither tumor growth by day 30 (or

after), nor a reduction in lifespan (Fig. 5A,B). We conclude that chromosome segregation is

defective in Nb-GMC hybrid cells but insufficient to form visible tumors in otherwise normal and

unperturbed larval fly brains.

Discussion

Cell – cell fusion can occur under normal physiological conditions and has been implicated in malignancy (Platt and Cascalho, 2019) but how hybrid cells recognize and separate endogenous and ectopic chromosomes during mitosis is unclear. Here we have shown *in vivo* that Nb-GMC derived hybrid cells separate endogenous chromosomes from the introduced ectopic GMC chromosomes until metaphase and potentially during anaphase. Chromosome separation is achieved through a dual-spindle mechanism; both endogenous and ectopic chromosomes nucleate two independent spindles, most likely through the canonical centrosome pathway. These dual spindles co-align during metaphase, thereby congressing the two chromosome clusters at the metaphase plate. Endogenous and ectopic chromosomes independently segregate during anaphase, manifested in delayed segregation onset of ectopic chromatin. Although chromosome missegregation is frequent in hybrid cells, potentially leading to aneuploidy, we failed to detect malignant tumor formation when hybrid cell – containing larval brains were grafted into wild type host flies.

Taken together, we propose that endogenous and ectopic chromosome separation is achieved through an early microtubule-dependent chromosome capture mechanism that retains endogenous chromosomes in close proximity to the apical neuroblast cortex during interphase. In addition to this geometrical separation, the nuclear envelope might impose a physical boundary, preventing random chromosome mixing prior to chromosome congression (Fig. 5C). In pre-mitotic neuroblasts, chromatin can be connected with centrosomes through the Linc complex (Lee and Burke, 2018), potentially implicating the SUN domain protein Klaroid (Kracklauer et al., 2007) and the KASH-domain protein Klarsicht (Razafsky and Hodzic, 2009; Lee and Burke, 2018) in asymmetric chromatin clustering and the prevention of chromatin mixing during interphase.

The chromatin separation mechanisms described here could be applicable to chromosome

separation occurring in the first cleavage after fertilization in insects, arthropods and vertebrates

(Reichmann et al., 2018; Kawamura, 2001). Similarly, biased chromatid segregation has been

observed in stem cells (Yadlapalli and Yamashita, 2013; Ranjan et al., 2019) and meiosis (Akera

et al., 2017). Since centromeres have also been found to be confined to specific nuclear locations

in many organisms (Weierich et al., 2003; Muller et al., 2019), it will be interesting to see whether

microtubule-dependent chromatin attachment provides an alternative mechanism for biased sister

chromatid segregation or other important cellular functions.

References

Aichel, O. 1911. Über Zellverschmelzung mit quantitativ abnormer Chromosomenverteilung als Ursache der Geschwulstbildung. W. Roux, editor. Verlag von Wilhelm Engelmann.

Akera, T., L. Chmátal, E. Trimm, K. Yang, C. Aonbangkhen, D.M. Chenoweth, C. Janke, R.M. Schultz, and M.A. Lampson. 2017. Spindle asymmetry drives non-Mendelian chromosome segregation. *Science*. 358:668 672. doi:10.1126/science.aan0092.

Akera, T., E. Trimm, and M.A. Lampson. 2019. Molecular Strategies of Meiotic Cheating by Selfish Centromeres. *Cell*. 178:1132-1144.e10. doi:10.1016/j.cell.2019.07.001.

Arco, A.G. del, B.A. Edgar, and S. Erhardt. 2018. In Vivo Analysis of Centromeric Proteins Reveals a Stem Cell-Specific Asymmetry and an Essential Role in Differentiated, Non-proliferating Cells. *Cell Reports*. 22:1982–1993. doi:10.1016/j.celrep.2018.01.079.

Ben-David, U., and A. Amon. 2020. Context is everything: aneuploidy in cancer. *Nat Rev Genet*. 21:44–62. doi:10.1038/s41576-019-0171-x.

Blachon, S., J. Gopalakrishnan, Y. Omori, A. Polyanovsky, A. Church, D. Nicastro, J. Malicki, and T. Avidor-Reiss. 2008. Drosophila asterless and vertebrate Cep152 Are orthologs essential for centriole duplication. *Genetics*. 180:2081 2094. doi:10.1534/genetics.108.095141.

Boveri, T. 2008. Concerning the Origin of Malignant Tumours by Theodor Boveri. Translated and annotated by Henry Harris. *J Cell Sci*. 121:1–84. doi:10.1242/jcs.025742.

Buszczak, M., S. Paterno, D. Lighthouse, J. Bachman, J. Planck, S. Owen, A.D. Skora, T.G. Nystul, B. Ohlstein, A. Allen, J.E. Wilhelm, T.D. Murphy, R.W. Levis, E. Matunis, N. Srivali, R.A. Hoskins, and A.C. Spradling. 2007. The carnegie protein trap library: a versatile tool for Drosophila developmental studies. *Genetics*. 175:1505 1531. doi:10.1534/genetics.106.065961.

Cabernard, C., and C.Q. Doe. 2009. Apical/basal spindle orientation is required for neuroblast homeostasis and neuronal differentiation in Drosophila. *Dev Cell*. 17:134 141. doi:10.1016/j.devcel.2009.06.009.

Cabernard, C., K.E. Prehoda, and C.Q. Doe. 2010. A spindle-independent cleavage furrow positioning pathway. *Nature*. 467:91 94. doi:10.1038/nature09334.

Caussinus, E., and C. Gonzalez. 2005. Induction of tumor growth by altered stem-cell asymmetric division in Drosophila melanogaster. *Nat Genet*. 37:1125 1129. doi:10.1038/ng1632.

Dietzl, G., D. Chen, F. Schnorrer, K.-C. Su, Y. Barinova, M. Fellner, B. Gasser, K. Kinsey, S. Oppel, S. Scheiblauer, A. Couto, V. Marra, K. Keleman, and B.J. Dickson. 2007. A genome-wide transgenic RNAi library for conditional gene inactivation in Drosophila. *Nature*. 448:151 156. doi:10.1038/nature05954.

Drpic, D., A.C. Almeida, F. Renda, J. Damas, H.A. Lewin, D.M. Larkin, A. Khodjakov, and H. Maiato.
2018. Chromosome Segregation Is Biased by Kinetochore Size. *Curr Biol*. 28:1344 1356.e5.
doi:10.1016/j.cub.2018.03.023.

Fukagawa, T., and W.C. Earnshaw. 2014. The Centromere: Chromatin Foundation for the Kinetochore
Machinery. *Dev Cell*. 30:496–508. doi:10.1016/j.devcel.2014.08.016.

Gallaud, E., A.R. Nair, N. Horsley, A. Monnard, P. Singh, T.T. Pham, D.S. Garcia, A. Ferrand, and C.
Cabernard. 2020. Dynamic centriolar localization of Polo and Centrobin in early mitosis primes
centrosome asymmetry. *Plos Biol*. 18:e3000762. doi:10.1371/journal.pbio.3000762.

Gallaud, E., T. Pham, and C. Cabernard. 2017. Drosophila melanogaster Neuroblasts: A Model for
Asymmetric Stem Cell Divisions. *Results Problems Cell Differ*. 61:183 210. doi:10.1007/978-3-319-
53150-2_8.

Gratz, S.J., A.M. Cummings, J.N. Nguyen, D.C. Hamm, L.K. Donohue, M.M. Harrison, J. Wildonger,
and K.M. O'Connor-Giles. 2013. Genome Engineering of Drosophila with the CRISPR RNA-Guided
Cas9 Nuclease. *Genetics*. 194:1029–1035. doi:10.1534/genetics.113.152710.

Heasley, L.R., S.M. Markus, and J.G. DeLuca. 2017. "Wait Anaphase" Signals Are Not Confined to the
Mitotic Spindle. *Mol Biol Cell*. 28:1186–1194. doi:10.1091/mbc.e17-01-0036.

Henikoff, S., K. Ahmad, J.S. Platero, and B. van Steensel. 2000. Heterochromatic deposition of
centromeric histone H3-like proteins. *Proc National Acad Sci*. 97:716–721.
doi:10.1073/pnas.97.2.716.

Januschke, J., and C. Gonzalez. 2010. The interphase microtubule aster is a determinant of asymmetric
division orientation in Drosophila neuroblasts. *J Cell Biology*. 188:693 706.
doi:10.1083/jcb.200905024.

Januschke, J., S. Llamazares, J. Reina, and C. Gonzalez. 2011. Drosophila neuroblasts retain the daughter
centrosome. *Nat Commun*. 2:243. doi:10.1038/ncomms1245.

Januschke, J., J. Reina, S. Llamazares, T. Bertran, F. Rossi, J. Roig, and C. Gonzalez. 2013. Centrobin
controls mother–daughter centriole asymmetry in Drosophila neuroblasts. *Nat Cell Biol*. 15:241 248.
doi:10.1038/ncb2671.

Katsani, K.R., R.E. Karess, N. Dostatni, and V. Doye. 2008. In Vivo Dynamics of Drosophila Nuclear
Envelope Components. *Mol Biol Cell*. 19:3652–3666. doi:10.1091/mbc.e07-11-1162.

Kawamura, N. 2001. Fertilization and the first cleavage mitosis in insects. *Dev Growth Differ*. 43:343–
349. doi:10.1046/j.1440-169x.2001.00584.x.

Kracklauer, M.P., S.M.L. Banks, X. Xie, Y. Wu, and J.A. Fischer. 2007. Drosophila klaroid Encodes a
SUN Domain Protein Required for Klarsicht Localization to the Nuclear Envelope and Nuclear
Migration in the Eye. *Fly*. 1:75–85. doi:10.4161/fly.4254.

Lee, Y.L., and B. Burke. 2018. LINC complexes and nuclear positioning. *Semin Cell Dev Biol*. 82:67 76.
doi:10.1016/j.semcdb.2017.11.008.

390 Maiato, H., A.M. Gomes, F. Sousa, and M. Barisic. 2017. Mechanisms of Chromosome Congression
391   during Mitosis. *Biology*. 6:13. doi:10.3390/biology6010013.

392 McIntosh, J.R. 2016. Mitosis. *Csh Perspect Biol*. 8:a023218. doi:10.1101/cshperspect.a023218.

393 Molina, O., M.A. Abad, F. Solé, and P. Menéndez. 2020. Aneuploidy in Cancer: Lessons from Acute
394   Lymphoblastic Leukemia. *Trends Cancer*. doi:10.1016/j.trecan.2020.08.008.

395 Muller, H., J. Gil, and I.A. Drinnenberg. 2019. The Impact of Centromeres on Spatial Genome
396   Architecture. *Trends Genet*. 35:565–578. doi:10.1016/j.tig.2019.05.003.

397 Musacchio, A. 2015. The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics. *Curr*
398   *Biol*. 25:R1002 18. doi:10.1016/j.cub.2015.08.051.

399 Platt, J.L., and M. Cascalho. 2019. Cell Fusion in Malignancy: A Cause or Consequence? A Provocateur
400   or Cure? *Cells*. 8:587. doi:10.3390/cells8060587.

401 Prosser, S.L., and L. Pelletier. 2017. Mitotic spindle assembly in animal cells: a fine balancing act. *Nat*
402   *Rev Mol Cell Bio*. 18:187–201. doi:10.1038/nrm.2016.162.

403 Quintyne, N.J., J.E. Reing, D.R. Hoffelder, S.M. Gollin, and W.S. Saunders. 2005. Spindle Multipolarity
404   Is Prevented by Centrosomal Clustering. *Science*. 307:127–129. doi:10.1126/science.1104905.

405 Ranjan, R., J. Snedeker, and X. Chen. 2019. Asymmetric Centromeres Differentially Coordinate with
406   Mitotic Machinery to Ensure Biased Sister Chromatid Segregation in Germline Stem Cells. *Cell Stem*
407   *Cell*. doi:10.1016/j.stem.2019.08.014.

408 Razafsky, D., and D. Hodzic. 2009. Bringing KASH under the SUN: the many faces of nucleo-
409   cytoskeletal connections. *J Cell Biology*. 186:461 472. doi:10.1083/jcb.200906068.

410 Reichmann, J., B. Nijmeijer, M.J. Hossain, M. Eguren, I. Schneider, A.Z. Politi, M.J. Roberti, L.
411   Hufnagel, T. Hiiragi, and J. Ellenberg. 2018. Dual-spindle formation in zygotes keeps parental
412   genomes apart in early mammalian embryos. *Science*. 361:189 193. doi:10.1126/science.aar7462.

413 Rieder, C.L., A. Khodjakov, L.V. Paliulis, T.M. Fortier, R.W. Cole, and G. Sluder. 1997. Mitosis in
414   vertebrate somatic cells with two spindles: Implications for the metaphase/anaphase transition
415   checkpoint and cleavage. *Proc National Acad Sci*. 94:5107–5112. doi:10.1073/pnas.94.10.5107.

416 Rossi, F., and C. Gonzalez. 2015. Studying tumor growth in Drosophila using the tissue allograft method.
417   *Nat Protoc*. 10:1525–1534. doi:10.1038/nprot.2015.096.

418 Roubinet, C., A. Tsankova, T.T. Pham, A. Monnard, E. Caussinus, M. Affolter, and C. Cabernard. 2017.
419   Spatio-temporally separated cortical flows and spindle geometry establish physical asymmetry in fly
420   neural stem cells. *Nat Commun*. 8:1383. doi:10.1038/s41467-017-01391-w.

421 Sunchu, B., and C. Cabernard. 2020. Principles and mechanisms of asymmetric cell division.
422   *Development*. 147:dev167650. doi:10.1242/dev.167650.

Thomas, G.E., M.R. Renjith, and T.K. Manna. 2017. Kinetochore–microtubule interactions in
chromosome segregation: lessons from yeast and mammalian cells. *Biochem J*. 474:3559–3577.
doi:10.1042/bcj20170518.

Weierich, C., A. Brero, S. Stein, J. von Hase, C. Cremer, T. Cremer, and I. Solovei. 2003. Three-
dimensional arrangements of centromeres and telomeres in nuclei of human and murine lymphocytes.
Chromosome Res. 11:485–502. doi:10.1023/a:1025016828544.

Yadlapalli, S., and Y.M. Yamashita. 2013. Chromosome-specific nonrandom sister chromatid segregation
during stem-cell division. *Nature*. 498:251 254. doi:doi:10.1038/nature12106.

Yu, K., N. Zhong, Y. Xiao, and Z. She. 2019. Mechanisms of kinesin-7 CENP-E in kinetochore–
microtubule capture and chromosome alignment during cell division. *Biol Cell*. 111:143–160.
doi:10.1111/boc.201800082.

434
435

Figures

 Figure 1: Nb-GMC hybrid cells align Nb and GMC chromatin independently at the

 metaphase plate.

 (A) Nb – GMC derived hybrid cells could (1) only align neuroblast chromosomes at the metaphase

 plate, (2) congress a mix of endogenous and ectopic chromosomes or (3) separately align Nb and

 GMC chromosomes at the metaphase plate. (B) Representative image sequence of a dividing third

 instar larval neuroblast Nb-GMC hybrid cell, expressing the histone marker His2A::GFP (dashed

 blue circle: endogenous chromatin; dashed orange circle: ectopic chromatin). (C) Alignment of

 endogenous and ectopic chromatin in hybrid cells derived from interphase (inter), prophase (pro),

 prometaphase (prometa) or metaphase (meta) fusions were quantified with angle measurements

 in metaphase and plotted in (D). (E) Chromosome alignment time for endogenous (blue spheres)

 and ectopic chromosomes (orange squares) compared to unfused control neuroblasts (grey

 triangles) Colored boxes represent corresponding cell cycle stages. One-way ANOVA was used for

 (D) and (E). Error bars correspond to standard deviation (SD). * p<0.05, **p<0.01,****

 p<0.000.1. For this and subsequent Figures, exact p values and complete statistical information

 can be found in Supplemental table 1. Time in mins:secs. Scale bar is 5 μm.

Figure 2: Dual spindles align endogenous and ectopic spindles separately at the metaphase plate.

(A) Nb – GMC derived hybrid cells could align neuroblast and GMC chromosomes either through a single or dual-spindle mechanism. (B) Representative third instar larval Nb-GMC hybrid cell, expressing the histone marker His2A::GFP (white in top row; cyan in merged channel below), the MT marker cherry::Jupiter (white in middle and bottom row). (C) Quantification of hybrid cells containing dual- or multiple spindles for hybrid cells derived from interphase (inter), prophase

(pro), prometaphase (prometa) or metaphase (meta) fusions. (D) The time difference between

endogenous and ectopic spindle formation was measured in hybrid cells and plotted in (E). (F)

Spindle angle and inter-spindle distances were measured during mitosis. A representative example

is shown in (G). Quantification of inter-spindle (H) distances and (I) angles at prometaphase and

metaphase. (J) A representative third instar larval Nb-GMC hybrid, expressing the centriole

marker, Asl::GFP (cyan; top row. White; bottom row) and the spindle marker cherry::Jupiter (MTs;

white in top row). Centrosomes were highlighted with green dashed circles. (K) Comparison of

centrosome number between unfused wild type and hybrid cells.

Colored boxes represent corresponding cell cycle stages. Error bars correspond to SDs. Figures (H)

and (I) paired t-test, Figures (K) unpaired t-test. * $p<0.05$, **$p<0.01$,**** $p<0.000$. Time in

mins:secs. Scale bar is 5 μm.

Figure 3: Asymmetric microtubule dependent centrosome-chromatin attachments contribute to the separation of endogenous and ectopic chromosomes.

Representative third instar larval (A) wild type or (D) *cnb* RNAi expressing Nb-GMC hybrid cell, expressing Cid::EGFP (top row; cyan, bottom row; white) and the microtubule marker cherry::Jupiter (white; top row). Neuroblast-derived and GMC-derived Cid clusters are outlined with a blue and orange dashed line, respectively. The white dashed line labels the cell outline. Indistinguishable GMC and Nb Cid clusters are highlighted with yellow dashed circles. The distance of (B) endogenous or (C) ectopic Cid from the wild type hybrid cell shown in (A) in relation to GMC and Nb centrosomes plotted over time. (E) Endogenous or (F) ectopic Cid – centrosome distance measurements of the cnb RNAi expressing hybrid cell shown in (D). The vertical solid line indicates the time point when endogenous and ectopic Cid can no longer be distinguished. Vertical

dashed line indicates nuclear envelope breakdown. Measurements of endogenous Cid – apical

(purple) or basal (yellow) endogenous centrosomes were plotted when (G) the endogenous basal

centrosome matured (0 mins) and 6 minutes thereafter (6mins) or (H) the ectopic centrosomes

matured ('0' mins) and 6 minutes thereafter. Corresponding measurements of *cnb* RNAi expressing

hybrid cells are shown in (J) and (K). Distance of ectopic Cid in relation to endogenous and ectopic

centrosomes shown for (I) wild type or (L) *cnb* RNAi expressing hybrid cells. (M) Time interval

between NEB and mixing of endogenous and ectopic Cid for wild type (blue circles) and *cnb* RNAi

expressing (green circles) hybrid cells. (N) Bar graph showing percentage of hybrid cells in which

the endogenous and ectopic CID mixed prior to NEB. (O) A third instar wild type hybrid cell

expressing the nuclear envelope (NE) marker, koi::GFP (cyan; top row, white; bottom row) and

cherry::Jupiter. (P) Schematic summary: separation of endogenous and ectopic chromosomes

depends on microtubule dependent, asymmetric chromatin-centrosome attachments and physical

separation through nuclear envelopes. *cnb* RNAi expressing neuroblasts release endogenous Cid

during interphase, partially randomizing re-attachment in the subsequent mitosis, which causes

premature mixing of endogenous and ectopic Cid. Colored boxes refer to the indicated cell cycle

stages. Error bars correspond to SDs. Figure (G) paired t-test and Figures (H-L) unpaired t-test. ns

– no signficiance, * $p<0.05$, **$p<0.01$,***$p<0.0001$, **** $p<0.0001$. Time in mins:secs. Scale

bar is 5 μm.

Figure 4: Ectopic chromosomes in hybrid cells segregate independently but erroneously.

(A) The time between chromosome alignment (metaphase) and separation (anaphase onset) was measured for endogenous and ectopic chromosomes in wild type hybrid cells and plotted in (C). (B) Representative examples of delayed (top row) and simultaneous (bottom row) segregation of endogenous (blue dashed circle) and ectopic (orange dashed circle) chromosomes in wild type hybrid cells expressing the chromatin marker His2A::GFP. (C) Quantification of metaphase to anaphase onset in control Nbs, compared to interphase and prophase Nb-GMC hybrids. (D) Average time difference between Nb and GMC chromatin anaphase onset in Nb-GMC hybrid cells. (E) Two representative third instar larval Nb-GMC hybrids expressing histone marker His2A::GFP showing missegregating chromatids (yellow arrows) during anaphase. Time stamps are in relation to NEB (=0:00). (F) Bar graph showing percentage of Nb-GMC hybrid cells with missegregating chromatids. (G) Bar graph quantifying the percentage of fused cells with micronuclei, heterokaryons or synkaryons. ΔT; time difference. Error bars correspond to SDs. One way ANOVA test was used in (C). ns – no significance, **** $p<0.0001$. Time in mins:secs. Scale bar is 5 μm.

Figure 5: Nb – GMC derived hybrid cells neither form visible tumors nor reduce the lifespan in adult hosts.

(A) Third instar larval brains expressing His2A::GFP (top row; negative control), co-expressing *brat* RNAi (middle row; positive control) or containing Nb – GMC hybrid cells (bottom row) were transplanted into wild type hosts and imaged at the day of transplantation (day 0) and Day 30. Yellow arrowheads highlight tumors in wild type hosts transplanted with brat RNAi expressing brains. (B) Scatter plot showing the lifespan of host flies after brain grafts. Grafted tissue either contained no fusions (wild type transplants), expressing *brat* RNAi without fusions (brat RNAi transplants), attempted but unsuccessful fusion (ablation without fusion transplants) and with brains containing hybrid cells (fusion transplants). Host flies without transplanted tissue were plotted for comparison (no transplantation). Error bars correspond to SDs. One way ANOVA. ns – no signficiance, **, ****$p<0.0001$. Scale bar, 1000 μm. (C) Model: Asymmetric MTOCs capture neuroblast chromatin during interphase, thereby establishing a first physical separation between endogenous and ectopic chromatin. Nuclear envelopes form an additional physical barrier. As hybrid cells enter mitosis, the neuroblast centrosomes and introduced ectopic GMC centrosomes nucleate two separate bipolar spindles, aligning their respective chromatin at the metaphase plate. This separation persists through anaphase. Ectopic chromosomes fail to segregate accurately.

Supplemental Data

Fig.S1: Acute induced Nb – GMC fusion in Drosophila larval brains.

(A) Schematic representation of a third instar *Drosophila* larval brain and neural stem cell division mode. Neural stem cells (neuroblasts (Nbs); blue circles) divide asymmetrically, generating a Prospero-positive, differentiating ganglion mother cell (GMC; Pros+) while self-renewing the Dpn+ neuroblast. GMCs cluster around Nbs and can be identified based on size and Prospero expression. Acute Nb – GMC fusion can result in hybrid cells containing two molecularly distinct nuclei. (B) Experimental outline and (C) representative example of a metaphase wild type neuroblast, expressing the cell cortex marker Sqh::EGFP (green), the mitotic spindle marker cherry::Jupiter (white) and the chromatin marker His2A::RFP (white). The site targeted by the ablation laser is highlighted with the white crosshair. The orange arrowhead labels a GMC nucleus moving into the Nb. This hybrid cell successfully completes cytokinesis. (D) Schematic and (E) representative example of a wild type early anaphase neuroblast expressing Sqh::EGFP (white; top, green; bottom), cherry::Jupiter (white) and Pros::EGFP (white). The orange arrowhead highlights a Pros+ GMC nucleus moving into the neuroblast. Cytokinesis completes, creating a hybrid cell containing a Pros+ and Pros- nucleus. Time in min:sec; scale bar: 10μm.

Fig.S2: Microtubule-dependent, asymmetric chromatin-centrosome attachments retain

chromosomes close to the apical neuroblast cortex during interphase.

(A) CID intensity measurements of endogenous and ectopic CID in Nb-GMC hybrid cells (p-value,

0.1294). (B) Representative third instar larval neuroblast expressing the centromere specific

histone-3 variant marker, Cid::EGFP (cyan; top, white; bottom) and the microtubule marker

cherry::Jupiter. Colored boxes represent corresponding cell cycle stages. (C) The distance between

the apical (purple) and basal centrosome (yellow) and individual Cid clusters (green circles), were

measured throughout the cell cycle and plotted in (D). Centrosome – Cid distance of an

unperturbed wild type neuroblast at the time of basal centrosome maturation (0 mins) and 6 mins

thereafter. (F) Representative wild type neuroblast expressing the canonical histone marker

His2A::GFP (white) and Cid::EGFP (cyan; top, white; below). The white dashed line highlights the

cell outline. (G) Representative example of a wild type neuroblast expressing Cid::EGFP (cyan)

and the microtubule marker cherry::Jupiter treated with colcemid. Blue and yellow arrowheads

highlight the position of Cid clusters and the disappearing apical MTOC, respectively. (H) MTOC

intensity and Cid location measurements for the cell shown in (G). (I) Representative third instar

larval neuroblast expressing *cnb* RNAi, cherry::Jupiter (white; top) and Cid::EGFP (cyan; top,

white; bottom row). Red arrowheads highlight the disappearing apical MTOC. Blue arrowheads

highlight the maturing basal MTOC. Blue dashed circle highlight Cid clusters. The white dashed

line highlights the cell outline. (J) Cid – centrosome distance measurement for the cell shown in

(I). 'Apical' centrosome refers to the centrosome destined to be positioned on the apical cortex,

whereas 'basal' centrosome will be inherited by the basal GMC. (K) Wild type neuroblasts maintain

apical CS – Cid attachments in prophase, due to asymmetric MTOC activity and microtubule

dependent interphase centrosome – Cid attachments. Cnb RNAi expressing neuroblasts lose MTOC

activity in interphase, randomizing the position of Cid clusters. When centrosomes mature again

in prophase, both centrosomes simultaneously attach to Cid clusters. (L, M) Cid – centrosome

distance measurement in *cnb* RNAi expressing neuroblasts. Error bars correspond to SDs. Fig.

(A,E,L,M) -paired t-test. ns – no signficiance, * $p<0.05$. Time in mins:secs. Scale bar is 5 μm.

Fig.S3: Asymmetric MTOCs and nuclear envelopes separate endogenous and ectopic chromatin in Nb – GMC hybrid cells.

(A) Averaged distance of basal centrosomes to *endogenous* Cid in wild type or (C) *cnb* RNAi expressing neuroblasts. (B) Averaged distance of basal centrosomes to *ectopic* Cid in wild type or (D) *cnb* RNAi expressing neuroblasts. Measurements are plotted for 0 mins (ectopic centrosome maturation for wild type; ectopic centrosome maturation in cnb RNAi) and 6 mins. Representative image sequence of a wild type (E) unfused neuroblasts or (F) hybrid cell, expressing the nuclear envelope marker koi::GFP (cyan; top, white; bottom) and the microtubule marker cherry::Jupiter (white; top). Error bars correspond to SDs. Fig. (A-D) unpaired t-test. ns – no signficiance, ** p<0.01, ***p<0.001, ****p<0.0001. Time in mins:secs. Scale bar is 5 μm.

Supplemental Table 1: Statistical information

Complete statistical information for the data shown in the corresponding Figures.

Methods

Fly Strains:

Transgenes and fluorescent markers: *worGal4, UAS-mCherry::Jupiter* (Cabernard and Doe, 2009)*;*

worGal4, UAS-mCherry::Jupiter, Sqh::GFP (Cabernard et al., 2010)*; His2A::GFP* (Bloomington

stock center); *UAS-mCherry::CAAX, UAS-iLID::CAAX:;mCherry (A. Monnard & C. Cabernard;*

unpublised); Cid::EGFP (Ranjan et al., 2019); *pUbq-Asl::GFP* (Blachon et al., 2008); *worgal4, UAS-*

mCherry::Jupiter, Asl::GFP (this work); *pros::EGFP* (endogenously tagged with CRISPR; this

work); *koi::GFP (CB04483)* (Buszczak et al., 2007), *cnb^{GD11735}* RNAi line (v28651) (Dietzl et al.,

2007).

Generation of pros::EGFP with CRISPR:

Target specific sequences with high efficiency were chosen using the CRISPR Optimal Target

Finder (http://tools.flycrispr.molbio.wisc.edu/targetFinder/), the DRSC CRISPR finder

(http://www.flyrnai.org/crispr/), and the Efficiency Predictor (http://

www.flyrnai.org/evaluateCrispr/) web tools. Sense and antisense primers for these chosen sites

were then cloned into pU6-BbsI-ChiRNA (Gratz et al., 2013) between BbsI sites. To generate the

replacement donor template, EGFP and 1 kb homology arms flanking the insertion site were cloned

into pHD-DsRed-attP (Addgene plasmid #51019) using Infusion technology (Takara/Clontech).

Injections were performed in house. Successful events were detected by DsRed-positive screening

in the F1 generation. Constitutively active Cre (BDSC#851) was then crossed in to remove the

DsRed marker. Positive events were then balanced, genotyped, and sequenced.

Live cell imaging acute cell-cell fusion:

Imaging medium consists of Schneider's insect medium (Sigma-Aldrich S0146) mixed with 10%

BGS (HyClone). Third instar larvae were dissected in imaging medium and the brains were

transferred into a μ-slide Angiogenesis or μ-slide 8 well (Ibidi). Live samples were imaged with an

Intelligent Imaging Innovations (3i) spinning disc confocal system, consisting of a Yokogawa CSU-

W1 spinning disc unit and two Prime 95B Scientific CMOS cameras. A 60x/1.4NA oil immersion

objective mounted on a Nikon Eclipse Ti microscope was used for imaging. Live imaging voxels

are 0.22 X 0.22 X 0.75-1μm (60x/1.4NA spinning disc).

Neuroblast-GMC fusions were induced using a 3i Ablate! ablation system, consisting of a 532nm

pulsed laser. We used a pulse width of 7 ns, targeting the membrane interface between the

neuroblast and the adjacent GMC.

Colcemid treatment:

Dissected brains were incubated with Colcemid (Sigma) in live imaging medium at a final

concentration of of 25 μgmL^{-1}.

Transplantation experiments:

Brain lobes containing the hybrid cells expressing *His2A::GFP* and *worGal4, UAS-mCherry::Jupiter*

were transplanted into 3 to 4 day old, well fed adult *w^{1118}* female host flies as described previously

(Rossi and Gonzalez, 2015). Custom made needles were prepared from Narishige GD-1 glass

capillaries using a Narishige, needle puller. Injection needles were shaped with forceps to have a

smooth, 45° opening. Transplanted flies were transferred into fresh vials each day for the first three

days, followed by biweekly flipping. The tumor growth was monitored and recorded with a Leica

MZ FLIII fluorescence stereomicroscope.

Image processing and measurements:

Live cell images were processed using imaris x64 8.3.1 and image J.

For angle and distance measurements, the coordinates for the two spindle poles were determined

in Imaris. From these coordinates, angles and distances between spindles were derived based on

the calculations outlined below.

Angle between spindles: $\theta = \cos^{-1}\dfrac{n \cdot e}{|n||e|}$

Dot product: $n \cdot e = (X1*X2) + (Y1*Y2) + (Z1*Z2)$

Magnitude of vectors: $|n| = \sqrt{X_1^2 + Y_1^2 + z_1^2}$ $|e| = \sqrt{X_2^2 + Y_2^2 + z_2^2}$

Where **n** corresponds to the spindle vector: **n** (x1,y1,z1) = (N1-N1', N2-N2', N3-N3') and **e** to the

ectopic spindle vector: **e** (x2,y2,z2) = (E1-E1', E2-E2', E3-E3')

N1, N2, N3 and N1', N2' and N3' are coordinates of the two poles of the Nb spindle. Similarly, E1,

E2, and E3 and E1', E2' and E3' are coordinates of the ectopic spindle poles.

Distance between spindle vectors:

The midpoints of the two spindle vectors are calculated from coordinates of the poles on either

side of the respective spindle. This is followed by calculating the distance between these midpoints.

Midpoint of the Nb spindle vector $= \left(\dfrac{N_1+N_1^1}{2}, \dfrac{N_2+N_2^1}{2}, \dfrac{N_3+N_3^1}{2}\right) = (M_1, M_2, M_3)$

Midpoint of the GMC spindle vector $= \left(\dfrac{G_1+G_1^1}{2}, \dfrac{G_2+G_2^1}{2}, \dfrac{G_3+G_3^1}{2}\right) = (m_1, m_2, m_3)$

Distance between these two points $= \sqrt{(M_1-m_1)^2 + (M_2-m_2)^2 + (M_3-m_3)^2}$

Centrosome - Cid distance:

The centrosome (CS) - Cid distance was calculated using Cid and CS coordinates.

CS – Cid distance: $= \sqrt{(x_1-x_2)^2 + (y_1-y_2)^2 + (z_1-z_2)^2}$

Where x_1,y_1,z_1 correspond to CS and x_2,y_2,z_2 to Cid coordinates, respectively.

Plotted values correspond to averaged values of all CS – Cid punctae distances and the

corresponding standard deviations.

0 and 6 mins corresponds to the appearance of the basal centrosome and 6 minutes thereafter.

'0' mins and '6' mins corresponds to the appearance of the ectopic centrosome appearance and 6

minutes thereafter.

Statistical analysis:

Statistical analysis was performed using Graphpad prism 8. Statistical significance was determined

using paired or unpaired t-test and one-way ANOVA. Significance was indicated as following: *;

$p < 0.05$, **;$p < 0.01$, ***;$p < 0.001$,****$p < 0.0001$, ns; not significant. Exact p values and complete

statistical information can be found in Extended data table 1.

Methods references

Blachon, S., J. Gopalakrishnan, Y. Omori, A. Polyanovsky, A. Church, D. Nicastro, J. Malicki,
and T. Avidor-Reiss. 2008. Drosophila asterless and vertebrate Cep152 Are orthologs
essential for centriole duplication. *Genetics*. 180:2081 2094.
doi:10.1534/genetics.108.095141.

Buszczak, M., S. Paterno, D. Lighthouse, J. Bachman, J. Planck, S. Owen, A.D. Skora, T.G.
Nystul, B. Ohlstein, A. Allen, J.E. Wilhelm, T.D. Murphy, R.W. Levis, E. Matunis, N.
Srivali, R.A. Hoskins, and A.C. Spradling. 2007. The carnegie protein trap library: a versatile
tool for Drosophila developmental studies. *Genetics*. 175:1505 1531.
doi:10.1534/genetics.106.065961.

Cabernard, C., and C.Q. Doe. 2009. Apical/basal spindle orientation is required for neuroblast
homeostasis and neuronal differentiation in Drosophila. *Dev Cell*. 17:134 141.
doi:10.1016/j.devcel.2009.06.009.

Cabernard, C., K.E. Prehoda, and C.Q. Doe. 2010. A spindle-independent cleavage furrow
positioning pathway. *Nature*. 467:91 94. doi:10.1038/nature09334.

Dietzl, G., D. Chen, F. Schnorrer, K.-C. Su, Y. Barinova, M. Fellner, B. Gasser, K. Kinsey, S.
Oppel, S. Scheiblauer, A. Couto, V. Marra, K. Keleman, and B.J. Dickson. 2007. A genome-
wide transgenic RNAi library for conditional gene inactivation in Drosophila. *Nature*.
448:151 156. doi:10.1038/nature05954.

Gratz, S.J., A.M. Cummings, J.N. Nguyen, D.C. Hamm, L.K. Donohue, M.M. Harrison, J.
Wildonger, and K.M. O'Connor-Giles. 2013. Genome Engineering of Drosophila with the
CRISPR RNA-Guided Cas9 Nuclease. *Genetics*. 194:1029–1035.
doi:10.1534/genetics.113.152710.

Ranjan, R., J. Snedeker, and X. Chen. 2019. Asymmetric Centromeres Differentially Coordinate
with Mitotic Machinery to Ensure Biased Sister Chromatid Segregation in Germline Stem
Cells. *Cell Stem Cell*. doi:10.1016/j.stem.2019.08.014.

Rossi, F., and C. Gonzalez. 2015. Studying tumor growth in Drosophila using the tissue allograft
method. *Nat Protoc*. 10:1525–1534. doi:10.1038/nprot.2015.096.

727
728

Acknowledgements

We thank Xin Chen for fly stocks, David Salvador Garcia for generating the Pros::EGFP transgenic

line, Sue Biggins and members of the Cabernard laboratory for helpful discussions and comments.

This work was supported by the National Institutes of Health (1R01GM126029-03) and a Research

Scholar grant from the American Cancer Society (130285-RSG-16253-01-CSM). Stocks obtained

from the Bloomington Drosophila Stock Center (NIH P40OD018537) and from the Vienna

Drosophila Resource Center (VDRC).

Author contributions

This study was conceived by B.S., N.L. and C.C. B.S and N.L performed all the experiments with

significant help from C.S.

B.S, N.L., C.S., and C.C analyzed the data. B.S and C.C. wrote the manuscript.

Competing interest declaration

The authors declare no competing financial interests.

Additional information

This paper contains three Fig. Ss, a supplementary table and 10 videos to support the main

conclusions.

748
749

Video legends

Video 1: Wild type neuroblast division; related to Fig. 1B

Wild type control (unfused) neuroblast expressing the microtubule binding protein Cherry::Jupiter

(white) and the canonical Histone marker His2A::GFP (cyan). Time scale is h:mm:ss and the scale

bar is 5 µm

Video 2: Wild type hybrid cell; related to Fig. 1B

Wild type hybrid cell derived from a neuroblast – GMC fusion *in vivo*, expressing the canonical

Histone marker His2A::GFP (white in single channel; cyan in merge) and the microtubule binding

protein Cherry::Jupiter (white). The blue and orange arrows mark endogenous and ectopic

chromatin, respectively. Time scale is h:mm:ss and the scale bar is 3 µm.

Video 3: Wild type hybrid cell; related to Fig. 2B

Wild type hybrid cell derived from a neuroblast – GMC fusion *in vivo*, expressing the canonical

Histone marker His2A::GFP (white in single channel; cyan in merge) and the microtubule binding

protein Cherry::Jupiter (white). The blue and orange arrows mark endogenous and ectopic

chromatin, respectively. Time scale is h:mm:ss and the scale bar is 2 µm.

Video 4: Wild type neuroblast division; related to Fig. S2B

Wild type control (unfused) neuroblast, expressing the microtubule binding protein

Cherry::Jupiter (white) and the centromere-specific H3 variant Cid::EGFP (Cyan). Purple and

yellow arrows point to the apical and basal centrosome, respectively. The blue arrow refers to

moving Cid clusters. Time scale is h:mm:ss and the scale bar is 1 µm.

Video 5: Wild type neuroblast division; related to Fig. S2F

Wild type control (unfused) neuroblast, expressing the membrane marker mCherry::CAAX

(white), the canonical Histone marker His2A::GFP (white) and Cid::EGFP (Cyan). The green arrow

points to Cid clusters. Time scale is h:mm:ss and the scale bar is 5 µm.

Video 6: Wild type neuroblast exposed to Colcemid; related to Fig. S2G

Wild type control (unfused) neuroblast exposed to the microtubule depolymerizing drug Colcemid,

expressing the membrane marker mCherry::CAAX (white), the microtubule binding protein

Cherry::Jupiter (white) and Cid::EGFP (Cyan). The yellow arrow points to the apical centrosome,

the blue arrow to Cid clusters. Time scale is h:mm:ss and the scale bar is 1 µm.

Video 7: Cnb RNAi expressing neuroblast; related to Fig. S2I

Cnb RNAi expressing (unfused) neuroblast, co-expressing the microtubule binding protein

Cherry::Jupiter (white) and Cid::EGFP (Cyan). The orange and blue arrow points to the apical and

basal centrosome, respectively. The green arrow highlights Cid clusters. Time scale is h:mm:ss and

the scale bar is 1 µm.

Video 8: wild type hybrid cell; related to Fig. S3A

Wild type hybrid cells expressing the microtubule binding protein Cherry::Jupiter (white) and

Cid::EGFP (white in single channel; Cyan in merge). The blue and orange arrow highlights

endogenous and ectopic Cid clusters, respectively. Time scale is h:mm:ss and the scale bar is 1 µm.

Video 9: Cnb RNAi expressing hybrid cell; related to Fig. S3D

Cnb RNAi expressing hybrid cell, expressing the microtubule binding protein Cherry::Jupiter

(white) and Cid::EGFP (white in single channel; Cyan in merge). The blue and orange arrow

highlights endogenous and ectopic Cid clusters, respectively. The yellow arrow indicates mixing of

endogenous and ectopic Cid clusters. Time scale is h:mm:ss and the scale bar is 1 µm.

Video 10: wild type hybrid cell; related to Fig. 4E

Wild type hybrid cell, expressing the canonical Histone marker His2A::GFP (white). The blue and

orange arrow highlights endogenous and ectopic chromosomes, respectively. The magenta

arrowhead highlights the fate of missegregated chromosomes. This hybrid cell forms a

heterokaryon. Time scale is h:mm:ss and the scale bar is 1 µm.