Asymmetric chromatin capture and nuclear envelopes separate chromosomes in

fused cells during mitosis

Bharath Sunchu [1], Nicole Lee [1,2], Roberto Carlos Segura[1], Chantal Roubinet[3] & Clemens

Cabernard [1]

[1] Department of Biology, University of Washington

Life Science Building

Seattle, Washington State, USA

Current address:

[2] Cancer Science Institute, National University of Singapore, Singapore

[3] MRC-Laboratory for Molecular Cell Biology, University College London, London, UK

[*] Lead and corresponding author: ccabern@uw.edu

Abstract

Hybrid cells derived through fertilization or somatic cell fusion recognize and separate chromosomes of different origin. The underlying mechanisms are unknown but could prevent aneuploidy and tumor formation. Here, we acutely induce fusion between *Drosophila* neural stem cells (neuroblasts; Nbs) and differentiating ganglion mother cells (GMCs) *in vivo* to define how epigenetically distinct chromatin is recognized and segregated. We find that Nb-GMC hybrid cells align both endogenous (neuroblast-origin) and ectopic (GMC-origin) chromosomes at the metaphase plate through centrosome derived dual-spindles. Mixing of endogenous and ectopic chromatin is prevented through an asymmetric, microtubule-dependent chromatin capture mechanism during interphase and physical boundaries imposed by nuclear envelopes. Although hybrid cells fail to accurately segregate ectopic chromatin, hybrid cells neither reduce the lifespan nor form visible tumors in host flies. We propose that Nb-GMC derived hybrid cells utilize asymmetric centrosome activity in interphase and nuclear envelopes to physically separate epigenetically distinct chromatin.

Introduction

Dividing cells equally distribute the replicated chromosomes between the two sibling cells through microtubule-dependent attachment and segregation mechanisms (Maiato et al., 2017; McIntosh, 2016). Microtubules of bipolar spindles are connected to chromosomes via kinetochore proteins, which are localized on centromeric DNA (Thomas et al., 2017; Yu et al., 2019). Mitotic metazoan cells usually only form a single bipolar spindle, but certain insect species, arthropods or mouse zygotes form two distinct mitotic spindles (dual-spindles hereafter), which physically separate the maternal from the paternal chromatin in the first division after fertilization (Kawamura, 2001; Reichmann et al., 2018; Snook et al., 2011). Chromosome separation also occurs in hybrid cells, derived from somatic cell-cell fusion events (Heasley et al., 2017; Rieder et al., 1997). Dual-spindle dependent chromosome separation suggests the presence of specific chromosome recognition mechanisms, distinguishing between epigenetically distinct chromatin. The molecular nature of these recognition mechanisms is not known but could entail asymmetries in centromere binding proteins, kinetochore size or kinetochore composition (Akera et al., 2019; Arco et al., 2018; Drpic et al., 2018; Ranjan et al., 2019).

Over a century ago, unregulated cell-cell fusion between different somatic cells has been proposed to initiate tumor formation (Aichel, 1911). Aichel's cell fusion model has the advantage that it can readily explain aneuploidy, a feature frequently observed at the early stages of tumor development (Ogle et al., 2005; Platt and Cascalho, 2019). Tetraploidy and supernumerary centrosomes – the natural products of cell fusion – predispose cells to aneuploidy through chromosome rearrangements (Fujiwara et al., 2005). Aichel's cell fusion model still remains to be experimentally validated, which requires a detailed characterization of chromosome dynamics in fused cells *in vivo*.

Here, we ask how hybrid cells derived through cell-cell fusion of molecularly distinguishable cell types accurately recognize, separate and segregate epigenetically distinct chromosomes. To this end, we acutely fused *Drosophila* neural stem cells (neuroblasts (NBs), hereafter) with differentiating ganglion mother cells (GMCs) in the intact larval fly brain to create hybrid cells containing both neuroblast and GMC chromosomes. Unperturbed *Drosophila* neuroblasts divide asymmetrically, self-renewing the neural stem cell while forming a differentiating GMC. Neuroblasts are twice the size of GMCs, express the transcription factor Deadpan (Dpn[+]) and divide asymmetrically with a rapid cell cycle. The smaller GMCs can also be identified based on Prospero (Pros[+]) expression and divide only once with a long cell cycle (Gallaud et al., 2017).

Neuroblasts are intrinsically polarized, consisting of an apically localized Par complex, which is

connected to the Pins complex, composed of Partner of Inscuteable (Pins; LGN/AGS3 in

vertebrates), Gαi and Mushroom body defects (Mud; NuMA in vertebrates, Lin-5 in C. elegans)

(Gallaud et al., 2017; Loyer and Januschke, 2020; Sunchu and Cabernard, 2020). The Pins

complex regulates spindle orientation during mitosis and biased centrosome asymmetry in

interphase, manifested in the establishment and maintenance of an apical interphase microtubule

organizing center (MTOC). The active interphase MTOC retains the daughter-centriole containing

centrosome close to the apical cell cortex and pre-establishes spindle orientation in the

subsequent mitosis (Gallaud et al., 2020; Gambarotto et al., 2019; Januschke et al., 2013, 2011;

Januschke and Gonzalez, 2010; Rebollo et al., 2007; Rusan and Peifer, 2007).

We found that hybrid cells derived from such Nb-GMC fusions independently align the

endogenous (neuroblast-origin) and ectopic (GMC-origin) chromosomes at the metaphase plate.

We propose that these hybrid cells utilize asymmetric centrosome activity in interphase to capture,

and nuclear envelopes to physically separate, epigenetically distinct chromatin *in vivo.* These

findings provide mechanistic insight into how metazoan cells recognize and isolate chromosomes

of different origin.

Results

Nb-GMC hybrid cells independently align Nb and GMC chromatin at the metaphase plate

To quantitatively describe chromosome dynamics in hybrid cells we developed an acute cell-cell fusion method in intact larval fly brains. We used a 532nm pulsed laser to induce a small lesion at the Nb – GMC interface, causing the GMC chromatin to enter the neuroblast cytoplasm. Neuroblasts can be distinguished from GMCs based on their size, molecular markers and cell cycle length (Figure 1 – figure supplement 1A). Targeted mitotic neuroblasts often retained the GMC chromosomes, creating a large apical hybrid cell containing one Dpn+ and one Pros+ nucleus (Nb – GMC hybrid). Most Nb – GMC hybrid cells normally localized the contractile ring marker non-muscle Myosin to the cleavage furrow and completed cytokinesis (Figure 1 – figure supplement 1B-E). Acute cell fusion can also result in the expulsion of the GMC chromatin, forming GMC – GMC hybrids (see Supplementary Figure 3c in (Roubinet et al., 2017)). Here, we are focusing on Nb – GMC hybrids (hybrid cells, hereafter) only.

To better characterize the dynamics of neuroblast (endogenous) and GMC (ectopic) chromosomes during mitosis, we induced cell fusion at different cell cycle stages in wild type neuroblasts, expressing the canonical chromosome marker His2A::GFP. We hypothesized that hybrid cells derived from Nb-GMC fusions early in the cell cycle could (1) align only the neuroblast chromosomes at the metaphase plate, (2) congress a mix of neuroblast and GMC chromosomes or (3) separate and align the two chromosome pools at the metaphase plate (Figure 1A). We found that the endogenous and ectopic chromatin was separated and distinguishable when fusions were induced in early mitosis. Both the ectopic and endogenous chromatin aligned at the metaphase plate (Figure 1B; video 1&2). Nb-GMC fusions could be induced at all cell cycle stages but GMC chromosomes aligned at the metaphase plate more accurately in hybrid cells derived from interphase or early prophase fusions (Figure 1C, D). All fusions reported here were performed with non-mitotic GMCs. We conclude that hybrid cells derived from fusions between interphase Nbs and non-mitotic GMCs accurately align ectopic and endogenous chromatin at the metaphase plate.

We next asked whether GMC chromatin congresses independently of neuroblast chromatin. To this end, we measured the time between nuclear envelope breakdown (NEB) and chromosome alignment at the metaphase plate for Nb and ectopic GMC chromatin in hybrid cells derived from interphase and early prophase fusions (Figure 1E). In most hybrid cells, ectopic and endogenous chromatin was distinguishable based on differences in location and intensity (see

Figure 1B; video 1&2). In untargeted control neuroblasts (unfused), Nb chromosomes aligned at

the metaphase plate within 6.5 minutes after NEB (SD = 1.55; n = 6), which is insignificantly faster

than the neuroblast chromosomes of hybrid cells (t = 6.9 mins; SD = 2.84; n = 11). GMC

chromosomes aligned within 7.63 mins (SD = 3.69; n = 11), statistically not significantly different

from unperturbed wild type chromosomes (Figure 1E). In most Nb-GMC hybrids, the endogenous

neuroblast and the ectopic GMC chromosomes aligned at the metaphase plate with no significant

time difference. However, in a few cases, ectopic chromatin aligned before or after the neuroblast

chromatin (Figure 1F). These results suggest that the neuroblast and GMC chromatin can move

independently to the metaphase plate in hybrid cells.

Ectopic spindles distinguish between Nb and GMC chromatin

We next investigated the mechanisms underlying independent Nb/GMC chromosome alignment,

considering the following possibilities: ectopic chromosomes could be aligned together with the

endogenous chromosomes via a single bipolar spindle. Alternatively, hybrid cells could form

multiple bipolar spindles, which attach to either the neuroblast's, GMC's, or chromosomes from

both cell types (Figure 2A). Live cell imaging showed that hybrid cells derived from interphase

Nb-GMC fusions contained double spindles in almost all cases, whereas the vast majority of cell

fusions induced in metaphase only formed one mitotic spindle (Figure 2B, C & video 3). Most

hybrid cells contained two mitotic spindles, which typically formed at the same time (Figure 2D,

E). We quantified spindle alignment and positioning dynamics by measuring the angle and

distance between the two spindles during metaphase (Figure 2F and methods). The two spindles

were initially misaligned and separated but decreased their inter-spindle distance and angle

during metaphase (Figure 2G-I). We conclude that Nb-GMC hybrid cells align neuroblast and

GMC chromosomes separately at the metaphase plate through the formation of independent

mitotic spindles.

Ectopic spindles are nucleated from GMC centrosomes

Mitotic spindles can be nucleated through the centrosome-dependent, chromatin or microtubule

pathway but when centrosomes are present, the centrosome pathway prevails (Prosser and

Pelletier, 2017). To elucidate the mechanisms underlying ectopic spindle formation, we induced

Nb-GMC fusions in interphase wild type neuroblasts expressing live centriole (Asterless;

Asl::GFP) and spindle (cherry::Jupiter) markers, and assayed centrosome dynamics and spindle

formation throughout mitosis. Normal wild type neuroblasts usually contained two Asl::GFP

positive centrioles in mitosis, forming a single bipolar spindle. However, in Nb-GMC hybrids, we

predominantly found four Asl::GFP positive centrioles, two of which were introduced from the

GMC (Figure 2J,K). The GMC centrosomes nucleated an ectopic bipolar spindle that

subsequently aligned with the main neuroblast spindle. Although multipolar spindle formation can

be prevented through centrosome clustering (Quintyne et al., 2005), we often observed that GMC-

derived and Nb-derived spindles were oriented in parallel to each other but remained separate.

These data suggest that ectopic spindles are formed through the centrosome pathway, using

centrioles originating from GMCs.

Microtubule-dependent, asymmetric chromatin-centrosome attachments retain

chromosomes close to the apical neuroblast cortex during interphase

Our data suggest that Nb and GMC chromatin are being separated through an endogenous, Nb-

derived and an ectopic, GMC-derived mitotic spindle. We next investigated how these spindles

distinguish between Nb and GMC chromosomes. During mitosis, microtubules emanate from

centrosomes and attach to sister chromatids via kinetochore proteins, localizing to the

centromeric region (Fukagawa and Earnshaw, 2014). *Drosophila* male germline stem cells

contain asymmetric levels of the centromere-specific H3 variant (Centromere identifier (Cid) in

flies (Henikoff et al., 2000) (Ranjan et al., 2019), prompting us to investigate whether hybrid cell

spindles differentiate between endogenous and ectopic chromosomes based on differing levels

of Cid. We induced Nb-GMC fusions of wild type cells expressing Cid::EGFP (Ranjan et al., 2019)

in interphase or early prophase and measured Cid intensity on both GMC and Nb chromatin.

These measurements did not reveal a significant intensity difference between Nb and GMC Cid

(Figure 3 – figure supplement 1A). However, we noticed that endogenous Cid::EGFP was

localized in very close proximity to the apical centrosome in unperturbed interphase and prophase

wild type neuroblasts (Figure 3A & video 4). Cid::EGFP remained associated with chromatin

throughout the neuroblast cell cycle, excluding the possibility that early Cid clusters are not

connected with chromatin (Figure 3 – figure supplement 1B & video 5).

Interphase wild type neuroblasts contain only one active apical microtubule organizing center

(MTOC), which retains the daughter centriole-containing centrosome close to the apical

neuroblast cortex. The mother-centriole-containing centrosome is inactive in interphase but

matures from prophase onward, positioning itself on the basal cell cortex (Gallaud et al., 2020;

Januschke et al., 2013, 2011; Januschke and Gonzalez, 2010). We measured the distance of

individual Cid::EGFP clusters to the apical and basal centrosome in unperturbed wild type

neuroblasts and found that during interphase Cid was always in close proximity to the apical

centrosome. After nuclear envelope breakdown (NEB), Cid moved progressively towards the

metaphase plate. Once the basal centrosome appeared (0 mins), Cid was still closer to the apical

than the basal centrosome and this distance asymmetry was also observed 6 minutes after the

appearance of the basal centrosome (Figure 3B, C, H).

The proximity of Cid::EGFP clusters to the active interphase MTOC suggests a microtubule-

dependent chromosome attachment mechanism. Indeed, wild type neuroblasts treated with the

microtubule-depolymerizing drug colcemid showed a strong correlation between apical MTOC

activity and Cid localization; as MTs depolymerized after colcemid addition, Cid progressively

moved away from the apical cortex towards the cell center (Figure 3 – figure supplement 1C, D &

video 6). To test whether MTOCs are connected to Cid clusters during interphase, we removed

the centriolar protein Centrobin (Cnb; CNTROB in humans). Neuroblasts lacking Cnb fail to

maintain an active apical interphase MTOC but regain normal MTOC activity during mitosis

(Januschke et al., 2013). Neuroblasts expressing *cnb* RNAi lost apical Cid localization after the

apical centrosome downregulated its MTOC activity. However, maturing centrosomes

reconnected with Cid in prophase (Figure 3D-G & video 7). Cid's proximity to the apical MTOC

('apical' refers to the centrosome destined to move to the apical cortex) in *cnb* RNAi expressing

neuroblasts was much more varied compared to wild type. At 6 minutes after centrosome

maturation onset, Cid – apical MTOC distance was comparable to wild type, as were Cid – basal

centrosome distance relationships (Figure 3I, J). This suggests that when centrosomes mature

during mitosis and MTOC activity is restored, Cid – MTOC attachments can be reestablished

during early mitosis in *cnb* RNAi expressing neuroblasts (Figure 3G). We conclude that in wild

type Nbs, Cid-containing chromatin is already attached to the apical centrosome prior to entry into

mitosis.

Asymmetric centrosome-chromatin attachments contribute to the separation of

endogenous and ectopic chromosomes

Based on these observations, we hypothesized that the separation between endogenous and

ectopic chromosomes could be due to a pre-attachment mechanism, preventing mixing of Nb and

GMC chromosomes. To test this hypothesis, we first analyzed Cid localization in relation to the

endogenous and ectopic centrosomes in wild type hybrid cells. Similar to unperturbed wild type

neuroblasts, endogenous Cid is also localized in close proximity to the endogenous apical

centrosome in wild type hybrid cells (Figure 4A-C, G, H & video 8; 'apical' refers to the centrosome

destined to segregate into the large apical sibling cell). Ectopic Cid, however, appeared closer to

ectopic centrosomes (Figure 4A-C, I & video 8; 0 mins refers to the appearance of the

endogenous basal centrosome; '0' refers to the appearance of the ectopic centrosomes).

We next attempted to randomize Cid – centrosome distance relationships by inducing fusions in

cnb RNAi expressing neuroblasts, since loss of interphase MTOC activity released endogenous

Cid from the apical centrosome (Figure 3D-J and Figure 3 - figure supplement 1C, D). In contrast

to wild type hybrid cells, endogenous Cid is roughly equidistant to the endogenous and ectopic

centrosomes in *cnb* RNAi expressing hybrid cells at 0 mins and '0' mins respectively (Figure 4D-

F, J, K & video 9). However, ectopic Cid was still closer to the ectopic centrosomes than to the

endogenous apical centrosome in *cnb* RNAi expressing hybrid cells (Figure 4D, F, L & video 9).

In both wild type and *cnb* RNAi expressing hybrid cells, ectopic and basal centrosomes were

about equidistant to endogenous Cid, but ectopic centrosomes were closer to ectopic Cid (Figure

4 - figure supplement 1A-D). Taken together, we conclude that in hybrid cells the apical MTOC

forms an asymmetric attachment to endogenous Cid-containing chromosomes prior to entry into

mitosis.

Nuclear envelopes separate ectopic and endogenous chromatin in hybrid cells

We next asked whether early MTOC – Cid attachments are sufficient to prevent endogenous and

ectopic chromosome separation and tracked endogenous and ectopic Cid::EGFP after induced

cell fusion. In wild type hybrid cells, endogenous and ectopic CID clusters started to congress at

the metaphase plate and became difficult to clearly separate 8.8 mins (SD = 5.63; n = 5; Figure

4M) after NEB. However, 50% of *cnb* RNAi hybrid cells showed endogenous and ectopic Cid

mixing prior to NEB (Figure 4D, N), although the time difference was not significantly different to

wild type hybrid cells (Figure 4M).

Neuroblasts undergo semi-closed mitosis, mostly retaining a matrix composed of nuclear

envelope proteins around the mitotic spindle (Katsani et al., 2008). We imaged hybrid cells with

the nuclear envelope marker Klaroid, using the protein-trap line koi::EGFP (Buszczak et al.,

2007). We confirmed that unperturbed wild type neuroblasts contain a nuclear envelope matrix

surrounding the mitotic spindle during mitosis (Figure 4 – figure supplement 1E). Similarly, wild

type hybrid cells contain two nuclear envelopes during mitosis (Figure 4O and Figure 4 – figure

supplement 1F). Taken together, these data suggest that both asymmetric MTOC-Cid

attachments in interphase and nuclear envelopes establish and maintain the physical separation

between endogenous and ectopic chromosomes in hybrid cells. Loss of biased interphase MTOC

activity removes asymmetric MTOC-Cid attachments and allows for cross-connections between

endogenous centrosomes and ectopic Cid in early mitosis (Figure 4P).

Hybrid cells segregate endogenous and ectopic chromosomes independently

We next investigated whether both bipolar spindles are functional in Nb-GMC hybrid cells.

Erroneous or incomplete microtubule-kinetochore attachments trigger the spindle assembly

checkpoint (SAC), preventing or delaying anaphase entry (Musacchio, 2015). Since the

kinetochore-derived 'wait anaphase' signal is diffusible (Heasley et al., 2017), ectopic spindles

should thus also obey the SAC in Nb-GMC hybrid cells. We tested whether hybrid cells contain

functional microtubule-kinetochore attachments by measuring the time between finished

chromosome alignment at the metaphase plate and chromosome separation in Nb-GMC hybrids

expressing His2A::GFP and cherry::Jupiter (Figure 5A, B). Unperturbed control neuroblasts

usually initiate anaphase onset within 2.63 minutes (SD=1.84; n=12) after chromosomes are

aligned at the metaphase plate. In hybrid cells derived from interphase fusions, endogenous and

ectopic chromatin entered anaphase 4.14 mins (SD= 2.23; n=14) and 5.12 minutes (SD= 1.85;

n=13) after metaphase alignment. Only ectopic chromatin for prophase-induced hybrid cells

showed a significantly delayed anaphase onset (Average: 10.25 mins; SD= 4.12; n=6) (Figure

5C). Ectopic chromatin never separated before endogenous chromosomes but entered anaphase

with a few minutes' delay (Figure 5D). We conclude that in Nb-GMC hybrids, endogenous and

ectopic chromosomes establish correct MT-kinetochore attachments, thereby fulfilling the spindle

assembly checkpoint necessary to enter anaphase. However, given the delays in ectopic

chromosome separation, we further conclude that ectopic spindles can initiate chromatid

separation independently from the endogenous neuroblast spindle.

Hybrid cells are insufficient to induce tumors in wild type host flies

Finally, we assessed the accuracy of chromosome segregation in wild type hybrid cells. Using

the canonical chromosome marker His2A::GFP we detected chromosome missegregation -

ranging from lagging chromosomes to chromosome bridges - in all wild type hybrid cells (Figure

5E,F & video 10). Chromosome segregation defects can result in aneuploidy and micronuclei

formation (Molina et al., 2020). We found a small percentage of hybrid cells containing

micronuclei, but more frequently discovered heterokaryons (hybrid cells containing two nuclei of

different origins). In most cases, hybrid cells fused both nuclei into one, forming synkaryons

(Figure 5G).

Aneuploidy has been proposed to be a hallmark of cancer (Molina et al., 2020) but appears

to be context dependent (Ben-David and Amon, 2020). To test whether neuroblast – GMC derived

hybrid cells are sufficient to induce tumor formation, we grafted His2A::GFP expressing larval fly

brains after successful induction of cell fusion into the abdomen of wild type adult hosts (Rossi

and Gonzalez, 2015) and monitored the host flies for tumor formation and life span changes. As

previously reported (Caussinus and Gonzalez, 2005), brat RNAi expressing brains formed visible

tumors in host flies by day 30 and caused a reduction in lifespan of the host. However, larval

brains without attempted fusions (wild type transplants), with attempted but unsuccessful fusions

(controlling for the effect of laser ablation), or with successful fusions, showed neither tumor

growth by day 30 (or after), nor a reduction in lifespan (Figure 6A, B, C). We conclude that

chromosome segregation is defective in Nb-GMC hybrid cells but insufficient to form visible

tumors in otherwise normal and unperturbed larval fly brains.

Discussion

Cell – cell fusion can occur under normal physiological conditions and has been implicated in malignancy (Platt and Cascalho, 2019) but how hybrid cells recognize and separate endogenous and ectopic chromosomes during mitosis is not known. Here, we acutely induce cell-cell fusions *in vivo* between Dpn+ stem cells and differentiating Pros+ GMCs in the developing larval fly brain. We showed that Nb-GMC derived hybrid cells physically separate endogenous neuroblast chromosomes from the introduced ectopic GMC chromosomes and align them independently at the metaphase plate. Chromosome separation is achieved through the formation of two distinct mitotic spindles, which are most likely formed through the canonical centrosome pathway. These dual spindles co-align during metaphase, thereby congressing the two chromosome clusters at the metaphase plate. Endogenous and ectopic chromosomes independently segregate during anaphase, manifested in delayed segregation onset of ectopic chromatin. Although chromosome missegregation is frequent in hybrid cells, potentially leading to aneuploidy, we failed to detect malignant tumor formation when hybrid cell – containing larval brains were grafted into wild type host flies.

We propose that endogenous and ectopic chromosome separation is achieved through an early microtubule-dependent chromosome capture or attachment mechanism that retains endogenous chromosomes in close proximity to the apical neuroblast cortex during interphase. Interphase neuroblasts contain one active MTOC that remains stably anchored close to the apical cell cortex (Rebollo et al., 2007; Rusan and Peifer, 2007). In wild type neuroblasts, chromatin associated Cid is localized in close proximity to the apical MTOC, but Cid's apical localization is lost upon microtubule-depolymerization or removal of interphase MTOC activity (*cnb* RNAi). Thus, unperturbed wild type neuroblast either retained, or establish microtubule-chromatin connections during interphase. In contrast to yeast, where chromosomes make dynamic attachments to microtubules in G1 (Dorn et al., 2005), this is not the case in other metazoan cells (Maiato et al., 2017). The functional significance of these interphase microtubule-chromatin attachments in neuroblasts are not known but are similar to *Drosophila* male germline stem cells (GSCs), where a single active centrosome connects to chromosomes in prophase, a potential mechanism for biased chromatid segregation (Ranjan et al., 2019).

The geometric separation of endogenous and ectopic chromatin in hybrid cells is further supported by the nuclear envelope, imposing a physical boundary that prevents random chromosome mixing prior to chromosome congression (Figure 6D). In pre-mitotic neuroblasts,

chromatin can be connected with centrosomes through the Linc complex (Lee and Burke, 2018),

potentially implicating the SUN domain protein Klaroid (Kracklauer et al., 2007) and the KASH-

domain protein Klarsicht (Lee and Burke, 2018; Razafsky and Hodzic, 2009) in asymmetric

chromatin clustering and the prevention of chromatin mixing during interphase.

The chromatin separation mechanisms described here could be applicable to chromosome

separation occurring in the first cleavage after fertilization in insects, arthropods and vertebrates

(Kawamura, 2001; Reichmann et al., 2018; Snook et al., 2011). Similarly, biased chromatid and

chromosome segregation has been observed in stem cells (Ranjan et al., 2019; Yadlapalli and

Yamashita, 2013) and during meiosis, respectively (Akera et al., 2017). Since centromeres have

also been found to be confined to specific nuclear locations in many organisms (Muller et al.,

2019; Weierich et al., 2003), it will be interesting to see whether microtubule-dependent chromatin

attachment provides an alternative mechanism for biased sister chromatid segregation or other

important cellular functions.

Figure 1: Nb-GMC hybrid cells align Nb and GMC chromatin independently at the

metaphase plate.

(A) Potential outcomes of Nb-GMC fusions: Nb – GMC derived hybrid cells could (1) only align

neuroblast chromosomes, (2) congress a mix of endogenous and ectopic chromosomes or (3)

separately align Nb and GMC chromosomes at the metaphase plate. **(B)** Representative image

sequence of a dividing third instar larval Nb-GMC hybrid cell obtained from an interphase fusion,

expressing the histone marker His2A::GFP (dashed blue circle; endogenous chromatin, dashed

orange circle; ectopic chromatin). **(C)** Alignment of endogenous and ectopic chromatin in hybrid

cells derived from interphase (inter), prophase (pro), prometaphase (prometa) or metaphase

(meta) fusions were quantified with angle measurements in metaphase and plotted in **(D)**. **(E)**

Chromosome alignment time for endogenous (blue circles) and ectopic chromosomes (orange

squares) compared to unfused control neuroblasts (grey triangles). **(F)** Scatter plot showing the

time difference (T_a; time of alignment) between endogenous and ectopic chromatin. Colored

boxes represent corresponding cell cycle stages. One-way ANOVA was used for (D) and (E).

Error bars correspond to standard deviation (SD). * p<0.05, **p<0.01, **** p<0.000.1. For this and

subsequent figures, exact p values and complete statistical information can be found in the

Extended data table 1. Time in mins:secs. Scale bar is 5 μm.

Figure 1 – figure supplement 1: Acute induced Nb – GMC fusion in *Drosophila* larval

brains.

(A) Schematic representation of a third instar *Drosophila* larval brain and neural stem cell division

mode. Neural stem cells (neuroblasts (Nbs); blue circles) divide asymmetrically, generating a

Prospero-positive, differentiating ganglion mother cell (GMC; Pros+) while self-renewing the Dpn+

neuroblast. GMCs cluster around Nbs and can be identified based on size and Prospero

expression. Acute Nb – GMC fusion can result in hybrid cells containing two molecularly distinct

nuclei. **(B)** Experimental outline and **(C)** representative example of a metaphase wild type

neuroblast, expressing the cell cortex marker Sqh::EGFP (green), the mitotic spindle marker

cherry::Jupiter (white) and the chromatin marker His2A::RFP (white). The site targeted by the

ablation laser is highlighted with the white crosshair. The orange arrowhead labels a GMC nucleus

moving into the Nb. This hybrid cell successfully completes cytokinesis. **(D)** Schematic and **(E)**

representative example of a wild type early anaphase neuroblast expressing Sqh::EGFP (white;

top, green; bottom), cherry::Jupiter (white) and Pros::EGFP (white). The orange arrowhead

highlights a Pros+ GMC nucleus moving into the neuroblast. Cytokinesis completes, creating a

hybrid cell containing a Pros+ and Pros- nucleus. Time in min:sec; scale bar: 10μm.

Figure 2: Dual spindles align endogenous and ectopic spindles separately at the

metaphase plate.

(A) Hypothetical outcomes of spindle organization after Nb-GMC fusions: hybrid cells could align

neuroblast and GMC chromosomes either through a single or dual-spindle mechanism. **(B)**

Representative third instar larval Nb-GMC hybrid cell, expressing the histone marker His2A::GFP

(white in top row; cyan in merged channel below) and the MT marker cherry::Jupiter (white in

middle and bottom row). 00:00 refers to nuclear envelope breakdown (NEB). **(C)** Quantification

of hybrid cells containing dual- or multiple spindles for hybrid cells derived from interphase (inter),

prophase (pro), prometaphase (prometa) or metaphase (meta) fusions. **(D)** The time difference

between endogenous and ectopic spindle formation was measured in hybrid cells and plotted in

(E). **(F)** Spindle angle and inter-spindle distances were measured during mitosis. A representative

example is shown in **(G)**. Quantification of inter-spindle **(H)** distances and **(I)** angles at

prometaphase and metaphase. **(J)** A representative third instar larval Nb-GMC hybrid, expressing

the centriole marker Asl::GFP (cyan; top row. White; bottom row) and the spindle marker

cherry::Jupiter (MTs; white in top row). Centrosomes were highlighted with green dashed circles.

(K) Comparison of centrosome number between unfused wild type and hybrid cells.

Colored boxes represent corresponding cell cycle stages. Error bars correspond to SDs. Figures

(H) and (I); two-sided paired t-test, figures (K); two-sided unpaired t-test. * $p<0.05$, **$p<0.01$, ****

$p<0.000$. Time in mins:secs. Scale bar is 5 μm.

Figure 3: Biased MTOC activity retains Cid in the apical neuroblast hemisphere during

interphase and early mitosis

(A) Representative third instar larval neuroblast expressing the centromere specific Histone-3

variant marker, Cid::EGFP (cyan; top, white; bottom) and the microtubule marker cherry::Jupiter

(white; top). Colored boxes represent corresponding cell cycle stages. **(B)** The distance (purple

and yellow dashed lines) between the apical (purple) and basal (yellow) centrosome, and

individual Cid clusters (green circles) were measured throughout the cell cycle and plotted in **(C)**.

(D) Representative third instar larval neuroblast expressing *cnb* RNAi, cherry::Jupiter (white; top)

and Cid::EGFP (cyan; top, white; bottom row). Red arrowheads highlight the apical MTOC. Blue

arrowheads highlight the maturing basal MTOC. The blue dashed circle highlights Cid clusters.

The cell outline is indicated with the white dashed. 'Apical' centrosome refers to the centrosome

destined to be positioned on the apical cortex, whereas 'basal' centrosome will be inherited by

the basal GMC. **(E)** CS – Cid distance measurements were performed in *cnb* RNAi expressing

Nbs. Once the apical CS disappeared in interphase, the last detectable position was used as a

reference point (orange cross; open circles). **(F)** CS – Cid measurements for *cnb* RNAi expressing

Nbs. Closed arrows refer to actual CS – Cid measurements. Open circles denote Cid – previous

active CS measurements. **(G)** Wild type neuroblasts maintain apical CS – Cid attachments in

prophase, due to asymmetric MTOC activity and microtubule-dependent interphase centrosome

– Cid attachments. *cnb* RNAi expressing neuroblasts lose MTOC activity in interphase,

randomizing the position of Cid clusters. When centrosomes mature again in prophase, both

centrosomes simultaneously attach to Cid clusters. **(H)** Centrosome – Cid distance of an

unperturbed wild type neuroblast at the time of basal centrosome maturation (0 mins) and 6 mins

thereafter. **(I, J)** Cid – centrosome distance measurement in *cnb* RNAi expressing neuroblasts.

Error bars correspond to SDs. Figure (H, I, J); two-sided paired t-test. ns; no signficiance. *

p<0.05. Time in mins:secs. Scale bar is 5 μm.

Figure 3 - figure supplement 1: Microtubule-dependent, asymmetric chromatin-

centrosome attachments retain chromosomes close to the apical neuroblast cortex

during interphase.

(A) CID intensity measurements of endogenous and ectopic CID in Nb-GMC hybrid cells (p-value,

0.1294). **(B)** Representative wild type neuroblast expressing the canonical histone marker

His2A::GFP (white) and Cid::EGFP (cyan; top, white; below). The white dashed line highlights the

cell outline. **(C)** Representative example of a wild type neuroblast expressing Cid::EGFP (cyan)

and the microtubule marker cherry::Jupiter treated with colcemid. Blue and yellow arrowheads

highlight the position of Cid clusters and the disappearing apical MTOC, respectively. **(D)** MTOC

intensity and Cid location measurements for the cell shown in (C). Time in mins:secs. Scale bar

is 5 μm.

Figure 4: Asymmetric microtubule dependent centrosome-chromatin attachments contribute to the separation of endogenous and ectopic chromosomes.

Representative third instar larval **(A)** wild type or **(D)** *cnb* RNAi expressing Nb-GMC hybrid cell, expressing Cid::EGFP (top row; cyan, bottom row; white) and the microtubule marker cherry::Jupiter (white; top row). Neuroblast-derived and GMC-derived Cid clusters are outlined with a blue and orange dashed line, respectively. The white dashed line labels the cell outline. Indistinguishable GMC and Nb Cid clusters are highlighted with yellow dashed circles. The distance of **(B)** endogenous or **(C)** ectopic Cid from the wild type hybrid cell shown in (A) in relation to GMC and Nb centrosomes (CS) plotted over time. **(E)** Endogenous or **(F)** ectopic Cid – centrosome distance measurements of the *cnb* RNAi expressing hybrid cell shown in (D). The vertical solid line indicates the time point when endogenous and ectopic Cid can no longer be distinguished. Vertical dashed line indicates nuclear envelope breakdown. For (B), (C), (E), (F): S refers to the time difference between NEB and when ectopic and endogenous CID becomes indistinguishable. Measurements of endogenous Cid – apical (purple) or basal (yellow) endogenous centrosomes were plotted when **(G)** the endogenous basal centrosome matured (0 mins) and 6 minutes thereafter (6mins) or **(H)** the ectopic centrosomes matured ('0' mins) and 6

minutes thereafter. Corresponding measurements of *cnb* RNAi expressing hybrid cells are shown

in **(J)** and **(K)**. Distance of ectopic Cid in relation to endogenous and ectopic centrosomes shown

for **(I)** wild type or **(L)** *cnb* RNAi expressing hybrid cells. **(M)** Time interval between NEB and

mixing of endogenous and ectopic Cid for wild type (blue circles) and *cnb* RNAi expressing (green

circles) hybrid cells. **(N)** Bar graph showing percentage of hybrid cells in which the endogenous

and ectopic CID mixed prior to NEB. **(O)** A third instar wild type hybrid cell expressing the nuclear

envelope (NE) marker, koi::GFP (cyan; top row, white; bottom row) and cherry::Jupiter. **(P)**

Schematic summary: separation of endogenous and ectopic chromosomes involves microtubule-

dependent asymmetric chromatin-centrosome attachments and physical separation through

nuclear envelopes. *cnb* RNAi expressing neuroblasts release endogenous Cid during interphase,

partially randomizing re-attachment in the subsequent mitosis, which causes premature mixing of

endogenous and ectopic Cid. Colored boxes refer to the indicated cell cycle stages. Error bars

correspond to SDs. Figure (G); two-sided paired t-test and Figures (H-L); two-sided unpaired t-

test. ns; no signficiance. * p<0.05, **p<0.01, ***p<0.0001, **** p<0.0001. Time in mins:secs. Scale

bar is 5 μm.

Figure 4 – figure supplement 1: Asymmetric MTOCs and nuclear envelopes separate endogenous and ectopic chromatin in Nb – GMC hybrid cells.

(A) Averaged distance of basal centrosomes to *endogenous* Cid in wild type or **(C)** *cnb* RNAi expressing neuroblasts. **(B)** Averaged distance of basal centrosomes to *ectopic* Cid in wild type or **(D)** *cnb* RNAi expressing neuroblasts. Measurements are plotted for '0' mins (ectopic centrosome maturation in wild type and *cnb* RNAi expressing hybrids) and 6 mins thereafter ('6'). Representative image sequence of a wild type **(E)** unfused neuroblasts or **(F)** hybrid cell, expressing the nuclear envelope (NE) marker koi::GFP (cyan; top, white; bottom) and the microtubule marker cherry::Jupiter (white; top). Error bars correspond to SDs. Figure (A-D) two-sided unpaired t-test. ns; no significance. ** p<0.01, ***p<0.001, ****p<0.0001. Time in mins:secs. Scale bar is 5 µm.

Figure 5: Ectopic chromosomes in hybrid cells segregate independently but erroneously.

(A) The time between chromosome alignment (metaphase) and separation (anaphase onset) was

measured for endogenous and ectopic chromosomes in wild type hybrid cells and plotted in (C).

(B) Representative examples of delayed (top row) and simultaneous (bottom row) segregation of

endogenous (blue dashed circle) and ectopic (orange dashed circle) chromosomes in wild type

hybrid cells expressing the chromatin marker His2A::GFP. **(C)** Quantification of metaphase to

anaphase onset in control Nbs, compared to interphase and prophase Nb-GMC hybrids. **(D)**

Average time difference between Nb and GMC chromatin anaphase onset in Nb-GMC hybrid

cells. **(E)** Representative third instar larval Nb-GMC hybrids expressing histone marker

His2A::GFP showing missegregating chromatids (yellow arrowheads) during anaphase, resulting

in micronuclei (top row), heterokaryon (middle row) or synkaryon (bottom row) formation. Time

stamps are in relation to NEB (=0:00). **(F)** Bar graph showing percentage of Nb-GMC hybrid cells

with missegregating chromatids. **(G)** Bar graph quantifying the percentage of fused cells with

micronuclei, heterokaryons or synkaryons. ΔT; time difference. Error bars correspond to SDs.

One way ANOVA test was used in (C). ns; no significance. **** p<0.0001. Time in mins:secs.

Scale bar is 5 μm.

Figure 6: Nb – GMC derived hybrid cells neither form visible tumors nor reduce the lifespan in adult hosts.

(A) Third instar larval brains expressing His2A::GFP (top row; negative control), co-expressing *brat* RNAi (middle row; positive control) or containing Nb – GMC hybrid cells (bottom row) were transplanted into wild type hosts and imaged at the day of transplantation (day 0) and Day 30. Yellow arrowheads highlight tumors in wild type hosts transplanted with brat RNAi expressing brains. **(B)** Bar graph showing the percentage of host flies containing visible tumors for the indicated grafted samples (colored boxes on top of the graph). **(C)** Scatter plot showing the lifespan of host flies after brain grafts. Grafted tissue either contained no fusions (wild type transplants), expressing *brat* RNAi without fusions (brat RNAi transplants), attempted but unsuccessful fusion (ablation without fusion transplants) and with brains containing hybrid cells (fusion transplants). Host flies without transplanted tissue were plotted for comparison (no transplantation). **(D)** Model: Asymmetric MTOCs capture neuroblast chromatin during interphase, thereby establishing a physical separation between endogenous and ectopic chromatin. Nuclear envelopes form an additional physical barrier. As hybrid cells enter mitosis, the neuroblast centrosomes and introduced ectopic GMC centrosomes nucleate two separate bipolar spindles, aligning their respective chromatin at the metaphase plate. This separation persists through anaphase. Ectopic chromosomes fail to segregate accurately. Error bars correspond to SDs. One way ANOVA. Ns; no signficiance, ****p<0.0001. Scale bar is 1mm.

References

Aichel O. 1911. Über Zellverschmelzung mit quantitativ abnormer Chromosomenverteilung als Ursache der Geschwulstbildung. Verlag von Wilhelm Engelmann.

Akera T, Chmátal L, Trimm E, Yang K, Aonbangkhen C, Chenoweth DM, Janke C, Schultz RM, Lampson MA. 2017. Spindle asymmetry drives non-Mendelian chromosome segregation. *Science* **358**:668 672. doi:10.1126/science.aan0092

Akera T, Trimm E, Lampson MA. 2019. Molecular Strategies of Meiotic Cheating by Selfish Centromeres. *Cell* **178**:1132-1144.e10. doi:10.1016/j.cell.2019.07.001

Arco AG del, Edgar BA, Erhardt S. 2018. In Vivo Analysis of Centromeric Proteins Reveals a Stem Cell-Specific Asymmetry and an Essential Role in Differentiated, Non-proliferating Cells. *Cell Reports* **22**:1982–1993. doi:10.1016/j.celrep.2018.01.079

Ben-David U, Amon A. 2020. Context is everything: aneuploidy in cancer. *Nat Rev Genet* **21**:44–62. doi:10.1038/s41576-019-0171-x

Blachon S, Gopalakrishnan J, Omori Y, Polyanovsky A, Church A, Nicastro D, Malicki J, Avidor-Reiss T. 2008. Drosophila asterless and vertebrate Cep152 Are orthologs essential for centriole duplication. *Genetics* **180**:2081 2094. doi:10.1534/genetics.108.095141

Buszczak M, Paterno S, Lighthouse D, Bachman J, Planck J, Owen S, Skora AD, Nystul TG, Ohlstein B, Allen A, Wilhelm JE, Murphy TD, Levis RW, Matunis E, Srivali N, Hoskins RA, Spradling AC. 2007. The carnegie protein trap library: a versatile tool for Drosophila developmental studies. *Genetics* **175**:1505 1531. doi:10.1534/genetics.106.065961

Cabernard C, Doe CQ. 2009. Apical/basal spindle orientation is required for neuroblast homeostasis and neuronal differentiation in Drosophila. *Dev Cell* **17**:134 141. doi:10.1016/j.devcel.2009.06.009

Cabernard C, Prehoda KE, Doe CQ. 2010. A spindle-independent cleavage furrow positioning pathway. *Nature* **467**:91 94. doi:10.1038/nature09334

Caussinus E, Gonzalez C. 2005. Induction of tumor growth by altered stem-cell asymmetric division in Drosophila melanogaster. *Nat Genet* **37**:1125 1129. doi:10.1038/ng1632

Dietzl G, Chen D, Schnorrer F, Su K-C, Barinova Y, Fellner M, Gasser B, Kinsey K, Oppel S, Scheiblauer S, Couto A, Marra V, Keleman K, Dickson BJ. 2007. A genome-wide transgenic RNAi library for conditional gene inactivation in Drosophila. *Nature* **448**:151 156. doi:10.1038/nature05954

Dorn JF, Jaqaman K, Rines DR, Jelson GS, Sorger PK, Danuser G. 2005. Yeast Kinetochore Microtubule Dynamics Analyzed by High-Resolution Three-Dimensional Microscopy. *Biophys J* **89**:2835–2854. doi:10.1529/biophysj.104.058461

Drpic D, Almeida AC, Renda F, Damas J, Lewin HA, Larkin DM, Khodjakov A, Maiato H. 2018. Chromosome Segregation Is Biased by Kinetochore Size. *Curr Biol* **28**:1344 1356.e5. doi:10.1016/j.cub.2018.03.023

Fujiwara T, Bandi M, Nitta M, Ivanova EV, Bronson RT, Pellman D. 2005. Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-null cells. *Nature* **437**:1043–1047. doi:10.1038/nature04217

Fukagawa T, Earnshaw WC. 2014. The Centromere: Chromatin Foundation for the Kinetochore Machinery. *Dev Cell* **30**:496–508. doi:10.1016/j.devcel.2014.08.016

Gallaud E, Nair AR, Horsley N, Monnard A, Singh P, Pham TT, Garcia DS, Ferrand A, Cabernard C. 2020. Dynamic centriolar localization of Polo and Centrobin in early mitosis primes centrosome asymmetry. *Plos Biol* **18**:e3000762. doi:10.1371/journal.pbio.3000762

Gallaud E, Pham T, Cabernard C. 2017. Drosophila melanogaster Neuroblasts: A Model for Asymmetric Stem Cell Divisions. *Results Problems Cell Differ* **61**:183 210. doi:10.1007/978-3-319-53150-2_8

Gambarotto D, Pennetier C, Ryniawec JM, Buster DW, Gogendeau D, Goupil A, Nano M, Simon A, Blanc D, Racine V, Kimata Y, Rogers GC, Basto R. 2019. Plk4 Regulates Centriole Asymmetry and Spindle Orientation in Neural Stem Cells. *Dev Cell* **50**:11-24.e10. doi:10.1016/j.devcel.2019.04.036

Gratz SJ, Cummings AM, Nguyen JN, Hamm DC, Donohue LK, Harrison MM, Wildonger J, O'Connor-
Giles KM. 2013. Genome Engineering of Drosophila with the CRISPR RNA-Guided Cas9 Nuclease.
Genetics **194**:1029–1035. doi:10.1534/genetics.113.152710

Heasley LR, Markus SM, DeLuca JG. 2017. "Wait Anaphase" Signals Are Not Confined to the Mitotic
Spindle. *Mol Biol Cell* **28**:1186–1194. doi:10.1091/mbc.e17-01-0036

Henikoff S, Ahmad K, Platero JS, Steensel B van. 2000. Heterochromatic deposition of centromeric
histone H3-like proteins. *Proc National Acad Sci* **97**:716–721. doi:10.1073/pnas.97.2.716

Januschke J, Gonzalez C. 2010. The interphase microtubule aster is a determinant of asymmetric division
orientation in Drosophila neuroblasts. *J Cell Biology* **188**:693 706. doi:10.1083/jcb.200905024

Januschke J, Llamazares S, Reina J, Gonzalez C. 2011. Drosophila neuroblasts retain the daughter
centrosome. *Nat Commun* **2**:243. doi:10.1038/ncomms1245

Januschke J, Reina J, Llamazares S, Bertran T, Rossi F, Roig J, Gonzalez C. 2013. Centrobin controls
mother–daughter centriole asymmetry in Drosophila neuroblasts. *Nat Cell Biol* **15**:241 248.
doi:10.1038/ncb2671

Katsani KR, Karess RE, Dostatni N, Doye V. 2008. In Vivo Dynamics of Drosophila Nuclear Envelope
Components. *Mol Biol Cell* **19**:3652–3666. doi:10.1091/mbc.e07-11-1162

Kawamura N. 2001. Fertilization and the first cleavage mitosis in insects. *Dev Growth Differ* **43**:343–349.
doi:10.1046/j.1440-169x.2001.00584.x

Kracklauer MP, Banks SML, Xie X, Wu Y, Fischer JA. 2007. Drosophila klaroid Encodes a SUN Domain
Protein Required for Klarsicht Localization to the Nuclear Envelope and Nuclear Migration in the Eye.
Fly **1**:75–85. doi:10.4161/fly.4254

Lee YL, Burke B. 2018. LINC complexes and nuclear positioning. *Semin Cell Dev Biol* **82**:67 76.
doi:10.1016/j.semcdb.2017.11.008

Loyer N, Januschke J. 2020. Where does asymmetry come from? Illustrating principles of polarity and
asymmetry establishment in Drosophila neuroblasts. *Curr Opin Cell Biol* **62**:70–77.
doi:10.1016/j.ceb.2019.07.018

Maiato H, Gomes AM, Sousa F, Barisic M. 2017. Mechanisms of Chromosome Congression during
Mitosis. *Biology* **6**:13. doi:10.3390/biology6010013

McIntosh JR. 2016. Mitosis. *Csh Perspect Biol* **8**:a023218. doi:10.1101/cshperspect.a023218

Molina O, Abad MA, Solé F, Menéndez P. 2020. Aneuploidy in Cancer: Lessons from Acute
Lymphoblastic Leukemia. *Trends Cancer*. doi:10.1016/j.trecan.2020.08.008

Muller H, Gil J, Drinnenberg IA. 2019. The Impact of Centromeres on Spatial Genome Architecture.
Trends Genet **35**:565–578. doi:10.1016/j.tig.2019.05.003

Musacchio A. 2015. The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics. *Curr*
Biol **25**:R1002 18. doi:10.1016/j.cub.2015.08.051

Ogle BM, Cascalho M, Platt JL. 2005. Biological implications of cell fusion. *Nat Rev Mol Cell Bio* **6**:567
575. doi:10.1038/nrm1678

Platt JL, Cascalho M. 2019. Cell Fusion in Malignancy: A Cause or Consequence? A Provocateur or
Cure? *Cells* **8**:587. doi:10.3390/cells8060587

Prosser SL, Pelletier L. 2017. Mitotic spindle assembly in animal cells: a fine balancing act. *Nat Rev Mol*
Cell Bio **18**:187–201. doi:10.1038/nrm.2016.162

Quintyne NJ, Reing JE, Hoffelder DR, Gollin SM, Saunders WS. 2005. Spindle Multipolarity Is Prevented
by Centrosomal Clustering. *Science* **307**:127–129. doi:10.1126/science.1104905

Ranjan R, Snedeker J, Chen X. 2019. Asymmetric Centromeres Differentially Coordinate with Mitotic
Machinery to Ensure Biased Sister Chromatid Segregation in Germline Stem Cells. *Cell Stem Cell*.
doi:10.1016/j.stem.2019.08.014

Razafsky D, Hodzic D. 2009. Bringing KASH under the SUN: the many faces of nucleo-cytoskeletal
connections. *J Cell Biology* **186**:461 472. doi:10.1083/jcb.200906068

Rebollo E, Sampaio P, Januschke J, Llamazares S, Varmark H, Gonzalez C. 2007. Functionally unequal
centrosomes drive spindle orientation in asymmetrically dividing Drosophila neural stem cells. *Dev*
Cell **12**:467 474. doi:10.1016/j.devcel.2007.01.021

Reichmann J, Nijmeijer B, Hossain MJ, Eguren M, Schneider I, Politi AZ, Roberti MJ, Hufnagel L, Hiiragi
T, Ellenberg J. 2018. Dual-spindle formation in zygotes keeps parental genomes apart in early
mammalian embryos. *Science* **361**:189 193. doi:10.1126/science.aar7462

Rieder CL, Khodjakov A, Paliulis LV, Fortier TM, Cole RW, Sluder G. 1997. Mitosis in vertebrate somatic
cells with two spindles: Implications for the metaphase/anaphase transition checkpoint and cleavage.
Proc National Acad Sci **94**:5107–5112. doi:10.1073/pnas.94.10.5107

Rossi F, Gonzalez C. 2015. Studying tumor growth in Drosophila using the tissue allograft method. *Nat*
Protoc **10**:1525–1534. doi:10.1038/nprot.2015.096

Roubinet C, Tsankova A, Pham TT, Monnard A, Caussinus E, Affolter M, Cabernard C. 2017. Spatio-
temporally separated cortical flows and spindle geometry establish physical asymmetry in fly neural
stem cells. *Nat Commun* **8**:1383. doi:10.1038/s41467-017-01391-w

Rusan NM, Peifer M. 2007. A role for a novel centrosome cycle in asymmetric cell division. *J Cell Biology*
177:13 20. doi:10.1083/jcb.200612140

Snook RR, Hosken DJ, Karr TL. 2011. The biology and evolution of polyspermy: insights from cellular and
functional studies of sperm and centrosomal behavior in the fertilized egg. *Reproduction* **142**:779–792.
doi:10.1530/rep-11-0255

Sunchu B, Cabernard C. 2020. Principles and mechanisms of asymmetric cell division. *Development*
147:dev167650. doi:10.1242/dev.167650

Thomas GE, Renjith MR, Manna TK. 2017. Kinetochore–microtubule interactions in chromosome
segregation: lessons from yeast and mammalian cells. *Biochem J* **474**:3559–3577.
doi:10.1042/bcj20170518

Weierich C, Brero A, Stein S, Hase J von, Cremer C, Cremer T, Solovei I. 2003. Three-dimensional
arrangements of centromeres and telomeres in nuclei of human and murine lymphocytes.
Chromosome Res **11**:485–502. doi:10.1023/a:1025016828544

Yadlapalli S, Yamashita YM. 2013. Chromosome-specific nonrandom sister chromatid segregation during
stem-cell division. *Nature* **498**:251 254. doi:doi:10.1038/nature12106

Yu K, Zhong N, Xiao Y, She Z. 2019. Mechanisms of kinesin-7 CENP-E in kinetochore–microtubule
capture and chromosome alignment during cell division. *Biol Cell* **111**:143–160.
doi:10.1111/boc.201800082

Methods

Fly Strains:

Transgenes and fluorescent markers: *worGal4, UAS-mCherry::Jupiter* (Cabernard and Doe,

2009)*; worGal4, UAS-mCherry::Jupiter, Sqh::GFP* (Cabernard et al., 2010)*; His2A::GFP*

(Bloomington stock center); *UAS-mCherry::CAAX, UAS-iLID::CAAX:;mCherry (A. Monnard & C.*

Cabernard; unpublised); Cid::EGFP (Ranjan et al., 2019); *pUbq-Asl::GFP* (Blachon et al., 2008);

worgal4, UAS-mCherry::Jupiter, Asl::GFP (this work); *pros::EGFP* (endogenously tagged with

CRISPR; this work); *koi::GFP (CB04483)* (Buszczak et al., 2007); *cnb^{GD11735}* RNAi line (v28651)

(Dietzl et al., 2007).

Generation of pros::EGFP with CRISPR:

Target specific sequences with high efficiency were chosen using the CRISPR Optimal Target

Finder (http://tools.flycrispr.molbio.wisc.edu/targetFinder/), the DRSC CRISPR finder

(http://www.flyrnai.org/crispr/), and the Efficiency Predictor (http://

www.flyrnai.org/evaluateCrispr/) web tools. Sense and antisense primers for these chosen sites

were then cloned into pU6-BbsI-ChiRNA (Gratz et al., 2013) between BbsI sites. To generate the

replacement donor template, EGFP and 1 kb homology arms flanking the insertion site were

cloned into pHD-DsRed-attP (Addgene plasmid #51019) using Infusion technology

(Takara/Clontech). Injections were performed in house. Successful events were detected by

DsRed-positive screening in the F1 generation. Constitutively active Cre (BDSC#851) was then

crossed in to remove the DsRed marker. Positive events were then balanced, genotyped, and

sequenced.

Live cell imaging acute cell-cell fusion:

Imaging medium consists of Schneider's insect medium (Sigma-Aldrich S0146) mixed with 10%

BGS (HyClone). Third instar larvae were dissected in imaging medium and the brains were

transferred into a μ-slide Angiogenesis or μ-slide 8 well (Ibidi). Live samples were imaged with an

Intelligent Imaging Innovations (3i) spinning disc confocal system, consisting of a Yokogawa CSU-

W1 spinning disc unit and two Prime 95B Scientific CMOS cameras. A 60x/1.4NA oil immersion

objective mounted on a Nikon Eclipse Ti microscope was used for imaging. Live imaging voxels

are 0.22 X 0.22 X 0.75-1 m (60x/1.4NA spinning disc).

Neuroblast-GMC fusions were induced using a 3i Ablate! ablation system, consisting of a 532nm

pulsed laser. We used a pulse width of 7 ns, targeting the membrane interface between the

neuroblast and the adjacent GMC.

Colcemid treatment:

Dissected brains were incubated with Colcemid (Sigma) in live imaging medium at a final

concentration of of 25 μgmL^{-1}.

Transplantation experiments:

Brain lobes containing the hybrid cells expressing *His2A::GFP* and *worGal4, UAS-*

mCherry::Jupiter were transplanted into 3 to 4 day old, well fed adult *w^{1118}* female host flies as

described previously (Rossi and Gonzalez, 2015). Custom made needles were prepared from

Narishige GD-1 glass capillaries using a Narishige, needle puller. Injection needles were shaped

with forceps to have a smooth, 45º opening. Transplanted flies were transferred into fresh vials

each day for the first three days, followed by biweekly flipping. The tumor growth was monitored

and recorded with a Leica MZ FLIII fluorescence stereomicroscope.

Image processing and measurements:

Live cell images were processed using imaris x64 8.3.1 and image J.

For angle and distance measurements, the coordinates for the two spindle poles were determined

in Imaris. From these coordinates, angles and distances between spindles were derived based

on the calculations outlined below.

Angle between spindles: $\theta = \cos^{-1} \dfrac{n \cdot e}{|n||e|}$

Dot product: $n \cdot e = (X1*X2) + (Y1*Y2) + (Z1*Z2)$

Magnitude of vectors: $|n| = \sqrt{X_1^2 + Y_1^2 + z_1^2}$ $|e| = \sqrt{X_2^2 + Y_2^2 + z_2^2}$

Where **n** corresponds to the spindle vector: **n** (x1,y1,z1) = (N1-N1', N2-N2', N3-N3') and **e** to the

ectopic spindle vector: **e** (x2,y2,z2) = (E1-E1', E2-E2', E3-E3')

N1, N2, N3 and N1', N2' and N3' are coordinates of the two poles of the Nb spindle. Similarly,

E1, E2, and E3 and E1', E2' and E3' are coordinates of the ectopic spindle poles.

Distance between spindle vectors:

The midpoints of the two spindle vectors are calculated from coordinates of the poles on either

side of the respective spindle. This is followed by calculating the distance between these

midpoints.

Midpoint of the Nb spindle vector $\quad = \left(\frac{N_1+N_1^1}{2}, \frac{N_2+N_2^1}{2}, \frac{N_3+N_3^1}{2}\right) = (M_1, M_2, M_3)$

Midpoint of the GMC spindle vector $\quad = \left(\frac{G_1+G_1^1}{2}, \frac{G_2+G_2^1}{2}, \frac{G_3+G_3^1}{2}\right) = (m_1, m_2, m_3)$

Distance between these two points $\quad = \sqrt{(M_1-m_1)^2 + (M_2 - m_2)^2 + (M_3 - m_3)^2}$

Centrosome - Cid distance:

The centrosome (CS) - Cid distance was calculated using Cid and CS coordinates.

CS – Cid distance: $= \sqrt{(x_1-x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$

Where x_1,y_1,z_1 correspond to CS and x_2,y_2,z_2 to Cid coordinates, respectively.

Plotted values correspond to averaged values of all CS – Cid punctae distances and the

corresponding standard deviations.

0 and 6 mins corresponds to the appearance of the basal centrosome and 6 minutes thereafter.

'0' and '6' mins corresponds to the appearance of the ectopic centrosome and 6 minutes

thereafter.

Statistical analysis:

Statistical analysis was performed using Graphpad prism 8. Statistical significance was

determined using paired or unpaired t-test and one-way ANOVA. Significance was indicated as

following: *; $p<0.05$, **;$p<0.01$, ***;$p<0.001$,****$p<0.0001$, ns; not significant. Exact p values and

complete statistical information can be found in Extended data table 1.

Measurements were taken from distinct samples and from several independent experiments.

Acknowledgements

We thank Xin Chen for fly stocks, David Salvador Garcia for generating the Pros::EGFP transgenic line, Sue Biggins and members of the Cabernard laboratory for helpful discussions and comments. This work was supported by the National Institutes of Health (1R01GM126029-03) and a Research Scholar grant from the American Cancer Society (130285-RSG-16253-01-CSM). Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) and from the Vienna Drosophila Resource Center (VDRC).

Author contributions

This study was conceived by B.S., N.L., and C.C. C.R provided some conceptual ideas early on.

Technical feasibility and was demonstrated by C.R & C.C.

B.S and N.L performed all the experiments with significant help from C.S.

B.S, N.L., C.S., and C.C analyzed the data.

B.S and C.C. wrote the manuscript.

Competing interest declaration

The authors declare no competing financial interests.

Data availability statement

The authors declare that all data supporting the findings of this study are available within the paper and its supplementary files. Source data are available upon request.

Supplementary Table 1: Statistical information

Complete statistical information for the data shown in the corresponding figures.

Video legends

Video 1: Wild type neuroblast division; related to Figure 1b

Wild type control (unfused) neuroblast expressing the microtubule binding protein Cherry::Jupiter

(white) and the canonical Histone marker His2A::GFP (cyan). Time scale is h:mm:ss and the

scale bar is 5 µm.

Video 2: Wild type hybrid cell; related to Figure 1b

Wild type hybrid cell derived from a neuroblast – GMC fusion *in vivo*, expressing the canonical

Histone marker His2A::GFP (white in single channel; cyan in merge) and the microtubule binding

protein Cherry::Jupiter (white). The blue and orange arrows mark endogenous and ectopic

chromatin, respectively. Time scale is h:mm:ss and the scale bar is 3 µm.

Video 3: Wild type hybrid cell; related to Figure 2b

Wild type hybrid cell derived from a neuroblast – GMC fusion *in vivo*, expressing the canonical

Histone marker His2A::GFP (white in single channel; cyan in merge) and the microtubule binding

protein Cherry::Jupiter (white). The blue and orange arrows mark endogenous and ectopic

chromatin, respectively. Time scale is h:mm:ss and the scale bar is 2 µm.

Video 4: Wild type neuroblast division; related to Extended Data Figure 2b

Wild type control (unfused) neuroblast, expressing the microtubule binding protein Cherry::Jupiter

(white) and the centromere-specific H3 variant Cid::EGFP (Cyan). Purple and yellow arrows point

to the apical and basal centrosome, respectively. The blue arrow refers to moving Cid clusters.

Time scale is h:mm:ss and the scale bar is 1 µm.

Video 5: Wild type neuroblast division; related to Extended Data Figure 2f

Wild type control (unfused) neuroblast, expressing the membrane marker mCherry::CAAX

(white), the canonical Histone marker His2A::GFP (white) and Cid::EGFP (Cyan). The green

arrow points to Cid clusters. Time scale is h:mm:ss and the scale bar is 5 µm.

Video 6: Wild type neuroblast exposed to Colcemid; related to Extended Data Figure 2g

Wild type control (unfused) neuroblast exposed to the microtubule depolymerizing drug Colcemid, expressing the membrane marker mCherry::CAAX (white), the microtubule binding protein Cherry::Jupiter (white) and Cid::EGFP (Cyan). The yellow arrow points to the apical centrosome, the blue arrow to Cid clusters. Time scale is h:mm:ss and the scale bar is 1 μm.

Video 7: Cnb RNAi expressing neuroblast; related to Extended Data Figure 2i

cnb RNAi expressing (unfused) neuroblast, co-expressing the microtubule binding protein Cherry::Jupiter (white) and Cid::EGFP (Cyan). The orange and blue arrow points to the apical and basal centrosome, respectively. The green arrow highlights Cid clusters. Time scale is h:mm:ss and the scale bar is 1 μm.

Video 8: wild type hybrid cell; related to Extended Data Figure 3a

Wild type hybrid cells expressing the microtubule binding protein Cherry::Jupiter (white) and Cid::EGFP (white in single channel; Cyan in merge). The blue and orange arrow highlights endogenous and ectopic Cid clusters, respectively. Time scale is h:mm:ss and the scale bar is 1 μm.

Video 9: Cnb RNAi expressing hybrid cell; related to Extended Data Figure 3d

cnb RNAi expressing hybrid cell, expressing the microtubule binding protein Cherry::Jupiter (white) and Cid::EGFP (white in single channel; Cyan in merge). The blue and orange arrow highlights endogenous and ectopic Cid clusters, respectively. The yellow arrow indicates mixing of endogenous and ectopic Cid clusters. Time scale is h:mm:ss and the scale bar is 1 μm.

Video 10: wild type hybrid cell; related to Figure 4e

Wild type hybrid cell, expressing the canonical Histone marker His2A::GFP (white). The blue and orange arrow highlights endogenous and ectopic chromosomes, respectively. The magenta arrowhead highlights the fate of missegregated chromosomes. This hybrid cell forms a heterokaryon. Time scale is h:mm:ss and the scale bar is 1 μm.