Sex-specific body mass ageing trajectories in adult Asian elephants

Lucas D. Lalande[1,2,3,a], Virpi Lummaa[1,a], Htoo H. Aung[4], Win Htut[4], U. Kyaw Nyein[4],
Vérane Berger[1,a,*], Michael Briga[1,a,*]

[1] Department of Biology, University of Turku, Turku, Finland;

[2] Université Bourgogne Franche-Comté, Dijon, France;

[3] Present address: Université de Lyon, Université Lyon 1, UMR CNRS 5558,
Villeurbanne CEDEX, France;

[4] Myanma Timber Enterprise, Ministry of Natural Resources and Environmental
Conservation, West Gyogone Forest Compound, Yangon, Myanmar;

[*] Equal contribution of these authors

[a] Corresponding authors: L. D. Lalande: lucas.lalande@univ-lyon1.fr, V. Lummaa:
virpi.lummaa@gmail.com, V. Berger: verane.berger@orange.fr, M. Briga:
michbriga@gmail.com

Abstract

In species with marked sexual dimorphism and where one sex undergoes stronger
intrasexual competition, that sex is expected to age earlier or quicker. Here, we utilise
a unique, longitudinal dataset of a semi-captive population of Asian elephants
(*Elephas maximus*), a species with marked male-biased intrasexual competition, with
males being larger and living shorter, and test the hypothesis that males show earlier
and/or faster body mass ageing than females. We show sex-specific body mass
ageing trajectories: adult males gained weight up to the age of 48 years old, followed
by a decrease in body mass until natural death. In contrast, adult females gained
body mass with age until a body mass decline in the last year of life. Our study shows
that sex-specific life-histories shape ageing patterns consistent with the predictions of
the classical theory of ageing.

Keywords: senescence, sex-differences, terminal decline, Elephas maximus,
long-term data

Introduction

Ageing – a decline in organismal functioning with age (Monaghan et al., 2008) – has been observed in many species (Jones et al., 2014). However, the onset and rates of ageing differ both between (Jones et al., 2014) and within species (Nussey et al., 2007) and between sexes (Douhard et al., 2017). A main challenge in ageing research is to quantify and explain such differences in the onset and rates of ageing (Rando & Wyss-Coray, 2021).

In species with sex-specific intrasexual competition, classic theory predicts that the sex with the highest intrasexual competition has a shorter lifespan and an earlier onset and/or higher rate of ageing (Maklakov & Lummaa, 2013). The rationale is that high intrasexual selection often results in one sex showing conspicuous displays or aggressive intrasexual behaviours, leading to increased mortality and a *live fast, die young* pace of life (Bonduriansky et al., 2008; Clutton-Brock & Isvaran, 2007; Maklakov & Lummaa, 2013). Accordingly, in polygynous species with male-biased intrasexual competition, males often die earlier (Lemaître et al., 2020) and age earlier or faster than females (Beirne et al., 2015; Clutton-Brock & Isvaran, 2007; Douhard et al., 2017; Nussey et al., 2009; Tidière et al., 2015). However, recent conceptual developments have shown that this association can be disrupted. This can occur for example because of condition-dependent extrinsic mortality selecting particularly high-performing individuals in the population (Chen & Maklakov, 2014) or canalisation (*i.e.* the more a trait contributes to fitness, the less it should deviate from optimal trait value, with respect to environmental variation (Flatt, 2005)), thereby contradicting the theoretically expected earlier or faster ageing in males. The extent to which such phenomena occur in nature remains unknown.

Here, we used a unique long-term dataset to describe sex-specific body mass ageing

trajectories in a nutritionally unsupplemented semi-captive timber population of Asian

elephants (*Elephas maximus*) living in their natural environment in Myanmar. Body

mass is of interest in the study of ageing because it is positively associated with key

life-history traits such as reproduction and lifespan in many non-human species

(Hämäläinen et al., 2014). Accordingly, in Asian elephants, seasonal variation in body

mass was positively associated with survival the following month (*e.g.* low body mass

was associated with low survival during dry season) (Mumby, Mar, et al., 2015).

However, we know almost nothing about how body mass changes with age in

elephants despite the interest in studying ageing in such a long-lived, social and

sexually dimorphic non-human species. While females live in kin groups, adult males

often roam solitary, undergo a more intense intrasexual competition for dominance

and mating (Sukumar, 2003) and hence are bigger, heavier (Mumby, Chapman, et

al., 2015), more aggressive and less sociable (Seltmann et al., 2019) and shorter-

lived than females (respective median lifespans: 30.8 and 44.7 years) (Lahdenperä

et al., 2018). Based on this male-biased intrasexual competition and shorter lifespan,

and following the classic theory of ageing, we expected males to display earlier

and/or faster body mass ageing than females (Bonduriansky et al., 2008; Maklakov &

Lummaa, 2013).

Material and methods

Study population

We studied the world's largest semi-captive Asian elephant population

consisting of around 3,000 individually-marked elephants owned by the government-

run Myanma Timber Enterprise (MTE) (Leimgruber et al., 2008). Their birth, death, pedigree details, and morphological measurements have been recorded for almost a century by local veterinarians. These elephants are distributed across Myanmar in forest camps and used as riding, transport and drafting animals. Elephants work during the day and, at night, they socialise, mate and forage freely, unsupervised in forests (Oo, 2010; Zaw, 1997). There are no husbandry procedures and timber elephants are never culled. Calves born in captivity are cared for and nursed by their biological mother and allomothers (Lahdenperä et al., 2016; Lynch et al., 2019). Therefore, breeding rates are natural with no reproductive management. Moreover, there is minimal food provisioning, but elephants benefit from veterinary care that consists of treatment of simple injuries and monitoring of working conditions.

Both males and females are used in the workforce, and each working group of six elephants is composed of both sexes. Males and females follow the same government set limitations on taming age, working and retirement age, working-days per year, hours of work per day and tonnage pulled annually apply to both sexes, although it is possible that males might be used for somewhat different working tasks at times (*e.g.* when use of tusks is required; only males can possess long tusks in Asian elephants). Pregnant females are given a rest period from mid-pregnancy (around 11 months into gestation) until the calf is 1-year-old (Toke Gale, 1974), while they and their calf are being monitored by their mahouts (individual caretakers and riders) throughout this period. Following this break, mothers are used for light work but are kept with calves at heel and able to suckle on demand until the calf is four or five years old (Oo, 2010) at which point calves are assigned a rider, name, logbook and registration number. After the training period, elephants are used for light work

duties until the age of 17, when they enter the full workforce until retirement around

age 50. The MTE maintains their care and logbooks until death.

Data collection and selection

Our analyses focused on age- and sex-specific variation in adult body mass

from age 18 onwards, omitting the phase during which elephants grow in height

(Mumby, Chapman, et al., 2015). From the logbooks, we extracted 1,901 body

masses of 347 elephants with known sex, year of birth ('YOB'), alive or dead status,

origin (captive-born or wild-caught) and measurement season (hot: Feb-May,

monsoon: Jun-Sep, cold: Oct-Jan (Mumby, Mar, et al., 2015)) from 325 working

localities ('township'). Study elephants were aged 18 – 72 years (mean = 39.3) and

born 1941 – 1999. Age and cohort information were comparable between sexes, with

171 males (n = 1,316) born 1954 – 1999 and aged 18 – 64 years (mean = 37.4), and

322 females (n = 2,570) born 1941 – 1999 and aged 18 – 72 years (mean = 40.2).

When body mass measurements were not available, we estimated body

masses using height to the shoulder and chest girth (n = 1,985 measurements on

342 individuals, method in Supplementary Information 1). In Asian elephants, chest

girth reflects changes in mass in addition to the fixed effect of structural size during

adulthood (details in SI2). In total, we obtained a median of 4.0

measurements/individual [2.5 – 97.5th percentiles: 1.0 – 36.4], followed for a median

period of 2.8 years [2.5 – 97.5th: 0.0 – 36.6] on 493 elephants (n = 3,886). Two

influential observations measured at age 18 and 23 were removed for one male

because of particularly low Δage (Cook's distance = 0.61 and 0.25, mean of 0.001 on

all males). Other observations for this male, all after age 50, were included.

Statistical analyses

We investigated the age- and sex-specific variation in body mass in R version

4.0.5 (R Core Team, 2020), using the log-transformed body mass as a dependent

variable with a normal error distribution. We first tested whether there were sex-

specific ageing trajectories (table S2) and then compared the sex-specific ageing

trajectories using general additive mixed models (GAMs) and general linear mixed

models (GLMs), with respectively the functions 'gamm' of the package 'mgcv' (Wood,

2017) and the function 'lmer' of the package 'lme4' (Bates et al., 2015). GAMs allow

more flexible ageing trajectories than GLMs, but the more constrained ageing

trajectories in GLMs allow a less descriptive identification of differences in ageing

trajectories (figure S1) and both approaches gave consistent conclusions (see results

section). We identified the best fitting models using the model selection approach

based on the second order Akaike Information Criterion (AICc) as implemented in the

package 'MuMIn' (Bartoń, 2019). In brief, the best fitting model has the lowest AICc

value, with other models within 4 ΔAICc being plausible and models become

increasingly equivocal up to 14 ΔAICc, after which they become implausible

(Burnham et al., 2011). Visual inspection of model residuals confirmed that these

fulfilled all assumptions of distribution and homogeneity without any influential data

points or outliers (see above).

Within- vs. between-individual change

In all models, we accounted for non-independence of data due to repeated

measurements from the same individual by including elephant identity ('ID') as a

random intercept. The composition of the population can change with age for

example due to selective disappearance of certain (*e.g.* lighter or heavier) individuals,

which can affect the age trajectory. In order to alleviate this problem to the best

possible in this long-lived species, we decomposed body mass changes with age into

between- and within-individual changes following the approach developed by van

de Pol & Verhulst, 2006 and van de Pol & Wright, 2009 using two terms: *i)* the age at

last measurement for each individual, which captures the between-individual

variations and *ii)* a 'Δage' term (age at measurement minus the individual's mean age

for all measurements) capturing the within-individual changes with age. We mean-

centred and standardised 'Δage' so that *i)* individuals measured once all get a Δage

= 0 and hence contribute to the variance of the Δage intercept but not to its slope and

ii) to avoid collinearity and to have comparable variance for Δage and Δage² (Bolker,

2008; Zuur et al., 2009). Our dataset included 105 individuals with one measurement

after the age of 18 and we included these individuals by giving them Δage = 0 (*i.e.*

mean-centred) so they do not contribute to the coefficient but do contribute to the

variance along the Y axis on Δage = 0, diminishing the likelihood of a false positive,

and ii) do contribute to the coefficient of the age at last measurement term, thereby

avoiding a bias in the dataset from selecting only the longer-lived or most monitored

individuals.

Testing ageing trajectories

We tested several within-individual ageing trajectories: GAMs can have any shape

and for GLMs we tested linear, quadratic, threshold and terminal models (figure S1)

and we selected the ageing trajectory with the lowest AICc. For GAMs, we identified

the age at which maxima occurred based on the first-order derivative (= 0) using the

function 'fderiv' of the package 'gratia' (Simpson & Singmann, 2021). For threshold

models (figure S1C), we followed the approaches previously developed in Briga et

al., 2019 and Douhard et al., 2017. In brief, we first identified the best-fitting threshold

age in a series of models, varying the threshold in the 'Δage' term between -35 to 22

years with intervals of one Δage (1 mean-centered Δage = 4.4 and 4.5 years for

males and females respectively) and estimated the threshold and its confidence

intervals using ± 4 ΔAIC age range. Then we compared the best-fitting threshold

model with the other ageing trajectories. Sometimes, declines in trait value appear

shortly before death (terminal decline). We coded a 'terminal' change (figure S1D) as

a binomial factor for whether an individual died during the year following the

measurement. We used a one-year-window to avoid a possible seasonal covariation

in weight and because it was the best fitting time-window, but note that models with

other time-windows gave consistent conclusions (figure S3).

Accounting for temporal and spatial variation in body mass

As body mass variation can be influenced by seasonal, spatial and within-individual

factors, we tested whether body mass values were affected by (i) measured or

estimated, (ii) individuals were alive or dead, (iii) captive- or wild-born and (iv) the

measurement season. To this end we used a combination of backward elimination

and forward selection approaches, *i.e.* first we analysed the models including these

covariates using the most complex ageing trajectory, identified that none of these

factors improved the model, then we eliminated these co-variates from the model,

identified the best fitting ageing trajectory, and then we repeated the final models

obtained with these covariates and identified again that none of these improved the

model fit or altered the previously obtained conclusions (table S3). In our models, we

included as random intercepts individual identity to account for the repeated

measurement of the same individual. We also included 'township' to account for the

spatial clustering of individuals across Myanmar (except in male GLMs which lacked

power), although actually adding township worsened the model fit (male GAM: ΔAICc

= +1.5; female GAM: ΔAICc = +0.6).

Results

At the measurements' starting age of 18 years, males were on average 253 kg

At the measurements' starting age of 18 years, males were on average 253 kg

heavier than females, weighing respectively 2,559 kg [95%CI: 2,431 – 2,694] and

2,306 kg [95%CI: 2,258 – 2,355] and this difference was statistically significant

(ΔAICc = -122.6 in a GLM with vs without sex as a fixed effect).

We identified the elephants body mass ageing trajectories using general additive

models (GAMs) and general linear models (GLMs) and both approaches gave

consistent results. Both analyses showed that sexes have different body mass

ageing trajectories (GAM: ΔAICc = -65.7, figure 1, table S2; GLM: ΔAICc = -47.0,

figure 2, table S2) and hence, we identified the ageing trajectories for both sexes

separately.

For males, both GAMs and GLMs indicated a body mass gain from age 18 years until

their late-forties or early fifties (GAM maximum: 54 years, 95%CI: 53 – 56, figure S2;

GLM maximum: 48.3 years, 4 AICc CI: 46.6 – 52.3, figure 2A), followed by a decline

until death (figure 1A, figure 2A, Table 1). In GLMs, this maximum was shown

through the best fit of a threshold model (ΔAICc = -30.4 compared to a linear

trajectory): males gained mass at a rate of 21 kg/year [95%CI: 19.1 – 23.4] or 0.9%

[95%CI: 0.9 – 1.0] of males' mean body mass and then lost mass at a rate of 28

kg/year [95%CI: 14.6 – 41.3] or 1.3% [95%CI: 0.7 – 1.8] of males' mean body mass

(figure 2A). Regarding the decline, neither GAMs nor GLMs showed confident

statistical support that it was terminal (*i.e.* determined by time before death rather

than age): in GAMs, models with and without the terminal term were almost

equivalent (ΔAICc=+0.01, Table 1) and in GLMs adding a terminal term worsened the

model fit (ΔAICc=+5.0, Table 1, figure S3A).

For females, both GAMs and GLMs indicated a body mass gain throughout their lives

until a terminal decline during their last year of life (figure 1B, figure 2B, Table 1).

GLMs indicated a mass gain of 8 kg/year [95%CI: 7.4 – 10.3] or 0.3% [95%CI: 0.3 –

0.4] of females' mean body mass (ΔAICc = -6.7, Table 1). Loss of body mass

occurred in the last year of life (GAM: ΔAICc = -12.2, Table 1, figure 1B; GLM: ΔAICc

= -4.9, Table 1, figure 2B, figure S3B) and consisted of 173 kg [95%CI: 80 – 263] or

6.8% [95%CI: 3.2 – 10.4] of their mean body mass. For comparison, the extent of the

terminal decline in males, if any, is just over half that in females at 97 kg [95%CI: -20

– 207] or 4.3% [95%CI: -0.9 – 9.2] of males' mean body mass (quantified in the

aforementioned best-fitting threshold model).

Figure 1. Body mass ageing trajectories of (A) males (n=1,316 measurements on 171 individuals) and (B) females (n=2,570 measurements on 322 individuals) with solid lines showing predictions of the best-fitting GAM models (table 1) and grey areas 95%CI. For females, measurements in the terminal year (red) are significantly lower than measurements at other ages (grey), but note that the association (slope) with Δage is for illustration purposes only and was not statistically tested.

Figure 2. Body mass ageing trajectories of (A) males (n = 1,316 measurements on 171 individuals) and (B) females (n = 2,570 measurements on 322 individuals) with predictions of the best-fitting GLMs (table 1) with grey areas 95%CI. For males, the thick dashed-line shows the threshold age at onset of the body mass decline (1.9 or 48.3 years) with thin dashed-lines the 4 ΔAICc-CI [46.6, 52.3]. For females, measurements in the terminal year (red) are significantly lower (intercept) than measurements at other ages (blue). Note that the terminal slope is for illustration purposes only and was not statistically tested.

Table 1. Best fitting body mass ageing trajectories (bold) for males and females, using GAMs (top) and GLMs (bottom) with for each model ageing trajectories ranked from the least to the most complex. AICc: second-order Akaike Information Criterion; ΔAICc: change in AICc relative to the best fitting model; k: degrees of freedom.

Model type	Model	Males			Females		
		k	AICc	ΔAICc	k	AICc	ΔAICc
GAM							
null	log(bm) ~ 1	4.0	-2,835.8	386.1	4.0	-5,422.6	217.1
smooth1	**log(bm) ~ s(Δage) + age-last**	**8.1**	**-3,221.9**	**0.0**	8.7	-5,613.9	25.8
+terminal	log(bm) ~ terminal + s(Δage) + age-last	9.1	-3,221.9	0.0	10.0	-5,627.3	12.4
smooth2	log(bm) ~ s(Δage) + age-last + age-last^2	9.1	-3,221.2	0.7	9.7	-5,627.5	12.2
+terminal	**log(bm) ~ terminal + s(Δage) + age-last + age-last^2**	10.1	-3,221.3	0.6	**10.9**	**-5,639.7**	**0.0**
GLM							
null	log(bm) ~ 1	3.0	-2,830.3	364.0	4.0	-5,442.9	160.7
linear	log(bm) ~ Δage + age-last	5.0	-3,163.9	30.4	6.0	-5,586.1	17.5
+terminal	log(bm) ~ terminal + Δage + age-last	6.0	-3,158.7	35.6	7.0	-5,592.3	11.3
age-last^2	log(bm) ~ Δage + age-last + age-last^2	6.0	-3,156.4	37.9	7.0	-5,598.7	4.9
+terminal	**log(bm) ~ terminal + Δage + age-last + age-last^2**	7.0	-3,151.2	43.1	**8.0**	**-5,603.6**	**0.0**
Δage^2	log(bm) ~ Δage + Δage^2 + age-last	6.0	-3,178.5	15.8	7.0	-5,571.6	32.0
+terminal	log(bm) ~ terminal + Δage + Δage^2 + age-last	7.0	-3,172.9	21.4	8.0	-5,577.9	25.7
quadratic	log(bm) ~ Δage + Δage^2 + age-last + age-last^2	7.0	-3,171.0	23.3	8.0	-5,584.3	19.3
+terminal	log(bm) ~ terminal + Δage + Δage^2 + age-last + age-last^2	8.0	-3,165.4	28.9	9.0	-5,589.3	14.3
threshold	**log(bm) ~ Δage1 + Δage2 + age-last**	**7.0**	**-3,194.3**	**0.0**	8.0	-5,580.5	23.1
+terminal	log(bm) ~ terminal + Δage1 + Δage2 + age-last	8.0	-3,189.3	5.0	9.0	-5,586.0	17.6
threshold (age-last²)	log(bm) ~ Δage1 + Δage2 + age-last + age-last^2	8.0	-3,186.7	7.6	9.0	-5,592.6	11.0
terminal	log(bm) ~ terminal + Δage1 + Δage2 + age-last + age-last^2	9.0	-3,181.8	12.5	10.0	-5,596.9	6.7

Discussion

We tested whether in a species with marked male-biased intrasexual competition, males showed an earlier and/or faster rate of body mass ageing than females. Both sexes gained mass during early adulthood. However, males began to lose mass from 48.3 years old onwards. In contrast, females lost body mass generally at an older age, namely in their last year of life. Here we discuss the implications of our results in the light of theories of ageing (Williams, 1957) and of the management of Asian elephants.

Asian elephants show male-biased intra-sexual competition, with males being more aggressive (Seltmann et al., 2019), fighting more for dominance and showing higher rates of mortality at all ages than females, including during early development, as calves and during adulthood (Lahdenperä et al., 2018). In such species, classic theory predicts that males should show an earlier onset or accelerated ageing (Williams, 1957). Indeed, in several polygynous mammals, males display higher rates of body mass declines than females, suggested to be due to their stronger intrasexual competition (Beirne et al., 2015; Clutton-Brock & Isvaran, 2007). Our results are consistent with those studies and with the prediction of the classic theory of ageing. Here, the prediction of the classic theory is not disrupted by high early-life condition-dependent mortality in males (Chen & Maklakov, 2014) or by canalisation (Flatt, 2005).

Our study is subject to a number of limitations when it comes down to identifying why sexes may differ in their ageing trajectories. First, it is possible that male elephants in our timber elephant population are used more for tasks requiring strength or tusks, thereby causing an earlier onset of body mass declines in males than in females.

However, both sexes fall under the same government-set workload, care and
retirement regulation, except for females' maternity leave. One substantial difference
between sexes is that parental care is concentrated on females, with for example
only females being given 'parental leave' following reproduction (Toke Gale, 1974).
However, since maternity is energetically expensive and no more favourable than
timber working, this is unlikely to lead to the delayed onset of body mass declines in
females. An ideal test would be to analyse the effect of timber work and maternity
leave on body mass dynamics.

Second, elephants have a specific dentition that consists of molar teeth that
eventually wear down at the end of their lives, and pathologic malocclusions or lack
of molars can lead to weight loss and death by starvation. Although both sexes have
the same molar dental anatomy, it is possible that the earlier onset of body mass
declines in males may reflect sex-specific differences in tooth wear rather than sex-
specific sexual selection. Indeed, in captive elephants, dental problems have been
described well before the age at onset of the terminal decline found in this study
(Gaillard et al., 2015) and, if there is sex-specific tooth wear, this could be associated
with the earlier onset of body mass decline in males. Third, male elephants have
recurring periods of physiological "musth" throughout their adult lives, which can
temporarily but profoundly impact the body mass of individual males (Eisenberg et
al., 1971) thereby affecting the body mass ageing trajectory. Fourth, in our study, we
did not find any evidence for body mass-based selective disappearance, but, as it is
often the case in long-lived species, the average longitudinal individual monitoring is
short relative to the lifespan of this species (*e.g.* Global BMI Mortality Collaboration et
al., 2016; Prospective Studies Collaboration, 2009), and hence we only have limited
power to detect such association. It is possible that there are sex-specific dynamics

of selective disappearance, but whether that is the case in Asian elephants remains

to be shown. An analysis with more longitudinal data would be useful to tackle this

question. Fifth, for both sexes, it is possible that maximum body mass is set by

physiological and ecological constraints as indicated by the weight growth curves

found earlier in this population (Mumby, Chapman, et al., 2015). These constraints

could be to some extent sex-specific, although at this point, we can only speculate as

to why these constraints may drive sex-specific ageing trajectory. Finally, we found a

maximum body mass in males but not in females. This sex-specific differences could

be driven by the fact that male elephants benefit more than females from being heavy

during intrasexual competition (Sukumar, 2003). However, another factor to take into

account is that retirement occurs at around 50 years in both sexes, which likely

diminishes physical exercise and allows more time for foraging, thereby continuing

the weight gain. The reduced intrasexual competition in females relative to males,

together with this retirement, could lead to the continued mass gain of females. Given

that elephants in the wild do not experience timber labour and retirement, we cannot

exclude that the sex-specific body mass ageing trajectories could be different in a

wild (non-working) population of Asian elephants compared to those found in our

study.

We found that females experienced a terminal body mass decline in the last year of

life. Our data contain both males and females among the oldest ages (>50), hence

sex-specific terminal decline is unlikely to emerge from differences in lifespan. In

European badgers (*Meles meles*), a species in which females outlive males, both

sexes displayed terminal body mass declines (Beirne et al., 2015). It is possible that

the sex-specific terminal declines in our study resulted from differences in power, with

5 old dead males and 18 old dead females. Indeed, for both males and females, the

coefficient and effect size of the terminal terms were negative, but the effect size in

males remained about half of that in females (Cohen's d_{males} = -0.045 [-0.10, 0.01],

Cohen's $d_{females}$ = -0.071 [-0.11, -0.03]). Hence, it is possible that the sex-specific

terminal effect is driven by power issues and we look forward to testing that with

several more years of monitoring.

Terminal declines emphasise that the chronological age is rarely a perfect estimation

of the biological age which can better describe the 'true biological state' of an

organism (Klemera & Doubal, 2006). In that sense, terminal decline is a biomarker of

health and remaining lifespan. The 'terminal illness' hypothesis refers to the age-

independent decrease of a trait value, related to the imminent death of the individual

(Coulson & Fairweather, 2001) and such terminal effects were shown for example for

body mass and sexual signals in mammals and birds (Beirne et al., 2015; Simons et

al., 2016). For which traits or under which conditions to expect terminal declines

remains yet poorly understood but our study highlights the importance of studying

sex-specific differences in ageing and illustrates the need to improve our

understanding of the mechanisms driving the diversity of ageing patterns in the wild.

References

Bartoń, K. (2019). *MuMIn : Multi-Model Inference* (1.43.15) [Computer software].

https://CRAN.R-project.org/package=MuMIn

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software*, *67*(1), 1 48.

https://doi.org/10.18637/jss.v067.i01

Beirne, C., Delahay, R., & Young, A. (2015). Sex differences in senescence : The role of

intra-sexual competition in early adulthood. *Proceedings of the Royal Society B:*

Biological Sciences, *282*(1811), 20151086. https://doi.org/10.1098/rspb.2015.1086

Bolker, B. M. (2008). *Ecological Models and Data in R*. Princeton University Press.

https://press.princeton.edu/books/hardcover/9780691125220/ecological-models-and-

data-in-r

Bonduriansky, R., Maklakov, A., Zajitschek, F., & Brooks, R. (2008). Sexual selection, sexual

conflict and the evolution of ageing and life span. *Functional Ecology*, *22*(3),

443□453. https://doi.org/10.1111/j.1365-2435.2008.01417.x

Briga, M., Jimeno, B., & Verhulst, S. (2019). Coupling lifespan and aging? The age at onset

of body mass decline associates positively with sex-specific lifespan but negatively

with environment-specific lifespan. *Experimental Gerontology*, *119*, 111□119.

https://doi.org/10.1016/j.exger.2019.01.030

Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and

multimodel inference in behavioral ecology□: Some background, observations, and

comparisons. *Behavioral Ecology and Sociobiology*, *65*(1), 23□35.

https://doi.org/10.1007/s00265-010-1029-6

Chen, H., & Maklakov, A. A. (2014). Condition Dependence of Male Mortality Drives the

Evolution of Sex Differences in Longevity. *Current Biology*, *24*(20), 2423□2427.

https://doi.org/10.1016/j.cub.2014.08.055

Clutton-Brock, T. H., & Isvaran, K. (2007). Sex differences in ageing in natural populations of

vertebrates. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1629),

3097□3104. https://doi.org/10.1098/rspb.2007.1138

Coulson, J. C., & Fairweather, J. A. (2001). Reduced reproductive performance prior to death

in the Black-legged Kittiwake□: Senescence or terminal illness? *Journal of Avian*

Biology, *32*(2), 146□152. https://doi.org/10.1034/j.1600-048X.2001.320207.x

Douhard, F., Gaillard, J.-M., Pellerin, M., Jacob, L., & Lemaître, J.-F. (2017). The cost of

growing large□: Costs of post-weaning growth on body mass senescence in a wild

mammal. *Oikos*, *126*(9), 1329□1338. https://doi.org/10.1111/oik.04421

Eisenberg, J. F., McKay, G. M., & Jainudeen, M. R. (1971). Reproductive behavior of the

Asiatic elephant (Elephas maximus maximus L.). *Behaviour*, *38*(3), 193□225.

https://doi.org/10.1163/156853971x00087

Flatt, T. (2005). The evolutionary genetics of canalization. *The Quarterly Review of Biology*,

80(3), 287□316. https://doi.org/10.1086/432265

Gaillard, J.-M., Berger, V., Tidière, M., Duncan, P., & Lemaître, J.-F. (2015). Does tooth wear

influence ageing? A comparative study across large herbivores. *Experimental*

Gerontology, *71*, 48□55. https://doi.org/10.1016/j.exger.2015.09.008

Global BMI Mortality Collaboration, Di Angelantonio, E., Bhupathiraju, S., Wormser, D., Gao,

P., Kaptoge, S., Berrington de Gonzalez, A., Cairns, B., Huxley, R., Jackson, C.,

Joshy, G., Lewington, S., Manson, J., Murphy, N., Patel, A., Samet, J., Woodward,

M., Zheng, W., Zhou, M., … Hu, F. (2016). Body-mass index and all-cause

mortality□: Individual-participant-data meta-analysis of 239 prospective studies in four

continents. *The Lancet*, *388*(10046), 776□786. https://doi.org/10.1016/S0140-

6736(16)30175-1

Hämäläinen, A., Dammhahn, M., Aujard, F., Eberle, M., Hardy, I., Kappeler, P. M., Perret,

M., Schliehe-Diecks, S., & Kraus, C. (2014). Senescence or selective disappearance?

Age trajectories of body mass in wild and captive populations of a small-bodied

primate. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1791),

20140830. https://doi.org/10.1098/rspb.2014.0830

Jones, O. R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C. G., Schaible, R., Casper,

B. B., Dahlgren, J. P., Ehrlén, J., García, M. B., Menges, E. S., Quintana-Ascencio, P.

F., Caswell, H., Baudisch, A., & Vaupel, J. W. (2014). Diversity of ageing across the

tree of life. *Nature*, *505*(7482), 169□173. https://doi.org/10.1038/nature12789

Klemera, P., & Doubal, S. (2006). A new approach to the concept and computation of

biological age. *Mechanisms of Ageing and Development*, *127*(3), 240□248.

https://doi.org/10.1016/j.mad.2005.10.004

Lahdenperä, M., Mar, K. U., Courtiol, A., & Lummaa, V. (2018). Differences in age-specific mortality between wild-caught and captive-born Asian elephants. *Nature Communications*, *9*(1), 3023. https://doi.org/10.1038/s41467-018-05515-8

Lahdenperä, M., Mar, K. U., & Lummaa, V. (2016). Nearby grandmother enhances calf survival and reproduction in Asian elephants. *Scientific Reports*, *6*(1), 27213. https://doi.org/10.1038/srep27213

Leimgruber, P., Senior, B., Aung, M., Songer, M., Mueller, T., Wemmer, C., & Ballou, J. (2008). Modeling population viability of captive elephants in Myanmar (Burma): Implications for wild populations. *Animal Conservation*, *11*, 198–205. https://doi.org/10.1111/j.1469-1795.2008.00172.x

Lemaître, J.-F., Ronget, V., Tidière, M., Allainé, D., Berger, V., Cohas, A., Colchero, F., Conde, D. A., Garratt, M., Liker, A., Marais, G. A. B., Scheuerlein, A., Székely, T., & Gaillard, J.-M. (2020). Sex differences in adult lifespan and aging rates of mortality across wild mammals. *Proceedings of the National Academy of Sciences*, *117*(15), 8546–8553. https://doi.org/10.1073/pnas.1911999117

Lynch, E. C., Lummaa, V., Htut, W., & Lahdenperä, M. (2019). Evolutionary significance of maternal kinship in a long-lived mammal. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *374*(1780), 20180067. https://doi.org/10.1098/rstb.2018.0067

Maklakov, A. A., & Lummaa, V. (2013). Evolution of sex differences in lifespan and aging: Causes and constraints. *BioEssays*, *35*, 717–724. https://doi.org/10.1002/bies.201300021

Monaghan, P., Charmantier, A., Nussey, D. H., & Ricklefs, R. E. (2008). The evolutionary ecology of senescence. *Functional Ecology*, *22*(3), 371–378. https://doi.org/10.1111/j.1365-2435.2008.01418.x

Mumby, H. S., Chapman, S. N., Crawley, J. A. H., Mar, K. U., Htut, W., Thura Soe, A., Aung, H. H., & Lummaa, V. (2015). Distinguishing between determinate and indeterminate

growth in a long-lived mammal. *BMC Evolutionary Biology*, *15*(1), 214.

https://doi.org/10.1186/s12862-015-0487-x

Mumby, H. S., Mar, K. U., Thitaram, C., Courtiol, A., Towiboon, P., Min-Oo, Z., Htut-Aung, Y.,

Brown, J. L., & Lummaa, V. (2015). Stress and body condition are associated with

climate and demography in Asian elephants. *Conservation Physiology*, *3*(1), cov030.

https://doi.org/10.1093/conphys/cov030

Nussey, D. H., Kruuk, L. E. B., Morris, A., & Clutton-Brock, T. H. (2007). Environmental

conditions in early life influence ageing rates in a wild population of red deer. *Current*

Biology, *17*(23), R1000⬚R1001. https://doi.org/10.1016/j.cub.2007.10.005

Nussey, D. H., Kruuk, L., Morris, A., Clements, M., Pemberton, J., & Clutton-Brock, T. (2009).

Inter- and Intrasexual Variation in Aging Patterns across Reproductive Traits in a Wild

Red Deer Population. *The American Naturalist*, *174*, 342⬚357.

https://doi.org/10.1086/603615

Oo, Z. M. (2010). The training methods used in Myanma Timber Enterprise. *Gajah*, *33*,

58⬚61.

Prospective Studies Collaboration. (2009). Body-mass index and cause-specific mortality in

900 000 adults⬚: Collaborative analyses of 57 prospective studies. *The Lancet*,

373(9669), 1083⬚1096. https://doi.org/10.1016/S0140-6736(09)60318-4

R Core Team. (2020). *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing. https://www.R-project.org

Rando, T. A., & Wyss-Coray, T. (2021). Asynchronous, contagious and digital aging. *Nature*

Aging, *1*(1), 29⬚35. https://doi.org/10.1038/s43587-020-00015-1

Seltmann, M. W., Helle, S., Htut, W., & Lahdenperä, M. (2019). Males have more aggressive

and less sociable personalities than females in semi-captive Asian elephants.

Scientific Reports, *9*(1), 2668. https://doi.org/10.1038/s41598-019-39915-7

Simons, M. J. P., Briga, M., & Verhulst, S. (2016). Stabilizing survival selection on

presenescent expression of a sexual ornament followed by a terminal decline.

Journal of Evolutionary Biology, *29*(7), 1368⬚1378. https://doi.org/10.1111/jeb.12877

Simpson, G. L., & Singmann, H. (2021). *gratia□: Graceful 'ggplot'-based graphics and other*

functions for GAMs fitted using « mgcv » (0.6.0) [Computer software].

Sukumar, R. (2003). *The living elephants□: Evolutionary ecology, behavior, and*

conservation. Oxford University Press.

Tidière, M., Gaillard, J.-M., Müller, D. W. H., Lackey, L. B., Gimenez, O., Clauss, M., &

Lemaître, J.-F. (2015). Does sexual selection shape sex differences in longevity and

senescence patterns across vertebrates? A review and new insights from captive

ruminants. *Evolution*, *69*(12), 3123□3140. https://doi.org/10.1111/evo.12801

Toke Gale, U. (1974). *Burmese timber elephant*. Trade Corporation.

van de Pol, M., & Verhulst, S. (2006). Age□dependent traits□: A new statistical model to

separate within□ and between□individual effects. *The American Naturalist*, *167*(5),

766□773. https://doi.org/10.1086/503331

van de Pol, M., & Wright, J. (2009). A simple method for distinguishing within- versus

between-subject effects using mixed models. *Animal Behaviour*, *77*(3), 753□758.

https://doi.org/10.1016/j.anbehav.2008.11.006

Williams, G. C. (1957). Pleiotropy, natural selection, and the evolution of senescence.

Evolution, *11*(4), 398□411. https://doi.org/10.2307/2406060

Wood, S. (2017). Generalized Additive Models□: An Introduction With R. In *Generalized*

Additive Models□: An Introduction with R, Second Edition (2nd éd., Vol. 66).

Chapman and Hall/CRC Press. https://doi.org/10.1201/9781315370279

Zaw, U. K. (1997). Utilization of elephants in timber harvesting in Myanmar. *Gajah*, *17*, 9□22.

Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed Effects*

Models and Extensions in Ecology with R. Springer-Verlag.

https://doi.org/10.1007/978-0-387-87458-6

Ethics

The study was performed with the permission from the Myanmar Ministry of Natural Resources and Environmental Conservation and following the University of Turku ethical guidelines. Note that the University of Turku does not provide project numbers.

Data accessibility

The dataset supporting the conclusions of this article will be published on a digital repository and made publicly available upon acceptance.

Authors' contributions

Lucas D. Lalande: Formal analysis, Data curation, Writing – Original draft, Writing – Review & Editing, Visualization, **Virpi Lummaa:** Conceptualization, Supervision, **Htoo H. Aung:** Investigation, Resources, **Win Htut:** Investigation, Resources, **U. Kyaw Nyein:** Investigation, Resources, **Vérane Berger:** Conceptualization, Writing – Review and Editing, Supervision, **Michael Briga:** Conceptualization, Writing – Review and Editing, Supervision.

Competing interests

None. We declare we have no competing interests.

Funding

This work was supported by the Academy of Finland [292368], the European Research Council [ERC-2014-CoG 648766] and the Ella & Georg Ehrnrooth Foundation.

Acknowledgements

We thank the Myanma Timber Enterprise (MTE) and the Myanmar Ministry of Natural Resources and Environmental Conservation for their collaboration and all of the MTE staff enabling this study by recording elephant life events and morphology for so long. Particularly, we thank Mu Mu Thein and Nina Aro for their patience in helping translating countless elephant morphological measurements from burmese to english. We also thank Carly Lynsdale for ensuring our english was fluent. We thank *the Academy of Finland, the European Research Council and the Ella & Georg Ehrnrooth Foundation for fundings.*