ZIKV disrupts placental ultrastructure and drug transporter expression in mice

Cherley Borba Vieira de Andrade[1], Victoria Regina de Siqueira Monteiro[1], Sharton

Vinicius Antunes Coelho[2], Hanailly Ribeiro Gomes[1], Ronny Paiva Campos de Sousa[1],

Veronica Muller de Oliveira Nascimento[1], Flavia Fonseca Bloise[1], Stephen

Matthews[3,4,5,6], Enrrico Bloise[7], Luciana Barros de Arruda[2], Tania Maria Ortiga-

Carvalho[1]*.

[1]Institute of Biophysics Carlos Chagas Filho, Federal University of Rio de Janeiro, Rio

de Janeiro, RJ, Brazil.

[2] Institute of Microbiology Paulo de Góes, Federal University of Rio de Janeiro, Rio de

Janeiro, RJ, Brazil.

[3] Department of Physiology, Faculty of Medicine, University of Toronto, Toronto,

Canada.

[4] Department of Obstetrics & Gynecology, Faculty of Medicine, University of Toronto,

Toronto, Canada.

[5] Department of Medicine, Faculty of Medicine, University of Toronto, Toronto,

Canada.

[6] Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Toronto, Canada.

[7] Department of Morphology, Federal University of Minas Gerais, Belo Horizonte, BH,

Brazil.

* Correspondence:

Tania Maria Ortiga-Carvalho (TMO-C)

taniaort@biof.ufrj.br

Key words: ZIKV, placenta, P-glycoprotein (P-gp), breast cancer resistance

protein (Bcrp), Abca1, Abcg1, ultrastructure, cytokine, chemokine

Abstract

Congenital Zika virus (ZIKV) infection can induce fetal brain abnormalities. Here, we investigated whether maternal ZIKV infection affects placental physiology and metabolic transport potential and impacts the fetal outcome, regardless of viral presence in the fetus at term. Low (10^3 PFU-ZIKVPE243; low ZIKV) and high (5×10^7 PFU-ZIKVPE243; high ZIKV) virus titers were injected into immunocompetent (ICompetent C57BL/6) and immunocompromised (ICompromised A129) mice at gestational day (GD) 12.5 for tissue collection at GD18.5 (term). High ZIKV elicited fetal death rates of 66% and 100%, whereas low ZIKV induced fetal death rates of 0% and 60% in C57BL/6 and A129 dams, respectively. All surviving fetuses exhibited intrauterine growth restriction (IUGR) and decreased placental efficiency. High-ZIKV infection in C57BL/6 and A129 mice resulted in virus detection in maternal spleens and placenta, but only A129 fetuses presented virus RNA in the brain. Nevertheless, pregnancies in both strains produced fetuses with decreased head sizes ($p<0.05$). Low-ZIKV-A129 dams had higher IL-6 and CXCL1 levels ($p<0.05$), and their placentas showed increased CCL-2 and CXCL-1 contents ($p<0.05$). In contrast, low-ZIKV-C57BL/6 dams had an elevated CCL2 serum level and increased type I and II IFN expression in the placenta. Notably, less abundant microvilli and mitochondrial degeneration were evidenced in the placental labyrinth zone (Lz) of ICompromised and high-ZIKV-ICompetent mice but not in low-ZIKV-C57BL/6 mice. In addition, decreased placental expression of the drug transporters P-glycoprotein (P-gp) and breast cancer resistance protein (Bcrp) and the lipid transporter Abca1 was detected in all ZIKV-infected groups, but Bcrp and Abca1 were only reduced in ICompromised and high-ZIKV ICompetent mice. Our data indicate that gestational ZIKV infection triggers specific proinflammatory responses and affects placental turnover and transporter expression in a manner dependent on

virus concentration and maternal immune status. Placental damage may impair proper fetal-maternal exchange function and fetal growth/survival, likely contributing to congenital Zika syndrome.

1. Introduction

Congenital Zika virus (ZIKV) infection can be associated with adverse pregnancy outcomes. Neonates born from ZIKV-positive pregnancies may develop severe neurological abnormalities, placental pathologies and intrauterine growth restriction (IUGR), among other complications (1). ZIKV vertical transmission has become a major public health issue worldwide, especially in Brazil, where more than 200,000 ZIKV-positive cases have been confirmed and over 2,000 congenital microcephaly births have been reported (2–6). These numbers represent a 20-fold rise in the incidence of congenital microcephaly in Brazil during the years of the ZIKV pandemic, with similar increases reported elsewhere in Latin America (2,3,7). Importantly, while the ZIKV pandemic is currently thought to be controlled, evidence points to a possible silent ZIKV spread across the Americas (8,9), highlighting the need for improved knowledge of the possible routes of vertical ZIKV transmission and its association with disruptive inflammatory and developmental phenotypes and the need for new avenues of prevention and treatment.

Previous studies have investigated the possible pathways involved in vertical ZIKV transmission. Miranda and colleagues (10) showed that in humans, ZIKV infection changed the pattern of tight junction proteins, such as claudin-4, in syncytiotrophoblasts. Jurado et al. (2016) suggested that the migratory activities of Hofbauer cells (feto-placental macrophages) could help disseminate ZIKV to the fetal brain (11). Other recent studies have shown that placental villous fibroblasts, cytotrophoblasts, endothelial cells and Hofbauer cells are permissive to ZIKV, and

placentae from ZIKV-infected women had chorionic villi with a high mean diameter

(11–14). Furthermore, in 2019, Rathore et al. demonstrated that pregnant mice carrying

high levels of antibodies against dengue virus (DENV) exhibited increased ZIKV

vertical transmission associated with severe microcephaly-like syndrome,

demonstrating another possible mechanism of antibody-dependent vertical ZIKV

transmission (15). However, at present, further studies are required to identify the

precise mechanism of maternal-fetal ZIKV transmission.

Many mouse models have been developed to identify how ZIKV overcomes

placental defenses. Initially, limited information was obtained due to the apparent

inability of the virus to infect wild-type (WT) mice (16). ZIKV NS5 targets the

interferon signaling pathway in humans but not in mice (17). Thus, WT mice show no

clear evidence of clinical disease (17,18) and are of limited use in modeling the disease.

However, mice lacking an interferon signaling response show evidence of disease and

have been widely used to investigate ZIKV infection during pregnancy (8,17,19).

The interferon system, especially type III interferon, is a key mechanism of host

defense and a viral target for immune evasion (20). Type III interferons have a role in

protection against ZIKV infection in human syncytiotrophoblasts from term placenta

(21). Luo et al. have shown that inhibition of Toll-like receptors 3 and 8 inhibits the

cytokine output of ZIKV-infected trophoblasts (22). In addition, viral replication

coincides with the induction of proinflammatory cytokines, such as interleukin [IL]-6.

This cytokine has a crucial role in inflammation and affects the homeostatic processes

related to tissue injury and activation of stress-related responses (23,24). ZIKV infection

can trigger an inflammatory response with IL-6 release (11,25).

Maternal infection has profound effects on placental permeability to drugs and

environmental toxins. Changes in the expression and function of specific ABC

transporters in the placenta and yolk sac following infective and inflammatory stimuli

have been demonstrated (26–30). ABC transporters are efflux transporters that control

the biodistribution of several endogenous and exogenous substrates, including

xenobiotics (antiretrovirals and synthetic glucocorticoids), steroid hormones (estrogens

and androgens), nutrients (folate and cholesterol) and immunological factors

(chemokines and cytokines) within the maternal-fetal interface (31). The best described

ABC transporters in the placenta are P-glycoprotein (P-gp; also known as multidrug

resistance protein 1, MDR1), breast cancer resistance protein (Bcrp) and the lipid Abca1

and Abcg1 transporters. P-gp and Bcrp transporters are responsible for preventing fetal

accumulation of xenobiotics and environmental toxins that may be present in the

maternal circulation, whereas Abca1 and Abcg1 control the placental exchange of

cytotoxic oxysterol and lipid permeability throughout pregnancy; therefore, they play an

important role in fetal protection and placental lipid homeostasis (26).

Despite the limited number of studies showing ZIKV infection in

immunocompetent mice, intrauterine inoculation with a high virus titer was previously

demonstrated to result in decreased fetal viability, with worse outcomes following

infection in early gestation (32). In another report, intravenous infection on a very early

embryonic day resulted in fetal demise even though the virus was not found in the fetal

compartment in most of the treated animals (33). In the present study, we hypothesize

that maternal exposure to ZIKV affects placental function, including placental

ultrastructure and ABC transporter (P-gp, Bcrp, Abca1 and Abcg1) protein expression,

even in the absence of vertical transmission and that these effects are dependent on viral

infective titers and maternal immune status.

2. Materials and methods

2.1 Virus preparation and storage

The Brazilian ZIKV$_{PE243}$ (GenBank ref. number KX197192) strain was isolated

from a febrile case during the ZIKV outbreak in the state of Pernambuco, Brazil and

was kindly provided by Dr Ernesto T. Marques Jr. (Centro de Pesquisa Aggeu

Magalhães, FIOCRUZ, PE). Viruses were propagated in C6/36 cells, and viral titers

were determined by plaque assays in Vero cells, as previously described (34).

Supernatants of noninfected C6/36 cells cultured under the same conditions were used

as mock controls.

2.2 Animal experimentation and study design

Two mouse strains were used in the study: immunocompetent (ICompetent)

C57BL/6 and immunocompromised (ICompromised) (type 1 *Ifnr*-deficient) A129

strains. Since we were unable to consistently produce viable pregnancies by mating

A129 males and females in our experimental settings, we mated A129 females (n=15)

with C57BL/6 males (n=4) to produce ICompromised C57BL6/A129 pregnancies,

whereas ICompetent C57BL/6 pregnancies were obtained by mating male (n=6) and

female (n=35) C57BL/6 mice (8-10 weeks old). Animals were kept in a controlled

temperature room (23°C) with a light/dark cycle of 12 hours and *ad libitum* access to

water and food. After detection of the proestrous/estrous phase via vaginal cytology,

copulation was confirmed by visualization of the vaginal plug and considered

gestational day 0.5 (GD0.5). Maternal weight was monitored for confirmation of

pregnancy; thus, females were weighed on GD0.5 and GD12.5, and females with a

weight gain greater than 3 g were considered pregnant and entered randomly in the

experimental groups. Experimental protocols were approved by the Animal Care

Committee of the Health Sciences Center, Federal University of Rio de Janeiro (CEUA-

036/16 and 104/16) and registered with the Brazilian National Council for Animal

Experimentation Control.

On GD12.5, pregnant mice (ICompetent and ICompromised pregnancies) were

injected with a single intravenous (i.v.) titer of ZIKV or mock control. ICompetent and

ICompromised pregnant mice were randomly subdivided into three experimental

groups: the mock (control) group, which received an injection of supernatant from

noninfected C6/36 cells (ICompetent mock and ICompromised mock); the high-ZIKV-

titer group, inoculated with $5x10^7$ plaque-forming units (PFU) of ZIKV$_{PE243}$

(ICompetent high and ICompromised high); and the ZIKV low-titer group, injected

with 10^3 PFU of ZIKV$_{PE243}$ (ICompetent low and ICompromised low).

On the morning of GD18.5, all animals were euthanized with a sodium

phenobarbital overdose of 300 mg/kg. Maternal blood was collected via cardiac

puncture, centrifuged (10 min, 4000 g) and stored at -20°C. The maternal brain and

spleen and all placentae and all fetuses were dissected, collected and weighed, followed

by fetal head isolation and measurement. The three placentae closest to the mean weight

in a litter were selected for further analysis and cut in half using umbilical cord insertion

as a reference (35–37). One-half of the placental disk was frozen in liquid nitrogen for

qPCR, and the other half was fixed overnight in buffered paraformaldehyde (4%,

Sigma-Aldrich, Brazil) for ultrastructural and protein expression/localization analysis.

Matched fetal heads, maternal brains and spleens were frozen in liquid nitrogen for

qPCR. Of important note, all fetuses obtained from ICompromised pregnancies were

heterozygous.

2.3 ZIKV RNA quantification via RT-qPCR

ZIKV load was evaluated in maternal blood, brains and spleens and in the

placentae and fetal heads. Brains, spleens, placentae and fetal heads were macerated in

RPMI medium (Gibco™ RPMI 1640 Medium) normalized by the ratio of 0.2 mg of

tissue to every 1 µl of medium and plotted per gram of tissue. The macerated volume

was centrifuged at 4500 g for 5 min to remove tissue residues, and then, 500 µl of the

centrifuged volume was used for RNA extraction using 1 mL of TRIzol reagent (Life

Technologies, Thermo Fischer, USA). Treatment with DNase I (Ambion, Thermo

Fischer, USA) was performed to prevent contamination by genomic DNA. cDNA was

synthesized using a cDNA High Capacity Kit (Applied Biosystems, Thermo Fischer,

USA) according to the manufacturer's instructions by subjecting the samples to the

following cycle: 25°C for 10 min, 37°C for 120 min and 85°C for 5 min. qPCR was

performed using a StepOnePlus Real-Time qPCR system, TaqMan Master Mix

Reagents (Applied Biosystems, Thermo Fischer, USA) and primers and probes specific

for the protein E sequence (38). Samples were then subjected to the following cycle:

50°C for 2 min, followed by 40 cycles of 95°C for 10 min, 95°C for 15 sec, and 60°C

for 1 min.

2.3.1 RT-qPCR

The placenta was macerated in 1.5 mL of TRIzol reagent (Life Technologies,

Thermo Fischer, USA). RNA extraction was performed following the manufacturer's

protocol. cDNA was prepared using Power SYBR Green PCR Master Mix (Life

Technologies, Thermo Fisher, USA). The reaction was carried out for selected genes

using intron-spanning primers (Table 1) and the StepOnePlus Real-Time PCR system

(Life Technologies, Thermo Fischer, USA). Samples were subjected to the following

cycle: 95°C for 10 min, followed by 40 amplification cycles consisting of DNA

denaturation for 30 sec at 95°C and annealing of primers for 30 sec at 60°C. The

threshold cycle (Ct) was determined for each gene of interest and for the reference

genes glycerol 3-phosphate dehydrogenase (*Gapdh*) and RNA Polymerase II Subunit A

(*Polr2a)*. The relative expression of each gene was calculated using $2^{-\Delta\Delta CT}$ (39) and

graphically expressed as the fold-increase. The efficiency was calculated using the

standard curve method. The melting curves were analyzed for each sample.

Table 1: Primer sequences for the real-time PCR assay.

Gene	Primer sequences	GenBank accession no.
ZIKV	5'CCGCTGCCCAACACAAG3' 5'CCACTAACGTTCTTTTGCAGACAT3'	
Il6	5'TCATATCTTCAACCAAGAGGTA3' 5'CAGTGAGGAATGTCCACAAACTG3'	NM_031168.2
Il1b	5'GTAATGAAAGACGGCACACC3' 5'ATTAGAAACAGTCCAGCCCA3'	XM_006498795.4
Il10	5'TAAGGGTTACTTGGGTTGCCAAG3' 5'CAAATGCTCCTTGATTTCTGGGC3'	NM_010548.2
Ifng	5'AGCAACAGCAAGGCGAAAA3' 5'CTGGACCTGTGGGTTGTTGA3'	NM_008337.4
Ifn1	5'CTGGAGCAGCTGAATGGAAAG3' 5'CTTGAAGTCCGCCCTGTAGGT3'	NM_010510.1
Gapdh	5'CTTTGTCAAGCTCATTTCCTGG3' 5'TCTTGCTCAGTGTCCTTGC3'	XM_017321385.2
Tnf	5'CCTCACACTCAGATCATCTTCTCA3' 5'TGGTTCTCTTTGAGATCCATGC3'	NM_013693.3
Polr2a	5'TCTGCCAAGAATGTGACGCT3' 5'CCAAGCGGCAAAGAATGTCC3'	NM_001291068.1

2.4 Detection of cytokines and chemokines in maternal serum and the placenta

Initially, placental tissue was homogenized in extraction buffer (50 mM Tris,

150 mM NaCl, 1X Triton, 0.1% SDS, 5 mM EDTA, 5 mM NaF, 50 mM sodium

pyrophosphate, 1 mM sodium orthovanadate, pH 7.4) containing complete protease

inhibitor cocktail (Roche Applied Science, Germany) with TissueLyser LT (Qiagen,

Germany). The protein concentration of each sample was analyzed using a Pierce™

BCA Protein Assay Kit (Thermo Scientific, USA) according to the manufacturer's

instructions. Analysis of the cytokines IL-6 and IL-1β and the chemokines monocyte

chemoattractant protein-1 (MCP-1/CCL2) and chemokine (C-X-C motif) ligand 1

(CXCL1) in maternal serum and placenta was performed with MILLIPLEX MAP

Mouse Cytokine/Chemokine Magnetic Bead Panel – Immunology Multiplex Assays

(MCYTOMAG-70K, Merck Millipore, Germany) following the manufacturer's

recommendations. The plate with samples and magnetic beads was analyzed on a

MAGPIX® System (Merck Millipore, Germany). The analyses were performed using

Luminex xPonent®□ for MAGPIX®□ v software. 4.2 (Luminex Corp., USA). For

each reaction well, the MAGPIX Luminex® platform reports the median fluorescence

intensity (MFI) for each of the analytes in the sample. The levels of each analyte were

then calculated against the standard curve. The ratio between the value obtained and the

protein quantification for each sample was determined and plotted.

2.5 Virus titration using plaque assays

Blood from mock- and $ZIKV_{PE243}$-infected mice was collected from the base of

the tail at 4 hours, 48 hours and 144 hours following the appropriate treatments and

subsequently centrifuged at 400 g for 30 min for plasma separation. Samples obtained at

different periods post infection were titrated using a plaque assay. Vero cells (obtained

from ATCC® CCL81™) (African green monkey kidney epithelial cell line) were plated

in 24-well plates at 4×10^4 cells per well in Dulbecco's modified Eagle's medium

(DMEM) (GIBCO, Thermo Fisher, USA) supplemented with 5% fetal bovine serum

(FBS) (GIBCO, Thermo Fisher, USA) and 1% gentamicin (10 µg/ml) (GIBCO, Thermo

Fisher, USA) and cultured overnight for complete adhesion at 37°C with 5% CO_2.

Then, the medium was removed, and the cells were washed with 1x PBS and incubated

with serial (base 10) dilutions of virus in FBS-free medium. After 90 min of incubation

under gentle shaking, the medium was removed, and the cells were washed with 1x PBS

and cultured with 1.5% carboxymethylcellulose (CMC) supplemented with 1% FBS

(GIBCO, Thermo Fisher, USA). After 5 days, the cells were fixed overnight with 4%

formaldehyde and stained with 1% crystal violet in 20% methanol (ISOFAR, Brazil) for

1 hour. Plaques were counted, and the virus yield was calculated and expressed as

plaque-forming units per milliliter (PFU/ml).

2.6 Histological, immunohistochemistry and TUNEL analyses of the placenta

Placental fragments were fixed overnight and subjected to dehydration (increasing ethanol series; ISOFAR, Brazil), diaphanization with xylol (ISOFAR, Brazil) and paraffin (Histopar, Easypath, Brazil). Sections (5 μm) were prepared using a Rotatory Microtome CUT 5062 (Slee Medical GmbH, Germany) and subjected to immunohistochemistry and TUNEL analyses.

For immunohistochemistry, blocking of endogenous peroxidase was performed with 3% hydrogen peroxide diluted in PBS, followed by microwave antigenic recovery in Tris-EDTA (pH=9) and sodium citrate (pH=6) buffers (15 min for Tris-EDTA buffer and 8 min for citrate buffer). Sections were washed in PBS + 0.2% Tween and exposed to 3% PBS/BSA for 1 hour. Sections were then incubated overnight at 4°C with the following primary antibodies: anti-Ki-67 (1:100 – [M3064]; Spring Bioscience, USA), anti-P-gp (1:500 – Mdr1[sc-55510]; Santa Cruz Biotechnology, USA), anti-Bcrp (1:100 – Bcrp [MAB4146]; Merck Millipore, USA), anti-Abcg1 (1:100 – [PA5-13462]; Thermo Fisher Scientific, USA) or anti-Abca1 (1:100 – [ab18180]; Abcam Plc, UK). The next day, sections were incubated with the biotin-conjugated secondary antibody SPD-060 (Spring Bioscience, USA) for 1 hour at room temperature. Three washes were performed with PBS + 0.2% Tween followed by incubation with streptavidin (SPD-060 - Spring Bioscience, USA) for 30 min. Sections were stained with 3,3-diamino-benzidine (DAB) (SPD-060 - Spring Bioscience, USA), counterstained with hematoxylin (Proquímios, Brazil), dehydrated, diaphanized and mounted with a coverslip and Entellan (Merck, Germany).

For analysis of apoptotic nuclei, terminal deoxynucleotidyl transferase dUTP nick-end labeling (TUNEL) staining was performed using an ApopTag® In Situ Peroxidase Detection Kit (S7100, Merck Millipore, USA) according to the

manufacturer's recommendations and as previously described (36). All negative controls

were prepared with omission of the primary antibody.

Image acquisition was performed using a high-resolution Olympus DP72

(Olympus Corporation, Japan) camera coupled to an Olympus BX53 light microscope

(Olympus Corporation, Japan). For nuclear quantification of Ki-67 and TUNEL

immunolabeling, Stepanizer software (40) was used. For this analysis, we evaluated 15

images from different random fields of the Lz (labyrinth zone) and Jz (junctional zone)

for each animal, in a total of five animals from each ICompetent group and three

animals from each ICompromised group. A total of 360 digital images (40X) randomly

captured per placental region (Lz and Jz) were evaluated in each experimental group.

The total number of immunolabeled Ki-67 or TUNEL nuclei in each digital image was

normalized by the total image area to obtain an index of the estimated number of

proliferative and apoptotic nuclei in the entire histological section analyzed. Analysis

was undertaken by two investigators blinded to the treatment.

Quantification of P-gp, Bcrp, Abca1 and Abcg1 staining was performed using

the Image-Pro Plus, version 5.0 software (Media Cybernetics, USA) mask tool. The

percentage of viable tissue area was considered upon exclusion of negative spaces. A

total of 360 digital images (40X) randomly captured per placental region (Lz and Jz)

were evaluated in each experimental group. Analysis was undertaken by two

investigators blinded to the treatment.

2.7 Transmission electron microscopy (TEM)

Sections of the placental Lz and Jz were fixed in paraformaldehyde 4% (Sigma-Aldrich,

Brazil) for 48 hours, postfixed with osmium tetroxide (Electron Microscopy Sciences,

USA) and potassium ferrocyanide (Electron Microscopy Sciences, USA) for 60 min and

dehydrated with an increasing series of acetone (30%, 50%, 70%, 90% and two of

100%) (ISOFAR, Brazil). Sections were subsequently embedded with EPOXI resin

(Electron Microscopy Sciences, USA) and acetone (1:2, 1:1 and 2:1, respectively).

After polymerization, ultrafine sections (70 nm) were prepared (Leica Microsystems,

USA) and collected into 300 mesh copper grids (Electron Microscopy Sciences, USA).

Tissue was contrasted with uranyl acetate and lead citrate and visualized using a JEOL

JEM-1011 transmission electron microscope (JEOL, Ltd., Akishima, Tokyo, Japan).

Digital micrographs were captured using an ORIUS CCD digital camera (Gatan, Inc.,

Pleasanton, California, EUA) at $6000\times$ magnification. An overall qualitative analysis of

the Lz and Jz in different groups was performed by investigating the ultrastructural

characteristics of the mitochondria and the ER cisterns. The qualitative evaluation

consisted of analyzing disruption of the mitochondrial membranes, mitochondrial

morphology, preservation of mitochondrial cristae and matrix intensity (41).

Ultrastructural analysis of nuclear morphology and the presence of microvilli in

trophoblast sinusoidal giant cells was also undertaken. Analysis of ER cisterns was

performed by evaluating the dilation of their lumen (42).

2.8 Statistical analysis

GraphPad Prism 8 software (GraphPad Software, Inc., USA) was used for

statistical analysis. A D'Agostino & Pearson normality test was used to evaluate normal

distribution, and outliers were identified using a Grubbs test. The data are expressed as

the mean☐±SEM or individual values. One-way ANOVA followed by Tukey's posttest

was used for comparisons between different inbred groups, whereas Student's t-test or a

nonparametric Mann-Whitney test was performed to compare the outbred groups.

Differences were considered significant when $p<0.05$. Pregnancy parameters were

evaluated using the mean value of all fetuses and placentae in a litter per dam and not

the individual conceptus, i.e., the mean value. In Figures 1 and 2, "n" represents the

number of dams. For MET and immunostaining data, placentae closest to the mean

weight of all placentae were selected from each litter; "n" represents the number of

litters (35–37).

3. Results

3.1 Weight gain during pregnancy is dependent on maternal immune status in ZIKV-

infected mice.

To determine the effect of ZIKV infection on fetal and placental phenotypes at

term (GD18.5), we infected ICompetent C57BL/6 and immunocompromised

(ICompromised) A129 mice with ZIKV at GD12.5 (Figure 1A). Given the very distinct

susceptibility of C57BL/6 and A129 mice to ZIKV, systemic infection models were

established by injecting high ($5x10^7$ PFU) and low (10^3 PFU) virus inoculum titers. As

shown in Figure 1B, ICompetent C57BL/6 mice in all groups and ICompromised A129

dams inoculated with mock and low ZIKV titers exhibited higher maternal weight at

GD18.5 than at GD12.5 and GD0.5 ($p<0.05$). On the other hand, ICompromised A129

mice presented significant weight loss at GD18.5, despite showing an increase at

GD12.5 in relation to GD0.5 (Figure 1B).

3.2 Immunocompetent and immunodeficient mice have distinct term placental and

fetal phenotypes in response to high and low ZIKV titer challenges in mid-pregnancy.

The fetuses from C57BL/6 dams and sires were called ICompetent. The fetuses

from the mating of A129 dams and C57BL/6 sires were called ICompromised. High-

ZIKV ICompetent mice exhibited 34% fetal loss, whereas high-ZIKV-A129 mice had

100% fetal loss (Figure 1C). In the low-ZIKV groups, C57BL/6 mice had no (0%) fetal

death, while A129 mice exhibited a 42% fetal death rate (Figure 1C). Fetal and fetal

head sizes were decreased in A129 mice compared to those in C57BL/6 dams infected

with the high ZIKV titer (p=0.05; Figure 1 D-G). However, no changes in fetal weight

or fetal head size were observed when the mice were infected with the low ZIKV titer

(Figure 1 D-G).

Figure 1: ZIKV infection induced fetal changes during pregnancy. A) Experimental
design: matings of C57BL/6 dams x C57BL/6 sires (mock n=11 dams; high ZIKV n=15
dams; low ZIKV n=9 dams) and A129 dams x C57BL/6 sires (n=3 dams/group). The
mock control consisted of noninfected C6/36 cell supernatants; high ZIKV titers
consisted of 5×10^7 plaque-forming units (PFU) of ZIKV$_{PE243}$, and low ZIKV titers
consisted of 10^3 PFU of ZIKV$_{PE243}$. Fetuses from C57BL/6 dams and sires were termed
ICompetent, whereas fetuses from A129 dams and C57BL/6 sires were termed
ICompromised. **B)** Maternal weight gain throughout pregnancy, *=p<0.05, one-way
ANOVA. **C)** Uterine horn and survival rates following ZIKV exposure (arrows show
resorption sites). **D)** Fetal/reabsorption images (#=example of fetal reabsorption). **E)**
Fetal weight, **F)** fetal head weight and **G)** fetal head weight/fetal weight ratio. One-way
ANOVA followed by Tukey's posttest was used to assess changes among ICompetent
groups, whereas an unpaired Student's t-test or nonparametric Mann-Whitney test was
used to assess differences between ICompromised groups. Values are the mean of
individual plotted values.

3.3 ZIKV is detected in the fetal brain of ICompromised, but not ICompetent mice.

ZIKV RNA was detected in the spleens of pregnant ICompetent C57BL/6 mice

inoculated with the highest ZIKV titer, confirming acute systemic infection. Viral RNA

was also detected in the majority of the placentae of those mice (Figure 2A-B) but not

in the maternal and fetal C57BL/6 brains (Figure 2A-B), suggesting that the virus was

not transmitted to the fetuses. Although low ZIKV inoculation resulted in virus

detection in the spleens of pregnant C57BL/6 mice, infection was not evidenced in the

placentae or fetal brains. In contrast, ZIKV RNA was detected in all analyzed organs

from ICompromised A129 dams, including the maternal brain and spleen and the

placenta and fetal brain (Figure 2A-B). Viremia in maternal plasma was evaluated at 4,

48 and 144 hours after infection. Within 4 hours, the presence of the virus was verified

in the serum (high ICompetent=637.5 PFU/mL, low ICompetent=740 PFU/mL and low

ICompromised=325 PFU/mL), indicating that the virus was correctly inoculated.

Afterwards, ZIKV RNA was detected in ICompromised dams at 48 and 144 hours post

inoculation but not in ICompetent dams (Figure 2C).

Figure 2: Viral load detection in maternal and fetal tissues after ZIKV infection. Pregnant ICompetent and ICompromised mice were inoculated i.v. with low and high doses of ZIKV. A-B) ZIKV RNA was measured via RT-qPCR in tissue samples obtained from maternal spleen and brain (A) and from placenta and fetal brain (B). C) The presence of ZIKV infectious particles in the plasma of infected dams was evaluated with a plaque assay 4, 48, and 144 hours post infection in ICompetent (mock n=11 dams; high ZIKV n=15 dams; low ZIKV n=9 dams) and ICompromised (mock n=3 dams; high ZIKV n=3 dams; low ZIKV n=3 dams) mice.

3.4 ZIKV infection induces distinct systemic and placental inflammatory responses in

ICompetent and ICompromised mice.

The maternal serum and placental protein levels of specific cytokines and chemokines

related to fetal death and preterm delivery (43–45) were evaluated to probe whether

midgestation ZIKV infection would induce a maternal inflammatory response at term in

our two distinct models. Since A129 infected with 10^7 ZIKV-PFU showed 100% fetal

loss, we proceeded using 10^7 PFU inoculation in C57BL/6 mice and 10^3 PFU

inoculation in both C57BL/6 and A129 mice.

CCL2 was elevated in the serum of low-ZIKV ICompetent mice, but no other

alteration was systemically detected in any ICompetent mice at this time point (Figure

3A-B). On the other hand, ICompromised dams showed significantly increased CXCL1

and IL-6 levels in the serum and a strong trend for an enhancement in CCL2 (Figure

3C). Analysis of cytokine and chemokine expression in the placenta demonstrated that

the CXCL1 and CCL2 chemokines were also upregulated in ICompromised but not

ICompetent mice (Figure 3D-F). Surprisingly, IL-6 protein expression was augmented

in some of the low-ZIKV ICompetent mice (58%; p<0.05) but not in high-ZIKV mice

or ICompromised mice (Figure 3D-F).

We also assessed the placental mRNA expression of a range of cytokines related

to placental infective responses: *Ifng, Il6, Ifn1, Tnf, Il1b and Il10*. All the infected mice

showed a significant increase in *Tnf* mRNA expression, with higher levels detected in

ICompetent mice (Figure 3G-I). The low-ZIKV ICompetent mice presented modest but

significant *Ifn1* expression, which was not detected in the high-ZIKV group (Figure

3G). Additionally, both low-ZIKV-infected groups (ICompetent and ICompromised)

presented increased *Ifng* mRNA expression (Figure 3G and 3I, p< 0.05). Interestingly,

placental *Il6* mRNA levels were only elevated in ICompromised pregnancies compared

to mock pregnancies (p=0.05) (Figure 3I). *Il1b* and *Il10* remained unchanged in all

groups analyzed (Figure 3 G-I).

Figure 3: ZIKV during pregnancy promotes an inflammatory response in the maternal serum and in the placenta. **A)** Levels of IL-1β, IL-6, CCL-2 and CXCL-1 in the maternal serum (A-C) and placenta (D-F) at GD18.5 in matings of C57BL/6 dams x C57BL/6 sires (ICompetent fetuses - mock n=11 dams; high ZIKV n=15 dams; low ZIKV n=9 dams) and A129 dams x C57BL/6 sires (ICompromised fetuses n=3 dams/group). Placental mRNA expression (G-I) of *Ifng, Ifn1, Il1b, Il6, Il10 and Tnf*. Broken lines show the expression levels in both lineages in the mock group. One-way ANOVA followed by Tukey's posttest was used to assess changes among ICompetent groups, whereas an unpaired Student's t-test or nonparametric Mann-Whitney test was

424   used to assess differences between ICompromised groups. The values are expressed as
425   the mean ± SEM.
426

3.5 ZIKV affects placental proliferation, apoptosis and ultrastructure in a viral load-

and maternal immune status-dependent manner.

We did not observe changes in placental weight or in the fetal:placental weight

ratio in any of the groups investigated (data not shown). However, since we detected the

presence of ZIKV RNA in the placentas, we investigated the cellular proliferation (Ki-

67^+ cells) and apoptotic ratio in the Lz and Jz of the mouse placenta. Increased Ki-67

staining was observed in the Lz of high- and low-ZIKV-treated ICompetent animals

compared to mock animals (Figure 4A-D, p=0.005 and p=0.015), whereas the apoptotic

ratio in the Lz was increased in high-ZIKV dams and decreased in low-ZIKV dams

(p=0.01, Figure 4E-H). Although no differences were observed in Ki67 staining (Figure

5A-D), a similar apoptotic pattern was detected in the Jz of ICompetent pregnancies

(p=0.008 and p=0.004, respectively; Figure 5E-H). In contrast, in ICompromised dams,

Lz Ki-67 staining was decreased (p=0.001; Figure 4I-L), while the apoptotic reaction

was increased in ZIKV-infected animals (p=0.05; Figure 4M-P). Jz from ICompromised

offspring exhibited increased Ki-67 labeling and no differences in the apoptotic reaction

(p=0.001; Figure 5 I-L and Figure 5M-P).

Figure 4: Labyrinthine remodeling in ICompetent and ICompromised placentae is affected by gestational ZIKV infection. A total of 180 digital images (40X) randomly captured from the whole labyrinth zone (Lz) of each placenta per dam were evaluated. Immunolabeled nuclei from each digital image were quantified and normalized by the total digital image area to obtain an index of the estimated number of proliferative and apoptotic nuclei in the entire histological section. (**A and I**) Quantification and (**B-D and J-K**) representative photomicrographs of Ki-67[+] stained nuclei in the Lz of ICompetent (n=6 placentae from 6 independent dams/group) and ICompromised (n=3 placentae from 3 independent dams/group) placentae, respectively. (**E and M**) Quantification and (**F-H and N-O**) representative photomicrographs of apoptotic nuclei (TUNEL) in the Lz of ICompetent (n=5 placentae from 5 independent dams/group) and ICompromised (n=3 placentae from 3 independent dams/group) placentae, respectively. (**L and P**) Negative controls. One-way ANOVA followed by Tukey's posttest was used to assess changes among ICompetent groups, whereas an unpaired Student's t-test or nonparametric Mann-Whitney test was used to assess differences between ICompromised groups. The values are expressed as the mean ± SEM. Images were captured at 40X. Scale bar=50 µm.

Figure 5: Junctional zone remodeling in ICompetent and ICompromised placentae is affected by gestational ZIKV infection. A total of 180 digital images (40X) randomly captured from the whole junctional zone (Jz) of each placenta per dam were evaluated. Immunolabeled nuclei from each digital image were quantified and normalized by the total digital image area to obtain an index of the estimated number of proliferative and apoptotic nuclei in the entire histological section. (**A and I**) Quantification and (**B-D and J-K**) representative photomicrographs of Ki-67[+] stained nuclei in the Jz of ICompetent (n=6 placentae from 6 independent dams/group) and ICompromised (n=3 placentae from 3 independent dams/group) placentae, respectively. (**E and M**) Quantification and (**F-H and N-O**) representative photomicrographs of apoptotic nuclei (TUNEL) in the Jz of ICompetent (n=5 placentae from 5 independent dams/group) and ICompromised (n=3 placentae from 3 independent dams/group) placentae, respectively. (**L and P**) Negative controls. One-way ANOVA followed by Tukey's posttest was used to assess changes among ICompetent groups, whereas an unpaired Student's t-test or nonparametric Mann-Whitney test was used to assess differences between ICompromised groups. The values are expressed as the mean ± SEM. Images were captured at 40X. Scale bar=50 μm.

3.6 Placental ultrastructure is differently impacted by high- and low-titer ZIKV

infection in ICompetent and ICompromised strains.

Lz ultrastructural analyses of ICompetent and ICompromised-mock animals

detected sinusoidal trophoblastic giant cells exhibiting regular microvilli, euchromatic

nuclei, preserved mitochondrial ultrastructure and regular narrow ER cisternae (Figure

6A and 6B). In sharp contrast, high-ZIKV ICompetent (Figure 6C) infected placentae

showed fewer villi in the sinusoidal giant trophoblastic cells, degenerated mitochondria,

granular ER with dilated cisterns and euchromatic nuclei. The sinusoidal giant

trophoblastic cells in the low-ZIKV ICompetent mice (Figure 6D) also had fewer villi

and degenerated mitochondria than those in the mock placentae, but no effect on the ER

or euchromatic nuclei observed. Low-ZIKV ICompromised infected placentae (Figure

6E) showed fewer villi in the sinusoidal giant trophoblastic cells, degenerated

mitochondria, granular ER with dilated cisterns and euchromatic nuclei.

Figure 6: Associated ultrastructural changes in the placental Lz after ZIKV
infection. Transmission electron photomicrographs of ICompetent mock (A),
ICompetent high (B), ICompetent low (C), ICompromised mock (D) and
ICompromised low (E) groups (n=5/group). We observed dilatation in the ER cisterns
of ICompetent high placentas. Additionally, there was a reduction in the microvilli in
both the ICompetent high and ICompetent low placentas. In the ICompromised low
group, we found fragmented ER and microvillus reduction. All infected groups showed
degenerate mitochondria. GER=granular endoplasmic reticulum; Δ=dilated granular
endoplasmic   reticulum;   *=fragmented   granular   endoplasmic   reticulum;

Mt=mitochondria; MtD=degenerate mitochondria; Mv=microvilli; EN=euchromatic nuclei; SS=sinusoidal space; TGC=trophoblastic giant cell. Scale bar=2 µm.

The Jz of mock ICompetent and ICompromised placentae (Figure 7A and 7B) exhibited euchromatic nuclei, with evident heterochromatin, preserved mitochondria and narrow cisternae in a granular ER. High-ZIKV ICompetent Jz had degenerated mitochondria, granular ER with dilated cisterns and euchromatic nuclei (Figure 7C). Low-ZIKV ICompetent (Figure 7D) placentae exhibited euchromatic nuclei with evident heterochromatin, preserved mitochondria and narrow cisternae in a granular ER, whereas low-ZIKV ICompromised placentae had degenerated mitochondria, granular ER with dilated cisterns and euchromatic nuclei (Figure 7E).

Figure 7: Associated ultrastructural changes in the placental Jz after ZIKV infection. Transmission electron photomicrographs of ICompetent mock (A), ICompetent high (B), ICompetent low (C), ICompromised mock (D) and ICompromised low (E) groups (n=5/group). We found deteriorating mitochondria and dilated reticulum endoplasmic cisterns in both the high ICompetent and low ICompromised groups. GER=granular endoplasmic reticulum; Δ=dilated granular endoplasmic reticulum; Mt=mitochondria; MtD=degenerate mitochondria;

Mv=microvilli; EN=euchromatic nuclei; SS=sinusoidal space; TGC=trophoblastic giant
cell; HA=heterochromatin area. Scale bar=2 µm.

3.7 ZIKV differentially affects placental expression of drug and lipid ABC

transporter systems.

Evaluation of key ABC transporters in the Lz of mock and ZIKV-infected

ICompetent and ICompromised placentae revealed that immunolabeling of the drug P-

gp and Bcrp efflux transporter systems was primarily present at the cellular membranes

of the sinusoidal trophoblastic giant cells, with diffuse cytoplasmic Bcrp staining.

Labeling of the Abca1 and Abcg1 lipid efflux transporters was moderately and

heterogeneously distributed within the Lz. Less Lz-P-gp was observed in ICompetent

mice infected with both high- and low-ZIKV infective regimens than in mock-treated

animals ($p=0.001$ and $p=0.002$, respectively; Figure 8A-E), whereas reduced Bcrp and

Abca1 staining was observed in high-ZIKV-infected mice ($p=0.003$ and $p=0.004$,

Figure 8F-J and Figure 8K-O, respectively). No changes in Abcg1 were observed in any

of the ICompetent experimental groups (Figure 8P-T). P-gp, Bcrp and Abca1

transporter immunostaining was downregulated in ICompromised low ZIKV-treated

animals ($p=0.001$, $p=0.05$ and $p=0.05$, Figure 9A-D, Figure 9E-H and Figure 9I-L,

respectively). No changes in Abcg1 were observed in any of the ICompromised

experimental groups (Figure 9M-P).

Figure 8: ZIKV infection decreases P-gp, Bcrp and Abca1 expression in the placental Lz of infected mice in the ICompetent groups. A total of 180 digital images (40X) randomly captured from the whole labyrinth zone (Lz) of each placenta per dam were evaluated. Immunolabeling in each digital image was quantified by calculating the percentage area of the total stained labyrinthine tissue after exclusion of the total negative space. (**A**) Quantification and (**B-D**) representative photomicrographs of P-gp staining in the Lz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**F**) Quantification and (**G-I**) representative photomicrographs of Bcrp staining in the Lz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**K**) Quantification and (**L-N**) representative photomicrographs of Abca1 staining in the Lz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**P**) Quantification and (**Q-S**) representative photomicrographs of Abcg1 staining in the Lz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**E, J, O, T**) Negative controls. One-way ANOVA followed by Tukey's post-test. The values are expressed as the mean ± SEM. Images were captured at 40X. Scale bar=50 µm.

Figure 9: ZIKV infection decreases P-gp, Bcrp and Abca1 protein expression in the placental Lz of infected mice in the ICompromised groups. A total of 180 digital images (40X) randomly captured from the whole labyrinth zone (Lz) of each placenta per dam were evaluated. Immunolabeling in each digital image was quantified by calculating the percentage area of the total stained labyrinth zone tissue after exclusion of the total negative space. (**A**) Quantification and (**B-C**) representative photomicrographs of P-gp staining in the Lz of ICompromised (n=3 placentae from 3 independent dams/group) placenta. (**E**) Quantification and (**F-G**) representative photomicrographs of Bcrp staining in the Lz of ICompromised (n=3 placentae from 3 independent dams/group) placenta. (**I**) Quantification and (**J-K**) representative photomicrographs of Abca1 staining in the Lz of ICompromised (n=3 placentae from 3 independent dams/group) placenta (**M**) quantification and (**N-O**) representative photomicrographs of Abcg1 staining in the Lz of ICompromised (n=3 placentae from 3 independent dams/group) placenta. (**D, H, L, P**) Negative controls. Unpaired Student's t-test or nonparametric Mann-Whitney test was used to assess differences between ICompromised groups. The values are expressed as the mean ± SEM. Images were captured at 40X. Scale bar=50 µm.

Next, the impact of ZIKV on ABC transporters in the Jz layer (structural and endocrine layers of the mouse placenta) was assessed. P-gp and Bcrp were predominantly localized at the cellular membranes of spongiotrophoblast cells, whereas Abca1 and Abcg1 exhibited membrane and cytoplasmic staining. P-gp staining was

582 decreased in Jz cells from the low-ZIKV ICompromised placentae (p=0.006), with no

583 other alterations observed (Figure 10A-T and Figure 11A-P).

Figure 10: ZIKV infection did not impact P-gp, Bcrp, Abca1 or Abcg1 protein expression in the placental Jz of infected mice in the ICompetent groups. A total of 180 digital images (40X) randomly captured from the whole junctional zone (Jz) of each placenta per dam were evaluated. Immunolabeling in each digital image was quantified by calculating the percentage area of the total stained junctional zone tissue after exclusion of the total negative space. (**A**) Quantification and (**B-D**) representative photomicrographs of P-gp staining in the Jz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**F**) Quantification and (**G-I**) representative photomicrographs of Bcrp staining in the Jz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**K**) Quantification and (**L-N**) representative photomicrographs of Abca1 staining in the Jz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**P**) Quantification and (**Q-S**) representative photomicrographs of Abcg1 staining in the Jz of ICompetent (n=6 placentae from 6 independent dams/group) placenta. (**E, J, O, T**) Negative controls. One-way ANOVA followed by Tukey's post-test. The values are expressed as the mean ± SEM. Images were captured at 40X. Scale bar=50 µm.

Figure 11: ZIKV infection decreases Bcrp protein expression in the placental Jz of infected mice in the ICompromised group. A total of 180 digital images (40X) randomly captured from the whole junctional zone (Jz) of each placenta per dam were evaluated. Immunolabeling in each digital image was quantified by calculating the percentage area of the total stained junctional zone tissue after exclusion of the total negative space. (**A**) Quantification and (**B-C**) representative photomicrographs of P-gp staining in the Jz of ICompromised (n=3 placentae from 3 independent dams/group) placenta. (**E**) Quantification and (**F-G**) representative photomicrographs of Bcrp staining in the Jz of ICompromised (n=3 placentae from 3 independent dams/group) placenta. (**I**) Quantification and (**J-K**) representative photomicrographs of Abca1 staining in the Jz of ICompromised (n=3 placentae from 3 independent dams/group) placenta. (**M**) Quantification and (**N-O**) representative photomicrographs of Abcg1 staining in the Jz of ICompromised (n=3 placentae from 3 independent dams/group) placenta. (**D, H, L, P**) Negative controls. Unpaired Student's t-test or nonparametric Mann-Whitney test was used to assess differences between ICompromised groups. The values are expressed as the mean ± SEM. Images were captured at 40X. Scale bar=50 µm.

4. Discussion

In this study, we investigated several fetal and placental features at term (GD18.5) in ICompetent (C57BL/6) and ICompromised (A129) mice exposed to ZIKV

at mid-pregnancy (GD12.5). Fetal survival rates, systemic and placental inflammatory

responses, placental ultrastructure and cell turnover, as well as the expression of key

drug (P-gp and Bcrp) and lipid (Abca1) efflux transporter systems in the placenta, were

consistently impacted by ZIKV in both strains. The magnitude of the effects was clearly

related to the infective titer (high and low) of ZIKV and maternal immune status

(ICompetent-C57BL/6 x and ICompromised-A129), and fetal alterations were not

exclusively dependent on virus detection in the fetuses.

Infection of ICompetent mice with ZIKV did not result in viremia in the initial

postinoculation phase, although viral RNA was detected in the maternal spleen in both

the high- and low-ZIKV groups, confirming systemic infection. This is consistent with a

previous report (46). Our data demonstrate that pregnant ICompetent C57BL/6 mice

were more susceptible to high ZIKV titers than to low ZIKV infective. Since viral RNA

was only detected in the placentae of high ZIKV-infected mice, fetal survival rates and

weights were impacted to a greater extent in those mice. Strikingly, even though the

virus was not present in the fetal brain (at least at term), fetal and fetal head weights

were lower in mice subjected to the high-ZIKV titer regimen, suggesting that high

infective viral load in mid-pregnancy, even in ICompetent individuals, can induce

IUGR and lower fetal head weight despite a lack of transmission to the fetal brain (47).

On the other hand, ICompromised placentae and fetal brains had detectable viral

transcripts, with no changes in weight, which is consistent with previous data (19). In

fact, in our models, the presence (ICompromised) or absence (ICompetent) of the virus

in the fetal brain did not correspond to fetal head size (decreased in only high

ICompetent). The data from ICompetent and ICompromised placentae demonstrate how

important the maternal immunological status is to control viremia, fetal survival and

accessibility of the virus to the fetal brain. The reason for the reduction in fetal brain

size in C57BL/6 mice in the absence of fetal brain infection requires further

investigation. It is possible that fetal brains in the ICompetent mice may have been

exposed to ZIKV earlier in pregnancy, when viremia was present in the maternal blood,

and this may have severely compromised brain development. Of note, one limitation of

our study is that we measured fetal head weight instead of cortical thickness. Future

studies should investigate whether high- and low-ZIKV exposure alters cortical

thickness in ICompetent and ICompromised offspring.

A distinct inflammatory profile was also detected in the three analyzed groups.

At the protein level, low-ZIKV-ICompromised dams exhibited increased maternal IL-6

and CXCL-1 and placental CCL-2 and CXCL-1, whereas low-ZIKV-ICompetent dams

had increased maternal CCL2 and placental IL-6 levels. CCCL-2 and CXCL-1 are

related to fetal death and preterm delivery (43–45) and could be associated with

pronounced fetal injury detected upon ICompromised pregnancy. In addition, IL-6 was

previously demonstrated (48) to be related to fetal response syndrome, characterized by

activation of the fetal immune system. This syndrome is known to increase fetal

morbidity and affect several organs, such as the adrenal gland, brain and heart (32,48–

51). At the mRNA level, *Il6* expression was only detected in ICompromised placentae

at term and may indicate a sustained harmful response in these mice until term. The IFN

signaling pathway may be triggered by ZIKV (17) and is one of the key mechanisms of

host defense and a viral target for immune evasion (20), but we only detected a slight

increase in *Ifn1* in low-ZIKV ICompetent mice at term. However, we cannot rule out

the possibility that these cytokines might have been produced earlier. Our findings

showed that *Ifng* expression was significantly enhanced in both ICompetent and

ICompromised low-ZIKV-derived placentae but not in high-ZIKV-infected mice.

Although we could not assess cytokine expression in high-ZIKV ICompromised

placentas, one may extrapolate that low-ZIKV infection could result in stimulation of

Ifng producing cells, which has been previously shown to be protective for ZIKV-

infected mice (52).

Both ICompromised and ICompetent mice showed increased expression of

placental *Tnf* mRNA, which has been demonstrated to be directly related to placental

damage, abortion and premature birth (53–56). In addition, an increased *Tnf* response is

related to impaired placental hormone production and trophoblastic invasion and

increased apoptosis in pregnancy (57,58). Although we did not assess TNF-α protein

levels in the placenta and maternal blood, this response could be implicated in the

overall damage detected.

Although differences in placental weight were not observed, ZIKV infection

mid-pregnancy had a profound effect on placental cellular turnover, dependent on titer,

strain and/or placental compartment. The Lz is responsible for fetal and maternal

nutrient, gas and waste exchange, while the Jz provides structural support, nutrient

storage and hormone synthesis (35). ZIKV induced a consistent increase in Lz

proliferation in all groups. However, the Lz apoptotic rate was increased only in the

high-ZIKV-ICompetent and ICompromised groups and decreased in low-ZIKV-

ICompetent mice. The mechanisms underlying these differences are unknown but may

be related to the distinct maternal and placental proinflammatory responses and/or to the

direct effect of the virus on the placenta (59). Increased Lz apoptosis in the high-

ICompetent-ZIKV group may be one of the mechanisms driving the lower fetal and

fetal head weight detected in this group. In this context, changes in placental turnover

can determine placental maturation and function and lead to fetal distress and

developmental abnormalities (60). An increase in the Lz apoptotic ratio may signify

damage to this placental layer, which is consistent with the fact that diverse pathological

lesions associated with congenital disorders were described in placentae from women

infected by ZIKV at different stages of pregnancy (61). Conversely, no proliferative

changes were observed in the Jz in high-ZIKV and low-ZIKV ICompetent mice, while

increased and decreased apoptotic rates were detected. It follows that the lack of Jz-Ki-

67 induction may suggest that this layer is less capable of restoring proliferation in

response to high-ZIKV challenge, and this may be related to decreased fetal growth.

Our placental ultrastructural analysis detected consistent differences across

ZIKV-exposed groups. The Lz and Jz layers from both strains exhibited signs of ER

stress, i.e., dilated ER cisterns or fragmented ER granular structures. These alterations

may result from the accumulation of folded or poorly folded viral proteins in the ER

lumen (42,62,63). The *Flaviridae* family uses the ER to replicate (64), and according to

Offerdahl et al. (2017)(63), there is evidence of ZIKV interacting with this organelle,

promoting an increased release of Ca^{+2} from the ER to the cell cytoplasm, thereby

causing an increase in the production of reactive oxygen species (ROS) (65,66).

The mitochondrial ultrastructure in the Lz and Jz layers was severely impacted

by ZIKV exposure. We found evidence of mitochondrial degeneration, i.e.,

mitochondrial membrane rupture, absence of mitochondrial ridges and a less electron-

dense mitochondrial matrix, in all the treated groups. Placental mitochondrial

dysfunction is associated with IUGR (67,68) and may be related, at least in part, to the

lower fetal weight observed in high-ZIKV-ICompetent fetuses along with the altered

placental apoptotic and proliferative patterns. Furthermore, mitochondrial dysfunction

together with ER stress is likely to modify the placental ROS balance and generate local

oxidative stress (69), which is associated with impaired fetal development (70). Of

importance, associations between mitochondrial disruption, ER stress and placental cell

senescence have been reported. Senescence is characterized as an irreversible

interruption of the cell cycle and acquisition of a senescence-associated secretory

phenotype (SASP) that promotes the release of cytokines, such as IL-1, IL-6, IL-8 and

proinflammatory proteases (70). Therefore, the increased expression of IL-6 detected in

the placentas of ICompromised mice suggests a SASP profile, which may be related to

changes in the ER and mitochondrial ultrastructure, accompanied by important changes

in apoptosis and cell proliferation. The interactions between mitochondria and the ER

are critical for homeostasis and cell signaling (71). In conjunction with the ER,

mitochondria can regulate cell death mediators in response to hypoxia and inflammation

(72). The increase in apoptosis observed in the high-titer ICompetent groups and the

low-titer ICompromised group may be related to the mitochondrial damage and ER

stress observed. In fact, we observed an important decrease in microvillus abundance in

sinusoidal giant trophoblast cells. Previously, we observed a decrease in microvillus

density in the Lz of pregnancies exposed to malaria in pregnancy (MiP) (36). Together,

our data show that different gestational infective stimuli (MiP and ZIKV) are capable of

damaging placental microvillus abundance and impairing proper fetal-maternal

exchange function and fetal growth/survival.

Next, to investigate whether maternal ZIKV exposure may influence fetal

protection, we evaluated the placental localization and expression (semiquantitative) of

the ABC efflux transporter systems P-gp, Bcrp, Abca1 and Abcg1, which are highly

enriched in labyrinthine microvilli and in human syncytiotrophoblasts. These efflux

transporters exchange drugs, environmental toxins, cytotoxic oxysterols and lipids

within the maternal-fetal interface (26). We found a consistent decrease in labyrinthine

P-gp expression in all ZIKV-exposed groups, demonstrating that ZIKV infection during

pregnancy has the potential to increase fetal exposure to P-gp substrates, such as

synthetic glucocorticoids, antibiotics, antiretrovirals, antifungals, stomach-protective

drugs, and nonsteroidal anti-inflammatory drugs (26). Furthermore, Jz-P-gp was

decreased in ICompromised placentae. Although little is known about the function of

ABC transporters in the Jz, our data highlight the need for further studies investigating

the biological importance of ABC transporters in the placental endocrine and structural

zones of the rodent hemochorial placenta under normal and infective conditions.

ZIKV impaired Lz Bcrp and Abca1 expression in ICompetent (high) and

ICompromised (low) mice. However, no effects were observed in ICompetent animals

at a low ZIKV titer or in Abcg1 in any experimental setting. Thus, ZIKV also likely

increases fetal accumulation of Bcrp substrates (antibiotics, antiretrovirals,

sulfonylureas, folate, mercuric species, estrogenic mycotoxins, carcinogens and

phototoxic compounds, among others) and disrupts placental lipid homeostasis (lipids,

cholesterol, and cytotoxic oxysterols) by reducing placental Abca1 expression (26,73–

76). We can speculate that the increased fetal accumulation of the P-gp, Bcrp and

Abca1 substrates during ZIKV infection may contribute to the establishment of

congenital Zika syndrome, although additional studies are clearly required to answer

this important question. The present data are in agreement with previous publications

showing that bacterial, viral and protozoan inflammation alters the expression and/or

function of P-gp, Bcrp and Abca1 in biological barriers, such as the placenta, yolk sac

and blood-brain barriers (26,27,36,64,77–79).

5. Conclusion

Our data show that gestational ZIKV impacts the fetal phenotype independently

of term fetal viremia. Abnormal placental cell turnover, ultrastructure and transporter

expression may result from specific proinflammatory responses that depend on the

ZIKV infective load and maternal immune status. Fetal accumulation of drugs,

environmental toxins and lipids within the fetal compartment may potentially be

increased in ZIKV-infected pregnancies due to altered levels of key ABC transporters.

Conflict of Interest:

778 The authors declare that the research was conducted in the absence of any commercial

779 or financial relationships that could be construed as a potential conflict of interest.

Author contributions

781 CBVA, FFB, EB, LBA and TMOC conceived and designed the experiments. CBVA,

782 VRSM, SVAC, HRG, RPCS and VMON performed the experiments. CBVA, SVAC,

783 FFB, EB, SGM, LBA and TMOC analyzed the data. CBVA, VRSM, EB, LBA and

784 TMOC wrote the paper and edited the manuscript. All authors contributed to the article

785 and approved the submitted version.

Funding

This study was supported by the Bill & Melinda Gates Foundation

(MCTI/CNPq/MS/SCTIE/Decit/Bill and Melinda Gates 05/2013; OPP1107597), the

Canadian Institutes for Health Research (SGM: Foundation-148368), Conselho

Nacional de Desenvolvimento Científico e Tecnológico (CNPq; 304667/2016□1,

422441/2016□3, 303734/2012□4, 422410/2016□0), Coordenação de Aperfeiçoamento

Pessoal de Nível Superior (CAPES, finance Code 001), Fundação de Amparo à

Pesquisa do Estado do Rio de Janeiro (FAPERJ, CNE 2015/E26/203.190/2015, PDR-

26/2002/010/2016), and PRPq-Universidade MG Federal MGPEFARJ, CNE

2015/E26/203.190/2015).

Acknowledgments

We would like to thank Alan Moraes for supporting the acquisition of electron

microscopy images and the electron microscopy laboratory at the UFF Biology Institute

for allowing the use of the JEM1011 transmission electron microscope. We would also

like to thank Mauro Jorge Castro Cabral for the use of the MAGPIX® System

equipment at the Paulo de Góes Institute of Microbiology/UFRJ; Ernesto T. Marques Jr.

(Centro de Pesquisa Aggeu Magalhães, FIOCRUZ, PE) for providing ZIKV to the

Institute of Microbiology Paulo de Góes, Federal University of Rio de Janeiro; and the

technicians Juliana Gonçalves and Rakel Alves for their support during all the

experiments.

References
807

1.   Garcez PP, Loiola EC, Madeiro da Costa R, Higa LM, Trindade P, Delvecchio R, et
al. Zika virus impairs growth in human neurospheres and brain organoids. Science.
2016;352(6287):816–8.
2.   Brasil P, Calvet GA, Siqueira AM, Wakimoto M, de Sequeira PC, Nobre A, et al.
Zika Virus Outbreak in Rio de Janeiro, Brazil: Clinical Characterization, Epidemiological
and Virological Aspects. Powers AM, editor. PLoS Negl Trop Dis. 2016;10(4):e0004636.
3.   Brasil P, Nielsen-Saines K. More pieces to the microcephaly–Zika virus puzzle in
Brazil. The Lancet Infectious Diseases. 2016;16(12):1307–9.
4.   Brasil P, Pereira JP, Moreira ME, Ribeiro Nogueira RM, Damasceno L, Wakimoto
M, et al. Zika Virus Infection in Pregnant Women in Rio de Janeiro. N Engl J Med. 2016
Mar 4;375(24):2321–34.
5.   Jaenisch T, Rosenberger KD, Brito C, Brady O, Brasil P, Marques ET. Risk of
microcephaly after Zika virus infection in Brazil, 2015 to 2016. Bull World Health
Organ. 2017;95(3):191–8.
6.   Proenca-Modena JL, Milanez GP, Costa ML, Judice CC, Maranhão Costa FT. Zika
virus: lessons learned in Brazil. Microbes and Infection. 2018;20(11–12):661–9.
7.   Jouannic J-M, Friszer S, Leparc-Goffart I, Garel C, Eyrolle-Guignot D. Zika virus
infection in French Polynesia. The Lancet. 2016;387(10023):1051–2.
8.   Musso D, Ko AI, Baud D. Zika Virus Infection — After the Pandemic. Longo DL,
editor. N Engl J Med. 2019;381(15):1444–57.
9.   Kasprzykowski JI, Fukutani KF, Fabio H, Fukutani ER, Costa LC, Andrade BB, et al.
A recursive sub-typing screening surveillance system detects the appearance of the
ZIKV African lineage in Brazil: Is there a risk of a new epidemic? International Journal of
Infectious Diseases. 2020;96:579–81.
10.   Miranda J, Martín-Tapia D, Valdespino-Vázquez Y, Alarcón L, Espejel-Nuñez A,
Guzmán-Huerta M, et al. Syncytiotrophoblast of Placentae from Women with Zika
Virus Infection Has Altered Tight Junction Protein Expression and Increased
Paracellular Permeability. Cells. 2019;8(10):1174.
11.   Jurado KA, Simoni MK, Tang Z, Uraki R, Hwang J, Householder S, et al. Zika virus
productively infects primary human placenta-specific macrophages. JCI Insight.
2016;1(13):e88461.

12.   Quicke KM, Bowen JR, Johnson EL, McDonald CE, Ma H, O'Neal JT, et al. Zika
Virus Infects Human Placental Macrophages. Cell Host & Microbe. 2016;20(1):83–90.

13.   Tabata T, Petitt M, Puerta-Guardo H, Michlmayr D, Wang C, Fang-Hoover J, et
al. Zika Virus Targets Different Primary Human Placental Cells, Suggesting Two Routes
for Vertical Transmission. Cell Host & Microbe. 2016;20(2):155–66.

14.   Simoni MK, Jurado KA, Abrahams VM, Fikrig E, Guller S. Zika virus infection of
Hofbauer cells. Am J Reprod Immunol. 2017;77(2):e12613.

15.   Rathore APS, Saron WAA, Lim T, Jahan N, John ALS. Maternal immunity and
antibodies to dengue virus promote infection and Zika virus–induced microcephaly in
fetuses. Science Advances. 2019; 5(2):eaav3208.

16.   Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, et al. A Mouse
Model of Zika Virus Pathogenesis. Cell Host & Microbe. 2016;19(5):720–30.

17.   Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, et al.
Zika Virus Targets Human STAT2 to Inhibit Type I Interferon Signaling. Cell Host &
Microbe. 2016;19(6):882–90.

18.   Yockey LJ, Varela L, Rakib T, Khoury-Hanold W, Fink SL, Stutz B, et al. Vaginal
Exposure to Zika Virus during Pregnancy Leads to Fetal Brain Infection. Cell.
2016;166(5):1247-1256.e4.

19.   Garcez PP, Stolp HB, Sravanam S, Christoff RR, Ferreira JCCG, Dias AA, et al. Zika
virus impairs the development of blood vessels in a mouse model of congenital
infection. Sci Rep. 2018;8(1):12774.

20.   Elong Ngono A, Shresta S. Immune Response to Dengue and Zika. Annu Rev
Immunol. 2018;36(1):279–308.

21.   Bayer A, Lennemann NJ, Ouyang Y, Bramley JC, Morosky S, Marques ETDAJ, et
al. Type III Interferons Produced by Human Placental Trophoblasts Confer Protection
against Zika Virus Infection. Cell Host Microbe. 2016;19(5):705–12.

22.   Luo H, Winkelmann ER, Fernandez-Salas I, Li L, Mayer SV, Danis-Lozano R, et al.
Zika, dengue and yellow fever viruses induce differential anti-viral immune responses
in human monocytic and first trimester trophoblast cells. Antiviral Research.
2018;151:55–62.

23.   Tanaka T, Narazaki M, Kishimoto T. IL-6 in Inflammation, Immunity, and
Disease. Cold Spring Harb Perspect Biol. 2014;6(10):a016295.

24.   Hunter CA, Jones SA. IL-6 as a keystone cytokine in health and disease. Nature
Immunology. 2015;16(5):448–57.

25.   Rabelo K, Gonçalves AJ da S, Souza LJ de, Sales AP, Lima SMB de, Trindade GF,
et al. Zika Virus Infects Human Placental Mast Cells and the HMC-1 Cell Line, and
Triggers Degranulation, Cytokine Release and Ultrastructural Changes. Cells.
2020;9(4):975–90.

26.   Bloise E, Ortiga-Carvalho TM, Reis FM, Lye SJ, Gibb W, Matthews SG. ATP-
binding cassette transporters in reproduction: a new frontier. Hum Reprod Update.
2015;22(2):164–81.

27.   do Imperio GE, Bloise E, Javam M, Lye P, Constantinof A, Dunk C, et al.
Chorioamnionitis Induces a Specific Signature of Placental ABC Transporters Associated
with an Increase of miR-331-5p in the Human Preterm Placenta. Cell Physiol Biochem.
2018;45(2):591–604.

28.   Martinelli LM, Fontes KN, Reginatto MW, Andrade CBV, Monteiro VRS, Gomes
HR, et al. Malaria in pregnancy regulates P-glycoprotein (P-gp/Abcb1a) and ABCA1
efflux transporters in the mouse visceral yolk sac. J Cell Mol Med. 2020;00:1–12.
29.   Reginatto MW, Fontes KN, Monteiro VRS, Silva NL, Andrade CBV, Gomes HR, et
al. Effect of sub-lethal prenatal endotoxemia on murine placental transport systems
and lipid homeostasis. bioRxiv. 2020 Aug 5;2020.08.04.236745.
30.   Martinelli LM, Reginatto MW, Fontes KN, Andrade CBV, Monteiro VRS, Gomes
HR, et al. Breast cancer resistance protein (Bcrp/Abcg2) is selectively modulated by
lipopolysaccharide (LPS) in the mouse yolk sac. Reprod Toxicol. 2020; 98:82-91.
31.   Imperio GE, Javam M, Lye P, Constantinof A, Dunk CE, Reis FM, et al.
Gestational age-dependent gene expression profiling of ATP-binding cassette
transporters in the healthy human placenta. J Cell Mol Med. 2019;23(1):610–8.
32.   Vermillion MS, Lei J, Shabi Y, Baxter VK, Crilly NP, McLane M, et al. Intrauterine
Zika virus infection of pregnant immunocompetent mice models transplacental
transmission and adverse perinatal outcomes. Nat Commun. 2017;8(1):14575.
33.   Szaba FM, Tighe M, Kummer LW, Lanzer KG, Ward JM, Lanthier P, et al. Zika
virus infection in immunocompetent pregnant mice causes fetal damage and placental
pathology in the absence of fetal infection. Coyne CB, editor. PLoS Pathog.
2018;14(4):e1006994.
34.   Coelho SVA, Neris RLS, Papa MP, Schnellrath LC, Meuren LM, Tschoeke DA, et
al. Development of standard methods for Zika virus propagation, titration, and
purification. Journal of Virological Methods. 2017;246:65–74.
35.   Connor KL, Kibschull M, Matysiak-Zablocki E, Nguyen TT-TN, Matthews SG, Lye
SJ, et al. Maternal malnutrition impacts placental morphology and transporter
expression: an origin for poor offspring growth. The Journal of Nutritional
Biochemistry. 2020;78:108329.
36.   Fontes KN, Reginatto MW, Silva NL, Andrade CBV, Bloise FF, Monteiro VRS, et
al. Dysregulation of placental ABC transporters in a murine model of malaria-induced
preterm labor. Sci Rep. 2019;9(1):11488.
37.   Bloise E, Lin W, Liu X, Simbulan R, Kolahi KS, Petraglia F, et al. Impaired
Placental Nutrient Transport in Mice Generated by *in Vitro* Fertilization. Endocrinology.
2012;153(7):3457–67.
38.   Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic
and Serologic Properties of Zika Virus Associated with an Epidemic, Yap State,
Micronesia, 2007. Emerg Infect Dis. 2008;14(8):1232–9.
39.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. Methods. 2001;25(4):402–8.
40.   Tschanz SA, Burri PH, Weibel ER. A simple tool for stereological assessment of
digital images: the STEPanizer: TOOL FOR STEREOLOGICAL ASSESSMENT. Journal of
Microscopy. 2011;243(1):47–59.
41.   Sesso A, Belizário JE, Marques MM, Higuchi ML, Schumacher RI, Colquhoun A,
et al. Mitochondrial Swelling and Incipient Outer Membrane Rupture in Preapoptotic
and Apoptotic Cells. Anatomical Record. 2012;295(10):1647.
42.   Montalbano R, Waldegger P, Quint K, Jabari S, Neureiter D, Illig R, et al.
Endoplasmic Reticulum Stress Plays a Pivotal Role in Cell Death Mediated by the Pan-
Deacetylase Inhibitor Panobinostat in Human Hepatocellular Cancer Cells.
Translational Oncology. 2013;6(2):143-IN6.

43.   Hamilton SA, Tower CL, Jones RL. Identification of Chemokines Associated with
the Recruitment of Decidual Leukocytes in Human Labour: Potential Novel Targets for
Preterm Labour. PLOS ONE. 2013;8(2):e56946.
44.   Bloise E, Bhuiyan M, Audette MC, Petropoulos S, Javam M, Gibb W, et al.
Prenatal Endotoxemia and Placental Drug Transport in The Mouse: Placental Size-
Specific Effects. PLOS ONE. 2013;8(6):e65728.
45.   Romero R, Mazor M, Brandt F, Sepulveda W, Avila C, Cotton DB, et al.
Interleukin-1α and Interleukin-1 β in Preterm and Term Human Parturition. American
Journal of Reproductive Immunology. 1992;27(3-4):117–23.
46.   Larocca RA, Abbink P, Peron JPS, de A. Zanotto PM, Iampietro MJ, Badamchi-
Zadeh A, et al. Vaccine protection against Zika virus from Brazil. Nature.
2016;536(7617):474–8.
47.   Souza IN de O, Frost PS, França JV, Nascimento-Viana JB, Neris RLS, Freitas L, et
al. Acute and chronic neurological consequences of early-life Zika virus infection in
mice. Science Translational Medicine. 2018;10(444):eaar2749.
48.   Kalagiri RR, Carder T, Choudhury S, Vora N, Ballard AR, Govande V, et al.
Inflammation in Complicated Pregnancy and Its Outcome. Am J Perinatol.
2016;33(14):1337–56.
49.   Barbeito-Andrés J, Pezzuto P, Higa LM, Dias AA, Vasconcelos JM, Santos TMP,
et al. Congenital Zika syndrome is associated with maternal protein malnutrition. Sci
Adv. 2020;6(2):eaaw6284.
50.   Saito S, Nakashima A, Shima T, Ito M. REVIEW ARTICLE: Th1/Th2/Th17 and
Regulatory T-Cell Paradigm in Pregnancy. American Journal of Reproductive
Immunology. 2010;63(6):601–10.
51.   Gotsch F, Romero R, Kusanovic JP, Mazaki-Tovi S, Pineles BL, Erez O, et al. The
fetal inflammatory response syndrome. Clin Obstet Gynecol. 2007;50(3):652–83.
52.   Lucas CGO, Kitoko JZ, Ferreira FM, Suzart VG, Papa MP, Coelho SVA, et al.
Critical role of CD4 + T cells and IFNγ signaling in antibody-mediated resistance to Zika
virus infection. Nature Communications. 2018;9(1):3136.
53.   Burdet J, Sacerdoti F, Cella M, Franchi AM, Ibarra C. Role of TNF-α in the
mechanisms responsible for preterm delivery induced by Stx2 in rats. British Journal of
Pharmacology. 2013;168(4):946.
54.   Carpentier PA, Dingman AL, Palmer TD. Placental TNF-α Signaling in Illness-
Induced Complications of Pregnancy. The American Journal of Pathology.
2011;178(6):2802–10.
55.   Romanowska-Próchnicka K, Felis-Giemza A, Olesińska M, Wojdasiewicz P,
Paradowska-Gorycka A, Szukiewicz D. The Role of TNF-α and Anti-TNF-α Agents during
Preconception, Pregnancy, and Breastfeeding. International Journal of Molecular
Sciences. 2021;22(6): 2922.
56.   Sarr D, Bracken TC, Owino SO, Cooper CA, Smith GM, Nagy T, et al. Differential
roles of inflammation and apoptosis in initiation of mid-gestational abortion in
malaria-infected C57BL/6 and A/J mice. Placenta. 2015;36(7):738.
57.   Galinsky R, Polglase GR, Hooper SB, Black MJ, Moss TJM. The consequences of
chorioamnionitis: preterm birth and effects on development. J Pregnancy.
2013;2013:412831.
58.   Mor G, Cardenas I, Abrahams V, Guller S. Inflammation and pregnancy: the role
of the immune system at the implantation site. Ann N Y Acad Sci. 2011;1221(1):80–7.

59. Ribeiro MR, Moreli JB, Marques RE, Papa MP, Meuren LM, Rahal P, et al. Zika-
virus-infected human full-term placental explants display pro-inflammatory responses
and undergo apoptosis. Arch Virol. 2018;163(10):2687–99.

60. Gown AM, Willingham MC. Improved Detection of Apoptotic Cells in Archival
Paraffin Sections: Immunohistochemistry Using Antibodies to Cleaved Caspase 3. J
Histochem Cytochem. 2002;50(4):449–54.

61. Noronha L de, Zanluca C, Burger M, Suzukawa AA, Azevedo M, Rebutini PZ, et
al. Zika Virus Infection at Different Pregnancy Stages: Anatomopathological Findings,
Target Cells and Viral Persistence in Placental Tissues. Front Microbiol.
2018;9(2266):1–11.

62. Lojpur T, Easton Z, Raez-Villanueva S, Laviolette S, Holloway AC, Hardy DB. Δ9-
Tetrahydrocannabinol leads to endoplasmic reticulum stress and mitochondrial
dysfunction in human BeWo trophoblasts. Reproductive Toxicology. 2019;87:21–31.

63. Offerdahl DK, Dorward DW, Hansen BT, Bloom ME. Cytoarchitecture of Zika
virus infection in human neuroblastoma and Aedes albopictus cell lines. Virology.
2017;501:54–62.

64. Desuzinges-Mandon E, Arnaud O, Martinez L, Huché F, Di Pietro A, Falson P.
ABCG2 Transports and Transfers Heme to Albumin through Its Large Extracellular Loop.
J Biol Chem. 2010;285(43):33123–33.

65. Ledur PF, Karmirian K, Pedrosa C da SG, Souza LRQ, Assis-de-Lemos G, Martins
TM, et al. Zika virus infection leads to mitochondrial failure, oxidative stress and DNA
damage in human iPSC-derived astrocytes. Scientific Reports. 2020;10(1):1218.

66. Reemst K, Noctor SC, Lucassen PJ, Hol EM. The Indispensable Roles of Microglia
and Astrocytes during Brain Development. Front Hum Neurosci. 2016;10:566.

67. Guitart-Mampel M, Gonzalez-Tendero A, Niñerola S, Morén C, Catalán-Garcia
M, González-Casacuberta I, et al. Cardiac and placental mitochondrial characterization
in a rabbit model of intrauterine growth restriction. Biochimica et Biophysica Acta
(BBA) - General Subjects. 2018;1862(5):1157–67.

68. Guitart-Mampel M, Juarez-Flores DL, Youssef L, Moren C, Garcia-Otero L, Roca-
Agujetas V, et al. Mitochondrial implications in human pregnancies with intrauterine
growth restriction and associated cardiac remodelling. J Cell Mol Med.
2019;23(6):3962–73.

69. Zhang Z, Rong L, Li Y-P. Flaviviridae Viruses and Oxidative Stress: Implications
for Viral Pathogenesis. Oxid Med Cell Longev. 2019;2019:1409582.

70. Martínez F, Kiriakidou M, Strauss JF. Structural and Functional Changes in
Mitochondria Associated with Trophoblast Differentiation: Methods to Isolate
Enriched Preparations of Syncytiotrophoblast Mitochondria. Endocrinology.
1997;138(5):2172–83.

71. Burton GJ, Yung HW, Murray AJ. Mitochondrial – Endoplasmic reticulum
interactions in the trophoblast: Stress and senescence. Placenta. 2017;52:146–55.

72. Holland O, Dekker Nitert M, Gallo LA, Vejzovic M, Fisher JJ, Perkins AV. Review:
Placental mitochondrial function and structure in gestational disorders. Placenta.
2017;54:2–9.

73. Bridges CC, Zalups RK, Joshee L. Toxicological significance of renal Bcrp:
Another potential transporter in the elimination of mercuric ions from proximal
tubular cells. Toxicol Appl Pharmacol. 2015;285(2):110–7.

74.   Szilagyi JT, Gorczyca L, Brinker A, Buckley B, Laskin JD, Aleksunes LM. Placental
BCRP/ABCG2 Transporter Prevents Fetal Exposure to the Estrogenic Mycotoxin
Zearalenone. Toxicol Sci. 2019;168(2):394–404.
75.   Scialis RJ, Aleksunes LM, Csanaky IL, Klaassen CD, Manautou JE. Identification
and Characterization of Efflux Transporters That Modulate the Subtoxic Disposition of
Diclofenac and Its Metabolites. Drug Metab Dispos. 2019;47(10):1080–92.
76.   Mao Q, Unadkat JD. Role of the breast cancer resistance protein (BCRP/ABCG2)
in drug transport--an update. AAPS J. 2015;17(1):65–82.
77.   Lye P, Bloise E, Javam M, Gibb W, Lye SJ, Matthews SG. Impact of Bacterial and
Viral Challenge on Multidrug Resistance in First- and Third-Trimester Human Placenta.
The American Journal of Pathology. 2015;185(6):1666–75.
78.   Girard S, Sebire G. Transplacental Transfer of Interleukin-1 Receptor Agonist
and Antagonist Following Maternal Immune Activation. American journal of
reproductive immunology (New York, NY : 1989). 2016;75(1):8–12.
79.   Bloise E, Petropoulos S, Iqbal M, Kostaki A, Ortiga-Carvalho TM, Gibb W, et al.
Acute Effects of Viral Exposure on P-Glycoprotein Function in the Mouse Fetal Blood-
Brain Barrier. Cellular physiology and biochemistry : international journal of
experimental cellular physiology, biochemistry, and pharmacology. 2017;41(3):1044–
50.

1043