1   **ZIKV disrupts fetal outcome, placental ultrastructure and drug transporter**

2   **expression in mice**

Cherley Borba Vieira de Andrade[1], Victoria Regina de Siqueira Monteiro[1], Sharton

Vinicius Antunes Coelho[2], Hanailly Ribeiro Gomes[1], Ronny Paiva Campos de Sousa[1],

Veronica Muller de Oliveira Nascimento[1], Flavia Fonseca Bloise[1], Stephen

Matthews[3,4,5,6], Enrrico Bloise[7], Luciana Barros de Arruda[2], Tania Maria Ortiga-

Carvalho[1]*.

[1]Institute of Biophysics Carlos Chagas Filho, Federal University of Rio de Janeiro, Rio

de Janeiro, RJ, Brazil.

[2] Institute of Microbiology Paulo de Góes, Federal University of Rio de Janeiro, Rio de

Janeiro, RJ, Brazil.

[3] Department of Physiology, Faculty of Medicine, University of Toronto, Toronto,

Canada.

[4] Department of Obstetrics & Gynecology, Faculty of Medicine, University of Toronto,

Toronto, Canada.

[5] Department of Medicine, Faculty of Medicine, University of Toronto, Toronto,

Canada.

[6] Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Toronto, Canada.

[7] Department of Morphology, Federal University of Minas Gerais, Belo Horizonte, BH,

Brazil.

* Correspondence:

Tania Maria Ortiga-Carvalho (TMO-C)

taniaort@biof.ufrj.br

Key words: ZIKV, placenta, P-glycoprotein (P-gp) breast cancer resistance protein

(BCRP), Abca1, Abcg1, ultrastructure, cytokine, chemokine

Abstract

Congenital Zika virus (ZIKV) infection can induce fetal brain abnormalities. Here, we investigated whether maternal ZIKV infection may affect placental physiology and metabolic transport potential, and impact the fetal outcome, regardless of viral presence in the fetus at term. Low (103 PFU-ZIKVPE243; low-ZIKV) and high ($5x10^7$ PFU-ZIKVPE243; high-ZIKV) virus titers were injected into immunocompetent (ICompetent C57BL/6) and immunocompromised (ICompromised A129) mice at gestational day (GD)12.5 for tissue collection at GD18.5 (term). High-ZIKV elicited fetal death rates of 66% and 100%, whereas low-ZIKV induced fetal death rates of 0% and 60% in C57BL/6 and A129 dams, respectively. All surviving fetuses exhibited intrauterine growth restriction (IUGR) and decreased placental efficiency. High-ZIKV infection in C57BL/6 and A129 mice resulted in virus detection in maternal spleens and placenta, but only A129 fetuses presented virus RNA in the brains. Nevertheless, pregnancies of both strains produced fetuses with decreased head sizes ($p<0.05$). Low-ZIKV-A129 dams had higher IL-6 and CXCL1 levels ($p<0.05$) and their placentas showed increased CCL-2 and CXCL-1 contents ($p<0.05$). In contrast, low-ZIKV-C57BL/6 dams had an elevated CCL2 serum level, and increased type I and II IFN expression in the placenta. Notably, increased apoptotic rates, less abundant microvilli and mitochondrial degeneration were evidenced in the placental labyrinth (Lz) of ICompromised and high-ZIKV-ICompetent mice, but not in low-ZIKV-C57BL/6. In addition, decreased placental expression of the drug P-glycoprotein (P-gp), breast cancer resistance protein (BCRP) and the lipid ABCA1 transporters was detected in all ZIKV-infected groups, but BCRP and ABCA-1 were only reduced in ICompromised and high-ZIKV ICompetent mice. Our data indicate that gestational ZIKV infection triggers specific proinflammatory responses and affects placental turnover and transporter expression, in a way dependent on virus concentration

and maternal immune status. Placental damage may impair proper fetal-maternal exchange function and fetal growth/survival, likely contributing to congenital Zika syndrome.

1. Introduction

Congenital Zika virus (ZIKV) infection can be associated with adverse pregnancy outcomes. As such, neonates born from ZIKV-positive pregnancies may develop severe neurological abnormalities, placental pathologies and intrauterine growth restriction (IUGR), among other complications (1). ZIKV vertical transmission has become a major public health issue worldwide, especially in Brazil, where more than 200,000 ZIKV-positive cases have been confirmed, and over 2,000 congenital microcephaly births have been reported (2–6). These numbers represented a 20-fold rise in the incidence of congenital microcephaly in Brazil during the years of the ZIKV pandemic, with similar increases reported elsewhere in Latin America (2,3,7). Importantly, while the ZIKV pandemic is currently thought to be controlled, evidence points to a possible silent ZIKV spread across the Americas (8,9), highlighting the need for improved knowledge of the possible routes of ZIKV vertical transmission and its association with disruptive inflammatory and developmental phenotypes and the need for new avenues of prevention and treatment.

Studies have investigated the possible pathways involved in ZIKV vertical transmission. Miranda and colleagues (10) showed that in humans, ZIKV infection changed the pattern of tight junction proteins, such as claudin-4, in syncytiotrophoblasts. Jurado et al. (2016) suggested that the migratory activities of Hofbauer cells (feto-placental macrophages) could help disseminate ZIKV to the fetal brain (11). Other recent studies have shown that placental villous fibroblasts, cytotrophoblasts, endothelial cells

and Hofbauer cells are permissive to ZIKV, and placentae from ZIKV-infected women had chorionic villi with a higher mean diameter (11–14). However, at present, there is limited information on the precise mechanism of maternal-fetal ZIKV transmission.

Many mouse models have been developed to identify how ZIKV overcomes placental defenses. First, limited information was obtained due to the apparent inability of the virus to infect wild-type (WT) mice (15). ZIKV NS5 targets the interferon signaling pathway in humans but not in mice (16). As such, WT mice develop no clear evidence of clinical disease (16,17) and are of limited use in modeling the disease. However, mice lacking an interferon signaling response show evidence of disease and have been widely used to investigate ZIKV infection during pregnancy (8,16,18).

The interferon system, especially type III interferon, is one of the key mechanisms of host defense and viral targets for immune evasion (19). Type III interferons have a role in protection against ZIKV infection in human syncytiotrophoblasts from term placenta (20). Luo et al. have shown that the inhibition of Toll-like receptors 3 and 8 inhibits the cytokine output of ZIKV-infected trophoblasts (21). In addition, viral replication coincides with the induction of proinflammatory cytokines such as IL-6. This cytokine has a crucial role in inflammation and affects the homeostatic processes related to tissue injury and activation of stress-related responses (22,23). ZIKV infection can trigger an inflammatory response with IL-6 release (11,24).

Maternal infection has profound effects on placental permeability to drugs and environmental toxins. Changes in the expression and function of specific ABC transporters in the placenta and yolk sac following infective and inflammatory stimuli have been demonstrated (25–29). ABC transporters are efflux transporters that control the biodistribution of several endogenous and exogenous substrates, including xenobiotics (antiretrovirals, synthetic glucocorticoids), steroid hormones (estrogens,

androgens), nutrients (folate, cholesterol) and immunological factors (chemokines and

cytokines) within the maternal-fetal interface (30). The best described ABC transporters

in the placenta are the multidrug resistance P-glycoprotein (P-gp), the breast cancer

resistance protein (Bcrp) and the lipid Abca1 and Abcg1 transporters. P-gp and BCRP

transporters are responsible for preventing fetal accumulation of xenobiotics and

environmental toxins that may be present in the maternal circulation, whereas ABCA1

and ABCG1 control the placental exchange of cytotoxic oxysterol and lipid permeability

throughout pregnancy; therefore, they play an important role in fetal protection and

placental lipid homeostasis (25).

Despite a limited number of studies showing ZIKV infection in

immunocompetent mice, it was previously demonstrated that intrauterine inoculation of

high virus titer resulted in decreased fetal viability, with worse outcomes following

infection in early gestation (31). In another report, intravenous infection at very early

embryonic day resulted in fetal demise even though the virus was not found in the fetal

compartment in most treated animals (32). In the present study, we hypothesize that

maternal exposure to ZIKV affects placental function, including placental ultrastructure

and ABC transporter (P-gp, Bcrp, Abca1 and Abcg1) protein expression, even in the

absence of vertical transmission and that these effects are dependent on viral infective

titers and maternal immune status.

2. Materials and methods

2.1 Virus preparation and storage

The Brazilian $ZIKV_{PE243}$ (GenBank ref. number KX197192) strain was isolated

from a febrile case during the ZIKV outbreak in the state of Pernambuco, Brazil and was

kindly provided by Dr Ernesto T. Marques Jr. (Centro de Pesquisa Aggeu Magalhães,

FIOCRUZ, PE). Viruses were propagated in C6/36 cells with viral titers determined by

plaque assays in Vero cells, as previously described (33). Supernatants of noninfected

C6/36 cells cultured under the same conditions were used as mock controls.

2.2 Animal experimentation and study design

Two mouse strains were used in the study: ICompetent C57BL/6 and

ICompromised (type 1 *Ifnr*-deficient) A129 strains. Since we were unable to consistently

produce viable pregnancies by mating A129 males and females in our experimental

settings, we mated A129 females (n=15) with C57BL/6 males (n=4) to produce

ICompromised C57BL6/A129 pregnancies, whereas ICompetent C57BL/6 pregnancies

were obtained by mating male (n=6) and female (n=35) C57BL/6 mice (8-10 weeks old).

Animals were kept in a controlled temperature room (23°C) with a light/dark cycle of 12

hours and *ad libitum* access to water and food. After detecting the proestrous/estrous

phase by vaginal cytology, copulation was confirmed by visualization of the vaginal plug

and considered gestational day 0.5 (GD0.5). Maternal weight was monitored for

confirmation of pregnancy; thus, females were weighed on GD0.5 and GD12.5, and

females with a weight gain greater than 3 g were considered pregnant and entered

randomly in the experimental groups. Experimental protocols were approved by the

Animal Care Committee of the Health Sciences Center, Federal University of Rio de

Janeiro (CEUA-036/16 and 104/16) and registered within the Brazilian National Council

for Animal Experimentation Control.

On GD12.5, pregnant mice (ICompetent and ICompromised pregnancies) were

injected with a single intravenous (i.v.) titer of ZIKV or mock control. ICompetent and

ICompromised pregnant mice were randomly subdivided into three experimental groups:

the mock (control) group, which received supernatants of noninfected C6/36 cell injection

(ICompetent mock and ICompromised mock); the high-ZIKV-titer group, inoculated

with 5×10^7 plaque-forming units (PFU) of ZIKV$_{PE243}$ (ICompetent high and

ICompromised high); and the ZIKV low-titer group, injected with 10^3 PFU of ZIKV$_{PE243}$

(ICompetent low and ICompromised low).

On the morning of GD18.5, all animals were euthanized with a sodium

phenobarbital overdose of 300 mg/kg. Maternal blood was collected by cardiac puncture,

centrifuged (10 min, 4000 g) and stored at -20°C. The maternal brain and spleen, all

placentae and all fetuses were dissected, collected and weighed, followed by fetal head

isolation and measurement. The three placentae closest to the mean weight in a litter were

selected for further analysis and cut in half using umbilical cord insertion as a reference

(34–36). One-half of the placental disk was frozen in liquid nitrogen for qPCR, and the

other half was fixed overnight in buffered paraformaldehyde (4%, Sigma-Aldrich, Brazil)

for ultrastructural and protein expression/localization analysis. Matched fetal heads,

maternal brains and spleens were frozen in liquid nitrogen for qPCR.

2.3 ZIKV RNA quantification by qPCR

ZIKV load was evaluated in maternal blood, brains and spleens and in the

placentae and fetal heads. Brains, spleens, placentae and fetal heads were macerated in

RPMI medium (Gibco™ RPMI 1640 Medium) normalized by the ratio of 0.2 mg of tissue

to every 1 µl of medium and plotted per gram of tissue. The macerated volume was

centrifuged at 4500 g for 5 min to remove tissue residues, and then 500 µl of the

centrifuged volume was used for RNA extraction using 1 mL of TRIzol reagent (Life

Technologies, Thermo Fischer, USA). Treatment with DNase I (Ambion, Thermo

Fischer, USA) was performed to prevent contamination by genomic DNA. cDNA was

synthesized using a cDNA High Capacity Kit (Applied Biosystems, Thermo Fischer,

USA) according to the manufacturer's instructions by subjecting the samples to the

following cycle: 25°C for 10 min, 37°C for 120 min and 85°C for 5 min. qPCR was

performed using a StepOnePlus Real-Time PCR system and TaqMan Master Mix

Reagents (Applied Biosystems, Thermo Fischer, USA) using primers and probes specific

for the protein E sequence (37). Samples were then subjected to the following cycle: 50°C

for 2 min, 40x (95°C for 10 min, 95°C for 15 sec, 60°C for 1 min).

2.3.1 qPCR

The placenta was macerated in 1.5 mL of TRIzol reagent (Life Technologies,

Thermo Fischer, USA). RNA extraction was performed following the manufacturer's

protocol. cDNA was prepared using Power SYBR Green PCR Master Mix (Life

Technologies, Thermo Fisher, USA). The reaction was carried out for selected genes

using intron-spanning primers (Table 1) and the StepOnePlus Real-Time PCR system

(Life Technologies, Thermo Fischer, USA). Samples were subjected to the following

cycle: 95°C for 10 min followed by 40 amplification cycles consisting of DNA

denaturation for 30 sec at 95°C and annealing of primers for 30 sec at 60°C. The threshold

cycles (Ct) for each gene of interest and for the reference gene glycerol 3-phosphate

dehydrogenase (*Gapdh*) and RNA Polymerase II Subunit A (*Polr2a)* were determined.

The relative expression for each gene was calculated using $2^{-\Delta\Delta CT}$ (38) and graphically

expressed as the fold-increase. The efficiency was calculated using the standard curve

method. The melting curves were analyzed for each sample.

Table 1: Primer sequences for the real-time PCR assay.

Gene	Primer sequences	GenBank accession no.
ZIKV	5'CCGCTGCCCAACACAAG3'	
	5'CCACTAACGTTCTTTTGCAGACAT3'	
mIL-6	5'TCATATCTTCAACCAAGAGGTA3'	NM_031168.2

	5'CAGTGAGGAATGTCCACAAACTG3'	
mIL-1b	5'GTAATGAAAGACGGCACACC3'	XM_006498795.4
	5'ATTAGAAACAGTCCAGCCCA3'	
mIL-10	5'TAAGGGTTACTTGGGTTGCCAAG3'	NM_010548.2
	5'CAAATGCTCCTTGATTTCTGGGC3'	
mIFN-γ	5'AGCAACAGCAAGGCGAAAA3'	NM_008337.4
	5'CTGGACCTGTGGGTTGTTGA3'	
m-IFN-I	5'CTGGAGCAGCTGAATGGAAAG3'	NM_010510.1
	5'CTTGAAGTCCGCCCTGTAGGT3'	
mGAPDH	5'CTTTGTCAAGCTCATTTCCTGG3'	XM_017321385.2
	5'TCTTGCTCAGTGTCCTTGC3'	
TNF-a	5'CCTCACACTCAGATCATCTTCTCA3'	NM_013693.3
	5'TGGTTCTCTTTGAGATCCATGC3'	
Polr2a	5'TCTGCCAAGAATGTGACGCT3'	NM_001291068.1
	5'CCAAGCGGCAAAGAATGTCC3'	

2.4 Detection of cytokines and chemokines in the maternal serum and placenta

Initially, placental tissue was homogenized in extraction buffer (50 mM Tris, 150 mM NaCl, 1X Triton, 0.1% SDS, 5 mM EDTA, 5 mM NaF, 50 mM sodium pyrophosphate, 1 mM sodium orthovanadate, pH 7.4) containing complete protease inhibitor cocktail (Roche Applied Science, Germany) with TissueLyser LT (Qiagen, Germany). The protein concentration of each sample was analyzed using the Pierce™ BCA Protein Assay Kit (Thermo Scientific, USA) according to the manufacturer's instructions. Analysis of the cytokines IL-6 and IL1-β and the chemokines monocyte chemoattractant protein-1 (MCP-1/CCL2) and chemokine (C-X-C motif) ligand 1

(CXCL1) in maternal serum and placenta was performed with the MILLIPLEX MAP

Mouse Cytokine/Chemokine Magnetic Bead Panel – Immunology Multiplex Assays

(MCYTOMAG-70K, Merck Millipore, Germany), following the manufacturer's

recommendations. The plate with samples and magnetic beads was analyzed on a

MAGPIX® System (Merck Millipore, Germany). The analyses were performed using the

Luminex xPonent® for MAGPIX® v software. 4.2 (Luminex Corp., USA). For each

reaction well, the MAGPIX Luminex® platform reports the median fluorescence

intensity (MFI) for each of the analytes in the samples. The levels of each analyte were

then calculated against the standard curve. The ratio between the value obtained and the

protein quantification for each sample was determined and plotted.

2.5 Virus titration by plaque assays

Blood from mock- and $ZIKV_{PE243}$-infected mice was collected from the base of

the tail at 4 hours, 48 hours and 144 hours following the appropriate treatments and

subsequently centrifuged at 400 g for 30 min for plasma separation. Samples obtained at

different periods post infection were titrated by a plaque assay. Vero cells (obtained from

ATCC® CCL81™) (African green monkey kidney epithelial cell line) were plated in 24-

well plates at $4x10^4$ cells per well in DMEM (GIBCO, Thermo Fisher, USA)

supplemented with 5% FBS (GIBCO, Thermo Fisher, USA) and 1% gentamicin (10

μg/ml) (GIBCO, Thermo Fisher, USA) and cultured overnight for complete adhesion at

37°C with 5% CO_2. Then, the medium was removed, and the cells were washed with 1x

PBS and incubated with serial (base 10) dilutions of virus in FBS-free medium. After 90

min of incubation under gentle shaking, the medium was removed, and the cells were

washed with 1x PBS and cultured with 1.5% carboxymethylcellulose (CMC)

supplemented with 1% FBS (GIBCO, Thermo Fisher, USA). After 5 days, the cells were

fixed overnight with 4% formaldehyde and stained with 1% crystal violet in 20%

methanol (ISOFAR, Brazil) for 1 hour. Plaques were counted, and the virus yield was

calculated and expressed as plaque-forming units per milliliter (PFU/ml).

2.6 Histological, immunohistochemistry and TUNEL analyses of the placenta

Placental fragments were fixed overnight and subjected to dehydration (increasing

ethanol series, ISOFAR, Brazil), diaphanization with xylol (ISOFAR, Brazil) and

paraffin (Histopar, Easypath, Brazil). Sections (5 μm) were prepared using the Rotatory

Microtome CUT 5062 (Slee Medical GmbH, Germany) and subjected to

immunohistochemistry and TUNEL analyses.

For immunohistochemistry, blocking endogenous peroxidase was performed with

3% hydrogen peroxide diluted in PBS, followed by microwave antigenic recovery in Tris-

EDTA (pH=9) and sodium citrate (pH=6) buffers (15 min for Tris-EDTA buffer and 8

min for citrate buffer). Sections were washed in PBS + 0.2% Tween and exposed to 3%

PBS/BSA for 1 hour. Sections were then incubated overnight at 4°C with the following

primary antibodies: anti-Ki-67 (1:100 – [M3064]; Spring Bioscience, USA), anti-P-gp

(1:500 – Mdr1[sc-55510]; Santa Cruz Biotechnology, USA), anti-Bcrp (1:100 – Bcrp

[MAB4146]; Merck Millipore, USA), anti-Abcg1 (1:100 – [PA5-13462]; Thermo Fisher

Scientific, USA) or anti-Abca1 (1:100 – [ab18180]; Abcam Plc, UK). The next day,

sections were incubated with the biotin-conjugated secondary antibody SPD-060 (Spring

Bioscience, USA) for 1 hour at room temperature. Three washes were performed with

PBS + 0.2% Tween followed by incubation with streptavidin (SPD-060 - Spring

Bioscience, USA) for 30 min. Sections were stained with 3,3-diamino-benzidine (DAB)

(SPD-060 - Spring Bioscience, USA), counterstained with hematoxylin (Proquímios,

Brazil), dehydrated, diaphanized and mounted with coverslip and entellan (Merck,

Germany).

For analysis of apoptotic nuclei, terminal deoxynucleotidyl transferase dUTP

nick-end labeling (TUNEL) staining was performed using the ApopTag® In Situ

Peroxidase Detection Kit (S7100, Merck Millipore, USA) according to the manufacturer's

recommendations and as previously described (35). All negative controls were performed

with the omission of the primary antibody.

Image acquisition was performed using a high-resolution Olympus DP72

(Olympus Corporation, Japan) camera coupled to an Olympus BX53 light microscope

(Olympus Corporation, Japan). For nuclear quantification of Ki-67 and TUNEL

immunolabeling, Stepanizer software (39) was used. For this analysis, we evaluated 15

images from different random fields of the Lz and Jz for each animal, in a total of five

animals from each ICompetent group and three animals from each ICompromised group.

A total of 360 digital images per placental region (Lz and Jz) were evaluated in each

experimental group. Total numbers of immunolabeled Ki-67 or TUNEL nuclei from each

image were normalized by the total image area to obtain an index of estimated numbers

of proliferative and apoptotic nuclei in the whole histological section analyzed. Analysis

was undertaken by two investigators blinded to the treatment.

Quantification of P-gp, Bcrp, Abca1 and Abcg1 staining was performed using the

Image-Pro Plus, version 5.0 software (Media Cybernetics, USA) mask tool. The

percentage of viable tissue area was considered upon exclusion of negative spaces. A total

of 360 digital images per placental region (Lz and Jz) were evaluated in each experimental

group. Analysis was undertaken by two investigators blinded to the treatment.

2.7 Transmission electron microscopy (TEM)

Sections of placental Lz and Jz were fixed in paraformaldehyde (4%, Sigma-Aldrich, Brazil) for 48 hours, postfixed with osmium tetroxide (Electron Microscopy Sciences, USA) and potassium ferrocyanide (Electron Microscopy Sciences, USA) for 60 min and dehydrated with an increasing series of acetone (30%, 50%, 70%, 90% and two of 100%) (ISOFAR, Brazil). Sections were subsequently embedded with EPOXI resin (Electron Microscopy Sciences, USA) and acetone (1:2, 1:1 and 2:1, respectively). After polymerization, ultrafine sections (70 nm) were prepared (Leica Microsystems, USA) and collected into 300 mesh copper grids (Electron Microscopy Sciences, USA). Tissue was contrasted with uranyl acetate and lead citrate and visualized using a JEOL JEM-1011 transmission electron microscope (JEOL, Ltd., Akishima, Tokyo, Japan). Digital micrographs were captured using an ORIUS CCD digital camera (Gatan, Inc., Pleasanton, California, EUA) at 6000× magnification. An overall qualitative analysis of Lz and Jz from different groups was performed by investigating the ultrastructural characteristics of the mitochondria as well as the ER cisterns. The qualitative evaluation consisted of analyzing the disruption of the mitochondrial membranes, mitochondrial morphology, preservation of mitochondrial cristae and matrix intensity (40). Ultrastructural analysis of nuclear morphology and the presence of microvilli in trophoblast sinusoidal giant cells was also undertaken. The analysis of the cisterns of the ER was performed by evaluating the dilation of their lumen (41).

2.8 Statistical analysis

GraphPad Prism 8 software (GraphPad Software, Inc., USA) was used for

statistical analysis. The D'Agostino & Pearson normality test was used to evaluate normal

distribution, and outliers were identified using the Grubbs test. Data are expressed as the

mean ±SEM or individual values. One-way ANOVA followed by Tukey's posttest was

used for comparisons between different inbred groups, whereas Student's t-test or

nonparametric Mann-Whitney test was performed between the outbred groups.

Differences were considered significant when p<0.05. Pregnancy parameters were

evaluated using the mean value of all fetuses and placentae in a litter per dam and not the

individual conceptus, i.e., the mean value. "n" represents the number of litters in Figures

1 and 2 (34–36).

3.   Results

3.1 Weight gain across pregnancy is dependent on maternal immune status in ZIKV-

infected mice.

To determine the effect of ZIKV infection on fetal and placental phenotypes at

term (GD18.5), we infected ICompetent C57BL/6 and immunocompromised

(ICompromised) A129 mice with ZIKV at GD12.5 (Figure 1A). Given the very distinct

susceptibility of C57BL/6 and A129 mice to ZIKV, systemic infection models were

established by injecting high (5×10^7 PFU) and low (10^3 PFU) virus inoculum titers. As

shown in Table 2, ICompetent C57BL/6 mice from all groups and ICompromised A129

dams inoculated with mock and low exhibited higher maternal weight at GD18.5 than at

GD12.5 and GD0.5 (p<0.05). On the other hand, ICompromised A129 mice presented

significant weight loss at GD18.5, despite showing an increase at GD12.5 in relation to

GD0.5.

Table 2: Maternal weight gain throughout pregnancy

Maternal weight (g)	ICompetent			ICompromised		
	Mock (n=11)	High (n=15)	Low (n=9)	Mock (n=3)	High (n=3)	Low (n=3)
GD0.5	22.4 ± 0.7^a	23.0 ± 0.6^a	21.9 ± 0.7^a	21.7 ± 1.9^a	$21.7\pm0.9^{a,b}$	25.0 ± 1.0^a
GD12.5	25.5 ± 0.6^b	26.2 ± 0.8^b	25.1 ± 0.5^b	25.3 ± 1.2^a	26.0 ± 1.0^a	28.0 ± 0.6^a
GD18.5	31.6 ± 0.7^c	30.3 ± 0.9^c	31.4 ± 1.1^c	32.0 ± 1.5^b	20.0 ± 1.5^b	30.7 ± 0.3^b
ΔGD12.5-0.5 (Δ)	4	3	3	3	4	3
ΔGD18.5-0.5 (Δ)	9	7	10	10	-2	5

GD=gestational day; Δ=delta; values are expressed as the mean±standard error of the
mean (SEM). One-way ANOVA followed by Tukey's test analyzed differences among
GD0.5, 12.5 and 18.5 in each group; different letters indicate p<0.05.

3.2 Immunocompetent and immunodeficient mice have distinct term placental and fetal

phenotypes in response to high and low ZIKV titer challenges in mid-pregnancy.

The fetuses from C57BL/6 dams and sires were called ICompetent. The fetuses

from the mates of A129 dams and C57BL/6 sires were called ICompromised. High-ZIKV

ICompetent mice exhibited 34% fetal loss, whereas high-ZIKV-A129 mice had 100%

fetal loss (Figure 1B). In the low-ZIKV groups, C57BL/6 mice had no (0%) fetal death,

while A129 mice exhibited a 42% fetal death rate (Figure 1B). Fetal and fetal head sizes

were decreased in A129 mice compared to those in C57BL/6 dams infected with the high

ZIKV titer (p=0.05; Figure 1 C-F). However, no changes in fetal weight and fetal head

sizes were observed when the mice were infected with the lower ZIKV titer (Figure 1 C-

F).

Figure 1: ZIKV infection induced fetal changes during pregnancy. A) Experimental schematic. The fetuses from C57BL/6 dams and sires were termed ICompetent. The fetuses from the mates of A129 dams and C57BL/6 sires were termed ICompromised. B) Uterine horn images and survival images (arrows show resorption sites). C) Fetal/reabsorption images. Measurement of fetal biometric parameters at GD18.5. D) Fetal weight, (E) fetal head weight and (F) fetal head weight/fetal weight ratio in ICompetent (n=11/15/9 dams in the mock, high and low groups, respectively) and ICompromised (n=3 dams/group). * example reabsorption. The values are plotted as individual values, and the bar represents the mean. Unpaired Student's t-test or nonparametric Mann-Whitney test was used for ICompromised groups, and one-way ANOVA followed by Tukey's post-test was used for ICompetent comparisons between different groups.

3.3 ZIKV is detected in the fetal brain of ICompromised, but not ICompetent mice.

ZIKV RNA was detected in the spleens of pregnant ICompetent C57BL/6 mice inoculated with the highest ZIKV titer, confirming acute systemic infection. Viral RNA was also detected in the majority of placentae of those mice (Figure 2A-B) but not in the maternal and fetal C57BL/6 brains (Figure 2A-B), suggesting that the virus was not transmitted to the fetuses. Although low ZIKV inoculation resulted in virus detection in

the spleens of pregnant C57BL/6 mice, infection was not evidenced in the placentae or

fetal brains. In contrast, ZIKV RNA was detected in all analyzed organs from

ICompromised A129 dams, including the maternal brain and spleen and the placenta and

fetal brain (Figure 2A-B). Viremia in maternal serum was evaluated at 4, 48 and 144

hours after infection. Within 4 hours, the presence of the virus was verified in the serum

(high ICompetent = 637.5 PFU/mL, low ICompetent = 740 PFU/mL and low

ICompromised = 325 PFU/mL), indicating that the virus was correctly inoculated.

Afterwards, ZIKV RNA was detected in ICompromised dams at 48 and 144 hours

postinoculation but not in ICompetent dams.

Figure 2: Viral load present in the spleen, placenta, fetal and maternal brains after ZIKV infection. A) Maternal spleen and brain viral load and B) placenta and fetal brain viral load of ZIKV-infected dams. C) Animal viral serology of all groups after 4, 48 and 144 infection hours. ICompetent (n=11/15/9 dams in the mock, high and low ZIKV titer groups, respectively) and ICompromised (n=3 dams/group).

3.4 ZIKV infection induces distinct systemic and placental inflammatory responses in

ICompetent and ICompromised mice.

The maternal serum and placental protein levels of specific cytokines and chemokines

related to fetal death and preterm delivery (42–44) were evaluated to probe whether

midgestation ZIKV infection would induce a maternal inflammatory response at term in

our two distinct models. Since A129 infected with 10^7 ZIKV-PFU showed 100% fetal

loss, we proceeded using 10^7 PFU inoculation in C57BL/6 and 10^3 PFU inoculation in

both C57BL/6 and A129 mice.

CCL2 was elevated in the serum of low-ZIKV ICompetent mice, but no other

alteration was systemically detected in any ICompetent mice at this time point (Figure

3A-B). On the other hand, ICompromised dams showed significantly increased CXCL1

and IL-6 levels in the serum, and a strong trend for an enhancement in CCL2 (Figure 3C).

Analysis of cytokines and chemokines expression at the placenta demonstrated that

CXCL1 and CCL2 chemokines were also upregulated in the ICompromised, but not of

ICompetent mice (Figure 3 D-F). Surprisingly, IL-6 protein expression was augmented

in some of the low-ZIKV ICompetent mice (58%; p<0.05), but not in high-ZIKV nor

ICompromised mice (Figure 3D-F).

We also assessed the placental mRNA expression of a range of cytokines related

to placental infective responses: Ifn-γ, Il-6, Ifn-1, Tnf-α, Il-1β, and Il-10. All the infected

mice showed a significantly increase in TNF- α mRNA expression, with higher levels

detected in ICompetent mice (Figure 3G-I). The low-ZIKV ICompetent mice presented a

modest, but significant type I IFN expression, which was not detected in the high-

ZIKVgroup (Figure 3G). Additionally, both low-ZIKV infected groups (ICompetent and

ICompromised) presented increased Ifn-γ mRNA expression (Figure 3G and 3I, p< 0.05).

Interestingly, placental Il-6 mRNA levels were only elevated in ICompromised

pregnancies compared to the mock (p=0.05) (Figure 3I). Il-1b and IL-10 remained

unchanged in all groups analyzed (Figure 3 G-I).

Figure 3: ZIKV promotes an inflammatory response in the placenta, maternal spleen and serum during pregnancy.
A) Cytokine and chemokine serum levels at GD18.5 in the ICompetent mock (n=9 dams), high (n=8) and low (n=9) and ICompromised mock and low (n=3/groups) ZIKV titer groups. A-C) IL-1β, IL-6, CCL-2 and CXCL-1 maternal serum levels. D-F) Placental IL-1β, IL-6, CCL-2 and CXCL-1 levels. G-I) Placental RNA levels of Ifnγ, *Ifn1, Il-1β, Il-6 and Tnf-α.* The broken line shows the expression levels of both lineages in the mock group. The values are expressed as the mean ± SEM. Unpaired Student's t-test or nonparametric Mann-Whitney test was used for comparisons between ICompetent groups, and one-way ANOVA followed by Tukey's posttest was used for comparisons between different ICompetent groups. * *p<0.05.*

3.5 ZIKV affects placental proliferation, apoptosis and ultrastructure in a viral load-

and maternal immune status-dependent manner.

We did not observe changes in placental weight or in the fetal:placental weight

ratio in any of the groups investigated (data not shown). However, since we detected the

presence of ZIKV RNA in the placentas, we investigated the cellular proliferation (Ki-

67$^+$ cells) and apoptotic ratio in the labyrinth zone (Lz) and junctional zone (Jz) of the

mouse placenta. Increased Ki-67 staining was observed in the Lz of high- and low-ZIKV-

treated ICompetent animals compared to mock animals (Figure 4A, p=0.005 and

p=0.015), whereas the Lz apoptotic ratio was increased in high- and decreased in low-

ZIKV dams (p=0.01, Figure 4B). A similar apoptotic pattern was detected in the Jz of

ICompetent pregnancies (p=0.008 and p=0.004, respectively; Figure 5B), although no

differences were observed in Ki67 staining (Figure 5A). In contrast, in ICompromised

dams, Lz Ki-67 staining was decreased (p=0.001; Figure 4C), while the apoptotic reaction

was increased in ZIKV-infected animals (p=0.05; Figure 5D). Jz from ICompromised

offspring exhibited increased Ki-67 labeling and no differences in the apoptotic reaction

(p=0.001; Figure 5 C-D).

Figure 4: Placental remodeling of ICompetent groups induced by ZIKV infection.
A) Quantification of Ki-67$^+$ nuclei in the mouse placenta Lz and representative images of
nuclei immunostained by Ki-67$^+$-positive nuclei in the ICompetent group (n=6/group).
B) The image shows the apoptotic nuclei score in the Lz and representative images of
nuclei immunostained by TUNEL in the ICompetent group (n=5/group). C)
Quantification of Ki-67+ nuclei in the mouse placenta Lz and representative images of
nuclei immunostained by Ki-67$^+$ nuclei in the ICompromised group (n=3/group). D) The
image shows the apoptotic nuclei score in the LZ and representative images of nuclei
immunostained by TUNEL in the ICompromised group (n=3/group). The values are
expressed as the mean ± SEM. One-way ANOVA followed by Tukey's post hoc test was
used for comparisons between different ICompetent groups. Unpaired Student's t-tests or
nonparametric Mann-Whitney tests were used for the ICompromised groups. CTR(-)=All
negative controls were performed with the omission of the primary antibody. Scale bar =
50 µm.

Figure 5: ZIKV induces hyperproliferation of the JZ area in infected placentas of
ICompetent groups. A) Quantification of Ki-67+ nuclei in the placental Jz of mice and
representative images of immunostained nuclei in the ICompetent (n=6/group). B) The
image shows apoptotic nuclei quantification in the Jz area and representative images of
nuclei immunostained by TUNEL in the ICompetent (n=5/group). C) Quantification of
Ki-67+ nuclei in the placental Jz area and representative images of immunostained nuclei
in the ICompromised group (n=3/group). D) The image shows apoptotic nuclei
quantification in the Jz area and representative images of nuclei immunostained by
TUNEL in the ICompromised (n=3/group). The values are expressed as the mean ± SEM.

One-way ANOVA followed by Tukey's post hoc test was used for comparisons between
different ICompetent groups. Unpaired Student's t-tests or nonparametric Mann-Whitney
tests were used for the ICompromised groups. CTR(-)=All negative controls were
performed with the omission of the primary antibody. Scale bar = 50 µm.

3.6 Placental ultrastructure is differently impacted by high- and low-titer ZIKV

infection in ICompetent and ICompromised strains.

Ultrastructural analyses in the Lz of ICompetent and ICompromised-mock

animals detected sinusoidal trophoblastic giant cells exhibiting regular microvilli,

euchromatic nuclei, preserved mitochondrial ultrastructure and regular narrow ER

cisternae (Figure 6A and 6D). In sharp contrast, high-ZIKV ICompetent (Figure 6B) and

low-ZIKV ICompromised infected placentae (Figure 6E) showed fewer villi in the

sinusoidal giant trophoblastic cells, degenerated mitochondria, granular ER with dilated

cisterns and euchromatic nuclei. The sinusoidal giant trophoblastic cells from the low-

ZIKV ICompetent (Figure 6C) also had fewer villi and degenerated mitochondria than

the mock placentae, but there and had no effect on the ER or euchromatic nuclei.

Figure 6: Associated ultrastructural changes in the placental LZ after ZIKV infection. Transmission electron photomicrographs of ICompetent mock (A); ICompetent high (B); ICompetent low (C); ICompromised mock (D) and ICompromised low (E) groups (N=5 dams/group). We observed dilatation in the ER cisterns of ICompetent high placentas. Additionally, there was a reduction in the microvilli from both ICompetent high and ICompetent low placentas. In the ICompromised low group, we found fragmented ER and microvillus reduction. All infected groups showed degenerate mitochondria. GER= Granular Endoplasmic Reticulum; ⊗= Dilated Granular Endoplasmic Reticulum; *= Fragmented Granular Endoplasmic Reticulum; Mt= Mitochondria; MtD= Degenerate Mitochondria; Mv= Microvilli; EN= Euchromatic Nuclei; SS= Sinusoidal Space; TGC= Trophoblastic Giant Cell. Scale bar = 2 µm.

The Jz of mock ICompetent and ICompromised (Figure 7A and 7D) and of low-ZIKV ICompetent (Figure 7C) placentae exhibited euchromatic nuclei, with evident heterochromatin, preserved mitochondria and narrow cisternae granular ER. High-ZIKV ICompromised and low-ZIKV ICompromised placentae had degenerated mitochondria, granular ER with dilated cisterns and euchromatic nuclei (Figure 7B and 7E).

Figure 7: Associated ultrastructural changes in the placental JZ after ZIKV infection. Transmission electron photomicrographs of ICompetent mock (A); ICompetent high (B); ICompetent low (C); ICompromised mock (D) and ICompromised low (E) groups (N=5 dams/group). We found deteriorating mitochondria and dilated reticulum endoplasmic cisterns in both the high ICompetent and low ICompromised groups. GER= Granular Endoplasmic Reticulum; ⊗= Dilated Granular Endoplasmic Reticulum; Mt= Mitochondria; MtD= Degenerate Mitochondria; Mv= Microvilli; EN= Euchromatic Nuclei; SS= Sinusoidal Space; TGC= Trophoblastic Giant Cell. White arrows HA = Heterochromatin Area. Scale bar = 2 µm.

3.7 ZIKV differentially affects placental expression of drug and lipid ABC transporter systems.

Evaluation of key ABC transporters in the Lz of mock and ZIKV-infected ICompetent and ICompromised placentae revealed that immunolabeling of the drug P-gp and Bcrp efflux transporter systems was primarily present at the cellular membranes of the sinusoidal trophoblastic giant cells, with diffuse cytoplasmic Bcrp staining. Labeling of the Abca1 and Abcg1 lipid efflux transporters was moderately and heterogeneously distributed within the Lz. Less Lz-P-gp was observed in ICompetent mice infected with

both high- and low-ZIKV infective regimens, in comparison to mock-treated animals

(p=0.001 and p=0.002, respectively; Figure 8A), whereas Bcrp and Abca1 staining was

only reduced was observed in high-ZIKV-infected mice (p=0.003 and p=0.004,

respectively; Figure 8B-C). P-gp, Bcrp and Abca1 transporter immunostaining was

downregulated in ICompromised low ZIKV-treated animals (p=0.001, p=0.05 and

p=0.05, respectively; Figure 9A-C). No changes in Abcg1 were observed in any of the

experimental groups (Figure 8D and Figure 9D).

Figure 8: ZIKV infection decreases P-gp, Bcrp and Abca1 expression in the
placental LZ of infected mice in the ICompetent groups. Immunostaining
quantification and representative immunohistochemistry images of A) P-gp, B) Bcrp, C)
Abca1 and D) Abcg1. The images represent the percentage of stained cells in each
analyzed field. The values are expressed as the mean ± SEM. One-way ANOVA followed
by Tukey's post hoc test was used for comparisons between different ICompetent groups.
Scale bar = 50 µm. N=6/ group. CTR(-)=All negative controls were performed with the
omission of the primary antibody.

Figure 9: ZIKV infection decreases P-gp, Bcrp and Abca protein expression in the placental LZ of infected mice in the ICompromised groups. Immunostaining quantification and representative immunohistochemistry images of A) P-gp, B) Bcrp, C) Abca1 and D) Abcg1. The images represent the percentage of stained cells in each analyzed field. The values are expressed as the mean ± SEM. Unpaired Student's t-tests or nonparametric Mann-Whitney tests were used for the ICompromised groups. Scale bar = 50 µm. N=3 for each group. CTR(-)=All negative controls were performed with the omission of the primary antibody.

Next, the impact of ZIKV on ABC transporters in the Jz layer (structural and endocrine layers of the mouse placenta) was assessed. P-gp and Bcrp were predominantly localized at the cellular membranes of spongiotrophoblast cells, whereas Abca1 and Abcg1 exhibited membrane and cytoplasmic staining. P-gp staining was decreased in Jz cells from the low-ZIKV ICompromised placentae (p=0.006), with no other alterations observed (Figure 10A-D and Figure 11A-D).

Figure 10: ZIKV infection did not impact P-gp, Bcrp, Abca1 or ABcg1 expression in the JZ area in infected placentas of ICompetent groups. Immunostaining quantification and representative immunohistochemistry images of A) P-gp, B) Bcrp, C) Abca1 and D) Abcg1. The images represent the percentage of stained cells/viable tissue. The values are expressed as the mean ± SEM. One-way ANOVA followed by Tukey's post hoc test was used for comparisons between different ICompetent groups. Scale bar = 50 µm. N=6/each group. CTR(-)=All negative controls were performed with the omission of the primary antibody.

Figure 11: ZIKV infection decreases Bcrp expression in the JZ area in infected placentas of the ICompromised group. Immunostaining quantification and representative immunohistochemistry images of A) P-gp, B) Bcrp, C) Abca1 and D) Abcg1. The images represent the percentage of stained cells/viable tissue. The values are expressed as the mean ± SEM. Unpaired Student's t-tests or nonparametric Mann-Whitney tests were used for the ICompromised groups. Scale bar = 50 µm. N=3 for each group. CTR(-)=All negative controls were performed with the omission of the primary antibody.

4. Discussion

In this study, we investigated several fetuses and placentas feature at term (GD18.5) in ICompetent (C57BL/6) and ICompromised (A129) mice exposed to ZIKV at mid-pregnancy (GD12.5). Fetal survival rates, systemic and placental inflammatory responses, placental ultrastructure and cell turnover, as well as the expression of key drug (P-gp and Bcrp) and lipid (Abca1) efflux transporter systems in the placenta, were consistently impacted by ZIKV in both strains. The magnitude of the effects was is clearly

related to the infective titer (high and low) of ZIKV and maternal immune status

(ICompetent-C57BL/6 x and ICompromised-A129) and fetal alterations were not

exclusively dependent on the virus detection in the fetuses.

Infection of ICompetent mice with ZIKV did not result in viremia in the initial

postinoculation phase, although viral RNA was detected in the maternal spleen in both

the high- and low-ZIKV groups, confirming systemic infection. This is consistent with a

previous report (45). Our data demonstrate that pregnant ICompetent C57BL/6 mice were

more susceptible to higher ZIKV than to lower ZIKV infective titers. Since viral RNA

was only detected in the placentae of high ZIKV, and fetal survival rates and weight were

impacted to a greater extent in those mice. Strikingly, even though the virus was not

present in the fetal brain (at least at term),fetal and fetal head weights were lower in the

high-ZIKV titer regimen, suggesting that high infective viral load in mid-pregnancy, even

in ICompetent individuals, can induce IUGR and lower fetal head weight despite a lack

of transmission to the fetal brain (46). On the other hand, ICompromised placentae and

fetal brains had detectable viral transcripts, with no changes in weight, which is consistent

with previous data (18). In fact, in our models, the presence (ICompromised) or the

absence (ICompetent) of the virus in the fetal brain did not correspond to fetal head size

(decreased in only high ICompetent). The data from ICompetent and ICompromised

placentae demonstrate how important the maternal immunological status is to control

viremia, fetal survival and accessibility of the virus into the fetal brain. The reason for the

reduction in fetal brain size in C57BL/6 mice in the absence of fetal brain infection

requires further investigation. It is possible that fetal brains of the ICompetent mice may

have been exposed to ZIKV earlier in pregnancy when viremia was present in the

maternal blood, and this may have severely compromised brain development.

A distinct inflammatory profile was also detected in the three analyzed groups. At

the protein level, low-ZIKV-ICompromised dams exhibited increased maternal IL-6 and

CXCL-1 and placental CCL-2 and CXCL-1, whereas low-ZIKV-ICompetent dams had

increased maternal CCL2 and placental IL-6 levels. CCCL-2 and CXCL-1 are related to

fetal death and preterm delivery (42–44) and could be associated with pronounced fetal

injure detected upon ICompromised pregnancy. In addition, it was previously

demonstrated (47) that IL-6 is related to fetal response syndrome, characterized by

activation of the fetal immune system. This syndrome is known to increase fetal morbidity

and affect several organs, such as the adrenal gland, brain and heart (31,47–50). At the

mRNA level, IL6 expression was only detected in ICompromised placentae at term and

may indicate a sustained harmful response in these mice until term. IFN signaling

pathway may be triggered by ZIKV (16) and represents one of the key mechanisms of

host defense and viral targets for immune evasion (19), but we did only detected a slight

increase in type I IFN in low-ZIKV ICompentent mice at term. However, we cannot rule

out the possibility that these cytokines might have been produced earlier. IFN-γ

expression was significantly enhanced in both ICompetent and ICompromised low-ZIKV

derived placentas, but not in the high-ZIKV infected mice. Although we could not assess

cytokine expression in high-ZIKV ICompromised placentas, one may extrapolate that

low-ZIKV infection could result in stimulation of IFN-γ-producing cells, which has been

previously showed to be protective for ZIKV-infected mice (51).

Both ICompromised and ICompetent mice showed increased expression of

placental *Tnf-α* mRNA, which has been demonstrated to be is directly related to placental

damage, abortion and premature birth (REF). In addition increased TNF-α response is

relatated to impaired placental hormone production and trophoblastic invasion and

increased apoptosis in pregnancy (52,53). Although we did not assessed TNF-α protein

levels in the placenta and maternal blood, this response could be implicated in the overall

damage detected.

Although differences in placental weight were not observed, ZIKV infection mid-

pregnancy had a profound effect on placental cellular turnover, dependent on titer, strain

and/or placental compartment. Lz is responsible for fetal and maternal nutrient, gas and

waste exchange, while Jz provides structural support, nutrient storage and hormone

synthesis (34). ZIKV induced a consistent increase in Lz proliferation in all groups.

However, the Lz apoptotic rate was increased only in the high-ZIKV-ICompetent and

ICompromised groups and decreased in low-ZIKV-ICompetent mice. The mechanisms

underlying these differences are unknown but may be related to the distinct maternal and

placental proinflammatory responses and/or to the direct effect of the virus on the

placenta (54). Increased Lz apoptosis in the high-ICompetent-ZIKV group may be one of

the mechanisms driving the lower fetal and fetal head weight detected in this group. In

this context, changes in placental turnover can determine placental maturation and

function and lead to fetal distress and developmental abnormalities (55). An increase in

the Lz apoptotic ratio may signify damage to this placental layer, which is consistent with

the fact that diverse pathological lesions associated with congenital disorders were

described in placentae from women infected by ZIKV at different stages of pregnancy

(56). Conversely, no proliferative changes were observed in Jz in high-ZIKV and low-

ZIKV ICompetent mice, while increased and decreased apoptotic rates were detected. It

follows that the lack of Jz-Ki-67 induction may suggest that this layer is less capable of

restoring proliferation in response to high-ZIKV challenge, and this may be related to

decreased fetal growth.

Our placental ultrastructural analysis detected consistent differences across

ZIKV-exposed groups. The Lz and Jz layers from both strains exhibited signs of ER

stress, i.e., dilated ER cisterns or fragmented ER granular structures. These alterations

may result from the accumulation of folded or poorly folded viral proteins in the ER

lumen (41,57,58). It has been demonstrated that the *Flaviridae* family uses the ER to

replicate (59), and according to Offerdahl et al. (2017)(58), there is evidence of ZIKV

interacting with this organelle, promoting an increased release of Ca^{+2} from the ER to the

cell cytoplasm, thereby causing an increase in the production of reactive oxygen species

(ROS) (60,61).

The mitochondrial ultrastructure in the Lz and Jz layers was severely impacted by

ZIKV exposure. We found evidence of mitochondrial degeneration, i.e., mitochondrial

membrane rupture, absence of mitochondrial ridges and less electron-dense

mitochondrial matrix in all treated groups. Placental mitochondrial dysfunction is

associated with IUGR (62,63) and may be related, at least in part, to the lower fetal weight

observed in high-ZIKV-ICompetent fetuses along with the altered placental apoptotic and

proliferative patterns. Furthermore, mitochondrial dysfunction together with ER stress is

likely to modify the placental ROS balance and generate local oxidative stress (64), which

is associated with impaired fetal development (65). Of importance, associations between

mitochondrial disruption, ER stress and placental cell senescence have been reported.

Senescence is characterized as an irreversible interruption of the cell cycle and acquisition

of a senescence-associated secretory phenotype (SASP) that promotes the release of

cytokines such as Il-1, Il-6, Il-8 and proinflammatory proteases (65). Therefore, the

increased expression of IL-6 detected in the placentas of ICompromised mice suggest a

SASP profile, which may be related to changes in the ER and mitochondrial

ultrastructure, accompanied by important changes in apoptosis and cell proliferation. The

interactions between mitochondria and the ER are critical for homeostasis and cell

signaling (66). In conjunction with the ER, mitochondria can regulate cell death mediators

in response to hypoxia and inflammation (67). The increase in apoptosis observed in the

high-titer ICompetent groups and the low-titer ICompromised group may be related to

the mitochondrial damage and ER stress observed. In fact, we observed an important

decrease in microvillus abundance in sinusoidal giant trophoblast cells. Previously, we

observed a decrease in microvillus density in the Lz of pregnancies exposed to malaria in

pregnancy (MiP) (35). Together, our data show that different gestational infective stimuli

(MiP and ZIKV) are capable of damaging placental microvillus abundance and impairing

proper fetal-maternal exchange function and fetal growth/survival.

Next, to investigate whether maternal ZIKV exposure may influence fetal

protection, we evaluated the placental localization and expression (semiquantitative) of

the ABC efflux transporter systems P-gp, Bcrp, Abca1 and Abcg1, which are highly

enriched in labyrinthine microvilli and in human syncytiotrophoblasts. These efflux

transporters exchange drugs, environmental toxins, cytotoxic oxysterols and lipids within

the maternal-fetal interface (25). We found a consistent decrease in labyrinthine P-gp

expression in all ZIKV-exposed groups, demonstrating that ZIKV infection during

pregnancy has the potential to increase fetal exposure to P-gp substrates such as synthetic

glucocorticoids, antibiotics, antiretrovirals, antifungals, stomach-protective drugs, and

nonsteroidal anti-inflammatory drugs (25). Furthermore, Jz-P-gp was decreased in

ICompromised placentae. Although little is known about the Jz function of ABC

transporters, our data highlight the need for further studies investigating the biological

importance of ABC transporters in the placental endocrine and structural zones of the

rodent hemochorial placenta under normal and infective conditions.

ZIKV impaired Lz Bcrp and Abca1 expression in ICompetent (high) and

ICompromised (low) mice. However, no effects were observed in ICompetent animals at

a low ZIKV titer or in Abcg1 in any experimental setting. As such, ZIKV also likely

increases the fetal accumulation of Bcrp substrates (antibiotics, antiretrovirals,

sulfonylureas, folate, mercuric species, estrogenic mycotoxins, carcinogens phototoxic

compounds, among others) and disrupts placental lipid homeostasis (lipids, cholesterol,

cytotoxic oxysterols) by reducing placental Abca1 expression (25,68–71). We can

speculate that the increased fetal accumulation of the P-gp, Bcrp and Abca1 substrates

during ZIKV infection may contribute to the establishment of congenital Zika syndrome,

although additional studies are clearly required to answer this important question. The

current data are in agreement with previous publications showing that bacterial, viral and

protozoan inflammation alters the expression and/or function of P-gp, Bcrp and Abca1 in

biological barriers such as the placenta, yolk sac and blood-brain barriers

(25,26,35,59,72–74).

5. Conclusion

Our data show that gestational ZIKV impacts the fetal phenotype independently

of term fetal viremia. Abnormal placental cell turnover, ultrastructure and transporter

expression may result from the specific proinflammatory responses that depend on the

ZIKV infective load and maternal immune status. Fetal accumulation of drugs,

environmental toxins and lipids within the fetal compartment may be potentially

increased in ZIKV-infected pregnancies due to altered levels of key ABC transporters.

Conflict of Interest:

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

Author contributions

CBVA, FFB, EB, LBA and TMOC conceived and designed the experiments. CBVA, VRSM, SVAC, HRG, RPCS and VMON performed the experiments. CBVA, SVAC, FFB, EB, SGM, LBA and TMOC analyzed the data. CBVA, VRSM, EB, LBA and TMOC wrote the paper and edited the manuscript. All authors contributed to the article and approved the submitted version.

Funding

This study was supported by the Bill & Melinda Gates Foundation (MCTI/CNPq/MS/SCTIE/Decit/Bill and Melinda Gates 05/2013; OPP1107597), the Canadian Institutes for Health Research (SGM: Foundation-148368), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq; 304667/2016-1, 422441/2016-3, 303734/2012-4, 422410/2016-0), Coordenação de Aperfeiçoamento Pessoal de Nível Superior (CAPES, finance Code 001), Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ, CNE 2015/E26/203.190/2015, PDR-26/2002/010/2016), and PRPq-Universidade MG Federal MGPEFARJ, CNE 2015/E26/203.190/2015).

Acknowledgments

We would like to thank Alan Moraes for supporting the acquisition of electron microscopy images and the electron microscopy laboratory at the UFF Biology Institute for allowing the use of the JEM1011 transmission electron microscope. Mauro Jorge Castro Cabral for the use of the MAGPIX® System equipment from the Paulo de Góes Institute of Microbiology/UFRJ. Ernesto T. Marques Jr. (Centro de Pesquisa Aggeu

754 **References**
755

1.   Garcez PP, Loiola EC, Madeiro da Costa R, Higa LM, Trindade P, Delvecchio R, et
al. Zika virus impairs growth in human neurospheres and brain organoids. Science.
2016;352(6287):816–8. doi: 10.1126/science.aaf6116.
2.   Brasil P, Calvet GA, Siqueira AM, Wakimoto M, de Sequeira PC, Nobre A, et al.
Zika Virus Outbreak in Rio de Janeiro, Brazil: Clinical Characterization, Epidemiological
and Virological Aspects. PLoS Negl Trop Dis. 2016;10(4):e0004636. doi:
10.1371/journal.pntd.0004636.
3.   Brasil P, Nielsen-Saines K. More pieces to the microcephaly–Zika virus puzzle in
Brazil. The Lancet Infectious Diseases. 2016;16(12):1307–9. doi: 10.1016/S1473-
3099(16)30372-3.
4.   Brasil P, Pereira JP, Moreira ME, Ribeiro Nogueira RM, Damasceno L, Wakimoto
M, et al. Zika Virus Infection in Pregnant Women in Rio de Janeiro. N Engl J Med. 2016
Mar 4;375(24):2321–34. doi: 10.1056/NEJMoa1602412
5.   Jaenisch T, Rosenberger KD, Brito C, Brady O, Brasil P, Marques ET. Risk of
microcephaly after Zika virus infection in Brazil, 2015 to 2016. Bull World Health
Organ. 2017;95(3):191–8. doi: 10.2471/BLT.16.178608.
6.   Proenca-Modena JL, Milanez GP, Costa ML, Judice CC, Maranhão Costa FT. Zika
virus: lessons learned in Brazil. Microbes and Infection. 2018;20(11–12):661–9. doi:
10.1016/j.micinf.2018.02.008.
7.   Jouannic J-M, Friszer S, Leparc-Goffart I, Garel C, Eyrolle-Guignot D. Zika virus
infection in French Polynesia. The Lancet. 2016;387(10023):1051–2. doi:
10.1016/S0140-6736(16)00625-5.
8.   Musso D, Ko AI, Baud D. Zika Virus Infection — After the Pandemic. N Engl J
Med. 2019;381(15):1444–57. doi: 10.1056/NEJMra1808246.
9.   Kasprzykowski JI, Fukutani KF, Fabio H, Fukutani ER, Costa LC, Andrade BB, et al.
A recursive sub-typing screening surveillance system detects the appearance of the
ZIKV African lineage in Brazil: Is there a risk of a new epidemic? International Journal of
Infectious Diseases. 2020;96:579–81. doi: 10.1016/j.ijid.2020.05.090.
10.   Miranda J, Martín-Tapia D, Valdespino-Vázquez Y, Alarcón L, Espejel-Nuñez A,
Guzmán-Huerta M, et al. Syncytiotrophoblast of Placentae from Women with Zika
Virus Infection Has Altered Tight Junction Protein Expression and Increased
Paracellular Permeability. Cells. 2019;8(10):1174. doi: 10.3390/cells8101174.
11.   Jurado KA, Simoni MK, Tang Z, Uraki R, Hwang J, Householder S, et al. Zika virus
productively infects primary human placenta-specific macrophages. JCI Insight.
2016;1(13):e88461. doi: 10.1172/jci.insight.88461.

12.   Quicke KM, Bowen JR, Johnson EL, McDonald CE, Ma H, O'Neal JT, et al. Zika
Virus Infects Human Placental Macrophages. Cell Host & Microbe. 2016;20(1):83–90.
doi: 10.1016/j.chom.2016.05.015.

13.   Tabata T, Petitt M, Puerta-Guardo H, Michlmayr D, Wang C, Fang-Hoover J, et
al. Zika Virus Targets Different Primary Human Placental Cells, Suggesting Two Routes
for Vertical Transmission. Cell Host & Microbe. 2016;20(2):155–66. doi:
10.1016/j.chom.2016.07.002

14.   Simoni MK, Jurado KA, Abrahams VM, Fikrig E, Guller S. Zika virus infection of
Hofbauer cells. Am J Reprod Immunol. 2017;77(2):e12613. doi: 10.1111/aji.12613.

15.   Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, et al. A Mouse
Model of Zika Virus Pathogenesis. Cell Host & Microbe. 2016;19(5):720–30. doi:
10.1016/j.chom.2016.03.010.

16.   Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, et al.
Zika Virus Targets Human STAT2 to Inhibit Type I Interferon Signaling. Cell Host &
Microbe. 2016;19(6):882–90. doi: 10.1016/j.chom.2016.05.009.

17.   Yockey LJ, Varela L, Rakib T, Khoury-Hanold W, Fink SL, Stutz B, et al. Vaginal
Exposure to Zika Virus during Pregnancy Leads to Fetal Brain Infection. Cell.
2016;166(5):1247-1256.e4. doi: 10.1016/j.cell.2016.08.004.

18.   Garcez PP, Stolp HB, Sravanam S, Christoff RR, Ferreira JCCG, Dias AA, et al. Zika
virus impairs the development of blood vessels in a mouse model of congenital
infection. Sci Rep. 2018;8(1):12774. doi: 10.1038/s41598-018-31149-3.

19.   Elong Ngono A, Shresta S. Immune Response to Dengue and Zika. Annu Rev
Immunol. 2018;36(1):279–308. doi: 10.1146/annurev-immunol-042617-053142.

20.   Bayer A, Lennemann NJ, Ouyang Y, Bramley JC, Morosky S, Marques ETDAJ, et
al. Type III Interferons Produced by Human Placental Trophoblasts Confer Protection
against Zika Virus Infection. Cell Host Microbe. 2016;19(5):705–12. doi:
10.1016/j.chom.2016.03.008

21.   Luo H, Winkelmann ER, Fernandez-Salas I, Li L, Mayer SV, Danis-Lozano R, et al.
Zika, dengue and yellow fever viruses induce differential anti-viral immune responses
in human monocytic and first trimester trophoblast cells. Antiviral Research.
2018;151:55–62. doi: 10.1016/j.antiviral.2018.01.003.

22.   Tanaka T, Narazaki M, Kishimoto T. IL-6 in Inflammation, Immunity, and
Disease. Cold Spring Harb Perspect Biol. 2014;6(10):a016295. doi:
10.1101/cshperspect.a016295.

23.   Hunter CA, Jones SA. IL-6 as a keystone cytokine in health and disease. Nature
Immunology. 2015;16(5):448–57. doi: 10.1038/ni.3153.

24.   Rabelo K, Gonçalves AJ da S, Souza LJ de, Sales AP, Lima SMB de, Trindade GF,
et al. Zika Virus Infects Human Placental Mast Cells and the HMC-1 Cell Line, and
Triggers Degranulation, Cytokine Release and Ultrastructural Changes. Cells.
2020;9(4):975–90. doi: 10.3390/cells9040975.

25.   Bloise E, Ortiga-Carvalho TM, Reis FM, Lye SJ, Gibb W, Matthews SG. ATP-
binding cassette transporters in reproduction: a new frontier. Hum Reprod Update.
2015;22(2):164–81. doi: 10.1093/humupd/dmv049.

26.   do Imperio GE, Bloise E, Javam M, Lye P, Constantinof A, Dunk C, et al.
Chorioamnionitis Induces a Specific Signature of Placental ABC Transporters Associated
with an Increase of miR-331-5p in the Human Preterm Placenta. Cell Physiol Biochem.
2018;45(2):591–604. doi: 10.1159/000487100.

27.   Martinelli LM, Fontes KN, Reginatto MW, Andrade CBV, Monteiro VRS, Gomes
HR, et al. Malaria in pregnancy regulates P-glycoprotein (P-gp/Abcb1a) and ABCA1
efflux transporters in the mouse visceral yolk sac. J Cell Mol Med. 2020;00:1–12. doi:
10.1111/jcmm.15682.

28.   Reginatto MW, Fontes KN, Monteiro VRS, Silva NL, Andrade CBV, Gomes HR, et
al. Effect of sub-lethal prenatal endotoxemia on murine placental transport systems
and lipid homeostasis. bioRxiv. 2020 Aug 5;2020.08.04.236745.

29.   Martinelli LM, Reginatto MW, Fontes KN, Andrade CBV, Monteiro VRS, Gomes
HR, et al. Breast cancer resistance protein (Bcrp/Abcg2) is selectively modulated by
lipopolysaccharide (LPS) in the mouse yolk sac. Reprod Toxicol 2020 Dec;98:82-91. doi:
10.1016/j.reprotox.2020.09.001

30.   Imperio GE, Javam M, Lye P, Constantinof A, Dunk CE, Reis FM, et al.
Gestational age-dependent gene expression profiling of ATP-binding cassette
transporters in the healthy human placenta. J Cell Mol Med. 2018/11/08 ed.
2019;23(1):610–8. doi: 10.1111/jcmm.13966.

31.   Vermillion MS, Lei J, Shabi Y, Baxter VK, Crilly NP, McLane M, et al. Intrauterine
Zika virus infection of pregnant immunocompetent mice models transplacental
transmission and adverse perinatal outcomes. Nat Commun. 2017;8(1):14575. doi:
10.1038/ncomms14575.

32.   Szaba FM, Tighe M, Kummer LW, Lanzer KG, Ward JM, Lanthier P, et al. Zika
virus infection in immunocompetent pregnant mice causes fetal damage and placental
pathology in the absence of fetal infection. PLoS Pathog. 2018;14(4):e1006994. doi:
10.1371/journal.ppat.1006994.

33.   Coelho SVA, Neris RLS, Papa MP, Schnellrath LC, Meuren LM, Tschoeke DA, et
al. Development of standard methods for Zika virus propagation, titration, and
purification. Journal of Virological Methods. 2017;246:65–74. doi:
10.1016/j.jviromet.2017.04.011.

34.   Connor KL, Kibschull M, Matysiak-Zablocki E, Nguyen TT-TN, Matthews SG, Lye
SJ, et al. Maternal malnutrition impacts placental morphology and transporter
expression: an origin for poor offspring growth. The Journal of Nutritional
Biochemistry. 2020;78:108329. doi: 10.1016/j.jnutbio.2019.108329.

35.   Fontes KN, Reginatto MW, Silva NL, Andrade CBV, Bloise FF, Monteiro VRS, et
al. Dysregulation of placental ABC transporters in a murine model of malaria-induced
preterm labor. Sci Rep. 2019;9(1):11488. doi: 10.1038/s41598-019-47865-3.

36.   Bloise E, Lin W, Liu X, Simbulan R, Kolahi KS, Petraglia F, et al. Impaired
Placental Nutrient Transport in Mice Generated by *in Vitro* Fertilization. Endocrinology.
2012;153(7):3457–67. doi: 10.1210/en.2011-1921

37.   Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic
and Serologic Properties of Zika Virus Associated with an Epidemic, Yap State,
Micronesia, 2007. Emerg Infect Dis. 2008 Aug;14(8):1232–9. doi:
10.3201/eid1408.080287.

38.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. Methods. 2001;25(4):402–8. doi:
10.1006/meth.2001.1262.

39.   Tschanz SA, Burri PH, Weibel ER. A simple tool for stereological assessment of
digital images: the STEPanizer: TOOL FOR STEREOLOGICAL ASSESSMENT. Journal of
Microscopy. 2011;243(1):47–59. doi: 10.1111/j.1365-2818.2010.03481.x.

40.   Sesso A, Belizário JE, Marques MM, Higuchi ML, Schumacher RI, Colquhoun A,
et al. Mitochondrial Swelling and Incipient Outer Membrane Rupture in Preapoptotic
and Apoptotic Cells. Anatomical Record (Hoboken, N.j : 2007). 2012 Oct;295(10):1647.
doi: 10.1002/ar.22553.
41.   Montalbano R, Waldegger P, Quint K, Jabari S, Neureiter D, Illig R, et al.
Endoplasmic Reticulum Stress Plays a Pivotal Role in Cell Death Mediated by the Pan-
Deacetylase Inhibitor Panobinostat in Human Hepatocellular Cancer Cells.
Translational Oncology. 2013;6(2):143-IN6. doi: 10.1593/tlo.12271
42.   Hamilton SA, Tower CL, Jones RL. Identification of Chemokines Associated with
the Recruitment of Decidual Leukocytes in Human Labour: Potential Novel Targets for
Preterm Labour. PLOS ONE. 2013 Feb 22;8(2):e56946. doi:
10.1371/journal.pone.0056946.
43.   Bloise E, Bhuiyan M, Audette MC, Petropoulos S, Javam M, Gibb W, et al.
Prenatal Endotoxemia and Placental Drug Transport in The Mouse: Placental Size-
Specific Effects. PLOS ONE. 2013 Jun 10;8(6):e65728. doi:
10.1371/journal.pone.0065728
44.   Romero R, Mazor M, Brandt F, Sepulveda W, Avila C, Cotton DB, et al.
Interleukin-1α and Interleukin-1 β in Preterm and Term Human Parturition. American
Journal of Reproductive Immunology. 1992;27(3-4):117–23. doi: 10.1111/j.1600-
0897.1992.tb00737.x.
45.   Larocca RA, Abbink P, Peron JPS, de A. Zanotto PM, Iampietro MJ, Badamchi-
Zadeh A, et al. Vaccine protection against Zika virus from Brazil. Nature.
2016;536(7617):474–8. doi: 10.1038/nature18952.
46.   Souza IN de O, Frost PS, França JV, Nascimento-Viana JB, Neris RLS, Freitas L, et
al. Acute and chronic neurological consequences of early-life Zika virus infection in
mice. Science Translational Medicine. 2018;10(444):eaar2749. doi:
10.1126/scitranslmed.aar2749.
47.   Kalagiri RR, Carder T, Choudhury S, Vora N, Ballard AR, Govande V, et al.
Inflammation in Complicated Pregnancy and Its Outcome. Am J Perinatol.
2016;33(14):1337–56. doi: 10.1055/s-0036-1582397.
48.   Barbeito-Andrés J, Pezzuto P, Higa LM, Dias AA, Vasconcelos JM, Santos TMP,
et al. Congenital Zika syndrome is associated with maternal protein malnutrition. Sci
Adv. 2020;6(2):eaaw6284. doi: 10.1126/sciadv.aaw6284.
49.   Saito S, Nakashima A, Shima T, Ito M. Th1/Th2/Th17 and Regulatory T-Cell
Paradigm in Pregnancy. American Journal of Reproductive Immunology.
2010;63(6):601–10. doi: 10.1111/j.1600-0897.2010.00852.x.
50.   Gotsch F, Romero R, Kusanovic JP, Mazaki-Tovi S, Pineles BL, Erez O, et al. The
fetal inflammatory response syndrome. Clin Obstet Gynecol. 2007;50(3):652–83. doi:
10.1097/GRF.0b013e31811ebef6.
51.   Lucas CGO, Kitoko JZ, Ferreira FM, Suzart VG, Papa MP, Coelho SVA, et al.
Critical role of CD4 + T cells and IFNγ signaling in antibody-mediated resistance to Zika
virus infection. Nature Communications. 2018;9(1):3136. doi: 10.1038/s41467-018-
05519-4.
52.   Galinsky R, Polglase GR, Hooper SB, Black MJ, Moss TJM. The consequences of
chorioamnionitis: preterm birth and effects on development. J Pregnancy.
2013;2013:412831. doi: 10.1155/2013/412831.

53.   Mor G, Cardenas I, Abrahams V, Guller S. Inflammation and pregnancy: the role
of the immune system at the implantation site. Ann N Y Acad Sci. 2011;1221(1):80–7.
doi: 10.1111/j.1749-6632.2010.05938.x.
54.   Ribeiro MR, Moreli JB, Marques RE, Papa MP, Meuren LM, Rahal P, et al. Zika-
virus-infected human full-term placental explants display pro-inflammatory responses
and undergo apoptosis. Arch Virol. 2018;163(10):2687–99. doi: 10.1007/s00705-018-
3911-x.
55.   Gown AM, Willingham MC. Improved Detection of Apoptotic Cells in Archival
Paraffin Sections: Immunohistochemistry Using Antibodies to Cleaved Caspase 3. J
Histochem Cytochem. 2002;50(4):449–54. doi: 10.1177/002215540205000401.
56.   Noronha L de, Zanluca C, Burger M, Suzukawa AA, Azevedo M, Rebutini PZ, et
al. Zika Virus Infection at Different Pregnancy Stages: Anatomopathological Findings,
Target Cells and Viral Persistence in Placental Tissues. Front Microbiol.
2018;9(2266):1–11. doi: 10.3389/fmicb.2018.02266.
57.   Lojpur T, Easton Z, Raez-Villanueva S, Laviolette S, Holloway AC, Hardy DB. Δ9-
Tetrahydrocannabinol leads to endoplasmic reticulum stress and mitochondrial
dysfunction in human BeWo trophoblasts. Reproductive Toxicology. 2019;87:21–31.
doi: 10.1016/j.reprotox.2019.04.008.
58.   Offerdahl DK, Dorward DW, Hansen BT, Bloom ME. Cytoarchitecture of Zika
virus infection in human neuroblastoma and Aedes albopictus cell lines. Virology.
2017;501:54–62. doi: 10.1016/j.virol.2016.11.002.
59.   Desuzinges-Mandon E, Arnaud O, Martinez L, Huché F, Di Pietro A, Falson P.
ABCG2 Transports and Transfers Heme to Albumin through Its Large Extracellular Loop.
J Biol Chem. 2010;285(43):33123–33. doi: 10.1074/jbc.M110.139170
60.   Ledur PF, Karmirian K, Pedrosa C da SG, Souza LRQ, Assis-de-Lemos G, Martins
TM, et al. Zika virus infection leads to mitochondrial failure, oxidative stress and DNA
damage in human iPSC-derived astrocytes. Scientific Reports. 2020;10(1):1218. doi:
10.1038/s41598-020-57914-x.
61.   Reemst K, Noctor SC, Lucassen PJ, Hol EM. The Indispensable Roles of Microglia
and Astrocytes during Brain Development. Front Hum Neurosci 2016 Nov 8;10:566.
doi: 10.3389/fnhum.2016.00566.
62.   Guitart-Mampel M, Gonzalez-Tendero A, Niñerola S, Morén C, Catalán-Garcia
M, González-Casacuberta I, et al. Cardiac and placental mitochondrial characterization
in a rabbit model of intrauterine growth restriction. Biochimica et Biophysica Acta
(BBA) - General Subjects. 2018;1862(5):1157–67. doi: 10.1016/j.bbagen.2018.02.006.
63.   Guitart-Mampel M, Juarez-Flores DL, Youssef L, Moren C, Garcia-Otero L, Roca-
Agujetas V, et al. Mitochondrial implications in human pregnancies with intrauterine
growth restriction and associated cardiac remodelling. J Cell Mol Med.
2019;23(6):3962–73. doi: 10.1111/jcmm.14282.
64.   Zhang Z, Rong L, Li Y-P. Flaviviridae Viruses and Oxidative Stress: Implications
for Viral Pathogenesis. Oxid Med Cell Longev. 2019;2019:1409582. doi:
10.1155/2019/1409582.
65.   Martínez F, Kiriakidou M, Strauss JF. Structural and Functional Changes in
Mitochondria Associated with Trophoblast Differentiation: Methods to Isolate
Enriched Preparations of Syncytiotrophoblast Mitochondria. Endocrinology.
1997;138(5):2172–83. doi: 10.1210/endo.138.5.5133.

66. Burton GJ, Yung HW, Murray AJ. Mitochondrial – Endoplasmic reticulum
interactions in the trophoblast: Stress and senescence. Placenta. 2017;52:146–55. doi:
10.1016/j.placenta.2016.04.001.
67. Holland O, Dekker Nitert M, Gallo LA, Vejzovic M, Fisher JJ, Perkins AV. Review:
Placental mitochondrial function and structure in gestational disorders. Placenta.
2017;54:2–9. doi: 10.1016/j.placenta.2016.12.012
68. Bridges CC, Zalups RK, Joshee L. Toxicological significance of renal Bcrp:
Another potential transporter in the elimination of mercuric ions from proximal
tubular cells. Toxicol Appl Pharmacol. 2015;285(2):110–7. doi:
10.1016/j.taap.2015.03.027.
69. Szilagyi JT, Gorczyca L, Brinker A, Buckley B, Laskin JD, Aleksunes LM. Placental
BCRP/ABCG2 Transporter Prevents Fetal Exposure to the Estrogenic Mycotoxin
Zearalenone. Toxicol Sci. 2019;168(2):394–404. doi: 10.1093/toxsci/kfy303.
70. Scialis RJ, Aleksunes LM, Csanaky IL, Klaassen CD, Manautou JE. Identification
and Characterization of Efflux Transporters That Modulate the Subtoxic Disposition of
Diclofenac and Its Metabolites. Drug Metab Dispos. 2019;47(10):1080–92. doi:
10.1124/dmd.119.086603.
71. Mao Q, Unadkat JD. Role of the breast cancer resistance protein (BCRP/ABCG2)
in drug transport--an update. AAPS J. 2015;17(1):65–82. doi: 10.1208/s12248-014-
9668-6.
72. Lye P, Bloise E, Javam M, Gibb W, Lye SJ, Matthews SG. Impact of Bacterial and
Viral Challenge on Multidrug Resistance in First- and Third-Trimester Human Placenta.
The American Journal of Pathology. 2015;185(6):1666–75. doi:
10.1016/j.ajpath.2015.02.013.
73. Girard S, Sebire G. Transplacental Transfer of Interleukin-1 Receptor Agonist
and Antagonist Following Maternal Immune Activation. American journal of
reproductive immunology (New York, NY : 1989). 2016;75(1):8–12. doi:
10.1111/aji.12444
74. Bloise E, Petropoulos S, Iqbal M, Kostaki A, Ortiga-Carvalho TM, Gibb W, et al.
Acute Effects of Viral Exposure on P-Glycoprotein Function in the Mouse Fetal Blood-
Brain Barrier. Cell Physiol Biochem . 2017;41(3):1044-1050. doi: 10.1159/000461569.