ZIKV disrupts fetal outcome, placental ultrastructure and drug transporter

expression in mice

Cherley Borba Vieira de Andrade[1], Victoria Regina de Siqueira Monteiro[1], Sharton

Vinicius Antunes Coelho[2], Hanailly Ribeiro Gomes[1], Ronny Paiva Campos de Sousa[1],

Veronica Muller de Oliveira Nascimento[1], Flavia Fonseca Bloise[1], Stephen

Matthews[3,4,5,6], Enrrico Bloise[7], Luciana Barros de Arruda[2], Tania Maria Ortiga-

Carvalho[1]*.

[1]Institute of Biophysics Carlos Chagas Filho, Federal University of Rio de Janeiro, Rio

de Janeiro, RJ, Brazil.

[2] Institute of Microbiology Paulo de Góes, Federal University of Rio de Janeiro, Rio de

Janeiro, RJ, Brazil.

[3] Department of Physiology, Faculty of Medicine, University of Toronto, Toronto,

Canada.

[4] Department of Obstetrics & Gynecology, Faculty of Medicine, University of Toronto,

Toronto, Canada.

[5] Department of Medicine, Faculty of Medicine, University of Toronto, Toronto,

Canada.

[6] Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Toronto, Canada.

[7] Department of Morphology, Federal University of Minas Gerais, Belo Horizonte, BH,

Brazil.

* Correspondence:

Tania Maria Ortiga-Carvalho (TMO-C)

taniaort@biof.ufrj.br

Key words: ZIKV, placenta, P-glycoprotein (P-gp) breast cancer resistance protein

(BCRP), Abca1, Abcg1, ultrastructure, cytokine, chemokine

Abstract

Congenital Zika virus (ZIKV) infection can induce fetal brain abnormalities. Here, we

investigated whether maternal ZIKV infection may affect placental physiology and

metabolic transport potential, and impact the fetal outcome, regardless of viral presence

in the fetus at term. Low (103 PFU-ZIKVPE243; low-ZIKV) and high ($5x10^7$ PFU-

ZIKVPE243; high-ZIKV) virus titers were injected into immunocompetent (ICompetent

C57BL/6) and immunocompromised (ICompromised A129) mice at gestational day

(GD)12.5 for tissue collection at GD18.5 (term). High-ZIKV elicited fetal death rates of

66% and 100%, whereas low-ZIKV induced fetal death rates of 0% and 60% in

C57BL/6 and A129 dams, respectively. All surviving fetuses exhibited intrauterine

growth restriction (IUGR) and decreased placental efficiency. High-ZIKV infection in

C57BL/6 and A129 mice resulted in virus detection in maternal spleens and placenta,

but only A129 fetuses presented virus RNA in the brains. Nevertheless, pregnancies of

both strains produced fetuses with decreased head sizes ($p<0.05$). Low-ZIKV-A129

dams had higher IL-6 and CXCL1 levels ($p<0.05$) and their placentas showed increased

CCL-2 and CXCL-1 contents ($p<0.05$). In contrast, low-ZIKV-C57BL/6 dams had an

elevated CCL2 serum level, and increased type I and II IFN expression in the placenta.

Notably, increased apoptotic rates, less abundant microvilli and mitochondrial

degeneration were evidenced in the placental labyrinth (Lz) of ICompromised and high-

ZIKV-ICompetent mice, but not in low-ZIKV-C57BL/6. In addition, decreased

placental expression of the drug P-glycoprotein (P-gp), breast cancer resistance protein

(BCRP) and the lipid ABCA1 transporters was detected in all ZIKV-infected groups,

but BCRP and ABCA-1 were only reduced in ICompromised and high-ZIKV

ICompetent mice. Our data indicate that gestational ZIKV infection triggers specific

proinflammatory responses and affects placental turnover and transporter expression, in

a way dependent on virus concentration and maternal immune status. Placental damage

may impair proper fetal-maternal exchange function and fetal growth/survival, likely

contributing to congenital Zika syndrome.

1. Introduction

Congenital Zika virus (ZIKV) infection can be associated with adverse

pregnancy outcomes. As such, neonates born from ZIKV-positive pregnancies may

develop severe neurological abnormalities, placental pathologies and intrauterine

growth restriction (IUGR), among other complications (1). ZIKV vertical transmission

has become a major public health issue worldwide, especially in Brazil, where more

than 200,000 ZIKV-positive cases have been confirmed, and over 2,000 congenital

microcephaly births have been reported (2–6). These numbers represented a 20-fold rise

in the incidence of congenital microcephaly in Brazil during the years of the ZIKV

pandemic, with similar increases reported elsewhere in Latin America (2,3,7).

Importantly, while the ZIKV pandemic is currently thought to be controlled, evidence

points to a possible silent ZIKV spread across the Americas (8,9), highlighting the need

for improved knowledge of the possible routes of ZIKV vertical transmission and its

association with disruptive inflammatory and developmental phenotypes and the need

for new avenues of prevention and treatment.

Studies have investigated the possible pathways involved in ZIKV vertical

transmission. Miranda and colleagues (10) showed that in humans, ZIKV infection

changed the pattern of tight junction proteins, such as claudin-4, in

syncytiotrophoblasts. Jurado et al. (2016) suggested that the migratory activities of

Hofbauer cells (feto-placental macrophages) could help disseminate ZIKV to the fetal

brain (11). Other recent studies have shown that placental villous fibroblasts,

cytotrophoblasts, endothelial cells and Hofbauer cells are permissive to ZIKV, and

placentae from ZIKV-infected women had chorionic villi with a higher mean diameter

(11–14). However, at present, there is limited information on the precise mechanism of

maternal-fetal ZIKV transmission.

Many mouse models have been developed to identify how ZIKV overcomes

placental defenses. First, limited information was obtained due to the apparent inability

of the virus to infect wild-type (WT) mice (15). ZIKV NS5 targets the interferon

signaling pathway in humans but not in mice (16). As such, WT mice develop no clear

evidence of clinical disease (16,17) and are of limited use in modeling the disease.

However, mice lacking an interferon signaling response show evidence of disease and

have been widely used to investigate ZIKV infection during pregnancy (8,16,18).

The interferon system, especially type III interferon, is one of the key

mechanisms of host defense and viral targets for immune evasion (19). Type III

interferons have a role in protection against ZIKV infection in human

syncytiotrophoblasts from term placenta (20). Luo et al. have shown that the inhibition

of Toll-like receptors 3 and 8 inhibits the cytokine output of ZIKV-infected trophoblasts

(21). In addition, viral replication coincides with the induction of proinflammatory

cytokines such as IL-6. This cytokine has a crucial role in inflammation and affects the

homeostatic processes related to tissue injury and activation of stress-related responses

(22,23). ZIKV infection can trigger an inflammatory response with IL-6 release (11,24).

Maternal infection has profound effects on placental permeability to drugs and

environmental toxins. Changes in the expression and function of specific ABC

transporters in the placenta and yolk sac following infective and inflammatory stimuli

have been demonstrated (25–29). ABC transporters are efflux transporters that control

the biodistribution of several endogenous and exogenous substrates, including

xenobiotics (antiretrovirals, synthetic glucocorticoids), steroid hormones (estrogens, androgens), nutrients (folate, cholesterol) and immunological factors (chemokines and cytokines) within the maternal-fetal interface (30). The best described ABC transporters in the placenta are the multidrug resistance P-glycoprotein (P-gp), the breast cancer resistance protein (Bcrp) and the lipid Abca1 and Abcg1 transporters. P-gp and BCRP transporters are responsible for preventing fetal accumulation of xenobiotics and environmental toxins that may be present in the maternal circulation, whereas ABCA1 and ABCG1 control the placental exchange of cytotoxic oxysterol and lipid permeability throughout pregnancy; therefore, they play an important role in fetal protection and placental lipid homeostasis (25).

Despite a limited number of studies showing ZIKV infection in immunocompetent mice, it was previously demonstrated that intrauterine inoculation of high virus titer resulted in decreased fetal viability, with worse outcomes following infection in early gestation (31). In another report, intravenous infection at very early embryonic day resulted in fetal demise even though the virus was not found in the fetal compartment in most treated animals (32). In the present study, we hypothesize that maternal exposure to ZIKV affects placental function, including placental ultrastructure and ABC transporter (P-gp, Bcrp, Abca1 and Abcg1) protein expression, even in the absence of vertical transmission and that these effects are dependent on viral infective titers and maternal immune status.

2. Materials and methods

2.1 Virus preparation and storage

The Brazilian $ZIKV_{PE243}$ (GenBank ref. number KX197192) strain was isolated from a febrile case during the ZIKV outbreak in the state of Pernambuco, Brazil and was kindly provided by Dr Ernesto T. Marques Jr. (Centro de Pesquisa Aggeu

Magalhães, FIOCRUZ, PE). Viruses were propagated in C6/36 cells with viral titers

determined by plaque assays in Vero cells, as previously described (33). Supernatants of

noninfected C6/36 cells cultured under the same conditions were used as mock controls.

2.2 Animal experimentation and study design

Two mouse strains were used in the study: ICompetent C57BL/6 and

ICompromised (type 1 *Ifnr*-deficient) A129 strains. Since we were unable to

consistently produce viable pregnancies by mating A129 males and females in our

experimental settings, we mated A129 females (n=15) with C57BL/6 males (n=4) to

produce ICompromised C57BL6/A129 pregnancies, whereas ICompetent C57BL/6

pregnancies were obtained by mating male (n=6) and female (n=35) C57BL/6 mice (8-

10 weeks old). Animals were kept in a controlled temperature room (23°C) with a

light/dark cycle of 12 hours and *ad libitum* access to water and food. After detecting the

proestrous/estrous phase by vaginal cytology, copulation was confirmed by

visualization of the vaginal plug and considered gestational day 0.5 (GD0.5). Maternal

weight was monitored for confirmation of pregnancy; thus, females were weighed on

GD0.5 and GD12.5, and females with a weight gain greater than 3 g were considered

pregnant and entered randomly in the experimental groups. Experimental protocols

were approved by the Animal Care Committee of the Health Sciences Center, Federal

University of Rio de Janeiro (CEUA-036/16 and 104/16) and registered within the

Brazilian National Council for Animal Experimentation Control.

On GD12.5, pregnant mice (ICompetent and ICompromised pregnancies) were

injected with a single intravenous (i.v.) titer of ZIKV or mock control. ICompetent and

ICompromised pregnant mice were randomly subdivided into three experimental

groups: the mock (control) group, which received supernatants of noninfected C6/36

cell injection (ICompetent mock and ICompromised mock); the high-ZIKV-titer group,

inoculated with 5×10^7 plaque-forming units (PFU) of ZIKV$_{PE243}$ (ICompetent high and

ICompromised high); and the ZIKV low-titer group, injected with 10^3 PFU of

ZIKV$_{PE243}$ (ICompetent low and ICompromised low).

On the morning of GD18.5, all animals were euthanized with a sodium

phenobarbital overdose of 300 mg/kg. Maternal blood was collected by cardiac

puncture, centrifuged (10 min, 4000 g) and stored at -20°C. The maternal brain and

spleen, all placentae and all fetuses were dissected, collected and weighed, followed by

fetal head isolation and measurement. The three placentae closest to the mean weight in

a litter were selected for further analysis and cut in half using umbilical cord insertion as

a reference (34–36). One-half of the placental disk was frozen in liquid nitrogen for

qPCR, and the other half was fixed overnight in buffered paraformaldehyde (4%,

Sigma-Aldrich, Brazil) for ultrastructural and protein expression/localization analysis.

Matched fetal heads, maternal brains and spleens were frozen in liquid nitrogen for

qPCR.

2.3 ZIKV RNA quantification by qPCR

ZIKV load was evaluated in maternal blood, brains and spleens and in the

placentae and fetal heads. Brains, spleens, placentae and fetal heads were macerated in

RPMI medium (Gibco™ RPMI 1640 Medium) normalized by the ratio of 0.2 mg of

tissue to every 1 µl of medium and plotted per gram of tissue. The macerated volume

was centrifuged at 4500 g for 5 min to remove tissue residues, and then 500 µl of the

centrifuged volume was used for RNA extraction using 1 mL of TRIzol reagent (Life

Technologies, Thermo Fischer, USA). Treatment with DNase I (Ambion, Thermo

Fischer, USA) was performed to prevent contamination by genomic DNA. cDNA was

synthesized using a cDNA High Capacity Kit (Applied Biosystems, Thermo Fischer,

USA) according to the manufacturer's instructions by subjecting the samples to the

following cycle: 25°C for 10 min, 37°C for 120 min and 85°C for 5 min. qPCR was

performed using a StepOnePlus Real-Time PCR system and TaqMan Master Mix

Reagents (Applied Biosystems, Thermo Fischer, USA) using primers and probes

specific for the protein E sequence (37). Samples were then subjected to the following

cycle: 50°C for 2 min, 40x (95°C for 10 min, 95°C for 15 sec, 60°C for 1 min).

2.3.1 qPCR

The placenta was macerated in 1.5 mL of TRIzol reagent (Life Technologies,

Thermo Fischer, USA). RNA extraction was performed following the manufacturer's

protocol. cDNA was prepared using Power SYBR Green PCR Master Mix (Life

Technologies, Thermo Fisher, USA). The reaction was carried out for selected genes

using intron-spanning primers (Table 1) and the StepOnePlus Real-Time PCR system

(Life Technologies, Thermo Fischer, USA). Samples were subjected to the following

cycle: 95°C for 10 min followed by 40 amplification cycles consisting of DNA

denaturation for 30 sec at 95°C and annealing of primers for 30 sec at 60°C. The

threshold cycles (Ct) for each gene of interest and for the reference gene glycerol 3-

phosphate dehydrogenase (*Gapdh*) and RNA Polymerase II Subunit A (*Polr2a)* were

determined. The relative expression for each gene was calculated using $2^{-\Delta\Delta CT}$ (38) and

graphically expressed as the fold-increase. The efficiency was calculated using the

standard curve method. The melting curves were analyzed for each sample.

2.4 Detection of cytokines and chemokines in the maternal serum and placenta

Initially, placental tissue was homogenized in extraction buffer (50 mM Tris,

150 mM NaCl, 1X Triton, 0.1% SDS, 5 mM EDTA, 5 mM NaF, 50 mM sodium

pyrophosphate, 1 mM sodium orthovanadate, pH 7.4) containing complete protease

inhibitor cocktail (Roche Applied Science, Germany) with TissueLyser LT (Qiagen,

Germany). The protein concentration of each sample was analyzed using the Pierce™

BCA Protein Assay Kit (Thermo Scientific, USA) according to the manufacturer's

instructions. Analysis of the cytokines IL-6 and IL1-β and the chemokines monocyte

chemoattractant protein-1 (MCP-1/CCL2) and chemokine (C-X-C motif) ligand 1

(CXCL1) in maternal serum and placenta was performed with the MILLIPLEX MAP

Mouse Cytokine/Chemokine Magnetic Bead Panel – Immunology Multiplex Assays

(MCYTOMAG-70K, Merck Millipore, Germany), following the manufacturer's

recommendations. The plate with samples and magnetic beads was analyzed on a

MAGPIX® System (Merck Millipore, Germany). The analyses were performed using

the Luminex xPonent® for MAGPIX® v software. 4.2 (Luminex Corp., USA). For

each reaction well, the MAGPIX Luminex® platform reports the median fluorescence

intensity (MFI) for each of the analytes in the samples. The levels of each analyte were

then calculated against the standard curve. The ratio between the value obtained and the

protein quantification for each sample was determined and plotted.

2.5 Virus titration by plaque assays

Blood from mock- and $ZIKV_{PE243}$-infected mice was collected from the base of

the tail at 4 hours, 48 hours and 144 hours following the appropriate treatments and

subsequently centrifuged at 400 g for 30 min for plasma separation. Samples obtained at

different periods post infection were titrated by a plaque assay. Vero cells (obtained

from ATCC® CCL81™) (African green monkey kidney epithelial cell line) were plated

228 in 24-well plates at 4×10^4 cells per well in DMEM (GIBCO, Thermo Fisher, USA)

229 supplemented with 5% FBS (GIBCO, Thermo Fisher, USA) and 1% gentamicin (10

230 μg/ml) (GIBCO, Thermo Fisher, USA) and cultured overnight for complete adhesion at

231 37°C with 5% CO_2. Then, the medium was removed, and the cells were washed with 1x

232 PBS and incubated with serial (base 10) dilutions of virus in FBS-free medium. After

233 90 min of incubation under gentle shaking, the medium was removed, and the cells

234 were washed with 1x PBS and cultured with 1.5% carboxymethylcellulose (CMC)

235 supplemented with 1% FBS (GIBCO, Thermo Fisher, USA). After 5 days, the cells

236 were fixed overnight with 4% formaldehyde and stained with 1% crystal violet in 20%

237 methanol (ISOFAR, Brazil) for 1 hour. Plaques were counted, and the virus yield was

238 calculated and expressed as plaque-forming units per milliliter (PFU/ml).

240 ***2.6 Histological, immunohistochemistry and TUNEL analyses of the placenta***

241  Placental fragments were fixed overnight and subjected to dehydration

242 (increasing ethanol series, ISOFAR, Brazil), diaphanization with xylol (ISOFAR,

243 Brazil) and paraffin (Histopar, Easypath, Brazil). Sections (5 μm) were prepared using

244 the Rotatory Microtome CUT 5062 (Slee Medical GmbH, Germany) and subjected to

245 immunohistochemistry and TUNEL analyses.

246  For immunohistochemistry, blocking endogenous peroxidase was performed

247 with 3% hydrogen peroxide diluted in PBS, followed by microwave antigenic recovery

248 in Tris-EDTA (pH=9) and sodium citrate (pH=6) buffers (15 min for Tris-EDTA buffer

249 and 8 min for citrate buffer). Sections were washed in PBS + 0.2% Tween and exposed

250 to 3% PBS/BSA for 1 hour. Sections were then incubated overnight at 4°C with the

251 following primary antibodies: anti-Ki-67 (1:100 – [M3064]; Spring Bioscience, USA),

anti-P-gp (1:500 – Mdr1[sc-55510]; Santa Cruz Biotechnology, USA), anti-Bcrp (1:100

– Bcrp [MAB4146]; Merck Millipore, USA), anti-Abcg1 (1:100 – [PA5-13462];

Thermo Fisher Scientific, USA) or anti-Abca1 (1:100 – [ab18180]; Abcam Plc, UK).

The next day, sections were incubated with the biotin-conjugated secondary antibody

SPD-060 (Spring Bioscience, USA) for 1 hour at room temperature. Three washes were

performed with PBS + 0.2% Tween followed by incubation with streptavidin (SPD-060

- Spring Bioscience, USA) for 30 min. Sections were stained with 3,3-diamino-

benzidine (DAB) (SPD-060 - Spring Bioscience, USA), counterstained with

hematoxylin (Proquímios, Brazil), dehydrated, diaphanized and mounted with coverslip

and entellan (Merck, Germany).

For analysis of apoptotic nuclei, terminal deoxynucleotidyl transferase dUTP

nick-end labeling (TUNEL) staining was performed using the ApopTag® In Situ

Peroxidase Detection Kit (S7100, Merck Millipore, USA) according to the

manufacturer's recommendations and as previously described (35). All negative controls

were performed with the omission of the primary antibody.

Image acquisition was performed using a high-resolution Olympus DP72

(Olympus Corporation, Japan) camera coupled to an Olympus BX53 light microscope

(Olympus Corporation, Japan). For nuclear quantification of Ki-67 and TUNEL

immunolabeling, Stepanizer software (39) was used. For this analysis, we evaluated 15

images from different random fields of the Lz and Jz for each animal, in a total of five

animals from each ICompetent group and three animals from each ICompromised

group. A total of 360 digital images per placental region (Lz and Jz) were evaluated in

each experimental group. Total numbers of immunolabeled Ki-67 or TUNEL nuclei

from each image were normalized by the total image area to obtain an index of

estimated numbers of proliferative and apoptotic nuclei in the whole histological section

analyzed. Analysis was undertaken by two investigators blinded to the treatment.

Quantification of P-gp, Bcrp, Abca1 and Abcg1 staining was performed using

the Image-Pro Plus, version 5.0 software (Media Cybernetics, USA) mask tool. The

percentage of viable tissue area was considered upon exclusion of negative spaces. A

total of 360 digital images per placental region (Lz and Jz) were evaluated in each

experimental group. Analysis was undertaken by two investigators blinded to the

treatment.

2.7 Transmission electron microscopy (TEM)

Sections of placental Lz and Jz were fixed in paraformaldehyde (4%, Sigma-Aldrich,

Brazil) for 48 hours, postfixed with osmium tetroxide (Electron Microscopy Sciences,

USA) and potassium ferrocyanide (Electron Microscopy Sciences, USA) for 60 min and

dehydrated with an increasing series of acetone (30%, 50%, 70%, 90% and two of

100%) (ISOFAR, Brazil). Sections were subsequently embedded with EPOXI resin

(Electron Microscopy Sciences, USA) and acetone (1:2, 1:1 and 2:1, respectively).

After polymerization, ultrafine sections (70 nm) were prepared (Leica Microsystems,

USA) and collected into 300 mesh copper grids (Electron Microscopy Sciences, USA).

Tissue was contrasted with uranyl acetate and lead citrate and visualized using a JEOL

JEM-1011 transmission electron microscope (JEOL, Ltd., Akishima, Tokyo, Japan).

Digital micrographs were captured using an ORIUS CCD digital camera (Gatan, Inc.,

Pleasanton, California, EUA) at 6000× magnification. An overall qualitative analysis of

Lz and Jz from different groups was performed by investigating the ultrastructural

characteristics of the mitochondria as well as the ER cisterns. The qualitative evaluation

consisted of analyzing the disruption of the mitochondrial membranes, mitochondrial

morphology, preservation of mitochondrial cristae and matrix intensity (40).

Ultrastructural analysis of nuclear morphology and the presence of microvilli in

trophoblast sinusoidal giant cells was also undertaken. The analysis of the cisterns of

the ER was performed by evaluating the dilation of their lumen (41).

2.8 Statistical analysis

GraphPad Prism 8 software (GraphPad Software, Inc., USA) was used for

statistical analysis. The D'Agostino & Pearson normality test was used to evaluate

normal distribution, and outliers were identified using the Grubbs test. Data are

expressed as the mean±SEM or individual values. One-way ANOVA followed by

Tukey's posttest was used for comparisons between different inbred groups, whereas

Student's t-test or nonparametric Mann-Whitney test was performed between the

outbred groups. Differences were considered significant when $p<0.05$. Pregnancy

parameters were evaluated using the mean value of all fetuses and placentae in a litter

per dam and not the individual conceptus, i.e., the mean value. "n" represents the

number of litters in Figures 1 and 2 (34–36).

3. Results

3.1 Weight gain across pregnancy is dependent on maternal immune status in ZIKV-

infected mice.

To determine the effect of ZIKV infection on fetal and placental phenotypes at

term (GD18.5), we infected ICompetent C57BL/6 and immunocompromised

(ICompromised) A129 mice with ZIKV at GD12.5 (Figure 1A). Given the very distinct

susceptibility of C57BL/6 and A129 mice to ZIKV, systemic infection models were

established by injecting high (5×10^7 PFU) and low (10^3 PFU) virus inoculum titers. As

shown in Table 2, ICompetent C57BL/6 mice from all groups and ICompromised A129

dams inoculated with mock and low exhibited higher maternal weight at GD18.5 than at

GD12.5 and GD0.5 (p<0.05). On the other hand, ICompromised A129 mice presented

significant weight loss at GD18.5, despite showing an increase at GD12.5 in relation to

GD0.5

3.2 Immunocompetent and immunodeficient mice have distinct term placental and

fetal phenotypes in response to high and low ZIKV titer challenges in mid-pregnancy.

335   The fetuses from C57BL/6 dams and sires were called ICompetent. The fetuses

from the mates of A129 dams and C57BL/6 sires were called ICompromised. High-

ZIKV ICompetent mice exhibited 34% fetal loss, whereas high-ZIKV-A129 mice had

100% fetal loss (Figure 1B). In the low-ZIKV groups, C57BL/6 mice had no (0%) fetal

death, while A129 mice exhibited a 42% fetal death rate (Figure 1B). Fetal and fetal

head sizes were decreased in A129 mice compared to those in C57BL/6 dams infected

with the high ZIKV titer (p=0.05; Figure 1 C-F). However, no changes in fetal weight

and fetal head sizes were observed when the mice were infected with the lower ZIKV

titer (Figure 1 C-F).

3.3 ZIKV is detected in the fetal brain of ICompromised, but not ICompetent mice.

346   ZIKV RNA was detected in the spleens of pregnant ICompetent C57BL/6 mice

inoculated with the highest ZIKV titer, confirming acute systemic infection. Viral RNA

was also detected in the majority of placentae of those mice (Figure 2A-B) but not in

the maternal and fetal C57BL/6 brains (Figure 2A-B), suggesting that the virus was not

transmitted to the fetuses. Although low ZIKV inoculation resulted in virus detection in

the spleens of pregnant C57BL/6 mice, infection was not evidenced in the placentae or

fetal brains. In contrast, ZIKV RNA was detected in all analyzed organs from

ICompromised A129 dams, including the maternal brain and spleen and the placenta

and fetal brain (Figure 2A-B). Viremia in maternal serum was evaluated at 4, 48 and

144 hours after infection. Within 4 hours, the presence of the virus was verified in the

serum (high ICompetent = 637.5 PFU/mL, low ICompetent = 740 PFU/mL and low

ICompromised = 325 PFU/mL), indicating that the virus was correctly inoculated.

Afterwards, ZIKV RNA was detected in ICompromised dams at 48 and 144 hours

postinoculation but not in ICompetent dams.

3.4 ZIKV infection induces distinct systemic and placental inflammatory responses in

ICompetent and ICompromised mice.

The maternal serum and placental protein levels of specific cytokines and chemokines

related to fetal death and preterm delivery (42–44) were evaluated to probe whether

midgestation ZIKV infection would induce a maternal inflammatory response at term in

our two distinct models. Since A129 infected with 10^7 ZIKV-PFU showed 100% fetal

loss, we proceeded using 10^7 PFU inoculation in C57BL/6 and 10^3 PFU inoculation in

both C57BL/6 and A129 mice.

CCL2 was elevated in the serum of low-ZIKV ICompetent mice, but no other

alteration was systemically detected in any ICompetent mice at this time point (Figure

3A-B). On the other hand, ICompromised dams showed significantly increased CXCL1

and IL-6 levels in the serum, and a strong trend for an enhancement in CCL2 (Figure

3C). Analysis of cytokines and chemokines expression at the placenta demonstrated that

CXCL1 and CCL2 chemokines were also upregulated in the ICompromised, but not of

ICompetent mice (Figure 3 D-F). Surprisingly, IL-6 protein expression was augmented

in some of the low-ZIKV ICompetent mice (58%; p<0.05), but not in high-ZIKV nor

ICompromised mice (Figure 3D-F).

We also assessed the placental mRNA expression of a range of cytokines related

to placental infective responses: Ifn-γ, Il-6, Ifn-1, Tnf-α, Il-1β, and Il-10. All the

infected mice showed a significantly increase in TNF- α mRNA expression, with higher

levels detected in ICompetent mice (Figure 3G-I). The low-ZIKV ICompetent mice

presented a modest, but significant type I IFN expression, which was not detected in the

high-ZIKVgroup (Figure 3G). Additionally, both low-ZIKV infected groups

(ICompetent and ICompromised) presented increased Ifn-γ mRNA expression (Figure

3G and 3I, p< 0.05). Interestingly, placental Il-6 mRNA levels were only elevated in

ICompromised pregnancies compared to the mock (p=0.05) (Figure 3I). Il-1b and IL-10

remained unchanged in all groups analyzed (Figure 3 G-I).

3.5 ZIKV affects placental proliferation, apoptosis and ultrastructure in a viral load-

and maternal immune status-dependent manner.

We did not observe changes in placental weight or in the fetal:placental weight

ratio in any of the groups investigated (data not shown). However, since we detected the

presence of ZIKV RNA in the placentas, we investigated the cellular proliferation (Ki-

67^+ cells) and apoptotic ratio in the labyrinth zone (Lz) and junctional zone (Jz) of the

mouse placenta. Increased Ki-67 staining was observed in the Lz of high- and low-ZIKV-treated ICompetent animals compared to mock animals (Figure 4A, p=0.005 and p=0.015), whereas the Lz apoptotic ratio was increased in high- and decreased in low-ZIKV dams (p=0.01, Figure 4B). A similar apoptotic pattern was detected in the Jz of ICompetent pregnancies (p=0.008 and p=0.004, respectively; Figure 5B), although no differences were observed in Ki67 staining (Figure 5A). In contrast, in ICompromised dams, Lz Ki-67 staining was decreased (p=0.001; Figure 4C), while the apoptotic reaction was increased in ZIKV-infected animals (p=0.05; Figure 5D). Jz from ICompromised offspring exhibited increased Ki-67 labeling and no differences in the apoptotic reaction (p=0.001; Figure 5 C-D).

3.6 Placental ultrastructure is differently impacted by high- and low-titer ZIKV infection in ICompetent and ICompromised strains.

Ultrastructural analyses in the Lz of ICompetent and ICompromised-mock animals detected sinusoidal trophoblastic giant cells exhibiting regular microvilli, euchromatic nuclei, preserved mitochondrial ultrastructure and regular narrow ER cisternae (Figure 6A and 6D). In sharp contrast, high-ZIKV ICompetent (Figure 6B) and low-ZIKV ICompromised infected placentae (Figure 6E) showed fewer villi in the sinusoidal giant trophoblastic cells, degenerated mitochondria, granular ER with dilated cisterns and euchromatic nuclei. The sinusoidal giant trophoblastic cells from the low-ZIKV ICompetent (Figure 6C) also had fewer villi and degenerated mitochondria than the mock placentae, but there and had no effect on the ER or euchromatic nuclei.

The Jz of mock ICompetent and ICompromised (Figure 7A and 7D) and of low-ZIKV ICompetent (Figure 7C) placentae exhibited euchromatic nuclei, with evident

heterochromatin, preserved mitochondria and narrow cisternae granular ER. High-

ZIKV ICompromised and low-ZIKV ICompromised placentae had degenerated

mitochondria, granular ER with dilated cisterns and euchromatic nuclei (Figure 7B and

7E).

3.7 ZIKV differentially affects placental expression of drug and lipid ABC

transporter systems.

Evaluation of key ABC transporters in the Lz of mock and ZIKV-infected

ICompetent and ICompromised placentae revealed that immunolabeling of the drug P-

gp and Bcrp efflux transporter systems was primarily present at the cellular membranes

of the sinusoidal trophoblastic giant cells, with diffuse cytoplasmic Bcrp staining.

Labeling of the Abca1 and Abcg1 lipid efflux transporters was moderately and

heterogeneously distributed within the Lz. Less Lz-P-gp was observed in ICompetent

mice infected with both high- and low-ZIKV infective regimens, in comparison to

mock-treated animals   (p=0.001 and p=0.002, respectively; Figure 8A), whereas Bcrp

and Abca1 staining was only reduced was observed in high-ZIKV-infected mice

(p=0.003 and p=0.004, respectively; Figure 8B-C). P-gp, Bcrp and Abca1 transporter

immunostaining was downregulated in ICompromised low ZIKV-treated animals

(p=0.001, p=0.05 and p=0.05, respectively; Figure 9A-C). No changes in Abcg1 were

observed in any of the experimental groups (Figure 8D and Figure 9D).

Next, the impact of ZIKV on ABC transporters in the Jz layer (structural and

endocrine layers of the mouse placenta) was assessed. P-gp and Bcrp were

predominantly localized at the cellular membranes of spongiotrophoblast cells, whereas

Abca1 and Abcg1 exhibited membrane and cytoplasmic staining. P-gp staining was

decreased in Jz cells from the low-ZIKV ICompromised placentae (p=0.006), with no

other alterations observed (Figure 10A-D and Figure 11A-D).

4. Discussion

In this study, we investigated several fetuses and placentas feature at term

(GD18.5) in ICompetent (C57BL/6) and ICompromised (A129) mice exposed to ZIKV

at mid-pregnancy (GD12.5). Fetal survival rates, systemic and placental inflammatory

responses, placental ultrastructure and cell turnover, as well as the expression of key

drug (P-gp and Bcrp) and lipid (Abca1) efflux transporter systems in the placenta, were

consistently impacted by ZIKV in both strains. The magnitude of the effects was is

clearly related to the infective titer (high and low) of ZIKV and maternal immune status

(ICompetent-C57BL/6 x and ICompromised-A129) and fetal alterations were not

exclusively dependent on the virus detection in the fetuses.

Infection of ICompetent mice with ZIKV did not result in viremia in the initial

postinoculation phase, although viral RNA was detected in the maternal spleen in both

the high- and low-ZIKV groups, confirming systemic infection. This is consistent with a

previous report (45). Our data demonstrate that pregnant ICompetent C57BL/6 mice

were more susceptible to higher ZIKV than to lower ZIKV infective titers. Since viral

RNA was only detected in the placentae of high ZIKV, and fetal survival rates and

weight were impacted to a greater extent in those mice. Strikingly, even though the

virus was not present in the fetal brain (at least at term),fetal and fetal head weights

were lower in the high-ZIKV titer regimen, suggesting that high infective viral load in

mid-pregnancy, even in ICompetent individuals, can induce IUGR and lower fetal head

weight despite a lack of transmission to the fetal brain (46). On the other hand,

ICompromised placentae and fetal brains had detectable viral transcripts, with no

changes in weight, which is consistent with previous data (18). In fact, in our models,

the presence (ICompromised) or the absence (ICompetent) of the virus in the fetal brain

did not correspond to fetal head size (decreased in only high ICompetent). The data

from ICompetent and ICompromised placentae demonstrate how important the maternal

immunological status is to control viremia, fetal survival and accessibility of the virus

into the fetal brain. The reason for the reduction in fetal brain size in C57BL/6 mice in

the absence of fetal brain infection requires further investigation. It is possible that fetal

brains of the ICompetent mice may have been exposed to ZIKV earlier in pregnancy

when viremia was present in the maternal blood, and this may have severely

compromised brain development.

A distinct inflammatory profile was also detected in the three analyzed groups.

At the protein level, low-ZIKV-ICompromised dams exhibited increased maternal IL-6

and CXCL-1 and placental CCL-2 and CXCL-1, whereas low-ZIKV-ICompetent dams

had increased maternal CCL2 and placental IL-6 levels. CCCL-2 and CXCL-1 are

related to fetal death and preterm delivery (42–44) and could be associated with

pronounced fetal injure detected upon ICompromised pregnancy. In addition, it was

previously demonstrated (47) that IL-6 is related to fetal response syndrome,

characterized by activation of the fetal immune system. This syndrome is known to

increase fetal morbidity and affect several organs, such as the adrenal gland, brain and

heart (31,47–50). At the mRNA level, IL6 expression was only detected in

ICompromised placentae at term and may indicate a sustained harmful response in these

mice until term. IFN signaling pathway may be triggered by ZIKV (16) and represents

one of the key mechanisms of host defense and viral targets for immune evasion (19),

but we did only detected a slight increase in type I IFN in low-ZIKV ICompentent mice

at term. However, we cannot rule out the possibility that these cytokines might have

been produced earlier. IFN-γ expression was significantly enhanced in both ICompetent

and ICompromised low-ZIKV derived placentas, but not in the high-ZIKV infected

mice. Although we could not assess cytokine expression in high-ZIKV ICompromised

placentas, one may extrapolate that low-ZIKV infection could result in stimulation of

IFN-γ-producing cells, which has been previously showed to be protective for ZIKV-

infected mice (51).

Both ICompromised and ICompetent mice showed increased expression of

placental *Tnf-α* mRNA, which has been demonstrated to be   is directly related to

placental damage, abortion and premature birth (REF). In addition increased TNF-α

response is relatated to impaired placental hormone production and trophoblastic

invasion and increased apoptosis in pregnancy (52,53). Although we did not assessed

TNF-α protein levels in the placenta and maternal blood, this response could be

implicated in the overall damage detected.

Although differences in placental weight were not observed, ZIKV infection

mid-pregnancy had a profound effect on placental cellular turnover, dependent on titer,

strain and/or placental compartment. Lz is responsible for fetal and maternal nutrient,

gas and waste exchange, while Jz provides structural support, nutrient storage and

hormone synthesis (34). ZIKV induced a consistent increase in Lz proliferation in all

groups. However, the Lz apoptotic rate was increased only in the high-ZIKV-

ICompetent and ICompromised groups and decreased in low-ZIKV-ICompetent mice.

The mechanisms underlying these differences are unknown but may be related to the

distinct maternal and placental proinflammatory responses and/or to the direct effect of

the virus on the placenta (54). Increased Lz apoptosis in the high-ICompetent-ZIKV

group may be one of the mechanisms driving the lower fetal and fetal head weight

detected in this group. In this context, changes in placental turnover can determine

placental maturation and function and lead to fetal distress and developmental

abnormalities (55). An increase in the Lz apoptotic ratio may signify damage to this

placental layer, which is consistent with the fact that diverse pathological lesions

associated with congenital disorders were described in placentae from women infected

by ZIKV at different stages of pregnancy (56). Conversely, no proliferative changes

were observed in Jz in high-ZIKV and low-ZIKV ICompetent mice, while increased

and decreased apoptotic rates were detected. It follows that the lack of Jz-Ki-67

induction may suggest that this layer is less capable of restoring proliferation in

response to high-ZIKV challenge, and this may be related to decreased fetal growth.

Our placental ultrastructural analysis detected consistent differences across

ZIKV-exposed groups. The Lz and Jz layers from both strains exhibited signs of ER

stress, i.e., dilated ER cisterns or fragmented ER granular structures. These alterations

may result from the accumulation of folded or poorly folded viral proteins in the ER

lumen (41,57,58). It has been demonstrated that the *Flaviridae* family uses the ER to

replicate (59), and according to Offerdahl et al. (2017)(58), there is evidence of ZIKV

interacting with this organelle, promoting an increased release of Ca^{+2} from the ER to

the cell cytoplasm, thereby causing an increase in the production of reactive oxygen

species (ROS) (60,61).

The mitochondrial ultrastructure in the Lz and Jz layers was severely impacted

by ZIKV exposure. We found evidence of mitochondrial degeneration, i.e.,

mitochondrial membrane rupture, absence of mitochondrial ridges and less electron-

dense mitochondrial matrix in all treated groups. Placental mitochondrial dysfunction is

associated with IUGR (62,63) and may be related, at least in part, to the lower fetal

weight observed in high-ZIKV-ICompetent fetuses along with the altered placental

apoptotic and proliferative patterns. Furthermore, mitochondrial dysfunction together

with ER stress is likely to modify the placental ROS balance and generate local

oxidative stress (64), which is associated with impaired fetal development (65). Of

importance, associations between mitochondrial disruption, ER stress and placental cell

senescence have been reported. Senescence is characterized as an irreversible

interruption of the cell cycle and acquisition of a senescence-associated secretory

phenotype (SASP) that promotes the release of cytokines such as Il-1, Il-6, Il-8 and

proinflammatory proteases (65). Therefore, the increased expression of IL-6 detected in

the placentas of ICompromised mice suggest a SASP profile, which may be related to

changes in the ER and mitochondrial ultrastructure, accompanied by important changes

in apoptosis and cell proliferation. The interactions between mitochondria and the ER

are critical for homeostasis and cell signaling (66). In conjunction with the ER,

mitochondria can regulate cell death mediators in response to hypoxia and inflammation

(67). The increase in apoptosis observed in the high-titer ICompetent groups and the

low-titer ICompromised group may be related to the mitochondrial damage and ER

stress observed. In fact, we observed an important decrease in microvillus abundance in

sinusoidal giant trophoblast cells. Previously, we observed a decrease in microvillus

density in the Lz of pregnancies exposed to malaria in pregnancy (MiP) (35). Together,

our data show that different gestational infective stimuli (MiP and ZIKV) are capable of

damaging placental microvillus abundance and impairing proper fetal-maternal

exchange function and fetal growth/survival.

Next, to investigate whether maternal ZIKV exposure may influence fetal

protection, we evaluated the placental localization and expression (semiquantitative) of

the ABC efflux transporter systems P-gp, Bcrp, Abca1 and Abcg1, which are highly

enriched in labyrinthine microvilli and in human syncytiotrophoblasts. These efflux

transporters exchange drugs, environmental toxins, cytotoxic oxysterols and lipids

within the maternal-fetal interface (25). We found a consistent decrease in labyrinthine

P-gp expression in all ZIKV-exposed groups, demonstrating that ZIKV infection during

pregnancy has the potential to increase fetal exposure to P-gp substrates such as

synthetic glucocorticoids, antibiotics, antiretrovirals, antifungals, stomach-protective

drugs, and nonsteroidal anti-inflammatory drugs (25). Furthermore, Jz-P-gp was

decreased in ICompromised placentae. Although little is known about the Jz function of

ABC transporters, our data highlight the need for further studies investigating the

biological importance of ABC transporters in the placental endocrine and structural

zones of the rodent hemochorial placenta under normal and infective conditions.

ZIKV impaired Lz Bcrp and Abca1 expression in ICompetent (high) and

ICompromised (low) mice. However, no effects were observed in ICompetent animals

at a low ZIKV titer or in Abcg1 in any experimental setting. As such, ZIKV also likely

increases the fetal accumulation of Bcrp substrates (antibiotics, antiretrovirals,

sulfonylureas, folate, mercuric species, estrogenic mycotoxins, carcinogens phototoxic

compounds, among others) and disrupts placental lipid homeostasis (lipids, cholesterol,

cytotoxic oxysterols) by reducing placental Abca1 expression (25,68–71). We can

speculate that the increased fetal accumulation of the P-gp, Bcrp and Abca1 substrates

during ZIKV infection may contribute to the establishment of congenital Zika

syndrome, although additional studies are clearly required to answer this important

question. The current data are in agreement with previous publications showing that

bacterial, viral and protozoan inflammation alters the expression and/or function of P-

gp, Bcrp and Abca1 in biological barriers such as the placenta, yolk sac and blood-brain

barriers (25,26,35,59,72–74).

5. Conclusion

Our data show that gestational ZIKV impacts the fetal phenotype independently

of term fetal viremia. Abnormal placental cell turnover, ultrastructure and transporter

expression may result from the specific proinflammatory responses that depend on the

ZIKV infective load and maternal immune status. Fetal accumulation of drugs,

environmental toxins and lipids within the fetal compartment may be potentially

increased in ZIKV-infected pregnancies due to altered levels of key ABC transporters.

Conflict of Interest:

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

Author contributions

CBVA, FFB, EB, LBA and TMOC conceived and designed the experiments. CBVA,

VRSM, SVAC, HRG, RPCS and VMON performed the experiments. CBVA, SVAC,

FFB, EB, SGM, LBA and TMOC analyzed the data. CBVA, VRSM, EB, LBA and

TMOC wrote the paper and edited the manuscript. All authors contributed to the article

and approved the submitted version.

Funding

This study was supported by the Bill & Melinda Gates Foundation

(MCTI/CNPq/MS/SCTIE/Decit/Bill and Melinda Gates 05/2013; OPP1107597), the

Canadian Institutes for Health Research (SGM: Foundation-148368), Conselho

Nacional de Desenvolvimento Científico e Tecnológico (CNPq; 304667/2016□1,

422441/2016☐3, 303734/2012☐4, 422410/2016☐0), Coordenação de Aperfeiçoamento

Pessoal de Nível Superior (CAPES, finance Code 001), Fundação de Amparo à

Pesquisa do Estado do Rio de Janeiro (FAPERJ, CNE 2015/E26/203.190/2015, PDR-

26/2002/010/2016), and PRPq-Universidade MG Federal MGPEFARJ, CNE

2015/E26/203.190/2015).

Acknowledgments

We would like to thank Alan Moraes for supporting the acquisition of electron

microscopy images and the electron microscopy laboratory at the UFF Biology Institute

for allowing the use of the JEM1011 transmission electron microscope. Mauro Jorge

Castro Cabral for the use of the MAGPIX® System equipment from the Paulo de Góes

Institute of Microbiology/UFRJ. Ernesto T. Marques Jr. (Centro de Pesquisa Aggeu

Magalhães, FIOCRUZ, PE) for providing ZIKV to the Institute of Microbiology Paulo

de Góes, Federal University of Rio de Janeiro and the technicians Juliana Gonçalves

and Rakel Alves for their support during all the experiments.

References
631
1.   Garcez PP, Loiola EC, Madeiro da Costa R, Higa LM, Trindade P, Delvecchio R, et
al. Zika virus impairs growth in human neurospheres and brain organoids. Science.
2016;352(6287):816–8. doi: 10.1126/science.aaf6116.
2.   Brasil P, Calvet GA, Siqueira AM, Wakimoto M, de Sequeira PC, Nobre A, et al.
Zika Virus Outbreak in Rio de Janeiro, Brazil: Clinical Characterization, Epidemiological
and Virological Aspects. PLoS Negl Trop Dis. 2016;10(4):e0004636. doi:
10.1371/journal.pntd.0004636.
3.   Brasil P, Nielsen-Saines K. More pieces to the microcephaly–Zika virus puzzle in
Brazil. The Lancet Infectious Diseases. 2016;16(12):1307–9. doi: 10.1016/S1473-
3099(16)30372-3.
4.   Brasil P, Pereira JP, Moreira ME, Ribeiro Nogueira RM, Damasceno L, Wakimoto
M, et al. Zika Virus Infection in Pregnant Women in Rio de Janeiro. N Engl J Med. 2016
Mar 4;375(24):2321–34. doi: 10.1056/NEJMoa1602412

5.   Jaenisch T, Rosenberger KD, Brito C, Brady O, Brasil P, Marques ET. Risk of
microcephaly after Zika virus infection in Brazil, 2015 to 2016. Bull World Health
Organ. 2017;95(3):191–8. doi: 10.2471/BLT.16.178608.
6.   Proenca-Modena JL, Milanez GP, Costa ML, Judice CC, Maranhão Costa FT. Zika
virus: lessons learned in Brazil. Microbes and Infection. 2018;20(11–12):661–9. doi:
10.1016/j.micinf.2018.02.008.
7.   Jouannic J-M, Friszer S, Leparc-Goffart I, Garel C, Eyrolle-Guignot D. Zika virus
infection in French Polynesia. The Lancet. 2016;387(10023):1051–2. doi:
10.1016/S0140-6736(16)00625-5.
8.   Musso D, Ko AI, Baud D. Zika Virus Infection — After the Pandemic. N Engl J
Med. 2019;381(15):1444–57. doi: 10.1056/NEJMra1808246.
9.   Kasprzykowski JI, Fukutani KF, Fabio H, Fukutani ER, Costa LC, Andrade BB, et al.
A recursive sub-typing screening surveillance system detects the appearance of the
ZIKV African lineage in Brazil: Is there a risk of a new epidemic? International Journal of
Infectious Diseases. 2020;96:579–81. doi: 10.1016/j.ijid.2020.05.090.
10.   Miranda J, Martín-Tapia D, Valdespino-Vázquez Y, Alarcón L, Espejel-Nuñez A,
Guzmán-Huerta M, et al. Syncytiotrophoblast of Placentae from Women with Zika
Virus Infection Has Altered Tight Junction Protein Expression and Increased
Paracellular Permeability. Cells. 2019;8(10):1174. doi: 10.3390/cells8101174.
11.   Jurado KA, Simoni MK, Tang Z, Uraki R, Hwang J, Householder S, et al. Zika virus
productively infects primary human placenta-specific macrophages. JCI Insight.
2016;1(13):e88461. doi: 10.1172/jci.insight.88461.
12.   Quicke KM, Bowen JR, Johnson EL, McDonald CE, Ma H, O'Neal JT, et al. Zika
Virus Infects Human Placental Macrophages. Cell Host & Microbe. 2016;20(1):83–90.
doi: 10.1016/j.chom.2016.05.015.
13.   Tabata T, Petitt M, Puerta-Guardo H, Michlmayr D, Wang C, Fang-Hoover J, et
al. Zika Virus Targets Different Primary Human Placental Cells, Suggesting Two Routes
for Vertical Transmission. Cell Host & Microbe. 2016;20(2):155–66. doi:
10.1016/j.chom.2016.07.002
14.   Simoni MK, Jurado KA, Abrahams VM, Fikrig E, Guller S. Zika virus infection of
Hofbauer cells. Am J Reprod Immunol. 2017;77(2):e12613. doi: 10.1111/aji.12613.
15.   Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, et al. A Mouse
Model of Zika Virus Pathogenesis. Cell Host & Microbe. 2016;19(5):720–30. doi:
10.1016/j.chom.2016.03.010.
16.   Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, et al.
Zika Virus Targets Human STAT2 to Inhibit Type I Interferon Signaling. Cell Host &
Microbe. 2016;19(6):882–90. doi: 10.1016/j.chom.2016.05.009.
17.   Yockey LJ, Varela L, Rakib T, Khoury-Hanold W, Fink SL, Stutz B, et al. Vaginal
Exposure to Zika Virus during Pregnancy Leads to Fetal Brain Infection. Cell.
2016;166(5):1247-1256.e4. doi: 10.1016/j.cell.2016.08.004.
18.   Garcez PP, Stolp HB, Sravanam S, Christoff RR, Ferreira JCCG, Dias AA, et al. Zika
virus impairs the development of blood vessels in a mouse model of congenital
infection. Sci Rep. 2018;8(1):12774. doi: 10.1038/s41598-018-31149-3.
19.   Elong Ngono A, Shresta S. Immune Response to Dengue and Zika. Annu Rev
Immunol. 2018;36(1):279–308. doi: 10.1146/annurev-immunol-042617-053142.
20.   Bayer A, Lennemann NJ, Ouyang Y, Bramley JC, Morosky S, Marques ETDAJ, et
al. Type III Interferons Produced by Human Placental Trophoblasts Confer Protection

against Zika Virus Infection. Cell Host Microbe. 2016;19(5):705–12. doi:
10.1016/j.chom.2016.03.008
21.   Luo H, Winkelmann ER, Fernandez-Salas I, Li L, Mayer SV, Danis-Lozano R, et al.
Zika, dengue and yellow fever viruses induce differential anti-viral immune responses
in human monocytic and first trimester trophoblast cells. Antiviral Research.
2018;151:55–62. doi: 10.1016/j.antiviral.2018.01.003.
22.   Tanaka T, Narazaki M, Kishimoto T. IL-6 in Inflammation, Immunity, and
Disease. Cold Spring Harb Perspect Biol. 2014;6(10):a016295. doi:
10.1101/cshperspect.a016295.
23.   Hunter CA, Jones SA. IL-6 as a keystone cytokine in health and disease. Nature
Immunology. 2015;16(5):448–57. doi: 10.1038/ni.3153.
24.   Rabelo K, Gonçalves AJ da S, Souza LJ de, Sales AP, Lima SMB de, Trindade GF,
et al. Zika Virus Infects Human Placental Mast Cells and the HMC-1 Cell Line, and
Triggers Degranulation, Cytokine Release and Ultrastructural Changes. Cells.
2020;9(4):975–90. doi: 10.3390/cells9040975.
25.   Bloise E, Ortiga-Carvalho TM, Reis FM, Lye SJ, Gibb W, Matthews SG. ATP-
binding cassette transporters in reproduction: a new frontier. Hum Reprod Update.
2015;22(2):164–81. doi: 10.1093/humupd/dmv049.
26.   do Imperio GE, Bloise E, Javam M, Lye P, Constantinof A, Dunk C, et al.
Chorioamnionitis Induces a Specific Signature of Placental ABC Transporters Associated
with an Increase of miR-331-5p in the Human Preterm Placenta. Cell Physiol Biochem.
2018;45(2):591–604. doi: 10.1159/000487100.
27.   Martinelli LM, Fontes KN, Reginatto MW, Andrade CBV, Monteiro VRS, Gomes
HR, et al. Malaria in pregnancy regulates P-glycoprotein (P-gp/Abcb1a) and ABCA1
efflux transporters in the mouse visceral yolk sac. J Cell Mol Med. 2020;00:1–12. doi:
10.1111/jcmm.15682.
28.   Reginatto MW, Fontes KN, Monteiro VRS, Silva NL, Andrade CBV, Gomes HR, et
al. Effect of sub-lethal prenatal endotoxemia on murine placental transport systems
and lipid homeostasis. bioRxiv. 2020 Aug 5;2020.08.04.236745.
29.   Martinelli LM, Reginatto MW, Fontes KN, Andrade CBV, Monteiro VRS, Gomes
HR, et al. Breast cancer resistance protein (Bcrp/Abcg2) is selectively modulated by
lipopolysaccharide (LPS) in the mouse yolk sac. Reprod Toxicol 2020 Dec;98:82-91. doi:
10.1016/j.reprotox.2020.09.001
30.   Imperio GE, Javam M, Lye P, Constantinof A, Dunk CE, Reis FM, et al.
Gestational age-dependent gene expression profiling of ATP-binding cassette
transporters in the healthy human placenta. J Cell Mol Med. 2018/11/08 ed.
2019;23(1):610–8. doi: 10.1111/jcmm.13966.
31.   Vermillion MS, Lei J, Shabi Y, Baxter VK, Crilly NP, McLane M, et al. Intrauterine
Zika virus infection of pregnant immunocompetent mice models transplacental
transmission and adverse perinatal outcomes. Nat Commun. 2017;8(1):14575. doi:
10.1038/ncomms14575.
32.   Szaba FM, Tighe M, Kummer LW, Lanzer KG, Ward JM, Lanthier P, et al. Zika
virus infection in immunocompetent pregnant mice causes fetal damage and placental
pathology in the absence of fetal infection. PLoS Pathog. 2018;14(4):e1006994. doi:
10.1371/journal.ppat.1006994.
33.   Coelho SVA, Neris RLS, Papa MP, Schnellrath LC, Meuren LM, Tschoeke DA, et
al. Development of standard methods for Zika virus propagation, titration, and

purification. Journal of Virological Methods. 2017;246:65–74. doi:
10.1016/j.jviromet.2017.04.011.
34.   Connor KL, Kibschull M, Matysiak-Zablocki E, Nguyen TT-TN, Matthews SG, Lye
SJ, et al. Maternal malnutrition impacts placental morphology and transporter
expression: an origin for poor offspring growth. The Journal of Nutritional
Biochemistry. 2020;78:108329. doi: 10.1016/j.jnutbio.2019.108329.
35.   Fontes KN, Reginatto MW, Silva NL, Andrade CBV, Bloise FF, Monteiro VRS, et
al. Dysregulation of placental ABC transporters in a murine model of malaria-induced
preterm labor. Sci Rep. 2019;9(1):11488. doi: 10.1038/s41598-019-47865-3.
36.   Bloise E, Lin W, Liu X, Simbulan R, Kolahi KS, Petraglia F, et al. Impaired
Placental Nutrient Transport in Mice Generated by *in Vitro* Fertilization. Endocrinology.
2012;153(7):3457–67. doi: 10.1210/en.2011-1921
37.   Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic
and Serologic Properties of Zika Virus Associated with an Epidemic, Yap State,
Micronesia, 2007. Emerg Infect Dis. 2008 Aug;14(8):1232–9. doi:
10.3201/eid1408.080287.
38.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. Methods. 2001;25(4):402–8. doi:
10.1006/meth.2001.1262.
39.   Tschanz SA, Burri PH, Weibel ER. A simple tool for stereological assessment of
digital images: the STEPanizer: TOOL FOR STEREOLOGICAL ASSESSMENT. Journal of
Microscopy. 2011;243(1):47–59. doi: 10.1111/j.1365-2818.2010.03481.x.
40.   Sesso A, Belizário JE, Marques MM, Higuchi ML, Schumacher RI, Colquhoun A,
et al. Mitochondrial Swelling and Incipient Outer Membrane Rupture in Preapoptotic
and Apoptotic Cells. Anatomical Record (Hoboken, N.j▯: 2007). 2012 Oct;295(10):1647.
doi: 10.1002/ar.22553.
41.   Montalbano R, Waldegger P, Quint K, Jabari S, Neureiter D, Illig R, et al.
Endoplasmic Reticulum Stress Plays a Pivotal Role in Cell Death Mediated by the Pan-
Deacetylase Inhibitor Panobinostat in Human Hepatocellular Cancer Cells.
Translational Oncology. 2013;6(2):143-IN6. doi: 10.1593/tlo.12271
42.   Hamilton SA, Tower CL, Jones RL. Identification of Chemokines Associated with
the Recruitment of Decidual Leukocytes in Human Labour: Potential Novel Targets for
Preterm Labour. PLOS ONE. 2013 Feb 22;8(2):e56946. doi:
10.1371/journal.pone.0056946.
43.   Bloise E, Bhuiyan M, Audette MC, Petropoulos S, Javam M, Gibb W, et al.
Prenatal Endotoxemia and Placental Drug Transport in The Mouse: Placental Size-
Specific Effects. PLOS ONE. 2013 Jun 10;8(6):e65728. doi:
10.1371/journal.pone.0065728
44.   Romero R, Mazor M, Brandt F, Sepulveda W, Avila C, Cotton DB, et al.
Interleukin-1α and Interleukin-1 β in Preterm and Term Human Parturition. American
Journal of Reproductive Immunology. 1992;27(3-4):117–23. doi: 10.1111/j.1600-
0897.1992.tb00737.x.
45.   Larocca RA, Abbink P, Peron JPS, de A. Zanotto PM, Iampietro MJ, Badamchi-
Zadeh A, et al. Vaccine protection against Zika virus from Brazil. Nature.
2016;536(7617):474–8. doi: 10.1038/nature18952.
46.   Souza IN de O, Frost PS, França JV, Nascimento-Viana JB, Neris RLS, Freitas L, et
al. Acute and chronic neurological consequences of early-life Zika virus infection in

mice. Science Translational Medicine. 2018;10(444):eaar2749. doi:
10.1126/scitranslmed.aar2749.
47.   Kalagiri RR, Carder T, Choudhury S, Vora N, Ballard AR, Govande V, et al.
Inflammation in Complicated Pregnancy and Its Outcome. Am J Perinatol.
2016;33(14):1337–56. doi: 10.1055/s-0036-1582397.
48.   Barbeito-Andrés J, Pezzuto P, Higa LM, Dias AA, Vasconcelos JM, Santos TMP,
et al. Congenital Zika syndrome is associated with maternal protein malnutrition. Sci
Adv. 2020;6(2):eaaw6284. doi: 10.1126/sciadv.aaw6284.
49.   Saito S, Nakashima A, Shima T, Ito M. Th1/Th2/Th17 and Regulatory T-Cell
Paradigm in Pregnancy. American Journal of Reproductive Immunology.
2010;63(6):601–10. doi: 10.1111/j.1600-0897.2010.00852.x.
50.   Gotsch F, Romero R, Kusanovic JP, Mazaki-Tovi S, Pineles BL, Erez O, et al. The
fetal inflammatory response syndrome. Clin Obstet Gynecol. 2007;50(3):652–83. doi:
10.1097/GRF.0b013e31811ebef6.
51.   Lucas CGO, Kitoko JZ, Ferreira FM, Suzart VG, Papa MP, Coelho SVA, et al.
Critical role of CD4 + T cells and IFNγ signaling in antibody-mediated resistance to Zika
virus infection. Nature Communications. 2018;9(1):3136. doi: 10.1038/s41467-018-
05519-4.
52.   Galinsky R, Polglase GR, Hooper SB, Black MJ, Moss TJM. The consequences of
chorioamnionitis: preterm birth and effects on development. J Pregnancy.
2013;2013:412831. doi: 10.1155/2013/412831.
53.   Mor G, Cardenas I, Abrahams V, Guller S. Inflammation and pregnancy: the role
of the immune system at the implantation site. Ann N Y Acad Sci. 2011;1221(1):80–7.
doi: 10.1111/j.1749-6632.2010.05938.x.
54.   Ribeiro MR, Moreli JB, Marques RE, Papa MP, Meuren LM, Rahal P, et al. Zika-
virus-infected human full-term placental explants display pro-inflammatory responses
and undergo apoptosis. Arch Virol. 2018;163(10):2687–99. doi: 10.1007/s00705-018-
3911-x.
55.   Gown AM, Willingham MC. Improved Detection of Apoptotic Cells in Archival
Paraffin Sections: Immunohistochemistry Using Antibodies to Cleaved Caspase 3. J
Histochem Cytochem. 2002;50(4):449–54. doi: 10.1177/002215540205000401.
56.   Noronha L de, Zanluca C, Burger M, Suzukawa AA, Azevedo M, Rebutini PZ, et
al. Zika Virus Infection at Different Pregnancy Stages: Anatomopathological Findings,
Target Cells and Viral Persistence in Placental Tissues. Front Microbiol.
2018;9(2266):1–11. doi: 10.3389/fmicb.2018.02266.
57.   Lojpur T, Easton Z, Raez-Villanueva S, Laviolette S, Holloway AC, Hardy DB. Δ9-
Tetrahydrocannabinol leads to endoplasmic reticulum stress and mitochondrial
dysfunction in human BeWo trophoblasts. Reproductive Toxicology. 2019;87:21–31.
doi: 10.1016/j.reprotox.2019.04.008.
58.   Offerdahl DK, Dorward DW, Hansen BT, Bloom ME. Cytoarchitecture of Zika
virus infection in human neuroblastoma and Aedes albopictus cell lines. Virology.
2017;501:54–62. doi: 10.1016/j.virol.2016.11.002.
59.   Desuzinges-Mandon E, Arnaud O, Martinez L, Huché F, Di Pietro A, Falson P.
ABCG2 Transports and Transfers Heme to Albumin through Its Large Extracellular Loop.
J Biol Chem. 2010;285(43):33123–33. doi: 10.1074/jbc.M110.139170
60.   Ledur PF, Karmirian K, Pedrosa C da SG, Souza LRQ, Assis-de-Lemos G, Martins
TM, et al. Zika virus infection leads to mitochondrial failure, oxidative stress and DNA

damage in human iPSC-derived astrocytes. Scientific Reports. 2020;10(1):1218. doi:
10.1038/s41598-020-57914-x.
61.   Reemst K, Noctor SC, Lucassen PJ, Hol EM. The Indispensable Roles of Microglia
and Astrocytes during Brain Development. Front Hum Neurosci 2016 Nov 8;10:566.
doi: 10.3389/fnhum.2016.00566.
62.   Guitart-Mampel M, Gonzalez-Tendero A, Niñerola S, Morén C, Catalán-Garcia
M, González-Casacuberta I, et al. Cardiac and placental mitochondrial characterization
in a rabbit model of intrauterine growth restriction. Biochimica et Biophysica Acta
(BBA) - General Subjects. 2018;1862(5):1157–67. doi: 10.1016/j.bbagen.2018.02.006.
63.   Guitart-Mampel M, Juarez-Flores DL, Youssef L, Moren C, Garcia-Otero L, Roca-
Agujetas V, et al. Mitochondrial implications in human pregnancies with intrauterine
growth restriction and associated cardiac remodelling. J Cell Mol Med.
2019;23(6):3962–73. doi: 10.1111/jcmm.14282.
64.   Zhang Z, Rong L, Li Y-P. Flaviviridae Viruses and Oxidative Stress: Implications
for Viral Pathogenesis. Oxid Med Cell Longev. 2019;2019:1409582. doi:
10.1155/2019/1409582.
65.   Martínez F, Kiriakidou M, Strauss JF. Structural and Functional Changes in
Mitochondria Associated with Trophoblast Differentiation: Methods to Isolate
Enriched Preparations of Syncytiotrophoblast Mitochondria. Endocrinology.
1997;138(5):2172–83. doi: 10.1210/endo.138.5.5133.
66.   Burton GJ, Yung HW, Murray AJ. Mitochondrial – Endoplasmic reticulum
interactions in the trophoblast: Stress and senescence. Placenta. 2017;52:146–55. doi:
10.1016/j.placenta.2016.04.001.
67.   Holland O, Dekker Nitert M, Gallo LA, Vejzovic M, Fisher JJ, Perkins AV. Review:
Placental mitochondrial function and structure in gestational disorders. Placenta.
2017;54:2–9. doi: 10.1016/j.placenta.2016.12.012
68.   Bridges CC, Zalups RK, Joshee L. Toxicological significance of renal Bcrp:
Another potential transporter in the elimination of mercuric ions from proximal
tubular cells. Toxicol Appl Pharmacol. 2015;285(2):110–7. doi:
10.1016/j.taap.2015.03.027.
69.   Szilagyi JT, Gorczyca L, Brinker A, Buckley B, Laskin JD, Aleksunes LM. Placental
BCRP/ABCG2 Transporter Prevents Fetal Exposure to the Estrogenic Mycotoxin
Zearalenone. Toxicol Sci. 2019;168(2):394–404. doi: 10.1093/toxsci/kfy303.
70.   Scialis RJ, Aleksunes LM, Csanaky IL, Klaassen CD, Manautou JE. Identification
and Characterization of Efflux Transporters That Modulate the Subtoxic Disposition of
Diclofenac and Its Metabolites. Drug Metab Dispos. 2019;47(10):1080–92. doi:
10.1124/dmd.119.086603.
71.   Mao Q, Unadkat JD. Role of the breast cancer resistance protein (BCRP/ABCG2)
in drug transport--an update. AAPS J. 2015;17(1):65–82. doi: 10.1208/s12248-014-
9668-6.
72.   Lye P, Bloise E, Javam M, Gibb W, Lye SJ, Matthews SG. Impact of Bacterial and
Viral Challenge on Multidrug Resistance in First- and Third-Trimester Human Placenta.
The American Journal of Pathology. 2015;185(6):1666–75. doi:
10.1016/j.ajpath.2015.02.013.
73.   Girard S, Sebire G. Transplacental Transfer of Interleukin-1 Receptor Agonist
and Antagonist Following Maternal Immune Activation. American journal of

reproductive immunology (New York, NY⬜: 1989). 2016;75(1):8–12. doi:
10.1111/aji.12444
74.   Bloise E, Petropoulos S, Iqbal M, Kostaki A, Ortiga-Carvalho TM, Gibb W, et al.
Acute Effects of Viral Exposure on P-Glycoprotein Function in the Mouse Fetal Blood-
Brain Barrier. Cell Physiol Biochem . 2017;41(3):1044-1050. doi: 10.1159/000461569.

884
885
886
887   **Table 1:** Primer sequences for the real-time PCR assay.

Gene	Primer sequences	GenBank accession no.
ZIKV	5'CCGCTGCCCAACACAAG3'	
	5'CCACTAACGTTCTTTTGCAGACAT3'	
mIL-6	5'TCATATCTTCAACCAAGAGGTA3'	NM_031168.2
	5'CAGTGAGGAATGTCCACAAACTG3'	
mIL-1b	5'GTAATGAAAGACGGCACACC3'	XM_006498795.4
	5'ATTAGAAACAGTCCAGCCCA3'	
mIL-10	5'TAAGGGTTACTTGGGTTGCCAAG3'	NM_010548.2
	5'CAAATGCTCCTTGATTTCTGGGC3'	
mIFN-γ	5'AGCAACAGCAAGGCGAAAA3'	NM_008337.4
	5'CTGGACCTGTGGGTTGTTGA3'	
m-IFN-I	5'CTGGAGCAGCTGAATGGAAAG3'	NM_010510.1
	5'CTTGAAGTCCGCCCTGTAGGT3'	
mGAPDH	5'CTTTGTCAAGCTCATTTCCTGG3'	XM_017321385.2
	5'TCTTGCTCAGTGTCCTTGC3'	
TNF-a	5'CCTCACACTCAGATCATCTTCTCA3'	NM_013693.3
	5'TGGTTCTCTTTGAGATCCATGC3'	
Polr2a	5'TCTGCCAAGAATGTGACGCT3'	NM_001291068.1
	5'CCAAGCGGCAAAGAATGTCC3'	

888
889
890
891

900   **Table 2:** Maternal weight gain throughout pregnancy

	ICompetent			ICompromised		
Maternal weight (g)	**Mock (n=11)**	**High (n=15)**	**Low (n=9)**	**Mock (n=3)**	**High (n=3)**	**Low (n=3)**
GD0.5	22.4 ± 0.7^a	23.0 ± 0.6^a	21.9 ± 0.7^a	21.7 ± 1.9^a	$21.7\pm0.9^{a,b}$	25.0 ± 1.0^a
GD12.5	25.5 ± 0.6^b	26.2 ± 0.8^b	25.1 ± 0.5^b	25.3 ± 1.2^a	26.0 ± 1.0^a	28.0 ± 0.6^a
GD18.5	31.6 ± 0.7^c	30.3 ± 0.9^c	31.4 ± 1.1^c	32.0 ± 1.5^b	20.0 ± 1.5^b	30.7 ± 0.3^b
ΔGD12.5-0.5 (Δ)	4	3	3	3	4	3
ΔGD18.5-0.5 (Δ)	9	7	10	10	-2	5

901   GD=gestational day; Δ=delta; values are expressed as the mean±standard error of the
902   mean (SEM). One-way ANOVA followed by Tukey's test analyzed differences among
903   GD0.5, 12.5 and 18.5 in each group; different letters indicate p<0.05.

904
905   **Figure Captions**

906
Figure 1: ZIKV infection induced fetal changes during pregnancy. A) Experimental
schematic. The fetuses from C57BL/6 dams and sires were termed ICompetent. The
fetuses from the mates of A129 dams and C57BL/6 sires were termed ICompromised.
B) Uterine horn images and survival images (arrows show resorption sites). C)
Fetal/reabsorption images. Measurement of fetal biometric parameters at GD18.5. D)
Fetal weight, (E) fetal head weight and (F) fetal head weight/fetal weight ratio in
ICompetent (n=11/15/9 dams in the mock, high and low groups, respectively) and
ICompromised (n=3 dams/group). * example reabsorption. The values are plotted as
individual values, and the bar represents the mean. Unpaired Student's t-test or
nonparametric Mann-Whitney test was used for ICompromised groups, and one-way
ANOVA followed by Tukey's post-test was used for ICompetent comparisons between
different groups.
Figure 2: Viral load present in the spleen, placenta, fetal and maternal brains after
ZIKV infection. A) Maternal spleen and brain viral load and B) placenta and fetal brain
viral load of ZIKV-infected dams. C) Animal viral serology of all groups after 4, 48 and
144 infection hours. ICompetent (n=11/15/9 dams in the mock, high and low ZIKV titer
groups, respectively) and ICompromised (n=3 dams/group).
924
Figure 3: ZIKV promotes an inflammatory response in the placenta, maternal
spleen and serum during pregnancy.
A) Cytokine and chemokine serum levels at GD18.5 in the ICompetent mock (n=9
dams), high (n=8) and low (n=9) and ICompromised mock and low (n=3/groups) ZIKV
titer groups. A-C) IL-1β, IL-6, CCL-2 and CXCL-1 maternal serum levels. D-F)

Placental IL-1β, IL-6, CCL-2 and CXCL-1 levels. G-I) Placental RNA levels of Ifnγ,
Ifn1, Il-1β, Il-6 and Tnf-α. The broken line shows the expression levels of both lineages
in the mock group. The values are expressed as the mean ± SEM. Unpaired Student's t-
test or nonparametric Mann-Whitney test was used for comparisons between
ICompetent groups, and one-way ANOVA followed by Tukey's posttest was used for
comparisons between different ICompetent groups. * $p<0.05$.
936
937

Figure 4: Placental remodeling of ICompetent groups induced by ZIKV infection.
A) Quantification of Ki-67$^+$ nuclei in the mouse placenta Lz and representative images
of nuclei immunostained by Ki-67$^+$-positive nuclei in the ICompetent group
(n=6/group). B) The image shows the apoptotic nuclei score in the Lz and
representative images of nuclei immunostained by TUNEL in the ICompetent group
(n=5/group). C) Quantification of Ki-67+ nuclei in the mouse placenta Lz and
representative images of nuclei immunostained by Ki-67$^+$ nuclei in the ICompromised
group (n=3/group). D) The image shows the apoptotic nuclei score in the LZ and
representative images of nuclei immunostained by TUNEL in the ICompromised group
(n=3/group). The values are expressed as the mean ± SEM. One-way ANOVA followed
by Tukey's post hoc test was used for comparisons between different ICompetent
groups. Unpaired Student's t-tests or nonparametric Mann-Whitney tests were used for
the ICompromised groups. CTR(-)=All negative controls were performed with the
omission of the primary antibody. Scale bar = 50 µm.
952
953

Figure 5: ZIKV induces hyperproliferation of the JZ area in infected placentas of
ICompetent groups. A) Quantification of Ki-67+ nuclei in the placental Jz of mice and
representative images of immunostained nuclei in the ICompetent (n=6/group). B) The
image shows apoptotic nuclei quantification in the Jz area and representative images of
nuclei immunostained by TUNEL in the ICompetent (n=5/group). C) Quantification of
Ki-67+ nuclei in the placental Jz area and representative images of immunostained
nuclei in the ICompromised group (n=3/group). D) The image shows apoptotic nuclei
quantification in the Jz area and representative images of nuclei immunostained by
TUNEL in the ICompromised (n=3/group). The values are expressed as the mean ±
SEM. One-way ANOVA followed by Tukey's post hoc test was used for comparisons
between different ICompetent groups. Unpaired Student's t-tests or nonparametric
Mann-Whitney tests were used for the ICompromised groups. CTR(-)=All negative
controls were performed with the omission of the primary antibody. Scale bar = 50 µm.
967
968

Figure 6: Associated ultrastructural changes in the placental LZ after ZIKV
infection. Transmission electron photomicrographs of ICompetent mock (A);
ICompetent high (B); ICompetent low (C); ICompromised mock (D) and
ICompromised low (E) groups (N=5 dams/group). We observed dilatation in the ER
cisterns of ICompetent high placentas. Additionally, there was a reduction in the
microvilli from both ICompetent high and ICompetent low placentas. In the
ICompromised low group, we found fragmented ER and microvillus reduction. All
infected groups showed degenerate mitochondria. GER= Granular Endoplasmic
Reticulum; ⊗= Dilated Granular Endoplasmic Reticulum; *= Fragmented Granular
Endoplasmic Reticulum; Mt= Mitochondria; MtD= Degenerate Mitochondria; Mv=

Microvilli; EN= Euchromatic Nuclei; SS= Sinusoidal Space; TGC= Trophoblastic
Giant Cell. Scale bar = 2 µm.
981
Figure 7: Associated ultrastructural changes in the placental JZ after ZIKV
infection. Transmission electron photomicrographs of ICompetent mock (A);
ICompetent high (B); ICompetent low (C); ICompromised mock (D) and
ICompromised low (E) groups (N=5 dams/group). We found deteriorating mitochondria
and dilated reticulum endoplasmic cisterns in both the high ICompetent and low
ICompromised groups. GER= Granular Endoplasmic Reticulum; ⊗= Dilated Granular
Endoplasmic Reticulum; Mt= Mitochondria; MtD= Degenerate Mitochondria; Mv=
Microvilli; EN= Euchromatic Nuclei; SS= Sinusoidal Space; TGC= Trophoblastic
Giant Cell. White arrows HA = Heterochromatin Area. Scale bar = 2 µm.
991
Figure 8: ZIKV infection decreases P-gp, Bcrp and Abca1 expression in the
placental LZ of infected mice in the ICompetent groups. Immunostaining
quantification and representative immunohistochemistry images of A) P-gp, B) Bcrp, C)
Abca1 and D) Abcg1. The images represent the percentage of stained cells in each
analyzed field. The values are expressed as the mean ± SEM. One-way ANOVA
followed by Tukey's post hoc test was used for comparisons between different
ICompetent groups. Scale bar = 50 µm. N=6/ group. CTR(-)=All negative controls were
performed with the omission of the primary antibody.
1000
Figure 9: ZIKV infection decreases P-gp, Bcrp and Abca protein expression in the
placental LZ of infected mice in the ICompromised groups. Immunostaining
quantification and representative immunohistochemistry images of A) P-gp, B) Bcrp, C)
Abca1 and D) Abcg1. The images represent the percentage of stained cells in each
analyzed field. The values are expressed as the mean ± SEM. Unpaired Student's t-tests
or nonparametric Mann-Whitney tests were used for the ICompromised groups. Scale
bar = 50 µm. N=3 for each group. CTR(-)=All negative controls were performed with
the omission of the primary antibody.
1009
Figure 10: ZIKV infection did not impact P-gp, Bcrp, Abca1 or ABcg1 expression
in the JZ area in infected placentas of ICompetent groups. Immunostaining
quantification and representative immunohistochemistry images of A) P-gp, B) Bcrp, C)
Abca1 and D) Abcg1. The images represent the percentage of stained cells/viable tissue.
The values are expressed as the mean ± SEM. One-way ANOVA followed by Tukey's
post hoc test was used for comparisons between different ICompetent groups. Scale bar
= 50 µm. N=6/each group. CTR(-)=All negative controls were performed with the
omission of the primary antibody.
1018
Figure 11: ZIKV infection decreases Bcrp expression in the JZ area in infected
placentas of the ICompromised group. Immunostaining quantification and
representative immunohistochemistry images of A) P-gp, B) Bcrp, C) Abca1 and D)
Abcg1. The images represent the percentage of stained cells/viable tissue. The values
are expressed as the mean ± SEM. Unpaired Student's t-tests or nonparametric Mann-
Whitney tests were used for the ICompromised groups. Scale bar = 50 µm. N=3 for
each group. CTR(-)=All negative controls were performed with the omission of the
primary antibody.
1027

1028
1029
1030

ZUNCTIONAL ZONE

Mock | High | Low | CTR (-)

Mock | Low | CTR (-)

A — P-gp (%) Labyrinth area — p=0.001 — Mock, Low — 50μm

B — Bcrp (%) Labyrinth area — p=0.05 — Mock, Low — 50μm

C — Abca1 (%) Labyrinth area — p=0.05 — Mock, Low — 50μm

D — Abcg1 (%) Labyrinth area — Mock, Low — 50μm