# Motor planning brings human primary somatosensory cortex into movement-specific preparatory states

Giacomo Ariani[1,2,*], J. Andrew Pruszynski[1,4,5,6], Jörn Diedrichsen[1,2,3]

[1]The Brain and Mind Institute, [2]Department of Computer Science, [3]Department of Statistical and Actuarial Sciences, [4]Department of Physiology and Pharmacology, [5]Department of Psychology, [6]Robarts Research Institute, Western University, London, ON N6A3K7, Canada.

*Correspondence should be addressed to Giacomo Ariani at gariani@uwo.ca, Western Interdisciplinary Research Building, London, ON N6A 3K7, Canada.

**Running title.** Motor planning in human sensorimotor cortex.

**Word count.** Abstract: 220; Significance statement: 117; Introduction: 432; Results: 1399; Discussion: 910. 28 pages, 4 figures, 0 tables, 1 supplementary figure.

**Author contributions.** G.A. and J.D. designed research; G.A. performed research; G.A. analyzed data; G.A. drafted the manuscript; G.A., J.A.P, and J.D. edited the manuscript.

**Acknowledgements.** This work was supported by a NSERC Discovery Grant (RGPIN-2016-04890) awarded to J.D., and the Canada First Research Excellence Fund (BrainsCAN). The authors wish to thank Eva Berlot for helpful discussions and contributions to data analysis.

**Disclosures.** The authors declare no conflicts of interest.

# Abstract

Motor planning plays a critical role in producing fast and accurate movement. Yet, the neural processes that occur in human primary motor and somatosensory cortex during planning, and how they relate to those during movement execution, remain poorly understood. Here we used 7T functional magnetic resonance imaging (fMRI) and a delayed movement paradigm to study single finger movement planning and execution. The inclusion of no-go trials and variable delays allowed us to separate what are typically overlapping planning and execution brain responses. Although our univariate results show widespread deactivation during finger planning, multivariate pattern analysis revealed finger-specific activity patterns in contralateral primary somatosensory cortex (S1), which predicted the planned finger movements. Surprisingly, these activity patterns were similarly strong to those found in contralateral primary motor cortex (M1). Control analyses ruled out the possibility that the detected information was an artifact of subthreshold movements during the preparatory delay. Furthermore, we observed that finger-specific activity patterns during planning were highly correlated to those during movement execution. These findings reveal that motor planning activates the specific S1 and M1 circuits that are engaged during the execution of a finger movement – while activity in S1 and M1 is overall suppressed. We propose that preparatory states in S1 may improve movement control through changes in sensory processing or via direct influence of spinal motor neurons.

# Key words

Motor planning; Sensorimotor cortex; Finger control; fMRI; Representational geometry.

## Significance statement

Motor planning is important for good motor performance, yet it is unclear which neural processes underlie the preparation of the nervous system for an upcoming movement. Using high-resolution functional neuroimaging, we investigated how the planning of finger movements changes the activity state in primary motor (M1) and somatosensory (S1) cortex, and how brain responses during planning and execution relate to each other. We show that planning leads to finger-specific activation in both M1 and S1, which is highly similar to the finger-specific activity patterns elicited during movement execution. Our findings suggest that even S1 is being specifically prepared for an upcoming movement, either to actively contribute to the outflowing motor command or to enable movement-specific sensory gating.

## Introduction

Animals are capable of generating a wide variety of dexterous behaviors accurately and effortlessly on a daily basis. This remarkable ability relies on the motor system reaching the appropriate preparatory state before each movement is initiated. At the level of behavior, the process of motor programming, or planning, has long been shown to be beneficial to performance (1–3), leading to faster reaction times (4–6) and more accurate response selection (7–10). The behavioral study of motor planning has spurred the neurophysiological investigation of what movement parameters are specified in the neuronal firing of a number of cortical regions including the dorsal premotor cortex, PMd (11–13), the supplementary motor area, SMA (14), and the posterior parietal cortex, PPC (15–17). Building on this work, human neuroimaging studies have shown that activity in parieto-frontal brain regions during planning of prehension movements can be used to decode several movement properties such as grip type (18, 19), action order (20), and effector (21–24). At the level of neural population dynamics (25), motor planning can be understood as bringing the neuronal state towards an ideal preparatory point. Once this state is reached and the execution is triggered, the intrinsic dynamics of the system then let the movement unfold (26, 27). While the neuronal correlates of movement planning have largely been studied in non-human primates using upper limb movements, such as reaching and grasping, motor planning also plays a crucial role in fine hand control (7).

Despite their importance for everyday dexterous behaviors such as typing, writing, or tying a knot, finger movements have not been closely investigated at the single neuron level. In humans, previous fMRI studies of finger movements have not separated planning from execution (28–35). Therefore, we have an incomplete picture of how motor planning readies the human sensorimotor system for the production of individuated finger movements. Based on previous work in reaching, we expected that the primary motor cortex (M1) should represent movements during both planning and execution (36–38). What is unclear, however, is whether the primary somatosensory cortex (S1) also receives information about the planned movement before movement onset. Furthermore, we currently do not know how brain representations of planned finger movements relate to those during movement execution. To address these gaps, we designed a high-field (7T) fMRI experiment to study what brain regions underlie the planning of individual finger movements. We used no-go trials

66 and variable delays to temporally separate the evoked responses to movement planning and

67 execution, and advanced multivariate pattern analyses to examine the correspondence

68 between fMRI correlates of planned and executed finger movements.

## 70 Results

*Deactivation in sensorimotor regions during planning of finger movements*

We instructed 22 participants to plan and execute repeated keypresses with individual

fingers of their right hand on a keyboard device while being scanned with 7T fMRI. The key

to be pressed corresponded to one of three fingers (thumb, middle, and little) and was cued

during the preparation phase by numbers on the screen (1=thumb, 3=middle, 5=little, e.g.,

Fig. 1A).

Figure 1 | fMRI task and BOLD responses. A. Example trial with timing information. Background colors indicate different experimental phases (yellow = preparation; green = movement (go) or wait (no-go); purple = reward; gray = inter-trial interval, ITI). B. Group-averaged BOLD response (N = 22) for go (blue) and no-go (orange) trials in a region that shows no planning evoked activity (Left M1, top), and one that shows some planning evoked activity (Left aSPL, bottom). Shaded areas indicate standard error of the mean (SEM). Background colors correspond to trial phases as in A.

After a variable delay (4–8 seconds), participants received a color cue indicating whether to

press the planned finger (go trials), or whether to withhold the response (no-go trials). Upon

the go cue, participants had to initiate the correct response as fast as possible and make 6

presses of the designated finger, before receiving accuracy points for reward (see Methods).

To control for involuntary overt movements during the preparation phase, we required

participants to maintain a steady force on all of the keys during the delay, which was closely

monitored online. To ensure that planning results would not be biased by the subsequent

execution, we restricted all our analyses to no-go trials without a subsequent movement (see

Methods). First, we asked which brain regions showed an evoked response during the

planning of finger movements (e.g., Fig. 1B). We focused our analysis on the lateral aspect

of the contralateral (left) hemisphere (purple and white areas of the Fig. 2 inset) which

included premotor, primary motor, primary somatosensory, and anterior parietal cortical

regions. To examine brain activation during finger planning, we performed a univariate

contrast of the preparation phase (across the three fingers) versus the resting baseline (Fig.

2A). Overall, the instruction stimulus evoked little to no activation in our regions of interest

(ROIs, Fig. 2A). In fact, compared to resting baseline, we observed significant deactivation

(Fig. 2E) in dorsal premotor cortex (PMd, $t_{21}$= -2.642, $p$ = 0.015), primary motor cortex (M1,

$t_{21}$ = -7.592, $p$ = 1.887e-07), and primary somatosensory cortex (S1, $t_{21}$ = -6.618, $p$ = 1.491e-

06). In comparison, movement execution strongly activated M1 and S1 (Fig. 2C), with

activation being significant in all our ROIs (Fig. 2E, all $t_{21}$ > 14.469, all $p$ < 2.153e-12).

*Planning induces informative patterns in primary motor and somatosensory cortex*

Although we found little univariate activation in our main ROIs, preparatory processes need

not increase the overall activation in a region. Rather, the region could converge to a specific

preparatory neural state (26), while activity increments and decrements within the region (i.e.,

at a finer spatial scale) average each other out. In this case, information about planned

movements would be present in the multivoxel activity patterns in that region. To test this

idea, we calculated the crossnobis dissimilarity (see Methods) between activity patterns.

Systematically positive values of this dissimilarity measure indicate that the patterns reliably

differentiate between the different planned actions (39, 40). Indeed, a surface-based

searchlight approach (41) revealed reliably positive crossnobis dissimilarity between the

activity patterns related to planning of individual finger movements (Fig. 2B) in both M1 ($t_{21}$ =

2.734, $p$ = 0.012) and S1 ($t_{21}$ = 2.987, $p$ = 0.007, Fig. 2F).

**Figure 2 | Activation and distance analyses of movement planning and execution.** The inset shows the inflated cortical surface of the contralateral (left) hemisphere, highlighting the area of interest (purple) and the strip used for cross-section analysis (white). Major sulci are indicated by white dotted lines. **A.** Univariate activation map (percent signal change) for the contrast planning>baseline (no-go trials only). **B.** Multivariate searchlight map of the mean crossnobis distance between the planning of the three fingers (no-go trials only). **C.** Same as A, but for the univariate contrast execution>baseline. **D.** Same as B, but for the mean crossnobis distance between fingers during movement execution. **E.** Cross-section analysis of the mean percent signal change (± SEM) during planning (orange) and execution (blue). Horizontal bars indicate significance ($p$ < 0.05) in a 2-sided one-sample $t$-test vs zero for

selected ROIs. **F.** Same as E, but for the mean crossnobis distance (± SEM). Pre-CS = precentral sulcus; CS = central sulcus; Post-CS = postcentral sulcus; PMd = dorsal premotor cortex; M1 = primary motor cortex; S1 = primary somatosensory cortex; aSPL = anterior superior parietal lobule.

Surprisingly, the distribution of these informative patterns was highly similar to the distribution

of information during movement execution (Fig. 2D). Visual inspection suggested that the

informative patterns during planning may be concentrated more dorsally in M1 and S1

relative to execution. However, a Hotelling $T^2$ test revealed no systematic difference in the

location of the peak vertex between planning and execution across subjects (M1: $T^2_{2,20}$ =

0.725, $p$ = 0.712; S1: $T^2_{2,20}$ = 2.424, $p$ = 0.335). Thus, our results show that information about

single finger movements is already represented during motor planning in the same parts of

the primary motor and somatosensory cortices that are engaged during execution of the

movements. Given that we only used the activity estimates from no-go trials (~40% of total

trials), this information cannot be explained by a spill-over from subsequent execution-related

activity. An analysis using the estimates of planning activity from all trials yielded very similar

results (see Fig. S1), demonstrating that we could separate planning from execution-related

signals.

*Activity patterns are not caused by small movements during the preparation phase*

The presence of planning-related information in primary sensorimotor regions was surprising,

especially in S1, where it had not previously been reported in comparable fMRI studies (18,

20). To ensure that these results were not caused by overt movement, participants were

instructed to maintain a steady force on the keyboard during the preparation phase, such

that we could monitor even the smallest involuntary preparatory movements. Inspection of

the average force profiles (Fig. 3A) revealed that participants were successful in maintaining

a stable force between 0.2 and 0.4 N during preparation. However, averaging forces across

trials may obscure small, idiosyncratic patterns visible during individual trials (Fig. 3B) that

could be used to distinguish the different movements. To test for the presence of such

patterns, we submitted both the mean and standard deviation of the force traces on each

finger to a multivariate dissimilarity analysis (see Methods).

**Figure 3 | Small involuntary movements do not explain preparatory activity patterns in M1 and S1. A.** Mean finger force (± SEM) plotted in 10 ms bins, time-aligned to instruction onset (dotted vertical line) and end of the preparation phase (dashed vertical lines), separately for the three fingers and go (blue) and no-go (orange) trials. **B.** Example of an individual trial with a 6 s preparation phase, followed six presses of the little finger (green). Horizontal solid line denotes press threshold (1 N). Dash-dotted lines denote the boundaries of the finger pre-activation red area in Fig. 1A (see Methods). RT = reaction time; MT = movement time. **C.** Pearson's correlation (r) between behavioral and neural distances in M1 and S1 (see Methods) during the preparation phase (planning, orange). Each dot represents an individual participant (N = 22). Solid line shows linear regression, $p$-values refers to the slope of the line. **D.** Same as C, but during the movement phase (execution, blue).

Indeed, this sensitive analysis revealed that some participants showed small movement

patterns predictive of the upcoming finger (Fig. 3C). These differences, however, were ~200-

300 times smaller than the average differences during execution (Fig. 3D). More importantly,

the magnitude of the behavioral differences for the preparation phase was uncorrelated to

the amount of planning information present in sensory-motor regions (both $p$-values for the

slope of the linear fit > 0.3). Even after correcting for the influence behavioral patterns, the

informative patterns in M1 and S1 remained significant, as shown by a significantly positive

intercept in the linear fit in Fig. 3C (M1: $p = 0.032$; S1: $p = 0.007$). Thus, the finding of finger-

specific activity patterns in M1 and S1 cannot be explained by small involuntary movements

during the preparation phase.

*Single finger activity patterns from planning to execution are positively correlated*

How do planning-related activity patterns in M1 and S1 relate to the activity patterns observed

during execution? Neurophysiological experiments have suggested that patterns of

movement preparation are orthogonal – or uncorrelated – to the patterns underlying active

movement (42). This arrangement allows movement preparation to occur without causing

overt movement. When we compared the planning- and execution-related activity patterns

as measured with fMRI, a technique that samples neuronal activity at a much coarser spatial

resolution, we found the opposite result. Planning- and movement-related patterns for the

same finger were tightly related. Inspection of the representational dissimilarity matrices

(RDMs) for M1 and S1 (Fig. 4A), clearly shows that the biggest difference was between

planning and execution patterns, which can also be seen in a 3D representation of the

representational geometry (PC1 in Fig. 4B). Within each phase, the pattern for the thumb was

more distinct than those for the other fingers, replicating previous results from execution

alone (28, 29).

**Figure 4 | Correlated representations of single fingers across planning and execution. A.** Representational dissimilarity matrices (RDMs) of the activity patterns evoked by the three fingers during the preparation (no-go planning, orange) and movement (execution, blue) phases, separately for the two main ROIs (M1, left; S1, right). **B.** Multidimensional scaling (MDS) projection of the RDMs in A highlighting the first principal component (PC 1, difference between planning and execution). The black cross denotes resting baseline. **C.** Same as B, but rotated view to highlight the representational geometries for PC 2 and PC 3. **D.** Pattern component modelling (PCM) evaluation of models of different correlations between planning- and execution-related activity patterns (x-axis). Shown is the group-average of the individual log-likelihood (± SEM) curve expressed as a difference from the mean log-likelihood across models. Horizontal gray bars indicate models that perform statistically equivalently ($p > 0.05$) to the best fitting model (determined in a cross-validated fashion, see Methods). Red dots indicate points of individually best fitting correlations (N = 22). Red dashed lines denote the average winning models across participants. Dotted lines show the projections of horizontal bars and dashed lines on the respective axes.

Importantly, however, a rotated view of the representational geometry (Fig. 4C) showed that

the finger patterns were arranged in a congruent way, with planning and execution related

activity patterns for the same finger being closer to one another. To more precisely quantify

the correspondence between planning and execution pattern for each finger, we used

Pattern Component Modeling (PCM, 43) to evaluate the likelihood of the data, assuming a

true correlation between 0 and 1. This approach automatically corrects for the biasing effect

of measurement noise, which would lead to simple empirical pattern correlations to be lower

than the true correlation. From the individual fits, we found that the average best correlation

model was at 0.83 (± 0.053 SEM) for M1 and at 0.81 (± 0.061 SEM) for S1 (Fig. 4D, dashed

lines). Using a cross-validated approach (see Methods), we compared the log-likelihoods to

test whether the best fitting model was significantly better than any of the other correlation

models. In both ROIs, the best fitting model was significantly better than the zero-correlation

model (i.e., activity patterns across planning and execution totally uncorrelated, both $p <$

0.005), but not better than the one-correlation model (i.e., activity patterns are scaled version

of each other, both $p > 0.1$). By applying this method to every other model, we have evidence

that the true (i.e., noiseless) correlation between planning and execution finger-specific

activity pattern was between 0.41–1.0 in M1 and between 0.54–1.0 in S1 (Fig. 4D, horizontal

bars). In sum, at the resolution of fMRI, the process of movement planning seems to induce

a finger-specific pattern which very similar, and possibly identical, to the pattern activated

during movement execution.

## 190 Discussion

In the present study, we asked participants to produce repeated single finger movements

while undergoing 7T fMRI. We used a variable preparatory delay and no-go trials to cleanly

dissociate the brain responses to the consecutive preparation and movement phases. We

found that information about planned finger movements is present in both S1 and M1 before

movement onset, even though the overall level of activation in these regions was below

resting baseline. Moreover, while execution elicited much higher brain activation, the fine-

grained, finger-specific activity patterns were highly similar across planning and execution.

Control analyses confirmed that the observed results were not caused by pre-movement
finger activity.

Our finding that motor planning activates M1 in a finger-specific fashion was not
necessarily surprising given many neurophysiological studies reporting anticipatory activity
of M1 neurons related to movement intentions (37, 44, 45), as well as human neuroimaging
showing shared information between delayed and immediate movement plans (36). In
contrast, the robust activity patterns related to single finger planning in S1 were more
surprising, given that this region is often considered to be mostly concerned with processing
incoming sensory information from tactile and proprioceptive receptors arising after
movement onset (18, 20–22). So, what could then be the role of S1 during movement
planning?

First, it is worth noting that there are substantial projections from S1 (Brodmann area 3a)
that terminate in the ventral horn of the cortico-spinal tract (46, 47). Although stimulation of
area 3a in macaques typically fails to evoke overt movements (48), it has been suggested
that this population of cortico-motoneurons specifically projects to gamma motoneurons that
control the sensitivity of muscle spindle afferents (47). Thus, it is possible that S1 plays an
active role in movement generation by preparing the spindle apparatus in advance of the
movement.

Second, the finger-specific preparatory state in S1 may reflect the anticipation or
prediction of the upcoming sensory stimulation, allowing for a movement-specific sensory
gain control (49). Some sensory stimuli could become attenuated to maintain movement
stability and filter out irrelevant or self-generated signals. Indeed, multiple studies have
shown that both somatosensation and somatosensory-evoked potentials in S1 decrease
during voluntary movement (50–53). Alternatively, sensory processing of the expected salient
signals could be enhanced to improve movement execution and adaptation.

Very recently, a human electrocorticography (ECoG) study suggested a possible role for
S1 in cognitive-motor imagery (54). The authors recorded neural activity from S1 while a
tetraplegic participant imagined reaching movements and found that S1 neurons encoded
movement direction during motor imagery in the absence of actual sensations. While the
authors raise the concern that their results may be unique to individuals who have lost their
main peripheral input, our finding of encoding of movement-related information in S1 before
the onset of a movement suggests that movement-specific information in S1 without actual

movement is a general phenomenon in the intact human motor system. Our findings are also consistent with a second recent ECoG study in non-human primates (55). During a delayed reaching and grasping task, the authors showed movement-specific information in the ECoG signals from S1 well before movement initiation, and only slightly later than in M1. Together, these results suggest that the somatosensory system not only passively receives signals from the external world but also actively processes them via interactions with anticipatory information from the motor system.

One may wonder why such movement-specific encoding in S1 during planning has not been reported in previous human fMRI studies. One contributing reason may be that we had higher sensitivity to detect these signals than previous studies, as we used finger rather than arm movements (the former having more distinct cortical representation in S1), higher field strength (7T) and spatial resolution, and a more sensitive multivariate analysis method (crossnobis dissimilarity vs. pattern classification, Walther et al., 2016).

Our second main finding – the close correspondence between finger-specific activity patterns across planning and execution – appears to be at odds with the idea that these two processes occupy orthogonal neural subspaces to avoid overt movement during planning (42, 56). We think that there are at least two possible explanations for this. First, the divergence of results could be caused by the difference between spatially directed arm movements and non-spatial finger movements. If for single finger movements even single-neuron activity patterns are highly correlated between planning and execution, then overt movement during planning would need to be actively suppressed, for example through the deactivation that we observed around the central sulcus. An alternative and perhaps more likely explanation of the discrepancy lies in the different measurement modalities. While the orthogonality was observed in electrophysiological recordings of individual neurons, the fMRI measurements we employed here mainly reflect excitatory postsynaptic potentials (57) and average metabolic activity across hundreds of thousands of cortical neurons. Thus, it is possible that planning pre-activates the specific cortical columns in M1 and S1 that are also most active during movement of that finger. Within each of these cortical micro-circuits, however, planning-related activity could still be orthogonal to the activity observed during execution at the single neuron level (e.g., see Arbuckle et al., 2020, for a similar observation for cortical representations of flexion and extension finger movements). This would suggest a new hypothesis for the architecture of the sensory-motor system where movement planning

262 pre-activates the movement-specific circuits in M1 and S1. However, it does so in a fashion

263 that the induced planning-related activity is, in terms of the firing output of neurons,

264 orthogonal to the patterns during movement execution.

## Methods

*Participants*

Twenty-three right-handed neurologically healthy participants volunteered to take part in the

experiment (13 F, 10 M; age 20–31 years, mean 23.43 years, SD 4.08 years). Criteria for

inclusion were right-handedness and no prior history of psychiatric or neurological disorders.

Handedness was assessed with the Edinburgh Handedness Inventory (mean 82.83, SD

9.75). All experimental procedures were approved by the Research Ethics Committee at

Western University. Participants provided written informed consent to procedures and data

usage and received monetary compensation for their participation. One participant withdrew

before study completion and was excluded from data analysis (final N = 22).

*Apparatus*

Repeated presses of right-hand finger movements were performed on a custom-made MRI-

compatible keyboard device (Fig. 1A). The keys of the device did not move but force

transducers underneath each key measured isometric force production at an update rate of

2 ms (Honeywell FS series; dynamic range 0-25 N; sampling 200 Hz). A keypress/release

was detected when the force crossed a threshold of 1 N. The forces measured from the

keyboard were low-pass filtered to reduce noise induced by the MRI environment, amplified,

and sent to PC for online task control and data recording.

*Task*

We used a task in which participants produced repeated keypresses with their right-hand

fingers in response to numerical cues appearing on the screen (white outline, Fig. 1A). On

each trial, a string of 6 numbers (instruction) instructed which fingers to plan (1 = thumb, 3 =

middle, 5 = little). After a variable delay (4, 6, or 8 s randomly sampled from a geometric

distribution with $p$ = 0.5; preparation phase, yellow background), participants received a

color cue (go/no-go cue) indicating whether to perform the planned finger movements (blue

outline = go, $p$ = 0.6), or not (orange outline = no-go, $p$ = 0.4). The role of no-go trials was to

de-couple the hemodynamic response to the successive planning and execution events. To

encourage planning during the delay period, at the go/no-go cue the digits were masked

with asterisks, and participants had to perform the movements from memory (movement

phase, green background). Participants had 2.5 seconds to complete the movement phase,

and a vanishing white bar under the asterisks indicated how much time was left to complete

all of the keypresses. To limit and monitor unwanted movements during the preparation

phase, we required participants to pre-activate their fingers by maintaining a steady force of

around 0.2-0.3 N on all of the keys during the Preparation phase. As a visual aid, we

displayed a red area (between 0 and 0.5 N) and asked participants to remain within the

middle of that range with all the fingers (touching either boundary of the red area would incur

an error, counting as unwanted movement). In the case of a no-go trial, participants were

instructed to remain as still as possible maintaining the finger pre-activation until the end of

the movement phase (i.e., releasing any of the keys would incur an error). During the

movement phase participants also received online feedback on the correctness of each

press with asterisks turning either green, for a correct press, or red, for incorrect presses.

After the movement phase, participants received points based on their performance (reward

phase, 0.5 s, purple background). Participants were instructed to perform the movements as

accurately as possible. As long as they remained within task constraints (i.e., 6 keypresses

in less than 2.5 seconds), an exact movement speed was not enforced. On a trial-by-trial

basis, during the reward phase participants received points for their performance according

to the following scheme: -1 point in case of no-go error or go cue anticipation (timing errors);

0 points for pressing any wrong key (press error); 1 point in case of a correct no-go trial; and

2 points in case of a correct go trial. Inter-trial-intervals (ITIs) varied between 1 and 16

seconds within the domain ITI = {1, 2, 4, 8, 16}. To reduce the overlap in brain responses

from one trial to the next, actual ITIs were randomly picked from a geometric distribution with

$p$ = 0.5. This meant a higher probability of shorter intervals (1 and 2 s) and occasional very

long intervals (8 and 16 s). The design was optimized to minimize the variance inflation factor

(VIF):

322                        $$VIF = var(E) / var(X),$$

the ratio of the mean estimation variance of all the regression weights (planning- and
execution-related regressors for each finger), and the mean estimation variance had these
regressors been estimated in isolation. For our design, the average VIF was quite low, on
average between 1 and 1.2, indicating that we could separate planning and execution
related activity without a large loss of experimental power.

*Design*
Participants underwent one fMRI session consisting of 10 functional runs and 1 anatomical
scan. In an event-related design, we randomly interleaved 3 types of repeated single finger
movements involving the thumb (1), the middle (3), and the little (5) fingers (e.g., 111111 for
thumb movement, Fig. 1A) and 3 types of multi finger sequences (e.g., 135315). The day
before the fMRI scan, participants familiarized themselves with the experimental apparatus
and the go/no-go paradigm in a short behavioral session of practice outside the scanner (5
blocks, about 15-30 min in total). For the behavioral practice, inter-trial intervals were kept to
a fixed 1 s to speed up the task, and participants were presented with different sequences
from what they would see while in the scanner. These 6-item sequences were randomly
selected from a pool of all possible permutations of the numbers 1, 3, and 5, with the
exclusion of sequences that contained consecutive repetitions of the same number. Given
that the current paper is concerned with the relationship between representations of simple
planning and execution, here we will focus only on the results for single finger movements.
The results for multi finger sequences will be reported in a future paper. Each single finger
trial type (e.g., 111111) was repeated 5 times (2 no-go and 3 go trials), totalling 30 trials per
functional run. Two periods of 10 s rests were added at the beginning and at the end of each
functional run to allow for signal relaxation and provide a better estimate of baseline
activation. Each functional run took about 5.5 minutes and the entire scanning session
(including the anatomical scan and setup time) lasted for about 75 minutes.

*Imaging data acquisition*
High-field functional magnetic resonance imaging (fMRI) data were acquired on a 7T
Siemens Magnetom scanner with a 32-channel head coil at Western University (London
Ontario, Canada). The anatomical T1-weighted scan of each participant was acquired
halfway through the scanning session (after the first 5 functional runs) using a Magnetization-

Prepared Rapid Gradient Echo sequence (MPRAGE) with voxel size of 0.75x0.75x0.75 mm

isotropic (field of view = 208 x 157 x 110 mm [A-P; R-L; F-H], encoding direction coronal). To

measure the blood-oxygen-level dependent (BOLD) responses in human participants, each

functional scan (330 volumes) used the following sequence parameters: GRAPPA 3, multi-

band acceleration factor 2, repetition time [TR] = 1.0 s, echo time [TE] = 20 ms, flip angle

[FA] = 30 deg, slice number: 44, voxel size: 2x2x2 mm isotropic. To estimate and correct for

magnetic field inhomogeneities, we also acquired a gradient echo field map with the following

parameters: transversal orientation, field of view: 210 x 210 x 160 mm, 64 slices, 2.5 mm

thickness, TR = 475 ms, TE = 4.08 ms, FA = 35 deg.

*Preprocessing and univariate analysis*

Preprocessing of the functional data was performed using SPM12 (fil.ion.ucl.ac.uk/spm) and

custom MATLAB code. This included correction for geometric distortions using the gradient

echo field map (58), and motion realignment to the first scan in the first run (3 translations: x,

y, z; 3 rotations: pitch, roll yaw). Due to the short TR, no slice timing corrections were applied.

The functional data were co-registered to the anatomical scan, but no normalization to a

standard template or smoothing was applied. To allow magnetization to reach equilibrium,

the first four volumes of each functional run were discarded. The pre-processed images were

analyzed with a general linear model (GLM). We defined separate regressors for each

combination of the 6 finger movements (single, multi) x 2 phases (preparation, movement).

To control for the effect of potential spill-over of movement execution activity on the preceding

planning activity, we also estimated a separate GLM with separate regressors for the

preparation phases of go and no-go trials, resulting in a total of 18 regressors (12 go + 6 no-

go), plus the intercept, for each run. Each regressor consisted of a boxcar function (on for 2

s of each phase duration and off otherwise) convolved with a two-gamma canonical

hemodynamic response function with a peak onset at 5 s and a post-stimulus undershoot

minimum at 10 s (Fig. 1B). Given the relatively low error rates (single: 8.51 ± 1.52 %, multi:

17.21 ± 3.38 %; timing errors, single: 7.58 ± 0. 62 %, multi: 10.23 ± 0.85 %; press errors,

single: 1.18 ± 0.26 %, multi: 9.04 ± 1.03 %), all trials were included to estimate the regression

coefficients, regardless of whether the execution was correct or erroneous. Ultimately, the

first-level analysis resulted in activation images (beta maps) for each of the 18 conditions per

run, for each of the participants.

*Surface reconstruction and ROI definition*

Individual subject's cortical surfaces were reconstructed using Freesurfer (59). First, we extracted the white-gray matter and pial surfaces from each participant's anatomical image. Next, we inflated each surface into a sphere and aligned it using sulcal depth and curvature information to the Freesurfer average atlas (Fischl et al., 1999). Both hemispheres in each participant were then resampled into Workbench's 164k vertex grid. This allowed us to compare similar areas of the cortical surface in each participant by selecting the corresponding vertices on the group atlas. Anatomical regions of interest (ROIs) were defined using a probabilistic cytoarchitectonic atlas (61) projected onto the common group surface. Our main ROIs were defined bilaterally as follows: primary motor cortex (M1) was defined by including nodes with the highest probability of belonging to Brodmann area (BA) 4 within 2 cm above and below the hand knob anatomical landmark (62); primary somatosensory cortex (S1) was defined by the nodes related to BA 1, 2 and 3; dorsal premotor cortex (PMd) was defined as the lateral part of the middle frontal gyrus; finally, the anterior part of the superior parietal lobule (aSPL) included areas anterior, superior and ventral to the intraparietal sulcus (IPS). ROI definition was carried out separately in each subject using FSL's subcortical segmentation. When resampling functional onto the surface, to avoid contamination between M1 and S1 activities, we excluded voxels with more than 25% of their volume in the grey matter on the opposite side of the central sulcus.

*Multivariate distance analysis*

To detect single finger representations across the cortical surface, we used representational similarity analysis (RSA; Diedrichsen and Kriegeskorte, 2017; Walther et al., 2016) with a surface-based searchlight approach (64). For each node, we selected a region (the searchlight) corresponding to 100 voxels (12 mm disc radius) in the gray matter and computed cross-validated Mahalanobis (crossnobis, Walther et al., 2016) dissimilarities between pairs of evoked activity patterns (beta estimates from first level GLM) of single finger sequences, during both preparation and movement phases. Prior to calculating the dissimilarities, beta weights for each condition were spatially pre-whitened (i.e., weighted by the matrix square root of the noise covariance matrix, as estimated from the residuals of the GLM. The noise covariance matrix was slightly regularized towards a diagonal matrix (65).

Multivariate pre-whitening has been shown to increase the reliability of dissimilarity estimates

(39). The resulting analyses (one RDM per participant containing the dissimilarities between

the three single fingers during planning and execution: 6 conditions, 15 dissimilarity pairs)

were then assigned to the central node and the searchlight was moved across all nodes

across the surface sheet obtaining a cortical map (Fig. 2B-2D). Cross-validation ensures the

distances estimates are unbiased, such that if two patterns differ only by measurement noise,

the mean of the estimated value would be zero. This also means that estimates can

sometimes become negative. Therefore, dissimilarities significantly larger than zero indicate

that two patterns are reliably distinct, similar to an above-chance performance in a cross-

validated pattern-classification analysis. Additionally, to the searchlight analysis, the

multivariate analysis was conducted separately for each anatomically defined ROI (e.g., Fig.

4A).

*Correlation between behavioral and neural distances*

To ensure that our planning results were not contaminated by unwanted micro-movements

during the preparation phase, we calculated the behavioral distance between sequences on

the basis of keyboard force data and correlated behavioral and neural distances. For

behavioral distances, we first extracted force data (2 ms temporal resolution, smoothed with

a gaussian kernel of 9.42 full width at half maximum, FWHM) and binned it in 10 ms steps

(down sampling largely due to memory constraints) for both the preparation and movement

phases (Fig. 3A). Next, for each subject, we calculated the mean (5) and the standard

deviation (5) of the time-averaged force of each finger for each condition (3 sequences x 2

phases = 6) and block (10). These subject-specific finger force patterns (60 x 10) were

multivariately pre-whitened using their covariance matrix. Finally, we calculated the cross-

validated squared Euclidean distances for each condition (6 x 6 RDM) and averaged

distances between the 3 finger movements for each phase (preparation, movement). These

mean finger force distances for each subject were correlated with the mean voxel activity

distances from the two phases for 2 ROIs (M1 and S1, Fig. 3C-3D).

*Pattern component modelling correlation models*

We used pattern component modelling (PCM) to quantify the correspondence of sequence-

specific activity patterns across planning and execution (43). This method has been shown

to be advantageous in estimating correlations. In contrast to simple Pearson's or cross-

validated correlation estimated from raw activity patterns, PCM separately models the noise

and signal components. We created 100 correlation models with correlations in the range [0–

1] in equal step sizes and assessed the likelihood of the observed data from each participant

under each model. Fig. 4D shows average log-likelihoods for each model, relative to the

mean log-likelihood across models. Differences between the log-likelihoods can be

interpreted as log-Bayes factors. Group inferences were performed using a simple t-test on

log-likelihoods. To compare each model to the best fitting model, we had to correct for the

bias arising from picking the best model and testing it on the same data: We used n-1

subjects to determine the group winning model, and then chose the log-likelihood of this

model for the left-out subject (for whom this model may not be the best one) as the likelihood

for the "best" model. This was repeated across all subjects and a one-sided paired-sample

$t$-test was performed on the recorded log-likelihood and every other model. This test revealed

which of the correlation models were significantly worse (i.e., associated with a lower log-

likelihood) than the winning model that was independently estimated via cross-validation.

## References

1.   S. W. Keele, J. J. Summers, "The Structure of Motor Programs" in *Motor Control*,
(Elsevier, 1976), pp. 109–142.

2.   S. W. Keele, Movement control in skilled motor performance. *Psychological Bulletin*
**70**, 387–403 (1968).

3.   D. A. Rosenbaum, Human movement initiation: Specification of arm, direction, and
extent. *Journal of Experimental Psychology: General* **109**, 444–474 (1980).

4.   S. T. Klapp, Motor Response Programming During Simple and Choice Reaction
Time: The Role of Practice. *Journal of Experimental Psychology: Human Perception*
*and Performance* **21**, 1015–1027 (1995).

5.   S. T. Klapp, C. I. Erwin, Relation between programming time and duration of the
response being programmed. *Journal of Experimental Psychology: Human*
*Perception and Performance* **2**, 591–598 (1976).

6.   A. M. Haith, J. Pakpoor, J. W. Krakauer, Independence of Movement Preparation and
Movement Initiation. *Journal of Neuroscience* **36**, 3007–3015 (2016).

7.   G. Ariani, J. Diedrichsen, Sequence learning is driven by improvements in motor
planning. *Journal of Neurophysiology* **121**, 2088–2100 (2019).

8.   A. L. Wong, A. M. Haith, Motor planning flexibly optimizes performance under
uncertainty about task goals. *Nature Communications* **8**, 14624 (2017).

9.   R. M. Hardwick, A. D. Forrence, J. W. Krakauer, A. M. Haith, Time-dependent
competition between goal-directed and habitual response preparation. *Nature*
*Human Behaviour* **3**, 1252–1262 (2019).

10.  C. Ghez, *et al.*, Discrete and continuous planning of hand movements and isometric
force trajectories. *Experimental Brain Research* **115**, 217–233 (1997).

11.  P. Cisek, J. F. Kalaska, Neural correlates of mental rehearsal in dorsal premotor
cortex. *Nature* **431**, 993–6 (2004).

12.  P. Cisek, J. F. Kalaska, Neural mechanisms for interacting with a world full of action
choices. *Annual review of neuroscience* **33**, 269–298 (2010).

13.  E. Hoshi, J. Tanji, Differential involvement of neurons in the dorsal and ventral
premotor cortex during processing of visual signals for action planning. *Journal of*
*neurophysiology* **95**, 3596–616 (2006).

14.    E. Hoshi, J. Tanji, Differential roles of neuronal activity in the supplementary and
500            presupplementary motor areas: from information retrieval to motor planning and
501            execution. *Journal of neurophysiology* **92**, 3482–99 (2004).

502    15.    R. A. Andersen, H. Cui, Intention, Action Planning, and Decision Making in Parietal-
503            Frontal Circuits. *Neuron* **63**, 568–83 (2009).

504    16.    H. Cui, R. A. Andersen, Posterior Parietal Cortex Encodes Autonomously Selected
505            Motor Plans. *Neuron* **56**, 552–559 (2007).

506    17.    H. Cui, R. A. Andersen, Different Representations of Potential and Selected Motor
507            Plans by Distinct Parietal Areas. *Journal of Neuroscience* **31**, 18130–18136 (2011).

508    18.    J. P. Gallivan, D. A. McLean, K. F. Valyear, C. E. Pettypiece, J. C. Culham, Decoding
509            Action Intentions from Preparatory Brain Activity in Human Parieto-Frontal Networks.
510            *Journal of Neuroscience* **31**, 9599–9610 (2011).

511    19.    G. Ariani, M. F. Wurm, A. Lingnau, Decoding Internally and Externally Driven
512            Movement Plans. *Journal of Neuroscience* **35**, 14160–14171 (2015).

513    20.    J. P. Gallivan, I. S. Johnsrude, J. R. Flanagan, Planning Ahead: Object-Directed
514            Sequential Actions Decoded from Human Frontoparietal and Occipitotemporal
515            Networks. *Cerebral Cortex* **26**, bhu302 (2015).

516    21.    J. P. Gallivan, D. A. McLean, F. W. Smith, J. C. Culham, Decoding Effector-
517            Dependent and Effector-Independent Movement Intentions from Human Parieto-
518            Frontal Brain Activity. *Journal of Neuroscience* **31**, 17149–17168 (2011).

519    22.    J. P. Gallivan, D. A. McLean, J. R. Flanagan, J. C. Culham, Where One Hand Meets
520            the Other: Limb-Specific and Action-Dependent Movement Plans Decoded from
521            Preparatory Signals in Single Human Frontoparietal Brain Areas. *Journal of*
522            *Neuroscience* **33**, 1991–2008 (2013).

523    23.    L. Turella, *et al.*, Beta band modulations underlie action representations for
524            movement planning. *NeuroImage* **136**, 197–207 (2016).

525    24.    F. T. M. Leoné, T. Heed, I. Toni, W. P. Medendorp, Understanding effector selectivity
526            in human posterior parietal cortex by combining information patterns and activation
527            measures. *The Journal of neuroscience* **34**, 7102–12 (2014).

528    25.    S. Vyas, M. D. Golub, D. Sussillo, K. V Shenoy, Computation Through Neural
529            Population Dynamics. *Annual Review of Neuroscience* **43**, 249–275 (2020).

26.    M. M. Churchland, J. P. Cunningham, M. T. Kaufman, S. I. Ryu, K. V. Shenoy,

Cortical Preparatory Activity: Representation of Movement or First Cog in a

Dynamical Machine? *Neuron* **68**, 387–400 (2010).

27.   K. V Shenoy, M. Sahani, M. M. Churchland, Cortical control of arm movements: a

dynamical systems perspective. *Annual Review of Neuroscience* **36**, 337–359

(2013).

28.   N. Ejaz, M. Hamada, J. Diedrichsen, Hand use predicts the structure of

representations in sensorimotor cortex. *Nature Neuroscience* **18**, 1034–1040 (2015).

29.   A. Yokoi, S. A. Arbuckle, J. Diedrichsen, The role of human primary motor cortex in

the production of skilled finger sequences. *Journal of Neuroscience* **38**, 1430–1442

(2018).

30.   J. Diedrichsen, T. Wiestler, J. W. Krakauer, Two distinct ipsilateral cortical

representations for individuated finger movements. *Cerebral Cortex* **23**, 1362–1377

(2013).

31.   T. Wiestler, S. Waters-Metenier, J. Diedrichsen, Effector-independent motor

sequence representations exist in extrinsic and intrinsic reference frames. *Journal of*

*Neuroscience* **34**, 5054–5064 (2014).

32.   A. Yokoi, J. Diedrichsen, Neural Organization of Hierarchical Motor Sequence

Representations in the Human Neocortex. *Neuron*, 1–13 (2019).

33.   S. A. Arbuckle, *et al.*, Structure of population activity in primary motor cortex for

single finger flexion and extension. *The Journal of Neuroscience*, JN-RM-0999-20

(2020).

34.   S. A. Arbuckle, A. Yokoi, J. A. Pruszynski, J. Diedrichsen, Stability of representational

geometry across a wide range of fMRI activity levels. *NeuroImage* **186**, 155–163

(2019).

35.   L. Huber, *et al.*, Sub-millimeter fMRI reveals multiple topographical digit

representations that form action maps in human motor cortex. *NeuroImage* **208**,

116463 (2020).

36.   G. Ariani, N. N. Oosterhof, A. Lingnau, Time-resolved decoding of planned delayed

and immediate prehension movements. *Cortex* **99** (2018).

37.   J. Tanji, E. V. Evarts, Anticipatory activity of motor cortex neurons in relation to

direction of an intended movement. *Journal of Neurophysiology* **39**, 1062–1068

(1976).

38.    D. J. Crammond, J. F. Kalaska, Prior information in motor and premotor cortex:

564           activity during the delay period and effect on pre-movement activity. *Journal of*

565           *neurophysiology* **84**, 986–1005 (2000).

566    39.    A. Walther, *et al.*, Reliability of dissimilarity measures for multi-voxel pattern analysis.

567           *NeuroImage* **137**, 188–200 (2016).

568    40.    J. Diedrichsen, *et al.*, Comparing representational geometries using whitened

569           unbiased-distance-matrix similarity (2020).

570    41.    N. N. Oosterhof, *et al.*, Surface-Based Information Mapping Reveals Crossmodal

571           Vision – Action Representations in Human Parietal and Occipitotemporal Cortex

572           Surface-Based Information Mapping Reveals Crossmodal Vision – Action

573           Representations in Human Parietal and Occipitotemporal. *Journal of neurophysiology*

574           **104**, 1077–89 (2012).

575    42.    M. T. Kaufman, M. M. Churchland, S. I. Ryu, K. V Shenoy, Cortical activity in the null

576           space: permitting preparation without movement. *Nature Neuroscience* **17**, 440–448

577           (2014).

578    43.    J. Diedrichsen, A. Yokoi, S. A. Arbuckle, Pattern component modeling: A flexible

579           approach for understanding the representational structure of brain activity patterns.

580           *NeuroImage* **180**, 119–133 (2018).

581    44.    G. E. Alexander, M. D. Crutcher, Neural representations of the target (goal) of

582           visually guided arm movements in three motor areas of the monkey. *Journal of*

583           *Neurophysiology* **64**, 164–178 (1990).

584    45.    A. Riehle, J. Requin, Monkey primary motor and premotor cortex: single-cell activity

585           related to prior information about direction and extent of an intended movement.

586           *Journal of neurophysiology* **61**, 534–49 (1989).

587    46.    J. D. Coulter, E. G. Jones, Differential distribution of corticospinal projections from

588           individual cytoarchitectonic fields in the monkey. *Brain Research* **129**, 335–340

589           (1977).

590    47.    J.-A. Rathelot, P. L. Strick, Muscle representation in the macaque motor cortex: An

591           anatomical perspective. *Proceedings of the National Academy of Sciences* **103**,

592           8257 LP – 8262 (2006).

593    48.    G. L. Widener, P. D. Cheney, Effects on Muscle Activity From Microstimuli Applied to

594           Somatosensory and Motor Cortex During Voluntary Movement in the Monkey. *Journal*

*of Neurophysiology* **77**, 2446–2465 (1997).

49.   E. Azim, K. Seki, Gain control in the sensorimotor system. *Current Opinion in*

*Physiology* **8**, 177–187 (2019).

50.   A. Starr, L. G. Cohen, 'Gating' of somatosensory evoked potentials begins before the

onset of voluntary movement in man. *Brain Research* **348**, 183–186 (1985).

51.   C. E. Chapman, M. C. Bushnell, D. Miron, G. H. Duncan, J. P. Lund, Sensory

perception during movement in man. *Experimental brain research* **68**, 516–524

(1987).

52.   W. Jiang, Y. Lamarre, C. E. Chapman, Modulation of cutaneous cortical evoked

potentials during isometric and isotonic contractions in the monkey. *Brain Research*

**536**, 69–78 (1990).

53.   K. Seki, E. E. Fetz, Gating of sensory input at spinal and cortical levels during

preparation and execution of voluntary movement. *Journal of Neuroscience* **32**, 890–

902 (2012).

54.   M. Jafari, *et al.*, The human primary somatosensory cortex encodes imagined

movement in the absence of sensory information. *Communications biology* **3**, 1–7

(2020).

55.   T. Umeda, T. Isa, Y. Nishimura, The somatosensory cortex receives information about

motor output. *Science Advances* **5** (2019).

56.   G. F. Elsayed, A. H. Lara, M. T. Kaufman, M. M. Churchland, J. P. Cunningham,

Reorganization between preparatory and movement population responses in motor

cortex. *Nature Communications*, 13239 (2016).

57.   N. K. Logothetis, J. Pauls, M. Augath, T. Trinath, A. Oeltermann, Neurophysiological

investigation of the basis of the fMRI signal. *Nature* **412**, 150–7 (2001).

58.   C. Hutton, *et al.*, Image Distortion Correction in fMRI: A Quantitative Evaluation.

*NeuroImage* **16**, 217–240 (2002).

59.   A. M. Dale, B. Fischl, M. I. Sereno, Cortical Surface-Based Analysis: I. Segmentation

and Surface Reconstruction. *NeuroImage* **9**, 179–194 (1999).

60.   B. Fischl, M. I. Sereno, R. B. H. Tootell, A. M. Dale, High-resolution intersubject

averaging and a coordinate system for the cortical surface. *Human Brain Mapping* **8**,

272–284 (1999).

61.   B. Fischl, *et al.*, Cortical folding patterns and predicting cytoarchitecture. *Cerebral*

627            *Cortex* **18**, 1973–1980 (2008).

628   62.   T. Yousry, Localization of the motor hand area to a knob on the precentral gyrus. A

629            new landmark. *Brain* **120**, 141–157 (1997).

630   63.   J. Diedrichsen, N. Kriegeskorte, Representational models: A common framework for

631            understanding encoding, pattern-component, and representational-similarity

632            analysis. *PLOS Computational Biology* **13**, e1005508 (2017).

633   64.   N. N. Oosterhof, T. Wiestler, P. E. Downing, J. Diedrichsen, A comparison of volume-

634            based and surface-based multi-voxel pattern analysis. *NeuroImage* **56**, 593–600

635            (2011).

636   65.   O. Ledoit, M. Wolf, Honey, I shrunk the sample covariance matrix. *The Journal of*

637            *Portfolio Management* **30**, 110–119 (2004).

## Supplementary figures

**Figure S1 | Activation and distance analyses using planning of both go and no-go trials. A.** Activation map (percent signal change) for the contrast planning>baseline. The selected area of interest is the same as shown in purple in the inset of Fig. 2A. **B.** Crossnobis distance searchlight map for movement planning. **C.** Same as A, but for the contrast execution>baseline. **D.** Same as B, but for movement execution. **E-F.** Cross-section analysis corresponding to the same area shown in white in the inset of Fig. 2A. **E.** Mean percent signal change (± SEM) during planning (orange) and execution (blue). **F.** mean crossnobis distance (± SEM). Horizontal bars indicate significance ($p < 0.05$) in a 2-sided one-sample $t$-test against zero. All other figure conventions are the same as in Fig. 2.