Motor planning brings human primary somatosensory cortex into action-specific preparatory states

Giacomo Ariani[1,2,*], J. Andrew Pruszynski[1,4,5,6], Jörn Diedrichsen[1,2,3]

[1]The Brain and Mind Institute, [2]Department of Computer Science, [3]Department of Statistical and Actuarial Sciences, [4]Department of Physiology and Pharmacology, [5]Department of Psychology, [6]Robarts Research Institute, Western University, London, ON N6A3K7, Canada.

*Correspondence should be addressed to Giacomo Ariani at gariani@uwo.ca, Western Interdisciplinary Research Building, London, ON N6A 3K7, Canada.

Running title. Motor planning in human sensorimotor cortex.

Word count. Abstract: 218; Significance statement: 114; Introduction: 601; Discussion: 810. 28 pages, 4 figures, 0 tables, 1 supplementary figure.

Author contributions. G.A. and J.D. designed research; G.A. performed research; G.A. analyzed data; G.A. drafted the manuscript; G.A., J.A.P, and J.D. edited the manuscript.

Acknowledgements. This work was supported by a NSERC Discovery Grant (RGPIN-2016-04890) awarded to J.D., and the Canada First Research Excellence Fund (BrainsCAN). The authors wish to thank Eva Berlot for helpful discussions and contributions to data analysis.

Disclosures. The authors declare no conflicts of interest.

1 Abstract

Motor planning plays a critical role in producing fast and accurate movement. Yet, the neural
processes that occur in human primary motor and somatosensory cortex during planning,
and how they relate to those during movement execution, remain poorly understood. Here
we used 7T functional magnetic resonance imaging (fMRI) and a delayed movement
paradigm to study single finger movement planning and execution. The inclusion of no-go
trials and variable delays allowed us to separate what are typically overlapping planning and
execution brain responses. Although our univariate results show widespread deactivation
during finger planning, multivariate pattern analysis revealed finger-specific activity patterns
in contralateral primary somatosensory cortex (S1), which predicted the planned finger
action. Surprisingly, these activity patterns were similarly strong to those found in
contralateral primary motor cortex (M1). Control analyses ruled out the possibility that the
detected information was an artifact of subthreshold movements during the preparatory
delay. Furthermore, we observed that finger-specific activity patterns during planning were
highly correlated to those during execution. These findings reveal that motor planning
activates the specific S1 and M1 circuits that are engaged during the execution of a finger
press, while activity in both regions is overall suppressed. We propose that preparatory states
in S1 may improve movement control through changes in sensory processing or via direct
influence of spinal motor neurons.
20

21 Key words

22 Motor planning; Sensorimotor cortex; Finger control; fMRI; Representational geometry.
23

Significance statement

Motor planning is important for good behavior performance, yet it is unclear which neural processes underlie the preparation of the nervous system for an upcoming movement. Using high-resolution functional neuroimaging, we investigated how motor planning for finger presses changes the activity state in primary motor and primary somatosensory cortex, and how brain responses during planning and execution relate to each other. We show that planning leads to finger-specific activation in both M1 and S1, which is highly similar to the finger-specific activity patterns elicited during execution. Our findings suggest that S1 is being specifically prepared for an upcoming action, either to actively contribute to the outflowing motor command or to enable action-specific sensory gating.

Introduction

Animals are capable of generating a wide variety of dexterous behaviors accurately and effortlessly on a daily basis. This remarkable ability relies on the motor system reaching the appropriate preparatory state before each movement is initiated. At the level of behavior, the process of motor programming, or planning, has long been shown to be beneficial to performance (Keele, 1968; Keele and Summers, 1976; Rosenbaum, 1980), leading to faster reaction times (Klapp and Erwin, 1976; Klapp, 1995; Haith et al., 2016) and more accurate response selection (Ghez et al., 1997; Wong and Haith, 2017; Ariani and Diedrichsen, 2019; Hardwick et al., 2019). The behavioral study of motor planning led to neurophysiological investigations showing the presence of preparatory signals in the patterns of neuronal firing in the dorsal premotor cortex, PMd (Cisek and Kalaska, 2004, 2010; Hoshi and Tanji, 2006), the supplementary motor area, SMA (Hoshi and Tanji, 2004), and the posterior parietal cortex, PPC (Cui and Andersen, 2007, 2011; Andersen and Cui, 2009). Building on this work, human neuroimaging studies have shown that activity in parieto-frontal brain regions during planning of prehension movements can be used to decode several movement properties such as grip type (Gallivan et al., 2011b; Ariani et al., 2015), action order (Gallivan et al., 2015), and effector (Gallivan et al., 2011a, 2013; Leoné et al., 2014; Turella et al., 2016). At the level of neural population dynamics (Vyas et al., 2020), motor planning can be understood as bringing the neuronal state towards an ideal preparatory point. Once this state is reached and the execution is triggered, the intrinsic dynamics of the system then let the movement unfold (Churchland et al., 2010; Shenoy et al., 2013).

Preparatory neural processes have not only been observed in premotor and parietal areas, but also in primary motor cortex (M1, Tanji and Evarts, 1976; Crammond and Kalaska, 2000; Ariani et al., 2018). In contrast, the degree to which primary somatosensory cortex (S1) receives information about the planned movement before movement onset is less clear. S1 is often considered to be mostly concerned with processing incoming sensory information from tactile and proprioceptive receptors arising after movement onset. Consistent with this notion, previous fMRI studies have not been able to detect the presence of planning-related information in this area (Gallivan et al., 2011a, 2011b, 2013, 2015). Very recently, however, a human electrocorticography (ECoG) study challenged this notion, suggesting a possible role for S1 in cognitive-motor imagery (Jafari et al., 2020). The authors recorded neural activity

from S1 while a tetraplegic participant imagined reaching movements and found that S1

neurons encoded movement direction during motor imagery in the absence of actual

sensations. Another recent ECoG study in non-human primates (Umeda et al., 2019) showed

grasp-specific information in the signals from S1 well before movement initiation, and only

slightly later than in M1. However, it remains unknown whether S1 plays an role during motor

planning in human participants with an intact sensory system. Furthermore, we currently do

not know how the signals during action preparation relate to those during execution, a fact

that could provide important insight into the role these signals may play.

Therefore, here we designed a high-field (7T) fMRI experiment to study what brain

regions underlie the planning of individual finger presses and how these brain

representations relate to those during execution. We used variable delays between an

instructing cue and a go signal, as well randomly interspersed no-go trials, to temporally

separate the evoked responses to movement planning and execution. Using advanced

multivariate pattern analyses we were able to examine the relationship between the fMRI

patterns related to planned and executed finger actions.

Results

Deactivation in sensorimotor regions during planning of finger actions

We instructed 22 participants to plan and execute repeated keypresses with individual

fingers of their right hand on a keyboard device while being scanned with 7T fMRI. The key

to be pressed corresponded to one of three fingers and was cued during the preparation

phase by numbers (1 = thumb, 3 = middle, 5 = little, e.g., Fig. 1A) presented on a computer

screen that was visible to the participants lying in the scanner through an angled mirror. After

a variable delay (4–8 seconds), participants received a color cue indicating whether to press

the planned finger (go trials), or whether to withhold the response (no-go trials). Upon the go

cue, participants had to initiate the correct response as fast as possible and make 6 presses

of the designated finger, before receiving accuracy points for reward (see Methods). To

control for involuntary overt movements during the preparation phase, we required

participants to maintain a steady force on all of the keys during the delay, which was closely

monitored online. To ensure that planning results would not be biased by the subsequent

execution, we restricted all our analyses of the preparation phase to no-go trials only (see

Methods). First, we asked which brain regions showed an evoked response during the

planning of finger presses (e.g., Fig. 1B).

Figure 1 | fMRI task and BOLD responses. A. Example trial with timing information. Background colors indicate different experimental phases (yellow = preparation; green = move (go) or stay (no-go); purple = reward; gray = inter-trial interval, ITI). B. Group-averaged BOLD response (N = 22) for go (blue) and no-go (orange) trials in a region that shows no planning evoked activity (Left M1, top), and one that shows some planning evoked activity (Left aSPL, bottom). Shaded areas indicate standard error of the mean (SEM). Background colors correspond to trial phases as in A.

103 We focused our analysis on the lateral aspect of the contralateral (left) hemisphere (purple

104 and white areas of the Fig. 2 inset) which included primary motor and somatosensory cortex,

105 as well as the premotor cortex and anterior parietal cortical regions. To examine brain

106 activation during finger planning, we performed a univariate contrast of the preparation

phase (across the three fingers) versus the resting baseline (Fig. 2A). Overall, the instruction stimulus evoked little to no activation in our regions of interest (ROIs, Fig. 2A). In fact, compared to resting baseline, we observed significant deactivation (Fig. 2E) in the primary motor cortex (M1, t_{21} = -7.592, p = 1.887e-07), the primary somatosensory cortex (S1, t_{21} = -6.618, p = 1.491e-06), and the dorsal premotor cortex (PMd, t_{21}= -2.642, p = 0.015), In comparison, execution strongly activated M1 and S1 (Fig. 2C), with activation being significant in all our ROIs (Fig. 2E, all t_{21} > 14.469, all p < 2.153e-12).

Planning induces informative patterns in primary somatosensory and motor cortex

Although we found little univariate activation in our main ROIs, preparatory processes need not increase the overall activation in a region. Rather, the region could converge to a specific preparatory neural state (Churchland et al., 2010), while activity increments and decrements within the region (i.e., at a finer spatial scale) average each other out. In this case, information about planned movements would be present in the multivoxel activity patterns in that region. To test this idea, we calculated the crossnobis dissimilarity (see Methods) between activity patterns. Systematically positive values of this dissimilarity measure indicate that the patterns reliably differentiate between the different planned actions (Walther et al., 2016; Diedrichsen et al., 2020). Indeed, a surface-based searchlight approach (Oosterhof et al., 2012) revealed reliably positive crossnobis dissimilarity between the activity patterns related to planning of individual finger presses (Fig. 2B) in both M1 (t_{21} = 2.734, p = 0.012) and S1 (t_{21} = 2.987, p = 0.007, Fig. 2F). Surprisingly, the distribution of these informative patterns was highly similar to the distribution of information during execution (Fig. 2D). Visual inspection suggested that the informative patterns during planning may be concentrated more dorsally in M1 and S1 relative to execution. However, a Hotelling T^2 test revealed no systematic difference in the location of the peak vertex between planning and execution across subjects (M1: $T^2_{2,20}$ = 0.725, p = 0.712; S1: $T^2_{2,20}$ = 2.424, p = 0.335). Thus, our results show that information about single finger actions is already represented during motor planning in the same parts of the primary motor and somatosensory cortices that are engaged during execution of the presses. Given that we only used the activity estimates from no-go trials (~40% of total trials), this information cannot be explained by a spill-over from subsequent execution-related activity. An analysis using the estimates of planning activity from all trials yielded very similar results (see Fig. S1), demonstrating that we could separate planning from execution-related signals.

Figure 2 | Activation and distance analyses of movement planning and execution. The inset shows the inflated cortical surface of the contralateral (left) hemisphere, highlighting the area of interest (purple) and the strip used for cross-section analysis (white). Major sulci are indicated by white dotted lines. **A.** Univariate activation map (percent signal change) for the contrast planning>baseline (no-go trials only). **B.** Multivariate searchlight map of the mean crossnobis distance between the planning of the three fingers (no-go trials only). **C.** Same as A, but for the univariate contrast execution>baseline (go trials). **D.** Same as B, but for the mean crossnobis distance between fingers during execution. **E.** Cross-section analysis of the mean percent signal change (± SEM) during planning (orange) and execution (blue). The x-axis corresponds to the x coordinate (in mm) on the whole-brain flat map of the cortical surface. Horizontal bars indicate significance ($p < 0.05$) in a 2-sided one-sample t-test vs zero for selected ROIs. **F.** Same as E, but for the mean crossnobis distance (± SEM). Vertical dotted lines mark the approximate location of the main anatomical landmarks according to x coordinate. Pre-CS = precentral sulcus; CS = central sulcus; Post-CS = postcentral sulcus; Vertical dashed lines denote approximate boundaries of the main regions of interest (ROIs).

PMd = dorsal premotor cortex; M1 = primary motor cortex; S1 = primary somatosensory cortex; aSPL = anterior superior parietal lobule.

Activity patterns are not caused by small movements during the preparation phase

The presence of planning-related information in primary sensorimotor regions was surprising,

especially in S1, where it had not previously been reported in comparable fMRI studies

(Gallivan et al., 2011b, 2015). To ensure that these results were not caused by overt

movement, participants were instructed to maintain a steady force on the keyboard during

the preparation phase, such that we could monitor even the smallest involuntary preparatory

movements. Inspection of the average force profiles (Fig. 3A) revealed that participants were

successful in maintaining a stable force between 0.2 and 0.4 N during preparation. However,

averaging forces across trials may obscure small, idiosyncratic patterns visible during

individual trials (Fig. 3B) that could be used to distinguish the different movements. To test

for the presence of such patterns, we submitted both the mean and standard deviation of the

force traces on each finger to a multivariate dissimilarity analysis (see Methods). Indeed, this

sensitive analysis revealed that some participants showed small movement patterns

predictive of the upcoming finger (Fig. 3C). These differences, however, were ~200-300

times smaller than the average differences during execution (Fig. 3D). More importantly, the

magnitude of the behavioral differences for the preparation phase was uncorrelated to the

amount of planning information present in sensory-motor regions (both p-values for the slope

of the linear fit > 0.3). Even after correcting for the influence behavioral patterns, the

informative patterns in M1 and S1 remained significant, as shown by a significantly positive

intercept in the linear fit in Fig. 3C (M1: $p = 0.032$; S1: $p = 0.007$). Thus, the finding of finger-

specific activity patterns in M1 and S1 cannot be explained by small involuntary movements

during the preparation phase.

Figure 3 | Small involuntary movements do not explain preparatory activity patterns in M1 and S1. A. Mean finger force (± SEM) plotted in 10 ms bins, time-aligned to instruction onset (dotted vertical line) and end of the preparation phase (dashed vertical lines), separately for the three fingers and go (blue) and no-go (orange) trials. **B.** Example of an individual trial with a 6 s preparation phase, followed six presses of the little finger (green). Horizontal solid line denotes press threshold (1 N). Dash-dotted lines denote the boundaries of the finger pre-activation red area in Fig. 1A (see Methods). RT = reaction time; MT = movement time. **C.** Pearson's correlation (r) between behavioral and neural distances in M1 and S1 (see Methods) during the preparation phase (planning, orange). Each dot represents an individual participant (N = 22). Solid line shows linear regression, *p*-values refers to the slope of the line. **D.** Same as C, but during the movement phase (execution, blue).

Single finger activity patterns from planning to execution are positively correlated

How do planning-related activity patterns in M1 and S1 relate to the activity patterns observed during execution? Neurophysiological experiments have suggested that patterns of movement preparation are orthogonal – or uncorrelated – to the patterns underlying active movement (Kaufman et al., 2014). This arrangement allows movement preparation to occur without causing overt movement.

When we compared the planning- and execution-related activity patterns as measured with fMRI, a technique that samples neuronal activity at a much coarser spatial resolution, we found the opposite result. Planning- and execution-related patterns for the same finger were tightly related. Inspection of the representational dissimilarity matrices (RDMs) for M1 and S1 (Fig. 4A), clearly shows that the biggest difference was between planning and execution patterns, which can also be seen in a 3D representation of the representational geometry (principal component PC1 in Fig. 4B). Within each phase, the pattern for the thumb was more distinct than those for the other fingers, replicating previous results from execution alone (Ejaz et al., 2015; Yokoi et al., 2018). Importantly, however, a rotated view of the representational geometry (Fig. 4C) showed that the finger patterns were arranged in a congruent way, with planning and execution related activity patterns for the same finger being closer to one another.

To more precisely quantify the correspondence between planning and execution pattern for each finger, we used Pattern Component Modeling (PCM, Diedrichsen et al., 2018). Simple correlations between measured fMRI patterns are usually lower than the true correlation because of the biasing influence of measurement noise. PCM corrects for this effect by evaluating the likelihood of the data (taking into account the measurement noise), assuming true correlation between 0 and 1. From the individual fits, we found that the average best correlation model was at 0.83 (± 0.053 SEM) for M1 and at 0.81 (± 0.061 SEM) for S1 (Fig. 4D, dashed lines). Using a cross-validated approach (see Methods), we compared the log-likelihoods to test whether the best fitting model was significantly better than any of the other correlation models. In both ROIs, the best fitting model was significantly better than the zero-correlation model. From this we can conclude that the activity patterns across planning and execution were not completely uncorrelated (both $p < 0.005$). However, the best fitting model did not perform better than the one-correlation model (both $p > 0.1$). Thus, our data

198   us consistent with the idea that the activity patterns for planning and execution of finger

199   presses in S1 and M1 are simply a scaled version of each other.

Figure 4 | Correlated representations of single fingers across planning and execution. A. Representational dissimilarity matrices (RDMs) of the activity patterns evoked by the three fingers during the preparation (no-go planning, orange) and movement (execution, blue) phases, separately for the two main ROIs (M1, left; S1, right). **B.** Multidimensional scaling (MDS) projection of the RDMs in A highlighting the first principal component (PC 1, difference between planning and execution). The black cross denotes resting baseline (arbitrary units). **C.** Same as B, but rotated view to highlight the representational geometries for PC 2 and PC 3. **D.** Pattern component modelling (PCM) evaluation of models (x-axis) of different correlations between planning- and execution-related activity patterns. Shown is the group-average of the individual log-likelihood (± SEM) curve expressed as a difference from the mean log-likelihood across models. Horizontal gray bars indicate models that perform

statistically equivalently ($p > 0.05$) to the best fitting model (determined in a cross-validated fashion, see Methods). Red dots indicate points of individually best fitting correlations (N = 22). Red dashed lines denote the average winning models across participants. Dotted lines show the projections of horizontal bars and dashed lines on the respective axes.

202 By applying this method to every other model, we have evidence that the true (i.e., noiseless)

203 correlation between planning and execution finger-specific activity pattern was between

204 0.41–1.0 in M1 and between 0.54–1.0 in S1 (Fig. 4D, horizontal bars). In sum, at the resolution

205 of fMRI, the process of movement planning seems to induce a finger-specific pattern which

206 very similar, and possibly identical, to the pattern activated during movement execution.

209 Discussion

In the present study, we asked participants to produce repeated single finger presses while

undergoing 7T fMRI. We used variable preparatory delays and no-go trials to cleanly

dissociate the brain responses to the consecutive preparation and movement phases. We

found that information about planned finger actions is present in both S1 and M1 before

action onset, even though the overall level of activation in these regions was below resting

baseline. Moreover, while execution elicited much higher brain activation, the fine-grained,

finger-specific activity patterns were highly similar across planning and execution. Control

analyses confirmed that the observed results were not caused by pre-movement finger

activity.

Our finding that motor planning activates M1 in a finger-specific fashion was not

necessarily surprising given many neurophysiological studies reporting anticipatory activity

of M1 neurons related to movement intentions (Tanji and Evarts, 1976; Riehle and Requin,

1989; Alexander and Crutcher, 1990), as well as human neuroimaging showing shared

information between delayed and immediate movement plans (Ariani et al., 2018). In

contrast, the robust activity patterns related to single finger planning in S1 were more

surprising, given that this region has classically been associated with the passive processing

of somatosensory information from receptors in the skin, muscles, and tendons. So, what

could then be the role of S1 during movement planning? First, it is worth noting that there are

substantial projections from S1 (Brodmann area 3a) that terminate in the ventral horn of the

cortico-spinal tract (Coulter and Jones, 1977; Rathelot and Strick, 2006). Although stimulation

of area 3a in macaques typically fails to evoke overt movements (Widener and Cheney, 1997), it has been suggested that this population of cortico-motoneurons specifically projects to gamma motoneurons that control the sensitivity of muscle spindle afferents (Rathelot and Strick, 2006). Thus, it is possible that S1 plays an active role in movement generation by preparing the spindle apparatus in advance of the movement. Second, the finger-specific preparatory state in S1 may reflect the anticipation or prediction of the upcoming sensory stimulation, allowing for a movement-specific sensory gain control (Azim and Seki, 2019). Some sensory stimuli could become attenuated to maintain movement stability and filter out irrelevant or self-generated signals. Indeed, multiple studies have shown that both somatosensation and somatosensory-evoked potentials in S1 decrease during voluntary movement (Starr and Cohen, 1985; Chapman et al., 1987; Jiang et al., 1990; Seki and Fetz, 2012). Alternatively, sensory processing of the expected salient signals could be enhanced to improve movement execution and adaptation.

One may wonder why such action-specific encoding in S1 during planning has not been reported in previous fMRI studies. One answer may lie in types of movements: previous studies (Gallivan et al., 2011a, 2011b, 2013, 2015) have used either reaching or grasping movements, which may be associated with less distinguishable activity patterns in S1 than the individuated finger presses used here. Additionally, the sensitivity to detect preparatory signals may have been higher given the higher field strength (7T), spatial resolution, and the more sensitive multivariate analysis method (crossnobis dissimilarity vs. pattern classification, Walther et al., 2016).

Our second main finding – the close correspondence between finger-specific activity patterns across planning and execution – appears to be at odds with the idea that these two processes occupy orthogonal neural subspaces to avoid overt movement during planning (Kaufman et al., 2014; Elsayed et al., 2016). We think that there are at least two possible explanations for this. First, the divergence of results could be caused by the difference in behavioral paradigms. While the neuronal correlates of movement planning in non-human primates have largely been studied using upper limb movements, we used here individuated finger presses. If for single finger actions even single-neuron activity patterns are highly correlated between planning and execution, then overt movement during planning would need to be actively suppressed, for example through the deactivation that we observed around the central sulcus. An alternative and perhaps more likely explanation of the

discrepancy lies in the different measurement modalities. Orthogonality was observed in

electrophysiological recordings of individual neurons, whereas the fMRI measurements we

employed here mainly reflect excitatory postsynaptic potentials (Logothetis et al., 2001) and

average metabolic activity across hundreds of thousands of cortical neurons. Thus, it is

possible that planning pre-activates the specific cortical columns in M1 and S1 that are also

most active during movement of that finger. Within each of these cortical micro-circuits,

however, planning-related activity could still be orthogonal to the activity observed during

execution at the single neuron level (e.g., see Arbuckle et al., 2020, for a similar observation

for cortical representations of flexion and extension finger movements). This would suggest

a new hypothesis for the architecture of the sensory-motor system where movement planning

pre-activates the action-specific circuits in M1 and S1. However, it does so in a fashion that

the induced planning-related activity is, in terms of the firing output of neurons, orthogonal to

the patterns during execution.

Methods

Participants

Twenty-three right-handed neurologically healthy participants volunteered to take part in the experiment (13 F, 10 M; age 20–31 years, mean 23.43 years, SD 4.08 years). Criteria for inclusion were right-handedness and no prior history of psychiatric or neurological disorders. Handedness was assessed with the Edinburgh Handedness Inventory (mean 82.83, SD 9.75). All experimental procedures were approved by the Research Ethics Committee at Western University. Participants provided written informed consent to procedures and data usage and received monetary compensation for their participation. One participant withdrew before study completion and was excluded from data analysis (final N = 22).

Apparatus

Repeated right-hand finger presses were performed on a custom-made MRI-compatible keyboard device (Fig. 1A). Participants only used the tips of their fingers to press on the keys. The keys of the device did not move but force transducers underneath each key measured isometric force production at an update rate of 2 ms (Honeywell FS series; dynamic range 0-25 N; sampling 200 Hz). A keypress/release was detected when the force crossed a threshold of 1 N. The forces measured from the keyboard were low pass filtered to reduce noise induced by the MRI environment, amplified, and sent to PC for online task control and data recording.

Task

We used a task in which participants produced repeated keypresses with the tip of their right-hand fingers in response to numerical cues appearing on a computer screen (white outline, Fig. 1A). On each trial, a string of 6 numbers (instructing cue) instructed which finger press to plan (1 = thumb, 3 = middle, 5 = little).

The length of the preparation phase (yellow background in Fig. 1) was randomly sampled to be 4 s (56% of trials), 6 s (30%), or 8 s (14%). To limit and monitor unwanted movements during the preparation phase, we required participants to pre-activate their fingers by maintaining a steady force of around 0.2-0.3 N on all of the keys during the preparation phase. As a visual aid, we displayed a red area (between 0 and 0.5 N) and asked

participants to remain in the middle of that range with all the fingers (touching either boundary

of the red area would count as unwanted movement, thus incurring an error).

At the onset of the movement phase (green background), participants received a

color cue (go/no-go cue) indicating whether to perform the planned finger presses (blue

outline = go, p = 0.6), or not (orange outline = no-go, p = 0.4). The role of no-go trials was to

de-couple the hemodynamic response to the successive planning and execution events,

which would otherwise always overlap in fast fMRI designs due to the sluggishness of the

fMRI response. To encourage planning during the delay period, at the go cue the digits were

masked with asterisks, and participants had to perform the presses from memory.

Participants had 2.5 s to complete the movement phase, and a vanishing white bar under the

asterisks indicated how much time was left to complete all of the keypresses. Participants

received online feedback on the correctness of each press with asterisks turning either

green, for a correct press, or red, for incorrect presses. As long as the participants remained

within task constraints (i.e., 6 keypresses in less than 2.5 s), an exact movement speed was

not enforced. In no-go trials, participants were instructed to remain as still as possible

maintaining the finger pre-activation until the end of the movement phase (i.e., releasing any

of the keys would incur an error).

During the reward phase (0.5 s, purple background) points were awarded based on

performance and according to the following scheme: -1 point in case of no-go error or go

cue anticipation (timing errors); 0 points for pressing any wrong key (press error); 1 point in

case of a correct no-go trial; and 2 points in case of a correct go trial.

Inter-trial-intervals (ITI, gray background) were randomly drawn from {1, 2, 4, 8, 16 s}

with the respective proportion of trials {52%, 26%, 13%, 6%, 3%}.

Experiment design and structure/procedure

Our chosen distribution of preparation times, inter-trial intervals, and no-go trials, was

determined by minimizing the variance inflation factor (VIF) for a given length of scan:

$$VIF = \frac{var(E)}{var(X)}$$

Where *var(E)* is the mean estimation variance of all the regression weights (planning- and

execution-related regressors for each finger), and *var(X)* the mean estimation variance had

these regressors been estimated in isolation. The VIF quantifies the severity of

multicollinearity between model regressors by providing an index of how much the variance

of an estimated regression coefficient is increased because of collinearity. Large values for

VIF mean that model regressors are not independent of each other, whereas a VIF of 1 means

no inflation of variance. After optimizing the design, the VIF was quite low, on average around

1.15, indicating that we could separate planning and execution related activity without a large

loss of experimental power.

Participants underwent one fMRI session consisting of 10 functional runs and 1

anatomical scan. In an event-related design, we randomly interleaved 3 types of repeated

single finger presses involving the tip of the thumb (1), the middle (3), and the little (5) fingers

(e.g., 111111 for thumb presses, Fig. 1A) and 3 types of multi finger sequences (e.g.,

135315). The day before the fMRI scan, participants familiarized themselves with the

experimental apparatus and the go/no-go paradigm in a short behavioral session of practice

outside the scanner (5 blocks, about 15-30 min in total). For the behavioral practice, inter-

trial intervals were kept to a fixed 1 s to speed up the task, and participants were presented

with different sequences from what they would see while in the scanner. These 6-item

sequences were randomly selected from a pool of all possible permutations of the numbers

1, 3, and 5, with the exclusion of sequences that contained consecutive repetitions of the

same number. Given that the current paper is concerned with the relationship between

representations of simple planning and execution, here we will focus only on the results for

single finger actions. The results for multi finger sequences will be reported in a future paper.

Each single finger trial type (e.g., 111111) was repeated 5 times (2 no-go and 3 go trials),

totalling 30 trials per functional run. Two periods of 10 s rests were added at the beginning

and at the end of each functional run to allow for signal relaxation and provide a better

estimate of baseline activation. Each functional run took about 5.5 minutes and the entire

scanning session (including the anatomical scan and setup time) lasted for about 75 minutes.

Imaging data acquisition

High-field functional magnetic resonance imaging (fMRI) data were acquired on a 7T

Siemens Magnetom scanner with a 32-channel head coil at Western University (London

Ontario, Canada). The anatomical T1-weighted scan of each participant was acquired

halfway through the scanning session (after the first 5 functional runs) using a Magnetization-

Prepared Rapid Gradient Echo sequence (MPRAGE) with voxel size of 0.75x0.75x0.75 mm

isotropic (field of view = 208 x 157 x 110 mm [A-P; R-L; F-H], encoding direction coronal). To

measure the blood-oxygen-level dependent (BOLD) responses in human participants, each

functional scan (330 volumes) used the following sequence parameters: GRAPPA 3, multi-

band acceleration factor 2, repetition time [TR] = 1.0 s, echo time [TE] = 20 ms, flip angle

[FA] = 30 deg, slice number: 44, voxel size: 2x2x2 mm isotropic. To estimate and correct for

magnetic field inhomogeneities, we also acquired a gradient echo field map with the following

parameters: transversal orientation, field of view: 210 x 210 x 160 mm, 64 slices, 2.5 mm

thickness, TR = 475 ms, TE = 4.08 ms, FA = 35 deg.

Preprocessing and univariate analysis

Preprocessing of the functional data was performed using SPM12 (fil.ion.ucl.ac.uk/spm) and

custom MATLAB code. This included correction for geometric distortions using the gradient

echo field map (Hutton et al., 2002), and motion realignment to the first scan in the first run

(3 translations: x, y, z; 3 rotations: pitch, roll yaw). Due to the short TR, no slice timing

corrections were applied. The functional data were co-registered to the anatomical scan, but

no normalization to a standard template or smoothing was applied. To allow magnetization

to reach equilibrium, the first four volumes of each functional run were discarded. The pre-

processed images were analyzed with a general linear model (GLM). We defined separate

regressors for each combination of the 6 finger-actions (single, multi) x 2 phases

(preparation, movement). To control for the effect of potential overlap between execution

activity and the preceding planning activity, we also estimated a separate GLM with separate

regressors for the preparation phases of go and no-go trials, resulting in a total of 18

regressors (12 go + 6 no-go), plus the intercept, for each run. Each regressor consisted of a

boxcar function (on for 2 s of each phase duration and off otherwise) convolved with a two-

gamma canonical hemodynamic response function with a peak onset at 5 s and a post-

stimulus undershoot minimum at 10 s (Fig. 1B). Given the relatively low error rates (i.e.,

number of error trials over total number of trials, timing errors: 7.58 ± 0. 62 %; press errors:

1.18 ± 0.26 %, see Task above), all trials were included to estimate the regression

coefficients, regardless of whether the execution was correct or erroneous. Ultimately, the

first-level analysis resulted in activation images (beta maps) for each of the 18 conditions per

run, for each of the participants.

Surface reconstruction and ROI definition

Individual subject's cortical surfaces were reconstructed using Freesurfer (Dale et al., 1999).

First, we extracted the white-gray matter and pial surfaces from each participant's anatomical

image. Next, we inflated each surface into a sphere and aligned it using sulcal depth and

curvature information to the Freesurfer average atlas (Fischl et al., 1999). Both hemispheres

in each participant were then resampled into Workbench's 164k vertex grid. This allowed us

to compare similar areas of the cortical surface in each participant by selecting the

corresponding vertices on the group atlas. Anatomical regions of interest (ROIs) were

defined using a probabilistic cytoarchitectonic atlas (Fischl et al., 2008) projected onto the

common group surface. Our main ROIs were defined bilaterally as follows: primary motor

cortex (M1) was defined by including nodes with the highest probability of belonging to

Brodmann area (BA) 4 within 2 cm above and below the hand knob anatomical landmark

(Yousry, 1997); primary somatosensory cortex (S1) was defined by the nodes related to BA

1, 2 and 3; dorsal premotor cortex (PMd) was defined as the lateral part of the middle frontal

gyrus; finally, the anterior part of the superior parietal lobule (aSPL) included areas anterior,

superior and ventral to the intraparietal sulcus (IPS). ROI definition was carried out separately

in each subject using FSL's subcortical segmentation. When resampling functional onto the

surface, to avoid contamination between M1 and S1 activities, we excluded voxels with more

than 25% of their volume in the grey matter on the opposite side of the central sulcus.

Multivariate distance analysis

To detect single finger representations across the cortical surface, we used representational

similarity analysis (RSA; Diedrichsen and Kriegeskorte, 2017; Walther et al., 2016) with a

surface-based searchlight approach (Oosterhof et al., 2011). For each node, we selected a

region (the searchlight) corresponding to 100 voxels (12 mm disc radius) in the gray matter

and computed cross-validated Mahalanobis (crossnobis, Walther et al., 2016) dissimilarities

between pairs of evoked activity patterns (beta estimates from first level GLM) of single finger

sequences, during both preparation and movement phases. Prior to calculating the

dissimilarities, beta weights for each condition were spatially pre-whitened (i.e., weighted by

the matrix square root of the noise covariance matrix, as estimated from the residuals of the

GLM. The noise covariance matrix was slightly regularized towards a diagonal matrix (Ledoit

and Wolf, 2004). Multivariate pre-whitening has been shown to increase the reliability of

dissimilarity estimates (Walther et al., 2016). The resulting analyses (one RDM per participant

containing the dissimilarities between the three single fingers during planning and execution:

6 conditions, 15 dissimilarity pairs) were then assigned to the central node and the

searchlight was moved across all nodes across the surface sheet obtaining a cortical map

(Fig. 2B-2D). Cross-validation ensures the distances estimates are unbiased, such that if two

patterns differ only by measurement noise, the mean of the estimated value would be zero.

This also means that estimates can sometimes become negative. Therefore, dissimilarities

significantly larger than zero indicate that two patterns are reliably distinct, similar to an

above-chance performance in a cross-validated pattern-classification analysis. Additionally,

to the searchlight analysis, the multivariate analysis was conducted separately for each

anatomically defined ROI (e.g., Fig. 4A).

Correlation between behavioral and neural distances

To ensure that our planning results were not contaminated by unwanted micro-movements

during the preparation phase, we calculated the behavioral distance between the different

fingers on the basis of keyboard force data and correlated behavioral and neural distances.

For behavioral distances, we first extracted force data (2 ms temporal resolution, smoothed

with a gaussian kernel of 9.42 full width at half maximum, FWHM) and binned it in 10 ms

steps (down sampling largely due to memory constraints) for both the preparation and

movement phases (Fig. 3A). Next, for each subject, we calculated the mean (5) and the

standard deviation (5) of the time-averaged force of each finger for each condition (3

sequences x 2 phases = 6) and block (10). These subject-specific finger force patterns (60

x 10) were multivariately pre-whitened using their covariance matrix. Finally, we calculated

the cross-validated squared Euclidean distances for each condition (6 x 6 RDM) and

averaged distances between the 3 finger presses for each phase (preparation, movement).

These mean finger force distances for each subject were correlated with the mean voxel

activity distances from the two phases for 2 ROIs (M1 and S1, Fig. 3C-3D).

Pattern component modelling correlation models

We used pattern component modelling (PCM) to quantify the correspondence of finger-

specific activity patterns across planning and execution (Diedrichsen et al., 2018). This

method has been shown to be advantageous in estimating correlations. In contrast to simple

Pearson's or cross-validated correlation estimated from raw activity patterns, PCM separately

models the noise and signal components. We created 100 correlation models with

correlations in the range [0–1] in equal step sizes and assessed the likelihood of the observed

data from each participant under each model. Fig. 4D shows average log-likelihoods for each

model, relative to the mean log-likelihood across models. Differences between the log-

likelihoods can be interpreted as log-Bayes factors. Group inferences were performed using

a simple t-test on log-likelihoods. To compare each model to the best fitting model, we had

to correct for the bias arising from picking the best model and testing it on the same data:

We used n-1 subjects to determine the group winning model, and then chose the log-

likelihood of this model for the left-out subject (for whom this model may not be the best one)

as the likelihood for the "best" model. This was repeated across all subjects and a one-sided

paired-sample *t*-test was performed on the recorded log-likelihood and every other model.

This test revealed which of the correlation models were significantly worse (i.e., associated

with a lower log-likelihood) than the winning model that was independently estimated via

cross-validation.

References

Alexander GE, Crutcher MD (1990) Neural representations of the target (goal) of visually guided arm movements in three motor areas of the monkey. Journal of Neurophysiology 64:164–178.

Andersen RA, Cui H (2009) Intention, Action Planning, and Decision Making in Parietal-Frontal Circuits. Neuron 63:568–583.

Arbuckle SA, Weiler J, Kirk EA, Rice CL, Schieber M, Pruszynski JA, Ejaz N, Diedrichsen J (2020) Structure of population activity in primary motor cortex for single finger flexion and extension. The Journal of Neuroscience:JN-RM-0999-20.

Ariani G, Diedrichsen J (2019) Sequence learning is driven by improvements in motor planning. Journal of Neurophysiology 121:2088–2100.

Ariani G, Oosterhof NN, Lingnau A (2018) Time-resolved decoding of planned delayed and immediate prehension movements. Cortex 99.

Ariani G, Wurm MF, Lingnau A (2015) Decoding Internally and Externally Driven Movement Plans. Journal of Neuroscience 35:14160–14171.

Azim E, Seki K (2019) Gain control in the sensorimotor system. Current Opinion in Physiology 8:177–187.

Chapman CE, Bushnell MC, Miron D, Duncan GH, Lund JP (1987) Sensory perception during movement in man. Experimental brain research 68:516–524.

Churchland MM, Cunningham JP, Kaufman MT, Ryu SI, Shenoy K V. (2010) Cortical Preparatory Activity: Representation of Movement or First Cog in a Dynamical Machine? Neuron 68:387–400.

Cisek P, Kalaska JF (2004) Neural correlates of mental rehearsal in dorsal premotor cortex. Nature 431:993–996.

Cisek P, Kalaska JF (2010) Neural mechanisms for interacting with a world full of action choices. Annual review of neuroscience 33:269–298.

Coulter JD, Jones EG (1977) Differential distribution of corticospinal projections from individual cytoarchitectonic fields in the monkey. Brain Research 129:335–340.

Crammond DJ, Kalaska JF (2000) Prior information in motor and premotor cortex: activity during the delay period and effect on pre-movement activity. Journal of neurophysiology 84:986–1005.

Cui H, Andersen RA (2007) Posterior Parietal Cortex Encodes Autonomously Selected

Motor Plans. Neuron 56:552–559.

Cui H, Andersen RA (2011) Different Representations of Potential and Selected Motor Plans

by Distinct Parietal Areas. Journal of Neuroscience 31:18130–18136.

Dale AM, Fischl B, Sereno MI (1999) Cortical Surface-Based Analysis: I. Segmentation and

Surface Reconstruction. NeuroImage 9:179–194.

Diedrichsen J, Berlot E, Mur M, Schütt HH, Shahbazi M, Kriegeskorte N (2020) Comparing

representational geometries using whitened unbiased-distance-matrix similarity.

Diedrichsen J, Kriegeskorte N (2017) Representational models: A common framework for

understanding encoding, pattern-component, and representational-similarity analysis

Cichy R, ed. PLOS Computational Biology 13:e1005508.

Diedrichsen J, Yokoi A, Arbuckle SA (2018) Pattern component modeling: A flexible

approach for understanding the representational structure of brain activity patterns.

NeuroImage 180:119–133.

Ejaz N, Hamada M, Diedrichsen J (2015) Hand use predicts the structure of

representations in sensorimotor cortex. Nature Neuroscience 18:1034–1040.

Elsayed GF, Lara AH, Kaufman MT, Churchland MM, Cunningham JP (2016)

Reorganization between preparatory and movement population responses in motor

cortex. Nature Communications:13239.

Fischl B, Rajendran N, Busa E, Augustinack J, Hinds O, Yeo BTT, Mohlberg H, Amunts K,

Zilles K (2008) Cortical folding patterns and predicting cytoarchitecture. Cerebral

Cortex 18:1973–1980.

Fischl B, Sereno MI, Tootell RBH, Dale AM (1999) High-resolution intersubject averaging

and a coordinate system for the cortical surface. Human Brain Mapping 8:272–284.

Gallivan JP, Johnsrude IS, Flanagan JR (2015) Planning Ahead: Object-Directed Sequential

Actions Decoded from Human Frontoparietal and Occipitotemporal Networks.

Cerebral Cortex 26:bhu302.

Gallivan JP, McLean DA, Flanagan JR, Culham JC (2013) Where One Hand Meets the

Other: Limb-Specific and Action-Dependent Movement Plans Decoded from

Preparatory Signals in Single Human Frontoparietal Brain Areas. Journal of

Neuroscience 33:1991–2008.

Gallivan JP, McLean DA, Smith FW, Culham JC (2011a) Decoding Effector-Dependent and

Effector-Independent Movement Intentions from Human Parieto-Frontal Brain Activity.

Journal of Neuroscience 31:17149–17168.

Gallivan JP, McLean DA, Valyear KF, Pettypiece CE, Culham JC (2011b) Decoding Action

Intentions from Preparatory Brain Activity in Human Parieto-Frontal Networks. Journal

of Neuroscience 31:9599–9610.

Ghez C, Favilla M, Ghilardi MF, Gordon J, Bermejo R, Pullman S (1997) Discrete and

continuous planning of hand movements and isometric force trajectories. Experimental

Brain Research 115:217–233.

Haith AM, Pakpoor J, Krakauer JW (2016) Independence of Movement Preparation and

Movement Initiation. Journal of Neuroscience 36:3007–3015.

Hardwick RM, Forrence AD, Krakauer JW, Haith AM (2019) Time-dependent competition

between goal-directed and habitual response preparation. Nature Human Behaviour

3:1252–1262.

Hoshi E, Tanji J (2004) Differential roles of neuronal activity in the supplementary and

presupplementary motor areas: from information retrieval to motor planning and

execution. Journal of neurophysiology 92:3482–3499.

Hoshi E, Tanji J (2006) Differential involvement of neurons in the dorsal and ventral

premotor cortex during processing of visual signals for action planning. Journal of

neurophysiology 95:3596–3616.

Hutton C, Bork A, Josephs O, Deichmann R, Ashburner J, Turner R (2002) Image Distortion

Correction in fMRI: A Quantitative Evaluation. NeuroImage 16:217–240.

Jafari M, Aflalo T, Chivukula S, Kellis SS, Salas MA, Norman SL, Pejsa K, Liu CY, Andersen

RA (2020) The human primary somatosensory cortex encodes imagined movement in

the absence of sensory information. Communications biology 3:1–7.

Jiang W, Lamarre Y, Chapman CE (1990) Modulation of cutaneous cortical evoked

potentials during isometric and isotonic contractions in the monkey. Brain Research

536:69–78.

Kaufman MT, Churchland MM, Ryu SI, Shenoy K V (2014) Cortical activity in the null space:

permitting preparation without movement. Nature Neuroscience 17:440–448.

Keele SW (1968) Movement control in skilled motor performance. Psychological Bulletin

70:387–403.

Keele SW, Summers JJ (1976) The Structure of Motor Programs. In: Motor Control, pp 109–

142. Elsevier.

Klapp ST (1995) Motor Response Programming During Simple and Choice Reaction Time:

The Role of Practice. Journal of Experimental Psychology: Human Perception and

Performance 21:1015–1027.

Klapp ST, Erwin CI (1976) Relation between programming time and duration of the

response being programmed. Journal of Experimental Psychology: Human Perception

and Performance 2:591–598.

Ledoit O, Wolf M (2004) Honey, I shrunk the sample covariance matrix. The Journal of

Portfolio Management 30:110–119.

Leoné FTM, Heed T, Toni I, Medendorp WP (2014) Understanding effector selectivity in

human posterior parietal cortex by combining information patterns and activation

measures. The Journal of neuroscience 34:7102–7112.

Logothetis NK, Pauls J, Augath M, Trinath T, Oeltermann A (2001) Neurophysiological

investigation of the basis of the fMRI signal. Nature 412:150–157.

Oosterhof NN, Wiestler T, Downing PE, Diedrichsen J (2011) A comparison of volume-

based and surface-based multi-voxel pattern analysis. NeuroImage 56:593–600.

Oosterhof NN, Wiggett AJ, Diedrichsen J, Tipper SP, Downing PE, Ramsey R, Hamilton

AFDC (2012) Surface-Based Information Mapping Reveals Crossmodal Vision –

Action Representations in Human Parietal and Occipitotemporal Cortex Surface-Based

Information Mapping Reveals Crossmodal Vision – Action Representations in Human

Parietal and Occipitotemporal. Journal of neurophysiology 104:1077–1089.

Rathelot J-A, Strick PL (2006) Muscle representation in the macaque motor cortex: An

anatomical perspective. Proceedings of the National Academy of Sciences 103:8257

LP – 8262.

Riehle A, Requin J (1989) Monkey primary motor and premotor cortex: single-cell activity

related to prior information about direction and extent of an intended movement.

Journal of neurophysiology 61:534–549.

Rosenbaum DA (1980) Human movement initiation: Specification of arm, direction, and

extent. Journal of Experimental Psychology: General 109:444–474.

Seki K, Fetz EE (2012) Gating of sensory input at spinal and cortical levels during

preparation and execution of voluntary movement. Journal of Neuroscience 32:890–

902.

Shenoy K V, Sahani M, Churchland MM (2013) Cortical control of arm movements: a

dynamical systems perspective. Annual Review of Neuroscience 36:337–359.

Starr A, Cohen LG (1985) 'Gating' of somatosensory evoked potentials begins before the

onset of voluntary movement in man. Brain Research 348:183–186.

Tanji J, Evarts E V. (1976) Anticipatory activity of motor cortex neurons in relation to

direction of an intended movement. Journal of Neurophysiology 39:1062–1068.

Turella L, Tucciarelli R, Oosterhof NN, Weisz N, Rumiati R, Lingnau A (2016) Beta band

modulations underlie action representations for movement planning. NeuroImage

136:197–207.

Umeda T, Isa T, Nishimura Y (2019) The somatosensory cortex receives information about

motor output. Science Advances 5.

Vyas S, Golub MD, Sussillo D, Shenoy K V (2020) Computation Through Neural Population

Dynamics. Annual Review of Neuroscience 43:249–275.

Walther A, Nili H, Ejaz N, Alink A, Kriegeskorte N, Diedrichsen J (2016) Reliability of

dissimilarity measures for multi-voxel pattern analysis. NeuroImage 137:188–200.

Widener GL, Cheney PD (1997) Effects on Muscle Activity From Microstimuli Applied to

Somatosensory and Motor Cortex During Voluntary Movement in the Monkey. Journal

of Neurophysiology 77:2446–2465.

Wong AL, Haith AM (2017) Motor planning flexibly optimizes performance under

uncertainty about task goals. Nature Communications 8:14624.

Yokoi A, Arbuckle SA, Diedrichsen J (2018) The role of human primary motor cortex in the

production of skilled finger sequences. Journal of Neuroscience 38:1430–1442.

Yousry T (1997) Localization of the motor hand area to a knob on the precentral gyrus. A

new landmark. Brain 120:141–157.

Figure S1 | Activation and distance analyses using planning of both go and no-go trials. A. Activation map (percent signal change) for the contrast planning>baseline. The selected area of interest is the same as shown in purple in the inset of Fig. 2A. **B.** Crossnobis distance searchlight map for movement planning. **C.** Same as A, but for the contrast execution>baseline. **D.** Same as B, but for movement execution. **E-F.** Cross-section analysis corresponding to the same area shown in white in the inset of Fig. 2A. **E.** Mean percent signal change (± SEM) during planning (orange) and execution (blue). **F.** mean crossnobis distance (± SEM). Horizontal bars indicate significance ($p < 0.05$) in a 2-sided one-sample t-test against zero. All other figure conventions are the same as in Fig. 2.