Genome architecture shapes evolutionary adaptation to DNA replication stress

Marco Fumasoni*, Andrew W. Murray*

Department of Molecular and Cellular Biology, Harvard University, Cambridge, United States

ABSTRACT

Evolutionary adaptation to perturbations in DNA replication follows reproducible trajectories that lead to changes in three important aspects of genome maintenance: DNA replication, the DNA damage checkpoint, and sister chromatid cohesion. We asked how these trajectories depend on a population's genome architecture by testing whether ploidy or the ability to perform homologous recombination influence the evolutionary fate of the budding yeast, *Saccharomyces cerevisiae*, as it adapts to constitutive DNA replication stress, a condition that characterizes many cancer cells. In all three genome architectures, adaptation happens within 1000 generations at rates that are linearly correlated with the initial fitness defect of the ancestors. Which genes are mutated depends on the frequency at which mutations occur and the selective advantage they confer. The recombination-deficient strain amplifies adaptive chromosomal regions less often, whereas the selective advantage of loss-of-function mutations, such as those that inactivate the DNA damage checkpoint, is reduced in diploids because of the presence of a second, wild-type copy of the gene. Despite these differences, selection targets the same three functional modules in all three architectures, suggesting that genome architecture controls which genes are mutated but not which modules are modified.

*For correspondence: marcofumasoni@fas.harvard.edu, awm@mcb.harvard.edu

INTRODUCTION

Organisms and cells adapt to selective pressures through the acquisition of beneficial mutations. The interactions between beneficial mutations that produce evolved phenotypes are complex and difficult to determine both in laboratory and natural evolution. These interactions are also likely to depend on two factors. The first, genetic background, refers to the collection of individual sequence variants in a strain or population [1–4]. The second is genome architecture, which we use to refer to functional features of the genome including its ploidy, rates of recombination, and the activity of mobile genetic elements. Previous work has revealed that different genetic backgrounds can select different spectra of mutations and different genome architectures lead do different rates of adaptation [5–14] but we lack a general, mechanistic explanation of these differences. Here, we take advantage of our earlier work on adaptation to DNA replication stress to investigate how different genome architectures control adaptive evolutionary trajectories.

We previously characterized the evolutionary adaptation of haploid cells of the budding yeast, *Saccharomyces cerevisiae*, experiencing constitutive DNA replication stress [15], a perturbation of DNA synthesis that produces lesions in DNA, increases genome instability, and decreases fitness [16–19]. DNA replication stress is often present in cancer cells as a consequence of oncogene activation [20–23] and can also be generated by the unrestrained proliferation of selfish genetic elements such as viruses or transposons [24–27]. In haploid *S. cerevisiae*, evolutionary adaptation to constitutive DNA replication stress is driven by mutations affecting three different functional modules that contribute to genome maintenance: gain-of-function mutations that stabilize the linkage between sister chromatids, mutations that improve the completion of DNA replication, most likely by stabilizing the progression of replication forks, and mutations that inactivate the DNA damage checkpoint, thus accelerating the cell division cycle. Full adaptation to the defects imposed by DNA replication stress requires a combination of mutations in all three modules, which are acquired in an order dictated by the epistatic interactions between the mutations [15].

We asked how genome architecture affected evolutionary adaptation to constitutive DNA replication stress by testing two additional architectures: i) diploid cells, which carry two copies of every chromosome and thus make fully recessive mutations phenotypically silent, ii) haploid, recombination-deficient cells which cannot produce gene amplifications or deletions that depend on homologous recombination between repeated sequences [28,29]. These architectures are relevant for cancer biology: cancer progression begins in diploid cells and defects in homologous recombination have been implicated in predisposition to tumorigenesis (i.e. BRCA2 mutations, [30,31]) and exploited for anticancer therapies [32–34].

We evolved cells with these two genome architectures and compared their responses with those of haploid, recombination-proficient cells. All three architectures experienced constitutive DNA replication stress caused by removing Ctf4 [15], a protein that physically links different activities at replication forks [35]. Adaptation to DNA replication stress occurs in all three genome

architectures at rates that are correlated with their initial fitness defects. Whole genome
sequencing of the evolved populations revealed diverse mutational spectra with only modest
convergence among the three different architectures. Despite the differences in which genes
were mutated, all three architectures accumulated mutations in the DNA replication machinery,
the DNA damage checkpoint, and sister chromatid cohesion. Individually engineering a subset of
these mutations into all three architectures demonstrated their ability to reduce the fitness cost
of DNA replication stress, even in architectures where they didn't appear. Combining mutations
in all three modules produced fitness increases that matched those of the evolved strains in all
three architectures. We conclude that although differences in the observed frequency of
mutations in different genes depend on genome architecture, the principles that govern
adaptation to constitutive DNA replications stress are largely architecture-independent. We
discuss the implication of our results for tumor evolution and for generating inter-species
molecular diversity in genome maintenance.

RESULTS

All populations adapt to constitutive DNA replication stress.

To induce constitutive DNA replication stress in *S. cerevisiae*, we deleted the *CTF4* gene, which
encodes a non-essential subunit of the DNA replication machinery (the replisome) [36]. Ctf4 is a
homo-trimer that functions as a structural hub within the replisome [35,37] by binding to the
replicative DNA helicase, primase (the enzyme that makes the RNA primers that initiate DNA
replication), and other accessory factors [35,38–40]. In the absence of Ctf4, cells experience DNA
replication stress [41–43] and have substantially reduced reproductive fitness [15]. We removed
CTF4 from wild-type (WT) haploid and diploid cells, as well as from haploids impaired in
homologous recombination due to the deletion of *RAD52* (Figure 1A), which encodes a conserved
enzyme required for pairing homologous DNA sequences during recombination [44]. Attempts
to generate homozygous *ctf4Δ/ctf4Δ rad52Δ/rad52Δ* mutants failed due to the lethality of the
resulting zygotes. We refer to the stains we analyzed as haploids (*ctf4Δ*, yellow), diploids
(*ctf4Δ/ctf4Δ*, blue) and recombination-deficient (*ctf4Δ rad52Δ*, red). Compared to the previously
reported *ctf4Δ* haploid strains, which manifested a fitness decrease of ~27% relative to isogenic
wild-type cells [15], constitutive DNA replication stress caused by the absence of Ctf4 was more
detrimental in diploid (~35% reduced fitness) and recombination-deficient strains (~55% reduced
fitness) (Figure 1B). We evolved eight parallel diploid and recombination-deficient populations
for 1000 generations in the same conditions and at the same time as the haploid populations we
previously analyze ([15] and see Materials and methods). In this regime, spontaneous mutations
that increase fitness and survive genetic drift will spread through the population [45,46]. In
addition, we evolved eight wild-type diploid populations like the previously evolved wild-type
haploids, to control for mutations that are selected by the culture conditions rather than by
ameliorating DNA replication stress [15].

After 1000 generations experiencing constitutive DNA replication stress, we asked how the
fitness of the evolved diploid and recombination-deficient populations compared to those of
evolved haploids. Regardless of their genome architecture, all populations lacking Ctf4 increased
in fitness compared to their respective ancestors: haploids from -27% to -9.6 ± 1.2% [15], diploids
from -35% to -14.4 ± 1.8%, and recombination-deficient haploids from -55% to -19 ± 1% (Figure
1B). The fitness of four clones isolated from each evolved population were similar to the fitness
of the population they were derived from (Figure 1-S1). Control evolved wild-type populations
showed only small fitness increases: 4 ± 1% for haploids and 6.7 ± 2.7% for diploids (Figure 1-S2).

Figure 1: Evolutionary adaptation to constitutive DNA replication stress in different genome architectures. (A)
Experimental scheme: Constitutive DNA replication stress was induced by removing Ctf4, a protein that coordinates
replisome activities. Haploid (orange), diploids (blue) and cells with a severely decreased ability to perform
homologous recombination (red, recombination-deficient), were subjected to 1000 generations of experimental
evolution. **(B)** The fitness of the 6 ancestral strains (2 for each genome architecture) and of 24 evolved populations
derived from them, relative to WT cells (*s* = 0). Fitness data of haploid strains (orange) is from [15]. The semi-
transparent gray dot represents a diploid strain that became haploid during the experiment and is thus excluded
from subsequent analysis. Error bars represent standard deviations. a and α refer to the strains' mating type (*MAT*
locus). a/α indicates diploid strains. The fitness values shown here are reported in Source data 1.

The rate of adaptation to DNA replication stress depends on the ancestor's fitness

Although all populations increased their fitness in response to constitutive DNA replication stress,
strains with different genome architectures increased in fitness to different extents (measured
as the absolute fitness gain of populations over 1000 generations). Recombination-deficient
populations showed the largest fitness gain (+34.6 ± 0.8%), followed by diploid (+20.3 ± 1.6%)
and haploid populations (+17.2 ± 1.0%, Figure 2A). We asked what could have generated different

adaptation rates in response to the same selective pressures. Less fit ancestral strains have been
proposed to increase in fitness faster than more fit ancestors, a phenomenon referred as
'declining adaptability' [47–50]. To test for this possibility, we plotted the final fitness of all
evolved populations against the fitness of their respective ancestors (Figure 2C). We found that
the absolute fitness gain over 1000 generations depended linearly on the ancestors' relative
fitness (R^2 = 0.95). These results agree with previous studies, which observed declining
adaptability across a wide range of organisms and selection conditions [9,10,51,52].

Figure 2: Fitness increase over 1000 generations. (A) The fitness increase over 1000 generations for strains
experiencing constitutive DNA replication stress. Δs is the difference between the fitness of the evolved populations
and the fitness of their ancestors, with both fitnesses measured relative to that of cells that contain Ctf4 (*CTF4*) and
have the same ploidy. **(B)**. The fitness increase over 1000 generations for strains that contained wild-type *CTF4*. **(C)**
shows the correlation between the fitness increase during evolution (Δs) and the fitness defect of the ancestors
relative to cells of the same architecture containing Ctf4 (the ancestor relative fitness). Fitness data of haploid strains
(orange and white) is from [15]. Error bars represent standard deviations. Grey line represents linear regression
between the datapoints. The data shown here are reported in or derived from Source data 1.

The majority of previous studies have suggested that diploids adapt more slowly than isogenic
haploids [13,53,54]. Our diploid *ctf4Δ/ctf4Δ* populations gained more fitness than *ctf4Δ* haploids
over the course of 1000 generations (Figure 2A). Wild-type diploids increased more in absolute
fitness than wild-type haploids (Figure 2B), suggesting that the discrepancy between our work
and previous studies is not due to the presence of replication stress. Diploid populations
recovered a slightly smaller fraction of their initial fitness defect compared to haploid and
recombination-deficient strains (Figure 2-S2A). Collectively, our results suggest that ploidy is a
poor predictor of the speed and extent of adaptation (Figure 2-S1). We conclude that the fitness
of the ancestral cells is the major determinant influencing how much populations adapt, adding
further support to the concept of 'declining adaptability'.

Limited parallelism in adaptively mutated genes across genome architectures.

We used whole-genome sequencing to understand the genetic basis of the evolutionary
adaptation to DNA replication stress. For each different genome architecture, we sequenced all
eight evolved populations, as well as four individual clones isolated from each of them. A total of
195, 276 and 905 genes or associated regulatory sequences were found mutated in evolved
haploid, diploid, and recombination-deficient populations respectively (Figure 3A,
Supplementary table 1). When normalized for DNA content, haploid (13 mutations/haploid
genome) and diploid clones (9 mutations/haploid genome) had similar but statistically different
mutation frequencies (Figure 3-S1) but recombination-deficient clones carried many more
mutations, both in total (50/haploid genome, Figure 3-S1A) and as synonymous mutations
(7/haploid genome, Figure 3-S1B). Our results agree with the reported increase in spontaneous
mutagenesis in *rad52Δ* cells [55–57].

We used statistical analysis to identify candidate adaptive mutations, by identifying genes that
were mutated more often than expected by chance in independent populations (see Materials
and methods for details). We performed this analysis both within individual genome
architectures and across all three architectures. Using a false discovery rate of 5% we found a
total 50 genes that met one or both criteria.

Only 8 of these genes were mutated in all three genome architectures (Figure 3A). This small
number could be due to different adaptive strategies adopted by the different architectures,
differences in the rate for the same mutation in different architectures, differences in the
selective benefit the mutation conferred, or a combination of these three effects. We found 28
genes with putative adaptive mutations that are specific to individual genome architectures, as
well as 14 genes with putative adaptive mutations in two architectures (Figure 3A, genes
highlighted in boxes and Supplementary table 2). Analyzing gene ontology (GO) terms associated
with the putative adaptive mutations revealed an enrichment of biological process terms
associated with genome maintenance (Supplementary table 3). The genes that were significantly
mutated in all genomic architectures included two that we had previously demonstrated to be
adaptive in haploid *ctf4Δ* cells: mutations in *RAD9* inactivate the DNA damage checkpoint and
those in *SLD5* alter the function of the replicative helicase [15]. All three architectures also had

mutations in *RAD61* and *PDS5*, which are implicated in chromosome segregation [58,59], a cellular module shown to play a role during the adaptation of haploid *ctf4Δ* populations [15].

Figure 3: Putative adaptive mutations in the evolved clones. (A) Venn diagram representing the mutations found in the evolved clones and populations. The area of the circles is proportional to the list of genes mutated at least once in the set of evolved clones for each genome architecture. The number within each sector represents number of genes which were mutated in one, two, or three genetic architectures. Tables include the genes whose mutations were significantly mutated in either global or architecture-specific analysis. Symbols next to the genes highlight that they were statistically significantly mutated in individual architectures (* refers to haploid, # to diploid and + to recombination-deficient architectures, e.g. *SLD5* +# means that the gene is found mutated in all three architectures, but is statistically significantly mutated only in diploid and recombination-deficient populations). The identities of all

the genes in the Venn diagram intersections are reported in Source data 3 **(B)** Copy number variations (CNVs)
affecting different chromosomes (roman numbers) which appeared in at least two independent populations. Arabic
numbers are the number of independent populations, in each genetic architecture (highlighted by a different color),
in which the CNVs were detected. Copy number change refers to the fragment's gain or loss during the evolution
experiment (i.e. +1 means that one copy was gained in haploid cells and that two copies were gained in diploid cells).
Frames around the plots have the same color as the sectors in the Venn diagram, and encircle CNVs that have been
found in one, two, or three genetic architectures. All the segmental amplifications detected are reported in
Supplementary table 4.

We also examined the evolved genomes for changes in gene copy numbers (Copy Number
Variations, CNVs) that might have played a role in adaptation. Many clones displayed an
increased copy number of defined chromosome segments (segmental amplification) compared
to the control populations (Supplementary table 4). Segmental amplifications were common in
diploid clones (Figure 3-S1C). We inferred that segmental amplifications were putatively adaptive
if the same chromosome region was found amplified in two clones isolated from independent
populations (Figure 3B). Among these segmental amplifications, four CNVs were found only in
evolved diploids, four were found in at least an evolved haploid and a diploid clone, and only one
was detected in populations of all three genome architectures (Figure 3B). This latter fragment is
located on chromosome IV and contains *SCC2*, which encodes a subunit of the cohesin loader
complex. This essential complex is responsible for loading the proteinaceous cohesin ring that
holds sister chromatids together until they segregate from each other during mitosis [60–62].
The gene encoding for the other subunit of the cohesin loader complex, *SCC4*, lies on the segment
of chromosome V that was amplified in both haploid and diploid clones. We previously showed
that extra copies of these genes alleviate the cohesion defects associated with the absence of
Ctf4 in haploid cells and increase their fitness [15].

Overall, genome sequencing reveals that the identity of the genes carrying selected mutations
varies depending on genome architecture. Nevertheless, we found evidence of common
evolutionary strategies across different populations: Mutations affecting sister chromatid
cohesion, DNA replication, and the DNA damage checkpoint, which were previously shown to
drive adaptation to DNA replication stress in haploid cells, are significantly mutated regardless of
the genome architecture of the ancestral cells.

Evolved diploids show adaptive loss of heterozygosity

We analyzed the frequency of mutated reads in evolved diploid clones to estimate whether
mutations were homozygous or heterozygous (Figure 4-S1). We detected 36 homozygous
mutations across the evolved diploid clones (Figure 4A, Figure 4-S1B, Supplementary table 5).
Homozygous mutations were evenly distributed across the evolved clones analyzed and thus
were not the product of a single unrepresentative event (Figure 4-S1C). Individual clones often
carried multiple homozygous mutations on the same chromosome arm (Supplementary table 5)
suggesting that the loss-of-heterozygosity (LOH) was likely the consequence of mitotic
recombination events affecting large chromosomal segments [63]. In addition to homozygous
mutations, three genes were found to carry two different mutations, most likely one in each of

the two copies of the gene (double hits, Figure 4A, Supplementary table 5). Ten of the homozygous or double hit mutations were abundant within their respective populations (mutation penetrance > 70%, Figure 4B and 4-S2). Gene ontology analysis of the 39 genes with homozygous mutations or double hit alleles revealed the enrichment of many terms associated with genome maintenance (Supplementary table 6). One cluster of genes, linked by genetic and physical interactions, affected components of the replisome and factors involved in the DNA damage checkpoint (Figure 4B). Because these genes had been changed by two events (either two mutations, or a mutation followed by LOH) and the altered forms had risen to high frequency in the evolved populations (Figure 4B), we argue that the inactivation or modification of many of these genes was selected for by constitutive DNA replication stress.

Figure 4: Homozygous mutations in diploids evolved in the presence of constitutive DNA replication stress. (A) Homozygous mutations detected in evolved *ctf4Δ/ctf4Δ* diploids clones distributed by chromosome (roman numbers). Dark blue diamonds represent loss of heterozygosity events (LOH). Light-blue triangles represent genes which contain two independent mutations (double hits) and the gray circles are the centromeres. The identities of the mutations shown here are reported in Supplementary table 5. **(B)** DNA replication and DNA damage checkpoint genes with mutations in both copies in evolved *ctf4Δ/ctf4Δ* diploid clones. Gray lines represent evidence of genetic and physical interactions from the literature (https://string-db.org). Node diameter is proportional to the number of populations in which the gene was mutated. Dark blue represents LOH events, light blue represents double hits . Percentages refer to the average estimated frequency within populations of the clones carrying the respective mutation in homozygosity (LOH) or carrying both mutations (double hits).

Genome architecture controls the distribution of adaptive mutants

In addition to altering which genes get adaptive mutations, we hypothesize that genome architecture could affect evolutionary trajectories in two ways: changing the frequency at which different modules are altered and changing the distribution of the types of mutations that are selected (for example by altering the ratio of adaptive point mutations to copy number variation [57,64–68]). Our analysis revealed that all three architectures acquired putative adaptive mutations in three genome maintenance modules: sister chromatid cohesion, DNA replication

and the DNA damage checkpoint. Out of the 23 populations analyzed, 18 of them contained at
least one clone with mutations affecting all three of these modules, while all of them contain a
clone with mutations in two of the three modules (Figure 5A). Despite the ubiquity of mutations
in genome maintenance, the frequency of mutations in genes known to affect individual cellular
modules varied between genome architectures: mutations affecting sister chromatid cohesion
were underrepresented in recombination-deficient strains (p=2.7e-04, χ^2 test) and there were
fewer mutations affecting the DNA damage checkpoint in diploid clones (p=0.03, χ^2 test, Figure
6A).

If amplifying the genes for the cohesin loader, *SCC2* and *SCC4*, depended on homologous
recombination between repeated sequences, the rate of amplification should be dramatically
decreased in the recombination-deficient strain, which lacks Rad52. Analysis of the segmental
amplifications containing *SCC2* or *SCC4* revealed that their boundaries correspond to three
classes of repeated sequences: transposons (Ty elements), transposon-associated long terminal
repeats (LTRs) and tRNA genes (Figure 5-S1 and Supplementary table 7). Recombination at these
repeated elements has been previously shown to produce segmental amplifications selected
under limited nutrient availability [69–71]. Amplification of *SCC2* thus occurs as a result of
homologous recombination between two different copies of the repeated sequences on
chromosome IV, one on either side of *SCC2* and amplification of *SCC4* is due to similar events on
chromosome V. This dependence on homologous recombination explains the absence of
segmental amplifications in all but one of the recombination-deficient clones (Figure 3B and
Supplementary table 4). Because changes affecting sister chromatid cohesion rely predominantly
on gene amplifications (Figure 5C), we detect fewer mutations affecting this module in
recombination-deficient strains (Figure 5A).

In haploid *ctf4Δ* cells, inactivating the DNA damage checkpoint increases fitness [15]. The primary
target for mutation is *RAD9*, both because of its large size (3927 bp) and the presence of a run of
11 As, which repeatedly gain an extra nucleotide causing a frameshift mutation. Because this is a
loss-of-function mutation [15], removing Rad9 function in diploids requires two events, thus
reducing the probability of adapting by destroying the DNA damage checkpoint. Despite this
difficulty, one diploid clone is homozygous for the frameshift mutation in *RAD9*, most likely as a
result of a mutation in one copy of the gene followed by mitotic recombination or gene
conversion that homozygosed this mutation and fully inactivated the checkpoint. The need to
follow a two-hit pathway to fully inactivate the DNA damage checkpoint explains why fewer RAD9
mutations were found in diploids.

We conclude that although mutations affecting sister chromatid cohesion, DNA replication and
DNA damage checkpoint are found in all the three genome architectures, the frequency at which
we detect them depends on the genome architecture of an evolving population: gene
amplification occurs much less frequently in recombination-deficient populations, making one
pathway for improving sister chromatid cohesion less accessible, and recessive null mutations

are rarer in diploid populations, blocking a major route by which haploids inactivate the DNA
damage checkpoint.

Figure 5: Mutations affecting sister chromatid cohesion, DNA replication and the DNA damage checkpoint. (A)
Heatmaps representing all the mutations (SNPs, InDels and segmental amplifications) causally implicated in these

302 phenotypes in the 92 sequenced evolved clones **(B)** Frequency of SNPs and small InDels. **(C)** Frequency of genes
303 associated with the three processes that are present on segmental amplifications. The number of hits and the
304 identities of the genes which were mutated are reported in Source data 5.

Mutations affecting sister chromatid cohesion, DNA replication and DNA damage checkpoint recapitulate evolution in all genome architectures.

We asked whether genome architecture affected the benefit that mutations conferred as well as
the frequency at which they were observed. To test this hypothesis, we chose a group of recurring
mutations to validate their causality and measure their fitness benefit in different genome
architectures. We mimicked the segmental amplification of chromosome IV (Figure 3B), affecting
the cohesin loader *SCC2*, by integrating an extra copy of the gene at the *URA3* locus (referred to
as *2xSCC2*). Ixr1 is a transcription factor that indirectly affects DNA replication by positively
regulating the production of deoxyribonucleotide triphosphates (dNTPs, [72]). The abundance of
mutations causing premature stop codons in *IXR1* suggested that evolution selected for the
inactivation of this gene (Supplementary table 8). We mimicked the loss of a functional Ixr1
protein by gene deletion (*ixr1Δ*). Another recurrently mutated gene affecting DNA replication
was *SLD5*, which encodes the subunit of the main replicative helicase (CMG, [73]) that is bound
in vivo by Ctf4 [74]. *SLD5* is an essential gene and our earlier analysis suggested that mutations
like *sld5-E130K* reduced or altered the helicase's function [15]. Finally, the majority of mutations
affecting the DNA damage checkpoint factor *RAD9* resulted in early stop codons (Supplementary
table 9), leading us to test the effect of the *rad9Δ* gene deletion. These mutations were all
previously demonstrated to be adaptive in haploid cells [15] and were used because they
occurred, albeit at different frequencies, in all three (point mutations in *SLD5* and *RAD9* and
amplification of *SCC2*) or at least two genome architectures (point mutations in *IXR1* in both
haploid populations).

We found that all four mutations were adaptive in all three genomic architectures. Remarkably,
mutations affecting individual genes were adaptive even in architectures where they were never
(*IXR1* in diploids) or rarely detected (*SLD5* in haploids, *RAD9* in diploids and *SCC2* in
recombination-deficient strains). Three of the mutations we tested (*SCC2* amplification, *sld5-
E130K*, and *rad9Δ*) conferred similar benefits in all three architectures, but the fourth, *ixr1Δ*, gave
a much larger benefit in the recombination-deficient strain. One possibility is that the absence of
Ixr1 reduces the frequency of double-stranded breaks associated with DNA replication in the
absence of Ctf4. These breaks are lethal in the absence of Rad52 but can be repaired in the other
two architectures.

For the diploid *ctf4Δ/ctf4Δ* strain, we tested each mutation as a heterozygote (e.g. *IXR1/ixr1Δ*)
and a homozygote (e.g. *ixr1Δ/ixr1Δ*). The homozygous deletion of *IXR1* gave a substantial fitness
benefit (Figure 6A, dark blue), but strains heterozygous for the deletion were slightly less fit than
their ancestors (Figure 6A, light blue). For the other three mutations (an extra copy of *SCC2*, *sld5-
E130K*, and *rad9*), the heterozygotes conferred a fitness benefit that was significant, albeit

weaker than that of the homozygote. These results suggest that homozygous mutations can be
selected by a two-step process in which both the first step, mutation in one copy of the gene,
and the second, mutation in the other copy or loss of heterozygosity, confer a selective
advantage. We conclude that, mutations affecting sister chromatid cohesion, DNA replication
and DNA damage checkpoint are beneficial in the presence of constitutive DNA replication stress
independently of the genome architecture of the strains.

Figure 6: Reconstructed adaptive mutations can produce evolved fitness increases. (A) A subset of mutations
affecting sister chromatid cohesion, DNA replication and DNA damage checkpoint were re-constructed in the
ancestor cells for all three genomic architectures. The fitness of the re-constructed strains, relative to their
respective *ctf4Δ* ancestor cells ($s = 0$) is depicted. For diploid strains, we replaced one (het) or two (hom) copies of
the wild-type gene with the mutant allele. The fitness values shown here are reported in Source data 6a. **(B)**
Reconstructed strains carrying combinations of adaptive mutations affecting sister chromatid cohesion, DNA
replication and DNA damage checkpoint recapitulate the fitness of their respective evolved populations after 1000
generations. The most frequently mutated gene affecting DNA replication was chosen for the reconstruction in
individual architectures (e.g. *IXR1* was mutated more often that *SLD5* in haploids whereas the reverse was true in

dioids). Fitness data of haploid strains (orange) is from [15]. The fitness values shown here are reported in Source
data 6b.

Are mutations in sister chromatid cohesion, DNA replication and the DNA damage checkpoint
enough to recapitulate the evolved fitness increase in all three genomic architectures? We
previously showed that a haploid *ctf4Δ* strain carrying a mutation in all three of these modules
recapitulated the fitness of the evolved strains (Figure 6B and [15]). We therefore asked if
mutations in these three modules were sufficient to recapitulate 1000 generations of evolution
in diploid and recombination-deficient strains. We constructed strains that carried mutations
affecting sister chromatid cohesion, DNA replication and DNA damage checkpoint (Figure 6B),
using mutations that had been found in the same genome architecture (Figure 3,4 and
Supplementary table 1,4) and shown to be adaptive in isolation (Figure 6A).

For all three genome architectures, the reconstructed strains had a fitness that was similar to the
evolved populations with the same architecture (Figure 6B). This result demonstrates that for all
three genome architectures, acquiring mutations in all three modules (sister chromatid cohesion,
DNA replication and DNA damage checkpoint) can recapitulate the fitness increase seen in
response to evolution in the presence of DNA replication stress.

DISCUSSION

We previously showed that haploid cells adapt to constitutive DNA replication stress by
accumulating mutations affecting DNA replication, sister chromatid cohesion, and the DNA
damage checkpoint [15]. We repeated this experiment in two other genomic architectures, a
diploid and a recombination-deficient haploid strain, to ask which features of this evolutionary
trajectory were general and which were specific to a particular genomic architecture. We find
that the genes that acquire adaptive mutations, the frequency at which they are mutated, and
the frequency at which these mutations are selected all differ between architectures but that
mutations that confer strong benefits always lie in the same three modules. Engineering one
mutation in each module into an ancestral strain lacking Ctf4 is enough to produce the evolved
fitness increase in all three genomic architectures, demonstrating the existence of a common
pathway for yeast cells to adapt to constitutive DNA replication stress.

The speed of adaptation depends on initial fitness not genome architecture

What determines the rate of evolutionary adaptation? One possibility is genome architecture:
beneficial, recessive mutations are harder to select in diploids [53,75] and gene amplification is
less frequent in recombination-defective cells [76,77], suggesting that recombination proficient
haploids should evolve faster than the other two genome architectures. Another explanation is
that the initial fitness of the population is the principal factor in setting the rate of adaptation,
with less fit ancestral populations increasing in fitness faster than more fit ones, a phenomenon
referred to as declining adaptability [47]. We observed that adaptation rates scale linearly with

the fitness difference between a mutant ancestor and a non-mutant wild-type, showing that the
recovery from DNA replication stress follows the declining adaptability documented in other
experimental conditions.

Previous work has produced a variety of results with the majority reporting that increasing ploidy
decreases the rate of evolution [13,53,54] and a minority reaching the opposite conclusion
[12,14]. We suggest that this variability arises because genome architecture affects two aspects
of the mutations that produce adaptation: the frequency at which they occur and the selective
advantage they confer. Thus, we find that defects in recombination reduce the rate of gene
amplification and diploidy makes the accumulation of fully recessive mutations require two
successive genetic events rather than a single mutation. Despite these barriers, our diploids and
recombination-defective haploids evolved faster than recombination-proficient haploids. We can
only speculate on why this is so but note that the recombination-proficient strains have a higher
mutation rate and diploids may be more resistant to the unbalanced gene dosage caused by
segmental amplifications.

We engineered the same mutations into all three genome architectures to ask if the benefit of
individual mutations depended on either genome architecture or the fitness of the recipient
strain. For three of the four mutations we tested (*sld5-E130K*, *rad9Δ*, and 2x*SCC2*) their effect
was independent of both these variables. The fourth (*ixr1Δ*) showed diminishing return epistasis,
defined as an adaptive mutation producing a smaller benefit when present in fitter genetic
backgrounds [49]. This effect has been invoked to explain declining adaptability and is thought
to arise from the complex pattern of epistasis between the adaptive mutations and the set of
alleles present in the genetic background [52,78–80]. Because most of the mutations did not
show diminishing returns epistasis, we speculate that the mutations that are selected during
evolutionary repair of genetic damage are likely to have fewer and less complex interactions than
those that accumulate when wild type strains are selected to grow faster in variants of standard
laboratory environments [8,52].

General principles of adaptation to constitutive DNA replication stress

Evolutionary change can be examined at multiple levels from the details of adaptive mutations
in individual genes, the identity of the functional modules in which the mutations lie, cellular
phenotypes, such as the integrity of the DNA damage checkpoint, to overall reproductive fitness.
For each of these levels, we examined the effect of genome architecture on the adaptation to
constitutive DNA replication stress. At both ends of this scale we see substantial variation:
genome architecture controls the frequency at which mutations occur, the fitness benefit they
confer, and the extent of overall adaptation. In the middle, we see more conservation, with all
three architectures accumulating mutations in the same three functional modules, suggesting
that they all follow similar phenotypic paths by increasing the stability of the linkage between
sister chromatids, stabilizing replication forks, and inactivating the DNA damage checkpoint. To
test this hypothesis, we reconstructed mutations from all three modules into the ancestral, Ctf4-
deficient strains of all three genome architectures. This manipulation reproduced fitness

observed after 1000 generations of evolution leading us to conclude that the principles that
govern evolutionary adaptation to constitutive DNA replication stress are conserved across the
genome architectures we examined.

Molecular mechanisms affecting adaptation in diploid cells.

The phenotype of heterozygous mutations in diploid cells depends on their dominance. Although
loss-of-function mutations are often assumed to be fully recessive, a substantial minority have
detectable phenotypes as heterozygotes (referred to as semi-dominance or haploinsufficiency),
typically much weaker than that of the homozygous loss-of-function mutations [81]. The
phenotype of the heterozygote determines whether mutations will be selected for in diploid
populations [82,83]. As an example, loss-of-function mutations in *IXR1* were repeatedly found in
haploid and recombination-deficient populations but were not present in evolved diploid strains.
This observation is consistent with the slight fitness cost of the heterozygous *IXR1* deletion,
despite the strong benefit provided by the homozygous deletion: the cost of the heterozygote
acts as a barrier that prevents access to the beneficial homozygous mutations. Like *IXR1*, deletion
of both copies of *RAD9* has a strong fitness benefit in diploids, but in this case there is a small
fitness benefit in the *rad9Δ/RAD9* heterozygotes providing a path to inactivating the DNA damage
checkpoint in which both steps increase fitness. This two-step path in diploids is less favorable
than the one-step path that inactivates the checkpoint in haploids and likely accounts for the
lower frequency of mutations affecting *RAD9* and other known checkpoint genes in evolved
diploid clones (Figure 5 and Source data 5).

Two-step paths affected the replication machinery or DNA damage checkpoint in 9 of the 28
diploid clones; 2 involved independent mutations in the same gene and 8 produced homozygous
mutations due to gene conversion or mitotic recombination (Figure 4, one clone contained two
homozygous mutations). The high frequency of these events suggests that these mutations
conferred some advantage even as heterozygotes. For one mutation, *sld5-E130K*, we verified the
fitness benefit of the *sld5-E130K/SLD5* heterozygote, which suggests that the mutant subunit is
present in a fraction of the genome-wide replicative helicases, where it helps to stabilize
replication forks in the absence of Ctf4. Mitotic recombination can then homozygose the
mutation, eliminating the wild-type Sld5 protein and stabilizing all replisomes. The genome
architecture of diploid cells thus explains both the reduced frequency of mutations in *RAD9* and
dictates which mutations affecting DNA replication are ultimately selected (e.g. *IXR1* versus
SLD5). We suggest that semi-dominant mutations are an important intermediate in the selection
of loss-of-heterozygosity during the evolution of diploid, somatic cells.

Molecular mechanisms affecting adaptation in recombination-deficient cells.

Recombination-deficient clones have much fewer segmental amplifications affecting the genes
for the cohesin loader, *SCC2* and *SCC4* (Figure 5C). The lack of the recombinase protein Rad52 in
recombination-deficient strains dramatically reduces their ability to achieve segmental
amplifications. Only one event, likely mediated by Rad52-independent recombination [84–87],

was found in the recombination-deficient evolved clones (Figure 3B and Supplementary table 4).
The other mutations in these strains that are likely to affect sister chromatid cohesion were point
mutations affecting other proteins involved in pairing and segregation of chromosomes such as
Pds5 and Rad61 (Figure 5B). The higher number of point mutations in this and other modules is
likely the consequence of increased spontaneous mutagenesis imposed by the lack of homology-
mediated repair of spontaneously occurring DNA lesions (Figure 3-S1,2, [88]). The genome
architecture of recombination-deficient strains thus explains both the reduced frequency of gene
amplification affecting sister chromatid cohesion and the increased number of point mutations
detected in evolved clones (Figure 6B).

Impact of genome architecture on evolutionary adaptation to DNA replication stress

Our analysis argues that although genome architecture doesn't affect the ability of cells to adapt
to DNA replication stress or the modules that alter to produce adaptation, it does affect the
mutational pathways by which different architectures adapt.

If the major adaptive mechanisms are shared, how do we explain the limited number of genes
that are mutated in all conditions and the genes that are uniquely and significantly mutated in
specific genome architectures (Figure 3A)? There are several hypotheses: some of these genes
may be false positives, mutations in these genes confer benefits in more than one architecture
but were only detected in a single architecture, and mutations in these genes are substantially
more, or exclusively, beneficial in one architecture. The fact that our reconstructed strains
recapitulate the fitness of the evolved populations with a minimal set of changes in genome
maintenance suggests that architecture-specific mutations play a marginal role in adaptation. Of
the 22 genes that were mutated in at least two architectures, 10 have known roles in DNA
replication, repair, or sister chromatid cohesion, whereas only 5 of the 28 that were uniquely
mutated in one architecture have such a role (p = 0.034, χ^2 test). The fitness of the reconstructed
strains (Figure 6A,B) also implies that the most important mutations for the adaptation to DNA
replication stress are those that were found in multiple architectures (Figure 3A). Mutations that
were never (*ixr1*), or rarely (*2xSCC2* and *rad9*) seen in a given genome architecture, nevertheless
increased fitness in all three architectures. This finding suggests that caution should be used in
inferring adaptive strategies from lists of genes that have been mutated more often than
expected by chance, especially when comparing populations that were evolved under different
conditions or have different ancestral genotypes. We argue that experimental dissection of
adaptive trajectories is needed to distinguish the mutations that make major contributions to
adaptation from minor players and mutational noise.

The distribution of mutations with known phenotypic effects suggests that genome architecture
controls how much adaptation in different functional modules contributes to the overall
response to selective pressure. Mutations that break the DNA damage checkpoint by inactivating
RAD9 are less frequent in diploid populations than they are in haploids, and amplifications of
cohesin loader genes are less frequent in recombination-deficient populations than they are in

recombination-proficient ones. Although we suspect that some of the diploid populations have
found alternate genetic routes to partially or fully inactivate the DNA damage checkpoint and
that some of the recombination-defective populations have selected other mutations that
improve sister chromatid cohesion, these changes are likely to be less effective than those that
were frequent in other genome architectures.

Implications for cancer and natural evolution

We previously speculated that evolutionary processes similar to the adaptation to constitutive
DNA replication stress could assist the evolution of cancer cells and organisms in nature [15].
Tumor cells often experience DNA replication stress as a consequence of oncogene activation,
promoting genetic instability and increasing the rate of cell death [17,20]. Evolutionary
adaptation to DNA replication stress could reduce its negative effects on cellular fitness and thus
assist tumor evolution. In natural populations, interference with DNA replication, through the
fixation of deleterious mutations, perturbation by selfish genetic elements or toxins secreted by
antagonistic organisms could generate similar evolutionary adaptation. The consequences of
these evolutionary processes could rewire genome maintenance modules to alter cancer cells'
physiology and generate molecular diversity in nature [89]. For these scenarios to be plausible,
the evolutionary adaptation to DNA replication stress needs to happen over short timescales. In
cancer, it would need to happen over the limited somatic evolution that accompanies cancer
progression. In nature, it would need to occur before the perturbed organisms are outcompeted
by unaffected ones or block the source of DNA replication stress. It could be argued that some
genome architectures, such as diploidy or reduced recombination, could reduce the likelihood of
evolving new strategies for genome replication and maintenance. In our hands, however, these
genome architectures do not preclude rapid, evolutionary adaptation to constitutive DNA
replication stress: loss of heterozygosity or increased mutation rates, which have both been
documented in cancer [90,91] and natural evolution [92,93], allow cells to circumvent these
obstacles and adapt quickly to a severe perturbation of the replication machinery. One rapidly
evolving clade of yeasts has lost *RAD9* [94], the primary target of checkpoint-inactivating
mutations in our experiments. Similarly, many cancers inactivate the DNA damage response
during tumor evolution [95,96]. We suggest that the careful examination of the genomic
sequences of tumors and fast-evolving species might reveal signs of adaptation that mirror the
molecular evolution we have described.

MATERIALS AND METHODS

Strains

All strains were derivatives of a modified version (Rad5$^+$) of S. cerevisiae strain W303 (*leu2-3,112*
trp1-1 can1-100 ura3-1 ade2-1 his3-11,15, RAD5). Supplementary table 10 lists each strain's
genotype. The ancestors of wild-type, *ctf4Δ* and *ctf4Δ rad52Δ* strains were obtained by
sporulating a *CTF4/ctf4Δ RAD52/rad52Δ* heterozygous diploid. This was done to minimize the
selection acting on the ancestor strains before the beginning of the experiment. *CTF4/CTF4* and

ctf4∆/ctf4∆ diploid strains were obtaining by mating freshly generated *MAT***a** and *MAT*α haploid
strains.

Media and growth conditions

550 Standard rich medium, YPD (1% Yeast-Extract, 2% Peptone, 2% D-Glucose) was used for the
551 evolution experiment and all fitness measurements.

Experimental evolution

All populations used in the evolution experiment (EVO1-40) were initially inoculated in glass
tubes containing 10 ml of YPD with yeast colonies of the relative genotype. Glass tubes were
placed in roller drums at 30˚C and grown for 24 hr. Daily passages were done by diluting 10 µl of
the previous culture into 10 ml of fresh YPD (1:1000 dilution, allowing for approximately 10
generations/cycle). All populations were passaged for a total of 100 cycles (≈1000 generations).
After 1000 generations four evolved clones were isolated from the populations evolved in the
absence of Ctf4 (EVO1-24) by streaking cells on a YPD plate (total of 96 clones isolated). All final
populations and clones isolated from them were mixed with 800 µl of 30% v/v glycerol and stored
at -70˚C for future analysis. The evolution of haploid, diploid, and recombination defective
strains, all lacking *CTF4*, and wild-type haploid and diploid strains were all carried out in parallel.

Whole genome sequencing

Genomic DNA library preparation was performed with an Illumina (RRID:SCR_010233, San Diego,
CA, US) Nextera DNA Library Prep Kit. Libraries were then pooled and sequenced either with an
Illumina HiSeq 2500 (125bp paired end reads) or an Illumina NovaSeq (150 bp paired end reads).
The SAMtools software package (RRID:SCR_002105, samtools.source- forge.net) was then used
to sort and index the mapped reads into a BAM file. GATK (RRID:SCR_ 001876,
www.broadinstitute.org/gatk; McKenna et al., 2010) was used to realign local indels, and VarScan
(RRID:SCR_006849, varscan.sourceforge.net) was used to call variants. Mutations were found
using a custom pipeline written in Python (RRID:SCR_008394, www.python.org). The pipeline
(github.com/koschwanez/mutantanalysis) compares variants between the reference strain, the
ancestor strain, and the evolved strains. A variant that occurs between the ancestor and an
evolved strain is labeled as a mutation if it either (1) causes a non-synonymous substitution in a
coding sequence or (2) occurs in a regulatory region, defined as the 500 bp upstream and
downstream of the coding sequence.

Identification of putative adaptive mutations

To identify genes and cellular modules targeted by selection we used the following methods:

Convergent evolution on genes: This method relies on the assumption that those genes that have
been mutated significantly more than expected by chance alone, represent cases of convergent
evolution among independent lines. The mutations affecting those genes are therefore

considered putatively adaptive. The same procedure was used independently on the mutations
found in wild-type and *ctf4Δ* evolved lines:

We first calculated per-base mutation rates as the total number of mutations in coding regions
occurring in a given background, divided by the size of the coding yeast genome in bp (including
1000bp per ORF to account for regulatory regions)

587
$$\lambda = \frac{SNPs + indels}{bp\text{coding}}$$

588   If the mutations were distributed randomly in the genome at a rate λ, the probability of finding
589   n mutations in a given gene of length N is given by the Poisson distribution:

590
$$\text{P} (n\ mutations | gene\ of\ length\ N) = \frac{(\lambda N)^n e^{-\lambda N}}{n!}$$

591   For each gene of length N, we then calculated the probability of finding ≥ n mutations if these
592   were occurring randomly.

593
$$P(\neq \geq n\ mutations | gene\ of\ length\ N) = \sum_{k=n}^{\infty} \frac{(\lambda N)^n e^{-\lambda N}}{k!} = 1 - \frac{\Gamma (n + 1, \lambda N)}{n!}$$

(Where Γ is the upper incomplete gamma function) which gives us the p-value for the comparison
of the observed mutations with the null, Poisson model. In order to decrease the number of false
positives, we then performed multiple-comparison corrections. The more stringent Bonferroni
correction (α=0.05) was applied on the wild-type evolved mutations dataset, while Benjamini-
Hochberg correction (α =0.05) was used for the *ctf4Δ* mutation dataset. Genes that were found
significantly selected in the evolved wild-type populations (after Bonferroni correction) were
removed from the list of evolved *ctf4Δ* strains (genes mutated in wild-type evolved populations
were removed from the *ctf4Δ* and *ctf4Δ rad52Δ* datasets and genes mutated in wild-type diploid
populations were removed from the *ctf4Δ/ctf4Δ* dataset). We argue that since these genes were
targets of selection even in wild-type cells, they are likely involved in processes that are unrelated
to DNA replication and are instead associated with adaptation to sustained growth by serial
dilutions. Supplementary table 2 lists the mutations detected in evolved clones and populations,
after filtering out those that occurred in genes that were significantly mutated in the respective
wild-type populations. Genes significantly selected are shown in dark grey (after Benjamini-
Hochberg correction with α=0.05).

Convergent evolution on cellular modules: Lists of genes that had been mutated more frequently
than expected by chance (Supplementary table 2) or found homozygously mutated in diploid
strains (Supplementary table 5) were further analyzed for gene ontology (GO) terms enrichment.
Lists of mutations were input as 'multiple proteins' in the STRING database, which reports on the
network of interactions between the input genes (https://string-db.org). The 'biological process'
terms enriched in the significantly or homozygously mutated genes are listed in Supplementary
tables 3 and 6 respectively.

Fitness assays

To measure relative fitness, we competed the ancestors and evolved strains against reference strains. Wild-type refence strains were used for the competition assays in all the figures, except for those in Figure 6A where *ctf4Δ* reference strains were used instead. A *pFA6a-prACT1-yCerulean-HphMX4* plasmid was digested with AgeI and integrated at one of the *ACT1* loci of the original heterozygous diploid (*CTF4/ctf4Δ RAD52/rad52Δ*) strain. This allows for the expression of fluorescent protein yCerulean under the strong actin promoter. The heterozygous diploid was then sporulated and dissected to obtain fluorescent wild-type, *ctf4Δ* or *ctf4Δ rad52Δ* reference haploid strains. *ctf4Δ/ctf4Δ* diploid reference strains were obtained by mating two *ctf4Δ* haploid strains, one of which expressed yCerulean under the *ACT1* promoter. For measuring a strain's relative fitness, 10 ml of YPD were inoculated in individual glass tubes with either the frozen reference or test strains. After 24 hrs. the strains were mixed in fresh 10 ml YPD tubes at a ratio dependent on the expected fitness of the test strain compared to the reference (i.e. 1:1 if believed to be nearly equally fit) and allowed to proliferate at 30°C for 24 hrs. 10 μl samples were taken from this mixed culture (day 0) and the ratio of the two starting strains was immediately measured. Tubes were then cultured following in the same conditions as the evolution experiment by diluting them 1:1000 into fresh medium every 24hrs for 2-4 days, monitoring the strain ratio at every passage. Samples were analyzed by flow cytometry (Fortessa, BD Bioscience, RRID:SCR_013311, Franklin Lakes, NJ, US). Strain ratios and the number of generations occurred between samples were calculated using a custom Python script utilizing the FlowCytometryTools package (https://pypi.org/project/FlowCytometryTools/). Ratios, *r*, were calculated based on the number of fluorescent and non-fluorescent events detected by the flow cytometer:

$$r = \frac{NonFluorescent_{events}}{Fluorescent_{events}}$$

Generations between time points, *g*, were calculated based on total events measured at time 0 hrs and time 24 hrs:

$$g = \frac{\log_{10}(Events_{t24}/events_{t0})}{\log_{10} 2}$$

Linear regression was performed between the $(g, \log_e r)$ points for each sample. Relative fitness was calculated as the slope of the resulting line. The mean relative fitness, *s*, was calculated from measurements obtained from at least three independent biological replicates. Error bars represent standard deviations. The P-values reported in figures are the result of t-tests assuming unequal variances (Welch's test).

Copy number variations (CNVs) detection by sequencing

Whole genome sequencing and read mapping was done as previously described. The read-depths for every unique 100 bp region in the genome were then obtained by using the VarScan copynumber tool. A custom pipeline written in python was used to visualize the genome-wide CNVs. First, the read-depths of individual 100 bp windows were normalized to the genome-wide median read-depth to control for differences in sequencing depths between samples. The coverage of the ancestor strains was then subtracted from the one of the evolved lines to reduce

the noise in read depth visualization due to the repeated sequences across the genome. The resulting CNVs were smoothed across five 100 bp windows for a simpler visualization. Final CNVs were then plotted relative to their genomic coordinate at the center of the smoothed window. Since the wild-type CNVs were subtracted from the evolved CNVs, the y axis refers to the copy number change occurred during evolution (i.e. +1 means that one an extra copy of a chromosome fragment has been gained in haploid cells, and that two extra copies have been gained in diploids). Segmental amplifications were identified as continuous chromosomal regions affected by the same change in copy number (Figure 3B and Supplementary table 4). Amplified genes (Figure 5A,C) were obtained from the lists of ORFs included within the boundaries of the segmental amplifications.

The correlation of chromosomal features with segmental amplification boundaries

We first identified the approximate boundaries (± 500bp) of all segmental amplification by manually inspecting the BAM files with the Integrative Genomic Viewer (https://software.broadinstitute.org/software/igv/). The coordinates of the sequence features that define the boundary between lower and higher read-depths are reported in Supplementary table 4. We then examined the sequences within these approximate boundaries to determine whether various chromosomal features were over- or underrepresented. We considered features that previous studies have found to be associated with hotspots for lesions and sources of genetic instability. We first counted how many times a given feature fell in the boundary zones of segmental amplifications. Then we calculated the expected number of features in these zones based on the total number of features in the genome and the percentage of the genome represented by segmental amplification boundaries zones. We compared these numbers by χ2 analysis and reported the associated p-values (Supplementary table 7). The number of tRNA genes, transposable elements, LTRs, ARS elements, snRNA and snoRNA genes and centromeres in the genome were determined using YeastMine (https://yeastmine.yeastgenome.org/). Coordinates of fragile sites were obtained from [97]. G4 sequences were obtained from [98]. Highly-(top 5%) and weakly-(least 5%) transcribed genes were identified from the data in [99]. Rrm3 binding sites and regions with high levels of γH2AX were derived from [100] and [101], respectively. Sites of DNA replication termination were derived from [15]. The tandemly repeated sequences, with a minimal repeat tract of twenty-four bases, were obtained from the tandem-repeat-database (TRDB; https://tandem.bu.edu/cgi-bin/trdb/trdb.exe).

<u>DATA AND CODE AVAILABILITY</u>

<u>ACKNOWLEDGEMENTS</u>

We thank Michael Laub, María Angélica Bravo Núñez, Sriram Srikant, Thomas LaBar and Andrea Giometto for critical reading of the manuscript; Andrea Giometto for assistance in data analysis; Claire Hartman from the Harvard Bauer Core Facility for technical assistance. We thank the

members of the Murray and Nelson labs for helpful discussions. MF acknowledges fellowship support from the Human Frontiers Science Program, EMBO and AIRC.

AUTHOR CONTRIBUTIONS

MF designed and performed the research, analyzed and interpreted the data, and wrote the paper. AWM designed the research, interpreted the data, and wrote the paper.

BIBLIOGRAPHY

1. Chandler CH, Chari S, Dworkin I. Does your gene need a background check? How genetic background impacts the analysis of mutations, genes, and evolution. Trends in Genetics. Elsevier Current Trends; 2013. pp. 358–366. doi:10.1016/j.tig.2013.01.009

2. Dowell RD, Ryan O, Jansen A, Cheung D, Agarwala S, Danford T, et al. Genotype to phenotype: A Complex problem. Science. 2010. p. 469. doi:10.1126/science.1189015

3. Weinreich DM, Lan Y, Jaffe J, Heckendorn RB. The Influence of Higher-Order Epistasis on Biological Fitness Landscape Topography. J Stat Phys. 2018;172: 208–225. doi:10.1007/s10955-018-1975-3

4. Fournier T, Schacherer J. Genetic backgrounds and hidden trait complexity in natural populations. Current Opinion in Genetics and Development. 2017. pp. 48–53. doi:10.1016/j.gde.2017.08.009

5. Filteau M, Hamel V, Pouliot M, Gagnon-Arsenault I, Dubé AK, Landry CR. Evolutionary rescue by compensatory mutations is constrained by genomic and environmental backgrounds. Mol Syst Biol. 2015;11: 832. doi:10.15252/msb.20156444

6. Kempher ML, Tao X, Song R, Wu B, Stahl DA, Wall JD, et al. Effects of Genetic and Physiological Divergence on the Evolution of a Sulfate-Reducing Bacterium under Conditions of Elevated Temperature. MBio. 2020;11. doi:10.1128/mBio.00569-20

7. Blount ZD, Borland CZ, Lenski RE. Historical contingency and the evolution of a key innovation in an experimental population of Escherichia coli. Proc Natl Acad Sci USA. 2008;105: 7899–7906. doi:10.1073/pnas.0803151105

8. Jerison ER, Kryazhimskiy S, Mitchell JK, Bloom JS, Kruglyak L, Desai MM. Genetic variation in adaptability and pleiotropy in budding yeast. Elife. 2017;6. doi:10.7554/eLife.27167

9. Rojas Echenique JI, Kryazhimskiy S, Nguyen Ba AN, Desai MM. Modular epistasis and the compensatory evolution of gene deletion mutants. PLOS Genet. 2019;15: e1007958. doi:10.1371/journal.pgen.1007958

10. Szamecz B, Boross G, Kalapis D, Kovács K, Fekete G, Farkas Z, et al. The Genomic Landscape of Compensatory Evolution. PLoS Biol. 2014;12: e1001935. doi:10.1371/journal.pbio.1001935

11. Fumasoni M. Tell Me Where You've Been and I'll Tell You How You'll Evolve. MBio. 2020;11: 2043–2063. doi:10.1128/mBio.02043-20

12. Selmecki AM, Maruvka YE, Richmond PA, Guillet M, Shoresh N, Sorenson AL, et al. Polyploidy can drive rapid adaptation in yeast. Nature. 2015;519: 349–352. doi:10.1038/nature14187

13. Gerstein AC, Cleathero LA, Mandegar MA, Otto SP. Haploids adapt faster than diploids across a range of environments. J Evol Biol. 2011;24: 531–540. doi:10.1111/j.1420-9101.2010.02188.x

14.   Paquin C, Adams J. Frequency of fixation of adaptive mutations is higher in evolving diploid than
haploid yeast populations. Nature. 1983;302: 495–500. doi:10.1038/302495a0

15.   Fumasoni M, Murray AW. The evolutionary plasticity of chromosome metabolism allows
adaptation to constitutive DNA replication stress. Elife. 2020;9. doi:10.7554/eLife.51963

16.   Zeman MK, Cimprich KA. Causes and consequences of replication stress. Nat Cell Biol. 2014;16:
2–9. doi:10.1038/ncb2897

17.   Muñoz S, Méndez J. DNA replication stress: from molecular mechanisms to human disease.
Chromosoma. 2016;126: 1–15. doi:10.1007/s00412-016-0573-x

18.   Branzei D, Foiani M. Maintaining genome stability at the replication fork. Nat Rev Mol Cell Biol.
2010;11: 208–219. doi:10.1038/nrm2852

19.   Bermejo R, Lai MS, Foiani M. Preventing Replication Stress to Maintain Genome Stability:
Resolving Conflicts between Replication and Transcription. Mol Cell. 2012;45: 710–718.
doi:10.1016/J.MOLCEL.2012.03.001

20.   Macheret M, Halazonetis TD. DNA replication stress as a hallmark of cancer. Annu Rev Pathol
Mech Dis. 2015;10: 425–48. doi:10.1146/annurev-pathol-012414-040424

21.   Di Micco R, Fumagalli M, Cicalese A, Piccinin S, Gasparini P, Luise C, et al. Oncogene-induced
senescence is a DNA damage response triggered by DNA hyper-replication. Nature. 2006;444:
638–642. doi:10.1038/nature05327

22.   Neelsen KJ, Zanini IMY, Herrador R, Lopes M. Oncogenes induce genotoxic stress by mitotic
processing of unusual replication intermediates. J Cell Biol. 2013;200: 699–708.
doi:10.1083/jcb.201212058

23.   Kotsantis P, Petermann E, Boulton SJ. Mechanisms of Oncogene-Induced Replication Stress:
Jigsaw Falling into Place. Cancer Discov. 2018;8: 537–555. doi:10.1158/2159-8290.CD-17-1461

24.   Weitzman MD, Fradet-Turcotte A. Virus DNA Replication and the Host DNA Damage Response.
Annu Rev Virol. 2018;5: 141–164. doi:10.1146/annurev-virology-092917-043534

25.   Moody CA. Impact of Replication Stress in Human Papillomavirus Pathogenesis. J Virol. 2018;93.
doi:10.1128/jvi.01012-17

26.   Lilley CE, Schwartz RA, Weitzman MD. Using or abusing: viruses and the cellular DNA damage
response. Trends in Microbiology. 2007. pp. 119–126. doi:10.1016/j.tim.2007.01.003

27.   Scheifele LZ, Cost GJ, Zupancic ML, Caputo EM, Boeke JD. Retrotransposon overdose and genome
integrity. Proc Natl Acad Sci USA. 2009;106: 13927–13932. doi:10.1073/pnas.0906552106

28.   San Filippo J, Sung P, Klein H. Mechanism of Eukaryotic Homologous Recombination. Annu Rev
Biochem. 2008;77: 229–257. doi:10.1146/annurev.biochem.77.061306.125255

29.   Symington LS, Rothstein R, Lisby M. Mechanisms and regulation of mitotic recombination in
saccharomyces cerevisiae. Genetics. 2014;198: 795–835. doi:10.1534/genetics.114.166140

30.   Powell SN, Kachnic LA. Roles of BRCA1 and BRCA2 in homologous recombination, DNA replication
fidelity and the cellular response to ionizing radiation. Oncogene. 2003;22: 5784–5791.
doi:10.1038/sj.onc.1206678

31.   Walsh CS. Two decades beyond BRCA1/2: Homologous recombination, hereditary cancer risk and
772   a target for ovarian cancer therapy? Gynecologic Oncology. 2015. pp. 343–350.
773   doi:10.1016/j.ygyno.2015.02.017

774   32.   Chernikova SB, Game JC, Brown JM. Inhibiting homologous recombination for cancer therapy.
775   Cancer Biol Ther. 2012;13: 61. doi:10.4161/CBT.13.2.18872

776   33.   Helleday T. Homologous recombination in cancer development, treatment and development of
777   drug resistance. Carcinogenesis. 2010;31: 955–960. doi:10.1093/carcin/bgq064

778   34.   Krajewska M, Fehrmann RSN, De Vries EGE, van Vugt MATM. Regulators of homologous
779   recombination repair as novel targets for cancer treatment. Frontiers in Genetics. 2015. p. 96.
780   doi:10.3389/fgene.2015.00096

781   35.   Villa F, Simon AC, Ortiz Bazan MA, Kilkenny ML, Wirthensohn D, Wightman M, et al. Ctf4 Is a Hub
782   in the Eukaryotic Replisome that Links Multiple CIP-Box Proteins to the CMG Helicase. Mol Cell.
783   2016;63: 385–96. doi:10.1016/j.molcel.2016.06.009

784   36.   Kouprina NYu, Pashina OB, Nikolaishwili NT, Tsouladze AM, Larionov VL. Genetic control of
785   chromosome stability in the yeast Saccharomyces cerevisiae. Yeast. 1988;4: 257–69.
786   doi:10.1002/yea.320040404

787   37.   Yuan Z, Georgescu R, Santos R de LA, Zhang D, Bai L, Yao NY, et al. Ctf4 organizes sister
788   replisomes and Pol α into a replication factory. Elife. 2019;8: e47405. doi:10.7554/eLife.47405

789   38.   Samora CP, Saksouk J, Goswami P, Wade BO, Singleton MR, Bates PA, et al. Ctf4 Links DNA
790   Replication with Sister Chromatid Cohesion Establishment by Recruiting the Chl1 Helicase to the
791   Replisome. Mol Cell. 2016;63: 371–84. doi:10.1016/j.molcel.2016.05.036

792   39.   Simon AC, Zhou JC, Perera RL, van Deursen F, Evrin C, Ivanova ME, et al. A Ctf4 trimer couples the
793   CMG helicase to DNA polymerase alpha in the eukaryotic replisome. Nature. 2014;510: 293–297.
794   doi:10.1038/nature13234

795   40.   Gambus A, van Deursen F, Polychronopoulos D, Foltman M, Jones RC, Edmondson RD, et al. A key
796   role for Ctf4 in coupling the MCM2-7 helicase to DNA polymerase alpha within the eukaryotic
797   replisome. EMBO J. 2009;28: 2992–3004. doi:10.1038/emboj.2009.226

798   41.   Abe T, Kawasumi R, Giannattasio M, Dusi S, Yoshimoto Y, Miyata K, et al. AND-1 fork protection
799   function prevents fork resection and is essential for proliferation. Nat Commun. 2018;9: 3091.
800   doi:10.1038/s41467-018-05586-7

801   42.   Fumasoni M, Zwicky K, Vanoli F, Lopes M, Branzei D. Error-Free DNA Damage Tolerance and
802   Sister Chromatid Proximity during DNA Replication Rely on the Polα/Primase/Ctf4 Complex. Mol
803   Cell. 2015;57: 812–823. doi:10.1016/j.molcel.2014.12.038

804   43.   Kouprina N, Kroll E, Bannikov V, Bliskovsky V, Gizatullin R, Kirillov A, et al. CTF4 (CHL15) mutants
805   exhibit defective DNA metabolism in the yeast Saccharomyces cerevisiae. Mol Cell Biol. 1992;12:
806   5736–5747. Available: http://www.ncbi.nlm.nih.gov/pubmed/1341195

807   44.   Pâques F, Haber JE. Multiple pathways of recombination induced by double-strand breaks in
808   Saccharomyces cerevisiae. Microbiol Mol Biol Rev. 1999;63: 349–404.

809   45.   Gerrish PJ, Lenski RE. The fate of competing beneficial mutations in an asexual population.

810   Genetica. 1998;102/103: 127–144. doi:10.1023/A:1017067816551

811   46.   Lang GI, Botstein D, Desai MM. Genetic variation and the fate of beneficial mutations in asexual
812   populations. Genetics. 2011;188: 647–661. doi:10.1534/genetics.111.128942

813   47.   Couce A, Tenaillon OA. The rule of declining adaptability in microbial evolution experiments.
814   Front Genet. 2015;6: 99. doi:10.3389/fgene.2015.00099

815   48.   Woods RJ, Barrick JE, Cooper TF, Shrestha U, Kauth MR, Lenski RE. Long-Term Dynamics of
816   Adaptation in Asexual Populations. Science. 2013;342: 1364–1367. doi:10.1126/science.1198914

817   49.   Chou H, Chiu H, Delaney NF, Segrè D, Marx CJ. Diminishing Returns Epistasis Among Beneficial
818   Mutations Decelerates Adaptation. Science. 2011;332: 1190–1193. doi:10.1126/science.1203799

819   50.   Khan AI, Dinh DM, Schneider D, Lenski RE, Cooper TF. Negative epistasis between beneficial
820   mutations in an evolving bacterial population. Science. 2011;332: 1193–1196.
821   doi:10.1126/science.1203801

822   51.   Barrick JE, Kauth MR, Strelioff CC, Lenski RE. Escherichia coli rpoB mutants have increased
823   evolvability in proportion to their fitness defects. Mol Biol Evol. 2010;27: 1338–1347.
824   doi:10.1093/molbev/msq024

825   52.   Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Microbial evolution. Global epistasis makes
826   adaptation predictable despite sequence-level stochasticity. Science. 2014;344: 1519–1522.
827   doi:10.1126/science.1250939

828   53.   Marad DA, Buskirk SW, Lang GI. Altered access to beneficial mutations slows adaptation and
829   biases fixed mutations in diploids. Nat Ecol Evol. 2018;2: 882–889. doi:10.1038/s41559-018-
830   0503-9

831   54.   Zeyl C, Vanderford T, Carter M. An Evolutionary Advantage of Haploidy in Large Yeast
832   Populations. Science. 2003;299: 555–558. doi:10.1126/SCIENCE.1078417

833   55.   Von Borstel RC, Cain KT, Steinberg CM. Inheritance of spontaneous mutability in yeast. Genetics.
834   1971;69: 17–27.

835   56.   Kunz BA, Peters MG, Kohalmi SE, Armstrong JD, Glattke M, Badiani K. Disruption of the RAD52
836   gene alters the spectrum of spontaneous SUP4-o mutations in Saccharomyces cerevisiae.
837   Genetics. 1989;122: 535–542.

838   57.   Puddu F, Herzog M, Selivanova A, Wang S, Zhu J, Klein-Lavi S, et al. Genome architecture and
839   stability in the Saccharomyces cerevisiae knockout collection. Nature. 2019; 1–5.
840   doi:10.1038/s41586-019-1549-9

841   58.   Rolef Ben-Shahar T, Heeger S, Lehane C, East P, Flynn H, Skehel M, et al. Eco1-dependent cohesin
842   acetylation during establishment of sister chromatid cohesion. Science. 2008/07/26. 2008;321:
843   563–566. doi:10.1126/science.1157774

844   59.   Panizza S, Tanaka T, Hochwagen A, Eisenhaber F, Nasmyth K. Pds5 cooperates with cohesion in
845   maintaining sister chromatid cohesion. Curr Biol. 2000;10: 1557–1564. doi:10.1016/S0960-
846   9822(00)00854-X

847   60.   Michaelis C, Ciosk R, Nasmyth K. Cohesins: chromosomal proteins that prevent premature
848   separation of sister chromatids. Cell. 1997;91: 35–45. doi:doi: 10.1016/s0092-8674(01)80007-6.

849   61.   Ciosk R, Shirayama M, Shevchenko A, Tanaka T, Toth A, Nasmyth K. Cohesin's binding to
850   chromosomes depends on a separate complex consisting of Scc2 and Scc4 proteins. Mol Cell.
851   2000;5: 243–254. doi:10.1016/S1097-2765(00)80420-7

852   62.   Srinivasan M, Fumasoni M, Petela NJ, Murray A, Nasmyth KA. Cohesion is established during DNA
853   replication utilising chromosome associated cohesin rings as well as those loaded de novo onto
854   nascent DNAs. Elife. 2020;9. doi:10.7554/eLife.56611

855   63.   St. Charles J, Petes TD. High-Resolution Mapping of Spontaneous Mitotic Recombination
856   Hotspots on the 1.1 Mb Arm of Yeast Chromosome IV. Klein H, editor. PLoS Genet. 2013;9:
857   e1003434. doi:10.1371/journal.pgen.1003434

858   64.   Sharp NP, Sandell L, James CG, Otto SP. The genome-wide rate and spectrum of spontaneous
859   mutations differ between haploid and diploid yeast. Proc Natl Acad Sci USA. 2018;115: E5046–
860   E5055. doi:10.1073/pnas.1801040115

861   65.   Serero A, Jubin C, Loeillet S, Legoix-Né P, Nicolas AG. Mutational landscape of yeast mutator
862   strains. Proc Natl Acad Sci USA. 2014;111: 1897–1902. doi:10.1073/pnas.1314423111

863   66.   Loeillet S, Herzog M, Puddu F, Legoix P, Baulande S, Jackson SP, et al. Trajectory and uniqueness
864   of mutational signatures in yeast mutators. Proc Natl Acad Sci USA. 2020;117: 24947–24956.
865   doi:10.1073/pnas.2011332117

866   67.   Fisher KJ, Buskirk SW, Vignogna RC, Marad DA, Lang GI. Adaptive genome duplication affects
867   patterns of molecular evolution in Saccharomyces cerevisiae. PLOS Genet. 2018;14: e1007396.
868   doi:10.1371/journal.pgen.1007396

869   68.   Scott AL, Richmond PA, Dowell RD, Selmecki AM. The Influence of Polyploidy on the Evolution of
870   Yeast Grown in a Sub-Optimal Carbon Source. Mol Biol Evol. 2017;34: 2690–2703.
871   doi:10.1093/molbev/msx205

872   69.   Dunham MJ, Badrane H, Ferea T, Adams J, Brown PO, Rosenzweig F, et al. Characteristic genome
873   rearrangements in experimental evolution of Saccharomyces cerevisiae. Proc Natl Acad Sci USA.
874   2002;99: 16144–16149. doi:10.1073/pnas.242624799

875   70.   Gresham D, Desai MM, Tucker CM, Jenq HT, Pai DA, Ward A, et al. The Repertoire and Dynamics
876   of Evolutionary Adaptations to Controlled Nutrient-Limited Environments in Yeast. PLoS Genet.
877   2008;4: e1000303. doi:10.1371/journal.pgen.1000303

878   71.   Lauer S, Avecilla G, Spealman P, Sethia G, Brandt N, Levy SF, et al. Single-cell copy number variant
879   detection reveals the dynamics and diversity of adaptation. PLOS Biol. 2018;16: e3000069.
880   doi:10.1371/journal.pbio.3000069

881   72.   Tsaponina O, Barsoum E, Åström SU, Chabes A. Ixr1 is required for the expression of the
882   ribonucleotide reductase Rnr1 and maintenance of dNTP pools. PLoS Genet. 2011;7.
883   doi:10.1371/journal.pgen.1002061

884   73.   Takayama Y, Kamimura Y, Okawa M, Muramatsu S, Sugino A, Araki H. GINS, a novel multiprotein
885   complex required for chromosomal DNA replication in budding yeast. Genes Dev. 2003;17: 1153–
886   1165. doi:10.1101/gad.1065903

887   74.   Tanaka H, Katou Y, Yagura M, Saitoh K, Itoh T, Araki H, et al. Ctf4 coordinates the progression of
888   helicase and DNA polymerase alpha. Genes to cells. 2009;14: 807–820. doi:10.1111/j.1365-

2443.2009.01310.x

75.   Orr HA, Otto SP. Does diploidy increase the rate of adaptation? Genetics. 1994;136: 1475–1480.

76.   Tlsty TD, Albertini AM, Miller JH. Gene amplification in the lac region of E. coli. Cell. 1984;37:
217–224. doi:10.1016/0092-8674(84)90317-9

77.   Slack A, Thornton PC, Magner DB, Rosenberg SM, Hastings PJ. On the Mechanism of Gene
Amplification Induced under Stress in Escherichia coli. PLoS Genet. 2006;2: e48.
doi:10.1371/journal.pgen.0020048

78.   Reddy G, Desai MM. Global epistasis emerges from a generic model of a complex trait. bioRxiv.
2020; 2020.06.14.150946. doi:10.1101/2020.06.14.150946

79.   Wünsche A, Dinh DM, Satterwhite RS, Arenas CD, Stoebel DM, Cooper TF. Diminishing-returns
epistasis decreases adaptability along an evolutionary trajectory. Nat Ecol Evol. 2017;1: 0061.
doi:10.1038/s41559-016-0061

80.   Schoustra S, Hwang S, Krug J, de Visser JAGM. Diminishing-returns epistasis among random
beneficial mutations in a multicellular fungus. Proc R Soc B Biol Sci. 2016;283: 20161376.
doi:10.1098/rspb.2016.1376

81.   Deutschbauer AM, Jaramillo DF, Proctor M, Kumm J, Hillenmeyer ME, Davis RW, et al.
Mechanisms of Haploinsufficiency Revealed by Genome-Wide Profiling in Yeast. Genetics.
2005;169: 1915–1925. doi:10.1534/GENETICS.104.036871

82.   Gerstein AC, Otto SP. Ploidy and the Causes of Genomic Evolution. J Hered. 2009;100: 571–581.
doi:10.1093/jhered/esp057

83.   Otto SP, Gerstein AC. The evolution of haploidy and diploidy. Current Biology. 2008. pp. R1121–
R1124. doi:10.1016/j.cub.2008.09.039

84.   Schacherer J, de Montigny J, Welcker A, Souciet JL, Potier S. Duplication processes in
Saccharomyces cerevisiae haploid strains. Nucleic Acids Res. 2005;33: 6319–6326.
doi:10.1093/nar/gki941

85.   Coïc E, Feldman T, Landman AS, Haber JE. Mechanisms of Rad52-independent spontaneous and
UV-induced mitotic recombination in Saccharomyces cerevisiae. Genetics. 2008;179: 199–211.
doi:10.1534/genetics.108.087189

86.   Prado F, González-Barrera S, Aguilera A. RAD52-dependent and -independent homologous
recombination initiated by Flp recombinase at a single FRT site flanked by direct repeats. Mol
Gen Genet. 2000;263: 73–80. doi:10.1007/PL00008677

87.   Payen C, Koszul R, Dujon B, Fischer G. Segmental Duplications Arise from Pol32-Dependent
Repair of Broken Forks through Two Alternative Replication-Based Mechanisms. PLoS Genet.
2008;4: e1000175. doi:10.1371/journal.pgen.1000175

88.   Naiman K, Pagès V, Fuchs RP. A defect in homologous recombination leads to increased
translesion synthesis in E. coli. Nucleic Acids Res. 2016;44: 7691–7699. doi:10.1093/nar/gkw488

89.   LaBar T, Phoebe Hsieh Y-Y, Fumasoni M, Murray AW. Evolutionary Repair Experiments as a
Window to the Molecular Diversity of Life. Curr Biol. 2020;30: R565–R574.
doi:10.1016/J.CUB.2020.03.046

90. Russo M, Crisafulli G, Sogari A, Reilly NM, Arena S, Lamba S, et al. Adaptive mutability of
colorectal cancers in response to targeted therapies. Science. 2019;366: 1473–1480.
doi:10.1126/science.aav4474

91. Thiagalingam S, Laken S, Willson JKV, Markowitz SD, Kinzler KW, Vogelstein B, et al. Mechanisms
underlying losses of heterozygosity in human colorectal cancers. Proc Natl Acad Sci USA. 2001;98:
2698–2702. doi:10.1073/pnas.051625398

92. D'Angiolo M, Chiara M De, Yue J-X, Irizar A, Stenberg S, Persson K, et al. A yeast living ancestor
reveals the origin of genomic introgressions. Nature. 2020; 1–6. doi:10.1038/s41586-020-2889-1

93. Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, et al. Genome evolution across
1,011 Saccharomyces cerevisiae isolates. Nature. 2018;556: 339–344. doi:10.1038/s41586-018-
0030-5

94. Steenwyk JL, Opulente DA, Kominek J, Shen X-X, Zhou X, Labella AL, et al. Extensive loss of cell-
cycle and DNA repair genes in an ancient lineage of bipolar budding yeasts. PLOS Biol. 2019;17:
e3000255. doi:10.1371/journal.pbio.3000255

95. Ghosal G, Chen J. DNA damage tolerance: a double-edged sword guarding the genome.
Translational Cancer Research. 2013. pp. 107–129. doi:10.3978/j.issn.2218-676X.2013.04.01

96. Hohl M, Mojumdar A, Hailemariam S, Kuryavyi V, Ghisays F, Sorenson K, et al. Modeling cancer
genomic data in yeast reveals selection against ATM function during tumorigenesis. PLOS Genet.
2020;16: e1008422. doi:10.1371/journal.pgen.1008422

97. Song W, Dominska M, Greenwell PW, Petes TD. Genome-wide high-resolution mapping of
chromosome fragile sites in Saccharomyces cerevisiae. Proc Natl Acad Sci USA. 2014;111: E2210-
8. doi:10.1073/pnas.1406847111

98. Capra JA, Paeschke K, Singh M, Zakian VA. G-Quadruplex DNA Sequences Are Evolutionarily
Conserved and Associated with Distinct Genomic Features in Saccharomyces cerevisiae. PLoS
Comput Biol. 2010;6: e1000861. doi:10.1371/journal.pcbi.1000861

99. Nagalakshmi U, Wang Z, Waern K, Shou C, Raha D, Gerstein M, et al. The Transcriptional
Landscape of the Yeast Genome Defined by RNA Sequencing. Science. 2008;320: 1344–1350.

100. Azvolinsky A, Dunaway S, Torres JZ, Bessler JB, Zakian VA. The S. cerevisiae Rrm3p DNA helicase
moves with the replication fork and affects replication of all yeast chromosomes. Genes Dev.
2006;20: 3104–16. doi:10.1101/gad.1478906

101. Szilard RK, Jacques P-E, Laramée L, Cheng B, Galicia S, Bataille AR, et al. Systematic identification
of fragile sites via genome-wide location analysis of gamma-H2AX. Nat Struct Mol Biol. 2010;17:
299–305. doi:10.1038/nsmb.1754