Title: Neural precursor grafts restore motor and non-motor functions in parkinsonian macaques

Authors: Florence Wianny[1, 2], Kwamivi Dzahini[1, 2], Karim Fifel[1, †], Charles R.E. Wilson[1], Agnieszka Bernat[1, ‡, §,] Virginie Dolmazon[1], Pierre Misery[1], Camille Lamy[1], Kenneth Knoblauch[1], Howard M. Cooper[1], Emmanuel Procyk[1], Henry Kennedy[1, 3], Pierre Savatier[1, 2], Colette Dehay[1, 2]* and Julien Vezoli[1, 4]*

Affiliations:

[1]Univ Lyon, Université Claude Bernard Lyon 1, Inserm U1208, Stem Cell and Brain Research Institute; Bron, France.

[2]Primastem; Bron, France.

[3]Institute of Neuroscience, State Key Laboratory of Neuroscience, Chinese Academy of Sciences (CAS) Key Laboratory of Primate Neurobiology; Shanghai, China.

[4]Ernst Strüngmann Institute (ESI) for Neuroscience in Cooperation with Max Planck Society; Frankfurt, Germany.

*Corresponding author. Email: julien.vezoli@inserm.fr (J.V.); colette.dehay@inserm.fr (C.D.).

[†]Present address: International Institute for Integrative Sleep Medicine (WPI-IIIS), University of Tsukuba; Tsukuba, Ibaraki 305-8575, Japan.

[‡]Present address: Laboratory of Molecular Diagnostics, Department of Biotechnology, Inter-collegiate Faculty of Biotechnology, University of Gdańsk and Medical University of Gdańsk; Gdańsk, Poland.

[§]Present address: Laboratory of Experimental Embryology, Institute of Genetics and Animal Biotechnology, Polish Academy of Sciences; Warsaw, Poland.

One Sentence Summary: Empirical evidence that cell therapy efficiently reverts cognitive and clinical motor symptoms in the non-human primate model of Parkinson's disease.

Abstract:

Cognitive deficits as well as sleep and biological rhythm disorders constitute the non-motor symptoms that significantly impact quality of life in Parkinson's disease. Few studies have evaluated the impact of cell replacement therapy on non-motor symptoms. Here we implement a multidisciplinary and multiparametric assessment of the therapeutic potential of bilateral grafts of neural precursors in a macaque model of Parkinson disease on both motor and non-motor markers of functional recovery. Integration and survival of grafted cells correlated with robust functional recovery in middle-aged animals. Histological examination showed that grafted NP cells acquired a dopaminergic and serotoninergic fate and promoted the dopaminergic phenotype in the host tissue. These findings point to complex cellular mechanisms underlying the capacity of the grafts to reverse the symptoms. We show unprecedented recovery from cognitive symptoms in addition to a strong clinical motor recuperation. Motor and cognitive recovery and partial circadian rhythm recovery correlated with the degree of graft integration into the host environment as well as with *in-vivo* levels of striatal dopaminergic innervation and function. Further, our findings suggest that positive benefits from grafts are much more likely to be obtained at early stages of the disease. This study underlines the importance of multidisciplinary assessment in view of clinical translation and provides empirical evidence that integration of neural precursors following transplantation in diverse functional domains of the basal ganglia efficiently restores function at multiple levels in parkinsonian non-human primates.

Main Text:

INTRODUCTION

Parkinson's disease (PD) is a neurodegenerative condition affecting up to 10M of the worldwide population with incidences increasing with age, making PD the fastest growing neurological disorder in a globally aging population [1]. The clinical manifestation is primarily motor with a typical parkinsonian syndrome including bradykinesia, rigidity and resting tremor and is confirmed by clinical diagnostic tools including *in-vivo* imaging of the denervation of the nigrostriatal dopaminergic (DA) axis [2]. Unfortunately, 60-80% of DA cells are lost prior to the onset of clinically diagnosed motor symptoms [3]. During this so-called premotor period preceding the clinical threshold, the DA-lesion is progressive and accompanied by the manifestation and further deterioration of non-motor symptoms including cognitive disorders as well as perturbation of circadian rhythm and sleep disorders [4, 5]. Although considered secondary, non-motor symptoms nonetheless have a significant impact on quality of life; the cognitive abilities of the vast majority of PD patients deteriorate leading to psychiatric disturbances [6] with circadian perturbations affecting biological rhythms [7].

DA lesion is the hallmark of PD and in addition to circadian regulation affects multiple frontal lobe functions including performance monitoring, motivational aspects of behavior and motricity. However, there is currently no cure for PD and palliative therapies such as levodopa and Deep-Brain Stimulation (DBS), mainly correct motor symptoms and often give rise to behavioral perturbation [8]. For example, DBS shows no long-lasting effect on axial symptoms e.g. freezing of gait and negatively impacts cognitive and neuropsychiatric symptoms [9, 10]. Importantly palliative therapies do not solve the issue of neuronal loss and the long-term prognosis. Consequently, cell replacement therapy for PD has been recently re-evaluated as a potential cure for PD [11], leading to recent trials [12] aimed at improved grafting procedures, and to pave the way for stem-cell based transplantation in humans. In this respect there is a recognized need for more detailed perspectives from pre-clinical investigations in animal models [13].

With the aim of optimizing the therapeutic use of cell replacement in PD we have examined the global consequence of cell grafts at different stages of the disease in a macaque model of PD. While numerous efforts have been deployed to use non-motor symptoms as premotor markers for the prognosis of PD [14], virtually no studies have addressed the impact of cell replacement on non-motor symptoms [15] leading to insufficient appraisal of their multi-parametric impact at either pre-clinical or clinical stages [8, 16]. This is problematic because advanced PD with an extensive nigrostriatal DA lesion constitutes a highly deleterious environment for the survival of grafted cells [17, 18]. Hence in the present study we use an experimental design that allows us to address the capacity of cell grafts to reverse the disease at disease onset in the motor phase and assess the impact of DA lesion extent on graft efficiency. This design also allowed us to characterize the dynamics of graft induced recovery by comparing it to spontaneous recovery in the premotor stage. Because non-motor symptoms have a gradual onset prior to clinical manifestations of DA denervation, evaluation of the effects of NP grafts in parkinsonian non-human primates (NHP) requires a longitudinal multi-parametric investigation, which we have implemented in the present study. We grafted neural precursors (NPs) because they hold the promise of enhanced efficiency compared to DA neurons [19-21], as in addition to their neuro-restorative potential NPs have the capacity to differentiate into neuroprotective astroglial cells [22-24].

Repeated systemic injections of low-doses of MPTP (0.2 mg/kg every 3-4 days) induced a parkinsonian syndrome with a slow and typically dorso-ventral progression of the nigrostriatal denervation, together with a premotor expression of cognitive and circadian *i.e.* non-motor symptoms [25-27]. Here, we implemented a multidisciplinary / multiparametric approach to assess the therapeutic potential of bilateral grafts of NPs in multiple functional domains of the basal ganglia of NHP (*macaca fascicularis*). For ethical reasons we reduced the number of animals used by monitoring individual macaques so that

each case constitutes its own control. This required continuous assessment of clinical motor and non-motor symptoms as well as *in-vivo* monitoring of the nigrostriatal lesion with Positron Emission Topography (PET) imaging of DA transporters (DAT). Monitoring was prior to and following the induction of stable motor symptoms [27], and subsequent to transplantation of NPs. In addition, Fluoro-DOPA (^{18}F-DOPA) imaging allowed assessment of the impact of the NP grafts on DA function [28]. Finally, *post-mortem* immunohistological examination of NP grafts was performed in order to evaluate the impact of the graft on host tissue as well the survival, integration and differentiation fate of the grafted cells into the host brain.

RESULTS

The experimental design (**Fig.1A, Table S1**) allowed full characterization of MPTP intoxication. In Stage I, all animals show non-motor symptoms [25, 26] and a nigrostriatal DA denervation superior to 70% [27] but, at this stage, spontaneous recovery from motor symptoms occurs and this was investigated in 3 out of the 6 individual cases [25, 27, 29]. Comparison of cognitive, circadian and DA function markers enabled characterization of spontaneous and graft-induced recovery. Subsequent to Stage I, resuming MPTP injections until a clinical score of 10 is reached on two consecutive days [27] leads to a Stage II, where motor symptoms become permanent. In order to implement this experimental strategy it is necessary to adapt cumulated MPTP-doses to individual clinical motor scores (**Table S2**) thereby taking into account the known variability of response to MPTP and adjustment of the intoxication protocol (**Fig.1A**, **Table S2**, see **Materials and Methods**). Clinical scores and behavioral measures were divided into quantiles (Q1-5, as described in [25], see **Materials and Methods**) in order to compare cases on the basis of the presumed DA-lesion rather than on the time spent in the premotor (MPTP), motor (post-MPTP/pre-graft) and post-graft periods. We compared groups (*recovered* vs. *non-recovered*) across control, premotor and motor quantiles (pre-/post-MPTP) to evaluate differences in behavioral markers between groups due to MPTP intoxication and, across pre- and post-grafts quantiles (pre-/post-graft) in order to evaluate differences between groups due to NP grafts (one-way anova: F-test (groups,df), p-value, effect-size (η_p^2), see **Materials and Methods**).

Behavioral and functional markers of chronic MPTP-induced Parkinsonism

During Stage II, all cases displayed a stable parkinsonian syndrome with clinical motor scores ranging from mild to severely symptomatic (**Figs.1B, S1A**). In two out of the five grafted cases, we failed to observe recovery from motor symptoms whereas the three other cases exhibited robust recovery from motor symptoms with a clinical motor score returning below the symptomatic threshold (pre-/post-graft: $F(2,10) = 126, p < 2E\text{-}16, \eta_p^2 = 0.50$ while effect-size was clearly reduced over pre-/post-MPTP quantiles: $F(2,11) = 11, p < 2E\text{-}16, \eta_p^2 = 0.14$, **Figs.1B, S1A, Table S8**). We accordingly assigned cases to the *non-recovered* or *recovered* group based on clinical motor outcome following NP grafts (**Figs.1A-B, S1A**).

During the pre-Graft period, non-motor symptoms emerged in all animals with progressive deterioration during the premotor period *i.e.* before motor scores reached clinical thresholds (**Figs.1B, S1**). Alterations of circadian rhythms appeared progressively during the premotor phase (**Figs.1B, S1C**) to reach peak levels during the symptomatic period as previously described [26]. This pattern of decreased (increased) activity during the Light (Dark) phase resembles circadian alterations of the sleep-wake cycle observed in PD patients which manifests as increased daytime sleepiness and fragmented sleep structure [30]. The premotor and motor alteration of rest-wake locomotor rhythms in all cases exhibited reduced light/dark (LD) ratios (minimal effect-size of group difference across pre-/post-MPTP quantiles: $F(2,11) = 4.3, p = 3.2E\text{-}16, \eta_p^2 = 0.07$, **Fig.1B, Table S8**) and relative amplitude, as well as increased rhythm variability and fragmentation (**Fig.S1C**). In all cases cognitive performance deteriorated during MPTP intoxication and after offset (pre-/post-MPTP quantiles, no group difference in Success: $F(2,11) =$

1.8, $p > 0.05$, $\eta_p^2 = 0.11$, and similarly minor effect-size of group for Errors: $F(2,11) = 2.1$, $p = 0.03$, $\eta_p^2 = 0.12$, **Figs.1B, S1B**).

Fig. 1. Motor, non-Motor and Functional Markers of Graft-induced Recovery. (**A**) Study design. At early stages of MPTP treatment (Stage I) there is spontaneous recovery from motor symptoms but not from cognitive and circadian symptoms. Continued MPTP intoxication leads to Stage II where motor symptoms now become permanent. (**B**) Cases are grouped in *recovered* (black) vs. *non-recovered* (grey) according to the level of parkinsonian motor-symptoms following cell therapy. Case2 excluded because of per-operative brain hemorrhage following graft. Motor symptoms clinically scored with the Parkinsonian Monkey Rating Scale (PMRS top panel), Cognitive performances (range normalized to control performance, -1 for success, middle panel) and, Chronobiological rhythms reflected by the ratio of spontaneous locomotor activity during the light and dark periods (normalized contrast with Light/Dark ratio in control conditions, bottom panel). Means ± SE, anova between groups, adjusted p-values for multiple comparisons * < 0.05, *** < 0.001, see **Table S8**. (**C**) Increase of nigrostriatal innervation. Parametric maps of DAT binding potential with axial (top), sagittal (bottom-left panel) and coronal (bottom-right panel) views of significant positive difference for the contrast Pre- vs. Post-Graft (color scale from 0.2-1, case1, recovered group). See **Fig.S1** for individual behavioral markers and **Fig.S9** for statistical thresholding in (**C**) and negative difference.

Nigrostriatal DA innervation evaluated by striatal DAT binding, decreased by more than 80% relative to control measures of striatal DAT levels with a relatively spared binding in the ventral striatum [27]. This

dorsoventral pattern of the nigrostriatal lesion was later confirmed by semi-quantitative evaluation of *post-mortem* immunostaining for DAT and TH (**Fig.2**) with a significantly more advanced lesion in *non-recovered* cases (**Fig.2C**).

Fig. 2. Dopaminergic nigrostriatal lesion. Immunostaining for DAT (upper row) and for TH (lower row). (**A**) Optical density (O.D.) measures of DA lesion (arbitrary units) is made on post-mortem tissue of controls, *recovered* MPTP, *non-recovered* MPTP. (**B**) Illustrative histological sections showing immunostaining. (**C**) Percent change in expression levels between MPTP and Controls provides DA lesion estimates by comparing MPTP sub-groups (two-sided Wilcoxon rank sum test, Bonferroni correction for multiple comparisons *** $p < 0.001$, $Z \geq 5.75$), *Recovered* (black) vs. *Non-recovered* (gray). Mean±SE. *Recovered* group: cases 1, 4 and 5; *Non-recovered* group: cases 3 and 6; Controls group: n=3.

Behavioral and functional markers following NP grafts

NPs were derived from a rhesus embryonic stem cell line (LYON-ES1) stably expressing the fluorescent marker tau-GFP [31, 32], which allows precise tracking of the grafted NPs and their integration into the host circuits. We chose to graft NPs given their inherent neuroprotective potential resulting from interaction with host-cells [33] and their observed higher survival rate compared to mature DA cell grafts [22, 34, 35]. Because *in-vitro* NPs show both astroglial and neuronal -including DA- fates (**Fig.S2**), it is possible to assess the influence of the host-environment on the differentiation fate of transplanted cells. NP grafting was delayed by 3 months on average following the last MPTP injection (13±6weeks, see **Materials and Methods**). Bilateral transplantations in multiple functional domains of the basal ganglia potentiated graft efficiency on both motor and non-motor symptoms. Grafts were targeted in striatal regions involved in cognitive (*i.e.* anterior caudate nucleus – ant-CdN) and motor functions (*i.e.* posterior putamen – post-Put.), as well as in the substantia nigra (SN), the latter providing the source of nigrostriatal DA innervation *i.e. pars compacta*, and part of the frontal DA innervation.

Following NP grafts, the reduction of the clinical motor score was associated with an additional cognitive and circadian recovery in the *recovered* group (**Figs.1B, S1**). In the *recovered* group, clinical scores progressively and significantly diverged at the 2nd quantile from those of the *non-recovered* group, whereas group differences were significant after the 3rd quantile for non-motor symptoms (**Fig.1B**). Clinical scores dropped below symptomatic thresholds on average 6 weeks following NP grafts, to ultimately achieve maximal motor recovery 5 weeks later (**Figs.1B, S1A**). In the *non-recovered* group, the initial cognitive and LD ratio recovery shortly after the graft was followed by a relapse on the last three to two quantiles (**Fig.1B**). However, the effect of the group difference of non-motor markers in pre-/post-graft was 3-5 times higher than in pre-/post-MPTP (with large effect-size for Success: $F(2,10) =$

10, $p = 3.1E\text{-}13, \eta_p^2 = 0.27$ and Errors: $F(2,10) = 11, p = 1.6E\text{-}14, \eta_p^2 = 0.29$, and moderate for LD ratio: $F(2,10) = 12, p < 2E\text{-}16, \eta_p^2 = 0.094$, **Fig.1B, Table S8**). Non-parametric circadian rhythm analyses (see **Material and Methods**) revealed high variability across cases, preventing post-graft group comparison on those markers (**Fig.S1C**). In the *recovered* group, markers of striatal DA function and DAT increased following NP grafts, in correlation with the observed behavioral recovery (**Figs.1B-C, 3B-C**). Contrasting with the beneficial effects of NP grafts in the *recovered* group, both clinical motor scores and non-motor symptoms persisted following NP graft in the *non-recovered* group (**Figs.1B, S1**), in agreement with the absence of change in markers of striatal DA function and DAT (**Fig.3B**).

Fig. 3. Functional markers of striatal DA function and DA terminals. (**A**) Images of striatal DAT binding potential (BP$_{ND}$, color scale 0-7, case1 – see **Fig.S3A** for all cases). (**B**) Functional markers of nigrostriatal innervation (normalized BP$_{ND}$ of ^{11}C-PE2i) and DA function (normalized Ki, uptake of ^{18}F-DOPA) in a *recovered* and a *non-recovered case*) during expression of stable symptoms (PD) and following transplantation (Post-Graft, minimum of 10weeks between measures). Min-max normalization per case. (**C**) Percent change in Striatal DAT binding in Grafted and Non-grafted regions for *recovered* (case1, black) and *non-recovered* (cases 3 and 6, grey). Purple, Striatal DAT binding following induced spontaneous recovery (case1). Two-sided Wilcoxon rank sum test, Bonferroni correction for multiple comparisons ** p < 0.01, Z = 2.92.

In the *recovered* group; we observed that PET-scan estimation of nigrostriatal innervation as revealed by DAT binding significantly increased bilaterally in the ventral striatum as early as 15 weeks post-graft (**Fig.1C**). This suggests a primarily protective influence of grafts on the remaining pool of nigrostriatal projection neurons. Post-graft F-DOPA scans confirmed a selective effect of NPs on DA synthesis (**Fig.3B**) that was associated with maximal motor recovery. Significant increases in DA function and DAT binding were also observed in grafted parts of the striatum when compared to the same regions in *non-recovered* cases (**Fig.3C**).

We performed sham transplantation (see **Materials and Methods**), in two out of the four cases following PET-scan imaging of striatal DAT innervation. No recovery from motor nor from non-motor symptoms was observed following sham grafts (**Fig.S3B**). Further, functional markers displayed no difference between *recovered* vs. *non-recovered* cases in striatal DAT binding and no specific increase in sham-grafted cases (**Fig.S3C**).

Graft-induced recovery differs from spontaneous recovery

We compared the time course of behavioral symptoms and functional markers following graft-induced recovery to those following spontaneous recovery. In half of the cases (n=3), spontaneous recovery was initially induced by halting MPTP intoxication as described previously (Stage I, **Fig.1A**) [25, 27]. Spontaneous recovery of clinical motor symptoms displayed faster dynamics and was more complete compared to graft-induced recovery (large effect-size: $F(2,13) = 24, p < 2.2E\text{-}16, \eta_p^2 = 0.40$, **Fig.4A**). Importantly, spontaneous recovery was not accompanied by changes in cognitive performance which contrasted with graft induced recovery that was invariably accompanied by progressive cognitive improvements (**Fig.4B**). These findings are intriguing because spontaneous motor recovery occurs despite an average 71.5±9.7% reduction in striatal DAT relative to control levels and that stable symptoms were observed when the lesion reached 82.3±7.9% of control [27]. The present findings show that graft-induced recovery occurs despite an average 78.04 ±12% reduction in striatal DAT binding (**Fig.3C**). These results confirm the specificity of graft-induced recovery compared to spontaneous recovery, particularly with respect to cognitive deficits (large effect-size for both Success: $F(2,5) = 39, p < 2.2E\text{-}16, \eta_p^2 = 0.40$ and Errors $F(2,5) = 35, p < 2.2E\text{-}16, \eta_p^2 = 0.38$, **Fig.4B**).

Fig. 4. Dynamics of spontaneous vs. graft-induced behavioral recovery. (**A**) Clinical motor score over weeks and (**B**) cognitive performances (over quantiles) comparison between spontaneous recovery (cases 1, 3 and 4) vs. graft-induced recovery (cases 1, 4, 5). Mean ±SE, anova between groups, adjusted p-values for multiple comparison * < 0.05, *** < 0.001.

Behavioral and functional recovery correlate with graft survival

We grafted NPs bilaterally in multiple sites including the ant-CdN, post-Put and SN. Each site received $1.5\times10^5\text{-}1\times10^6$ NP cells (see **Materials and Methods**). Following an average post-graft period of 30±8weeks, we performed a *post-mortem* immuno-histological assessment of tau-GFP+ cells survival and integration into the host milieu (**Fig.1A**).

Noteworthy, the *non-recovered* group is characterized by a complete absence of tau-GFP+ cells (**Fig.S4**). This could be due to the fact that the nigrostriatal DA lesion was most severe in these cases (**Fig.2C**). However, immunological rejection of the grafted NPs cannot be excluded as no immunosuppressive treatment was employed. The fact that non-motor symptoms initially improved in the *non-recovered*

group prior to an abrupt deterioration after the 2nd Quantile (**Fig.1B**) is coherent with a chronic graft
rejection in the *non-recovered* group (**Fig.S4**).
In sharp contrast, we found surviving grafted cells in all cases of the *recovered* group, at locations
including the targeted functional domains of the basal ganglia (**Figs.5, S5-8**). The average number of
surviving cells was: $1.23\pm0.9\times10^4$ cells in the ant-CdN, $1.95\pm1.1\times10^4$ cells in the post-Put and
$5.93\pm3.0\times10^4$ cells in the SN (**Table S3**) which corresponds to an average survival rate of: $6.9\pm6.2\%$ in
the ant-CdN, $13.0\pm7.3\%$ in the post-Put and $15.5\pm9.4\%$ in the SN (overall survival rate of $11.6\pm4.4\%$,
Mean±SE).

Fig. 5. Post-mortem localization of grafted NPs. (**A**) Digital reconstructions of post mortem grafts in case1 (case4
shown in **D** and case5 in **E** - scale bars 200μm). (**B-C**) Photomicrographs of brain slices with zoom on tau-GFP$^+$ grafts (scale
bar 2mm, case4 shown in **B** and case5 in **C**).

We observed unilateral survival of NPs in some of the grafted striatal regions (**Table S3**). We could not
link asymmetrical survival in the post-Put to a systematic difference in left vs. right motor recovery
(**Fig.S9A**) suggesting the induction of known cross-hemispheric compensatory processes [36].
Nevertheless, small differences were observed in cognitive performance linked to unilateral survival in
the ant-CdN (**Fig.S9B**); differences that were coupled with a minor unilateral DAT binding decrease in
the corresponding striatal region (**Fig.S9C**, right).
Surviving grafts displayed extensive tau-GFP positive projection fibers (**Fig.6A-C**) which extend several
millimeters from the graft core, considerably further than the local processes which are less than 1
millimeter (**Fig.6D**). These findings indicate that the *recovered* group is characterized by integration of
NPs into the host tissue (**Fig.6**).

Differentiation fates indicate protective and restorative effects of NP grafts

Detailed immuno-histological examination of graft locations reveals that grafted NPs differentiated
principally into astrocytes and mature neurons (**Fig.7A-C, D; Fig.S5**). Among surviving NPs, astrocytic
fate accounted for $42\pm5\%$ (GFAP$^+$/tau-GFP$^+$ co-expression ratio, **Fig.7A**, **Table S4**) and neuronal fate
for $16\pm4\%$ (MAP2$^+$/tau-GFP$^+$ co-expression ratio, **Fig.7A**, **Table S4**). Furthermore, we demonstrate that
NPs spontaneously differentiate into aminergic (TH+/DAT+) cells thereby potentially restoring a
significant proportion of the lost DA pool (**Fig.7C-D**).
On average, the TH$^+$/tau-GFP$^+$ co-expression ratio implies that $6\pm2\%$ of tau-GFP$^+$ cells differentiated
into aminergic cells (**Fig.7B**, **Table S4**). Up to 13% of tau-GFP$^+$ in SN graft co-localize with TH$^+$
expression (case4), which corresponds to an estimate of 1.4×10^4 TH$^+$/tau-GFP$^+$ cells and hence to a

restoration of nearly 18% of the original pool of DA-containing midbrain neurons in the same species
[37]. We qualitatively confirmed NP differentiation into both the serotoninergic and the dopaminergic
phenotypes (**Fig.7H**). Together with the protective effect of the graft on the survival or upregulation of
DA cells (**Figs.1C, 7F-G**), these results strongly support a role of the graft in the recovery from motor
and cognitive symptoms (**Figs.1**, **S1** and **Tables S4-S5**).

Fig.6. Grafted NPs project over long distances. Digital post-mortem reconstructions of NP grafts at the level of (**A**)
anterior CdN in case1 (left) and case5 (middle and right), (**B**) posterior Put for cases 1 (left) and 5 (right) and, (**C**) SN for
cases 1, 4 and 5 (from left to right). Tau-GFP+ projection fibers (clear blue) are found at relatively distant locations from the
graft core (dark green), much farther than local graft processes (clear green) emanating from the core. (**D**) Typical processes
emanating from the core have an average length superior to 340μm. Left, count histogram of processes' length for case5.
Right, typical tau-GFP+ photomicrograph showing graft-core and processes at the level of ant-CdN (case5, faint purple lines
from measurement overlay, scale bar 100μm).

At some graft sites, aggregates of host cells (*i.e.* tau-GFP negative, **Figs.S7E, G and S8D, H**) were observed in the graft-core of nigral transplants, exhibiting aminergic TH$^+$, dopaminergic DAT$^+$ and sparse serotoninergic SERT$^+$ processes (**Figs.7F-G, S7G-J**). Importantly, no post-graft overgrowth was observed (**Fig.S10A-C**) and few CD68$^+$ macrophages were detected in the vicinity of transplanted sites, despite the absence of immunosuppressive treatment (**Fig.S10D-F**).

Fig. 7. Differentiation Fate of grafted NPs. (**A**) Fraction of tau-GFP$^+$ co-localized with TH$^+$, MAP2$^+$ and GFAP+. Illustrative example of NPs differentiated into: (**B**) aminergic cells, SN, case4 (arrowheads); (**C**) aminergic cells, SN, case5 (arrowheads, left is enlarged view of region framed on the right); (**D**) neuronal cells (arrowheads) and processes (arrows), post-Put, case5; (**E**) astroglial cells (arrowheads) and processes (arrows), SN, case4. (**F-G**) Host tissue within the graft core presenting numerous dopaminergic (DAT$^+$, blue) and sparse serotoninergic (SERT$^+$, red) processes. (**H**) NPs also display serotoninergic and/or dopaminergic phenotype (arrowheads). Scale bars, **B** 20µm, **C** left 50 µm a, **D-E** 100µm, **F** 25µm, **G** 5µm, **H** 15µm.

Finally, we explored the extent to which the grafted NPs were structurally integrated, via contacts with host neurons. For this purpose, we examined synaptophysin (SYN) immunohistology. SYN is localized

in the membrane of pre-synaptic vesicles and provides a specific and sensitive marker for synapses. We explored regions at the periphery of the grafts, where the numerous local processes emanating from the core appear preferentially directed toward host DA cells (**Fig.8A, C**). Confocal microscopy shows that SYN co-localized at the junction between TH$^+$ and MAP2$^+$ host-cells and tau-GFP$^+$ processes from grafted-cells (**Figs.8E, S8E, G, I-K**), indicating that local processes stemming from the graft core establish synaptic contacts with the remaining host DA cells in the SN (**Fig.8**). Together with the presence of long-distance projections emanating from the grafted sites (**Fig.6**), this suggests functional integration of NPs into the host brain circuitry and supports the effect of NP grafts in the observed functional recovery (**Figs.1B-C, 3**).

Fig.8. Histological evidence of NP integration and interaction with host cells. (**A**) Photomicrographs (up-left panel) shows position of graft in SN, framed region zoomed in bottom-left panel. Right-panel show zoom of framed region in the bottom-left panel. In (**A**) and (**B**), Grafted NPs (tau-GFP$^+$, green) extend processes (arrowheads) toward surviving host DA cells (SN grafts). (**C**) NPs display numerous extensive processes projecting outside the graft-core. (**D**) NPs differentiate into aminergic phenotype displaying TH$^+$ processes (arrowhead). (**E**) Synaptophysin expression (SYN, blue) co-localized with grafted cell (tau-GFP$^+$) differentiated into mature neuron (MAP2$^+$) suggesting synaptic contact. Scale bars: **A-C** 100μm, **D-E** 5μm.

DISCUSSION

We transplanted macaque NPs into multiple functional domains of the basal ganglia, in both hemispheres of middle-aged parkinsonian macaques. Our study demonstrates that the successful integration of the grafts promotes a robust clinical and unprecedented cognitive recovery (**Figs.1B, S1**), as well as partial circadian recovery. Recovery is accompanied by quantitative effects on functional markers of nigrostriatal DA innervation and striatal DA function (**Figs.1C, 3B-C**). This study thus provides major empirical evidences in favor of a promising stem-cell derived cell therapy for PD patients. First, we show

for the first time that transplantation and successful engraftment of NPs efficiently promotes recovery from motor and non-motor symptoms. Second, we demonstrate the restorative and the neuroprotective potential of grafted NPs. Our results show that NPs have the capacity to restore a significant pool of mature neurons (**Fig.7A**), including dopaminergic and serotoninergic cells. In addition, NPs largely differentiate into astroglial cells that the present study confirms provide a neuroprotective environment for the surviving pool of DA neurons [22-24]. Third, our results also suggest that nigral transplantation of NPs provide neurotrophic effects on the host cells (**Figs.7F-G, S7E, G-J and S8D, H**), orienting endogenous cells towards a DA fate [38, 39]. Finally, we provide cellular evidences of structural and functional integration of grafted NPs (**Figs.6, 8 and S8E, G, I-K**) into the host brain network.

The consistent cognitive recovery following successful engraftment of NPs in ant-CdN and SN establishes the potential of cell therapy to alleviate non-motor cognitive symptoms in addition to motor symptoms in PD patients. The specificity of the graft-induced cognitive recovery (**Fig.1B**) was further confirmed by the absence of cognitive improvements following sham-grafts (**Fig.S3B**) and spontaneous motor recovery (**Fig.4B**). These results anticipate the outcome of cell therapy for the majority of PD patients experiencing mild cognitive impairment (MCI) that significantly affects their quality of life [40]. Cognitive functions in human and NHP depend on the integrity of the prefrontal cortex (PFC), the DA system and the DA innervation of PFC via direct projections from SN and ventral tegmental area [41-46]. In addition, fronto-striatal circuits are also extensively involved in cognitive function and more particularly the cognitive sectors of the striatum i.e. ant-CdN [47]. Many cognitive symptoms expressed in PD patients not only arise from DA degeneration but also from other monoamine neurotransmitter systems closely connected to the DA system i.e. serotonin and norepinephrine [48, 49]. These systems are also affected in the MPTP-monkey model [50]. With the cognitive task that was used here, increased errors and reduced success rates observed in Stages I and II can be directly related to the reduced control over motor planning and parallel executive dysfunctions observed in PD patients [4, 51]. Hence, the observed reduction of cognitive performances (**Figs.1B, S1B**) can be directly imputed to the nigrostriatal DA lesion, the altered DA innervation of frontal cortex [41] and the wider MPTP-induced alteration of the aminergic system including noradrenalin and serotonin [29, 52-55]. We relate the consistent cognitive recovery following NP grafts (**Fig.1B**) to the efficiency of the transplanted NPs in both SN and ant-CdN of which an average of, respectively $5.93\pm3\times10^4$ and $1.23\pm0.9\times10^4$ cells survived, differentiated (**Fig.7**) and integrated the host network through local and long-distance projections (**Figs.6, 8**). Importantly, we show that grafted NPs can differentiate into a substantial proportion of TH$^+$ expressing cells (**Fig.7A**), well within the range of previous studies reporting clinical improvement (**Table S4**), but also in the serotoninergic phenotype (**Fig.7H**) thereby supporting their role not only in motor but also in cognitive recovery.

In addition to these restorative effects, our results suggest that bilateral multi-site NP grafts exert a neuroprotective influence on the pool of surviving DA cells and of their ventro-striatal target. We show a significant and specific increase in striatal DAT binding following graft (**Fig.1C**) and not following sham-surgery (**Fig.S3C**). This increase is located in the ventral striatum that corresponds to the surviving pool of nigrostriatal innervation following induced spontaneous recovery [27]. The large proportion of astroglial differentiation (**Fig.7A**) might support this protective effect of NP grafts, in accordance with previous reports on human NP grafts in MPTP-monkeys [56]. Although we report some degree of recovery of DAT binding in the grafted caudate nucleus and putamen (**Fig.3B-C**), these effects were minimal compared to what was found in the ventral striatum. Longer post-graft survival could lead to greater increases in DAT binding. Indeed, transplanted PD patients display fast clinical improvements in motor function i.e. within 2-3 months after transplantation, whereas it can take 1-2 years to detect via functional imaging significant increase in DA function at the grafted locations [57-62].

The impact of cell therapy on chronobiological rhythms as far as we know has not been previously studied in the MPTP-monkey model. The initially observed circadian recovery following NP grafts

concerned both groups but was followed very quickly by a deterioration after the third quantile in *non-recovered* cases. However, in the *recovered* group the LD ratio recovery was significant but only partial (**Fig.1B**). Previously we showed that following MPTP-induced nigrostriatal DA lesion, environmental cues mask the failure of the intact SCN to drive striatal clock genes and DA functions that control rest-wake locomotor rhythms [26]. Here, the partial recovery of circadian rhythms in the presence of environmental timing cues (**Fig.1B**), could reflect incomplete restoration of DA function. However, following NP grafts, under continuous light conditions (n=4, cases 1, 3, 4 and 5) there was an absence of endogenous circadian rhythm as observed in the MPTP monkey [26]. This again points to NP grafts failing to fully restore the DA network involved in the SCN control of rest-wake rhythms. DA projection neurons from SN and VTA regulate sleep-wake rhythms through interaction with several structures from the circadian network including the locus coeruleus, the lateral hypothalamus and the pedunculopontine nucleus [63]. Those structures, which were not grafted in the present study, are impacted in PD and following MPTP intoxication and thus constitute future candidates for cell replacement protocols [64-66].

There are limitations in the use of cell-therapy for PD e.g. presence of Lewy-body formations in the grafted cells [67, 68]. However, the balance benefit-risk of cell-therapy, even in cases of Lewy-body development in the grafted cells, is clearly in favor of the transplanted patients [11]. Three decades after the first clinical trials of transplanting cells from fetal ventral mesencephalon in PD patients [69], cell replacement therapy in PD is still not a proposed treatment. However, this therapeutic approach has recently been shown to be efficient at different levels [61, 62, 70], leading to more recent trials [71]. In parallel, many strategies have been developed in recent years in order to improve the safety and efficiency of using ESCs or iPSCs as a source for the production of the to-be-transplanted cells [21]. Virtually no NHP studies have addressed the impact of cell replacement on non-motor circadian symptoms and only one recent reported beneficial effects on depressive symptoms (**Table S5**). Here we report for the first-time efficient recovery from cognitive and clinical motor symptoms, as well as partial circadian recovery following NP grafts in the NHP model of PD. The present study confirms the neuroprotective and restorative capacities of NP grafts and, underlines the crucial importance of longitudinal and multifaceted approaches for a better translation to the clinic and appraisal of early-phase clinical trials.

While there are doubtless limitations in the use of the MPTP-monkey model to study human-specific aspect of PD; this model remains nevertheless the gold standard for modeling PD and for the pre-clinical evaluation of new therapeutic approaches. Previous studies have shown that advanced nigrostriatal lesion is detrimental to the survival of grafted cells [17, 18], which is coherent with our observation of poor survival of grafted NPs in the *non-recovered* group. Further, in the *non-recovered* group immune rejection is likely to have occurred as previously observed following allogenic transplantation in non-immunosuppressed monkeys [72, 73]. These observations suggest that future therapeutic cell replacement procedures should be carried out preferably at early stages of PD.

MATERIALS AND METHODS

In agreement with the 3Rs [74], the rationale for the current study is that each case represents its own control through detailed follow-up of all the periods of the protocol i.e. CTR, MPTP, post-MPTP and Pre- vs. Post-Graft. We published precise and detailed reports for each clinical, behavioral and functional parameter followed as well as for the characterization of NPs grafted in the present study. These precise and detailed control procedures are described in **Table S1**.

Study Design

The objective of the research was to investigate the efficiency of NP grafts to restore motor and non-motor symptoms in parkinsonian macaque monkeys. All behavioral and functional measures were acquired by experimenters blinded to the outcome of the grafts integration as this was evaluated post-mortem. Post-mortem quantifications of graft survival, differentiation rate and length of graft's processes were done by investigators blinded to the precise evolution of parkinsonian symptoms of the animals following graft. The experimental findings include replication of the procedure in all individual cases separately and conclusions on the findings arise from difference in outcome based on post-mortem evaluation of grafts integration. Cases followed by PET imaging were chosen randomly before experiment starts as well as cases chosen for sham-grafts. Data collection was stopped at least 7 months following transplantation and after last possible PET-scan acquisition. Data-points of the locomotor activity acquired on the day of anesthesia for MRI, PET-scan or surgery were excluded from circadian analyses. Presented results include all other data-points, no outliers were excluded.

Nine macaque monkeys (Macaca fascicularis, n=9) were used in total for this study. The study was designed such that one third (n=3) were randomly selected as histological controls i.e. never received MPTP injections and, the other two third (n=6) entered the MPTP protocol. Whilst we selected control animals for histological comparison, we designed the study such that each animal would be its own control for clinical, behavioral and functional measures. **Figure 1A** displays a schematic representation of the study design and protocol. Two third of the MPTP group (n=4) were randomly selected to be followed by PET imaging of striatal DA transporters and DA function throughout the full study and, half of them (n=2) were randomly selected to evaluate sham-grafts effects on clinical, behavioral and functional measures. In addition, half of the animals entering the MPTP protocol were randomly selected (n=3) for a two-step MPTP procedure in which MPTP injections were first halted based on clinical score, in order to induce spontaneous recovery [25] for further comparison of this *spontaneous* recovery to the prospective *graft-induced* recovery. MPTP-intoxication was then resumed in order to induce stable motor symptoms [27] over the period preceding cell therapy. Study design is further detailed below.

Six late middle-aged – 11-13 (13-17) years old at protocol onset (end) female macaque monkeys (Macaca fascicularis, 4-5kg) were intoxicated with low-dose 6-methyl-1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridin injections (MPTP, 0.2mg/kg, i.m). Animals were housed in a room dedicated to MPTP experiments, with free access to water and received food twice a day. The neurotoxin was delivered at low-doses chronically each 3-4 days (slowly progressive lesion) during prolonged periods, followed by acute low-doses intoxication (daily injections) in order to ensure stable persistent motor symptoms, as described previously [27, 29]. Cases 1, 3 and 4, were selected to critically compare the evolution of clinical, behavioral and functional markers following spontaneous clinical motor recovery [25, 27] with those obtained during potential recovery induced by cell therapy. In order to do so, we suspended the slowly progressive lesion induced by chronic low-dose MPTP injections as soon as the clinical score reached symptomatic threshold (**Fig.S1** and **Table S2**). In all cases, acute (daily) low-dose MPTP injections induced persistent motor-symptoms. We cautiously stopped daily MPTP injections when the PMRS-motor score was above ten for two consecutive days following one MPTP injection. For results presentation, animals were grouped according to their clinical motor state at the end of the protocol following cell therapy (see **Fig.1**) i.e. 1st group – recovered (n=3, cases 1, 4, 5) and, 2nd group – non-

recovered (n=2, cases 3 and 6). Case2 had per operative brain hemorrhage following NPs transplantation
and was thus excluded from group comparison. Data are presented according to the following periods of
the protocol: 1- CTR (measures acquired before MPTP-onset); 2- MPTP (during MPTP intoxication
period); 3- post-MPTP (following arrest of last MPTP injections, clinical motor score remained stably
above 5 during this period i.e. motor-symptomatic); 4- post-Grafts (following transplantation, described
in **Supplementary Materials and Methods**). Delays between last MPTP injection and cell therapy were
4-23 weeks (n=6, all cases) and, delays between last MPTP injection and sham-grafts were 6-16 weeks
(n=2, cases 1 and 6).

Complete details concerning Parkinsonian Motor Rating Scale – PMRS, Surgical Procedures,
Transplanted cells, Cognitive behavior – detour task, Circadian rhythm follow-up, Dopamine function
imaging – [^{11}C]-PE2I and [^{18}F]-FDOPA, Immuno-histological and Quantification procedures, Statistics
can be found in the **Supplementary Materials and Methods**.

Supplementary Materials

Supplementary Material and Methods.

Fig.S1. Individual follow-up of behavioral and clinical markers.

Fig.S2. Characterization of grafted cells.

Fig.S3. Functional follow-up of striatal DAT binding and Sham-grafts consequences on behavioral and
functional markers.

Fig.S4. Grafted NPs did not survive in the *non-recovered* group.

Fig.S5. Post-mortem localization and characterization of transplanted NPs.

Fig.S6. Grafted NPs in case1.

Fig.S7. Grafted NPs in case4.

Fig.S8. Grafted NPs in case5.

Fig.S9. Hemispheric differences in graft-induced recovery.

Fig.S10. NPs show no overgrowth after transplantation and induce minor immune reaction.

Table S1. Longitudinal protocol design, repeated measures and case controls.

Table S2. Total cumulated MPTP doses per case.

Table S3. Cell survival following transplantation of NPs.

Table S4. Grafted cells number, survival rates, differentiation fates and comparison to literature.

Table S5. Clinical motor, cognitive and functional impact of grafts described in Table S4.

Table S6. Antibodies list.

Table S7. Primers for semi-quantitative RT-PCR.

Table S8. Group-level stats of Fig.1B.

References and Notes

1.   E.R. Dorsey, T. Sherer, M.S. Okun, and B.R. Bloem, The Emerging Evidence of the Parkinson Pandemic.
J.ParkinsonsDis.**8**(s1),S3-S8(2018).
2.   D.J. Brooks, Imaging approaches to Parkinson disease. *J.Nucl.Med*.**51**(4),596-609(2010).
3.   Y. Agid, Parkinson's disease: pathophysiology. *Lancet*.**337**(8753),1321-4(1991).

4.   M.C. Rodriguez-Oroz, M. Jahanshahi, P. Krack, I. Litvan, R. Macias, E. Bezard, and J.A. Obeso, Initial clinical
manifestations of Parkinson's disease: features and pathophysiological mechanisms. *LancetNeurol*.**8**(12),1128-
39(2009).
5.   Y. Leng, E.S. Musiek, K. Hu, F.P. Cappuccio, and K. Yaffe, Association between circadian rhythms and
neurodegenerative diseases. *LancetNeurol*.**18**(3),307-318(2019).
6.   E. Oikonomou and T. Paparrigopoulos, [Neuropsychiatric manifestations in Parkinson's disease].
Psychiatriki.**26**(2),116-30(2015).
7.   B. Bruguerolle and N. Simon, Biologic rhythms and Parkinson's disease: a chronopharmacologic approach to
considering fluctuations in function. *Clin.Neuropharmacol*.**25**(4),194-201(2002).
8.   M.J. Armstrong and M.S. Okun, Diagnosis and Treatment of Parkinson Disease: A Review. *JAMA*.**323**(6),548-
560(2020).
9.   I. Aviles-Olmos, Z. Kefalopoulou, E. Tripoliti, J. Candelario, H. Akram, I. Martinez-Torres, M. Jahanshahi, T.
Foltynie, M. Hariz, L. Zrinzo, and P. Limousin, Long-term outcome of subthalamic nucleus deep brain stimulation
for Parkinson's disease using an MRI-guided and MRI-verified approach.
J.Neurol.Neurosurg.Psychiatry.**85**(12),1419-25(2014).
10.   M.A. Volonte, G. Clarizio, S. Galantucci, P.G. Scamarcia, R. Cardamone, L.R. Barzaghi, M. Falautano, P. Mortini,
G. Comi, and M. Filippi, Long term follow-up in advanced Parkinson's disease treated with DBS of the
subthalamic nucleus. *J.Neurol*.(2021).
11.   A. Bjorklund and O. Lindvall, Replacing Dopamine Neurons in Parkinson's Disease: How did it happen?
J.ParkinsonsDis.**7**(s1),S21-S31(2017).
12.   S.F. Moore, N.V. Guzman, S.L. Mason, C.H. Williams-Gray, and R.A. Barker, Which patients with Parkinson's
disease participate in clinical trials? One centre's experiences with a new cell based therapy trial (TRANSEURO).
J.ParkinsonsDis.**4**(4),671-6(2014).
13.   R.A. Barker, Designing stem-cell-based dopamine cell replacement trials for Parkinson's disease.
Nat.Med.**25**(7),1045-1053(2019).
14.   D.M. Swallow, M.A. Lawton, K.A. Grosset, N. Malek, C.R. Smith, N.P. Bajaj, R.A. Barker, Y. Ben-Shlomo, D.J.
Burn, T. Foltynie, J. Hardy, H.R. Morris, N. Williams, N.W. Wood, and D.G. Grosset, Variation in Recent Onset
Parkinson's Disease: Implications for Prodromal Detection. *J Parkinsons Dis*.**6**(2),289-300(2016).
15.   A. Kirkeby, S. Nolbrant, K. Tiklova, A. Heuer, N. Kee, T. Cardoso, D.R. Ottosson, M.J. Lelos, P. Rifes, S.B.
Dunnett, S. Grealish, T. Perlmann, and M. Parmar, Predictive Markers Guide Differentiation to Improve Graft
Outcome in Clinical Translation of hESC-Based Therapy for Parkinson's Disease. *CellStemCell*.**20**(1),135-
148(2017).
16.   F. Wianny and J. Vezoli, Transplantation in the nonhuman primate MPTP model of Parkinson's disease: update
and perspectives. *PrimateBiol*.**4**(2),185-213(2017).
17.   P. Piccini, N. Pavese, P. Hagell, J. Reimer, A. Bjorklund, W.H. Oertel, N.P. Quinn, D.J. Brooks, and O. Lindvall,
Factors affecting the clinical outcome after neural transplantation in Parkinson's disease. *Brain*.**128**(Pt 12),2977-
86(2005).
18.   N. Breysse, T. Carlsson, C. Winkler, A. Bjorklund, and D. Kirik, The functional impact of the intrastriatal
dopamine neuron grafts in parkinsonian rats is reduced with advancing disease. *J.Neurosci*.**27**(22),5849-56(2007).
19.   I. Garitaonandia, R. Gonzalez, G. Sherman, A. Semechkin, A. Evans, and R. Kern, Novel Approach to Stem Cell
Therapy in Parkinson's Disease. *StemCellsDev*.**27**(14),951-957(2018).
20.   I. Madrazo, O. Kopyov, M.A. Avila-Rodriguez, F. Ostrosky, H. Carrasco, A. Kopyov, A. Avendano-Estrada, F.
Jimenez, E. Magallon, C. Zamorano, G. Gonzalez, T. Valenzuela, R. Carrillo, F. Palma, R. Rivera, R.E. Franco-
Bourland, and G. Guizar-Sahagun, Transplantation of Human Neural Progenitor Cells (NPC) into Putamina of
Parkinsonian Patients: A Case Series Study, Safety and Efficacy Four Years after Surgery.
CellTransplant.**28**(3),269-285(2019).
21.   P.A. Barbuti, R.A. Barker, P. Brundin, S. Przedborski, S.M. Papa, L.V. Kalia, and H. Mochizuki, Recent Advances
in the Development of Stem-Cell-Derived Dopaminergic Neuronal Transplant Therapies for Parkinson's Disease.
Mov.Disord.(2021).
22.   M.E. Emborg, Y. Liu, J. Xi, X. Zhang, Y. Yin, J. Lu, V. Joers, C. Swanson, J.E. Holden, and S.C. Zhang, Induced
pluripotent stem cell-derived neural cells survive and mature in the nonhuman primate brain. *CellRep*.**3**(3),646-
50(2013).
23.   L.H. Thompson and A. Bjorklund, Reconstruction of brain circuitry by neural transplants generated from
pluripotent stem cells. *Neurobiol.Dis*.**79**,28-40(2015).
24.   R. Gonzalez, I. Garitaonandia, M. Poustovoitov, T. Abramihina, C. McEntire, B. Culp, J. Attwood, A. Noskov, T.
Christiansen-Weber, M. Khater, S. Mora-Castilla, C. To, A. Crain, G. Sherman, A. Semechkin, L.C. Laurent, J.D.
Elsworth, J. Sladek, E.Y. Snyder, D.E. Redmond, Jr., and R.A. Kern, Neural Stem Cells Derived From Human

Parthenogenetic Stem Cells Engraft and Promote Recovery in a Nonhuman Primate Model of Parkinsons Disease. *CellTransplant*.**25**(11),1945-1966(2016).

25. J. Vezoli, K. Fifel, V. Leviel, C. Dehay, H. Kennedy, H.M. Cooper, C. Gronfier, and E. Procyk, Early Presymptomatic and Long-Term Changes of Rest Activity Cycles and Cognitive Behavior in a MPTP-Monkey Model of Parkinson's Disease. *PLoSOne*.**6**(8),e23952(2011).

26. K. Fifel, J. Vezoli, K. Dzahini, B. Claustrat, V. Leviel, H. Kennedy, E. Procyk, O. Dkhissi-Benyahya, C. Gronfier, and H.M. Cooper, Alteration of daily and circadian rhythms following dopamine depletion in MPTP treated non-human primates. *PLoSOne*.**9**(1),e86240(2014).

27. J. Vezoli, K. Dzahini, N. Costes, C.R. Wilson, K. Fifel, H.M. Cooper, H. Kennedy, and E. Procyk, Increased DAT binding in the early stage of the dopaminergic lesion: a longitudinal [11C]PE2I binding study in the MPTP-monkey. *Neuroimage*.**102 Pt 2**,249-61(2014).

28. M. Kanazawa, H. Ohba, N. Harada, T. Kakiuchi, S. Muramatsu, and H. Tsukada, Evaluation of 6-11C-Methyl-m-Tyrosine as a PET Probe for Presynaptic Dopaminergic Activity: A Comparison PET Study with beta-11C-l-DOPA and 18F-FDOPA in Parkinson Disease Monkeys. *J.Nucl.Med*.**57**(2),303-8(2016).

29. S. Mounayar, S. Boulet, D. Tande, C. Jan, M. Pessiglione, E.C. Hirsch, J. Feger, M. Savasta, C. Francois, and L. Tremblay, A new model to study compensatory mechanisms in MPTP-treated monkeys exhibiting recovery. *Brain*.**130**(Pt 11),2898-914(2007).

30. Y. Leng, T. Blackwell, P.M. Cawthon, S. Ancoli-Israel, K.L. Stone, and K. Yaffe, Association of Circadian Abnormalities in Older Adults With an Increased Risk of Developing Parkinson Disease. *JAMA Neurol*.(2020).

31. F. Wianny, A. Bernat, C. Huissoud, G. Marcy, S. Markossian, V. Cortay, P. Giroud, V. Leviel, H. Kennedy, P. Savatier, and C. Dehay, Derivation and cloning of a novel rhesus embryonic stem cell line stably expressing tau-green fluorescent protein. *StemCells*.**26**(6),1444-53(2008).

32. F. Wianny, P.Y. Bourillot, and C. Dehay, Embryonic stem cells in non-human primates: An overview of neural differentiation potential. *Differentiation*.**81**(3),142-52(2011).

33. L. Madhavan, V. Ourednik, and J. Ourednik, Neural stem/progenitor cells initiate the formation of cellular networks that provide neuroprotection by growth factor-modulated antioxidant expression. *StemCells*.**26**(1),254-65(2008).

34. L. Qiu, M.C. Liao, A.K. Chen, S. Wei, S. Xie, S. Reuveny, Z.D. Zhou, W. Hunziker, E.K. Tan, S.K.W. Oh, and L. Zeng, Immature Midbrain Dopaminergic Neurons Derived from Floor-Plate Method Improve Cell Transplantation Therapy Efficacy for Parkinson's Disease. *Stem.Cells.Transl.Med*.**6**(9),1803-1814(2017).

35. A. Morizane, J. Takahashi, Y. Takagi, Y. Sasai, and N. Hashimoto, Optimal conditions for in vivo induction of dopaminergic neurons from embryonic stem cells through stromal cell-derived inducing activity. *J.Neurosci.Res*.**69**(6),934-9(2002).

36. J. Blesa, C. Juri, M.A. Garcia-Cabezas, R. Adanez, M.A. Sanchez-Gonzalez, C. Cavada, and J.A. Obeso, Inter-hemispheric asymmetry of nigrostriatal dopaminergic lesion: a possible compensatory mechanism in Parkinson's disease. *Front.Syst.Neurosci*.**5**,92(2011).

37. D.C. German, M. Dubach, S. Askari, S.G. Speciale, and D.M. Bowden, 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced parkinsonian syndrome in Macaca fascicularis: which midbrain dopaminergic neurons are lost? *Neuroscience*.**24**(1),161-74(1988).

38. L. Madhavan, B.F. Daley, C.E. Sortwell, and T.J. Collier, Endogenous neural precursors influence grafted neural stem cells and contribute to neuroprotection in the parkinsonian rat. *Eur.J.Neurosci*.**35**(6),883-95(2012).

39. B. Marchetti, C. Tirolo, F. L'Episcopo, S. Caniglia, N. Testa, J.A. Smith, S. Pluchino, and M.F. Serapide, Parkinson's disease, aging and adult neurogenesis: Wnt/beta-catenin signalling as the key to unlock the mystery of endogenous brain repair. *AgingCell*.**19**(3),e13101(2020).

40. W. Maetzler, I. Liepelt, and D. Berg, Progression of Parkinson's disease in the clinical phase: potential markers. *LancetNeurol*.**8**(12),1158-71(2009).

41. A. Diamond, The development and neural bases of memory functions as indexed by the AB and delayed response tasks in human infants and infant monkeys. *Ann.NYAcad.Sci*.**608**,267-309(1990).

42. A.F. Arnsten, Catecholamine modulation of prefrontal cortical cognitive function. *TrendsCogn.Sci*.**2**(11),436-47(1998).

43. A.M. Thierry, S. Pirot, Y. Gioanni, and J. Glowinski, Dopamine function in the prefrontal cortex. *Adv.Pharmacol*.**42**,717-20(1998).

44. R. Cools, E. Stefanova, R.A. Barker, T.W. Robbins, and A.M. Owen, Dopaminergic modulation of high-level cognition in Parkinson's disease: the role of the prefrontal cortex revealed by PET. *Brain*.**125**(Pt 3),584-94(2002).

45. J.D. Elsworth, C. Leranth, D.E. Redmond, Jr., and R.H. Roth, Loss of asymmetric spine synapses in prefrontal cortex of motor-asymptomatic, dopamine-depleted, cognitively impaired MPTP-treated monkeys. *Int.J.Neuropsychopharmacol*.**16**(4),905-12(2013).

46.   A.F. Arnsten, M. Wang, and C.D. Paspalas, Dopamine's Actions in Primate Prefrontal Cortex: Challenges for
Treating Cognitive Disorders. *Pharmacol.Rev.***67**(3),681-96(2015).
47.   S.N. Haber, Corticostriatal circuitry. *DialoguesClin.Neurosci.***18**(1),7-21(2016).
48.   G.M. Halliday, J.B. Leverenz, J.S. Schneider, and C.H. Adler, The neurobiological basis of cognitive impairment
in Parkinson's disease. *Mov.Disord.***29**(5),634-50(2014).
49.   E.M. Vazey and G. Aston-Jones, The emerging role of norepinephrine in cognitive dysfunctions of Parkinson's
disease. *Front.Behav.Neurosci.***6**,48(2012).
50.   G. Porras, Q. Li, and E. Bezard, Modeling Parkinson's disease in primates: The MPTP model.
*ColdSpringHarb.Perspect.Med.***2**(3),a009308(2012).
51.   Y. Yang, B.S. Tang, and J.F. Guo, Parkinson's Disease and Cognitive Impairment.
*ParkinsonsDis.***2016**,6734678(2016).
52.   B. Ballanger, M. Beaudoin-Gobert, S. Neumane, J. Epinat, E. Metereau, S. Duperrier, E. Broussolle, S. Thobois, F.
Bonnefoi, C. Tourvielle, F. Lavenne, N. Costes, D. Lebars, L. Zimmer, V. Sgambato-Faure, and L. Tremblay,
Imaging Dopamine and Serotonin Systems on MPTP Monkeys: A Longitudinal PET Investigation of
Compensatory Mechanisms. *J.Neurosci.***36**(5),1577-89(2016).
53.   M. Gupta, D.L. Felten, and D.M. Gash, MPTP alters central catecholamine neurons in addition to the nigrostriatal
system. *BrainRes.Bull.***13**(6),737-42(1984).
54.   I. Perez-Otano, M.T. Herrero, C. Oset, M.L. De Ceballos, M.R. Luquin, J.A. Obeso, and J. Del Rio, Extensive loss
of brain dopamine and serotonin induced by chronic administration of MPTP in the marmoset.
*BrainRes.***567**(1),127-32(1991).
55.   J.D. Elsworth, A.Y. Deutch, D.E. Redmond, Jr., J.R. Sladek, Jr., and R.H. Roth, MPTP reduces dopamine and
norepinephrine concentrations in the supplementary motor area and cingulate cortex of the primate.
*Neurosci.Lett.***114**(3),316-22(1990).
56.   D.E. Redmond, Jr., K.B. Bjugstad, Y.D. Teng, V. Ourednik, J. Ourednik, D.R. Wakeman, X.H. Parsons, R.
Gonzalez, B.C. Blanchard, S.U. Kim, Z. Gu, S.A. Lipton, E.A. Markakis, R.H. Roth, J.D. Elsworth, J.R. Sladek,
Jr., R.L. Sidman, and E.Y. Snyder, Behavioral improvement in a primate Parkinson's model is associated with
multiple homeostatic effects of human neural stem cells. *Proc.Natl.Acad.Sci.USA.***104**(29),12175-80(2007).
57.   H. Widner, J. Tetrud, S. Rehncrona, B. Snow, P. Brundin, B. Gustavii, A. Bjorklund, O. Lindvall, and J.W.
Langston, Bilateral fetal mesencephalic grafting in two patients with parkinsonism induced by 1-methyl-4-phenyl-
1,2,3,6-tetrahydropyridine (MPTP). *N.Engl.J.Med.***327**(22),1556-63(1992).
58.   G.V. Sawle, P.M. Bloomfield, A. Bjorklund, D.J. Brooks, P. Brundin, K.L. Leenders, O. Lindvall, C.D. Marsden,
S. Rehncrona, H. Widner, and et al., Transplantation of fetal dopamine neurons in Parkinson's disease: PET
[18F]6-L-fluorodopa studies in two patients with putaminal implants. *Ann.Neurol.***31**(2),166-73(1992).
59.   O. Lindvall, G. Sawle, H. Widner, J.C. Rothwell, A. Bjorklund, D. Brooks, P. Brundin, R. Frackowiak, C.D.
Marsden, P. Odin, and et al., Evidence for long-term survival and function of dopaminergic grafts in progressive
Parkinson's disease. *Ann Neurol.***35**(2),172-80(1994).
60.   P. Piccini, D.J. Brooks, A. Bjorklund, R.N. Gunn, P.M. Grasby, O. Rimoldi, P. Brundin, P. Hagell, S. Rehncrona,
H. Widner, and O. Lindvall, Dopamine release from nigral transplants visualized in vivo in a Parkinson's patient.
*Nat.Neurosci.***2**(12),1137-40(1999).
61.   M. Politis and P. Piccini, Brain imaging after neural transplantation. *Prog.BrainRes.***184**,193-203(2010).
62.   Z. Kefalopoulou, M. Politis, P. Piccini, N. Mencacci, K. Bhatia, M. Jahanshahi, H. Widner, S. Rehncrona, P.
Brundin, A. Bjorklund, O. Lindvall, P. Limousin, N. Quinn, and T. Foltynie, Long-term clinical outcome of fetal
cell transplantation for Parkinson disease: two case reports. *JAMANeurol.***71**(1),83-7(2014).
63.   J.M. Monti and D. Monti, The involvement of dopamine in the modulation of sleep and waking.
*SleepMed.Rev.***11**(2),113-33(2007).
64.   L.S. Forno, J.W. Langston, L.E. DeLanney, I. Irwin, and G.A. Ricaurte, Locus ceruleus lesions and eosinophilic
inclusions in MPTP-treated monkeys. *Ann.Neurol.***20**(4),449-55(1986).
65.   R. Sandyk, R.P. Iacono, and S.R. Kay, The hypothalamus in MPTP-induced parkinsonism.
*Ital.J.Neurol.Sci.***11**(4),367-72(1990).
66.   K. Fifel and A. Videnovic, Circadian alterations in patients with neurodegenerative diseases: Neuropathological
basis of underlying network mechanisms. *Neurobiol.Dis.***144**,105029(2020).
67.   J.Y. Li, E. Englund, J.L. Holton, D. Soulet, P. Hagell, A.J. Lees, T. Lashley, N.P. Quinn, S. Rehncrona, A.
Bjorklund, H. Widner, T. Revesz, O. Lindvall, and P. Brundin, Lewy bodies in grafted neurons in subjects with
Parkinson's disease suggest host-to-graft disease propagation. *Nat.Med.***14**(5),501-3(2008).
68.   J.H. Kordower, Y. Chu, R.A. Hauser, T.B. Freeman, and C.W. Olanow, Lewy body-like pathology in long-term
embryonic nigral transplants in Parkinson's disease. *Nat.Med.***14**(5),504-6(2008).
69.   O. Lindvall, S. Rehncrona, B. Gustavii, P. Brundin, B. Astedt, H. Widner, T. Lindholm, A. Bjorklund, K.L.
Leenders, J.C. Rothwell, R. Frackowiak, C.D. Marsden, B. Johnels, G. Steg, R. Freedman, B.J. Hoffer, L. Seiger, I.

669		Stromberg, M. Bygdeman, and L. Olson, Fetal dopamine-rich mesencephalic grafts in Parkinson's disease.
670		*Lancet*.**2**(8626-8627),1483-4(1988).
671	70.	W. Li, E. Englund, H. Widner, B. Mattsson, D. van Westen, J. Latt, S. Rehncrona, P. Brundin, A. Bjorklund, O.
672		Lindvall, and J.Y. Li, Extensive graft-derived dopaminergic innervation is maintained 24 years after
673		transplantation in the degenerating parkinsonian brain. *Proc.Natl.Acad.Sci.USA*.**113**(23),6544-9(2016).
674	71.	M. Parmar, S. Grealish, and C. Henchcliffe, The future of stem cell therapies for Parkinson disease.
675		*Nat.Rev.Neurosci*.**21**(2),103-115(2020).
676	72.	A. Morizane, D. Doi, T. Kikuchi, K. Okita, A. Hotta, T. Kawasaki, T. Hayashi, H. Onoe, T. Shiina, S. Yamanaka,
677		and J. Takahashi, Direct comparison of autologous and allogeneic transplantation of iPSC-derived neural cells in
678		the brain of a non-human primate. *StemCellReports*.**1**(4),283-92(2013).
679	73.	Y. Tao, S.C. Vermilyea, M. Zammit, J. Lu, M. Olsen, J.M. Metzger, L. Yao, Y. Chen, S. Phillips, J.E. Holden, V.
680		Bondarenko, W.F. Block, T.E. Barnhart, N. Schultz-Darken, K. Brunner, H. Simmons, B.T. Christian, M.E.
681		Emborg, and S.C. Zhang, Autologous transplant therapy alleviates motor and depressive behaviors in parkinsonian
682		monkeys. *Nat.Med*.(2021).
683	74.	A.S. Mitchell, A. Thiele, C.I. Petkov, A. Roberts, T.W. Robbins, W. Schultz, and R. Lemon, Continued need for
684		non-human primate neuroscience research. *Curr.Biol*.**28**(20),R1186-R1187(2018).
685	75.	A. Benazzouz, T. Boraud, P. Dubedat, A. Boireau, J.M. Stutzmann, and C. Gross, Riluzole prevents MPTP-
686		induced parkinsonism in the rhesus monkey: a pilot study. *Eur.J.Pharmacol*.**284**(3),299-307(1995).
687	76.	P. Hantraye, M. Varastet, M. Peschanski, D. Riche, P. Cesaro, J.C. Willer, and M. Maziere, Stable parkinsonian
688		syndrome and uneven loss of striatal dopamine fibres following chronic MPTP administration in baboons.
689		*Neuroscience*.**53**(1),169-78(1993).
690	77.	M.E. Emborg, J.W. Tetrud, J. Moirano, W.W. McLaughlin, and K.S. Bankiewicz, Rest tremor in rhesus monkeys
691		with MPTP-induced parkinsonism. *Front Biosci*.**8**,a148-54(2003).
692	78.	H. Bergman, A. Raz, A. Feingold, A. Nini, I. Nelken, D. Hansel, H. Ben-Pazi, and A. Reches, Physiology of
693		MPTP tremor. *Mov.Disord*.**13 Suppl 3**,29-34(1998).
694	79.	G.R. Choudhury and M.M. Daadi, Charting the onset of Parkinson-like motor and non-motor symptoms in
695		nonhuman primate model of Parkinson's disease. *PLoSOne*.**13**(8),e0202770(2018).
696	80.	L. Conti, S.M. Pollard, T. Gorba, E. Reitano, M. Toselli, G. Biella, Y. Sun, S. Sanzone, Q.L. Ying, E. Cattaneo,
697		and A. Smith, Niche-independent symmetrical self-renewal of a mammalian tissue stem cell.
698		*PLoSBiol*.**3**(9),e283(2005).
699	81.	A.L. Perrier, V. Tabar, T. Barberi, M.E. Rubio, J. Bruses, N. Topf, N.L. Harrison, and L. Studer, Derivation of
700		midbrain dopamine neurons from human embryonic stem cells. *Proc.Natl.Acad.Sci.USA*.**101**(34),12543-8(2004).
701	82.	R. Araki, M. Uda, Y. Hoki, M. Sunayama, M. Nakamura, S. Ando, M. Sugiura, H. Ideno, A. Shimada, A. Nifuji,
702		and M. Abe, Negligible immunogenicity of terminally differentiated cells derived from induced pluripotent or
703		embryonic stem cells. *Nature*.**494**(7435),100-4(2013).
704	83.	Y.K. Wang, W.W. Zhu, M.H. Wu, Y.H. Wu, Z.X. Liu, L.M. Liang, C. Sheng, J. Hao, L. Wang, W. Li, Q. Zhou,
705		and B.Y. Hu, Human Clinical-Grade Parthenogenetic ESC-Derived Dopaminergic Neurons Recover Locomotive
706		Defects of Nonhuman Primate Models of Parkinson's Disease. *StemCellReports*.**11**(1),171-182(2018).
707	84.	G.G. Doxiadis, A.J. Rouweler, N.G. de Groot, A. Louwerse, N. Otting, E.J. Verschoor, and R.E. Bontrop,
708		Extensive sharing of MHC class II alleles between rhesus and cynomolgus macaques. *Immunogenetics*.**58**(4),259-
709		68(2006).
710	85.	S. Bunlungsup, S. Kanthaswamy, R.F. Oldt, D.G. Smith, P. Houghton, Y. Hamada, and S. Malaivijitnond, Genetic
711		analysis of samples from wild populations opens new perspectives on hybridization between long-tailed (Macaca
712		fascicularis) and rhesus macaques (Macaca mulatta). *Am.J.Primatol*.**79**(12)(2017).
713	86.	A. Morizane, T. Kikuchi, T. Hayashi, H. Mizuma, S. Takara, H. Doi, A. Mawatari, M.F. Glasser, T. Shiina, H.
714		Ishigaki, Y. Itoh, K. Okita, E. Yamasaki, D. Doi, H. Onoe, K. Ogasawara, S. Yamanaka, and J. Takahashi, MHC
715		matching improves engraftment of iPSC-derived neurons in non-human primates. *Nat.Commun*.**8**(1),385(2017).
716	87.	A. Diamond, Evidence for the importance of dopamine for prefrontal cortex functions early in life.
717		*Philos.Trans.R.Soc.Lond.B.Biol.Sci*.**351**(1346),1483-93; discussion 1494(1996).
718	88.	J.S. Schneider, M. Van Velson, F. Menzaghi, and G.K. Lloyd, Effects of the nicotinic acetylcholine receptor
719		agonist SIB-1508Y on object retrieval performance in MPTP-treated monkeys: comparison with levodopa
720		treatment. *Ann.Neurol*.**43**(3),311-7(1998).
721	89.	J.R. Taylor, J.D. Elsworth, R.H. Roth, T.J. Collier, J.R. Sladek, Jr., and D.E. Redmond, Jr., Improvements in
722		MPTP-induced object retrieval deficits and behavioral deficits after fetal nigral grafting in monkeys.
723		*Prog.BrainRes*.**82**,543-59(1990).
724	90.	J.R. Taylor, J.D. Elsworth, R.H. Roth, J.R. Sladek, Jr., and D.E. Redmond, Jr., Cognitive and motor deficits in the
725		acquisition of an object retrieval/detour task in MPTP-treated monkeys. *Brain*.**113 (Pt 3)**,617-37(1990).

726 91.   O. Dkhissi-Benyahya, C. Gronfier, W. De Vanssay, F. Flamant, and H.M. Cooper, Modeling the role of mid-
727   wavelength cones in circadian responses to light. *Neuron*.**53**(5),677-87(2007).

728 92.   P. Emond, L. Garreau, S. Chalon, M. Boazi, M. Caillet, J. Bricard, Y. Frangin, L. Mauclaire, J.C. Besnard, and D.
729   Guilloteau, Synthesis and ligand binding of nortropane derivatives: N-substituted 2beta-carbomethoxy-3beta-(4'-
730   iodophenyl)nortropane and N-(3-iodoprop-(2E)-enyl)-2beta-carbomethoxy-3beta-(3',4'-disubstituted
731   phenyl)nortropane. New high-affinity and selective compounds for the dopamine transporter.
732   *J.Med.Chem*.**40**(9),1366-72(1997).

733 93.   T. Poyot, F. Conde, M.C. Gregoire, V. Frouin, C. Coulon, C. Fuseau, F. Hinnen, F. Dolle, P. Hantraye, and M.
734   Bottlaender, Anatomic and biochemical correlates of the dopamine transporter ligand 11C-PE2I in normal and
735   parkinsonian primates: comparison with 6-[18F]fluoro-L-dopa. *J Cereb Blood Flow Metab*.**21**(7),782-92(2001).

736 94.   M.C. Shih, M.Q. Hoexter, L.A. Andrade, and R.A. Bressan, Parkinson's disease and dopamine transporter
737   neuroimaging: a critical review. *Sao.Paulo.Med.J*.**124**(3),168-75(2006).

738 95.   K. Dzahini, C. Dentresangle, M. Le Cavorsin, A. Bertrand, I. Detraz, M. Savasta, and V. Leviel, Pre-synaptic
739   glutamate-induced activation of DA release in the striatum after partial nigral lesion. *J.Neurochem*.**113**(6),1459-
740   70(2010).

741 96.   K.B. Bjugstad, Y.D. Teng, D.E. Redmond, Jr., J.D. Elsworth, R.H. Roth, S.K. Cornelius, E.Y. Snyder, and J.R.
742   Sladek, Jr., Human neural stem cells migrate along the nigrostriatal pathway in a primate model of Parkinson's
743   disease. *Exp.Neurol*.**211**(2),362-9(2008).

744 97.   D.R. Wakeman, S. Weiss, J.R. Sladek, J.D. Elsworth, B. Bauereis, C. Leranth, P.J. Hurley, R.H. Roth, and D.E.
745   Redmond, Survival and integration of neurons derived from human embryonic stem cells in MPTP-lesioned
746   primates. *CellTransplant*.**23**(8),981-94(2014).

747 98.   R. Gonzalez, I. Garitaonandia, A. Crain, M. Poustovoitov, T. Abramihina, A. Noskov, C. Jiang, R. Morey, L.C.
748   Laurent, J.D. Elsworth, E.Y. Snyder, D.E. Redmond, Jr., and R. Semechkin, Proof of concept studies exploring the
749   safety and functional activity of human parthenogenetic-derived neural stem cells for the treatment of Parkinson's
750   disease. *CellTransplant*.**24**(4),681-90(2015).

751 99.   Q. Xu, X. Jiang, Y. Ke, S. Zhang, R. Xu, and Y. Zeng, Gene therapy in hemiparkinsonian rhesus monkeys: long-
752   term survival and behavioral recovery by transplantation of autologous human tyrosine hydroxylase-expressing
753   neural stem cells. *Cytotherapy*.**12**(2),226-37(2010).

754 100.   Y. Zhou, M. Sun, H. Li, M. Yan, Z. He, W. Wang, and S. Lu, Recovery of behavioral symptoms in hemi-
755   parkinsonian rhesus monkeys through combined gene and stem cell therapy. *Cytotherapy*.**15**(4),467-80(2013).

756 101.   S. Kriks, J.W. Shim, J. Piao, Y.M. Ganat, D.R. Wakeman, Z. Xie, L. Carrillo-Reid, G. Auyeung, C. Antonacci, A.
757   Buch, L. Yang, M.F. Beal, D.J. Surmeier, J.H. Kordower, V. Tabar, and L. Studer, Dopamine neurons derived
758   from human ES cells efficiently engraft in animal models of Parkinson's disease. *Nature*.**480**(7378),547-51(2011).

759 102.   D. Doi, A. Morizane, T. Kikuchi, H. Onoe, T. Hayashi, T. Kawasaki, M. Motono, Y. Sasai, H. Saiki, M. Gomi, T.
760   Yoshikawa, H. Hayashi, M. Shinoyama, M.M. Refaat, H. Suemori, S. Miyamoto, and J. Takahashi, Prolonged
761   maturation culture favors a reduction in the tumorigenicity and the dopaminergic function of human ESC-derived
762   neural cells in a primate model of Parkinson's disease. *StemCells*.**30**(5),935-45(2012).

763 103.   M.M. Daadi, B.A. Grueter, R.C. Malenka, D.E. Redmond, Jr., and G.K. Steinberg, Dopaminergic neurons from
764   midbrain-specified human embryonic stem cell-derived neural stem cells engrafted in a monkey model of
765   Parkinson's disease. *PLoSONE*.**7**(7),e41120(2012).

766 104.   R. Sanchez-Pernaute, L. Studer, D. Ferrari, A. Perrier, H. Lee, A. Vinuela, and O. Isacson, Long-term survival of
767   dopamine neurons derived from parthenogenetic primate embryonic stem cells (cyno-1) after transplantation.
768   *StemCells*.**23**(7),914-22(2005).

769 105.   Y. Takagi, J. Takahashi, H. Saiki, A. Morizane, T. Hayashi, Y. Kishi, H. Fukuda, Y. Okamoto, M. Koyanagi, M.
770   Ideguchi, H. Hayashi, T. Imazato, H. Kawasaki, H. Suemori, S. Omachi, H. Iida, N. Itoh, N. Nakatsuji, Y. Sasai,
771   and N. Hashimoto, Dopaminergic neurons generated from monkey embryonic stem cells function in a Parkinson
772   primate model. *J.Clin.Invest*.**115**(1),102-9(2005).

773 106.   M. Sundberg, H. Bogetofte, T. Lawson, J. Jansson, G. Smith, A. Astradsson, M. Moore, T. Osborn, O. Cooper, R.
774   Spealman, P. Hallett, and O. Isacson, Improved cell therapy protocols for Parkinson's disease based on
775   differentiation efficiency and safety of hESC-, hiPSC-, and non-human primate iPSC-derived dopaminergic
776   neurons. *StemCells*.**31**(8),1548-62(2013).

777 107.   P.J. Hallett, M. Deleidi, A. Astradsson, G.A. Smith, O. Cooper, T.M. Osborn, M. Sundberg, M.A. Moore, E.
778   Perez-Torres, A.L. Brownell, J.M. Schumacher, R.D. Spealman, and O. Isacson, Successful function of autologous
779   iPSC-derived dopamine neurons following transplantation in a non-human primate model of Parkinson's disease.
780   *CellStemCell*.**16**(3),269-74(2015).

781 108.   S. Wang, C. Zou, L. Fu, B. Wang, J. An, G. Song, J. Wu, X. Tang, M. Li, J. Zhang, F. Yue, C. Zheng, P. Chan,
782   Y.A. Zhang, and Z. Chen, Autologous iPSC-derived dopamine neuron transplantation in a nonhuman primate
783   Parkinson's disease model. *Cell.Discov*.**1**,15012(2015).

784   109.   T. Kikuchi, A. Morizane, D. Doi, H. Onoe, T. Hayashi, T. Kawasaki, H. Saiki, S. Miyamoto, and J. Takahashi,
785   Survival of human induced pluripotent stem cell-derived midbrain dopaminergic neurons in the brain of a primate
786   model of Parkinson's disease. *J.ParkinsonsDis*.**1**(4),395-412(2011).

787   110.   S. Peng, Y. Ma, J. Flores, M. Cornfeldt, B. Mitrovic, D. Eidelberg, and D.J. Doudet, Modulation of Abnormal
788   Metabolic Brain Networks by Experimental Therapies in a Nonhuman Primate Model of Parkinson Disease: An
789   Application to Human Retinal Pigment Epithelial Cell Implantation. *J.Nucl.Med*.**57**(10),1591-1598(2016).

790   111.   E.F. Wolff, L. Mutlu, E.E. Massasa, J.D. Elsworth, D. Eugene Redmond, Jr., and H.S. Taylor, Endometrial stem
791   cell transplantation in MPTP- exposed primates: an alternative cell source for treatment of Parkinson's disease.
792   *J.Cell.Mol.Med*.**19**(1),249-56(2015).

793   112.   Z. Ren, J. Wang, S. Wang, C. Zou, X. Li, Y. Guan, Z. Chen, and Y.A. Zhang, Autologous transplantation of
794   GDNF-expressing mesenchymal stem cells protects against MPTP-induced damage in cynomolgus monkeys.
795   *Sci.Rep*.**3**,2786(2013).

796   113.   M.R. Luquin, M. Manrique, J. Guillen, J. Arbizu, C. Ordonez, and I. Marcilla, Enhanced GDNF expression in
797   dopaminergic cells of monkeys grafted with carotid body cell aggregates. *Brain.Res*.**1375**,120-7(2011).

798   114.   J. Bloch, J.F. Brunet, C.R. McEntire, and D.E. Redmond, Primate adult brain cell autotransplantation produces
799   behavioral and biological recovery in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced parkinsonian St. Kitts
800   monkeys. *J.Comp.Neurol*.**522**(12),2729-40(2014).

Acknowledgments: We acknowledge Dr. Vincent Leviel for discussions at different stages of the study. We thank Frank Lavenne for PET acquisition; Didier Le Bars and Frédéric Bonnefoi for radioligand synthesis; Mélissa Remy, Léo Pion and Luc Grinand for histological graft analyses; Claude Gronfier for advice during circadian analyses; Pascale Giroud for precious help during anesthesia, surgery and perfusion.

Funding: The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Région Rhône-Alpes (C.D, J.V)

Fondation de France (E.P, H.M.C, J.V)

Fondation Caisse d'Epargne Rhône-Alpes Lyon (C.D, J.V)

Rhône-Alpes cible 11-010869 (H.K)

Fondation Neurodis (C.R.E.W)

Cluster Handicap Vieillissement Neurosciences Rhône-Alpes and LabEX CORTEX : Agence Nationale de la Recherche ANR-11 LABX-0042 (H.K, E.P), P6-2005 IST-1583 (H.K), FP7-2007 ICT-216593 (H.K)

Agence Nationale de la Recherche ANR-11-BSV4-501 (H.K).

Author contributions:

Conceptualization: HMC, EP, HK, PS, CD, JV

Methodology: KK, HMC, EP, HK, PS, CD, JV

Investigation: FW, KD, KF, CREW, AB, VD, PM, CL, JV

Visualization: FW, JV

Funding acquisition: CREW, HMC, EP, HK, CD, JV

Supervision: CD, JV

Writing – original draft: FW, HK, JV

Writing – review & editing: FW, KF, CREW, KK, HMC, EP, HK, PS, CD, JV

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: The data, code, and materials that support the findings of this study are available from the corresponding author upon reasonable request.

Additional Information: Correspondence and requests for materials should be addressed to JV.