Main Manuscript for

Molecular insights into substrate recognition and discrimination by the N-

terminal domain of Lon AAA+ protease

Authors: Shiou-Ru Tzeng[1*], Yin-Chu Tseng[1], Chien-Chu Lin[2,5], Chia-Ying Hsu[1], Shing-Jong Huang[3], Yi-

Ting Kuo[1], Chung-I Chang[2,4*]

Affiliations:

1. Institute of Biochemistry and Molecular Biology, College of Medicine, National Taiwan University,

Taipei, Taiwan.

2. Institute of Biological Chemistry, Academia Sinica, Taipei 11529

3. Instrumentation Center, National Taiwan University

4. Institute of Biochemical Sciences, College of Life Science, National Taiwan University, Taipei, 10617

Taiwan

5. Present Address: Institute of Molecular Biology, Academia Sinica, Taipei 11529, Taiwan

*__Corresponding authors__

Shiou-Ru Tzeng: R907, No. 1 Jen- Ai Road Section 1, Taipei 10051, Taiwan.; TEL: (+886)-22312-3456

ext 88205; FAX: (+886)-2239-15295; E-mail: srtzeng@ntu.edu.tw

Chung-I Chang: 128, Section 2, Academia Road, Nankang 11529, Taipei, Taiwan.; TEL: (+886)-22785-

5696 ext 7110; E-mail: chungi@gate.sinica.edu.tw

Keywords

Lon AAA+ protease; N-terminal domain; degrons; misfolded proteins; protein aggregates; *Meiothermus*

taiwanensis.

This PDF file includes:

Main Text

Table 1

Figures 1 to 7

Abstract

The Lon AAA+ proteases (LonA) is a ubiquitous ATP-dependent proteolytic machine, which selectively

degrades damaged proteins or native proteins carrying exposed motifs (degrons). Here we characterize

the structural basis for substrate recognition and discrimination by the N-terminal domain (NTD) of LonA.

The results reveal that the six NTDs are attached to the hexameric LonA chamber by flexible linkers such

that the formers tumble independently of the latter. Further spectral analyses show that the NTD

selectively interacts with unfolded proteins, protein aggregates, and degron-tagged proteins by two

hydrophobic patches of its N-lobe, but not intrinsically disordered substrate, α-casein. Moreover, the NTD

selectively binds to protein substrates when they are thermally induced to adopt unfolded conformations.

Collectively, our findings demonstrate that NTDs enable LonA to perform protein quality control to

selectively degrade proteins in damaged states and suggest that substrate discrimination and selective

degradation by LonA are mediated by multiple NTD interactions.

Impact Statement

The N-terminal domains enable Lon protease to discriminate and capture selected protein species for

degradation by exposed hydrophobic patches and flexible linkages to the hexameric core complex.

Introduction:

The Lon AAA+ protease (LonA), previously known as the protease La, is an ATP-dependent

protease distributed in prokaryote and eukaryote[1]. It forms a homo-hexamer to execute its biological

function[2,3]. LonA belongs to the AAA+ (ATPases associated with various cellular activities) superfamily

and contains an N-terminal domain (NTD), a middle ATPase domain with conserved Walker motifs for

ATP hydrolysis, and a C-terminal protease domain (CTD) with a serine-lysine catalytic dyad in the active

site[4,5]. LonA is responsible for degrading damaged or unfolded proteins, as well as native proteins

bearing specific recognition elements, known as degradation tag or degrons[6,7]. How can LonA

discriminate its substrate from other non-substrate proteins in a cell? Substrate recognition of LonA is

thought to be mediated by its NTD[8-14]. The substrates are unfolded and translocated in an ATP-

dependent process into a secluded chamber formed by the ATPase domains and the protease domains.

Finally, the unfolded substrates inside the chamber are degraded into small peptide fragments[15,16]. The

ATP-dependent translocation process of substrates into the hexameric chamber, formed by the AAA+

and protease domains, has been well characterized[17,18]. Structures of different N-terminal fragments of

LonA from three species have been reported[19-22], including *Escherichia coli* (EcLon), *Mycobacterium*

avium complex (MacLon) and *Bacillus subtilis* (BsLon). However, no structural study has been conducted

to analyze substrate interaction by the NTD either in isolation or in the context of full-length LonA.

Here we address the question of substrate recognition and discrimination by the NTD domains of

LonA using nuclear magnetic resonance (NMR) spectroscopy. To understand how LonA selectively

recognizes protein substrates being in damaged states, we have used a thermal stable LonA isolated

from *Meiothermus taiwanensis* (termed MtaLonA henceforth), which allows temperature cycling

experiments to be applied to change the populations of a protein substrate in folded and unfolded states.

In this work, five various protein substrates have been employed for NMR experiments, which include: (i)

Ig2 (domains 5 and 6 of the gelation factor from *Dictyostelium discoideum*[23], hereafter abbreviated as Ig2)

with an immunoglobulin (Ig) fold forming a dimer that is natively folded up to 40 °C; (ii) α-casein, which is

an intrinsically disordered protein with no tertiary structure; (iii) native lysozyme with four disulfide

bridges and reduced lysozyme forming loose and flexible aggregates; (iv) Ig2D5 (domain 5 of the

gelation factor from *Dictyostelium discoideum*), which is used for characterization of substrate

conformation selection by NTD; and (v) a degron-tagged Ig2D5 designed to identify the residues of

degrons that are recognized and bound by NTD.

Here we show that the presence of NTD is required to degrade damaged proteins and native

proteins with degrons, but it does not play an active role in mediating the degradation by LonA of an

intrinsically disordered substrate, α-casein. We have also determined the crystal structure of an N-

terminal fragment of MtaLonA, and refined it to 2.1 Å resolution. The structure is homologous to the

structures of N-terminal fragment of both EcLon and MacLon[19-22] , but not to that of BsLon[20]. With the aid

of perdeuteration labeling and transverse relaxation optimized NMR spectroscopy (TROSY), we find that

the NTD appears to tumble independently of the hexameric core complex, suggesting that the NTD is

attached to the hexameric chamber by a flexible linker thus rendering it possible for detailed chemical

shift perturbation mapping of substrate binding in the context of full-length MtaLonA. We further

structurally characterize the NTD-mediated interactions with unfolded proteins, protein aggregates, or

degron-tagged proteins to elucidate its functional role in substrate discrimination. Our results demonstrate

that the substrate-binding site of NTD is located at hydrophobic patches of its N-lobe, which recognizes

exposed aromatic and hydrophobic residues of damaged proteins and degrons. Using temperature

cycling experiments, we investigate how the conformations of substrates are affected in the presence of

MtaLonA NTDs, which interact with the initial substrate before subsequent translocation and degradation.

Altogether, this work elucidates the substrate recognition mediated by the NTDs of MtaLonA at

substantially high resolution and suggests that the flexibly linked NTDs can help survey, discriminate, and

selectively capture damaged unfolded protein species or native protein substrates carrying exposed

degradation sequence motifs.

Results

The NTD is essential for efficient degradation of a thermally unfolded substrate but not

intrinsically disordered protein substrate, *α-casein.*

To evaluate the importance of the NTD in the proteolysis activity of MtaLonA, we constructed a

MtaLonA variant, AAAP, of which the NTD (residues 1-241) was removed (Fig. 1A), based on the

structure of *B. subtilis* LonA. Substrate degradation activity of full-length MtaLonA and AAAP were

evaluated using two protein substrates, namely α-casein and Ig2. Native α-casein is an intrinsically

disordered substrate of LonA. Ig2 is an all β-stranded protein that becomes partially unfolded at 55 °C

(supplementary Fig. 1A) and only then is susceptible to proteolysis by MtaLonA[6]. Full-length MtaLonA

degrades both α-casein as well as Ig2[24]. Compared to the full-length MtaLonA, AAAP exhibits a slightly

reduced degradation activity against α-casein (Fig. 1B) while its proteolytic activity for Ig2 is severely

impaired (Fig. 1C). These results showed that, despite the lack of the NTD, AAAP retains the proteolytic

activity for intrinsically disordered protein substrates like α-casein. However, the NTD is essential for

MtaLonA to recognize and degrade the thermally damaged substrate, Ig2, indicating the NTD may

mediate specific interactions with damaged Ig2 but not α-casein.

The NTDs tumble independently of the hexameric LonA core in solution

We obtained crystals of the N-terminal fragment NN206 of MtaLonA by *in situ* proteolysis during

crystallization (Fig. 2A). The structure was refined to 2.1 Å resolution with a free R factor of 24.9%

(crystallographic R factor is 20.9%) (Table 1). NN206 forms a bilobal fold: (i) the N-terminal lobe, which

consist of one α-helix and six β-strands, forming three β-sheets (β1/β3/β4/β5, β2/β5/β4 and β1/β6/β5); (ii)

the C-terminal lobe, which is is exclusively α-helical, consisting of a five-helix bundle (Figs. 2B). The two

lobes are jointed together by an 18-residue linker, which is similar to that of both EcLon and MacLon[19-22].

Based on the result, we also crystallized and solved the structure of the longer N-terminal fragment

NN291 (Fig. 2A). The structure of NN291 was refined to 1.7 Å resolution (Table 1), which is similar to

NN206 except for a longer C-terminal helix by 4 more residues, i.e., residues 207-210 (Fig. 2C). Despite

the lack of spontaneous proteolytic fragmentation in the crystals, no electron density was seen past

residue A210 (Fig. 2D).

Interestingly, the C-terminus of *E. coli* N-terminal fragment forms an extended 40-residue helix[21].

Therefore, we further investigate the structural feature of the corresponding residues 207-243 of MtaLonA

using solution-state NMR spectroscopy. Most of the well-dispersed cross-peaks in the two-dimensional

^{15}N-^{1}H HSQC NMR spectra of NN206 and NN243 can be superimposed, indicating that the well-folded

domain structure corresponding to residues 1-206 is not perturbed by the C-terminal extension, residues

207-243 (Fig. 2E). Nonetheless, NN243 exhibits extra intense and poorly dispersed cross-peaks around

7.5-8.5 ppm along the proton dimension that is indicative of a highly disordered peptide segment, which

most likely corresponds to the C-terminal extension of NN243. Together, our crystallographic and NMR

analyses suggest that the N-terminal domain of MtaLonA forms a bilobal globular structure (residues 1-

206) followed by a C-terminal flexible extension (residues 211-243).

LonA forms a large homo-hexameric complex resulting in a very high molecular weight of 0.5

MDa. To verify the dynamic feature of the NTD, we compared the backbone amide ^{15}N-^{1}H NMR

correlations of NN206 and those of full-length MtaLonA at 55 °C. With the aid of perdeuteration labeling

and transverse relaxation optimized NMR spectroscopy (TROSY), we could observe over 200 well-

resolved ^{15}N-^{1}H correlations of full-length hexameric MtaLonA at an apparent molecular weight of 0.5

MDa. Importantly, most of these correlations are superimposable to those observed in the NN206

spectrum (Fig. 2F). These results indicate that the NTD adopts the same structure as the isolated NTD

while loosely tethered to the near-megadalton hexameric LonA core via a long flexible linker thus leading

to an independent, fast tumbling motion to yield highly favorable NMR relaxation properties and thus high-

quality multidimensional NMR spectra. We further applied multidimensional heteronuclear NMR

experiments to assign the observed cross peaks in the ^{1}H–^{15}N TROSY HSQC spectrum of NN206

(supplementary Fig. 2). Together, these results suggest that in a LonA complex, the six N-terminal

domains may each connect to the hexameric core of fused AAA+ and protease domains by flexible ~40-

residue linkers. Therefore, each of the six 24-kDa N-terminal domains may tumble independently of the

350-kDa hexameric core (Fig. 2G).

The chemical shifts of NTDs are perturbed by thermally unfolded proteins, aggregates, and

degron tags

To directly observe discrimination of native or thermally damaged substrate mediated by NTD, we

treated NMR samples by a thermal cycle with temperature increasing from 32 °C to 55 °C and then

returning to 32 °C. With temperature rising, the exposed hydrophobic areas of thermally unfolded Ig2 is

increased, as detected by SYPRO Orange (supplementary Fig. 1A), indicating that the population of Ig2

in unfolded states is increased by thermal denaturation. By contrast, thermal cycling of isolated NTDs

exhibited no temperature effect on protein structure monitored by NMR spectroscopy, which may be

explained by the high melting temperature (T_m) of full-length MtaLonA and NN206 (68.0 °C and 85.5 °C,

respectively; supplementary Figs. 1A and 1B). Upon addition of a 2-fold molar excess of well-folded Ig2 at

32 °C, no significant chemical shift perturbation (CSP) is observed (Fig. 3A), where only the NTD is

isotopically enriched. By increasing temperature from 32 °C to 55 °C for the same NMR sample, a

dramatic variation in the 2D spectral features is detected and the resonances undergo a prominent

decrease in intensity (Fig. 3B), indicating that the NTD interacts with thermally unfolded Ig2. With

temperature dropping from 55 °C to 32 °C, line width of the expected bound-state resonances is too

broad to be detected in the presence of binding events (Fig. 3C), suggesting a certain portion of Ig2 still

stays in the unfolded or aggregated state. In contrast, the addition of α-casein results in little to no effect

on the cross-peaks of residues 1-206 of the NTD (supplementary Fig. 3) thus precluding specific

interaction mode between the intrinsically disordered protein α-casein with the NTD of MtaLonA.

To clarify whether overall broadening of the cross-peak signals is due to protein aggregation or

precipitation, we performed size-exclusion chromatography (SEC) experiments to examine the NMR

sample treated with thermal cycling. The analysis revealed that more than half of Ig2 protein becomes

unfolded and aggregated through a thermal cycle while the NTD is still well-folded (Fig. 3D). The NTD-Ig2

aggregate complex showed a dissociable SEC profile, suggesting the NTD-substrate interaction is quite

dynamic by nature and the fast substrate dissociation may facilitate the translocation process.

NMR analysis indicated that thermally damaged Ig2 induced significant CSPs and broadened

resonances in the NTD N-lobe, which comprise one α-helix and three β-sheets β1/β3/β4/β5, β2/β5/β4

and β1/β6/β5. The residues of NTD with the pronounced CSPs are part of the interaction interface

located mainly at helix α1, loop L1, L2, L3 and β-sheet β2/β5/β4 (Figs. 3E and 3F), which consist primarily

of hydrophobic residues and are decorated by a number of polar residues (supplementary Fig. 4A). To

characterize the binding sites of the NTD interaction with damaged Ig2, we chose to mutate two exposed

hydrophobic residues Pro-22 and Met-85 located at the center of β-sheet β2/β5/β4, which form a

contiguous hydrophobic surface in the N-lobe subdomain (Fig. 3G and supplementary Fig. 4B). We found

that NN206*, containing two specific mutations in the NTD (P22A and M85A; hereafter NN206*), may

influence the hydrophobic substrate-binding interaction. The NMR sample containing both NN206* and

Ig2 protein was also treated with one thermal cycle. NN206* showed much weak interactions with

thermal damaged Ig2 (Figs. 3H-J), confirming that this binding event is mainly contributed by the specific

hydrophobic residues with high hydrophobicity in a continuous hydrophobic surface of the N-lobe.

Collectively, our results show that the NTD of LonA selectively interacts with protein substrates, via critical

bulky hydrophobic residues on the hydrophobic surface of its N-lobe, demonstrating that NTD enables

LonA to perform protein quality control by selectively interacting with proteins in damaged or unfolded

states.

To further examine how the NTD is involved with protein aggregation, we probed the interactions

between the scrambled lysozyme aggregate and MtaLonA NTD by NMR spectroscopy. Native lysozyme

is a 14.3-kDa protein (129 amino acids) comprising four disulfide bridges while reduced lysozyme

forms loose and flexible aggregates[25]. TCEP (tris (2-carboxyethyl) phosphine; nonthiol-based reducing

agent) drives the formation of amorphous lysozyme aggregates observed in this study. Moreover, there is

no cysteine residue found in full-length MtaLonA, suggesting TCEP does not affect the structure of NTD.

Addition of TCEP-treated denatured lysozymes leaded to substantial spectral changes of NN206

resonances recorded at 32 °C while NN206* showed marginal CSPs marginal CSPs (supplementary Fig.

5A). The structural mapping shows that the affected residues caused by TCEP-treated denatured

lysozyme are also located mainly in the N-lobe shown in supplementary Figs. 5B and C. Furthermore, we

also examined the degradation activity of LonA against either native or denatured lysozyme, respectively.

A gel-based assay showed that full-length MtaLonA can efficiently degrade denatured-reduced lysozyme,

but not native lysozyme (supplementary Fig. 5D). Accordingly, AAAP, without the NTD, was inactive

against either the scrambled lysozyme aggregate or native lysozyme (supplementary Fig. 5E). Together,

these results highlight the key role of NTD, via the N-lobe region, in discriminating and recognizing protein

aggregates before subsequent translocation and degradation.

LonA can also degrade natively folded proteins with degradation tags such as sul20 degron[6,26].

Previously, the NTD of *E. coli* LonA was shown to be required for degron binding by chemical cross-

linking[7]. To understand how the NTD directly interacts with degrons, a 20-residue peptide (the C-terminal

20 residues of SulA; namely Sul20)[7] is synthesized and used for NMR titration experiments. Addition of

equimolar unlabeled Sul20 causes significant spectral changes of NN206 resonances (Fig. 4A) and the

structural mapping shows that the affected residues are again located at β-sheet β2/β5/β4, loop L1, L3

and helix α1 of the N-lobe subdomain (Fig. 4B). Mutations of P22 and M85 located at the center of the

hydrophobic surface to moderately hydrophobic Ala lead to much weak interactions of NN206* with Sul20

peptide (supplementary Fig. 6A), suggesting that specific hydrophobic residues of beta-stranded

β2/β5/β4 located at the N-lobe of MtaLon NTD play a critical role on degrons recognition.

To characterize the effect of MtaLon NTD on degrons further, we appended Sul20 to the C terminus

of a protein substrate Ig2D5 (domains 5 of the gelation factor from *Dictyostelium discoideum*, hereafter

abbreviated as Ig2D5) and investigated the conformational changes experienced by Sul20 upon binding

the NTD. The NMR spectra of Ig2D5 and Ig2D5-S20 can be well superimposed, indicating that the well-

folded domain structure corresponding to Ig2D5 is not perturbed by the C-terminal Sul20 (Fig. 4C).

Furthermore, Ig2D5-S20 exhibits additional cross-peaks that well correspond to Sul20. Elevated [^{1}H]–^{15}N

nuclear Overhauser effect (NOE) values for most residues are from Ig2D5 (Fig. 4D), along with the large

increase in chemical shift dispersion. Lower [^{1}H]–^{15}N NOEs for Sul20 residues are due to the increased

flexibility on ps-ns timescale. By addition of the unlabeled NN206 to isotopically enriched Ig2D5-S20, we

observed chemical shift changes and broadened resonances of C-terminal Sul20 (Figs. 4E and 4F),

suggesting that residues Y19 and H20 are strongly affected by the binding event based on the resonance

broadening beyond detection. The NMR spectrum of Ig2D5-S20 in the presence of an equimolar amount

of unlabeled NN206* shows a much less pronounced chemical shift effect (supplementary Fig. 6B) than

wild-type NN206 binding to Ig2D5-S20, indicating that Sul20 tag can be recognized by the hydrophobic

patches of N-lobe of MtaLon NTD.

The thermally unfolded conformations of substrate proteins are stabilized by engagement with

NTDs.

To understand further selective recognition of protein substrate in the unfolded states by the NTD,

we ask whether the unfolding-refolding process of protein may be modulated by interacting with the NTD.

To investigate how the conformations of substrates are influenced in the presence of NTDs, we

constructed a single-domain protein substrate Ig2D5 (domains 5 of the gelation factor from *Dictyostelium*

discoideum) for NMR experiments due to its high-quality NMR spectrum (supplementary Fig. 7A). About

96% of the backbone resonances of Ig2D5 can be assigned by multidimensional heteronuclear NMR

experiments. Ig2D5 is a β-sheet protein with Ig-like fold and the T_m of substrate Ig2D5 is similar to Ig2

(supplementary Figs. 7B and 7C). Here we applied NMR spectroscopy to investigate unfolding-refolding

equilibria of the substrate Ig2D5 induced by temperature cycling while MtaLon NTD remains stable

through the entire temperature cycling. The NMR samples containing isotopically enriched Ig2D5 in the

absence and presence of unlabeled NN206 were heated from 32 ºC to 60 ºC and then cooled to 32 ºC.

Based on the intensity changes of resonances corresponding to native Ig2D5, we estimated that isolated

Ig2D5 is largely unfolded at 60 ºC (Figs. 5A and B; the spectra are shown in green) and ~72% becomes

folded after a thermal cycling (Fig. 5C). However, in the presence of equimolar NN206, less than 15%

Ig2D5 is folded after one thermal cycle (Figs. 5C; the spectrum is shown in blue). Similar results were

obtained using equimolar full-length MtaLonA catalytic mutant S678A (LonA*) (Figs. 5C; the spectrum is

shown in red). However, addition of equimolar AAAP* (residues 242-793 with catalytic mutant S678A),

Ig2D5 is largely unfolded at 60 ºC (Fig. 5B; the spectrum is shown in black) and ~75% becomes folded

after thermal cycling (Fig. 5C; the spectrum is in black). The results reveal that the unfolded state of

Ig2D5 is engaged with MtaLon NTD and the temperature-induced unfolding-refolding of substrate is

significantly affected in the presence of MtaLon NTDs (Fig. 5D), potentially priming it for subsequent

translocation process. Accordingly, we also examined how the unfolding-refolding of Ig2D5 can be

affected by NN206*. After one thermal cycle, about 75% of Ig2D5 becomes folded in the presence of

equimolar NN206* (Fig. 5E and supplementary Figs. 7D, E and F). Upon addition of eightfold excess of

NN206*, about 45% of Ig2D5 becomes folded, suggesting that NN206* interacts with denatured protein

substrate weakly. These results demonstrate that the conformations of substrate are affected by the

hydrophobic substrate-binding interaction and the thermally unfolded states of substrate proteins can be

stabilized by engagement with the NTDs. Thus, by selectively interacting with hydrophobic residues

exposed in the thermally unfolded states of substrate proteins NTDs prominently perturb temperature-

induced unfolding-refolding process of substrate.

Discussion

In this work we used α-casein and Ig2, which represent two types of substrates. Native α-casein

is an intrinsically disordered protein; by contrast, the native Ig2 is a β-sheet protein with T_m at 46 °C

(Supplementary Fig. 1). We show that the construct AAAP devoid of the NTD (residues 242-793, ATPase

and protease domains), which retains wild-type ATPase and peptidase activities in LonA, lacks the

degradation activity against thermally denatured Ig2 while exhibits a wild-type-like activity against α-

casein (Fig.1). Therefore, the NTD is specific for denatured or damaged protein substrates, whose

hydrophobic core regions normally buried in the folded state become exposed, but not for intrinsically

disordered substrates like casein, which contains mainly charged or polar residues and lack hydrophobic

regions in the protein sequence. Indeed, it has been shown that LonA prefers substrates rich in

hydrophobic residues[26].

Our crystallographic results of NN206 and NN291 are consistent with previous findings that the

~40-residue region between the NTD and ATPase-Protease domains is structurally flexible with multiple

accessible protease sites[9,20,27,28]. This notion is also corroborated by our high-resolution NMR analyses

on the NTD both in isolation and as a part of the full-length protein, demonstrating the independent, fast

tumbling motion of the NTDs in the ~0.5 MDa hexameric assembly of LonA. Moreover, their highly

favorable NMR relaxation properties and resulting high-quality multidimensional NMR spectra have

allowed us to analyze the interaction of the NTD with unfolded proteins, protein aggregates, degron tags,

and intrinsically disordered substrates. Our work provides atomic details of the NTD-mediated substrate

discrimination by temperature-based switching of the native folded protein to its unfolded state in an NMR

sample. NMR characterization indicates NTD does not interact with well-folded Ig2, native lysozyme or

intrinsically disordered α-casein, whereas unfolded proteins, aggregates and degron tags can elicit

pronounced chemical shift changes to the N-lobe of the NTD, which is subsequently confirmed by

structure-based mutagenesis. Collectively, these results suggests that LonA has two substrate-interacting

modes: (1) the NTD-non-requiring mode for intrinsically disordered substrates lacking hydrophobic-rich

region, which may be engaged directly with the pore-loops in the LonA assembly without involving the six

NTDs; (2) the NTD-requiring mode for recognizing and trapping damaged/denatured protein substrates

with exposed hydrophobic regions before their engagement with the pore-loops of the LonA assembly.

The key role of NTD is to enable LonA to perform protein quality control to selectively capture and

degrade proteins in damaged unfolded states.

By analyzing the NTD of MtaLonA interactions with damaged Ig2, the scrambled lysozyme and

Sul 20 peptide, we conclude that these binding events are mediated by two hydrophobic patches, which

comprise (i) Leu-10, Ile-15, Pro-22, Val-23, and Met-85 (termed hpI in the following) ; (ii) Met-75, Leu-77,

Pro-78 and Leu-82 (termed hpII) (Fig. 6A). Our results also demonstrate that a double mutant NN206*

(Pro-22 and Met-85 replaced by Ala), resulting in decreased hydrophobicity of hpI, has a significant effect

on the ability of MtaLonA NTD to bind to damaged or misfolded proteins, indicating that the bulky

hydrophobic residues on hydrophobic patches are essential determinants for substrate recognition and

discrimination. We also examine the exposed hydrophobic residues of β-sheet β2/β5/β4, loop L1 and loop

L3 located at the N-lobes of other reported structures of LonA NTDs[19-22]. The N-terminal fragment from *B.

subtilis* LonA was previously reported to adopt a domain-swapped dimer where the N-lobe of one

monomer is positioned next to the C-lobe of the other monomer[20]. In contrast, the structures of the N-

terminal fragment from MtaLon, EcLon and MacLon showed that the N- and the C-lobes are joined

together via a short linker (supplementary Fig. 8A and B). Based on structural alignments (supplementary

Fig. 8C), the N-lobe of EcLon has almost the same pattern of exposed hydrophobic residues as that of

MtaLon NTD (Fig. 6B). Interestingly, The NTD of MacLon shows merged hp I and II while the hpI of

BsLon NTD is a little distant from hpII (Figs. 6C and D). The structural comparison reveals that, similarly

to MtaLon NTD, all three reported structures exhibit exposed hp I and II with slight variations in their

shapes and sizes, suggesting that the conserved hydrophobic patches at the N-lobes may be potentially

responsible for conferring substrate selectivity toward degrons, misfolded and damaged proteins.

It has previously been known that LonA can recognize a degron-tagged protein by binding to its

largely exposed hydrophobic residues. Here, our results show the C-terminal Sul20 of Ig2D5-S20 has

substantially low heteronuclear NOE values, reflecting this region undergoes rapid local motion and is

solvent-exposed in solution. Therefore, NTD of MtaLonA can easily recognize and interact with the C-

terminal aromatic and hydrophobic residues of Sul20. This finding indicates that a pair of aromatic

residues consisting of Y19 and H20 in Sul20 play crucial roles in the hydrophobic interactions with

residues P22 and M85 of LonA NTD, which is verified with a double mutant NN206*. Residues Pro and

Met exhibit higher values of hydrophobicity than moderately hydrophobic Ala and mutations of both P22

and M85, located at the center of the contiguous hydrophobic surface, result in much weak interactions of

NN206* with damaged Ig2, the scrambled lysozyme and Sul20 peptide, highlighting the importance of

direct interaction with exposed and contiguous surface with high hydrophobicity in the recognition of

protein substrates by the NTD of MtaLonA. Furthermore, by selectively binding to hydrophobic residues

exposed in the thermally unfolded states of proteins, the NTDs prominently stabilize substrate's unfolded

conformation. These results show that the NTD-substrates interactions involve the hydrophobic residues

on both partners, suggesting aromatic and hydrophobic side chains, which are usually buried in the

native proteins, are important recognition determinants for the NTD to enable MtaLonA to selectively

capture and degrade proteins in a damaged unfolded state or with degron tags.

Our results directly demonstrate that thermally damaged proteins and degrons induce chemical

shift changes and broadened resonances located mainly in the two hydrophobic patches of N-terminal

lobe but we do not find evidence that the C-lobe of the NTD is involved in substrate interaction. However,

addition of thermally unfolded Ig2 into U-^{2}H/^{15}N-labelled full-length LonA*, affected residues are shown in

both N- and C-lobes of LonA* NTD at 55 °C (Figs. 7A and B). Compared ^{1}H–^{15}N TROSY HSQC spectra

of NN206 and MtaLonA* in the presence of thermally unfolded protein Ig2, the signals from the C-lobe of

MtaLonA* NTD are missing due to line broadening, suggesting others NTDs of LonA* hexamer may join

to interact with substrates. All together, these results reveal the free molecular tumbling of the NTD and

its substrate binding site, based on which a model involving multi-NTD interactions, may be proposed for

substrate recognition. The flexibly linked NTD of the hexametric LonA is swaying from side to side and

back and forth to survey, recognize, and trap substrates with exposed hydrophobic sequences (Fig. 7C).

After initial binding, other NTDs may join to increase the avidity of the LonA-substrate interaction (Fig.

7D); the engagement of a substrate protein by multiple NTDs also serves to facilitate effective substrate

unfolding and translocation mediated by the coordinated movements of the pore-loops in the ATPase

modules of the LonA chamber, powered by cycles of ATP binding and hydrolysis. Finally, the substrate

polypeptide chain is pulled inside the chamber and undergoes proteolysis by protease modules of LonA.

Materials and Methods

Peptide

Sul20 peptide (Sequence: ASSHATRQLSGLKIHSNLYH) was synthesized by GenScript

(https://www.genscript.com/) at >95% purity.

Cloning, protein expression and purification

The plasmids expressing for full-length MtaLonA (1-793 residues) or AAAP (242-793 residues) with a C-

terminal 6xHis-tag were transformed into *E.coli* BL21(DE3) cells[18]. Site-directed mutagenesis was

performed using the Quickchange kit (Stratagene). NN206 (1-206 residues), NN243 (1-243 residues) and

NN291 (1-291 residues) were cloned into pET-modified vector with a tobacco etch virus (TEV) cleavage

site. The three resultant plasmids, encoding the proteins NN206, NN243 and NN291 with the N-terminal

His-tag, were transformed into *E.coli* BL21(DE3) cells. The target proteins synthesis was induced with 0.5

mM isopropyl-thio-β-D-galactoside (IPTG) at an absorbance at 600 nm (OD_{600}) ~0.6 at 28 °C. The target

proteins were purified with nickel chelate affinity resins (Ni-NTA, Qiagen). The protein samples were

collected and purified by a Superdex 200 (GE Healthcare) column[17]. MtaLonA and AAAP proteins were

collected and further purified on a MonoQ (GE Healthcare) chromatography. For NN206, NN243 and

NN291 purification, the N-terminal His-tagged proteins were cleaved by TEV protease for overnight at 4

°C and then reloaded onto Ni-NTA to remove TEV protease. The flow-through fraction containing target

proteins were collected and purified with Superdex 75 (GE Healthcare) chromatography. Ig2 (domains 5

and 6 of the gelation factor ABP-120 of *Dictyostelium discoideum*)[29] and Ig2D5-S20 (Ig2 fused with

Sul20) were cloned into pET28a(+)tev for generation of pET28a(+)tev-Ig2 and pET28a(+)tev-Ig2D5-20.

The recombinant protein was induced by 0.5 mM IPTG at OD_{600} ~0.6 for 16 hours. The purifications of

Ig2, Ig2D5 and Ig2D5-20 is the same as NN206, NN243 and NN291. Isotopically labeled samples for

NMR studies were prepared by growing the cells in minimal (M9) medium. The full-length MtaLonA were

prepared by supplementing the growing medium with 1 g l^{-1} of $^{15}NH_4Cl$ and 4 g l^{-1} of $^2H_7/^{12}C_6$-glucose in

99.8%-^2H_2O (Sigma-Aldrich).

Substrate degradation assays

α-casein (Sigma, USA) and Ig2 were used as the substrates in these assays[17]. 4 μM substrate proteins

were incubated with 0.4 μM MtaLonA (hexamer) or AAAP or mutations at 55 °C. The degradation

reactions were halted by protein sample dye and heat inactivation at 95 °C for 5 minutes. The treated

reaction mixtures were then analyzed by SDS-PAGE.

Protein crystallization

Crystallizations of NN206 and NN291 were performed at 295K by hanging drop vapor-diffusion method.

For Crystallization of NN206, in situ proteolysis was performed[30]. 1 μl of MtaLonA (10 mg/ml) plus trypsin

in ratio 1000:1 was mixed with 1 μl of 0.1 M Tris-HCl (pH 7.5), 0.2 M Sodium acetate and 30% PEG 4000.

The NN206 crystals appeared in the mixed drop after 3 days. The crystals of NN291 were grown by

mixing with 1 μl of NN291 protein sample (15 mg/mL) and 1 μl of well solution containing 0.1 M MES (pH

5.8) and 0.8 M ammonium sulfate. Crystals grew to 0.2-0.3 mm over two weeks. Crystals of NN206 and

NN291 were cryoprotected with 20% glycerol or 20% ethylene glycol before data collection.

Structure determination

The data sets of NN206 and NN291 were collected at the beamlines BL-13C1 of National Synchrotron

Radiation Research Center (Taiwan) and BL-1A of Photon Factory (Japan), respectively. All Data sets

were processed by HKL2000[31]. The structure of NN206 was solved by molecular replacement with the

program Phaser[32] using the structures of the N-terminal fragment of *B. subtilis* LonA (residues 4-116,

PDB code 3M65) and the N-terminal fragment of *E. coli* LonA (residues 120-210, PDB code 3LJC) as the

search models. The structure of NN291 was determined by molecular replacement using NN206

structure. All initial models were automatically rebuilt using the program AutoBuild[33]. Subsequently, these

structures were refined by manual refitting in Coot[34] and performance of refinement using the program

Refmac5[35]. Crystallographic and refinement statistics are listed in Table 1. All structure figures were

made using PyMOL (Version 1.3, Schrödinger, LLC). The atomic coordinates and structure factors for

NN206 and NN291 have been deposited in the Protein Data Bank (http://www.rcsb.org/) with the

accession numbers 7CR9 and 7CRA, respectively.

Thermal shift assay

Thermal shift assay was carried out in qPCR 8-strip tubes (Gunster Biotech) using SYBRO orange (Life

Technologies) as dye. For each reaction, 5 µM purified MtaLonA, NN206 or Ig2 was mixed with assay

buffer (50 mM NaPi, 2 mM β-Me, pH 6.5) and SYPRO Orange dye (5X final concentration) in 45 µL total

volume/well. 100 µM Ig2D5 was mixed with assay buffer and SYPRO Orange dye (10X final

concentration) in 45 µL total volume/well. Samples were heated at a rate of 0.5 °C per minute from 25 °C

to 95 °C and fluorescence signals were recorded with the CFX Connect Real-Time PCR Detection

System (Bio-Rad). The melting temperatures were analyzed using the derivative plots of the melting

curve.

NMR spectroscopy

NMR experiments were acquired on Bruker 800-MHz spectrometers (Bruker BioSpin, Karlsruhe,

Germany). The assignments of NN206 backbone ^{15}N, ^{1}H^{N}, ^{13}Cα, ^{13}Cβ, and ^{13}C′ chemical shifts were

obtained by ^{15}N-^{1}H HSQC, HNCACB, CBCA(CO)NH, HNCA, HN(CO)CA and HNCO spectra[36]. NMR

samples containing 50 µM [U-15N]-labeled NN206 in the absence and presence of 100 µM Ig2 are

prepared in the buffer: 50 mM sodium phosphate (pH 6.5) with 10% D2O. For temperature cycling

experiments, the first 2D ^{1}H–^{15}N TROSY HSQC spectrum was acquired at 32 °C; the same sample was

heated to 55 °C for recording the second spectra, and the temperature is decreased to 32 °C for

recording the third spectra. For characterization of substrate conformation selection by NTD, 2D ^{1}H-^{15}N

TROSY HSQC NMR spectra of 50 µM [U-^{15}N]-labeled Ig2D5 titrated with 50 µM unlabeled NN206,

MtaLonA or AAAP proteins were recorded during a thermal cycle starting from 32 to 60 °C, and then back

to 32 °C. 72 residues of Ig2D5 were selected for calculating folded ratio. The interaction of NN206 with

Sul20 peptide was recorded with 50 µM [U-15N]-labeled NN206 titrated with unlabeled 50 and 150 µM

unlabeled Sul20 peptide. The interaction of NN206 with TCEP-treated denatured lysozyme was recorded

with 50 µM [U-15N]-labeled NN206 titrated with 25 µM unlabeled denatured lysozyme. ^{1}H-^{15}N NOE data

were recorded in an interleaved manner: one spectrum with 4 s recycle delay followed by 4 s saturation

and another spectrum with no saturation and 8 s recycle delay. All NMR data were processed and

analyzed by XWIN-NMR (Bruker BioSpin), NMRPipe[37], and NMRView[38].

Data availability

The atomic coordinates and structure factors have been deposited in the Protein Data

Bank, www.pdb.org, with accession codes 7CR9 and 7CRA. Other relevant data are available from the

corresponding author upon reasonable requests.

References

1. Charette, M.F., Henderson, G.W. & Markovitz, A. ATP hydrolysis-dependent protease
activity of the lon (capR) protein of Escherichia coli K-12. *Proc Natl Acad Sci U S A* **78**, 4728-
32 (1981).
2. Park, S.C. et al. Oligomeric structure of the ATP-dependent protease La (Lon) of
Escherichia coli. *Mol Cells* **21**, 129-34 (2006).
3. Goldberg, A.L., Moerschell, R.P., Chung, C.H. & Maurizi, M.R. ATP-dependent protease La
(lon) from Escherichia coli. *Methods Enzymol* **244**, 350-75 (1994).
4. Rotanova, T.V. et al. Classification of ATP-dependent proteases Lon and comparison of
the active sites of their proteolytic domains. *European Journal of Biochemistry* **271**, 4865-
4871 (2004).
5. Botos, I. et al. The catalytic domain of Escherichia coli Lon protease has a unique fold and
a Ser-Lys dyad in the active site. *Journal of Biological Chemistry* **279**, 8140-8148 (2004).
6. Higashitani, A., Ishii, Y., Kato, Y. & Koriuchi, K. Functional dissection of a cell-division
inhibitor, SulA, of Escherichia coli and its negative regulation by Lon. *Mol Gen Genet* **254**,
351-7 (1997).
7. Wohlever, M.L., Baker, T.A. & Sauer, R.T. Roles of the N domain of the AAA+ Lon protease
in substrate recognition, allosteric regulation and chaperone activity. *Mol Microbiol* **91**,
66-78 (2014).
8. Ebel, W., Skinner, M.M., Dierksen, K.P., Scott, J.M. & Trempy, J.E. A conserved domain in
Escherichia coli Lon protease is involved in substrate discriminator activity. *J Bacteriol* **181**,
2236-43 (1999).
9. Roudiak, S.G. & Shrader, T.E. Functional role of the N-terminal region of the Lon protease
from Mycobacterium smegmatis. *Biochemistry* **37**, 11255-63 (1998).
10. Rudyak, S.G. & Shrader, T.E. Polypeptide stimulators of the Ms-Lon protease. *Protein Sci*
9, 1810-7 (2000).
11. Iyer, L.M., Leipe, D.D., Koonin, E.V. & Aravind, L. Evolutionary history and higher order
classification of AAA+ ATPases. *J Struct Biol* **146**, 11-31 (2004).
12. Melnikov, E.E. et al. Limited proteolysis of E. coli ATP-dependent protease Lon - a unified
view of the subunit architecture and characterization of isolated enzyme fragments. *Acta*
Biochim Pol **55**, 281-96 (2008).
13. Adam, C. et al. Biological roles of the Podospora anserina mitochondrial Lon protease and
the importance of its N-domain. *PLoS One* **7**, e38138 (2012).

14. Cheng, I. et al. Identification of a Region in the N-Terminus of Escherichia coli Lon That
Affects ATPase, Substrate Translocation and Proteolytic Activity. *Journal of Molecular*
Biology **418**, 208-225 (2012).
15. Gottesman, S. Proteolysis in bacterial regulatory circuits. *Annu Rev Cell Dev Biol* **19**, 565-
87 (2003).
16. Baker, T.A. & Sauer, R.T. ATP-dependent proteases of bacteria: recognition logic and
operating principles. *Trends Biochem Sci* **31**, 647-53 (2006).
17. Lin, C.C. et al. Structural Insights into the Allosteric Operation of the Lon AAA+ Protease.
Structure **24**, 667-675 (2016).
18. Su, S.C. et al. Structural Basis for the Magnesium-Dependent Activation and
Hexamerization of the Lon AAA+ Protease. *Structure* **24**, 676-686 (2016).
19. Li, M. et al. Crystal structure of the N-terminal domain of E. coli Lon protease. *Protein Sci*
14, 2895-900 (2005).
20. Duman, R.E. & Lowe, J. Crystal structures of Bacillus subtilis Lon protease. *J Mol Biol* **401**,
653-70 (2010).
21. Li, M. et al. Structure of the N-terminal fragment of Escherichia coli Lon protease. *Acta*
Crystallogr D Biol Crystallogr **66**, 865-73 (2010).
22. Chen, X. et al. Crystal structure of the N domain of Lon protease from Mycobacterium
avium complex. *Protein Sci* **28**, 1720-1726 (2019).
23. Hsu, S.T., Cabrita, L.D., Fucini, P., Dobson, C.M. & Christodoulou, J. Structure, dynamics
and folding of an immunoglobulin domain of the gelation factor (ABP-120) from
Dictyostelium discoideum. *J Mol Biol* **388**, 865-79 (2009).
24. Van Melderen, L. et al. ATP-dependent degradation of CcdA by Lon protease. Effects of
secondary structure and heterologous subunit interactions. *J Biol Chem* **271**, 27730-8
(1996).
25. Yang, M., Dutta, C. & Tiwari, A. Disulfide-bond scrambling promotes amorphous
aggregates in lysozyme and bovine serum albumin. *J Phys Chem B* **119**, 3969-81 (2015).
26. Gur, E. & Sauer, R.T. Recognition of misfolded proteins by Lon, a AAA(+) protease. *Genes*
Dev **22**, 2267-77 (2008).
27. Patterson, J. et al. Correlation of an adenine-specific conformational change with the ATP-
dependent peptidase activity of Escherichia coli Lon. *Biochemistry* **43**, 7432-42 (2004).
28. Vasilyeva, O.V., Kolygo, K.B., Leonova, Y.F., Potapenko, N.A. & Ovchinnikova, T.V. Domain
structure and ATP-induced conformational changes in Escherichia coli protease Lon
revealed by limited proteolysis and autolysis. *FEBS Lett* **526**, 66-70 (2002).
29. McCoy, A.J., Fucini, P., Noegel, A.A. & Stewart, M. Structural basis for dimerization of the
Dictyostelium gelation factor (ABP120) rod. *Nat Struct Biol* **6**, 836-41 (1999).
30. Wernimont, A. & Edwards, A. In situ proteolysis to generate crystals for structure
determination: an update. *PLoS One* **4**, e5094 (2009).
31. Otwinowski, Z. & Minor, W. [20] Processing of X-ray diffraction data collected in
oscillation mode. *Methods Enzymol* **276**, 307-326 (1997).
32. McCoy, A.J. Solving structures of protein complexes by molecular replacement with
Phaser. *Acta Crystallogr D Biol Crystallogr* **63**, 32-41 (2007).

33. Terwilliger, T.C. et al. Iterative model building, structure refinement and density
modification with the PHENIX AutoBuild wizard. *Acta Crystallogr D Biol Crystallogr* **64**, 61-
9 (2008).
34. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
Crystallogr D Biol Crystallogr **60**, 2126-32 (2004).
35. Murshudov, G.N. et al. REFMAC5 for the refinement of macromolecular crystal structures.
Acta Crystallogr D Biol Crystallogr **67**, 355-67 (2011).
36. Sattler M, S.J., Griesinger C. Heteronuclear multidimensional NMR experiments for the
structure determination of proteins in solution employing pulsed field gradients. *Progress*
in Nuclear Magnetic Resonance Spectroscopy. **34**, 93-158 (1999).
37. Delaglio, F. et al. NMRPipe: a multidimensional spectral processing system based on UNIX
pipes. *J Biomol NMR* **6**, 277-93 (1995).
38. Downing, A.K. *Protein NMR techniques*, xi, 487 p. (Humana Press, Totowa, N.J., 2004).
521

Acknowledgements

We thank Dr. Shang-Te Danny Hsu for helpful discussion. We also thank Ms S.-L. Huang of
Ministry of Science and Technology (National Taiwan University) for the assistance in NMR experiments.
We are grateful to the staff of the Biomedical Resource Core at the First Core Labs, National Taiwan
University College of Medicine, for technical assistance. Portions of this research were carried out at
beamlines 13B1 and 13C1 of the National Synchrotron Radiation Research Center (Taiwan). This work
was supported by National Taiwan University Grants 108L7809 and 109L7809; and the Ministry of
Science and Technology, Taiwan (grant numbers: MOST108-2113-M-002-006- and MOST109-2113-M-
002-018- to S.-R.T. and MOST105-2320-B-001-015-MY3 to C.-I.C.).
531

532 **Author contributions**
533 S.R.T., and C.I.C. conceived the study. S.R.T., and C.I.C. designed the experiments and analyzed the
534 data. Y.C.T., C.C.L., Y.T.K., and S.J.H. performed the experiments. S.R.T., and C.I.C. wrote the
535 manuscript with inputs from all other authors.

546   **Table 1.** Data collection and refinement statistics

	NN206	NN291
PDB entry	7CR9	7CRA
Data collection		
Space group	P212121	P6322
Cell dimensions		
a, b, c (Å)	32.830, 58.296, 198.492	86.016, 86.016, 110.126
α, β, γ (°)	90, 90, 90	90, 90, 120
Resolution (Å)	30 - 2.09 (2.16 - 2.09)*	30 - 1.7 (1.76 - 1.7)
R_{sym} or R_{merge}	0.106 (0.937)	0.069 (0.779)
$I/\sigma I$	17.9 (3.0)	49.1 (4.7)
Completeness (%)	99.7 (100)	100 (100)
Redundancy	6.5 (6.6)	20.9 (21.4)
Refinement		
Resolution (Å)	29.409 - 2.1 (2.149 -2.1)	30 - 1.7 (1.744 - 1.7)
R_{work}/ R_{free}	0.209/0.249	0.197/0.237
No. atoms		
Protein	3285	1684
Ligand (SO_4^{2-})	0	10
Water	107	244
B-factors		
Protein	53.14	16.18
Ligand (SO_4^{2-})		27.45
Water	52.66	28.21
R.m.s deviations		
Bond lengths (Å)	0.004	0.009
Bond angles (º)	0.74	1.34
Ramachandran		
Favored (%)	96.81	96.67
Allowed (%)	2.46	2.38
Outliers (%)	0.74	0.95

547   Number of crystals for each structure should be noted in footnote.
548   *Highest resolution shell is shown in parenthesis.
549

A

B

551   Fig. 1. ***The NTD of Meiothermus taiwanensis LonA (MtaLonA) is essential for efficient***

552   ***degradation of a thermally unfolded substrate, but not the intrinsically disordered substrate, α-***

553   ***casein.*** (A) Schematic diagrams illustrating domain organization of full-length MtaLonA and the

554   N-terminal domain truncated AAAP. (B) Degradation of α-casein by full-length MtaLonA (FL)

555   and AAAP at 55 °C. (C) Degradation of Ig2 by full-length and AAAP at 55 °C.

Fig. 2. ***Overall structures of the N-terminal domain of MtaLonA.*** (A) The three N-terminal constructs NN206, NN291, and NN243. The disordered region is shown in the dashed box. (B) Structure of NN206. (C) Superposition of NN206 (marine) and NN291 (pink) structures.

563 *Figure 2 continued*

564 (D) The refined *2F₀-Fc* electron density map around the C-terminal residue A210 of NN291,

565 contoured at 1.0 σ. (E) Overlay of 2D ^{1}H–^{15}N HSQC NMR spectra of NN206 (red) and NN243

566 (blue). (F) Comparison of the ^{1}H–^{15}N TROSY HSQC spectra of NN206 (red) and MtaLonA

567 (black) recorded at 55 °C. The well-dispersed resonances are labeled with residue numbers.

568 About 200 well-resolved correlations are superimposable, suggesting that NTD is loosely linked

569 to the hexameric core via a flexible linker. (G) The six NTD may each connect to the hexameric

570 core of fused AAA+ and protease domains by a flexible ~40-residue linker. The NTDs, and

571 ATPase-Protease chamber are illustrated in blue and green colors, respectively. Red dashed lines

572 represent the flexible linker regions.

Fig. 3. ***The N-lobe of MtaLonA NTD is involved in recognition of thermally unfolded substrate***

Ig2. (A) ^{1}H–^{15}N TROSY HSQC spectra of NTD in the absence (blue) and presence (red) of Ig2

recorded at 32 °C. (B) Comparison of the ^{1}H–^{15}N TROSY HSQC spectra of NN206 in the apo

(black) and Ig2-bound (magenta) states recorded at 55 °C. (C) After one thermal cycle, ^{1}H–^{15}N

TROSY HSQC spectra of NTD in the absence (green) and presence (orange) of Ig2 recorded at

32 °C. (D) Gel filtration profiles of NMR samples treated with thermal cycling shown in blue, and

folded Ig2 (red) analyzed by Superdex 200 10/300 GL column.

Figure 3 continued on next page

(E) Comparison of the ^1H–^{15}N TROSY HSQC spectra of NN206 in the apo (black) and Ig2-bound

(magenta) states. The residues of NTD with the CSPs are part of the interaction interface, located

mainly at β-sheet β2/β5/β4, loop L1, L2, L3 and helix α1. *Figure 3 continued on next page*

(F) Interaction interface of NTD: Ig2 mapped onto the structure of NTD, based on the spectrum recorded at 55 °C. Proline residue is shown in green. (G) The exposed hydrophobic residues that make up the damaged Ig2-binding sites are shown as balls and sticks. Mutations of two exposed hydrophobic residues P22 and M85, located at the center of β-sheet β2/β5/β4, lead to reduce the hydrophobicity of the contiguous hydrophobic surface in the N-lobe subdomain.

Figure 3 continued on next page

Figure 3 continued

(H) ^1H–^{15}N TROSY HSQC spectra of NTD* (blue) in the absence (blue) and presence (red) of

Ig2 recorded through thermal cycling from 32 °C (H) to 55 °C (I) and then returning to 32 °C (J).

Fig. 4. ***Interaction of MtaLonA NTDs with degron tag Sul20.*** (A) Overlay of the ^1H–^{15}N TROSY

HSQC spectra of NN206 in the absence (black) and presence (red) of Sul20 peptide. (B) Structural

mapping of the chemical shift perturbations (orange) of NN206 caused by Sul20 peptide.

Figure 4 continued on next page

 (C) Overlay of the ^{1}H–^{15}N TROSY HSQC spectra of Ig2D5 (green) and Ig2D5-S20 (black);

 overlay of the ^{1}H–^{15}N TROSY HSQC spectra of Ig2D5-S20 in the absence (black) and in the

 presence (orange) of an equimolar amount of NN206. (D) [^{1}H]–^{15}N nuclear Overhauser effect

628   (NOE) values of Ig2D5-S20. (E) Expanded region of the ^{1}H–^{15}N TROSY HSQC spectra of Ig2D5-

629   S20 in the absence (black) and in the presence (orange) of NN206. (F) CSPs of amide moieties of

630   50 µM Ig2D5-S20 after binding 50 µM unlabeled NN206, plotted against Sul20 residue number.

631   Residues Y19 and H20 are strongly affected by the binding event based on the resonance

632   broadening beyond detection.

Fig. 5. *The thermodynamics of Ig2D5 unfolding-refolding is affected in the presence of MtaLonA NTDs.* 2D ^{1}H-^{15}N TROSY HSQC NMR spectra of apo Ig2D5(green), Ig2D5 titrated with NN206(blue), LonA (red), or AAAP (black) are recorded with a thermal cycle starting from 32 °C (A) to 60 °C (B), and then returning to 32 °C (C). (D) Folded ratio of apo Ig2D5, Ig2D5 titrated with NN206(blue), LonA (red), or AAAP (black) after a thermal cycle. (E) Folded ratio of apo Ig2D5, Ig2D5 titrated with different concentrations of NN206* after one thermal cycle.

Fig. 6 ***The NTD of MtaLonA selectively interacts with unfolded proteins, protein aggregates,***

and degron-tagged proteins via two hydrophobic patches of its N-lobe. (A) Two hydrophobic

patches located at the N-lobe of MtaLonA: (i) Leu-10, Ile-15, Pro-22, Val-23, and Met-85

(termed hpI in the following); (ii) Met-75, Leu-77, Pro-78 and Leu-82 (termed hpII). (B) Hp I

and II of EcLonA: (i) Leu-16, Val-21, Pro-28, Leu-29, and Leu-91; (ii) Leu-81, Leu-83, Pro-84

and Val-88. (C) Hp I and II of BsLonA: (i) Leu-14, Leu-17, Leu-18, Val-19, Val-24, Leu-25,

Leu-27 and Leu-89; (ii) Leu-79, Leu-81, Pro-82 and Ile-86. (D) merged Hp I and II of

MacLonA: (i) Val-12, Val-17, Val-22, Pro-24, Ile-25, Leu-66 and Val-79; (ii) Ile-71, Ala-72 and

Ala-77. The conserved hydrophobic patches at the N-lobes may be potentially responsible for

conferring substrate selectivity toward degrons, misfolded and damaged proteins. The PDB

codes are indicated in parentheses.

Fig. 7. *Proposed model for NTD-mediated substrate interaction.*

(A) ^{1}H–^{15}N TROSY HSQC spectra of NTD in the absence (black) and presence (red) of

unlabeled Ig2 recorded at 55 °C. (B) ^{1}H–^{15}N TROSY HSQC spectra of LonA* in the absence

(black) and presence (green) of unlabeled Ig2 recorded at 55 °C. (C) The flexibly linked NTD of

the hexametric LonA is swaying from side to side and back and forth to survey, recognize, and

trap substrates with exposed hydrophobic sequences. Substrate, N-lobe, Clobe of NTDs, and the

ATPase-Protease chamber are illustrated in purple, marine, blue and green colors, respectively.

The substrate-binding patches are depicted in yellow. Red dashed lines represent the flexible

linker regions. (D) After initial binding, other NTDs may join to increase the avidity of the

LonA-substrate interaction. The substrate polypeptide chain is pulled inside the chamber and

undergoes proteolysis by protease modules of LonA.