Conformational dynamics of auto-inhibition in the ER calcium sensor STIM1

Stijn van Dorp[1], Ruoyi Qiu[1], Ucheor B. Choi[1,3], Minnie M. Wu[1], Michelle Yen[1], Michael Kirmiz[1], Axel T. Brunger[1,2], and Richard S. Lewis[1]*

[1]Department of Molecular and Cellular Physiology, Stanford University School of Medicine, Stanford, CA 94305

[2]Howard Hughes Medical Institute, Stanford University School of Medicine, Stanford, CA 94305, USA

[3]Present address: Department of Biochemistry, West Virginia University School of Medicine, Morgantown, WV 2650

*email: rslewis@stanford.edu

1   **Abstract**

The dimeric ER Ca^{2+} sensor STIM1 controls store-operated Ca^{2+} entry (SOCE) through the

regulated binding of its CRAC activation domain (CAD) to Orai channels in the plasma

membrane. In resting cells, the STIM1 CC1 domain interacts with CAD to suppress SOCE, but

the structural basis of this interaction is unclear. Using single-molecule Förster resonance energy

transfer (smFRET) and protein crosslinking approaches, we show that CC1 interacts dynamically

with CAD in a domain-swapped configuration with an orientation predicted to sequester its Orai-

binding region adjacent to the ER membrane. Following ER Ca^{2+} depletion and release from

CAD, cysteine crosslinking indicates that the two CC1 domains become closely paired along

their entire length in the active Orai-bound state. These findings provide a structural basis for

the dual roles of CC1: sequestering CAD to suppress SOCE in resting cells and propelling it

towards the plasma membrane to activate Orai and SOCE after store depletion.

Introduction

Store-operated Ca^{2+} entry (SOCE) is a nearly ubiquitous signaling pathway activated by

extracellular stimuli that deplete Ca^{2+} from the endoplasmic reticulum (ER). SOCE is essential

for many physiological functions of excitable and non-excitable cells, including gene expression,

secretion, and motility (Prakriya and Lewis, 2015). Accordingly, loss-of-function mutations in

the core SOCE components lead to serious human pathologies, including severe combined

immunodeficiency, autoimmunity, myopathy, ectodermal dysplasia and anhidrosis (Lacruz and

Feske, 2015), while gain-of-function mutations cause Stormorken syndrome, miosis, myopathy,

thrombocytopenia and excessive bleeding (Böhm and Laporte, 2018). These clinical

manifestations underscore the need for precise regulation of SOCE to ensure that it is silent when

ER Ca^{2+} stores are full yet reliably activated by stimuli that drive store depletion.

The ER membrane protein STIM1 (Figure 1A, top) controls SOCE through the regulated

interaction of its CRAC activation domain (CAD (Park et al., 2009); also known as SOAR (Yuan

et al., 2009) or CCb9 (Kawasaki et al., 2009)) with Orai1 Ca^{2+} channels in the plasma

membrane. In resting cells, Ca^{2+} bound to the luminal EF hands of the STIM1 dimer suppresses

STIM1 activity by promoting intramolecular association of the coiled-coil 1 (CC1) domains with

CAD, referred to as the 'inhibitory clamp' (Korzeniowski et al., 2010; Fahrner et al., 2014; Ma et

al., 2015; Yang et al., 2012; Yu et al., 2013). Upon ER Ca^{2+} depletion, dissociation of Ca^{2+} from

the luminal EF hands triggers conformational changes that release CC1 from CAD and promotes

STIM1 accumulation at ER-plasma membrane (ER-PM) junctions where the CAD domain binds,

traps, and activates Orai1 channels diffusing in the PM (Liou et al., 2007; Wu et al., 2014, 2006).

Thus, to elucidate the regulatory mechanism of SOCE, it is essential to understand how CC1

interacts with CAD to control STIM1 activity.

The CC1 domain comprises three helical segments, CC1α1-3 (Soboloff et al., 2012)

(Figure 1A). Current evidence suggests that CC1α1, the most ER-proximal segment, acts as a

bimodal switch to stabilize both the quiescent and active states of STIM1. When ER Ca^{2+} stores

are full, CC1α1 binds to the CAD CC3 helix to sequester CAD (Fahrner et al., 2014; Ma et al.,

2015); upon store depletion, the luminal EF-SAM domains dimerize, CC1α1 unbinds from CAD,

and the transmembrane and proximal CC1α1 regions form a coiled-coil (Stathopulos et al., 2006;

Schober et al., 2019; Hirve et al., 2018), projecting CAD towards the plasma membrane.

Although several residues in CC1α1 and CC3 have been identified as essential for forming the

inhibitory clamp (Fahrner et al., 2014; Muik et al., 2011; Zhou et al., 2013; Ma et al., 2015), the

helical arrangement of the CC1α1:CAD binding interface is completely unknown. In addition,

while CC1α3 is implicated in stabilizing the inactive state (Yang et al., 2012; Korzeniowski et

al., 2010), it is unclear whether this results from binding to CAD or another mechanism (Zhou et

al., 2013).

Attempts to determine the structure of full-length STIM1 in the inactive or active state

have been unsuccessful. Instead, structures of several fragments from the STIM1 cytosolic

region have offered potential clues, although it is not yet clear how they might relate to

physiological states of the full-length protein. The crystal structure of human STIM1 CAD (aa

345-444, with mutations L374M, V419A, and C437T) depicts a parallel V-shaped dimer

comprising CC2 and CC3 domains (Figure 1A, bottom left) (Yang et al., 2012). It is unknown

whether this structure represents an active or inactive conformation, and the absence of CC1

precludes inferences about its interaction with CAD. A solution NMR structure of a CC1α3-

CC2 fragment depicts a dimer of wedged antiparallel CC2 helices (Figure 1A, bottom right)

(Stathopulos et al., 2013). This NMR structure can bind Orai1 C-terminal fragments *in vitro* and

has been proposed to be an intermediate in the activation of STIM1 (Stathopulos et al., 2013),

but to our knowledge it has never been shown to bind intact Orai1 channels *in situ* and lacks the

CC3 helix needed to evoke SOCE (Covington et al., 2010). Finally, in a crystal structure of CC1

(aa 237-340, with mutations M244L and L321M), two elongated CC1 helices form an

antiparallel coiled-coil dimer within the CC1α2 and CC1α3 domains (see Figure 5D (Cui et al.,

2013), but it is unclear how this elongated structure with its widely separated C-termini could

attach to the closely paired N-termini of CAD.

Although the sequences of these three structures partially overlap, their conformations are

incompatible with each other, raising fundamental questions about their potential relation to the

inactive and active forms of STIM1. What is the conformation of CAD in the context of the

entire cytosolic domain? How is CC1 arranged relative to CAD in the inactive state, and how

does this change during activation? Which regions of STIM1 are stable, and which are flexible?

To address these questions, we examined the structure and dynamics of the full cytosolic domain

of STIM1 (ctSTIM1, aa 233-685) using single-molecule Förster resonance energy transfer

(smFRET) measurements at sites throughout CC1 and CAD. smFRET is commonly applied to

estimate intramolecular distances and reveal asynchronous molecular transitions that are

typically obscured in population measurements. Our results show that CAD in ctSTIM1

resembles the CAD crystal structure, but with unexpected structure and flexibility in an apical

region that is critical for Orai binding and activation. CC1α1 interacts with CAD in a domain-

swapped configuration to form the inhibitory clamp, enabled by the arrangement of the CC1α2

and CC1α3 domains. The Stormorken mutation R304W activates ctSTIM1 by releasing CC1α1

from CAD. In studies of full-length STIM1 in intact cells, state-dependent cysteine crosslinking

after store depletion suggests that after unbinding from CAD, the two CC1 domains pack

together along their entire length, in contrast to the CC1 crystal and CC1α3-CC2 NMR

structures. Together, these findings offer the first structural view of the STIM1 inhibitory clamp

and reveal the massive conformational changes evoked by store depletion, which serve to

reorient and translocate CAD towards the PM where it can bind Orai1 and activate SOCE.

Results

The CC2 domains in ctSTIM1 are oriented in a parallel configuration

We initially applied smFRET to test whether the conformation of CAD within the cytosolic

domain (ctSTIM1, aa 233-685) resembles that of the CC2-CC3 crystal structure (Yang et al.,

2012) or the CC1α3-CC2 NMR structure (Stathopulos et al., 2013), referred to below as the

'crystal structure' or 'NMR structure,' respectively. A major distinction between the two

structures is that the CC2 domains (aa 345-391) in CAD are parallel in the crystal structure but

antiparallel in the NMR structure (Figure 1A). After removing the single native cysteine in

ctSTIM1 with a C437S mutation, we created three mutants with cysteines at positions 337, 363,

and 378, for which the two structures predicted very different inter-subunit distances, particularly

for 337:337′ and 378:378′ (hereafter ':' is used to denote a pair of sites in the dimer and '′'

indicates a site on the adjacent subunit) (Figure 1A). The ctSTIM1 mutants were expressed in *E.*

coli and purified, yielding full-length protein of >90% purity (see Methods). After randomly

labeling the cysteine-mutant ctSTIM1 molecules with FRET donor (Alexa Fluor 555) and

acceptor (Alexa Fluor 647) dyes, single ctSTIM1 dimers were surface-tethered to a glass

coverslip for imaging by TIRF microscopy, either encapsulated in liposomes or attached directly

through a tag on the C-terminus (Figure 1B; the attachment method for each measurement is

shown in Supplementary Figure 1). Upon excitation of the donor dye, the donor and acceptor

emissions of individual molecules were ratioed to calculate FRET efficiencies, which were high

and stable at all three label sites (Figure 1C, D). These results are consistent with a CAD

conformation in which the CC2 domains are closely apposed and parallel, as in the crystal

structure. Inter-fluorophore distances calculated from simulations of the donor and acceptor dye

positions on the crystal structure (see Methods) closely matched those derived from smFRET,

while simulations based on the NMR structure deviated substantially (Figure 1E). We conclude

that in the context of the complete ctSTIM1 in solution, the CC2 domains in CAD adopt a

parallel conformation consistent with the CAD crystal structure and not the NMR structure.

The apex of CAD deviates from the crystal structure

We made additional smFRET measurements throughout the entire CAD to allow a more

complete comparison with the crystal structure. Near the C-terminal end of the CC3 domain (aa

408-437), the 431:431′ label pair produced a stable, narrowly defined high-FRET state (Figure

2A), and inter-subunit smFRET throughout the CAD exceeded 0.5, consistent with a compact

parallel dimer (Figure 2B). Near the dimer interface (hereafter termed the 'base' of CAD),

smFRET-derived distances quantitatively agreed with those from the crystal structure, and an

intra-subunit measurement 431:363 directly confirmed the compact folding of CC3 onto CC2

within each subunit (Figure 2C).

In contrast to the base of CAD, smFRET measurements of the distal CC2 and CC2-CC3

linker regions (the CAD 'apex') deviated from those indicated in the crystal structure (Figure

2D). This is particularly interesting as mutagenesis studies have identified this region as critical

for Orai1 binding and activation (Calloway et al., 2010; Korzeniowski et al., 2010; Thompson et

al., 2018; Butorac et al., 2019; Wang et al., 2014). To provide a rough indication of the extent of

the deviation implied by smFRET, we generated an alternative model of the CAD apex based on

the crystal structure by allowing rotation around G379 to bring the simulated FRET efficiencies

in line with our experimental results. In the resulting model the distal CC2 regions were rotated

outward, while the regions around aa 400 were pulled closer together, effectively compacting the

apical 'V-shape' (Figure 2E and see Methods). Whereas inter-subunit smFRET at aa 399-401

appeared as a well-defined peak, FRET levels at the distal end of CC2 were broadly distributed

and appeared multimodal, particularly at aa 388 (Figure 2B). Accordingly, smFRET

measurements in this region displayed occasional fluctuations (Figure 2F), suggesting

conformational flexibility that was not expected from the uninterrupted rigid helix of the crystal

structure. The smFRET-derived model in Figure 2E thus likely represents just one of a number

of potential apical conformations sampled by spontaneous fluctuations of the distal CC2 domain.

As a complementary structural test, we analyzed inter-subunit cysteine-cysteine (cys-cys)

crosslinking upon oxidation by copper phenanthroline (CuP, see Methods). Cysteines introduced

throughout the CAD apex formed inter-subunit crosslinks, detected as ctSTIM1 dimers on non-

reducing SDS-PAGE gels (Supplementary Figure 2A). This result shows that opposing residues

in the apical region at least transiently come within the ~6 Å range for cys-cys crosslinking (Qin

et al., 2015), again inconsistent with the CAD crystal structure. Moreover, crosslinking

efficiencies were similar for three adjacent apical cysteine pairs (399:399′, 400:400′, and

401:401′), suggesting significant rotational flexibility beyond the temporal resolution of our

smFRET recordings. Taken together, these data show that flexibility and spontaneous

fluctuations cause the CAD apical region to deviate significantly from the crystal structure,

unlike the stable basal region.

CC1α1 docks parallel to CC3 in a domain-swapped configuration

The CC1 domain controls the activation of STIM1 *in vivo* by releasing CAD for interaction with

Orai1 (Zhou et al., 2013; Ma et al., 2015). Although it is clear that the membrane-proximal

CC1α1 domain can interact with CAD to inhibit STIM1 (Ma et al., 2015), their relative

orientations remain undefined. We detected high inter-subunit FRET efficiencies between labels

at the N-termini of CC1α1 and CC3 (red histogram in Figure 3A, top), as well as between labels

at their C-termini (blue histogram in Figure 3A, bottom), indicating that the CC1α1 domain (a

continuous α-helix (Cui et al., 2013; Rathner et al., 2020)) is oriented parallel to CC3 in the

predominant ctSTIM1 conformation.

To determine whether CC1α1 associates with the CAD of the same subunit or the

adjacent one within the dimer, we compared inter-subunit and intra-subunit FRET efficiencies

between labels at the N-terminus of CC1α1 and three sites in CAD (Figure 3B). The

predominant inter-subunit FRET efficiencies were consistently higher than intra-subunit FRET

efficiencies, demonstrating that in the predominant ctSTIM1 conformation, CC1α1 engages

CAD in a domain-swapped configuration. We also observed spontaneous and reversible

transitions from the predominant FRET state, in which FRET efficiencies fluctuated between the

same two levels for both inter- and intra-subunit measurements (Figure 3C). Similar fluctuation

behavior and transition kinetics were observed at all three CAD sites (Supplementary Figure

3A), suggesting a conformational transition in which the two CC1α1 domains briefly switch

sides on CAD, probably using the same binding interface (Figure 3D).

The CC1α3 domains are compactly folded and directed away from CAD

In vivo, mutations or deletions of the CC1α3 domain activate STIM1 (Korzeniowski et al., 2010;

Yang et al., 2012) but the underlying mechanism, in particular the proposal that CC1α3 interacts

directly with CAD (Yang et al., 2012), has been questioned (Zhou et al., 2013). We measured

predominantly high inter-subunit FRET efficiencies of ~0.9 between labels at the N- and C-

terminal ends of the CC1α3 domains (aa 312 and 337, respectively), indicating that the two

domains are closely apposed and parallel (Figure 4A). To determine their orientation relative to

CAD, we measured intra-subunit FRET efficiencies between labels at either end of the CC1α3

domain to three sites in CC2. While FRET efficiency from the CC1α3 C-terminus to the

proximal CC2 domain was high as expected (red histogram in Figure 4B, top), low FRET

efficiencies were measured from the CC1α3 N-terminus, indicating that the N-terminus is

directed away from CAD (Figure 4B, bottom).

Stable high inter-subunit FRET efficiencies at the CC1α3 C-terminus (aa 337) were

consistent with its proximity to the compact, stable CAD base. However, label sites near the N-

terminus of CC1α3 displayed brief (0.8 s average lifetime), large transitions to a low FRET state,

suggesting that the N-termini of CC1α3 intermittently splay out from a stable pivot point at the

base of CAD (Figure 4C, D). While these large conformational transitions were most common,

further analysis revealed additional substates in the smFRET trajectories (Supplementary Figure

3B, C).

A model for the CC1-CAD complex

Through the integration of many inter-molecular distance measurements, smFRET can be

applied to build models of tertiary protein structure (Brunger et al., 2011). We used 36 unique

inter- and intra-subunit smFRET measurements to develop a structural model of the resting

conformation of CC1 subdomains relative to the optimized model of CAD depicted in Figure 2E.

Each CC1 domain in the ctSTIM1 dimer was modeled as a set of three rigid α-helical domains

connected by flexible linkers, consistent with bioinformatic analysis (Soboloff et al., 2012),

NMR measurements of the CC1 fragment (Rathner et al., 2020), and distances estimated from

intra-subunit smFRET efficiencies (239:274, 274:307, and 307:337; Supplementary Figure 1).

We selected randomly generated models that met the criteria of satisfying smFRET-derived

distance constraints while avoiding steric clashes with each other or with CAD (see Methods and

Supplementary Figure 4A, C for details of the modeling process). Two classes of models

emerged that were highly similar but distinguishable by the organization of the CC1α2/3

domains. In ~70% of solutions the CC1α2/3 domains were stacked parallel (representative

model in Figure 5A from an ensemble of models shown in Supplementary Figure 4Bi), while the

remaining solutions had an alternative, wedged CC1α2/3 topology (Figure 5A and

Supplementary Figure 4Bii). Importantly, both classes of solutions displayed the defining

features described above, including closely apposed CC1α3 domains directed away from CAD

and CC1α1 domains oriented parallel to CC3 in a domain-swapped configuration.

As an independent test of the smFRET-derived models, we applied the lysine

crosslinking agent BS^3 combined with mass spectrometry (XLMS) to identify residues in close

proximity to each other (Rappsilber, 2011). BS^3 XLMS confirmed key structural arrangements

of both models of the CC1-CAD complex, including the compact packing of CC1α2/3 domains

and the domain-swapped CC1α1-CAD interaction (Supplementary Figure 5A-E).

Although the models do not specify the orientation of side chains in the CC1 helices, they

highlight two regions for potential inter-subunit coiled-coil interactions in the CC1-CAD

complex: an antiparallel apposition of heptad repeats in CC1α2 and CC1α3′ (Figure 5B) and a

parallel apposition of heptad repeats in CC1α1 and CC3′ (Figure 5C). The CC1α2/α3′ interface

is reminiscent of the tight, antiparallel interaction of CC1α2 and CC1α3′ domains in the CC1

dimer crystal structure (Cui et al., 2013), an arrangement that was further supported by the

formation of amine-carboxy crosslinks in ctSTIM1 by 1-ethyl-3-(3-

dimethylaminopropyl)carbodiimide hydrochloride (EDC), a zero-length crosslinker (Figure 5D

and Supplementary Figure 6).

Interestingly, the CC1α1:CC3′ interface in the smFRET-derived models juxtaposes

multiple hydrophobic residues that have been previously implicated through mutagenesis in

stabilizing the CC1-CAD complex: L248, L251, L258, L261 in CC1α1 and L416, V419, and

L423 in CC3 (Fahrner et al., 2014; Muik et al., 2011; Zhou et al., 2013; Ma et al., 2015). To

refine the CC1α1:CC3′ interface, we aligned the crystal structure of the CC1α1 domain (aa 246-

271) to the smFRET-derived model and performed a computational docking simulation using

Rosetta software. The resulting model indicates an extensive pairing of hydrophobic residues

predicted to stabilize the CC1α1:CC3′ interface (Figure 5E), providing a plausible explanation

for the activating effects of disruptive mutations at these sites.

The Stormorken mutation R304W activates ctSTIM1 by releasing CC1α1 from CAD

The close packing of CC1α1 and CAD in our model suggests that ctSTIM1 is primarily inactive.

Accordingly, when expressed with Orai1 in HEK293 cells, mCh-ctSTIM1 only slightly

increased $[Ca^{2+}]_i$ compared to mCh-CAD (Supplementary Figure 7), in agreement with previous

evidence that ctSTIM1 is a relatively weak activator of Orai1 (Park et al., 2009; Korzeniowski et

al., 2010; Yuan et al., 2009; Muik et al., 2009).

Deletion of the CC1 domain (mCh-ctSTIM1-ΔCC1) restored ctSTIM1 activity to the

level seen with CAD, as did the introduction of R304W, a naturally occurring mutation that

causes Stormorken syndrome by constitutively activating STIM1 (Misceo et al., 2014; Morin et

al., 2014; Nesin et al., 2014) (Supplementary Figure 7). Interestingly, introduction of R304W

into ctSTIM1 almost completely eliminated the high inter-subunit FRET between CC1α1 sites

and the CC3 N-terminus *in vitro*, indicating the release of CC1α1 from CAD (Figure 6A, left and

center). A more moderate decrease in FRET efficiency was observed relative to the CC3 C-

terminus (aa 431), suggesting that CC1α1 pivoted around the CC1α1/2 linker (Figure 6A, right).

To determine how R304W releases the CC1α1 domain given its distant location at the C-

terminal end of CC1α2, we examined its effect on folding of the CC1α2 and CC1α3 domains.

Intra-subunit smFRET measurements indicated that R304W increases the separation between the

CC1α2 N-terminus and the CC1α3 C-terminus (274:337) (Figure 6A, bottom left), suggesting it

has opened the tight angle between CC1α2 and CC1α3. R304W also shifted CC1α2:CC1α2′ and

CC1α3:CC1α3′ FRET to lower levels (Figure 6B), again consistent with opening of the compact

inactive structure.

The R304W mutation also significantly altered the structural dynamics of CC1α2/α3. In

the wild-type protein, the predominant inter-subunit FRET efficiency at the CC1α3 N-terminus

was high, interrupted by occasional transitions to a low FRET state (Figure 4C and Figure 6C,

left). In contrast, the R304W mutation appeared to destabilize the CC1α3 pair, such that it

transitioned between high and low FRET through a series of intermediates states (Figure 6C,

right). Taken together, our data reveal several effects of R304W on the CC1 domain: it

destabilizes the tight packing of CC1α3 domains, causes CC1α2 to splay out from CC1α3, and

completely releases CC1α1 from CAD.

We also assessed effects of R304W on the structure of CAD. While the R304W mutation

or the deletion of CC1 (ΔCC1) did not affect inter-subunit FRET efficiencies at the base of CAD

(431:431′), they did alter the smFRET distributions at the CAD apex (389:389′ and 400:400′)

(Figure 6D). Together, these results suggest that the association of CC1α1 with CAD influences

the conformation of the flexible apex without affecting the arrangement of helices at the more

stable CAD base.

The CC1 domains refold into a parallel dimer during STIM1 activation

We applied cysteine crosslinking to obtain complementary structural information for the CC1

domains. Inter-subunit cysteine crosslinking in ctSTIM1 was notably robust in the CC1α2/α3

linker (aa 307) and the C-terminus of CC1α3 (aa 339) (Supplementary Figure 8A). Disulfide

formation at T307C prevented BS^3 from crosslinking CC1α1 and CAD, indicating that T307C

crosslinking is incompatible with the CC1α1-CAD inhibitory clamp (Supplementary Figure 5F).

Similar findings were obtained for the activated ctSTIM1 mutants R304W and L248S/L251S

(Supplementary Figure 5F), suggesting that close apposition and crosslinking at T307C resulted

from spontaneous, transient visits to an activated state of ctSTIM1.

Under physiological conditions, full-length STIM1 (flSTIM1) in cells is activated by Ca^{2+}

store depletion. To extend our *in vitro* ctSTIM1 results to flSTIM1, we examined cysteine

crosslinking of inactive and active flSTIM1 *in vivo*. Cysteine mutations in flSTIM1 were made

at selected sites throughout CC1, transiently expressed in HEK293 cells, and treated with the

cell-permeant oxidizer diamide before or after store depletion induced by cyclopiazonic acid

(CPA). Lysates of diamide-treated cells were then analyzed for the presence of crosslinked

flSTIM1 dimers. Prior to store depletion, when flSTIM1 is inactive, little crosslinking was

observed throughout CC1, with the exception of S339C near the junction with CAD (Figure 7A

and 7b). In contrast, after store depletion prominent crosslinking occurred at A268C and T307C

in addition to S339C (Figure 7A, B, Supplementary Figure 8). These results further support our

conclusion above that T307C crosslinking of ctSTIM1 *in vitro* is enabled by transient activation

events. More importantly, they suggest that activation *in vivo* brings all three CC1 subdomains

together.

To test this idea, we asked whether cysteine crosslinking after ER Ca^{2+} store depletion

could lock flSTIM1 in the activated state, thereby preventing deactivation of STIM1 and SOCE

upon store refilling. Ionomycin was added in Ca^{2+}-free Ringer's to deplete stores, after which

ionomycin was removed and Ca^{2+} readded to allow Ca^{2+} influx through SOCE. In HEK cells

expressing WT flSTIM1, $[Ca^{2+}]_i$ increased rapidly upon Ca^{2+} readdition and subsequently

declined due to store refilling and deactivation of STIM1 and Orai1, regardless of diamide

treatment (Figure 7C). We obtained a similar result with flSTIM1-S339C, indicating that

diamide-induced crosslinking at S339C does not impede flSTIM1 deactivation. In contrast,

crosslinking at A268C and T307C prevented the normal deactivation of SOCE as indicated by

the sustained elevation of $[Ca^{2+}]_i$ after Ca^{2+} readdition. These findings suggest that all three CC1

subdomains are brought close together during STIM1 activation *in vivo* and remain closely

associated in the final, Orai1-bound conformation (Figure 7D).

Discussion

Precise regulation of SOCE is essential for many physiological processes and is achieved through control of transitions between inactive and active states of STIM1. These transitions are triggered by changes in ER $[Ca^{2+}]$ that toggle reversible interactions of the STIM1 CC1 domain with CAD. Through a combination of smFRET measurements and crosslinking assays, we have determined the arrangements of CC1 and CAD helices in inactive and active states of STIM1. The results reveal how CC1 is oriented around CAD to stabilize the inactive state, and how its conformation changes upon activation to project CAD toward the PM to engage Orai1 and trigger SOCE.

In this study, we measured FRET efficiencies from single ctSTIM1 dimers to generate an overlapping set of intramolecular distance constraints for modeling tertiary structure (Choi et al., 2010; Brunger et al., 2011). Caution must be applied in using FRET signals to infer absolute distances, as FRET efficiency can also be affected by photophysical transitions or interactions with the local environment (Weiss, 1999, 2000). We employed several strategies to avoid misinterpretation of the data. First, by including a large number of different FRET pairs (36 pairs for CC1-CAD), no major feature of the CC1-CAD model was dependent on any single measurement, safeguarding against potential problems with individual label sites (Supplementary Figure 4A, C). Second, we independently verified the CC1-CAD model by mass spectrometry analysis of BS^3-crosslinked protein (Schmidt and Robinson, 2014) (Supplementary Figure 5A-E). Third, measurements of key smFRET amplitudes and spontaneous structural transitions were confirmed at multiple nearby sites (Figure 2 and Supplementary Figure 3). Finally, we applied relatively conservative distance error limits (± 1 nm) in constraining the CC1-CAD model. These strategies taken together help validate the tertiary structures represented in our models.

The conformation of CAD in the context of ctSTIM1

smFRET measurements demonstrated that CAD within the context of ctSTIM1 resembles the

CAD crystal structure rather than the NMR structure of CC1α3-CC2 (Figure 1). The base is

stable and compact (Figure 2), as expected given the hydrophobic interactions and hydrogen

bonding between the two CAD subunits in this region (Yang et al., 2012). However, the CAD

apex appeared flexible and diverged from the crystal structure (Figure 2). These differences

most likely arose from stabilization of an alternate apical conformation in the crystal by

hydrogen bonds between the apex and adjacent CAD subunits (Supplementary Figure 1B) (Yang

et al., 2012). The flexible structure of the CAD apex we observed is intriguing because

mutagenesis and cysteine-crosslinking studies suggest this region interacts with Orai1 to initiate

channel activation (Calloway et al., 2010; Korzeniowski et al., 2010; Thompson et al., 2018;

Butorac et al., 2019; Wang et al., 2014), and conformational flexibility can facilitate protein-

protein interactions through effects on the thermodynamics of binding (Grünberg et al., 2006).

Thus, it is likely that the CAD apex in complex with Orai adopts a structure different from that

depicted in the CAD crystal and distinct from the NMR structure of CC1α3-CC2 (see below).

Release of CC1α1 from CAD, or deletion of the CC1 domain altogether, did not cause a gross

change in CAD structure, although it did alter the conformation of the CAD apex (Figure 6).

This raises the possibility that upon CC1α1 dissociation the apex changes shape, perhaps

facilitating its binding to Orai1.

Defining structural roles for all three CC1 subdomains in the STIM1 inhibitory clamp

Our results reveal how the three helical regions within CC1 collaborate to regulate the activity of

STIM1. Prior evidence indicated that CC1 autoinhibits STIM1 in resting cells but left

unanswered fundamental questions surrounding its structure. Korzeniowski and colleagues

originally reported that neutralization of acidic residues in the CC1α3 domain activated STIM1

and proposed an 'inhibitory clamp' mechanism in which this region binds to a basic region

within CAD to shield it from Orai in resting cells (Korzeniowski et al., 2010). This hypothesis

potentially explained the activation of STIM1 by deletion of the CC1α3 domain (Yang et al.,

2012). However, subsequent studies showed that CC1α3 alone is not sufficient to maintain the

inactive state, as CC1α3-CAD is fully active (Zhou et al., 2013); moreover, intermolecular FRET

between ER-tethered STIM1 fragments showed that CC1α1 can bind to CC3 (Fahrner et al.,

2014), and ER-tethered CC1α1 by itself can bind CAD and keep it from binding to Orai1 in the

PM (Fahrner et al., 2014; Ma et al., 2015). These findings and the activating effects of mutations

in CC1α1 (Fahrner et al., 2014; Muik et al., 2011; Zhou et al., 2013; Ma et al., 2015) and CC3

(Muik et al., 2011) shifted the focus to a CC1α1-CC3 interaction as the basis for the inhibitory

clamp, and the requirement for CC1α3 has remained unexplained. In addition to defining the

CC1α1-CC3 interaction (below), our smFRET-derived structure can now clarify why deletion of

CC1α3 disables the inhibitory clamp. CC1α2 and CC1α3 are compactly folded and directed

away from CAD, allowing CC1α1 to engage CC3 (Figure 5); in the absence of CC1α3, the rigid

CC1α2 helix would prevent CC1α1 from binding to CAD, which would then be free to interact

with Orai1. Consistent with this explanation, deletion of CC1α2 in addition to CC1α3 allows the

inhibitory clamp to function normally (Fahrner et al., 2014).

Based largely on mutagenesis studies, Zhou and colleagues proposed that the inhibitory

clamp results from antiparallel binding of CC1α1 to CC3, involving interactions between L261

and L416 and between L258 and V419 (Ma et al., 2015). In contrast, our smFRET

measurements show that CC1α1 binds parallel to CC3 (Figure 3). Moreover, our model depicts

interactions of L248 and L251 (CC1α1) with L416 (CC3), and of L258 and L261 (CC1α1) with

L423 and L427 (CC3) (Figure 5), which may explain the activating effects of non-conservative

mutations at these sites (Fahrner et al., 2014; Muik et al., 2011; Zhou et al., 2013; Ma et al.,

2015). Interestingly, our data reveal that CC1α1 preferentially associates with CC3 of the

adjacent subunit in the CAD dimer (Figure 3). Such a domain-swapped configuration may

enhance the stability of the inactive state through inter-subunit interactions and could promote

cooperativity between subunits during structural transitions.

Consistent with a high degree of flexibility, CC1 displayed large spontaneous FRET

fluctuations. The CC1α2/α3 domains occasionally transitioned from a compact, high-FRET

resting state to a low-FRET open state (Figure 4), while the CC1α1 domains briefly switched

sides on CAD, probably trading binding interfaces (Figure 3). Differences in dwell times

suggest these two types of fluctuations occurred independently (Supplementary Figure 3), but

further studies will be needed to determine how they relate to structural changes during

activation of membrane-inserted flSTIM1. During the preparation of this manuscript, a report

appeared describing a solution NMR structure of the isolated CC1 peptide, in which the

CC1α1/α2/α3 helices form a monomeric three-helix bundle (Rathner et al., 2020). Rathner et al

have proposed that this structure represents a pre-activated state of CC1 in which the interaction

of CC1α1 and CC1α2 has destabilized the clamp formed by CC1α1 and CC3. Our smFRET

experiments show that when CC1α1 associates with CC3 in ctSTIM1, the CC1α2 and CC1α3

domains primarily interact with each other to form a compact bundle pointing away from CAD

(Figure 5); in the context of this recent report, we hypothesize that the fluctuations of CC1α2/α3

we observed may represent spontaneous transitions into such a pre-activated state.

The R304W Stormorken mutation activates ctSTIM1 (Supplementary Figure 7),

consistent with its ability to activate flSTIM1 (Fahrner et al., 2018; Nesin et al., 2014; Morin et

al., 2014; Misceo et al., 2014). smFRET measurements show that R304W destabilizes the

compact packing of CC1α2/3 domains, increasing the angle between CC1α2 and CC1α3 helices

and fully releasing CC1α1 from CAD (Figure 6). These results are consistent with NMR studies

showing that R304W causes N-terminal extension of the CC1α3 helix, thereby stiffening the

CC1α2/α3 linker (Fahrner et al., 2018; Rathner et al., 2020).

A structural view of quiescent STIM1 and its transition to the activated state

Our results provide a first view of how the CC1 domain of STIM1 sequesters CAD near the ER

membrane in resting cells to prevent it from interacting with Orai. The parallel arrangement of

CC1α1 and CC3 in our smFRET-derived structure predicts that the CAD apex of inactive STIM1

is juxtaposed to the ER membrane and pointed away from the PM *in vivo* (Figure 8A). This

orientation implies that to activate STIM1 and SOCE, the CAD apex must rotate through a large

angle (up to 180º) and translocate more than 10 nm towards the PM - a massive conformational

change reminiscent of the refolding and extension of viral envelope proteins like hemagglutinin

when stimulated to trigger membrane fusion (Harrison, 2008). Hirve et al. showed that

activation of the flSTIM1 cytosolic domain is initiated by formation of a coiled-coil in the TM

domains and the proximal part of CC1α1 up to L251 (Hirve et al., 2018). As an extension to

these findings, we found that after store depletion, inter-subunit disulfide bonds form at A268C

(C-terminus of CC1α1) and T307C and S339C (N- and C-termini of CC1α3, respectively; Figure

7 and Supplementary Figure 8). Taken together, these results suggest that all three helical

domains in CC1 are closely paired in the active dimer, which would effectively move CAD

across the ~15 nm gap of the ER-PM junction to reach Orai (Wu et al., 2006).

Our CC1-CAD model specifies the possible types of conformational changes that must

occur to activate flSTIM1 and present the CAD apex to Orai (Figure 8B). We can envision two

potential solutions. In a 'flip-out' mechanism, CAD rotates as a whole by 180°. Although our

smFRET data indeed indicate that release of CC1α1 does not cause a gross conformational

change in CAD (Figure 6), this kind of CAD rotation would imply a complex asymmetric

intermediate, in which one of the C-terminal domains (aa 449-685) downstream of CAD would

need to be threaded between the two CC1 domains. Alternatively, in a 'fold-out' model, CAD

undergoes a symmetric conformational change while moving its apex away from the ER

membrane. This model posits that during activation the CC2 domains rotate to transiently

assume an antiparallel state, reminiscent of the NMR structure (Figure 1A and 8B). It is

important to note that our cysteine crosslinking results indicate the CC1α3 C-termini are closely

apposed in the Orai-bound state (Figure 7D, S339C), which is incompatible with the splayed

configuration of CC1α3 domains in the NMR structure. Thus, if the antiparallel conformation

exists it is likely to represent a transient intermediate during the activation of STIM1

(Stathopulos et al., 2013).

The ultimate endpoint in delineating the mechanism of SOCE is an atomic-level

description of the pathway of STIM1 activation by store depletion. To reach this goal the

starting, intermediate, and final conformations of STIM1 after store depletion must be resolved,

as well as the structure of the activated STIM-Orai complex. While approaches like cryo-EM

certainly offer an effective strategy for determining structures, time-resolved measurements such

as those afforded by smFRET will be essential to identify structural transitions in the activation

pathway. The results obtained thus far through smFRET of ctSTIM1 have revealed critical

measurement sites that offer a starting point for these studies. Ultimately, the identification of

transient intermediate STIM1 conformations and rate-limiting steps in the activation process may

reveal critical transitions that could be targeted to manipulate SOCE for therapeutic benefit.

Materials and methods

DNA constructs

For samples that were encapsulated in liposomes for surface immobilization, DNA encoding

ctSTIM1 (aa 233-685) was amplified by PCR from full-length human STIM1 (Origene),

appending an N-terminal NcoI cleavage site, and a C-terminal TEV protease recognition

sequence (SENLYFQG) followed by a HindIII cleavage site. The ctSTIM1 insert contained a

silent T1764C mutation in H588 to remove an endogenous NcoI site, and a G1310C mutation to

replace the endogenous cysteine (C437S). ctSTIM1 inserts were ligated into the pET28a vector

which encoded a C-terminal 6-His tag. ctSTIM1 cysteine mutants were created by site-directed

mutagenesis (QuickChange II, Agilent). The translated ctSTIM1 protein retained the sequence

SENLYFQ at the C terminus after cleavage of the 6-His tag by TEV.

For samples that were directly attached to a PEG-coated surface, the pAC6 vector was

used with a C-terminal AviTag™ sequence GLNDIFEAQKIEWHE (Avidity). ctSTIM1 was

first ligated into a pTEV5 vector which encoded a TEV-cleavable N-terminal 6-His tag. DNA

encoding the 6-His tag, TEV recognition site and ctSTIM1 was then amplified by PCR,

appending an XhoI cleavage site to the N terminus, and a HindIII cleavage site to the C terminus

for ligation into the pAC6 vector. The resulting translated ctSTIM1 protein retained the N-

terminal sequence GAS after TEV cleavage of the 6-His tag, and it had the AviTag™ sequence

following a spacer KLPAGG on the C terminus. All constructs were verified by DNA

sequencing.

Protein expression, purification, and labeling

ctSTIM1 protein was expressed in *E. coli* based on standard methods described previously (Choi

et al., 2010). Briefly, plasmid containing the ctSTIM1 insert was transformed to BL21

competent cells (New England Biolabs #C25271) for pET28a plasmids, or CVB101 competent

cells (Avidity #CVB101) for pAC6 plasmids. Protein expression was induced with IPTG at OD

~0.4 in 450 ml Luria Broth (LB), and for pAC6 plasmids biotin was added to a concentration of

50 µM. Bacteria were spun down and then resuspended in denaturing PBS (8 M urea, 100 mM

NaH_2PO_4, 10 mM Tris (pH 7.4 with NaOH)) and lysed by sonication. TCEP (Thermo Fisher

Scientific #77720) was added to a concentration of 0.5 mM. For intra-subunit smFRET samples,

double-cysteine ctSTIM1 was mixed with cysteine-free ctSTIM1 in a 1:5 ratio to allow

formation of ctSTIM1 heterodimers. Samples were purified by NiNTA purification (Qiagen

#30210), and subsequently slowly dialysed using SnakeSkin™ (Life Technologies #68700) to

20/50 TBS (50 mM NaCl, 20 mM Tris (pH 7.4 with HCl)) containing 1 mM DTT. TEV

protease (MCLAB #TEV-200) was added during dialysis at 10 µg/ml. Cleavage of the 6-His tag

was usually almost 100%, as checked on SDS page gel. Samples were further purified by ion

exchange on a HiTrap Q column (GE Healthcare Life Sciences #17-1153-01) and eluted in 250

µl aliquots of 20/250 TBS. All buffers contained 0.5 mM TCEP. Resulting ctSTIM1 samples

were >90% pure and displayed the expected molecular weight (~52 kD) by SDS-PAGE. Dimer

concentration was typically 25-50 µM, measured by absorption photometry (NanoDrop 2000,

ThermoFisher Scientific).

Protein labeling was performed as described previously (Choi et al., 2010) using Alexa

Fluor 555 as donor fluorophore and Alexa Fluor 647 as acceptor fluorophore (Invitrogen Life

Technologies #A-20346 and #A20347). For symmetric inter-subunit smFRET and for intra-

subunit smFRET, donor and acceptor fluorophores were mixed in a 1:1 ratio before adding to the

protein. For asymmetric inter-subunit smFRET, two different ctSTIM1 cysteine mutants were

mixed separately with donor or acceptor fluorophores, to be recombined afterward into ctSTIM1

heterodimers. Labeling efficiency assessed by absorption photometry was typically ~70% but

varied for different locations; cysteine mutants with labeling efficiency below 50% were not

used for smFRET experiments.

For recombination of asymmetric inter-subunit samples, labeled ctSTIM1 homodimers

were denatured by adding urea to a concentration of 8 M. Donor and acceptor species were then

mixed in a 1:1 ratio, and refolding was allowed to take place during slow dialysis as described

above, resulting in a mixture containing 50% ctSTIM1 heterodimers. ctSTIM1 heterodimers

with a single donor and a single acceptor fluorophore were individually selected during posthoc

analysis of the smFRET experiments.

Sample preparation

Flow cells for single-molecule fluorescence imaging were prepared according to established

protocols (Roy et al., 2008). Briefly, strips of double-sided tape were put on a quartz microscope

slide (Finkenbeiner) to form channel walls, with holes at the ends of each channel to create entry

and exit points for buffer solutions. A microscope coverslip (Erie Scientific #24X40 1.5 001)

was pressed on the tape strips and edges of the channels were sealed with two-component epoxy

glue. For liposome experiments channels were coated with BSA-biotin (Sigma Aldrich

#A8549). For direct surface attachment microscope slides and coverslips were coated with a

100:1 PEG/PEG-biotin mixture (Laysan Bio #BIO-PEG-SVA-5K & MPEG-SVA-5K) prior to

flow cell construction. Neutravidin (Thermo Fisher CWA #31000) was flushed into the channels

before loading the protein samples. 20/150 TBS (150 mM NaCl, 20 mM Tris, pH 7.4 with HCl)

was used for rinsing in between steps.

For liposome encapsulation of fluorophore-labeled protein samples, biotinylated lipids

(18:1 Biotinyl Cap PE, Avanti Polar Lipids #870273C) were mixed with egg-PC lipids (Avanti

Polar Lipids #840051C) in a 1:100 ratio. Lipids were suspended in sample buffer and the

mixture was passed back-and-forth (Extruder Set, Avanti Polar Lipids #610023) through a

liposome extrusion membrane (PC Membranes 0.1μm, Avanti Polar Lipids #610005-1EA).

Liposomes were separated from free protein on a Sepharose® CL-4B column (Sigma Aldrich

#CL4B200). The conditions resulted in on average <<1 ctSTIM1 dimer per liposome.

For direct surface attachment, biotinylated ctSTIM1 samples were diluted to a

concentration of ~100 pM and flushed into a PEG-coated, neutravidin-treated channel.

Immediately prior to fluorescence imaging, channels were filled with 20/150 TBS containing

100 μM cyclooctatetraene (Sigma Aldrich #138924) (COT) and an oxygen scavenging system

consisting of 1% D-glucose, 1 mg/ml glucose oxidase (Sigma Aldrich #G2133), and 0.04 mg/ml

catalase (Sigma Aldrich #C9322).

TIRF microscopy and smFRET measurements

Single-molecule FRET recordings were performed using a home-built through-the-objective

total internal reflection fluorescence (TIRF) imaging system based on a Zeiss Axiovert S100 TV

microscope. Alexa Fluor 555 (donor) and Alexa Fluor 647 (acceptor) fluorophores were excited

by 532 and 637 nm wavelength lasers (OBIS 532 nm LS 150 mW, Coherent #1280719 and

OBIS 637 nm LX 140 mW, Coherent #1196626). A Fluar 100x 1.45 NA oil-immersion

objective (Zeiss) was used to establish TIR excitation and collect photon emission. Donor and

acceptor emission were separated by a dichroic beamsplitter (Semrock FF652-Di01-25x36) and

passed through band-pass filters (Semrock FF01-580/60-25-D for donor emission and Semrock

FF01-731/137-25 for acceptor emission) mounted in an OptoSplit-II (Cairn Research), projecting

the images side-by-side onto an EM-CCD camera (iXon DU897E, Andor). Hardware and image

acquisition were controlled by BeanShell scripts from μManager (Edelstein et al., 2010), and

image sequences were stored as stacked TIFF files.

Prior to image acquisition, the channel surface was scanned manually using low-intensity

donor excitation to identify a suitable location for smFRET imaging, where the density of

fluorescent spots appeared homogeneous and without contamination or aggregations. Donor and

acceptor emission were then recorded in response to donor excitation, typically for 60 s,

immediately followed by 1 s of acceptor excitation to directly identify acceptor fluorophores.

Images were acquired in frame-transfer mode with a 100-ms integration time, and laser intensity

was set to obtain a trade-off between good signal-to-noise ratio and sufficient active time for

both fluorophores (ideally tens of seconds), while inducing fluorophore bleaching within the

recording time window. Typically, 10-20 movies were recorded in each of four channels per

microscope slide. Measurements were performed at room temperature.

smFRET data analysis

Fluorescence movies were analyzed using custom Python scripts. Donor and acceptor images

were aligned using pre-recorded registration images, and fluorescing molecules were identified

in the donor and acceptor channels by detecting local signal maxima within a five-pixel-diameter

neighborhood. Donor signals that could not be matched to any acceptor signal were discarded.

Pixel values for each molecule were summed, and signals from sequential movie frames were

stacked to construct the fluorescence time series for each molecule. The local background for

each molecule was determined by calculating the median background value within a circular 35-

pixel-diameter region in each frame and subtracting it from the raw fluorescence signal.

Leakage of donor signal into the acceptor channel in our system was approximately seven

percent of the measured donor signal and was systematically subtracted from the measured

acceptor signal.

γ correction was used to correct for differences in photon emission and detection

probabilities of donor and acceptor fluorophores. The value of γ was empirically determined for

each individual molecule (McCann et al., 2010), and the correction was applied by multiplying

the donor fluorescence signal by γ before calculating the smFRET ratio. The average value from

all 12,252 molecules was $\gamma = 1.1$.

Traces were selected for final analysis if all of the following criteria were met: (1) the

signal-to-noise ratio was equal or greater than five; (2) the acceptor bleached in a single step

before the donor bleached; (3) the γ factor was between 0.5 and 2.5; (4) any spontaneous

fluctuations of donor and acceptor fluorescence were negatively correlated; and (5) if a donor

bleach event was recorded, it occurred in a single step.

The FRET ratio E was calculated for each point in the donor and acceptor fluorescence

traces (respectively I_D and I_A) where donor and acceptor were both active as $E = I_A/(I_A + \gamma I_D)$. A

FRET histogram was constructed for each trace by distributing FRET amplitudes into 50 bins in

a [-0.25, 1.25] range. Histograms were normalized by dividing each bin by the total number of

FRET points in the trace. An ensemble histogram for multiple molecules was constructed by

summing the normalized histograms of the individual traces and dividing each bin by the total

number of molecules.

The part of the fluorescence traces where both donor and acceptor were active were

parameterized by fitting a Hidden Markov model (HMM), using routines from the SMART

software package (Greenfeld et al., 2012). To determine the optimal kinetic model, the Bayesian

information criterium (BIC) was used to select between models with different numbers of states.

To compare FRET levels between different molecules in an ensemble, each HMM state was

assigned to a cluster based on its mean FRET value. Clustering was performed based on all

observed states in the ensemble, using a k-means algorithm with a predetermined number of

target clusters. The number of clusters was determined ad hoc depending on the shape of the

FRET histogram. For each cluster the peak of its histogram was reported as its representative

FRET level. The predominant FRET cluster was the cluster containing most FRET measurement

points.

Periods between cluster transitions were used for constructing dwell time histograms.

The first and last period of each trace were discarded, so that traces with fewer than two

transitions were not included in the dwell time analysis. For transition density plots each

transition was described by a coordinate given by the FRET level just before the transition

(horizontal axis) and the level after the transition (vertical axis). The local transition density was

then calculated on a 50x50 grid ranging from FRET -0.25 to 1.25, by convolution with a

Gaussian kernel with two-pixel standard deviation. To prevent individual molecules with many

transitions from dominating the plot, each molecule contributed only one data point for each kind

of transition. In this way the transition plot reports the number of molecules in which a certain

kind of transition appeared.

For each protein sample, single-molecule recordings from multiple experiments were

pooled to construct smFRET amplitude histograms. The total number of molecules sampled for

each histogram (see Supplementary Figure 1) was determined such that additional molecules did

not alter the position of the predominant peak in the histogram. Key smFRET results were

replicated by multiple authors (biological replications with newly purified protein samples).

Molecules that did not produce smFRET were excluded, as they could not be properly

interpreted.

Molecular modeling and simulation

Molecular dynamics simulations and model optimization procedures were adapted from

published procedures (Choi et al., 2010), and performed using Crystallography and NMR System

(CNS) (Brunger et al., 1998; Brunger, 2007). Structures were visualized and manipulated using

PyMOL (Version 1.8, Schrödinger, LLC.) and ProDy (Bakan et al., 2011). To predict distances

between fluorophores, we performed molecular dynamics simulations using atomic models of

donor and acceptor fluorophores. Because the structures of Alexa Fluor 555 and Alexa Fluor

647 are not available, we used the structures of the spectrally similar Cy3 and Cy5. Residues of

interest on the STIM structure were mutated to cysteine and fluorophore models were covalently

attached by CNS. While all protein atoms were held fixed in space, a simulated-annealing

protocol sampled possible fluorophore conformations until a local energy minimum was reached.

This procedure was repeated 100 times to obtain a distribution of fluorophore orientations, and

the average location of the fluorophore center (taken as the location of the CAO atom) was

computed from the resulting coordinates. When attached to sites in the CAD crystal structure the

overall mean dye center protruded from the Cα atom of the attachment residue by 0.99 ± 0.08 nm

for Cy3 and 1.10 ± 0.08 nm for Cy5. Since labeling was stochastic, the donor or acceptor could

be on either side in a pair of label sites. To account for this, we averaged Cy3 and Cy5 center

coordinates for each residue into a single "effective" center location. Fluorophores were

represented in the simulations by a pseudo-atom positioned at this effective center location, and

inter-fluorophore distances were then calculated as the distance between two pseudo-atoms.

According to Förster theory the distance between donor and acceptor fluorophores (R)

can be derived from experimentally determined smFRET efficiency (E) as $R = R_0 (1/E - 1)^{1/6}$,

where R_0 is a proportionality factor reflecting the inter-fluorophore distance at which $E = 0.5$ (ref

(Stryer and Haugland, 1967)). For Alexa Fluor 555 and Alexa Fluor 647 in water, the theoretical

value of R_0 is 5.1 nm, assuming fast isotropic re-orientation of the fluorophores. We used this

value for calculating smFRET-derived distances.

For structural modeling of the CAD apex, we first created a symmetric version of the

CAD crystal structure, as the original is slightly asymmetric in the apical region. We copied one

subunit and aligned the proximal CC2 (aa 345-378) and CC3 (aa 408-436) domains to the

adjacent subunit in the dimer, resulting in a symmetric pair. We then used CNS to allow the

apex to relax to a new conformation, guided by smFRET-derived distance constraints (inter-

subunit constraints at aa 388, 389, 399, 400, and 401, and intra-subunit constraint 431:389 on

both subunits). Helices were treated as rigid bodies, while unstructured regions were treated as

flexible chains with variable torsion angles. The distal CC2 helix (aa 379-391) was allowed to

rotate around residue G379. During the relaxation of the apex, the proximal CC2 and CC3

domains were held fixed in space.

To reconstruct the arrangement of CC1 helices around CAD, we used the symmetric

CAD structure with smFRET-optimized apical region and modeled CC1 as a chain of

dimensionless nodes using custom Python code. Each node on the chain represented the center

of a CC1 residue. CC1 chains were generated starting from aa 344 at the CAD N-terminus, by

progressively connecting new nodes until the final residue aa 233 was reached. In unstructured

regions, nodes were spaced every 0.38 nm and were generated at random angles $\geq 90^\circ$. Alpha-

helical regions were modeled as a straight chain of nodes at 0.15 nm intervals. To create a

symmetric pair of CC1 chains, nodes were generated for one subunit, and then copied and

mirrored in CAD's symmetry planes to create the chain for the other subunit. Each node was

subjected to two criteria: (1) it cannot overlap with any residues in CC1 or CAD (volume

exclusion in a 0.5 nm-radius sphere), and (2) it must obey smFRET-derived distance (D_{FRET})

constraints. Fluorophores positions were not explicitly modeled on the CC1 chains but were

accounted for by setting CC1:CC1′ distance constraints to the range of D_{FRET} to D_{FRET} - 2 nm,

due to fluorophores extending up to 1 nm on both sides of the helix backbone. Because

fluorophores were explicitly modeled in CAD, CC1:CAD distance constraints of D_{FRET} to D_{FRET}

- 1 nm were applied. In cases of very high or very low FRET the margins were occasionally

relaxed to account for the lack of sensitivity of the smFRET method at these extremes (see

Supplementary Figure 4A for a table of all distance constraints used for the CC1:CAD model).

If both criteria were met, the node was retained; otherwise it was rejected, and a new node was

generated. If no suitable node was found, a new search was started for the preceding node. This

process continued until the final N-terminal residue of CC1 (aa 233) was reached, and the

resulting model was then stored. Solutions obtained in this way together defined a solution space

within which CC1 obeyed the imposed constraints. We used the average of 50 solutions as a

model for presentation (Figure 5 and Supplementary Figure 4).

670   To obtain a more detailed picture for the molecular interface of the CC1α1-CAD

complex, we aligned the crystal structure of the CC1α1 helical segment (aa 246-271) to the

smFRET-derived CC1 model, directing the hydrophobic sidechains to point toward CAD.

Residue A419 in the CAD crystal structure was mutated back to the original valine using Pymol.

The CC1α1 helix was then docked onto CC3 using the flexpepdock server of Rosetta (London et

al., 2011), which performed a molecular dynamics simulation with a fully flexible backbone. An

initial docking was performed with distance constraints (2.5 ± 2.5 Å) between CG atoms of

residue pairs L248:L416, L251:L416, L258:L423, and L261:L427, which were directly apposed

in the smFRET-derived model. This was followed by an unconstrained simulation to obtain the

final model (Figure 5E).

Chemical crosslinking and mass spectrometry

For BS^3 crosslinking of ctSTIM1, buffer was changed from Tris- to HEPES-based (150 mM

NaCl, 20 mM HEPES (pH 8.0 with NaOH)) during the ion exchange procedure. ctSTIM1

samples at 10 μM (~1 mg/ml) were allowed to incubate with 0, 20 and 50 μM BS^3 (BS^3-d0,

Sigma Aldrich #21590 or BS^3-d4, Sigma Aldrich #21595) for 30 min at room temperature. The

reaction was then quenched for 15 min at room temperature with 20 mM Tris.

For crosslinking with EDC, the buffer was changed from Tris- to MES-based (250 mM

NaCl, 100 mM MES, pH 6.0 with HCl) during the ion exchange procedure. To increase

crosslinking efficiency, EDC (Sigma Aldrich #77149) was used in combination with sulfo-NHS

(Sigma Aldrich #24520) to convert carboxyl groups into stable amine-reactive sulfo-NHS esters.

ctSTIM1 samples at 100 μM (~10 mg/ml) were allowed to incubate with 0/0, 20/62, 50/156 and

100/312 mM EDC/sulfo-NHS for 15 min at room temperature. Two volumes of PBS (250 mM

NaCl, 100 mM Na$_2$HPO$_4$, pH 8.5 with NaOH) with 25 mM BME (pH 8.5) were then slowly

added to quench the EDC and increase the pH for the subsequent amine reaction, incubated 2 h

at room temperature and quenched with 40 mM Tris.

Crosslinked ctSTIM1 samples were run on SDS-PAGE gels, and monomer and dimer

bands were cut and analyzed separately by mass spectrometry. For comparative BS^3

crosslinking experiments, BS^3-d0- and BS^3-d4-treated samples were first mixed 1:1 before

electrophoresis. Gel bands at 20 and 50 μM BS^3 were combined for analysis (Supplementary

Figure 5). For EDC experiments, the monomer band at 10 mM EDC and the combined dimer

bands at 2, 5, and 10 mM were used (Supplementary Figure 6).

Mass spectrometry analysis of cross-linked samples was performed as described

previously (Komolov et al., 2017). Briefly, gel bands were diced into 1x1 mm squares, reduced

in 5 mM DTT at 55ºC for 30 min, and then alkylated with 10 mM acrylamide for 30 min at room

temperature to cap uncrosslinked cysteines. Following alkylation of free cysteines and washing

of gel pieces, proteolytic digestion was completed using trypsin/LysC protease (Promega) in the

presence of ProteaseMax (Promega) overnight at 37ºC. Peptides were then extracted and dried

under SpeedVac prior to LC/MS analysis.

Mass spectrometry experiments were performed using an Orbitrap Fusion Tribrid mass

spectrometer (Thermo Scientific, San Jose, CA) with an Acquity M-Class UPLC system (Waters

Corporation, Milford, MA) for reverse phase separations. The column was an in-house pulled-

and-packed fused silica column with an I.D. of 100 microns pulled to a nanospray emitter.

Packing material for the column was C18 reprosil Pur 1.8 micron stationary phase (Dr. Maisch)

with a total length of ~25 cm. The UPLC system was set to a flow rate of 300 nL/min, where

mobile phase A was 0.2% formic acid in water and mobile phase B was 0.2% formic acid in

acetonitrile. Gel-extracted peptides were directly injected onto the column with a gradient of 2-

45% mobile phase B, followed by a high-B wash over a total 90 minutes. The mass

spectrometer was operated in a data-dependent mode using HCD/ETD decision-tree

fragmentation in the orbitrap (HCD) or ion trap (ETD) for MS/MS spectral generation.

The collected mass spectra were analyzed using Byonic v 2.12.0 or v2.14.27 (Protein

Metrics) as the search engine for peptide ID and protein inference. The precursor and fragment

ion tolerance were both set to 12 ppm and 0.4 Da for HCD and ETD, respectively. The search

assumed tryptic digestion and allowed for up to two missed cleavages. Potential crosslinked

peptides were identified using Byonic's xlink feature, allowing for linker mass where

appropriate. Data were validated using the standard reverse-decoy technique at a 1% false

discovery rate, and inspected using Byologic for crosslink verification and assessment.

Cysteine crosslinking

Symmetric inter-subunit ctSTIM1 dimers (5 μM) were incubated with 300 μM $CuSO_4$ and 900

μM o-phenanthroline for 15 min at room temperature, after which the reaction was quenched

with 2x LDS buffer containing 50 mM EDTA. After samples were run on an SDS-PAGE gel

and stained with Coomassie Blue, gels were destained and scanned with a LI-COR Odyssey

infrared imaging system. Crosslinking efficiency was quantified by the integrated intensity of

the dimer band as a fraction of the summed intensities of the monomer and dimer bands.

For crosslinking flSTIM1 *in situ*, HEK293 cells were transfected with mCh-STIM1-

C437S constructs with selected residues replaced by cysteine. 48 h after transfection, cells were

exposed to 0.2 mM diamide for 10 min in either 2 Ca Ringer's (full Ca^{2+} stores) or in 0 Ca^{2+} + 1

mM EGTA + 20 μM cyclopiazonic acid (depleted Ca^{2+} stores; CPA, Sigma). Cells were then

lysed in RIPA containing 20 mM NEM, protease inhibitor cocktail (Cell Signaling Technology,

#5871S) and EDTA. Samples were run on SDS-PAGE and analyzed by Western blot using

mouse anti-mCherry primary antibody (1:2000, Takara Bio #632543) and Licor secondary

antibody (Licor #926032212) on a LI-COR Odyssey imaging system. 50 mM DTT was added to

duplicate samples to test for disulfide crosslinks. Sites of interest were replicated 2-3 times

(biological replications with newly transfected cells).

Calcium imaging

For experiments with STIM1 fragments (Supplementary Figure 7), HEK293 cells were

transfected with mCherry-labeled STIM1 cytosolic fragments (0.25 µg + 0.25 µg empty

pcDNA3 vector) and Orai1-GFP (0.5 µg) in a 35-mm dish using lipofectamine2000. Transfected

cells were grown overnight in DMEM supplemented with Glutamax, 10% FBS, pen/strep, and

20 µM LaCl$_3$ to minimize the toxic effects of constitutive Ca^{2+} influx. Cells were loaded with 1

µg/ml fura-2/AM for 30 min at room temperature, rinsed, and plated on poly-D-lysine-coated

coverslips for imaging.

For experiments with flSTIM1, HEK293 cells were transfected with flSTIM1 cysteine

mutants with Orai1-GFP. 24-48 h after transfection, cells were loaded with 1 µM fura-2/AM for

30 min at room temperature, rinsed, and plated on poly-D-lysine treated coverslip chambers.

Fura-2 imaging was conducted using a Zeiss 200M inverted microscope with a Fluar 40X

NA 1.3 objective; cells were excited alternately at 350 and 380 nm (Polychrome II, TILL

Photonics) and emission at 534 ± 30 nm (FF02-534/30-25, Semrock) or >480 nm (Chroma) was

collected with a Flash4.0 sCMOS camera (Hamamatsu Corp.) with 2x2 binning. Images were

background corrected before calculation of the 350/380 ratio for each individual cell. Standard

bath solution contained (in mM): 155 NaCl, 4.5 KCl, 2 CaCl$_2$, 1 MgCl$_2$, 10 D-glucose and 5 Na-

HEPES (pH 7.4). Ca^{2+}-free solution was prepared by replacing CaCl$_2$ with 2 mM MgCl$_2$ and 1

mM EGTA. For cysteine crosslinking experiments, cells were treated with 1 μM ionomycin +

0.25 mM diamide in Ca^{2+}-free Ringer's followed by Ca^{2+}-free Ringer's + 0.25 mM diamide +

0.1% BSA to wash out ionomycin prior to readding standard (2 mM Ca^{2+}) Ringer's + 0.1% BSA.

Calcium imaging was performed on a sufficient number of cells to obtain an error of the mean

that allowed a clear distinction among responses to different conditions. Cells with an

abnormally high or unstable baseline calcium level were excluded.

Acknowledgements

The authors thank members of the Lewis lab for helpful discussions during the course of this project. This work was supported by NIH grant R37GM45374, the Mathers Charitable Foundation, and a Stanford Medicine Discovery Innovation Award (R.S.L.), and by NIH grant R37MH63105 (A.T.B.). S.v.D. was supported by Rubicon postdoctoral fellowship 825.13.027 from the Dutch Research Council NWO, and postdoctoral fellowship 16POST30780015 from the American Heart Association. Mass spectrometry was performed in collaboration with the Vincent Coates Foundation Mass Spectrometry Laboratory, Stanford University Mass Spectrometry. We thank R. Leib and K. Singhal for mass spectrometry analysis of crosslinked STIM1. The work was supported in part by NIH P30 CA124435 utilizing the Stanford Cancer Institute Proteomics/Mass Spectrometry Shared Resource.

Competing interests

The authors declare no competing interests.

References

Bakan, A., L.M. Meireles, and I. Bahar. 2011. ProDy: Protein dynamics inferred from theory and

experiments. *Bioinformatics*. 27:1575–1577. doi:10.1093/bioinformatics/btr168.

Böhm, J., and J. Laporte. 2018. Gain-of-function mutations in STIM1 and ORAI1 causing

tubular aggregate myopathy and Stormorken syndrome. *Cell Calcium*. 76:1–9.

doi:10.1016/j.ceca.2018.07.008.

Brunger, A.T. 2007. Version 1.2 of the Crystallography and NMR system. *Nature protocols*.

2:2728–33. doi:10.1038/nprot.2007.406.

Brunger, A.T., P.D. Adams, G.M. Clore, W.L. Delano, P. Gros, R.W. Grossekunstleve, J.S.

Jiang, J. Kuszewski, M. Nilges, N.S. Pannu, R.J. Read, L.M. Rice, T. Simonson, and G.L.

Warren. 1998. Crystallography & NMR system: A new software suite for macromolecular

structure determination. *Acta Crystallographica Section D: Biological Crystallography*.

54:905–921. doi:10.1107/S0907444998003254.

Brunger, A.T., P. Strop, M. Vrljic, S. Chu, and K.R. Weninger. 2011. Three-dimensional

molecular modeling with single molecule FRET. *J Struct Biol*. 173:497–505.

doi:10.1016/j.jsb.2010.09.004.

Butorac, C., M. Muik, I. Derler, M. Stadlbauer, V. Lunz, A. Krizova, S. Lindinger, R. Schober, I.

Frischauf, R. Bhardwaj, M. Hediger, K. Groschner, and C. Romanin. 2019. A novel

STIM1-Orai1 gating interface essential for CRAC channel activation. *Cell Calcium*. 79:57–

67. doi:S0143416019300156.

Calloway, N.T., D.A. Holowka, and B.A. Baird. 2010. A basic sequence in STIM1 promotes

Ca2+ influx by interacting with the C-terminal acidic coiled-coil of Orai1. *Biochemistry*.

49:1067–1071. doi:10.1021/bi901936q.

Choi, U.B., P. Strop, M. Vrljic, S. Chu, A.T. Brunger, and K.R. Weninger. 2010. Single-

molecule FRET-derived model of the synaptotagmin 1-SNARE fusion complex. *Nat Struct*

Mol Biol. 17:318–324. doi:10.1038/nsmb.1763.

Covington, E.D., M.M. Wu, and R.S. Lewis. 2010. Essential role for the CRAC activation

domain in store-dependent oligomerization of STIM1. *Mol Biol Cell*. 21:1897–1907.

doi:10.1091/mbc.E10-02-0145.

Cui, B., X. Yang, S. Li, Z. Lin, Z. Wang, C. Dong, and Y. Shen. 2013. The inhibitory helix

controls the intramolecular conformational switching of the C-terminus of STIM1. *PLoS*

One. 8:e74735. doi:10.1371/journal.pone.0074735.

Edelstein, A., N. Amodaj, K. Hoover, R. Vale, and N. Stuurman. 2010. Computer control of

microscopes using μManager. *Current protocols in molecular biology*. 14:20.1-20.17.

doi:10.1002/0471142727.mb1420s92.

Fahrner, M., M. Muik, R. Schindl, C. Butorac, P. Stathopulos, L. Zheng, I. Jardin, M. Ikura, and

C. Romanin. 2014. A coiled-coil clamp controls both conformation and clustering of

stromal interaction molecule 1 (STIM1). *J Biol Chem*. 289:33231–33244.

doi:10.1074/jbc.M114.610022.

Fahrner, M., M. Stadlbauer, M. Muik, P. Rathner, P. Stathopulos, M. Ikura, N. Müller, and C.

Romanin. 2018. A dual mechanism promotes switching of the Stormorken STIM1 R304W

mutant into the activated state. *Nat Commun*. 9:825. doi:10.1038/s41467-018-03062-w.

Greenfeld, M., D.S. Pavlichin, H. Mabuchi, and D. Herschlag. 2012. Single Molecule Analysis

Research Tool (SMART): an integrated approach for analyzing single molecule data. *PLoS*

One. 7:e30024. doi:10.1371/journal.pone.0030024.

Grünberg, R., M. Nilges, and J. Leckner. 2006. Flexibility and Conformational Entropy in

Protein-Protein Binding. *Structure*. 14:683–693. doi:10.1016/j.str.2006.01.014.

Harrison, S.C. 2008. Viral membrane fusion. *Nature Structural and Molecular Biology*. 15:690–

698. doi:10.1038/nsmb.1456.

Hirve, N., V. Rajanikanth, P.G. Hogan, and A. Gudlur. 2018. Coiled-coil formation conveys a

STIM1 signal from ER lumen to cytoplasm. *Cell Rep*. 22:72–83.

doi:10.1016/j.celrep.2017.12.030.

Kawasaki, T., I. Lange, and S. Feske. 2009. A minimal regulatory domain in the C terminus of

STIM1 binds to and activates ORAI1 CRAC channels. *Biochem Biophys Res Commun*.

385:49–54. doi:10.1016/j.bbrc.2009.05.020.

Komolov, K.E., Y. Du, N.M. Duc, R.M. Betz, J.P.G.L.M. Rodrigues, R.D. Leib, D. Patra, G.

Skiniotis, C.M. Adams, R.O. Dror, K.Y. Chung, B.K. Kobilka, and J.L. Benovic. 2017.

Structural and Functional Analysis of a β2-Adrenergic Receptor Complex with GRK5. *Cell*.

169:407-421.e16. doi:10.1016/j.cell.2017.03.047.

Korzeniowski, M.K., I.M. Manjarrés, P. Varnai, and T. Balla. 2010. Activation of STIM1-Orai1

involves an intramolecular switching mechanism. *Sci Signal*. 3:ra82.

doi:10.1126/scisignal.2001122.

Lacruz, R.S., and S. Feske. 2015. Diseases caused by mutations in ORAI1 and STIM1. *Ann NY*

Acad Sci. 1356:45–79. doi:10.1111/nyas.12938.

Liou, J., M. Fivaz, T. Inoue, and T. Meyer. 2007. Live-cell imaging reveals sequential

oligomerization and local plasma membrane targeting of stromal interaction molecule 1

after Ca^{2+} store depletion. *Proc Natl Acad Sci USA*. 104:9301–9306.

London, N., B. Raveh, E. Cohen, G. Fathi, and O. Schueler-Furman. 2011. Rosetta FlexPepDock

web server - high resolution modeling of peptide-protein interactions. *Nucleic Acids*

Research. 39:249–253. doi:10.1093/nar/gkr431.

Ma, G., M. Wei, L. He, C. Liu, B. Wu, S.L. Zhang, J. Jing, X. Liang, A. Senes, P. Tan, S. Li, A.

Sun, Y. Bi, L. Zhong, H. Si, Y. Shen, M. Li, M.-S. Lee, W. Zhou, J. Wang, Y. Wang, and

Y. Zhou. 2015. Inside-out Ca^{2+} signalling prompted by STIM1 conformational switch. *Nat*

Commun. 6:7826. doi:10.1038/ncomms8826.

McCann, J.J., U.B. Choi, L. Zheng, K. Weninger, and M.E. Bowen. 2010. Optimizing methods

to recover absolute FRET efficiency from immobilized single molecules. *Biophys J*.

99:961–970. doi:10.1016/j.bpj.2010.04.063.

Misceo, D., A. Holmgren, W.E. Louch, P.A. Holme, M. Mizobuchi, R.J. Morales, A.M. De

Paula, A. Stray-Pedersen, R. Lyle, B. Dalhus, G. Christensen, H. Stormorken, G.E.

Tjønnfjord, and E. Frengen. 2014. A Dominant STIM1 Mutation Causes Stormorken

Syndrome. *Human Mutation*. 35:556–564. doi:10.1002/humu.22544.

Morin, G., N.O. Bruechle, A.R. Singh, C. Knopp, G. Jedraszak, M. Elbracht, D. Brémond-

Gignac, K. Hartmann, H. Sevestre, P. Deutz, D. Hérent, P. Nürnberg, B. Roméo, K.

Konrad, M. Mathieu-Dramard, J. Oldenburg, E. Bourges-Petit, Y. Shen, K. Zerres, H.

Ouadid-Ahidouch, and J. Rochette. 2014. Gain-of-Function Mutation in STIM1 (P.R304W)

Is Associated with Stormorken Syndrome. *Hum Mutat*. 35:1221–1232.

doi:10.1002/humu.22621.

Muik, M., M. Fahrner, I. Derler, R. Schindl, J. Bergsmann, I. Frischauf, K. Groschner, and C.

Romanin. 2009. A cytosolic homomerization and a modulatory domain within STIM1 C

terminus determine coupling to ORAI1 channels. *J Biol Chem*. 284:8421–8426.

doi:10.1074/jbc.C800229200.

Muik, M., M. Fahrner, R. Schindl, P. Stathopulos, I. Frischauf, I. Derler, P. Plenk, B. Lackner,

K. Groschner, M. Ikura, and C. Romanin. 2011. STIM1 couples to ORAI1 via an

intramolecular transition into an extended conformation. *EMBO J.* 30:1678–1689.

doi:10.1038/emboj.2011.79.

Nesin, V., G. Wiley, M. Kousi, E.-C. Ong, T. Lehmann, D.J. Nicholl, M. Suri, N. Shahrizaila, N.

Katsanis, P.M. Gaffney, K.J. Wierenga, and L. Tsiokas. 2014. Activating mutations in

STIM1 and ORAI1 cause overlapping syndromes of tubular myopathy and congenital

miosis. *Proc Natl Acad Sci USA*. 111:4197–4202. doi:10.1073/pnas.1312520111.

Park, C.Y., P.J. Hoover, F.M. Mullins, P. Bachhawat, E.D. Covington, S. Raunser, T. Walz, K.C.

Garcia, R.E. Dolmetsch, and R.S. Lewis. 2009. STIM1 clusters and activates CRAC

channels via direct binding of a cytosolic domain to Orai1. *Cell*. 136:876–890.

doi:10.1016/j.cell.2009.02.014.

Prakriya, M., and R.S. Lewis. 2015. Store-operated calcium channels. *Physiol Rev*. 95:1383–

1436. doi:10.1152/physrev.00020.2014.

Qin, M., W. Wang, and D. Thirumalai. 2015. Protein folding guides disulfide bond formation.

Proc Natl Acad Sci USA. 112:11241–11246. doi:10.1073/pnas.1503909112.

Rappsilber, J. 2011. The beginning of a beautiful friendship: cross-linking/mass spectrometry

and modelling of proteins and multi-protein complexes. *J Struct Biol*. 173:530–540.

doi:10.1016/j.jsb.2010.10.014.

Rathner, P., M. Fahrner, L. Cerofolini, H. Grabmayr, F. Horvath, H. Krobath, A. Gupta, E.

Ravera, M. Fragai, M. Bechmann, T. Renger, C. Luchinat, C. Romanin, and N. Müller.

2020. Interhelical interactions within the STIM1 CC1 domain modulate CRAC channel

activation. *Nature Chemical Biology*. doi:10.1038/s41589-020-00672-8.

Roy, R., S. Hohng, and T. Ha. 2008. A practical guide to single-molecule FRET. *Nat Methods*.

5:507–516. doi:10.1038/nmeth.1208.

Schmidt, C., and C. V Robinson. 2014. A comparative cross-linking strategy to probe

conformational changes in protein complexes. *Nat Protoc*. 9:2224–2236.

doi:10.1038/nprot.2014.144.

Schober, R., D. Bonhenry, V. Lunz, J. Zhu, A. Krizova, I. Frischauf, M. Fahrner, M.Q. Zhang, L.

Waldherr, T. Schmidt, I. Derler, P.B. Stathopulos, C. Romanin, R.H. Ettrich, and R.

Schindl. 2019. Sequential activation of STIM1 links Ca^{2+} with luminal domain unfolding.

Sci Signal. 12. doi:10.1126/scisignal.aax3194.

Soboloff, J., B.S. Rothberg, M. Madesh, and D.L. Gill. 2012. STIM proteins: dynamic calcium

signal transducers. *Nat Rev Mol Cell Biol*. 13:549–565. doi:10.1038/nrm3414.

Stathopulos, P., G. Li, M. Plevin, J. Ames, and M. Ikura. 2006. Stored Ca^{2+} depletion-induced

oligomerization of STIM1 via the EF-SAM region: An initiation mechanism for capacitive

Ca^{2+} entry. *J Biol Chem*. 281:35855–35862.

Stathopulos, P.B., R. Schindl, M. Fahrner, L. Zheng, G.M. Gasmi-Seabrook, M. Muik, C.

Romanin, and M. Ikura. 2013. STIM1/Orai1 coiled-coil interplay in the regulation of store-

operated calcium entry. *Nat Commun*. 4:2963. doi:10.1038/ncomms3963.

Stryer, L., and R.P. Haugland. 1967. Energy transfer: a apectroscopic ruler. 58:719–726.

Thompson, J.L., Y. Zhao, P.B. Stathopulos, A. Grossfield, and T.J. Shuttleworth. 2018.

Phosphorylation-mediated structural changes within the SOAR domain of STIM1 enable

specific activation of distinct Orai channels. *J Biol Chem*. 293:3145–3155.

doi:10.1074/jbc.M117.819078.

Wang, X., Y. Wang, Y. Zhou, E. Hendron, S. Mancarella, M.D. Andrake, B.S. Rothberg, J.

Soboloff, and D.L. Gill. 2014. Distinct Orai-coupling domains in STIM1 and STIM2 define

the Orai-activating site. *Nat Commun*. 5:3183. doi:10.1038/ncomms4183.

Weiss, S. 1999. Fluorescence spectroscopy of single biomolecules. *Science*. 283:1676–1683.

doi:10.1126/science.283.5408.1676.

Weiss, S. 2000. Measuring conformational dynamics of biomolecules by single molecule

fluorescence spectroscopy. *Nature Structural Biology*. 7:724–729. doi:10.1038/78941.

Wu, M.M., J. Buchanan, R.M. Luik, and R.S. Lewis. 2006. Ca^{2+} store depletion causes STIM1 to

accumulate in ER regions closely associated with the plasma membrane. *J Cell Biol*.

174:803–813. doi:10.1083/jcb.200604014.

Wu, M.M., E.D. Covington, and R.S. Lewis. 2014. Single-molecule analysis of diffusion and

trapping of STIM1 and Orai1 at endoplasmic reticulum-plasma membrane junctions. *Mol*

Biol Cell. 25:3672–3685. doi:10.1091/mbc.E14-06-1107.

Yang, X., H. Jin, X. Cai, S. Li, and Y. Shen. 2012. Structural and mechanistic insights into the

activation of Stromal interaction molecule 1 (STIM1). *Proc Natl Acad Sci USA*. 109:5657–

5662. doi:10.1073/pnas.1118947109.

Yu, F., L. Sun, S. Hubrack, S. Selvaraj, and K. Machaca. 2013. Intramolecular shielding

maintains the ER Ca^{2+} sensor STIM1 in an inactive conformation. *J Cell Sci*. 126:2401–

2410. doi:10.1242/jcs.117200.

Yuan, J., W. Zeng, M. Dorwart, Y. Choi, P. Worley, and S. Muallem. 2009. SOAR and the

polybasic STIM1 domains gate and regulate Orai channels. *Nat Cell Biol*. 11:337–343.

doi:10.1038/ncb1842.

Zhou, Y., P. Srinivasan, S. Razavi, S. Seymour, P. Meraner, A. Gudlur, P.B. Stathopulos, M.

Ikura, A. Rao, and P.G. Hogan. 2013. Initial activation of STIM1, the regulator of store-

operated calcium entry. *Nat Struct Mol Biol*. 20:973–981. doi:10.1038/nsmb.2625.

Figure 1. Parallel orientation of CC2 domains in ctSTIM1 is consistent with the CAD

crystal structure. (**A**) (*top*) Schematic overview of domains in ctSTIM1, comprising the

membrane-proximal coiled-coil 1 (CC1) domain (with helical regions α1, α2, and α3), and the

CC2 and CC3 domains which are part of CAD. (*bottom*) The orientations of the CC2 domains

are a prominent feature that distinguishes the crystal structure of CAD (*bottom left*, parallel CC2

domains; 3TEQ.pdb) and the NMR structure of CC1α3-CC2 (*bottom right*, anti-parallel CC2

domains; 2MAJ.pdb). Sites tested by inter-subunit smFRET are indicated. Although aa 337 is

not part of the crystal structure (aa 344-443), it is sufficiently close to permit comparison to the

NMR structure. (**B**) Two methods for immobilizing single ctSTIM1 dimers for smFRET.

ctSTIM1 peptides were encapsulated in liposomes and immobilized on a glass slide by a biotin-

neutravidin interaction or were attached directly via a C-terminal avitag. (**C**) Example

recordings of donor (*green*) and acceptor (*red*) fluorescence evoked by donor excitation for sites

337:337′, 363:363′, and 378:378′. Single-step bleaching events indicate a single molecule,

while the anti-correlated response of the donor to the acceptor bleach event is a hallmark of

smFRET. Calculated instantaneous FRET ratios (*black*) are shown below. (**D**) Ensemble

smFRET histograms for sites 337:337′, 363:363′, and 378:378′ displayed predominantly high

FRET, indicating close parallel apposition. (**E**) A comparison of smFRET-derived distances

(*black*) with simulated inter-fluorophore distances from the crystal (*white*) and NMR structures

(*gray*). The measured distances closely match the crystal structure but deviate strongly from the

NMR structure, particularly for 337:337′ and 378:378′. smFRET error bars indicate the

expected distance deviation corresponding to an uncertainty of ±0.05 in FRET measurement.

Simulation error bars indicate the s.e.m. of inter-fluorophore distance from 100 simulations of

dye position (see Methods).

Figure 2. The apex of CAD in ctSTIM1 deviates from the CAD crystal structure. (**A**)

Representative smFRET recording at the base of CAD (431:431´) showing stable, high FRET.

(**B**) Ensemble density plots of the initial 2 s of inter-subunit smFRET recordings for sites

throughout CAD. Predominant FRET levels are shown by arrowheads to the right of each plot.

(**C-D**) Comparison of inter-fluorophore distances from smFRET and the CAD crystal structure.

Simulated fluorophores were represented by the average position of a central atom, indicated

here as ball-on-stick models (see Methods). For sites near the dimer interface (the 'base' of

CAD, **C**), the smFRET-derived distances (*black bars)* agree closely with the crystal structure

(*white bars*). Measurements in the apex (**D**) deviated from the crystal structure. (**E**) Modified

structure (*red*) generated by a smFRET-constrained optimization of the crystal structure (*gray*).

Outward rotation of the distal CC2 region (aa 379-391) around G379 as a pivot point restored

982   close correspondence between smFRET-derived distances and the crystal structure (*bottom*). (**F**)

983   Representative example of smFRET fluctuations in the CAD apex (388:388´).

Figure 3. The CC1α1 domain is near and parallel to the CC3 domain of the opposing CAD

subunit. (**A**) Asymmetric inter-subunit smFRET measurements between CC1α1 and CC3 reveal

a parallel orientation, with high FRET levels at 242:400′ (*top, red*) and 274:431′ (*bottom, blue*)

indicating close apposition. The crystal structure of CAD is shown with schematic

representation of CC1α1 domains. (**B**) The predominant smFRET level between CC1α1 and

CC3 was consistently higher for inter-subunit measurements (*red*) than intra-subunit

measurements (*blue*). This pattern indicates a predominant domain-swapped configuration in

which CC1α1 docks close to the CC3 domain of its opposing subunit. (**C**) Inter- and intra-

subunit recordings (242:431′ and 242:431, respectively) displayed brief spontaneous smFRET

transitions between two states indicated by red and black bars. In both cases fluctuations

occurred between the same two FRET levels, suggesting the CC1α1 domains switched sides on

CAD (see also Supplementary Figure 3A). (**D**) Schematic illustration of spontaneous

conformational transitions of the CC1α1 domain. In the predominant conformation, CC1α1 is

closely apposed and parallel to CC3 on its opposing subunit in a domain-swapped configuration

('trans'). Occasionally, CC1α1 switches to interact briefly with CC3 on its own subunit ('cis').

1000   Rate constants were derived from dwell time analysis of smFRET traces (Supplementary Figure

1001   3A).

1002

Figure 4. The CC1α2 and CC1α3 domains are closely apposed and directed away from

CAD. (**A**) High inter-subunit smFRET at the N- and C-termini of the CC1α3 domain (aa 312

and 337, respectively) indicates close parallel apposition of the two helices. The crystal structure

of CAD is shown with a schematic depiction of CC1α2 and CC1α3 domains. (**B**) (*top*) High

intra-subunit smFRET from aa 337 to sites on CC2 declined progressively with distance, while

intra-subunit smFRET from aa 298 (near the CC1α2/3 linker) to sites in CC2 was low (*bottom*),

indicating that the CC1α2 and α3 domains are directed away from CAD. (**C**) Inter-subunit

recordings near the CC1α3 N-terminus showed brief transitions from the predominant high-

FRET state to a transient low-FRET state. Ensemble histograms show similar main and transient

FRET states for neighboring sites aa 309 and 312. (**D**) Schematic depiction of spontaneous

conformational transitions of the CC1α3 domains. CC1α3 domains are predominantly in close

parallel apposition (*left*) and occasionally transition to a transient open configuration (*right*).

Rate constants were derived from dwell time analysis of smFRET traces (Supplementary Figure

3B).

1017

Figure 5. A model of the configuration of CC1 and CAD in ctSTIM1. (**A**) 36 smFRET-

derived distances were used to reconstruct the orientations of CC1 domains relative to CAD,

yielding two classes of solutions with the CC1α2/3 domains in a 'stacked' (*left*) or 'wedged'

(*right*) configuration. In both classes, CC1α1 domains are in close parallel apposition to the CC3

domains on the adjacent subunit in a domain-swapped configuration, with the CC1α2 and CC1α3

domains forming a compact parallel structure directed away from CAD. The average of 50

model solutions is shown for each class (see Supplementary Figure 4B for individual solutions).

(**B**) smFRET-derived models suggest hydrophobic stabilization of the CC1α2/3 complex by

antiparallel apposition of CC1α2 and CC1α3′. (**C**) The crystal structure of CC1 peptides defines

an antiparallel interaction of CC1α2/α3′ domains (*left*, *dashed box*), in which heptad-like

hydrophobic sequences form a tightly packed dimer interface (*top right*, adapted from

4O9B.pdb). (*bottom right*) Tight antiparallel packing of CC1α2/α3′ in ctSTIM1 was confirmed

by inter-subunit crosslinks with EDC (*solid lines*) and BS^3 (*dashed line*) (see also Supplementary

Figure 6). (**D**) Parallel apposition of hydrophobic residues on CC1α1 and CC3′. Several of

these were previously identified by mutagenesis to stabilize the inactive state of STIM1,

including L248, L251, L416, and L423. (**E**) A model of the CC1α1:CC3′ interface obtained by

computational peptide docking (see Methods). In the lowest-energy solution, the helices engage

in direct mutual hydrophobic interactions.

1036

Figure 6. The Stormorken R304W mutation releases CC1 from CAD. (**A**) The R304W

mutation reduced inter-subunit smFRET between aa 242 on CC1α1 and sites 400′, 417′, and

431′ on CC3′, indicating separation of CC1α1 from CAD (*top*). The mutation also reduced

intra-subunit FRET for 274:337, suggesting a widening of the angle between CC1α2 and CC1α3

(*bottom*). (**B**) The predominant FRET levels for 298:298′ (*left*) and 312:312′ (*right*) were

diminished by R304W, suggesting the compact parallel structure of CC1α2/3 domains was

disrupted in favor of a dilated, low-FRET state. (**C**) Effects of R304W on inter-subunit FRET

fluctuations at 312:312′, near the CC1α3 N-terminus. The dominant large-amplitude

fluctuations of the wild-type (*top left*) were entirely absent in the mutant, which instead

displayed frequent small fluctuations at a range of FRET levels (*top right*). The different

fluctuation modes are summarized in ensemble transition density plots (see Methods) which

show a single dominant FRET transition for wild-type molecules (*bottom left*, 38 molecules), and

a multimodal density distribution for R304W mutants (*bottom right*, 25 molecules). (**D**) In the

1050   CAD apex, R304W or deletion of the entire CC1 region (ctSTIM1-ΔCC1) increased the FRET at

1051   389:389′ (*left*) and reduced it at 400:400′ (*center*), suggesting that the apex interacts with CC1α1

1052   in the WT conformation. Neither R304W nor ΔCC1 affected FRET at 431:431′ (*right*) in the

1053   CAD base.

1054

Figure 7. Close contacts form between CC1 domains along their entire length in the

activated state of STIM1. (A) Full-length STIM1 (flSTIM1) mutants were transiently over-

expressed in HEK293 cells for diamide-induced cysteine crosslinking *in vivo*. Western-blot

analysis of cysteine crosslinking of flSTIM1-T307C (*left*) and flSTIM1-S339C (*right*), under

resting (2 mM Ca^{2+}) or store-depleted (0 mM Ca^{2+} + CPA) conditions. **(B)** Summary of

flSTIM1 cysteine crosslinking before (*black*) and after (*white*) store depletion measured in

individual paired experiments. While crosslinking at aa 268 and aa 307 strongly increased in the

activated state, crosslinking at aa 339 occurred independent of STIM1 activation (see also **A** and

Supplementary Figure 8B). **(C)** WT flSTIM1 and cysteine mutants were co-expressed with

Orai1 for cytosolic calcium imaging (see Methods). For WT flSTIM1 cells, after store depletion

by transient exposure to ionomycin (io), readdition of Ca^{2+} elevated $[Ca^{2+}]_i$ due to SOCE,

followed by a decline as SOCE deactivated from store refilling (*top left, black*). Diamide (*red*)

did not affect the WT response. In contrast, diamide-induced crosslinking of T307C stabilized

the active state, as evidenced by persistent calcium influx after ionomycin wash-out and store

refilling (*bottom left, red*). Similar results were obtained for crosslinking A268C (*bottom right*,

see also Supplementary Figure 8). Crosslinking of S339C did not inhibit deactivation of SOCE

upon store refilling (*top right*). Each trace shows the mean and s.e.m. of the following numbers

of cells (control/diamide) from at least two independent experiments: WT (91/106), S339C

(44/41), T307C (38/46), A268C (31/47). (**D**) Schematic illustration of CC1 cysteine crosslinking

in the resting (*left*) and activated (*right*) states of flSTIM1. In the resting conformation, CC1α1

and CC1α3 domains are kept apart, preventing crosslinking at locations upstream of aa 339.

Upon store depletion, release of the CC1α1 domains from CAD promotes alignment of CC1

domains along their entire length, enabling crosslinking at aa 268 and 307.

1078

Figure 8. Possible conformational trajectories of flSTIM1 activation *in vivo*. (**A**) In the

resting state, the parallel orientation of CC1α1 and CC3 domains implies that CAD is held with

its apex close to and pointed towards the ER membrane, effectively shielding critical interaction

sites (*stars*) from engaging with Orai. The region C-terminal to CAD is abbreviated as its

conformation is not known. (**B**) Alternative models for transitions to the active state of flSTIM1.

In the "fold out" model (*top*), following the release of CC1α1, CAD undergoes a symmetric

conformational change in which the CC2 domains pass through a transient antiparallel state,

providing a free path for the region downstream of CAD to extend towards the plasma

membrane. In the "flip out" model (*bottom*), CAD maintains its parallel resting conformation

and undergoes an asymmetric outward rotation from between the CC1 domains. Assuming that

the initial resting conformation is symmetric, one of its C-terminal domains (*gray*) would have to

be carried through the gap between the two CC1 domains.