HUMAN ADENOVIRUS TYPE 4 COMPRISES TWO MAJOR PHYLOGROUPS WITH

DISTINCT PHENOTYPES RELEVANT TO PATHOGENESIS

Camden R. Bair[a][§], Wei Zhang[b], Gabriel Gonzalez[c], Arash Kamali[b], Daniel Stylos[b], Jorge

C. G. Blanco[b], and Adriana E. Kajon[a][#]

a. Lovelace Biomedical Research Institute (LBRI), Albuquerque, New Mexico, USA

b. Sigmovir Biosystems, Inc. (SBI), Rockville, Maryland, USA

c. UCD National Virus Reference Laboratory, Dublin, Ireland

Running head: Phenotypic differences between phylogroups of HAdV-E4

Address correspondence to Adriana E. Kajon, akajon@lovelacebiomedical.org

§ Present address: Camden R. Bair, Department of Microbiology and Immunology, Emory

University School of Medicine, Atlanta, Georgia, USA.

KEYWORDS: human adenovirus 4, phylogroup, fitness, virulence

Word count for the text: 6939

Word count for the abstract: 150

ABSTRACT

Human adenovirus type 4 (HAdV-E4) is the only type (and serotype) classified within species *Human mastadenovirus E* that has been isolated from a human host to the present. Recent phylogenetic analysis of whole genome sequences of strains representing the spectrum of intratypic genetic diversity described to date identified two major evolutionary lineages designated phylogroups I, and II, and validated the early clustering of HAdV-E4 genomic variants into two major groups by low resolution restriction fragment length polymorphism analysis. Viruses in phylogroups I and II carry distinct coding variations in proteins involved in viral replication and modulation of host responses to infection. They also differ in their growth, spread, and cell killing phenotypes in cell culture and in their pathogenic phenotypes in a cotton rat model of HAdV respiratory infection. This body of work is the first documenting phenotypic correlates of naturally occurring intratypic genetic variability for human adenovirus type 4.

IMPORTANCE

Human adenovirus type 4 (HAdV-E4) is a prevalent causative agent of acute respiratory illness of variable severity and of conjunctivitis. This is the first report of phenotypic divergence among naturally occurring known genetic variants of a HAdV type of medical importance. This research reveals readily detectable phenotypic differences between strains representing phylogroups I and II, and it introduces a unique experimental system for the elucidation of the genetic basis of adenovirus fitness and virulence. These data improve our understanding of the implications of intratypic genetic diversity in the presentation and course of HAdV-E4-associated disease.

INTRODUCTION

Human adenovirus type 4 (HAdV-E4), the only type (and serotype) classified within species *Human mastadenovirus E* (HAdV-E) isolated from humans to the present, was originally recovered from a respiratory specimen obtained from a military recruit in basic training in the winter of 1952-1953 in Fort Leonard Wood, MO, USA during an epidemic outbreak of acute respiratory disease (ARD) (1). The "new" virus was designated RI-67, as it was derived from Respiratory Illness case number 67 in the epidemic (1). The circulation of HAdV-E4 was subsequently documented between 1958 and 1959 in Taiwan in association with cases of conjunctivitis and ARD (2). By the 1980s this unique type had been widely recognized as a prevalent causative agent of respiratory illness of variable severity, and also of conjunctivitis with clinical presentations ranging from mild "pink eye" to severe epidemic keratoconjunctivitits (EKC) (3-11).

The pioneer investigations of intratypic genetic variability for HAdV-E4 conducted in the late 1980s - early 1990s using low resolution restriction fragment length polymorphism (RFLP) analysis demonstrated the existence of two major clusters of genomic variants readily distinguishable by their digestion profiles with BamHI, DraI, EcoRV, and SmaI, among other endonucleases (12, 13). Based on their BamHI digestion profiles, these variants were designated by Li and Wadell (12) with a letter "p" for prototype-like, or a letter "a" or "b" for the non-prototype like variants followed by numbers 1, 2, etc., designating additional variability detectable with other endonucleases. Our recent evaluation of a sizable collection of whole genome sequences (n=47) representing 45 strains isolated worldwide from clinical specimens between 1953 and 2015 (14) and the spectrum of genetic variability described to date by RFLP (11-13, 15) provided a global

view of the magnitude of genetic diversity for this medically important type, confirmed the

early groupings of related genomic variants, and identified two major evolutionary

lineages designated phylogroups (PG) I, and II (14). PG I and PG II comprise the

prototype-like and non-prototype-like genomic variants, respectively.

The two PGs differ significantly in their G + C content and apparent evolutionary rates,

and as we recently reported, they also exhibit marked sequence differences in coding

regions including but not limited to E1A, E1B, E3, L3, and L4 (14).

Interestingly, published data indicate that the representation of PG II strains among

collections of respiratory and ocular clinical specimens sampled from symptomatic

individuals in a variety of settings since 1965 has greatly exceeded that of PG I strains

(11, 15-18) suggesting a higher fitness and/or possibly a higher virulence and/or

transmissibility for these viruses.

To the present, no studies have formally identified genetic determinants of adenovirus

virulence and fitness. Importantly, the pathobiological implications for the intratypic

genetic variability documented for HAdV-E4 and other types remain unknown.

In this study, we expand our original analysis of intra- and inter-PG genetic variability

(14) and use a panel of viruses representative of the spectrum of genetic diversity

described for HAdV-E4 to examine the magnitude of inter- and intra-PG phenotypic

diversity using an array of cell-based assays and a cotton rat model of HAdV respiratory

infection.

Our data show that PG I and PG II display distinct coding variations in proteins involved

in viral replication and modulation of host responses to infection, and that they are

intrinsically different regarding their ability to grow and spread in culture and to cause

pulmonary disease in cotton rats.

RESULTS

Deeper analyses of genomic sequences for HAdV-E4 strains in PG I and PG II reveal

additional genetic differences between and within the two major clades. The use of

new analytical tools to further assess variation across the 45-genome collection originally

characterized by Gonzalez *et al.* (14) revealed genome-wide sequence differences

between strains clustering in PG I and PG II and significant diversity in protein-coding

sequences (Fig 1A). Proteotyping of HAdV-E strains using concatenated protein

sequences in selected coding regions including E1A, E1B 19K and 55K, DNA

polymerase, L4 100K, E3 proteins CR1α, gp19K, CR1β, CR1δ, RIDα, RIDβ and 14.7K,

and E4 34K (Fig 1B) confirmed that the two clades encode distinct variants/proteotypes

at most of these loci. Importantly, in each phylogroup additional divergence was identified

with the occurrence of minor proteotypes; for example, strains USA 2000 (AP014852),

USA 2004 (AP014853) and USA 2011 (EF371058) of PG-I show characteristic mutations

in E2B DNA polymerase, L4 100K, and proteins in the E3 region, reflecting the emergence

and maintained circulation in the USA of a distinct lineage. Intra-phylogroup II variations

were also identified, with multiple small groups of sequences showing characteristic

mutations across the DNA polymerase and E3 proteins.

PG I and PG II strains differ in their average plaque size and cell-to-cell spread

phenotype. A panel of HAdV-E4 strains representative of the spectrum of HAdV-E4

genomic variants were selected for comparison of *in vitro* growth characteristics.

Proteotyping results for coding regions in genome order for this subset of strains are shown in Table 1.

First, we assessed whether the stocks for the selected 7 strains exhibited any detectable differences in specific infectivity. The number of genome copies per PFU for each of the strains was determined by two independent quantitative PCR (qPCR) approaches: the DiaSorin qPCR platform described in the Methods section and the protocol developed by Lu *et al.* at the US Centers for Disease Control and Prevention (19). No significant differences in the number of genome copies per PFU were detectable between individual strains or PGs by either methodology (data not shown).

Next, we examined plaque phenotypes for the selected 7 HAdV-E4 strains in A549 human alveolar epithelial A549 cells. As shown in Figure 2, no obvious differences in plaque morphology were detected by visual examination of bright field images taken at 7 days post infection (dpi). However, PG I strains had larger plaque sizes than PG II strains. The vaccine strain CL68578 formed the largest plaques, while strains NY-15-4054, the cell culture-adapted RI-67 strain, and NHRC 90339 formed relatively smaller plaques. The PG II strains formed uniformly smaller plaque sizes that were not significantly smaller than those formed by the PG I strain NHRC 90339. One-way ANOVA with Holm-Sidak's multiple comparisons test showed significant differences between PG I strains CL68578, NY-15-4054, and RI-67, and all 3 examined PG II strains as well as PG I strain NHRC 90339.

The rate of plaque development was examined by calculating the percentage of plaques detectable on any given day post infection relative to the final total number of plaques over a period of 25 days, following the approach used by Tollefson *et al.* to

characterize HAdV-C ADP mutants (20, 21). As shown in Fig 3A, plaques formed by PG

I strains were visible one day earlier (day 4) than plaques formed by PG II strains (day 5).

PG I strains CL68578 and NY-15-4054 exhibited the fastest rate of plaque development.

At 10 dpi, about 90% and 70% of all detected plaques were visible for the CL68578 and

NY-15-4054, respectively, while only 30-40% of all detected plaques were visible for the

other examined strains, including NHRC 90339.

To evaluate cell-to-cell spread phenotypes, cytopathic effect (CPE) progression was

examined over a period of 7 days in A549 cell monolayers infected at different

multiplicities (MOI 0.001 - 10 PFU/cell) as described by Doronin *et al.* (22, 23). Overall,

the three PG II strains spread at a restricted rate compared to the PG I strains (Fig 3B).

Differences in replication kinetics and cell killing account for the observed

differences in plaque size and spread phenotypes. The larger plaque phenotype and

enhanced spread phenotype of PG I strains could either be a reflection of a faster

replication kinetics and/or of enhanced cytopathic effect resulting from infection. To

investigate this, we compared the growth kinetics of PG I and PG II strains in A549 cell

monolayers infected at a MOI of 1 PFU/cell. While infectious virus yields were similar at

48 and 72 hours post infection (hpi), infectious virus yields at 24 hpi were higher for PG I

strains (Fig 3C), suggesting faster growth kinetics. Next, we examined cell killing

phenotypes in A549 cell monolayers infected with PG I and PG II strains at a MOI of 10

PFU/cell. ATP levels were determined as indicators of cell viability. At 4 dpi intracellular

ATP levels in PG II-infected cells were relatively high, and similar to those detected in

mock-infected cells. In contrast, in PG I-infected cells ATP levels were significantly lower,

reflecting increased cell death. ATP levels were the lowest in RI-67- and CL68578-

infected cells and NY-15-4054- and NHRC 90339 exhibited intermediate cell killing

phenotypes. (Fig 3D). From the results of these experiments we conclude that both faster

growth kinetics and increased killing may contribute to the documented large plaque and

lateral spread phenotypes of PG I strains.

Examination of PG I and PG II strains in a cotton rat model of adenovirus

respiratory infection revealed differences. For an initial comparison of pathogenic

phenotypes of PG I and PG II *in vivo* we selected one strain from each PG to represent

the most extreme plaque size, spread, and cell killing phenotypes observed: CL68578,

(PG I) and NHRC 42606 (PG II). We first conducted some additional cell-based assays

to expand the characterization of these two strains. As shown in Figure 4A, and consistent

with our initial findings, PG I strain CL68578 exhibited a faster kinetics of viral progeny

release than NHRC 42606 in polarized human airway epithelial Calu-3 cells. Moreover

and consistent with the results of the cell viability assays based on intracellular ATP

detection, significantly higher levels of lactate dehydrogenase (LDH) activity indicative of

enhanced cell death (24) were detected at 48 and 72 hpi in supernatants of A549 cells

infected with CL68578 (Fig 4B). Both strains replicated in cotton rat lung epithelial cells -

albeit at modest levels compared to those observed in A549 cells, with CL68578

displaying faster growth kinetics (Fig 4C).

We infected female cotton rats intranasally (i.n.) with 10^7 PFU of strains CL68578 or

NHRC 42606 (n=3 rats per group) and found marked differences in the magnitude of the

resulting pulmonary pathology at day 3 pi (data not shown). In a second study, female

cotton rats in groups of 6 were infected with 10^7 PFU of each virus i.n. and euthanized at

1, 2, 3, and 6 dpi. The assessment of viral load in infected cotton rat lungs revealed

differences in the course of infection between the two examined viruses. Two of the

animals infected with NHRC 42606 (PG II) pre-segregated into two different cages died

at 3 dpi. These animals could not be dissected for tissue sampling and were therefore

excluded from subsequent analysis. Similar numbers of genome copies per gram of

tissue were detected by real-time quantitative PCR (qPCR) at 1 dpi in the lungs of rats

infected with the PG I or PG II strains (Fig. 5A). The viral load increased modestly by 2

dpi in animals infected with CL68578 (PG I) and gradually declined thereafter. Consistent

with the *in vitro* growth phenotypes documented for PG II strain NHRC 42606, lung viral

loads were almost a log unit lower at 2, 3, and 6 dpi in the animals infected with this virus.

Infectious virus was detectable by plaque assay in lung homogenates of infected rats

in both groups until day 3 pi in 4/6 animals infected with CL68578 and in 3/5 surviving

animals infected with NHRC 42606 (Fig. 5B). While titers at 1 dpi were higher in rats

infected with PG II strain NHRC 42606, a steady decline in viral load was documented for

this group supporting the qPCR results and suggesting a more rapid clearance of the

virus. HAdV hexon protein was readily detectable by immunohistochemistry in very

modest amounts in large and small airway epithelial cells and alveolar macrophages

during the first 3 days after infection (Fig. 5C) confirming that the cotton rat is a permissive

host for HAdV-E4. This assay revealed another difference between the examined PG I

and PG II strains, the localization of the signal in infected animals, particularly at 1 dpi

(Table 2). At this time, while the virus was almost exclusively detected in airway epithelia

in the rats infected with the PG I strain, the signal was predominantly found in alveolar

macrophages in the rats infected with the PG II strain. AdV hexon protein was also

detectable in alveolar macrophages of rats infected with the PG II strain on days 2 and 3

post infection.

To assess the extent of pulmonary inflammation resulting from HAdV-E4 respiratory

infection, lung sections from mock- and HAdV-E4-infected animals were stained with

hematoxylin and eosin, and scored for several metrics of pulmonary pathology as

described in Materials and Methods. Representative photomicrographs of stained lung

sections from rats euthanized at 1, 2, 3, and 6 dpi are shown in Figure 6A.

In agreement with the original descriptions of pulmonary pathology for cotton rats

infected with HAdV-C5 reported by Prince *et al.* (25), we observed histopathologic

changes in two phases, an early phase of monocyte-macrophage and neutrophil

infiltration affecting alveoli, bronchiolar epithelium, and peribronchiolar regions spanning

days 1-3; and a later phase of lymphocytic infiltration affecting peribronchiolar and

perivascular regions apparent at day 6 pi. Contrary to what we were expecting based on

the *in vitro* data, the average combined pathology scores were higher at all time points

sampled for rats infected with PG II strain NHRC 42606 (Figure 6B). Differences were

statistically significant at days 1, 3, and 6. On day 1 pi, rats infected with this strain

showed a stronger and more significant increase in interstitial pneumonia (inflammatory

cell infiltration and thickening of alveolar walls), and alveolitis (inflammatory cell infiltration

within the alveolar spaces) than rats infected with PG I strain CL68578. On day 3 pi there

was a significant enhancement in interstitial and alveolar cell infiltration in rats infected

with strain NHRC 42606, with some of them exhibiting complete consolidation of the lung

parenchyma. This increase in lung pathology correlated with the initiation of mortality at

3 dpi recorded in 2 animals infected with this virus.

Rats euthanized on day 6 presented the maximum overall pathology for both infected

groups. There was evidence of progression towards a marked increase in peribronchiolitis

and perivasculitis (particularly in the PG II strain NHRC 42606-infected rats), and a partial

resolution of the parenchymal infiltration (milder interstitial pneumonia and alveolitis).

The levels of pulmonary proinflammatory cytokine/chemokine gene expression were

determined by qPCR in lung homogenate supernatants from animals euthanized at 1, 2,

3, and 6 dpi. Statistically significant differences between groups were only detectable at

1 dpi when higher levels of IL-6, IFNβ, MCP-1, IP-10, and RANTES (CCL5) transcripts

were observed in samples from PG II strain NHRC 42606-infected animals. Although the

mean fold increase values for IFNγ, TNFα, and GRO were also higher in the PG II-infected

cotton rats, differences between groups were not statistically significant (Figure 7). These

results are consistent with the more pronounced cellular infiltration observed in the lungs

of rats infected with the PG II virus. Cytokine transcript levels at the other time points post

infection were not statistically different between groups (data not shown).

DISCUSSION

Evidence of intratypic genetic variably for HAdV-E4 has been reported in the literature

since the 1980s, with the description of multiple genomic variants identified by RFLP

analysis of serologically indistinguishable strains isolated from cases of respiratory illness

and conjunctivitis. However, the clinical significance of this naturally occurring variation

has remained elusive for this and other pathogenic HAdV types. In this paper, deeper

sequence analysis and proteotyping of a panel of 47 HAdV-E4 whole genome sequences

representing 45 strains enabled us to demonstrate additional differences between the two

major phylogroups (PG) of HAdV-E4 described by Gonzalez *et al.* (14), and also among

the examined genomic variants clustering within each of the two evolutionary lineages.

While the genomes of strains clustering into PG I and PG II are overall 94.5% identical,

we previously identified sequence differences between the two lineages mapping

throughout the entire genome. These included numerous single nucleotide

polymorphisms, insertions and deletions in regions of the genome that have the potential

to contribute to *in vitro* and *in vivo* phenotypes relevant to pathogenesis (14, 18, 26-29).

Our proteotyping analyses of selected coding regions confirmed that PG I and PG II

encode different variants not only for proteins involved in viral replication (E1A, E1B-55K,

DNA Pol, L4-100K, and E4-34K), but also for proteins with immunomodulatory functions

such as E1B-19K (30) and the E3-encoded gp19K, 14.7K, and CR1β (31-34). Using a

subset of previously sequenced strains that represent different genomic variants in the

two phylogroups, we conducted a comparative evaluation of *in vitro* and *in vivo*

phenotypes reflecting replicative fitness and pathogenicity. We demonstrated marked

differences between the two genetic lineages of HAdV-E4.

Compared to many RNA viruses for which genetic determinants of fitness and

virulence have been identified and links between genetic variability and phenotypic

variability have been demonstrated (35-39), relatively little is known about the implications

of genetic variability for DNA viruses of medical importance. Recent applications of

genomics approaches to the study of various aspects of human herpesvirus pathogenesis

have advanced the understanding of the magnitude and implications of genetic diversity

for members of this group (40-42). In addition, important studies have identified gene

variations of myxoma virus field strains that may be responsible for changes in virulence

(43). To our knowledge, this paper is the first to report phenotypic correlates of naturally

occurring intratypic genetic variability for human adenoviruses associated with disease.

The work presented here demonstrates distinct *in vitro* and *in vivo* phenotypes relevant

to pathogenesis for HAdV-E4 strains belonging to PG I and PG II. We show that in cell

culture, PG I strains are fitter viruses than PG II strains, exhibiting larger plaque sizes,

enhanced lateral spread, faster plaque development and replication, enhanced cell killing,

and earlier release of infectious viral progeny. The documented coding variations in the

E1, E2, and E4 regions among other shown in Table 1 segregate with these phenotypic

differences. A few of these regions have been previously identified experimentally to

contribute to replicative fitness. Through the generation of compensatory mutants with a

restored large plaque phenotype compared to the parental ADP-deficient mutant of

HAdV-C5 *dl327*, Subramanian *et al.* identified E1A, E1B19K, E1B55K, and E4Orf3 as

genes influencing the spread phenotype (44). Importantly, these authors also identified

compensatory mutations in the penton base and fiber genes that warrant further

investigation for HAdV-E4 strains.

As documented by our group and others, genomes of HAdV-E4 strains in PG I and

PG II also differ in the sequences of their respective inverted terminal repeats (ITRs) (14,

26, 27). Wunderlich *et al.* reported an increased genome replication rate for a HAdV-B35-

based vaccine vector carrying the alternative ITR sequence 5'-CTATCTAT-3' (45).

Interestingly, they also described the presence of this alternative ITR sequence in the

genomes of several HAdV types, including HAdV-E4 PG I strains RI-67 and CL68578,

which in our *in vitro* experiments had faster replication rates and enhanced lateral spread

compared to the PG II viruses.

The deeper examination of strains with apparent intermediate phenotypes for lateral

spread and plaque size (RI-67) or cell killing (NY-15-4054) as well as of strains encoding

different protein variants in loci of interest will facilitate the finer mapping of genetic

determinants of replicative fitness.

In the cotton rat model of HAdV respiratory infection the pathological response to

HAdV-E4 resembled that previously described for HAdV-C5 (25). Importantly, the model

allowed us to observe marked differences in induced pulmonary pathology between the

examined PG I and PG II viruses. This demonstrates the value of the cotton rat as a

permissive host for experimental *in vivo* examination of the effects of genetic variability of

HAdV-E4 on virulence and for the identification of candidate genetic determinants of

pathogenicity for this unique HAdV type.

Surprisingly, the differences in replicative fitness documented *in vitro* between PGs

did not correlate with the differences in virulence observed in the cotton rat model. These

data strongly implicate a differential involvement of the host immune response in the

pathogenesis of the documented pulmonary disease rather than a direct viral cytopathic

effect. The detection of significant differences in the levels of gene expression for

proinflammatory cytokines and of chemokines that regulate the migration and infiltration

of monocytes, macrophages, and T cells (MCP-1⁄CCL2, IP-10, and RANTES/CCL5) are

consistent with the observation of enhanced pulmonary pathology in rats infected with the

PG II virus.

With the exception of RIDα, all of the E3 proteins encoded in the genomes of CL68578

(PG I) and NHRC 42606 (PG II) are different, i.e. they represent different proteotypes, so

it is likely that the two examined viruses differ in their ability to elicit and modulate host

responses to infection.

There is a documented higher representation of PG II in HAdV-E4 positive clinical

specimens obtained from symptomatic patients (11, 14, 16-18) which suggests a higher

"epidemiologic fitness" and/or "transmission fitness" (46) for this clade. PG II strain NHRC

42606 was more virulent than PG I strain CL68578 in our *in vivo* experiments, as

evidenced by the more exacerbated inflammatory response in the lung of infected animals

and associated fatalities. Interestingly, the studies conducted by Russell *et al.* during 2004

in a cohort of unvaccinated military recruits in basic training at the Marine Corps Recruit

Training Command, San Diego, California revealed the efficient transmission of PG II

genomic variant 4a (represented in our in vitro studies by strain NHRC 22652) in that

unique setting, and collected data that implicated the presence of infectious virus in

various fomites and environmental samples in the process (47).

Further work using isogenic virus mutants carrying some of the identified mutations

that encode different proteotypes in various regions of the viral genome will be necessary

to dissect the contribution of individual coding regions to the differences in observed

pulmonary pathology phenotypes between PG I and PG II viruses. Of particular interest

are the variants encoded in regions implicated in the modulation of host responses to

infection, such as E1, L4, E3, and VA RNA (48-53).

Using the Syrian hamster model of HAdV respiratory infection, Radke *et al.* reported

enhanced pulmonary pathology for a naturally occurring genomic variant of HAdV-B14,

14p1, whose genomic sequence differs from that of the prototype strain de Wit in regions

of the genome including E1B and whose distinct *in vitro* phenotypes include a decreased

expression of E1B-19K, the viral Bcl2 homolog, and the formation of larger plaques (54).

Like the mouse, the cotton rat has been shown to be a powerful host to evaluate the

effect of early genes in HAdV-C5 pathogenesis (55-58). Our experiments demonstrate

that *Sigmodon hispidus* is permissive to HAdV-E4 infection and we expect the model to

be particularly useful for examination of the role of the species HAdV-E-specific E3-CR1

proteins that are predominantly expressed from late transcripts (59, 60). Importantly, the

biological function of E3CR1δ/E3-30K has not been identified to the present (59). The

work of Li and Wold (59) and preliminary studies in our laboratory (Kajon, personal

communication) indicate that HAdV-E4 does not encode an ADP homolog.

Our studies uncover a powerful experimental system and provide a strong foundation

for further exploration of specific genotype-phenotype associations for HAdV-E4 and

possibly for HAdVs in general. Our various low passage clinical isolates of HAdV-E4 and

their corresponding whole genome sequence provide a unique resource to examine other

phenotypes both *in vitro* and *in vivo* to define system-specific correlations of fitness and

virulence.

The identification of genetic determinants of fitness and virulence will be extremely

useful for the design of HAdV-E4-based vectors and for the genetic manipulation of

viruses for attenuation and use in live vaccine formulations. In addition, the identification

of genetic factors that influence pathogenicity can potentially contribute to the

development of molecular typing tools designed to identify mutations associated with

increased virulence at loci of special interest.

MATERIALS AND METHODS

Genetic analysis and proteotyping. To identify the distribution of nucleotide differences among HAdV-E4 genomes, the complete genomic sequences of HAdV-E4 strains in phylogroups (PG) I and II characterized by Gonzalez *et al.* (14) were aligned using MAFFT (61). Sequences were ordered following the branching pattern of the phylogenetic tree inferred with MEGA7 (62). Nucleotide sites were color-coded per genome in comparison to the consensus inferred from the multiple sequence alignment. The comparisons were performed with the R package APE (https://cran.r-project.org/web/packages/ape/) to identify by position the consensus nucleotide and the mutations. The predicted amino acid sequence divergence for selected coding regions among PG I and PG II HAdV-E4 strains was examined using the method of proteotyping adapted from Obenauer *et al.* (63). The concatenated protein sequences of E1A, E1B 19K and 55K, DNA polymerase, L4 100K, E3 proteins, and E4 34K were analyzed. Amino acid signatures were derived from a phylogenetic tree-guided sequence alignment, indicating amino acid sites with polymorphisms relative to the most frequently occurring residues with a frequency-based color coding.

Cell Culture. Human A549 type II alveolar epithelial cells (ATCC CL-185) were grown in Minimum Essential Medium (MEM, Gibco) supplemented with 8% heat-inactivated newborn calf serum (HI-NBCS, Rocky Mountain Biologicals), 1.5 g/ml Na_2CO_3 (Amresco), 2 mM L-glutamine (Corning), 10 U/ml penicillin (Corning), 10 µg/ml streptomycin (Corning), and 25 mM HEPES (Corning). Human Calu-3 airway epithelial cells (ATCC HTB-55) were grown in RPMI-1640 medium (Corning) supplemented with 10% fetal

bovine serum (FBS, Atlanta Biologicals), 2 mM L-glutamine, 10 U/ml penicillin, 10 μg/ml

streptomycin, and 25 mM HEPES. Cotton rat lung epithelial cells (CRLC, ATCC PTA-

3930) were grown in MEM (Gibco) supplemented as above except with 10% (v/v) heat-

inactivated FBS (Atlanta Biologicals) instead of HI-NBCS. For virus infections, all cell lines

were cultured in the appropriate medium with the serum content reduced to 2%. The cell

lines were confirmed to be Mycoplasma sp. free by testing using the ABM Mycoplasma

PCR Detection Kit (G238, Applied Biological Materials Inc., Canada).

Viruses. Low passage virus stocks of clinical isolates were produced in A549 cells. After

3 rounds of freeze/thawing, cell debris was removed by extraction with of H_2O-saturated

chloroform. After venting overnight at room temperature to remove traces of chloroform,

infectious titers were determined by standard plaque assay on A549 cell monolayers in

6-well plates. The prototype strain RI-67 of PG I was obtained from the collection of Goran

Wadell, Umeå University, Sweden with an unknown history of passage. The PG I 4p

vaccine strain CL68578 was derived from the pVQ WT #11183 genomic clone obtained

from ABL, Inc. Briefly, the genomic clone was released from the bacterial plasmid by

digestion with Pac I (New England Biolabs), and 1 μg of linearized DNA was transfected

into A549 cells using Effectene Transfection Reagent (Qiagen) following the

manufacturer's recommendations. The resulting infectious virus was passaged minimally

in A549 cells for stock production.

Cell-based assays

Plaque assay. Confluent A549 cell monolayers in 6 well plates were inoculated with 100

μl of serially diluted virus (1:10 in phosphate buffered saline (PBS). After adsorption for 1

hour at 37°C and 5% CO_2 with periodic rocking, monolayers were overlaid with 2 ml of a

solution consisting of equal parts of 1.4% (w/v) Low Melt Agarose (G-BioSciences) and

2X MEM (Gibco) supplemented with 4% (v/v) HI-NBCS, 4.0 mM L-glutamine, 20 U/ml

penicillin, 20 μg/ml streptomycin, and 25 mM $MgCl_2$. At 7 dpi, infected monolayers were

fixed with 1% formaldehyde (Sigma)/ 0.15 M NaCl and stained with crystal violet (EMD-

Millipore Sigma).

Plaque development assay. A549 cells were seeded in 15 mm x 60 mm dishes to be

confluent on the day of infection. Stocks were diluted in PBS so that approximately equal

number of PFUs of each strain were used to inoculate each dish in biological triplicate.

Growth medium was aspirated, and the cells were inoculated with diluted virus and

synchronized on ice for 10 minutes. Inoculated cells were then incubated for two hours at

37°C 5% CO_2. Following absorption, 5 ml of overlay medium (described above) were

added and allowed to solidify at room temperature, then incubated at 37°C and 5% CO_2.

At 3 dpi, an additional overlay containing 0.0015% neutral red dye (JT Baker Chemical

Co.) was added. Starting a 4 dpi, plaques were counted daily using a lighted background

and marked with a pen. The assay was terminated when no more plaques developed.

The data are presented as percentage of the final titer (total number of plaques observed

on a given day divided by the total number of plaques observed on the final day of the

assay) on each day (21).

Plaque size assays. A549 cells in 6-well plates were infected with approximately 20-40

PFU of each virus strain in a volume of 100 μl. After adsorption for 1 hour at 37°C with

occasional rocking, the infected monolayers were overlaid as described above. At 7 dpi,

prior to fixation, individual plaques were photographed using an Olympus CKX41

microscope equipped with an Infinity 2-3 digital camera (Teledyne Lumenera) and a 4X

objective. Images were captured using the Infinity Capture software V6.5.4 (Teledyne Lumenera). The plaque assays were then fixed, stained with crystal violet, and photographed using a copy stand and a transilluminator with the camera in a fixed position. The average plaque size in arbitrary units was determined using the FIJI image analysis software (64).

Growth kinetics studies. A549 cells seeded in 10 mm x 35 mm cell culture dishes at 80% confluence were inoculated with HAdV-E4 strains at a MOI of 1 PFU/cell in biological triplicate, incubated on ice for 10 minutes, then absorbed for one hour at 37°C and 5% CO_2 with periodic rocking. Unbound virus was removed by aspiration, and the infected cell monolayers were washed twice with PBS. Cells were then replenished with A549 Infection Medium (MEM + 2% HI-NBCI + supplements) and incubated at 37°C with 5% CO_2. At various times post infection, infected cell monolayers were frozen at -80°C and subjected to three freeze/thaw cycles. Cell debris was clarified by centrifugation at 300 x g for 10 minutes, and infectious virus titers were determined by plaque assay as described above.

CRLC monolayers were infected with strains CL68578 or NHRC 42606 at a MOI of 5 PFU/cell in biological triplicate, incubated on ice for 10 minutes, then absorbed for one hour at 37°C and 5% CO_2 with periodic rocking. At 0, 12, 24, 48 and 72 hpi, cell monolayers were processed as described above for titration of infectious virus yields.

Virus spread assays. A549 cells seeded on 48 well plates to be 100% confluent at the time of infection were inoculated with 100 μl of virus stock diluted in PBS for MOIs of 10, 1, 0.1, 0.01, and 0.001 PFU/cell. After adsorption for 1 hour at 37°C, the inoculum was removed by aspiration, cell monolayers were washed twice with PBS, replenished with

A549 Infection Medium, and incubated at 37°C and 5% CO_2. At 7 dpi cell monolayers

were fixed with 1% formaldehyde in 0.15 M NaCl, and stained with crystal violet.

Cell viability assays. Intracellular ATP levels were measured at various times post

infection as indicators of the presence of metabolically active cells and thus of cell

viability (65) using the luminescent cell viability CellTiter-Glo 2.0 assay (Promega)

according to the manufacturer's specifications. Briefly, A549 cells were seeded in 96 well

dishes to be 75% confluent on the day of infection. Cells were inoculated at a MOI of 10

PFU/cell with virus diluted in PBS or mock-infected with PBS in biological quadruplicate.

After adsorption for 2 hours at 37°C the inoculum was removed, monolayers were washed

twice with PBS, and then then replenished with 50 μl of A549 Infection Medium. Relative

luminescence was measured in a Luminoskan Ascent Type 392 (ThermoFisher

Scientific). Data are reported as mean relative luminescence units ± SEM.

To further characterize the cell killing phenotype of PG I and PG II strains, LDH release

from lysed cells was assessed using the colorimetric CytoTox 96 Non-radioactive

Cytotoxicity Assay Kit (Promega). Briefly, A549 cells seeded in 48-well plates to be

confluent at the time of infection were inoculated with virus diluted in PBS at a MOI of 1

PFU/cell in biological triplicates. After adsorption for 1 hour at 37°C the inoculum was

removed, monolayers were washed twice with PBS and then replenished with infection

medium. At 1, 2, and 3 dpi, the supernatant was removed and centrifuged at 300 x g for

10 minutes. LDH activity was then assessed according to the manufacturer's directions

in technical quadruplicate. Data are reported as mean absorbance at 490 nm ± the

standard error of the mean (SEM).

Viral progeny release assays in polarized Calu-3 cells. Calu-3 cells were seeded onto transwell inserts (0.4 μm pore size, Corning) at a density of 5 x10^5 cells/well, and Calu-3 growth medium was added to both the apical and basal compartments. Two days post seeding, the cells were washed with PBS++ (Dulbecco's Phosphate-Buffered Saline with Magnesium and Calcium, Gibco). All liquid was then removed from the apical chamber, and the cells were grown at an air/liquid interphase for 10 days with medium changes every 48 hours. Polarization of Calu-3 cells was evaluated by measuring the transepithelial resistance (TER) using a Millicell ERS meter (Millipore). Cells were considered polarized when TER values reached 500-600 Ωcm^{-2}.

For infection, medium was removed from the basal compartment, and Calu-3 infection medium was added. Virus diluted in PBS++ for a MOI of 10 PFU/cell was then added to the apical compartment. After absorption at 37°C, 5% CO_2 for 1 hour, the inoculum was removed, and the cells were washed with PBS++. All liquid in the apical and basal compartments was removed, and fresh Calu-3 infection medium was added to the basal compartment. The cells were then incubated at 37°C, 5% CO_2. At 1-3 dpi, the infected cells were processed by washing the apical compartment twice with a total of 400 μl of PBS ++. Infectious virus titers in the apical wash were determined by plaque assay.

Cotton rat studies

All animal studies were conducted under applicable laws and guidelines and after protocol approval by the Sigmovir Biosystems, Inc. (SBI) Institutional Animal Care and Use Committee. Cotton rats (*Sigmodon hispidus*) obtained from SBI's inbred colony were housed in large polycarbonate cages and fed a diet of standard rodent chow and water

ad libitum. The number of animals per experimental group required to observe differences

in pulmonary pathology scores was determined to be 6 by power analysis using G*power

software v3.1.9.2 (66), The calculations were carried out assuming a significance level of

0.05, 80% power, a signal/noise ratio of 1.5 pathology score units, and a standard

deviation of 1.

Four-week-old female cotton rats were inoculated with 5×10^6 PFUs of PG I strain

CL68578 or PG II strain NHRC 42606 in each nare under isoflurane anesthesia. Control

mock-infected animals were inoculated in the same manner with chloroform-extracted

supernatants of uninfected A549 cells. Cotton rats were sacrificed at 1, 2, 3, and 6 days

post infection by CO_2 asphyxiation. Lung tissue was collected for downstream analyses

as follows: left lobe for assessment of viral load; left lingular lobe for mRNA reverse

transcription for cytokine gene expression; and right lung for histopathology. All lung lobes

except for the right lobe were snap-frozen in liquid nitrogen and stored at -80°C until

processed. The right lung was inflated with 10% neutral buffered formalin.

Assessment of pulmonary viral load. Frozen left lobe sections were weighed and then

homogenized in 1mL of A549 infection medium using a TissueLyser II apparatus (Qiagen)

and 3 cycles of 30 hz for 30 seconds, chilling on ice between each cycle. Homogenates

were centrifuged at 16,000 X g for 10 minutes, and the supernatant was aliquoted into

three separate tubes of equal volume.

One aliquot of lung homogenate supernatant was processed for assessment of

infectious viral load by plaque assay. Supernatants were serially diluted ten-fold in PBS

and 100 μl were plated in triplicate onto monolayers of A549 cells in 6-well plates. The

assays were carried out as described above.

At DiaSorin Molecular (Cypress, CA), DNA was extracted from 200 µl of another

aliquot of lung homogenate supernatant using the MagNA Pure Total Nucleic Acid

Isolation protocol and reagents on a MagNA Pure LC 2.0 instrument (Roche Diagnostics).

Prior to extraction, 5 µl of the Simplexa Extraction and Amplification control DNA (SEAC,

DiaSorin Molecular) were added to each sample. Samples were eluted in 50 µl.

The real-time qPCR assay was carried on the LIAISON® MDX instrument (DiaSorin

Molecular) using the 96 well Universal Disc (DiaSorin Molecular) and DiaSorin Molecular

Adenovirus analyte-specific reagents. The adenovirus 3' Hexon primer pair and the

adenovirus 5' Hexon primer pair with FAM-labeled forward primers were used to amplify

and detect conserved regions at the 3' and 5' ends of the HAdV hexon gene. A Control

SEAC Primer Pair with a Quasar® 760-labeled forward primer was used to amplify and

detect the SEAC DNA fragment. The 5' and 3' hexon primer pairs are a proprietary bi-

functional fluorescent primer-probe set targeting the 3' and 5' regions of the HAdV hexon

gene. Each reaction contained 4 µl of 2.5X Universal Master Mix, 0.2 µl of Adenovirus 3'

Hexon primer pair, 0.2 µl of Adenovirus 5' Hexon primer pair, 0.2 µl of internal control

primer, 0.4 µl of water, and 5 µl of extracted template, in a total reaction volume of 10 µl.

PCR amplification with real-time detection was performed using the following cycling

conditions: denaturation at 97°C for 120 seconds (1 cycle); denaturation at 97°C for 10

seconds followed by annealing/extension at 60°C for 30 seconds (40 cycles).

Fluorescence was measured at the end of each annealing/extension cycle with

quantitative values calculated by the LIAISON® MDS software.

Standard curves were generated using highly pure genomic DNA extracted from either

PG I vaccine strain CL68578- or PG II clinical isolate NHRC 42606 -infected A549 cells

as previously described (67). The calibrator values ranged from $1.2x10^6$ to $1.2x10^2$

genome copies (cp) /ml for CL68578 and from $2.8x10^6$ to $2.8x10^2$ cp/ml for NHRC 42606.

Each standard was extracted in singlicate and each extract amplified in quadruplicate in

a single run. Quantitative analysis was performed by the LIAISON® MDX Software.

Standard curves were used for all viral load determinations. Viral loads were expressed

as genome copies per gram of tissue.

RNA Isolation and Reverse Transcription. Total RNA was extracted from the left

lingular lung lobes using the RNAqueous kit (ThermoFisher Scientific). Frozen lung

sections were homogenized in 1 ml of Lysis/Binding solution in a TissueLyser II (Qiagen)

with three cycles of at 30 Hz for 30 seconds, with chilling on ice between cycles. Samples

were centrifuged at 16,000 x g for 2 minutes at 4°C and the supernatant was used for

total RNA isolation according to the manufacturer's recommendations. Residual DNA was

removed from the extracted RNA with the TurboDNA-free kit (ThermoFisher Scientific).

The removal of contaminating DNA was verified by PCR amplification of the rig/S15 gene

using the forward (5' – 3') primer TTC CGC AAG TTC ACC TAC C and the reverse (3' –

5') primer CGG GCC GGC CAT GCT TTA CG with GoTaq Flexi DNA polymerase

(Promega) and the following cycling conditions: 1 cycle of 95°C, 5 minutes; 50 cycles of

95°C for 30 seconds, followed by 57°C for 30 seconds, then 72°C for 30 seconds; 1 cycle

of 72°C for 5 minutes. Reverse transcription was then performed on 1 μg of DNAse-

treated RNA using the RetroScript Reverse Transcription kit (Invitrogen/ThermoFisher

Scientific) and oligo(dT) primers following the manufacturer's recommendations.

Histological analysis. Dissected right lung lobes were inflated with 10% neutral buffered

formalin and immersed in formalin for fixation. Tissues were then shipped to Histoserv,

Inc. (Germantown, MD) for paraffin-embedding, sectioning, and staining with hematoxylin

and eosin. The stained lung sections were scored blindly by co-author J.C.G.B. on a scale

of 0 to 4 (absent, minimal, mild, moderate, and marked) for the following indices of

pulmonary inflammation: peribronchiolitis (peribronchiolar inflammatory cell infiltration),

perivasculitis (inflammatory cell infiltration around the small blood vessels), interstitial

pneumonia (inflammatory cell infiltration and thickening of alveolar walls), and alveolitis

(inflammatory cell infiltration of alveolar spaces). The average combined pathology score

was calculated from these four individual components.

Immunohistochemistry. Services were contracted from the Veterinary Diagnostic

Laboratory, University of Minnesota (https://www.vdl.umn.edu/) where paraffin-

embedded lung sections on glass slides were processed for staining with AdV Mab

MAB805 (Millipore) following an in-house protocol. Images from stained sections were

acquired in a Nikon Eclipse E600 microscope equipped with a Nikon DXM1200F digital

camera at 40X magnification.

Analysis of cytokine/chemokine gene expression in lung tissue. cDNA obtained as

described above was diluted to 0.1μg /ml and 3 μl were used as template for each 25 μl

qPCR reaction using primers and cycling conditions described previously for IL-6, IFNβ,

IFNγ, TNFα, MCP-1, IP-10, RANTES, and GRO (68-70). qPCR reactions were carried

out using the QuantiFast SYBR Green PCR kit (Qiagen, Valencia, CA) in a MyiQ single

color Real-time PCR cycler (Bio-Rad, Temecula, CA). Relative quantification of PCR

products was performed by normalizing to β-actin using the $\Delta\Delta C_T$ method. Data are

presented as mean fold induction ± SEM.

Statistical Analysis. Unless otherwise noted, all statistical analyses were carried out using GraphPad Prism v8.1.0 for Microsoft Windows (GraphPad, https://www.graphpad.com). p values < 0.05 were considered statistically significant.

ACKNOWLEDGMENTS

The authors wish to thank Bojana Rodic-Polic, DiaSorin Molecular (Cypress, CA) for the execution of qPCR assays to assess viral load in cotton rat lung homogenates and Xioyan Lu at CDC (Atlanta, GA) for the qPCR assessment of viral load for our infectious stocks. We also especially thank Kathy Spindler for critical reading of the manuscript and constructive input.

C.R.B. was supported by the University of New Mexico Infectious Diseases and Inflammation NIH Training Grant T32-AI007538.

A.E.K. and C.R.B conceived the study. C.R.B., G.G., W.Z, A.K., D.S., J.C.G.B. and A.E.K. performed investigations for this study. C.R.B. and A.E.K. wrote the manuscript, and all authors reviewed and edited the manuscript.

 FIGURE LEGENDS

Figure 1: A. Sequence differences among HAdV-E4 strains of phylogroups I and II

(PG I and PG II). The consensus was obtained from the multiple sequence alignment

built using MAFFT (61) with the set of whole genome sequences originally analyzed by

Gonzalez *et al.* (15), and summarized with the R package APE to identify by position the

consensus nucleotide and the mutations. The GenBank accession numbers for the

examined genomic sequences together with the place and year of isolation of the

corresponding strain are shown on the vertical axis on the left. Positions with nucleotide

mutations differing from the consensus are colored with red, green, blue or orange,

corresponding to mutations to adenine, cytosine, guanine or thymine, respectively. **B.**

Proteotyping of concatenated predicted protein sequences for selected coding

regions. The GenBank accession numbers for the examined genomic sequences

together with the place and year of isolation of the corresponding strain are shown on the

vertical axis on the left. The start, end and polypeptide name of the concatenated

sequences arranged in genome order are shown on the top of the panel. Sites with amino

acid polymorphisms are colored according to the frequency across sequences, with blank

for the consensus, blue and red for the first and second most frequent polymorphisms for

a position; gap sites are colored in black.

Figure 2: Plaque phenotypes. (A) Plaque assays on A549 cell monolayers in 6-well

plates. Monolayers infected with PG I (RI-67, CL68578, NY-15-4054, and NHRC 90339)

or PG II (NHRC 25562, NY-12-27440, and NHRC 42606) strains were fixed and stained

with crystal Violet at 7 days post infection. For each examined strain, representative

brightfield images of individual plaques are shown next to the corresponding images of a

representative crystal-violet stained monolayer. (B) Plaque areas (in arbitrary units) were

determined using FIJI image analysis software. Statistical significance was assessed by

one-way ANOVA (6, 484 = 37.5) and Holm-Sidak's multiple-comparison test (*, $p < 0.05$;

***, $p < 0.001$; ****, $p < 0.0001$).

Figure 3: Growth and spread phenotypes. A. Plaque development. Plaques newly

developed on infected A549 cell monolayers in 60 mm dishes were counted every one to

two days. The total number of plaques observed on a given day is plotted as a percentage

of the total number of plaques observed on the final day of the assay. **B. Lateral spread.**

A549 cell monolayers were infected at MOIs ranging from 0.001 to 10 PFU/cell with the

indicated PG I or PG II strains. At 7 days post infection, cell monolayers were fixed and

stained with crystal violet for visualization of the extent of cytopathic effect. **C. Replication**

kinetics. Submerged A549 cell monolayers were infected with PG I or PG II strains at a

MOI of 1 PFU/cell. Infectious virus yields were determined at the indicated time points

post infection by plaque assay. **D. Cell killing phenotype.** A549 cells were infected at a

MOI of 10 PFU/cell. At 96 hours post infection, intracellular ATP levels were measured

as indicators of cell viability. Error bars denote SEM. Statistical significance was accessed

by one-way ANOVA analysis, $F_{(7, 24)} = 140.9$, $p < .0001$. Tukey's post hoc comparisons

to mock-treated cells revealed a significant difference. ****, $p < .0001$.

Figure 4: Additional phenotypic characterization of strains CL68578 and NHRC42606 using cell-based assays. A. Kinetics of viral progeny release in polarized human airway epithelial cells. Polarized Calu-3 cells in transwell inserts were infected at a MOI of 10 PFU/cell. At the indicated times post infection, the apical compartment was washed twice with a total of 400 μl of PBS and infectious virus titers in the combined wash fluid were determined by plaque assay. **B. Kinetics of LDH release from infected A549 cells.** A549 cell monolayers in 48-well plates were infected at a MOI of 1 PFU/cell in biological triplicates. LDH activity in infected cell supernatants was determined as an indicator of cell lysis. Statistical significance was assessed using the Student's *t* test, *$p< 0.05$, **** $p < 0.0001$.

C. Growth kinetics in monolayers of cotton rat lung epithelial cells. CRLC were infected at a MOI of 5 PFU/cell in biological triplicates. Infectious virus yields were determined by plaque assay in A549 cells.

Figure 5: Progression and localization of infection in cotton rat lungs. A. Assessment of viral load in the lung of infected cotton rats by qPCR. The viral load in the supernatants of left lung lobe homogenates from PG I and PG II-infected cotton rats at the indicated times post infection was determined by qPCR using a DiaSorin Molecular proprietary platform for human adenovirus. Genome copy numbers were normalized to the mass of homogenized tissue. Horizontal bars denote mean values for each group. Each symbol represents a single animal. **B. Infectious virus load in the lungs of infected cotton rats.** Infectious virus loads in lung homogenate supernatants were determined by plaque assay. Titers were normalized to the mass of homogenized

tissue and loads expressed as PFU/gram of tissue. Horizontal bars denote mean values

for each group. Infectious virus was not detectable in samples collected at 6 dpi. **C.**

Immunohistochemical detection of hexon protein in lung sections of infected

cotton rats. Paraffin-embedded lung sections on glass slides were stained with AdV Mab

MAB805 (Millipore). Images from stained sections were acquired in a Nikon Eclipse E600

microscope equipped with a Nikon DXM1200F digital camera at 40X magnification.

Arrowheads indicate positive cells.

Figure 6: Pulmonary pathology in cotton rats infected intranasally. A. Microscopic

examination of lung sections. The right lung lobes of mock- or HAdV-E4-infected cotton

rats were harvested at 1, 2, 3 and 6 dpi and processed for paraffin-embedding, sectioning

and hematoxylin and eosin staining for microscopic comparative examination of lung

pathology. Representative photomicrographs for each experimental group and time post

infection were taken at a 40X magnification in an Olympus BH-2 microscope equipped

with an Olympus DP70 digital camera and Olympus DP Controller software with focus on

terminal bronchioles and surrounding parenchyma. The insert on panel XI is a 400X

image of an area of parenchymal consolidation. The white arrow on panel XII points to an

area of perivascular infiltration. The black arrow on panel XII points to an area of

peribronchiolar infiltration. **B. Combined pulmonary pathology scores.** Mean combined

pulmonary pathology scores of hematoxylin and eosin-stained lung sections were

calculated for experimental groups of 5 or 6 rats. Error bars denote SEM. Statistical

significance was assessed by Mann-Whittney U test. $*p < 0.05$, $**p < 0.01$.

Figure 7: Proinflammatory cytokine and chemokine gene expression in infected

cotton rat lungs at 1 dpi. The levels of gene expression in lung homogenates of cotton

rats infected with the PG I strain CL68578, PG II strain NHRC 42606, or mock-infected

were determined by RT-qPCR. n= 5 or 6 rats per group. Error bars denote SEM.

Statistical significance was assessed by Student's *t* test. $p < 0.05$**.**

REFERENCES

1. Hilleman MR, Werner JH. 1954. Recovery of new agent from patients with acute respiratory illness. Proc Soc Exp Biol Med 85:183-8.

2. Grayston JT, Yang YF, Johnston PB, Ko LS. 1964. Epidemic Keratoconjunctivitis on Taiwan: Etiological and Clinical Studies. Am J Trop Med Hyg 13:492-8.

3. Muzzi A, Rocchi G, Lumbroso B, Tosato G, Barbieri F. 1975. Acute haemorrhagic conjunctivitis during an epidemic outbreak of adenovirus-type-4 injection. Lancet 2:822-3.

4. D'Angelo LJ, Hierholzer JC, Keenlyside RA, Anderson LJ, Martone WJ. 1979. Pharyngoconjunctival fever caused by adenovirus type 4: report of a swimming pool-related outbreak with recovery of virus from pool water. J Infect Dis 140:42-7.

5. Faden H, Gallagher M. 1980. Disseminated infection due to adenovirus type 4. Clin Pediatr (Phila) 19:427-9.

6. Levandowski RA, Rubenis M. 1981. Nosocomial conjunctivitis caused by adenovirus type 4. J Infect Dis 143:28-31.

7. Aoki K, Kato M, Ohtsuka H, Ishii K, Nakazono N, Sawada H. 1982. Clinical and aetiological study of adenoviral conjunctivitis, with special reference to adenovirus types 4 and 19 infections. Br J Ophthalmol 66:776-80.

8. Ariga T, Shimada Y, Ohgami K, Tagawa Y, Ishiko H, Aoki K, Ohno S. 2004. New genome type of adenovirus serotype 4 caused nosocomial infections associated with epidemic conjunctivitis in Japan. J Clin Microbiol 42:3644-8.

9. Kandel R, Srinivasan A, D'Agata EM, Lu X, Erdman D, Jhung M. 2010. Outbreak of

adenovirus type 4 infection in a long-term care facility for the elderly. Infect Control

Hosp Epidemiol 31:755-7.

10. Narra R, Bono P, Zoccoli A, Orlandi A, Piconi S, Grasselli G, Crotti S, Girello A, Piralla

A, Baldanti F, Lunghi G. 2016. Acute respiratory distress syndrome in adenovirus

type 4 pneumonia: A case report. J Clin Virol 81:78-81.

11. Kajon AE, Lamson DM, Bair CR, Lu X, Landry ML, Menegus M, Erdman DD, St

George K. 2018. Adenovirus Type 4 Respiratory Infections among Civilian Adults,

Northeastern United States, 2011-2015(1). Emerg Infect Dis 24:201-209.

12. Li QG, Wadell G. 1988. The degree of genetic variability among adenovirus type 4

strains isolated from man and chimpanzee. Arch Virol 101:65-77.

13. Adrian T. 1992. Genome type analysis of adenovirus type 4. Intervirology 34:180-3.

14. Gonzalez G, Bair CR, Lamson DM, Watanabe H, Panto L, Carr MJ, Kajon AE. 2019.

Genomic characterization of human adenovirus type 4 strains isolated worldwide

since 1953 identifies two separable phylogroups evolving at different rates from their

most recent common ancestor. Virology 538:11-23.

15. Kajon AE, Moseley JM, Metzgar D, Huong HS, Wadleigh A, Ryan MA, Russell KL.

2007. Molecular epidemiology of adenovirus type 4 infections in US military recruits

in the postvaccination era (1997-2003). J Infect Dis 196:67-75.

16. Itakura S, Aoki K, Sawada H, Ishiguro N, Shinagawa M. 1991. Changes in subgenome

types of adenovirus type 4 isolated from patients with ocular disease between 1985

and 1989 in Sapporo, Japan. J Clin Microbiol 29:1740-3.

17. Wadell G, Hammarskjold ML, Winberg G, Varsanyi TM, Sundell G. 1980. Genetic
variability of adenoviruses. Ann N Y Acad Sci 354:16-42.

18. Zhang J, Kang J, Dehghan S, Sridhar S, Lau SKP, Ou J, Woo PCY, Zhang Q, Seto
D. 2019. A Survey of Recent Adenoviral Respiratory Pathogens in Hong Kong
Reveals Emergent and Recombinant Human Adenovirus Type 4 (HAdV-E4)
Circulating in Civilian Populations. Viruses 11.

19. Lu X, Trujillo-Lopez E, Lott L, Erdman DD. 2013. Quantitative real-time PCR assay
panel for detection and type-specific identification of epidemic respiratory human
adenoviruses. J Clin Microbiol 51:1089-93.

20. Tollefson AE, Ryerse JS, Scaria A, Hermiston TW, Wold WS. 1996. The E3-11.6-kDa
adenovirus death protein (ADP) is required for efficient cell death: characterization of
cells infected with adp mutants. Virology 220:152-62.

21. Tollefson AE, Scaria A, Hermiston TW, Ryerse JS, Wold LJ, Wold WS. 1996. The
adenovirus death protein (E3-11.6K) is required at very late stages of infection for
efficient cell lysis and release of adenovirus from infected cells. J Virol 70:2296-306.

22. Doronin K, Toth K, Kuppuswamy M, Krajcsi P, Tollefson AE, Wold WS. 2003.
Overexpression of the ADP (E3-11.6K) protein increases cell lysis and spread of
adenovirus. Virology 305:378-87.

23. Doronin K, Toth K, Kuppuswamy M, Ward P, Tollefson AE, Wold WS. 2000. Tumor-
specific, replication-competent adenovirus vectors overexpressing the adenovirus
death protein. J Virol 74:6147-55.

24. Decker T, Lohmann-Matthes ML. 1988. A quick and simple method for the quantitation

of lactate dehydrogenase release in measurements of cellular cytotoxicity and tumor

necrosis factor (TNF) activity. J Immunol Methods 115:61-9.

25. Prince GA, Porter DD, Jenson AB, Horswood RL, Chanock RM, Ginsberg HS. 1993.

Pathogenesis of adenovirus type 5 pneumonia in cotton rats (Sigmodon hispidus). J

Virol 67:101-11.

26. Dehghan S, Seto J, Liu EB, Walsh MP, Dyer DW, Chodosh J, Seto D. 2013.

Computational analysis of four human adenovirus type 4 genomes reveals molecular

evolution through two interspecies recombination events. Virology 443:197-207.

27. Houng HS, Clavio S, Graham K, Kuschner R, Sun W, Russell KL, Binn LN. 2006.

Emergence of a new human adenovirus type 4 (Ad4) genotype: identification of a

novel inverted terminal repeated (ITR) sequence from majority of Ad4 isolates from

US military recruits. J Clin Virol 35:381-7.

28. Kidd AH, Garwicz D, Oberg M. 1995. Human and simian adenoviruses: phylogenetic

inferences from analysis of VA RNA genes. Virology 207:32-45.

29. Crawford-Miksza LK, Nang RN, Schnurr DP. 1999. Strain variation in adenovirus

serotypes 4 and 7a causing acute respiratory disease. J Clin Microbiol 37:1107-12.

30. Radke JR, Grigera F, Ucker DS, Cook JL. 2014. Adenovirus E1B 19-kilodalton protein

modulates innate immunity through apoptotic mimicry. J Virol 88:2658-69.

31. Lichtenstein DL, Toth K, Doronin K, Tollefson AE, Wold WS. 2004. Functions and

mechanisms of action of the adenovirus E3 proteins. Int Rev Immunol 23:75-111.

32. Fessler SP, Delgado-Lopez F, Horwitz MS. 2004. Mechanisms of E3 modulation of

immune and inflammatory responses. Curr Top Microbiol Immunol 273:113-35.

33. Fu J, Li L, Bouvier M. 2011. Adenovirus E3-19K proteins of different serotypes and

subgroups have similar, yet distinct, immunomodulatory functions toward major

histocompatibility class I molecules. J Biol Chem 286:17631-9.

34. Martinez-Martin N, Ramani SR, Hackney JA, Tom I, Wranik BJ, Chan M, Wu J, Paluch

MT, Takeda K, Hass PE, Clark H, Gonzalez LC. 2016. The extracellular interactome

of the human adenovirus family reveals diverse strategies for immunomodulation.

Nat Commun 7:11473.

35. Gonzalez-Scarano F, Beaty B, Sundin D, Janssen R, Endres MJ, Nathanson N. 1988.

Genetic determinants of the virulence and infectivity of La Crosse virus. Microbial

Pathogenesis 4:1-7.

36. Tscherne DM, Garcia-Sastre A. 2011. Virulence determinants of pandemic influenza

viruses. J Clin Invest 121:6-13.

37. Muller V, Fraser C, Herbeck JT. 2011. A strong case for viral genetic factors in HIV

virulence. Viruses 3:204-16.

38. Yoshii K, Sunden Y, Yokozawa K, Igarashi M, Kariwa H, Holbrook MR, Takashima I.

2014. A critical determinant of neurological disease associated with highly pathogenic

tick-borne flavivirus in mice. J Virol 88:5406-20.

39. Coetzee P, Van Vuuren M, Stokstad M, Myrmel M, Venter EH. 2012. Bluetongue virus

genetic and phenotypic diversity: towards identifying the molecular determinants that

influence virulence and transmission potential. Vet Microbiol 161:1-12.

40. Mao H, Rosenthal KS. 2003. Strain-dependent structural variants of herpes simplex

virus type 1 ICP34.5 determine viral plaque size, efficiency of glycoprotein

processing, and viral release and neuroinvasive disease potential. J Virol 77:3409-

17.

41. Galitska G, Biolatti M, De Andrea M, Leone A, Coscia A, Bertolotti L, Ala U, Bertino E,

Dell'Oste V, Landolfo S. 2018. Biological relevance of Cytomegalovirus genetic

variability in congenitally and postnatally infected children. J Clin Virol 108:132-140.

42. Akhtar LN, Bowen CD, Renner DW, Pandey U, Della Fera AN, Kimberlin DW, Prichard

MN, Whitley RJ, Weitzman MD, Szpara ML. 2019. Genotypic and Phenotypic

Diversity of Herpes Simplex Virus 2 within the Infected Neonatal Population.

mSphere 4.

43. Kerr PJ, Cattadori IM, Rogers MB, Fitch A, Geber A, Liu J, Sim DG, Boag B, Eden JS,

Ghedin E, Read AF, Holmes EC. 2017. Genomic and phenotypic characterization of

myxoma virus from Great Britain reveals multiple evolutionary pathways distinct from

those in Australia. PLoS Pathog 13:e1006252.

44. Subramanian T, Vijayalingam S, Chinnadurai G. 2006. Genetic identification of

adenovirus type 5 genes that influence viral spread. J Virol 80:2000-12.

45. Wunderlich K, van der Helm E, Spek D, Vermeulen M, Gecgel A, Pau MG, Vellinga J,

Custers J. 2014. An alternative to the adenovirus inverted terminal repeat sequence

increases the viral genome replication rate and provides a selective advantage in

vitro. J Gen Virol 95:1574-1584.

46. Wargo AR, Kurath G. 2012. Viral fitness: definitions, measurement, and current

insights. Curr Opin Virol 2:538-45.

47. Russell KL, Broderick MP, Franklin SE, Blyn LB, Freed NE, Moradi E, Ecker DJ,

Kammerer PE, Osuna MA, Kajon AE, Morn CB, Ryan MA. 2006. Transmission

dynamics and prospective environmental sampling of adenovirus in a military recruit

setting. J Infect Dis 194:877-85.

48. Kalvakolanu DV, Bandyopadhyay SK, Harter ML, Sen GC. 1991. Inhibition of

interferon-inducible gene expression by adenovirus E1A proteins: block in

transcriptional complex formation. Proc Natl Acad Sci U S A 88:7459-63.

49. Chahal JS, Qi J, Flint SJ. 2012. The human adenovirus type 5 E1B 55 kDa protein

obstructs inhibition of viral replication by type I interferon in normal human cells. PLoS

Pathog 8:e1002853.

50. Chahal JS, Flint SJ. 2012. Timely synthesis of the adenovirus type 5 E1B 55-kilodalton

protein is required for efficient genome replication in normal human cells. J Virol

86:3064-72.

51. Andrade F, Bull HG, Thornberry NA, Ketner GW, Casciola-Rosen LA, Rosen A. 2001.

Adenovirus L4-100K assembly protein is a granzyme B substrate that potently inhibits

granzyme B-mediated cell death. Immunity 14:751-61.

52. Kitajewski J, Schneider RJ, Safer B, Munemitsu SM, Samuel CE, Thimmappaya B,

Shenk T. 1986. Adenovirus VAI RNA antagonizes the antiviral action of interferon by

preventing activation of the interferon-induced eIF-2 alpha kinase. Cell 45:195-200.

53. Darweesh M, Kamel W, Gavrilin MA, Akusjarvi G, Svensson C. 2019. Adenovirus VA

RNAI Blocks ASC Oligomerization and Inhibits NLRP3 Inflammasome Activation.

Front Immunol 10:2791.

54. Radke JR, Yong SL, Cook JL. 2016. Low-Level Expression of the E1B 20-Kilodalton

Protein by Adenovirus 14p1 Enhances Viral Immunopathogenesis. J Virol 90:497-

505.

55. Ginsberg HS, Lundholm-Beauchamp U, Horswood RL, Pernis B, Wold WS, Chanock

RM, Prince GA. 1989. Role of early region 3 (E3) in pathogenesis of adenovirus

disease. Proc Natl Acad Sci U S A 86:3823-7.

56. Ginsberg HS, Horswood RL, Chanock RM, Prince GA. 1990. Role of early genes in

pathogenesis of adenovirus pneumonia. Proc Natl Acad Sci U S A 87:6191-5.

57. Ginsberg HS, Moldawer LL, Sehgal PB, Redington M, Kilian PL, Chanock RM, Prince

GA. 1991. A mouse model for investigating the molecular pathogenesis of adenovirus

pneumonia. Proc Natl Acad Sci U S A 88:1651-5.

58. Sparer TE, Tripp RA, Dillehay DL, Hermiston TW, Wold WS, Gooding LR. 1996. The

role of human adenovirus early region 3 proteins (gp19K, 10.4K, 14.5K, and 14.7K)

in a murine pneumonia model. J Virol 70:2431-9.

59. Li Y, Wold WS. 2000. Identification and characterization of a 30K protein (Ad4E3-30K)

encoded by the E3 region of human adenovirus type 4. Virology 273:127-38.

60. Robinson CM, Rajaiya J, Zhou X, Singh G, Dyer DW, Chodosh J. 2011. The E3 CR1-

gamma gene in human adenoviruses associated with epidemic keratoconjunctivitis.

Virus Res 160:120-7.

61. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version

7: improvements in performance and usability. Mol Biol Evol 30:772-80.

62. Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics

Analysis Version 7.0 for Bigger Datasets. Mol Biol Evol 33:1870-4.

63. Obenauer JC, Denson J, Mehta PK, Su X, Mukatira S, Finkelstein DB, Xu X, Wang J,

Ma J, Fan Y, Rakestraw KM, Webster RG, Hoffmann E, Krauss S, Zheng J, Zhang

Z, Naeve CW. 2006. Large-scale sequence analysis of avian influenza isolates.

Science 311:1576-80.

64. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch

S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image

analysis. Nat Methods 9:676-82.

65. Crouch SP, Kozlowski R, Slater KJ, Fletcher J. 1993. The use of ATP

bioluminescence as a measure of cell proliferation and cytotoxicity. J Immunol

Methods 160:81-8.

66. Faul F, Erdfelder E, Lang AG, Buchner A. 2007. G*Power 3: a flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. Behav Res

Methods 39:175-91.

67. Kajon AE, Erdman DD. 2007. Assessment of genetic variability among subspecies b1

human adenoviruses for molecular epidemiology studies. Methods Mol Med 131:335-

55.

68. Blanco JC, Richardson JY, Darnell ME, Rowzee A, Pletneva L, Porter DD, Prince GA.

2002. Cytokine and chemokine gene expression after primary and secondary

respiratory syncytial virus infection in cotton rats. J Infect Dis 185:1780-5.

69. Blanco JC, Pletneva L, Boukhvalova M, Richardson JY, Harris KA, Prince GA. 2004.

The cotton rat: an underutilized animal model for human infectious diseases can now

be exploited using specific reagents to cytokines, chemokines, and interferons. J

Interferon Cytokine Res 24:21-8.

70. Pletneva LM, Haller O, Porter DD, Prince GA, Blanco JCG. 2008. Induction of type I

interferons and interferon-inducible Mx genes during respiratory syncytial virus

infection and reinfection in cotton rats. J Gen Virol 89:261-270.

Table 1: Genome-wide proteotyping results for the seven HAdV-E4 strains examined in this study

| Gene | Product | Amino acid diversity | No. of variants | PG I | | | | PG II | | |
				RI-67[a] AY594253[b]	CL68578 AY594254	NY-15-4054 KY996447	NHRC 90339 EF371058	NHRC 22652 AP014843	NY-12-27440 KY996451	NHRC 42606 AY599835
E1A	13S	0.04 ± 0.04	2	v1[f]	v1	v1	v1	v2	v2	v2
E1B	19K	0.03 ± 0.03	2	v1	v1	v1	v1	v2	v2	v2
	55K	0.01 ± 0.01	3	v1	v1	v1	v1	v2	v2	v2
IX	pIX	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
IVa2	IVa2	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
E2B	DNA pol[c]	0.01 ± 0.02	3	v1	v1	v2	v2	v3	v3	v3
	pTP[d]	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
L1	52-55KDa	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
	pIIIa	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
L2	pIII	0.01 ± 0.01	3	v1	v1	v1	v1	v2	v2	v2
	pV	0.02 ± 0.02	2	v1	v1	v1	v1	v2	v2	v2
	pVII	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
	pX	0 ± 0	1	v1	v1	v1	v1	v1	v1	v1
L3	pVI	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
	hexon	0.01 ± 0.01	3	v1	v1	v3	v1	v2	v2	v2
	protease	0.02 ± 0.02	2	v1	v1	v1	v1	v2	v2	v2
E2A	DBP[e]	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
L4	100KDa	0.01 ± 0.02	6	v1	v1	v1	v2	v3	v3	v3
	33KDa	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
	pVIII	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
E3	12.1KDa	0.02 ± 0.03	3	v1	v1	v2	v2	v3	v3	v3
	CR1α	0.02 ± 0.03	2	v1	v1	v1	v1	v2	v2	v2
	gp19K	0.02 ± 0.02	3	v1	v1	v1	v1	v2	v2	v2
	CR1β	0.04 ± 0.05	3	v1	v1	v1	v2	v3	v3	v3
	CR1δ	0.04 ± 0.05	3	v1	v1	v1	v1	v2	v2	v2
	RIDα	0 ± 0	1	v1	v1	v1	v1	v1	v1	v1
	RIDβ	0.03 ± 0.03	3	v1	v1	v1	v2	v3	v3	v3
	14.7K	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
L5	fiber	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
E4	34K	0.01 ± 0.01	2	v1	v1	v1	v1	v2	v2	v2
	13.5K	0.03 ± 0.03	2	v1	v1	v1	v1	v2	v2	v2
	13.7K	0 ± 0	2	v1	v1	v1	v1	v1	v1	v1
	14.1K	0.01 ± 0.02	2	v1	v1	v1	v1	v2	v2	v2
	14.6K	0.03 ± 0.04	2	v1	v1	v1	v1	v2	v2	v2
	7.4K	0.03 ± 0.04	2	v1	v1	v1	v1	v2	v2	v2

a. Strain designation; b. GenBank accession number for whole genome sequence; c. DNA polymerase; d. precursor of the terminal protein; e. DNA binding protein; f. protein variant/ 1, 2, 3. variant number

Table 2: Localization of viral infection in cotton rat lungs during the first 3 days post infection[a]

virus	dpi[b]	Detection of HAdV hexon protein by immunohistochemistry:		
		Alveolar macrophages	Large airways	Small airways
vehicle	1-6	0/6 rats	0/6 rats	0/6 rats
PG I CL68578	1	1/6 rats	6/6 rats	6/6 rats
	2	0/6 rats	6/6 rats	6/6 rats
	3	0/6 rats	6/6 rats	6/6 rats
PG II NHRC 42606	1	6/6 rats	6/6 rats	6/6 rats
	2	6/6 rats[c]	6/6 rats	6/6 rats
	3[d]	2/5 rats	5/5 rats	4/5 rats

a. Representative images are shown in Figure 5B
b. dpi: days post infection
c. Fewer positive cells than at day 1
d. One rat in this group was found dead at 3 dpi

Bair et al., Figure 1

Bair et al., Figure 2

Bair et al., Figure 3

Bair et al., Figure 4

Bair et al., Figure 5

Bair et al., Figure 6

Bair et al., Figure 7