SARS-CoV-2 spike-glycoprotein processing at S1/S2 and S2' and shedding of the ACE2 viral

receptor: roles of Furin and TMPRSS2 and implications for viral infectivity and cell-to-cell

fusion

Rachid Essalmani[1], Jaspreet Jain[2,7], Delia Susan-Resiga[1,7], Ursula Andréo[1,2,7], Alexandra

Evagelidis[1], Rabeb Mouna Derbali[1], David N. Huynh[1], Frédéric Dallaire[2], Mélanie Laporte[2],

Adrien Delpal[3], Priscila Sutto-Ortiz[3], Bruno Coutard[4], Claudine Mapa[5], Keith Wilcoxen[5], Etienne

Decroly[3], Tram NQ Pham[2], Éric A. Cohen[2,6]* & Nabil G. Seidah[1]*

[1] **Laboratory of Biochemical Neuroendocrinology Montreal Clinical Research Institute**

(IRCM, affiliated to the Univ. of Montreal) 110 Pine Ave west, Montreal, QC, H2W1R7, Canada

[2] **Laboratory of Human Retrovirology Montreal Clinical Research Institute** (IRCM, affiliated

to the University of Montreal) 110 Pine Ave west, Montreal, QC, H2W1R7, Canada

[3] **AFMB, CNRS**, Université Aix-Marseille, CNRS, UMR 7257, Case 925, 163 Avenue de Luminy,

13288 Marseille Cedex 09, France, Marseille, France

[4] **Unité des Virus Émergents** (UVE: Aix-Marseille Univ - IRD 190 - **Inserm** 1207 - IHU

Méditerranée Infection), Marseille, France

[5] **Boston Pharmaceuticals,** Translational Research, 55 Cambridge Parkway, Suite 400,

Cambridge, MA 02142, USA

[6] **Department of Microbiology, Infectiology and Immunology**, Université de Montréal,

Montréal, Québec, Canada

[7] Each of these authors contributed equally: Jaspreet Jain, Delia Susan-Resiga and Ursula Andreo

*Corresponding authors: seidahn@ircm.qc.ca; eric.cohen@ircm.qc.ca

Short title: **Furin & TMPRSS2: spike cleavage & SARS-CoV-2 infection**

ABSTRACT

The spîke (S)-protein of SARS-CoV-2 binds ACE2 and requires proteolytic "priming" at P$\underline{R}$RA$\underline{R}_{685}\downarrow$ into S1 and S2 (cleavage at S1/S2), and "fusion-activation" at a S2' site for viral entry. *In vitro,* Furin cleaved peptides mimicking the S1/S2 cleavage site more efficiently than at the putative S2', whereas TMPRSS2 inefficiently cleaved both sites. In HeLa cells Furin-like enzymes mainly cleaved at S1/S2 during intracellular protein trafficking, and S2' processing by Furin at KPS$\underline{KR}_{815}\downarrow$ was strongly enhanced by ACE2, but not for the optimized S2' K$\underline{RR}$KR$_{815}\downarrow$ mutant (μS2'), whereas individual/double KR815AA mutants were retained in the endoplasmic reticulum. Pharmacological Furin-inhibitors (Boston Pharmaceuticals, BOS-inhibitors) effectively blocked endogenous S-protein processing in HeLa cells. Furthermore, we show using pseudotyped viruses that while entry by a "pH-dependent" endocytosis pathway in HEK293 cells did not require Furin processing at S1/S2, a "pH-independent" viral entry in lung-derived Calu-3 cells was sensitive to inhibitors of Furin (BOS) and TMPRSS2 (Camostat). Consistently, these inhibitors potently reduce infectious viral titer and cytopathic effects, an outcome enhanced when both compounds were combined. Quantitative analyses of cell-to-cell fusion and spîke processing revealed the key importance of the Furin sites for syncytia formation. Our assays showed that TMPRSS2 enhances fusion and proteolysis at S2' in the absence of cleavage at S1/S2, an effect that is linked to ACE2 shedding by TMPRSS2. Overall, our results indicate that Furin and TMPRSS2 play synergistic roles in generating fusion-competent S-protein, and in promoting viral entry, supporting the combination of Furin and TMPRSS2 inhibitors as potent antivirals against SARS-CoV-2.

IMPORTANCE

SARS-CoV-2 is the etiological agent of COVID-19 that resulted in >5 million deaths. The spike protein (S) of the virus directs infection of the lungs and other tissues by binding the angiotensin-

converting enzyme 2 (ACE2) receptor. For effective infection, the S-protein is cleaved at two sites: S1/S2 and S2'. Cleavage at S1/S2, induces a conformational change favoring the recognition of ACE2. The S2' cleavage is critical for cell-to-cell fusion and virus entry into host cells. Our study contributes to a better understanding of the dynamics of interaction between Furin and TMPRSS2 during SARS-CoV-2 entry and suggests that the combination of a non-toxic Furin inhibitor with a TMPRSS2 inhibitor could significantly reduce viral entry in lung cells, as evidenced by an average synergistic ~95% reduction of viral infection. This represents a powerful novel antiviral approach to reduce viral spread in individuals infected by SARS-CoV-2 or future related coronaviruses.

INTRODUCTION

Epidemics date from prehistoric times but are exacerbated by overcrowding and human impact on the ecosystem (1). The RNA coronaviruses (CoV) are zoonotic pathogens that occasionally spread in the human population, causing respiratory, enteric, renal, and neurological diseases (2). Electron microscopy of CoV revealed that the lipid envelope of each virion is surrounded by a "crown"-like structure (3), composed of multiple copies of a viral surface glycoprotein known as "spike" (S), which is essential for receptor binding and virus entry. Severe Acute Respiratory Syndrome coronavirus (SARS-CoV-1) and Middle East Respiratory Syndrome coronavirus (MERS-CoV) are infectious pathogenic viruses that appeared in humans at the beginning of the 21st century (2, 4). At the end of 2019, a third CoV, namely SARS-CoV-2, emerged causing widespread respiratory and vascular illnesses (5), coined COVID-19 (6).

Like envelope glycoproteins of many infectious viruses (7-9), the secretory type-I membrane-bound S of SARS-CoV-2 is synthesized as a precursor (proS) that undergoes post-transcriptional cleavages by host cell proteases at specific sites to allow viral entry. During infection, the trimeric proS (monomer, 1,272 residues) is first processed at an S1/S2 cleavage site (Fig. 1A). Unlike SARS-CoV-

1, the S-protein of SARS-CoV-2 exhibits an insertion of four critical amino acids (PRRA) at the S1/S2

junction (10-12), forming a canonical P**R**RA**R**$_{685}\downarrow$ Furin-like cleavage site (FCS). Such "priming"

step divides the protein into two subunits S1 and S2 held together by non-covalent interactions.

Following S-protein priming, the N-terminal S1-ectodomain undergoes a conformational change that

exposes its receptor-binding-domain (RBD) (13), which recognizes the ACE2 entry receptor (11). The

S2-subunit, which is responsible for the fusogenic activity of the spike-S glycoprotein, contains an

additional "fusion-activation" proteolytic site (S2') followed by an α-helical fusion peptide (FP) and

two heptad-repeat domains (HR1 and HR2) preceding the transmembrane domain (TM) and cytosolic

tail (CT) (Fig. 1A). It is thought that cleavage at S2' triggers large-scale rearrangements, including a

refolding step that is associated with the separation of S1- and S2-subunits and exposure of the

hydrophobic α-helix FP, favoring fusion of viral and host cell membranes leading to virus entry (14).

Fusion with host cells can occur either at the cell surface (pH-independent) or with internal

membranes following endocytosis (pH-dependent) (15). However, the cognate host-cell proteases

responsible for the S1/S2 and S2' cleavages vary between coronaviruses and cell types (11, 12, 16-

19).

The proprotein convertases (PCs; genes *PCSKs*) constitute a family of nine secretory serine

proteases that regulate various processes in both health and disease states (20). Through

proteolysis, PCs are responsible for the activation and/or inactivation of many secretory precursor

proteins, including virus/pathogen surface glycoproteins (9, 20). Seven PCs, including the widely

expressed Furin, PC5A, PACE4 and PC7, cleave secretory substrates at specific single/paired basic

amino acids (aa) within the motif $(K/R)-X_n-(K/R)\downarrow$, where X_n= 0, 2, 4 or 6 spacer X residues (20).

Because of their critical functions, PCs, especially Furin (21), are implicated in many viral

infections by inducing specific cleavages of envelope glycoproteins, a condition that allows not

only the fusion of the viral lipid envelope with host cell membranes (9, 20), but can also lead to

cell-to-cell fusion (syncytia), especially for viruses that undergo pH-independent fusion (22, 23).

As the S1/S2 cleavage of SARS-CoV-2 is thought to play a critical role for cellular receptor

recognition and virus entry, the efficacy and extent of this activation step by host proteases might

be a key determinant regulating cellular tropism, viral pathogenesis, and human-to-human

transmission. In contrast to SARS-CoV-1, the proS of SARS-CoV-2 contains a structurally

exposed P**R**RA**R**$_{685}$↓S**V** motif (10, 11) (Fig. 1A), which corresponds to a canonical FCS (9, 10,

20). This Furin-like motif is presumably cleaved during *de novo* virus egress (23) for S-protein

priming and may play a key role for the efficient spread of SARS-CoV-2 to various human tissues

compared to the more limited tropism of other lineage B β-coronaviruses (10, 24). Furthermore,

based on the predicted S2' KPS**KR**$_{815}$↓S**F** sequence of SARS-CoV-2, we proposed (10) that Furin-

like enzymes could also cleave at this S2' site (Fig. 1A). Indeed, various reports have since

supported the implication of Furin in the S1/S2 priming of the S-protein in human cell culture

models (12, 25, 26) and *in vivo* in mice, hamsters and ferrets (27, 28). In addition, it was also

suggested that the cell surface type-II transmembrane serine protease 2 (TMPRSS2) can enhance

fusion by cleavage at S2', but that S1/S2 cleavage is mostly Furin-dependent (18). The ability of

the Arg/Lys-specific TMPRSS2 (29, 30) to directly cleave at S2' was suggested based on the viral

entry blockade by the TMPRSS2 inhibitor Camostat (31-33), and through silencing of TMPRSS2

expression using a morpholino oligomer (18), but direct evidence of its involvement in such spike

protein processing at S2' is still lacking. Thus, it is likely that one or more proteases regulate

SARS-CoV-2 entry into human airway epithelial cells (18, 24). Furthermore, since the tissue-

expression of TMPRSS2 is restricted to a limited set of cell types compared to that of the

ubiquitously expressed Furin, the activity of the latter may widen viral tropism (34).

Thus, the major goals of the present study were to precisely define the respective roles of Furin and

TMPRSS2 in the fusion activation, and to test the consequences of their inhibition on SARS-CoV-2

infectivity and S-mediated cell-to-cell fusion. Herein, using a multi-disciplinary approach, we provide

mechanistic evidence supporting a critical role of the proprotein convertase Furin in the processing of

SARS-CoV-2 spike protein. Specifically, we map the exact S2' processing site by proteomics and

highlight by mutagenesis the functional importance of S1/S2 and S2' regions in viral entry and cell-

to-cell fusion. For the first time, we demonstrate that three novel cell-permeable small molecules

inhibitors of proprotein convertases developed by Boston Pharmaceuticals, referred hereafter as BOS-

inhibitors, can potently inhibit proS processing at S1/S2 and S2' by endogenous Furin-like proteases

leading to efficient inhibition of viral entry, viral replication, and cell-to-cell fusion. Finally, our work

sheds a new light on the role of TMPRSS2 in promoting ACE2 shedding and enabling S2' processing,

thereby leading to enhanced cell-to-cell fusion.

RESULTS

Comparative analysis of cleavage of SARS-CoV-2 peptides mimicking S1//S2 and S2'

processing sites by Furin and TMPRSS2. Furin is thought to be important in the processing of

SARS-CoV-2 spike-glycoprotein (S) at the S1/S2 site (10, 24) while TMPRSS2 has been proposed to

have an important role in activating S at S2' (18, 31-33) (Fig. 1A). Nevertheless, the relative

contributions of Furin and TMPRSS2 towards cleavage of SARS-CoV-2 S glycoprotein at both sites

remain poorly defined. Thus**,** the susceptibility of SARS-CoV-2 S glycoprotein to Furin-cleavage

was first assessed *in vitro*. Incubation of quenched fluorogenic peptides encompassing S1/S2 and

S2' sites (Supporting Information SI-Table 1) demonstrated that the S1/S2 site of SARS-CoV-2 S

was efficiently cleaved by 2 nM Furin at pH 7.5 (Fig. 1B), whereas the S1/S2 site of SARS-CoV-1,

which lacks an FCS, was not cleaved (Fig. 1B). Furin less efficiently cleaved the SARS-CoV-2-mimic

peptide at S2', requiring 50-fold higher enzyme concentrations (100 nM) to detect cleavage (inset Fig.

1B). The high specificity of the SARS-CoV-2 for processing at Furin-like motifs was next confirmed

by demonstrating that substitutions of basic residues at the S1/S2 cleavage site (RRA$\underline{A}_{685}\downarrow$S,

$\underline{A}$RA$\underline{A}_{685}\downarrow$S, $\underline{A}$RAR$_{685}\downarrow$S) dramatically impaired S1/S2 cleavage (Fig. 1B). Altogether, these data

demonstrate that *in vitro* Furin best cleaves at S1/S2 and less efficiently at S2'. In contrast, TMPRSS2

did not efficiently cleave the S1/S2 and S2' peptides (Fig 1C). The cleavage at S1/S2 became

detectable only when TMPRSS2 was present at high concentration (50 nM). However, different from

Furin, under this condition TMPRSS2 cleavage of peptides mimicking the S1/S2 Ala-mutants RRA$\underline{A}$,

$\underline{A}$RA$\underline{A}$ and $\underline{A}$RAR and S2' was also evident (Fig. 1C). Taken together, these data emphasize the

critical importance of the P1 and P4 Arg for Furin-mediated cleavage at S1/S2 and suggest that the

likely Arg-motif recognized by TMPRSS2 is either (Ala/Arg)-Arg$\downarrow$Ala or Ala-Arg$\downarrow$Ser with a

preference for Ala at P2 over Arg, and Ala or Ser at P1'.

Furin and Furin-like proteases can process proS at S1/S2 and S2' sites. To examine the ability

of Furin and Furin-like enzymes to process the precursor proS of SARS-CoV-2 *in cellulo,* we used a

HeLa cells model, which endogenously express Furin but not TMPRSS2 or ACE2 (*not shown*), as

reported earlier (35). Here, we found that endogenous enzymes efficiently processed a V5-tagged proS

(Fig. 1A), likely at the S1/S2 junction to generate a ~100 kDa S2-like fragment (Fig. 2A). Interestingly,

when the proprotein convertases PC5A, Furin, PC7 or PACE4 were transiently overexpressed,

cleavage at this site became more prominent. Of note, a partial knockdown of Furin decreased S2

levels by more than 60% (SI-Fig. 1A). Furthermore, it was only when Furin or PC5A were

overexpressed that cleavage at a potentially S2' site was noticeable, yielding a ~75 kDa fragment (Fig.

2A). The remaining ~200 kDa proS$_{im}$ corresponded to an immature precursor form that had not exited

the ER, as attested by its sensitivity to both endoglycosidase-F and endoglycosidase-H (SI-Fig. 1B)

and insensitivity to Furin-like convertases, which are only active in the TGN and/or cell

surface/endosomes (20, 36).

To precisely define the ~100-kDa fragment, we mutated the S1/S2 site **R**RA**R**$_{685}$↓S and found that

the double Ala-mutant [**A**RA**A**$_{685}$] (denoted μS1/S2) abrogated processing at S1/S2 and putative S2'

(Fig. 2B), highlighting once again the importance of the P4- and P1-Arg for recognition by Furin-like

enzymes (20). The loss of Furin-like cleavage at S1/S2 resulted in accumulation of a higher molecular

size band (~230 kDa), representing mature proS, that exited the endoplasmic reticulum (ER), as

confirmed by its resistance to endoglycosidase-H, while still sensitive to endoglycosidase-F digestion

(SI-Fig. 1B).

To further define the Arg-residues critical for processing at S1/S2, we assessed the effect of single

residue mutations: R682A, R685A and S686A and confirmed the critical importance of P1-Arg$_{685}$ or

P4-Arg$_{682}$ for the generation of S2 by endogenous Furin (SI-Fig. 1C). However, unlike μS1/S2 (Fig.

2B), these single mutants were partially cleaved by overexpressed Furin (SI-Fig. 1C), reflecting the

multi-basic nature of the S1/S2 recognition sequence and suggesting the importance of the P3 Arg$_{683}$

(37). The S686A mutant was based on the prediction that Ser$_{686}$ could be O-glycosylated (38), which

may hamper processing at S1/S2 (39). However, like the WT-S, the S686A mutant was efficiently

processed by Furin into S2 and S2' (SI-Fig. 1C), suggesting the lack of O-glycosylation at Ser$_{686}$ in

HeLa cells.

We next used site-directed mutagenesis to identify the exact S2' site cleaved by Furin. However,

K814A, R815A and K814R815A mutants at P1 and P2 residues (20) of the predicted S2' site

(KPS**KR**$_{815}$↓SF) altered S-protein trafficking resulting in a predominantly ER-retained proS$_{im}$

protein, especially for the R815A and the double mutant (SI-Fig. 1D). Therefore, we resorted to

mass spectrometry analysis of proteins migrating at the S2' position to unambiguously identify

this site. The peptides generated by a Lys-specific protease (K$_{814}$↓) allowed for the discrimination

between SFIEDLLFN$\underline{K}_{825}$ that would be generated if Furin cleaved at Arg$_{815}$, and $\underline{R}_{815}$SFIEDLLFN$\underline{K}_{825}$ that would be derived from N-terminally extended proteins, e.g., S2. Proteomic data (Fig. 2C) revealed a >50-fold higher ratio of $\underline{S}$FIEDLLFN$\underline{K}_{825}$ to $\underline{R}$SFIEDLLFN$\underline{K}_{825}$ for the S2' product, demonstrating that the N-terminus of S2' starts at Ser$_{816}$ and that Furin cleaves after Arg$_{815}\downarrow$ in the sequence KPSK$\underline{\mathbf{R}}_{815}\downarrow$SFIEDLLFNKVT (Fig. 1A). To further demonstrate the role of Furin in S2' cleavage we generated a Furin-optimized S2' site (called µS2') with a polybasic sequence K$\underline{RR}$KR$_{815}\downarrow$SF in proS and found that this derivative was very efficiently cleaved by endogenous and especially by overexpressed Furin, yielding a similar ~75 kDa fragment (Fig. 2D). Altogether, these data demonstrate that the S2' cleavage occurs at Arg$_{815}\downarrow$ and further reveal that this site can be partially processed by overexpressed Furin and/or PC5A.

Processing at S2' by Furin is enhanced in the presence of ACE2. Immunocytochemical analyses of HeLa cells co-expressing S or µS1/S2-S and ACE2 showed that both S-proteins and ACE2 co-localized at the cell surface independent of the state of proS processing (SI-Fig. 2). Given that binding of SARS-CoV-2 S-trimer to the dimeric ACE2 receptor has been proposed to trigger a conformational change in S1, promoting cleavage at S2' (13, 40), we next examined whether this phenomenon would effectively increase/promote S2' processing by Furin. To this end, we expressed the V5-tagged proS spike protein together with ACE2 and Furin in HeLa cells. We found that while not significantly affecting S1/S2 cleavage, ACE2 expression seemed to stabilize the S2 subunit and to strongly enhance the generation of S2' by endogenous and overexpressed Furin (Fig. 3A). Amazingly, in the presence of ACE2, the µS1/S2-S, which is otherwise resistant to cleavage at S1/S2 by endogenous or overexpressed Furin (Fig. 2B), can be partially cleaved directly into S2' by

overexpressed Furin (Fig. 3B, last lane). Taken together, we conclude that binding of S-protein to

ACE2 likely facilitates exposure of the S2' site (41), thereby enhancing Furin processing at S2'.

Furin-inhibitors block S1/S2 and S2' cleavages. Given the importance of Furin in proS

processing at the S1/S2 and S2' sites, we next evaluated the activity of three novel non-toxic, cell-

permeable Furin-like inhibitors developed by Boston Pharmaceuticals available as oral (BOS-981,

BOS-318) or inhalable (BOS-857) formulations (Fig. 4A). Accordingly, we first tested *in vitro* the

efficacy and selectivity of these inhibitors on purified soluble forms of Furin, PC5A, PACE4 and

PC7 using a quenched fluorogenic substrate FAM-Q**R**V**RR**AVGIDK-TAMRA. As shown, the

inhibitors effectively blocked substrate processing by all convertases with an IC_{50} of ~7-9 nM

compared to ~9-10 nM for the known cell-permeable PC-inhibitor decanoyl-RVKR-

chloromethylketone (dec-RVKR-cmk) (42, 43) (Fig. 4B). The Furin S1/S2 cleavage was also

validated using a 12-residue quenched fluorogenic substrate DABSYL/Glu-TNSP**RRAR**↓SVAS-

EDANS mimicking the S1/S2 priming site. The inhibition deduced after hill-plot curve fitting (Fig.

4C) gave an estimated IC_{50} of 4 ± 0.7 nM for BOS-981, 32 ± 4 nM for BOS-857 and 35 ± 5 nM

for BOS-318. As well, BOS-inhibitors inhibited endogenous Furin-like processing of a dibasic

bone morphogenic protein 10 (BMP10)-mimic (43) with an IC_{50} of ~8 nM *versus* 5 nM for the

dec-RVKR-cmk as determined by a cell-based Golgi imaging assay with U2OS cells (Fig. 4D).

We further showed that BOS-inhibitors efficiently blocked S1/S2 and S2' processing by

endogenous Furin-like enzymes, resulting in a near complete inhibition at 0.3 μM, also obtained

with 50 μM of dec-RVKR-cmk (RVKR; Fig. 4E). Overall, our data clearly demonstrate a role of

Furin in the processing of proS at the S1/S2 and S2' sites.

Furin-like inhibitors reduce virus production in SARS-CoV-2-infected cells. We next examined whether blocking the processing of proS by BOS-inhibitors modulates SARS-CoV-2 infection. Indeed, in lung derived Calu-3 cells pretreated with 1 µM BOS-inhibitors for 24h before infection, we observed significantly decreased viral titers at 12, 24 and 48h post-infection (Fig. 5A). Importantly, the inhibitory effect was dose-dependent, reducing viral burden up to >30-fold with 1 µM BOS-318 (Fig. 5B; left panel). As well, the IC_{50} and selectivity index (44) of BOS-318 were 0.2 µM and 475, respectively (Fig. 5B; right panel). Importantly, the levels of spike (full length and cleaved S) and nucleocapsid proteins in the supernatant and cells were decreased in a dose-dependent manner (Fig. 5C), underscoring the crucial role played by Furin-like convertases in SARS-CoV-2 infection in this lung epithelial cell model. A similar analysis with BOS-857 and BOS-981 revealed comparable antiviral effects and selectivity index (SI-Figs. 3A, B). In addition, BOS-inhibitors were also evaluated in Vero E6 cells where SARS-CoV-2 entry and infection is established primarily *via* the endocytic pathway (11, 24). In this system and as expected, treatment with BOS-inhibitors led to weaker effect since we observed a decreased virus production by only ~2.6-5.7-fold (SI-Fig. 4), possibly reflecting a role of Furin-like activity in early endosomes (36) for pH-dependent virus entry in Vero cells.

Since TMPRSS2 has been proposed to be important for viral entry at the plasma membrane, we next determined whether combining BOS-inhibitors and the TMPRSS2-inhibitor Camostat would have a synergistic effect, leading to a more pronounced antiviral effect in Calu-3. As shown, although these compounds could reduce viral replication individually, their co-treatment resulted in a synergistic inhibition of ~95 ± 2.5% (>70-fold) of progeny infectious viruses (Fig. 5D and SI-Fig. 5), reinforcing the importance of both Furin-like proteases and TMPRSS2 in promoting efficient SARS-CoV-2 infection of Calu-3 cells.

Furin-like inhibitors reduce viral entry by blocking processing of proS during biosynthesis and at the viral entry site. The more dramatic impact observed with Furin-like inhibitors on virus infection of Calu-3 cells *versus* Vero cells suggests that these inhibitors affect mainly the pH-independent entry mechanism. Thus, we next assessed the effect of BOS-inhibitors on viral entry. Using nanoluciferase-expressing HIV particles pseudotyped with WT, μS1/S2 or μS2' S-proteins, we observed that the viral entry of μS1/S2-S pseudovirions is ~10-fold reduced in Calu-3 cells (Fig. 6A). In contrast, all three pseudotyped viruses were at least 10-fold more infectious in HEK293T-ACE2 cells, suggesting that S-priming at S1/S2 is required for optimal viral entry in Calu-3 cells, but dispensable or perhaps even less desirable in HEK293T-ACE2 cells (Fig. 6A). Since these findings were similar to those in Vero E6 cells (45), we surmise that viral entry in HEK293T-ACE2 cells was through the same pH-dependent, endocytic route as reported for Vero cells (46, 47). This agrees with the fact that HEK293 cells allow endocytosis of SARS-CoV-2 pseudovirions *via* clathrin-coated vesicles (48). In the case of μS2'-S-expressing viral particles, entry was more efficient in both cell types (compare the same-coloured dots between WT-S and μS2'-S in absence of BOS, Fig. 6A), implying that S cleavage at the S2'optimized Furin-like site could enhance viral entry. When BOS-318 was present during biosynthesis of pseudovirions, processing of WT-S and μS2'-S was blocked (Fig. 6B), leading to reduced viral entry in Calu-3 by ~3.6- to ~12.5-fold, respectively (Fig. 6A). Thus, BOS-318 treatment phenocopied the effect of the μS1/S2 in both cell types. Nevertheless, under this condition viral entry in HEK293T-ACE2 cells was enhanced by ~10-fold for WT-S and ~2-fold for μS2'-S (Fig. 6A), suggesting that viral entry by the pH-dependent pathway does not require FCS processing.

Having observed the negative effect of BOS-318 on S processing by particle-producing cells, we asked whether pre-treating target cells with BOS-318 would also affect entry of SARS-CoV-2 pseudoparticles. In Calu-3 cells, where viral entry occurs primarily through fusion at the plasma

membrane (46), we observed reduced viral entry by ~3.8-fold for WT-S and ~14-fold for µS2'-S (Fig.

6C). This emphasizes a significant contribution of Furin to the processing and priming of S at the

plasma membrane of Calu-3 cells after ACE2 recognition. The fact that BOS-318 had a more

pronounced effect on entry by µS2'-S-containing viral particles was not surprising given the very

efficient S2' processing of this mutant by Furin (Fig. 2D). Of note, viral entry by WT-S pseudoparticles

was more affected by 100 µM Camostat compared to that by µS2'-S viral particles (Fig. 6C). Here,

entry was reduced by ~8.6-fold for WT-S and ~3.2-fold for µS2'-S, suggesting that Furin plays a more

prominent role in the entry of µS2'-S *versus* WT-S viral particles. Lastly, the combined pre-treatment

of both BOS-318 and Camostat led to a complete block of viral entry, highlighting the importance of

both Furin and TMPRSS2 in mediating viral fusion at the plasma membrane.

Furin-like inhibitors decrease cell-to-cell fusion and syncytia formation. To assess whether

BOS-inhibitors also affect cell-to-cell fusion, we developed a co-culture assay in which donor HeLa

cells express HIV Tat and the fusogenic S-protein, while acceptor HeLa TZM-bl cells express ACE2

and Tat-driven luciferase (49) (Figs. 7A). As a proof-of-principle, we showed that when donor HeLa

cells expressing HIV gp160 and Tat fused with acceptor cells expressing CD4 (SI-Fig. 6A, panel b),

luciferase activity was increased compared to that observed in TZM-bl control cells co-cultured with

donor Hela cells expressing only Tat (SI-Fig. 6B). The expression of S-protein alone in donor HeLa

cells did not induce fusion with acceptor TZM-bl control cells (SI-Fig. 6B). However, ACE2

expression in TZM-bl allowed fusion with HeLa-expressing S-protein (SI-Fig. 6A, panel c; 6B) in a

dose-dependent manner (SI-Fig. 6C), but no fusion was observed with µS1/S2-S (SI-Fig. 6A, panel

d). Indeed, the linearity of our assay (correlation coefficient of 0.87) validated the use of luminescence

as an indicator of cell-to-cell fusion (SI-Fig. 6C). Using this assay, we found that while donor cells

expressing WT-S led to syncytia formation (Fig. 6B) and a >10-fold increased cell-to-cell fusion

compared to control (empty vector V, no S) cells (Fig. 6C), donor cells expressing μS1/S2-S did not

promote any cell fusion even in the presence of ACE2 (Figs. 6C, D). Thus, the μS1/S2-S phenocopies

the effect of BOS-inhibitors on cell-to-cell-fusion (Fig. 6C), whereby absence of Furin-activity would

not allow fusion and demonstrates a key role of S1/S2 cleavage in S-mediated cell-to-cell fusion.

Consistent with this finding, we observed an almost complete loss of cell fusion when donor cells were

treated with BOS-inhibitors or the PC-inhibitor decanoyl-RVKR-cmk (RVKR; Fig. 6C) (43),

emphasizing the critical role of Furin-cleavage in promoting ACE2-dependent cell-to-cell fusion in the

context of acceptor cells that do not endogenously express TMPRSS2, such as HeLa cells. This fusion

assay also enabled the assessment of the effects of some worldwide-spreading S-protein variants of

SARS-CoV-2, which seem to affect viral traits such as transmissibility, pathogenicity, host range, and

antigenicity of the virus (50, 51). Among these, we selected mutants that modify the Pro at the P5

position of the S1/S2 site (Fig. 1A), i.e., the P681H and P681R associated with the α- and δ- variants,

respectively (9). Our data showed that while the μS2' mutant did not affect cell-to-cell fusion, the

P681H and P681R mutants significantly enhanced it by ~2-fold (Fig. 7D), in line with the higher

transmissibility of the associated α- and δ- SARS-CoV-2 variants (50, 51) and increased cell-to-cell

fusion (52).

TMPRSS2 promotes cell-to-cell fusion in the absence of Furin-mediated cleavage at S1/S2.

Having shown the critical role of Furin-cleavage in promoting ACE2-dependent cell-to-cell fusion in

HeLa cells that do not express TMPRSS2, we next examined the importance of TMPRSS2 in cell-to-

cell fusion given its significant role in viral entry and replication (Figs. 5D and 6C). Thus, we analyzed

fusion between donor cells expressing WT-S or μS1/S2-S (double tagged S-protein: HA at the N-

terminus and V5 at the C-terminus) and acceptor cells expressing ACE2 in the absence or the presence

of TMPRSS2 (Fig. 8). As shown, in the presence of ACE2, TMPRSS2 reduced fusion by ~2.1-fold in

the case of WT-S but increased it by ~2.4-fold in the case of μS1/S2-S (Fig. 8A). Under the same

conditions, WB analyses showed that for WT-S, the TMPRSS2 expression decreased by ~4-fold (3.8

versus 0.9) the relative levels of S2 and modestly increased by ~1.5-fold (0.6 *versus* 0.9) the relative

levels of S2'. In contrast, for μS1/S2-S, expression of TMPRSS2 increased S2' levels by ~7-fold (0.1

versus 0.7) without any detectable change in S2 (0.6 *versus* 0.5) (Fig. 8B). Therefore, in the presence

of ACE2, the levels of S2 correlate with cell-to-cell fusion in the case of WT-S that is well cleaved at

S1/S2 by Furin. In contrast, fusion correlates best with S2' levels, when cleavage at S1/S2 is limited

(μS1/S2-S). Interestingly, secretion of the HA-tagged, N-terminal S1 subunit (aa 14-685; Fig. 1A) was

more pronounced for WT-S compared to μS1/S2-S (Fig. 8B). Yet, in both cases, the extent of S1

release was not modulated by the presence of TMPRSS2, suggesting that TMPRSS2-mediated

enhanced fusion of μS1/S2 was not a consequence of altered S1/S2 cleavage (Fig. 8B).

Mechanism by which TMPRSS2 promotes cell-to-cell fusion. To assess whether TMPRSS2

promoted cell-to-cell fusion by enhancing S2' cleavage in the complete absence of S1/S2 priming, we

generated a new S-derivative lacking all Arg at S1/S2 (Fig. 1), namely μAS1/S2 (AAAA$_{685}$)-S, and

hence different from μS1/S2 (Fig. 1C; ARAA mutant), it would not be cleaved by TMPRSS2 or Furin.

Cell-to cell fusion was assessed following incubation of cells expressing WT-S or μAS1/S2-S with

acceptor cells expressing ACE2 and/or TMPRSS2. In this context, TMPRSS2 reduced fusion of WT-

S by ~2.6-fold and enhanced that of μAS1/S2-S by ~3.1-fold (Fig. 9A), as previously observed with

the μS1/S2 mutant (Fig. 8A). Camostat completely restored TMPRSS2-reduced fusion with WT-S and

largely attenuated the TMPRSS2-enhanced fusion with μAS1/S2 (Fig. 9A). This revealed that in the

presence of ACE2, TMPRSS2 could enhance fusion in absence of S1/S2 priming. We next assessed

whether this effect could be related to the differential processing of the S protein and/or to ACE2

receptor shedding by TMPRSS2. The WB of cell lysates shows that treatment of acceptor cells with

Camostat eliminated the TMPRSS2-induced reduction in S2 observed with WT-S (Fig. 9B), yet it did

not significantly alter S2' levels either with WT-S or μAS1/S2 (Figs. 9B, C). In contrast, the WB of

ACE2 reveal that TMPRSS2 strongly reduced the levels of mature membrane-bound ACE2 migrating

at ~120 KDa, without affecting the lower molecular weight protein which correspond to an immature

ER-retained form of ACE2 (ACE2$_{im}$, Figs. 9B, C) that is sensitive to endoH digestion (*not shown*).

These results show that ACE2 is shed by TMPRSS2, as proposed in earlier studies (33). As expected,

we observed that ACE2 shedding by TMPRSS2 is almost completely blocked by Camostat (compare

ACE2+TMP lanes DMSO to CAM) (Figs. 9B and C).

In contrast, the WB anti ACE2 reveal that TMPRSS2 strongly reduced the levels of mature

membrane-bound ACE2 migrating at ~120 KDa, without affecting the lower molecular weight protein

which correspond to an immature ER-retained form of ACE2 (ACE2$_{im}$, Figs. 9B, C) that is sensitive

to endoH digestion (*not shown*). These results suggest that ACE2 is shed due to TMPRSS2 cleavage

as proposed in earlier studies (33). As expected, we observed that this effect of TMPRSS2 on ACE2

processing is almost completely blocked by Camostat (compare ACE2+TMP lanes DMSO to CAM)

(Figs. 9B and C). Finally, by analysing the release of S1 in the cell culture media we observed that

TMPRSS2-induced release into the media of a longer ~175 kDa fragment from μAS1/S2, referred

hereafter as S1$_L$ (Fig. 9C), which is not detected in presence of Camostat. The molecular weight of the

S1$_L$ indicates that it corresponds to a TMPRSS2 cleavage of μAS1/S2-S at S2' (Fig. 9B).

These results suggest that TMPRSS2 modulation of fusion is complex, as TMPRSS2 alters both

the integrity of ACE2 receptor, but also participates in S-processing as attested by the release of S1$_L$

which corresponds to the N-terminal cleavage product of μAS1/S2-S at S2'. Interestingly, we did not

detect an associated increase in S2' levels, suggesting that after cleavage the fusion process occurs and

the S2 protein is subsequently degraded, as suggested by a small degradation product seen on WB

(arrow in Figs. 9B, C). We surmise that in the presence of ACE2, TMPRSS2 facilitates the secretion

of the N-terminal fragment S1$_L$ generated by cleavage at S2'.

Our next goal was to further decipher the contribution of ACE2 cleavage by TMPRSS2 and the

seemingly opposite effects of TMPRSS2 on fusion activity with WT-S and μAS1/S2-S (and μS1/S2-

S). Hence, since TMPRSS2 cleaves at single Arg and Lys residues (53), we tested the effect of multiple

Arg/Lys to Ala mutants in segments close to the transmembrane domain of ACE2 (C0 + C4 in Fig.

10A), which were previously proposed to limit the TMPRSS2-induced shedding (54). Accordingly,

WB analyses (Fig. 10B) revealed that TMPRSS2, but not its Ser* active site S441A mutant (μTMPR,

Fig. 10B; SI-Fig. 7) sheds the membrane-bound ~120 kDa WT ACE2, releasing a major ~95 kDa and

a minor ~80 kDa form into the media. In contrast, TMPRSS2 primarily generated the ~80 kDa

fragment from the ACE2 (C0+C4) mutant. It should be mentioned that in the absence of TMPRSS2

the extent of fusion for both WT-S and μAS1/S2-S was comparable between ACE2 (WT) and ACE2

(C0+C4). Interestingly, in the context of the ACE2 (C0+C4) receptor mutant, we found that TMPRSS2

no longer modulates the fusion of both WT-S and μAS1/S2-S (Fig. 10C). Importantly, and different

from WT ACE2 (Figs. 8, 9), TMPRSS2 did not reduce the levels of S2 when ACE2 (C0+C4) was

used (Fig. 10D), nor did it reduce fusion (Fig. 10C). Additionally, the TMPRSS2-enhanced secretion

of the ~ 175 kDa S1$_L$ fragment in the absence of S1/S2 cleavage (Fig. 9C; SI-Fig. 7), was no longer

detected in the media when ACE2 (C0+C4) was expressed on acceptor cells (Fig. 10D), correlating

with the lack of fusion of μAS1/S2-S (Fig. 10C). Altogether, these results suggest for the first time that

in the absence of Furin-mediated priming of S, TMPRSS2 promotes cell-to-cell fusion by generating

a ~95 kDa sACE2 that likely binds the S1$_L$ cap (55) and favors its release.

DISCUSSION

Furin and TMPRSS2 have both been proposed to process the SARS-CoV-2 spike protein and promote viral entry and infection. In this study, we functionally characterize the role of Furin and TMPRSS2 and show their complementary role in SARS-CoV-2 mechanism of entry and infection.

Consistent with previous data (18, 33, 56, 57), processing at S1/S2 is required for optimal viral entry in Calu-3 lung epithelial cells. Indeed, mutation of the arginines of the FCS or blocking S1/S2 cleavage by a series of novel Furin-inhibitors effectively reduced SARS-CoV-2 entry in Calu-3 cells. In the context of cell-to-cell fusion, mutations of the FCS impaired syncytia formation in HeLa cells in the absence of TMPRSS2 in acceptor cells, and Furin-inhibitors phenocopied the effect of these S1/S2 mutations. These results highlight the importance of Furin in promoting S-mediated cell-to-cell fusion. In contrast, in HEK293T cells, where SARS-CoV-2 likely enters *via* endocytosis in acidic endosomes, mutations of the FCS enhanced entry (Fig. 6), suggesting that cleavage at S1/S2 by Furin was not required for efficient S-processing by endosomal proteases (e.g., cathepsins) (58, 59). The pH-dependent entry pathway is sensitive to drugs increasing the pH of endosomal compartments, e.g., hydroxy-chloroquine (47). Nevertheless, this drug failed to show significant improvement in COVID-19 patients (60, 61) or infected animals (62), supporting the notion that the virus can effectively utilize alternative entry pathways that are critical for virus entry, transmission and pathogenicity. The fact that Furin-cleavage at S1/S2 is conserved in SARS-CoV-2 isolated from COVID-19 patients and that ΔPRRA viruses are poorly infectious in hamsters (60), suggest that blocking viral entry through the pH-independent pathway is a viable approach towards thwarting SARS-CoV-2 dissemination. In this context, the non-toxic, small molecules BOS-inhibitors that were analyzed in this study, which can be delivered orally or by inhalation, deserve consideration as potential antivirals against acute SARS-CoV-2 infection. As observed in adult animal models, short-term inhibition of Furin would not cause severe side effects, despite the many physiological functions of this enzyme (20).

Our results also demonstrate that Furin can also process S at S2', a site that we map by proteomics

to be at **KR**$_{815}\downarrow$SF. The latter S2' cleavage site was confirmed by introducing an optimized S2' site

(μS2', KRRKR$_{815}\downarrow$SF) that was efficiently cleaved by Furin, yielding a protein fully competent

for fusion in pseudotyped experiments. Importantly, we showed that the S2' processing is strongly

increased when ACE2 receptor is expressed. Our results suggest that the recognition of the ACE2

receptor by the spîke protein induces a conformational change of the S2 domain and enhances

cleavage of S2' site by cellular proteases such as Furin.

In contrast TMPRSS2 appears to cleave synthetic peptides encompassing the S1/S2 and S2'

cleavage sites and proS protein in Hela cells less efficiently compared to Furin. TMPRSS2 has been

reported to enhance the infectivity of SARS-CoV-1 and MERS-CoV *via* cleavages of the S-protein

(53, 63), as reviewed in (4, 15, 23). The S1/S2 site VSLL**R**$_{667}\downarrow$ST of SARS-CoV-1 contains an

Arg$\downarrow$Ser cleavage motif that is propitious to cleavage by TMPRSS2 (53) but not by Furin. However,

S1 production was minimal and fusion was observed only when donor cells expressing S-protein were

co-cultured with acceptor cells expressing ACE2 and TMPRSS2 (53). In SARS-CoV-2 the S1

released, presumably following cleavage of S at S1/S2 and ACE2 binding, is much reduced with

μS1/S2-S compared to WT-S, indicating that Furin-mediated cleavage at S1/S2 promotes a more

efficient release of S1 (Fig. 8B). Additionally, in a co-culture system the relative levels of soluble S1

fragment were not affected by the presence or absence of TMPRSS2 for WT and μS1/S2 (Fig. 8B).

This suggests that the differential effect of TMPRSS2 on cell-to-cell fusion could not be attributed to

its direct cleavage activity on S. Rather, we assert that TMPRSS2 primarily modulates cell-to-cell

fusion in part *via* shedding of ACE2. In the presence of TMPRSS2 the extent of fusion would depend

on a combination of reduced levels of full-length cell surface ACE2 and increased levels of sACE2

that could act as a decoy to inhibit fusion (64). In that context, a human recombinant sACE2 that

includes the collectrin domain but lacks the TM domain (hrsACE2, aa 1-740) (see Fig. 10A) effectively

blocked SARS-CoV-2 infection of Vero E6 cells (55). Indeed, our cell-to-cell fusion assay tested in

co-culture revealed that the ~95 kDa sACE2 primarily produced by TMPRSS2 cleavage of ACE2

resulted in lower levels of cellular ACE2 (Figs. 9B, 10B) and in impaired WT-S induced fusion (Figs.

9A, 10C). In contrast, the mutant ACE2 (C0+ C4) was no longer sensitive to TMPRSS2 impairment

of WT-S-induced fusion, suggesting that the ~80 kDa form of sACE2 primarily produced by

TMPRSS2 cleavage of ACE2 (C0+C4), likely upstream of Lys_{619}, may no longer be able to inhibit

WT-S-fusion (Fig. 10C), but may still bind S1 as shown by cryoEM studies that used a short ACE2

ectodomain (aa 19-615) (40). Altogether, these results suggest that the C-terminal collectrin-like

domain (65, 66) of ACE2 (aa 616-768) (Fig. 10A), which is lost in the ~80 kDa form, may be critical

for the ability sACE2 to inhibit cell-to-cell fusion of WT-S. Since Arg_{710} and Arg_{716} within the

collectrin-like domain (Fig. 10A) have been reported to be implicated in ACE2 dimerization (66), their

Ala mutation in the ACE2 (C0+C4) mutant should significantly impair ACE2 dimerization. Whether

dimerization of ACE2 and the ~95 kDa sACE2 (Fig. 10B) are needed for efficient inhibition of WT-

S-induced fusion (64) is yet to be confirmed.

In the context of endogenous expression of ACE2 and TMPRSS2, e.g., in Calu-3 cells (67, 68), our

data show that Camostat significantly reduces SARS-CoV-2 infectivity (Fig. 6C), revealing that a

TMPRSS2-like activity favors viral entry. Similarly, in the presence of BOS-inhibitors or absence of

Furin cleavage at S1/S2, TMPRSS2 enhances cell-to-cell-fusion (Figs. 8A, 9A). HeLa cells expressing

WT-S in donor cells and both ACE2 and TMPRSS2 in acceptor cells, exhibited reduced cell-to-cell

fusion, a process inhibited by Camostat. This suggests that the relative contribution of TMPRSS2 to

the shedding of ACE2 into sACE2 and cleavage of S-protein at S2' is cell-type dependent. Future

experiments may unravel the underlying mechanism that would explain the difference between the

HeLa and Calu-3 cells. Our study highlights a complex dynamic between spike, ACE2, Furin and

TMPRSS2 (Fig. 11) and points to a potential role of TMPRSS2, which in the absence of S1/S2

processing (e.g., µAS1/S2 or in presence of BOS-inhibitors) can by shedding ACE2 facilitate S2'

cleavage and cell-to-cell fusion.

The human airway epithelium is an important site of early SARS-CoV-2 infection (18, 24, 69). The

virus can then disseminate to other tissues/cells such as gut, liver, endothelial cells and macrophages

where ACE2, Furin and TMPRSS2 are co-expressed (70). While Furin is mostly found in the TGN, it

is also present in endosomes and on apical/basolateral plasma membranes in polarized cells such as

those of the lung, small intestine and kidney (71). In contrast, TMPRSS2 is mostly found at the apical

membrane of secretory epithelia (72), suggesting that both enzymes would be poised to process the S-

protein on the apical side. Their relative abundance may be an important factor governing which of the

two might cleave S at the S2' cleavage site. The complementarity and interchangeability of these

different proteases, together with those in endosomes, likely allows SARS-CoV-2 to exhibit a wider

tropism compared to SARS-CoV-1 (73). In this context, we note that randomized placebo controlled

clinical trials using an orally administered TMPRSS2 inhibitor Camostat mesilate three times a day for

treatment of COVID-19 patients did not show improvement in outcomes (74). We believe that the

maximal benefit of TMPRSS2 inhibition could be achieved when Furin activity is also inhibited, since

our results showed that TMPRSS2 activity is preponderant in the context of absence of cleavage at

S1/S2. Furthermore, the intranasal delivery of both agents would be expected to have less side effects

and be more effective, as it would directly target the airway epithelia of the nose and the lungs, the

major sites of SARS-CoV-2 entry.

Altogether, our results strongly support the notion that a combination of BOS- and selective

TMPRSS2-inhibitors would provide a more effective blockade against SARS-CoV-2 infection

(Figs. 5D, 11). It would now be interesting to validate *in vivo* whether a combination of a BOS-

inhibitor and a more potent/selective TMPRSS2-inhibitor (75) together with hydroxy-chloroquine

(56, 76) would synergize the antiviral effect of these entry inhibitors.

The availability for worldwide distribution of various SARS-CoV-2 vaccines that inhibit the

accessibility of the RBD of S-protein to ACE2 (https://www.raps.org/news-and-articles/news-

articles/2020/3/covid-19-vaccine-tracker) represents a major advance to limit the spread of SARS-

CoV-2 infections. However, it is still not known with certainty whether they will be effective in

patients with impaired immune systems, and whether they will confer a persistent protection

against new variants of SARS-CoV-2. If the protective effect of the vaccination remains

incomplete, effective antiviral drugs are still needed and could help with early diagnosis of the

disease. Ultimately, in case of new emerging coronavirus pandemics (77), the availability of such

treatments would constitute a powerful anti-viral arsenal to be used in pandemic preparedness.

MATERIALS AND METHODS

Enzymatic PC-inhibition by BOS-inhibitors

Biochemical assay: The proprotein convertases Furin (108-574-Tev-Flag-6His), PC5A (PCSK5; 115-63-Tev-Flag-6His), PACE4 (PCSK6; 150-693-Tev-Flag-6His), and PC7 (PCSK7; 142-634-Tev-Flag-6His) enzymes were purified from BacMam transduced CHO cells. Reactions were performed in black 384-well polystyrene low volume plates (Greiner) at a final volume of 10 μL. BOS-inhibitors (BOS-318, BOS-857 and BOS-981) were dissolved in DMSO (1 mM) and serially diluted 1 to 3 with DMSO through eleven dilutions to provide a final compound concentration range from 0.00017 to 10 μM. 0.05 μl of each concentration was transferred to the corresponding well of an assay plate, and then 5 μl of enzyme (Furin, PCSK5, PCSK6, and PCSK7) in assay buffer (100 mM HEPES pH7.5, 1 mM CaCl$_2$ and 0.005% Triton X-100) was added using a Multidrop Combi (Thermo) to the compound plates to give a final protein concentration of 0.02, 0.5, 2.5, and 1.0 nM respectively. The plates were mixed by inversion and following a 30 min preincubation of enzyme with compound at room temperature (~22°C), the substrate FAM-QRVRRAVGIDK-TAMRA (AnaSpec # 808143, 5 μl of a 1, 0.25, 0.20, and 0.5 μM solution in assay buffer for Furin, PCSK5, PCSK6, and PCSK7 respectively) was added using a Multidrop Combi. The plates were centrifuged at 500 x g for 1 minute and incubated at room temperature for two hours. Enzyme inhibition was then quantified using an Envision instrument (PerkinElmer). Data were normalized to maximal inhibition determined by 1 μM decanoyl-Arg-Val-Lys-Arg-chloromethylketone (Calbiochem #344930).

Golgi imaging assay: This assay uses an image-based platform to evaluate the intracellular activity of Furin inhibitors. Reactions were performed in black 384-well, tissue culture-treated, clear bottom plates (Greiner). Compounds dissolved in DMSO (1.0 mM) were serially three-fold diluted

to give a final compound concentration range from 0.00017 to 10 μM. Analyses were initiated by

the addition of U2OS cells simultaneously transduced with a BacMam-delivered construct

containing a Golgi-targeting sequence followed by a 12-amino acid Furin/PCSK cleavage site

from Bone Morphogenic Protein 10 (BMP10) and then GFP at the C terminus. The dibasic Furin

cleavage site sequence was flanked by glycine rich linkers (GalNAc-T2-GGGGS-

DSTARIRRNAKG-GGGGS-GFP). Briefly, frozen cells are thawed in assay media (Dulbecco's

Modified Eagles Medium Nutritional Mixture F-12 (Ham) without phenol red containing 5% FBS)

and diluted to deliver 6000 cells/well (50 μl) to the plate using a Multidrop Combi (Thermo). After

a 24-hour incubation period at 37°C, the cells are stained with Cell Mask Deep Red, fixed in

paraformaldehyde and the nuclei stained using Ho33342. The Golgi-targeted GFP forms bright

punctate clusters within the cell. In the absence of a Furin/PCSK inhibitor, the endogenous

protease cleaves GFP from its N-acetylgalactosaminyltransferase-2 Golgi tether, releasing GFP

into the Golgi lumen where fluorescence was diluted below the threshold of assay sensitivity. In

the presence of a cell permeable Furin/PCSK inhibitor, GFP fluorescence increases as intra-Golgi

protease activity was reduced. Cellular GFP intensity was determined by image-based acquisition

(Incell 2200, Perkin Elmer) at 40x magnification with 4 fields measured per well. Multi-scale top

hat segmentation was used to identify the GFP-tagged puncta and to quantitate the average

fluorescence of all puncta on a per cell basis. Cellular toxicity was determined in parallel.

<u>Furin and TMPRSS2 fluorogenic assays</u>: Recombinant Furin was purchased from BioLegend

(#719406), human recombinant TRMPSS2 from Cliniscience (ref LS-G57269-100), and the

DABCYLGlu-EDANS labelled peptides encompassing the different cleavage sites (SI Table 1)

were purchased from Genscript. Reactions were performed at room temperature in black 384-well

polystyrene low volume plates (CELLSTAR-Greiner Bio-One # 784476) at a final volume of 15

μL. The fluorescent peptides were used at 5 µM and the reactions were performed in 50 mM Tris

buffer (pH 6.5 or 7.5), 0.2% Triton X-100, 1mM CaCl$_2$ and Furin was added at a final

concentration of 0.2 to 100 nM. BOS-inhibitors (BOS-318, BOS-857 and BOS-981) were

dissolved in DMSO (1 mM) and serially diluted 1 to 2 with DMSO to provide a final compound

concentration range from 50 µM to 0.01 nM with 5% DMSO in the enzymatic assay. For

TMPRSS2, the fluorescent peptides were used at 5 µM and the reactions were performed in 50

mM Tris buffer (pH 8), 150 mM NaCl and TMPRSS2 was added at final concentrations of 50 nM.

Cleavage of the synthetic peptides was quantitated by determining the increase of EDANS (493

nM) fluorescence following release of the DABCYL quencher, which was excited at 335 nM using

a Safire 2 Tecan fluorimeter. The fluorescence was followed during 90 min, and the enzymatic

activity was deduced by measurement of the increase of fluorescence during the linear phase of

the reaction. Each reaction was performed in triplicate and the standard deviation was calculated

using Excel-ecart type function ($\sqrt{\frac{\sum (x-\bar{x})^2}{(n-1)}}$).

Plasmids

Single tagged (C-terminal V5 tag) or double tagged (N-terminal HA tag and C-terminal V5 tag)

spike-glycoprotein of SARS-CoV-2 (optimized sequence) and its mutants were cloned into the

pIRES2-EGFP vector. Site-directed mutagenesis was achieved using a Quick-Change kit

(Stratagene, CA) according to the manufacturer's instructions. The plasmids pCI-NEO-hACE2

received from DW Lambert (University of Leeds) and pIRES-NEO3-hTMPRSS2 from P Jolicoeur

(IRCM). The ΔEnv Vpr Luciferase Reporter Vector (pNL4-3.Luc.R-E-) was obtained from Dr.

Nathaniel Landau through the NIH AIDS Reagent Program whereas the pHIV-1NL4-3 ΔEnv-

NanoLuc construct was a kind gift from Dr. P Bieniasz. Plasmids encoding VSV-G, as HIV-1

Env and tat were previously described (78, 79).

Cell culture and transfection

Monolayers of HeLa, HEK293T, HEK293T17, Vero E6 and Calu-3 cells were cultured in 5% CO_2

at 37°C in Dulbecco's modified Eagle's medium (DMEM; Wisent) supplemented with 10% (v/v)

fetal bovine serum (FBS; Wisent). HEK293T-ACE2(80), a generous gift from Dr. Paul Bieniasz,

were maintained in DMEM containing 10% FBS, 1% nonessential amino acids (NEAA) and 50

μg/ml blasticidin (Invivogen). The cells were transfected with JetPrime transfection reagent

according to the manufacturer's instructions (Polyplus transfection, New York, USA). At 24h post

transfection, the culture media were changed to serum-free and cells incubated for an additional

24h. To establish the HeLa cells stably express human ACE2, transfected cells were selected using

media containing 500 μg/ml of neomycin (G418, Wisent).

For knockdown of Furin in HeLa cells, an optimized set of 4 small interfering RNAs (siRNAs;

SMARTPool) targeted against human Furin were purchased from Horizon Discoveries (Perkin

Elmer, Lafayette, LA, USA) and transfections of HeLa cells were carried out using INTERFERin

(PolyPlus) as recommended by the manufacturer.

To generate HIV particles pseudotyped with SARS-CoV-2 S, 293T17 cells (600,000 cells plated

in a 6-well vessel) were transfected with 1 μg pNL4-3 Luc.R-E- (or pHIV-1NLΔEnv-NanoLuc)

in the presence or absence of 0.3 μg pIR-2019-nCoV-S V5 plasmids using Lipofectamine-3000

(Life Technologies). In certain experiments, 293T17 cells were treated with BOS-inhibitors at 6 h

post transfection. Pseudovirions expressing the nano- or firefly-luciferase were collected at 24 h

or 48 h post transfection, respectively. Viral supernatants were clarified by centrifugation at 300 x

g, passed through a 0.45-μm pore-size polyvinylidene fluoride (PVDF; Millipore) syringe filter

(Millipore; SLGVR33RS), and aliquots frozen at −80°C. For WB analysis of purified

pseudovirions, viral supernatants were concentrated by ultracentrifugation on a 20% sucrose

cushion for 3h at 35,000 RPM; Beckman Coulter OPTIMA XE; Ti70.1 rotor). HIV particles

lacking the SARS-CoV-2 S glycoprotein served as a negative control in all experiments.

Cell viability assay using MTT

Cells, seeded in a 96-well plate, the day before, at 10,000 (HEK-293T and Vero E6) or 50,000

(Calu-3) cells, were treated with serial 10-fold dilutions of BOS-inhibitors for up to 48h. Cells

treated with vehicle alone were used as negative control. MTT was subsequently added to the

medium (final concentration: 2.5 mg/ml) and cells were further incubated for 4h at 37 ^{0}C. After

removal of the culture media, DMSO was added and absorbance read at 595 nm using a microplate

spectrophotometer. The data from two independent experiments done in triplicates was used to

calculate the CC50 by nonlinear regression using GraphPad Prism V5.0 software.

Western blots

The cells were washed with PBS and then lysed using RIPA buffer (1% Triton X-100, 150 mM

NaCl, 5 mM EDTA, and 50 mM Tris, pH 7.5) for 30 min at 4ºC. The cell lysates were collected

after centrifugation at 14,000 × g for 10 min. The proteins were separated on 7% tris-glycine or

8% tricine gels by SDS-PAGE and transferred to a PVDF membrane (Perkin Elmer). The proteins

were revealed using a V5-monoclonal antibody (V5-mAb V2660; 1:5000; Invitrogen), ACE2

antibody (rabbit monoclonal ab108252; 1:3,000; Abcam), Actin antibody (rabbit polyclonal

A2066; 1:5,000; Sigma), or HA-HRP antibody (12-013-819; 1:3,500; Roche). The antigen-

antibody complexes were visualized using appropriate HRP conjugated secondary antibodies and

enhanced chemiluminescence kit (ECL; Amersham or Bio-Rad) and normalization was reported

to β-actin. Quantification of immune-bands was performed using Image Lab software (Bio-Rad).

For analysis of SARS-CoV-2 S virions or pseudovirions, protein extracts of purified viral

particles and corresponding producing cells (Calu-3 or 293T17, respectively) were resolved on

10% tris-glycine gels and immunoblotted for spike, nucleocapsid, HIV-1 Gag p24 or actin using

anti-V5 (for pseudovirion detection; V2660)/anti-S2 (for virion detection; Sino Biologicals;

40590-T62), anti-N (Sino Biologicals; 40143-MM05), anti-p24 (MBS Hybridoma line 31-90-25)

or anti-actin (MP Biomedicals, SKU 08691001), respectively.

Glycosidase treatment

30 to 50 μg proteins were digested for 90 min at 37°C with endoglycosidase-H (Endo-H; P0702L)

or endoglycosidase-F (Endo-F; P0705S) as recommended by the manufacturer (New England

Biolabs).

Inhibitor treatment

At 24h post transfection, cells were incubated for 6h with two pan-PC inhibitors: the cell permeable

decanoyl-RVKR-chloromethylketone (cmk; 50 mM; 4026850.001; Bachem) or with the cell

surface PC-inhibitor hexa-D-arginine (D6R; 20 μM; 344931; EMD). Culture media were then

replaced with fresh ones containing the inhibitors for an additional 24h. For the selective cell-

permeable Furin-like inhibitors (BOS; Boston Pharmaceuticals), the cells were treated with the

inhibitors at the specified concentration starting at 5h pre-transfection and throughout the duration

of the experiment.

Cell-to-cell fusion assay

HeLa or HeLa TZM-bl cells were plated at 200,000 cells in 12-well plates. HeLa cells were

transiently transfected with different constructs of SARS-CoV-2 spike or NL4.3-HIV Env, or an

empty vector and 0.2 μg of CMV-Tat plasmid. HeLa TZM-bl cells were transfected with human

ACE2, TMPRSS2 or a combination of both. At 6h post-transfection, media were replaced with

fresh ones containing Furin-inhibitors, and 24h later the cells were detached with PBS-EDTA (1

μM). Different combinations of HeLa and HeLa-TZM-bl cells were placed in co-culture plate at a

ratio of 1:1 for a total of 60,000 cells/well of a 96 well place. After 18-24h the media were removed

and 50 µl of cell lysis reagent was added in each well. 20 µl of the cell lysate was used for luciferase

reading using 50 µl of Renilla luciferase reagent (Promega, Madison, WI, USA). Relative light

units (RLU) were measured using a Promega GLOMAX plate reader (Promega, Madison, WI,

USA) and values were reported as fold increase over the RLU measured in co-culture of HeLa

cells transfected an empty vector (V) with respective TZM-bl cells.

Protein immunoprecipitation from co-culture media

When indicated, a secreted form of double tagged spike-glycoprotein of SARS-CoV-2 (N-terminal

HA tag and C-terminal V5 tag) from media of co-cultured HeLa and HeLa-TZM-bl cells was

analyzed by immunoprecipitation. Namely, 0.3 ml of media were precipitated with 25 µl EZ view

Red anti-HA affinity gel (E 6779; Sigma-Aldrich) according to the manufacturers' protocol. Upon

SDS-PAGE separation and PVDF transfer, the proteins were detected using an HA-HRP antibody

(12-013-819; 1:3,500; Roche).

Microscopy

In our luciferase assay, cell co-cultures were plated on glass coverslips. After 18-24h, the cells

were incubated with 488 CellMask™ to stain the membrane and then fixed with 4% PFA for 15

min at 4ºC. The glass coverslips were mounted on glass slides using ProLong™ Gold Antifade

containing DAPI (Invitrogen). The number of syncytia were counted over 10 fields.

Immunofluorescence

Cell culture and transfection were performed on glass coverslips. Cells were washed twice with

PBS and fixed with fresh 4% paraformaldehyde for 10 min at room temperature. Following

washes, cells were either non-permeabilized or permeabilized with 0.2% Triton X-100 in PBS

containing 2% BSA for 5 min, washed, and then blocking was performed with PBS containing 2%

BSA for 1h. Cells were incubated with primary antibodies overnight at 4°C using an antibody

against V5 (mouse monoclonal R960-25; 1:1000; Invitrogen), spike (mouse monoclonal

GTX632604; 1:500; GeneTex) and ACE2 (goat polyclonal AF933; 1:500; R&D Systems).

Following wash, corresponding species-specific Alexa-Fluor (488 or 555)-tagged antibodies

(Molecular Probes) were incubated for 1h at room temperature. Coverslips were mounted on a

glass slide using ProLong Gold Reagent with DAPI (P36935, Life Technologies). Samples were

visualized using a confocal laser-scanning microscope (LSM710, Carl Zeiss) with Plan-

Apochromat 63x/1.40 Oil DIC M27 objective on ZEN software.

Pseudovirus entry

HEK293T-ACE2 or Calu-3 (10,000 cells/well plated in a 96-well dish 24 or 48h before,

respectively) were incubated with up to 200 µl filtered pseudovirions for overnight. In certain

experiments, target cells were pretreated with BOS-318 (1 µM) for 6h and/or Camostat (40 µM)

for 2h before transduction. The overnight incubation with pseudovirions was performed in the

presence of the inhibitors. Viral inoculum was removed, then fresh media were added, and the

cells cultured for up to 72h. Upon removal of spent media, 293T-ACE2 and Calu-3 cells were

gently washed twice with PBS and analyzed for firefly- or nano- luciferase activity, respectively

using Promega luciferase assay (Cat # E1501) or Nano-Glo luciferase system (Cat # N1110),

respectively.

Replication competent SARS-CoV-2 Viruses

SARS-CoV-2, which served as the viral source, was originally isolated from a COVID-19 patient

in Quebec, Canada and was designated as LSPQ1. The clinical isolate was amplified, tittered in

Vero E6 using a plaque assay as detailed below, and the integrity of the S-protein multi-basic

protein convertase site validated by sequencing. All experiments involving infectious SARS-CoV-

2 virus were performed in the designated areas of the Biosafety level 3 laboratory (IRCM)

previously approved for SARS-CoV-2 work.

Plaque assay in Vero E6 cells

Vero E6 cells (1.2 x 10^5 cells/well) were seeded in quadruplicate in 24-well tissue culture plates in

DMEM supplemented with 10% FBS two days before infection. Cells were infected with up to six

ten-fold serial dilutions (10^{-2}-10^{-6}) of viral supernatant containing SARS-CoV-2 for 1h at 37ºC

(200 µl infection volume). The plates were manually rocked every 15 min during the 1-hour period.

Subsequently, virus was removed, cells were washed and overlaying media (containing 0.6% low

melt agarose in DMEM with 10% FBS) was added and incubated undisturbed for 60-65h at 37ºC.

Post incubation, cells were fixed with 4% formaldehyde and stained with 0.25% crystal violet

(prepared in 30% methanol). High quality plaque pictures were taken using a high resolution

DLSR camera (Nikon model: D80, objective: "AF Micro-Nikkor 60mm f/2.8D"). Plaques were

counted manually and in parallel, imaged plaque plates were processed and plaques enumerated

using an automated algorithm based Matlab software. Virus titer is expressed as plaque-forming

units per ml (PFU/ml): (number of plaques x dilution factor of the virus) x 1000 / volume of virus

dilution used for infection (in µl). Multiplicity of infection (MOI) expressed as: MOI = PFU of

virus used for infection / number of cells.

Cell infections with fully replicative SARS-CoV-2

Vero E.6 and Calu-3 cells were seeded in duplicates in 12-well plates (2.3 x 10^5 cells/well) the day

before. Cells were pre-treated with various concentrations (0.1-1µM) of BOS-inhibitors and

vehicle alone (DMSO) for up to 24h. In certain experiments, Calu-3 were also pre-treated with

Camostat for 1h. Thereafter, the cells were infected with SARS-CoV-2 virus at MOI of 0.001 for

1h (Vero E6) or 0.01 for 3h (Calu-3 cells) in 350 µl of serum-free DMEM at 37ºC with occasional

manual rocking of plates. Cells plus media only were used as a control. After incubation, virus

was removed, and the cell monolayer was washed twice successively with PBS and serum-free

DMEM. New media (total 1ml) containing the concentrations of BOS-inhibitors was subsequently

added to cells. Cell-free supernatant (250 µl) was removed at 12, 24 and 48h post infection. The

drugs were replenished for 1 ml media at 24h post-infection. The virus supernatants were stored

at -80°C until further use. Viral production in the supernatant was quantified using a plaque assay

on Vero E6.1 cells as described above. In certain experiments, viral supernatants were harvested

at the end of infection and purified on a 20% sucrose cushion using ultracentrifugation as described

above. The resulting concentrated virus and corresponding infected cells were analyzed by

Western blotting as appropriate.

<u>Quantification and statistical analysis</u>: Virus titers quantified by plaque assay in triplicate were

shown as mean ± standard deviation. The results from experiments done with two biological

replicates and two technical replicates in triplicates were used to calculate the IC_{50} by nonlinear

regression using GraphPad Prism V5.0 software. The difference between the control cells (virus

with 0.001% DMSO) and the cells treated with BOS-inhibitors were evaluated by Student's t test.

The P values of 0.05 or lower were considered statistically significant (*, $p < 0.05$; **, $p < 0.01$;

***, $p < 0.001$).

DATA AVAILABILITY

734   Source data are provided with this paper. The data that support the findings of this study are

735   preserved at repositories of the Montreal Clinical Research Institute (IRCM), Montreal, QC,

736   Canada and available from the corresponding authors upon reasonable request.

ACKNOWLEDGEMENTS

This work was supported in part by CIHR Foundation grants (NGS: # 148363) and (ÉAC: # 154324), a Canada Research Chairs in Precursor Proteolysis (NGS: # 950-231335), a CIHR CHAMPS Team Grant # HAL 157986 (NGS and ÉAC), Réseau SIDA maladies infectieuses COVID-19 initiative (ÉAC and NGS), the European score project and ANR Reacting COVID-19 (ED and BC). The authors thank the Quebec public health laboratory for providing the infectious isolate LSPQ1 SARS-CoV-2. We thank Paul Bieniasz for the 293T-ACE2 cell line and the pHIV-1NL4-3ΔEnv-NanoLuc construct. The following reagents were obtained from the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: TZM-bl cells, from John C. Kappes, Xiaoyun Wu, and Tranzyme, Inc. and HIV-1 pNL4-3 ΔEnv Vpr Luciferase Reporter Vector (pNL4-3.Luc.R-E-) obtained from Nathaniel Landau. We are thankful to Dominic Filion for developing the algorithm for image-assisted plaque quantification. JJ is supported by the CIHR Postdoctoral Fellowship (HIV-435243-73284). We also thank Dr Annik Prat (IRCM) for the design of the summary model shown in Fig. 7. Finally, we would like to thank Mrs. Habiba Oueslati and Brigitte Mary for ther excellent editorial help and organization of the manuscript.

AUTHOR CONTRIBUTIONS

RE made all the original critical experiments revealing the role of the PCs in spike processing and the effect of their inhibitors. JJ performed all the cell assays with infectious SARS-CoV-2. DSR participated in the biochemical characterizations of TMPRSS2 processing of ACE2 and S1. UA performed all cell-to-cell fusion assays. AE made all the mutants used in the work. RMD generated the HeLa-ACE2 cells and prepared all the cells for *ex vivo* analyses. DNH performed all the immunocytochemical experiments. FD and ML performed experiments related to SARS-CoV-2 pseudovirions. AD and PSO performed all the Furin and TMPRSS2 *in vitro* kinetic cleavage analyses of peptides mimicking the S1/S2 and S2' sites. CM and KW provided the BOS-inhibitors

and their characterization. ED made seminal contributions to the possible role of Furin-like

enzymes in the processing of the spike-glycoprotein and actively contributed to the

conceptualization and writing of the manuscript. TNQP designed, performed, and analyzed

experiments related to viral entry and contributed to the writing of the manuscript. EAC (virology)

and NGS (biochemistry and cell biology) conceptualized the research program and provided the

intellectual contributions and funding for the whole project. All authors actively contributed to the

final version of the manuscript.

REFERENCES

1. Dobson AP, Carper ER. 1996. Infectious Diseases and Human Population History: Throughout history the establishment of disease has been a side effect of the growth of civilization. Bioscience 46:115-126.

2. Cui J, Li F, Shi ZL. 2019. Origin and evolution of pathogenic coronaviruses. Nat Rev Microbiol 17:181-192.

3. Almeida JD, Tyrrell DA. 1967. The morphology of three previously uncharacterized human respiratory viruses that grow in organ culture. J Gen Virol 1:175-8.

4. Belouzard S, Millet JK, Licitra BN, Whittaker GR. 2012. Mechanisms of coronavirus cell entry mediated by the viral spike protein. Viruses 4:1011-33.

5. Liu PP, Blet A, Smyth D, Li H. 2020. The Science Underlying COVID-19: Implications for the Cardiovascular System. Circulation 142:68-78.

6. Usul Afsar C. 2020. 2019-nCoV-SARS-CoV-2 (COVID-19) infection: Cruciality of Furin and relevance with cancer. Med Hypotheses 140:109770.

7. Braun E, Sauter D. 2019. Furin-mediated protein processing in infectious diseases and cancer. Clin Transl Immunology 8:e1073.

8. Izaguirre G. 2019. The Proteolytic Regulation of Virus Cell Entry by Furin and Other Proprotein Convertases. Viruses 11.

9. Seidah NG, Pasquato A, Andreo U. 2021. How Do Enveloped Viruses Exploit the Secretory Proprotein Convertases to Regulate Infectivity and Spread? Viruses 13:1229.

10. Coutard B, Valle C, de Lamballerie X, Canard B, Seidah NG, Decroly E. 2020. The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV of the same clade. Antiviral Res 176:104742.

11. Tang T, Bidon M, Jaimes JA, Whittaker GR, Daniel S. 2020. Coronavirus membrane

fusion mechanism offers as a potential target for antiviral development. Antiviral Res

doi:10.1016/j.antiviral.2020.104792:104792.

12. Hoffmann M, Kleine-Weber H, Pöhlmann S. 2020. A Multibasic Cleavage Site in the Spike

Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. Mol Cell 78:779-

784.e5.

13. Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, Zhang Q, Shi X, Wang Q, Zhang L, Wang X.

2020. Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2

receptor. Nature 581:215-220.

14. Lu G, Wang Q, Gao GF. 2015. Bat-to-human: spike features determining 'host jump' of

coronaviruses SARS-CoV, MERS-CoV, and beyond. Trends Microbiol 23:468-78.

15. Millet JK, Whittaker GR. 2018. Physiological and molecular triggers for SARS-CoV

membrane fusion and entry into host cells. Virology 517:3-8.

16. Ou X, Liu Y, Lei X, Li P, Mi D, Ren L, Guo L, Guo R, Chen T, Hu J, Xiang Z, Mu Z,

Chen X, Chen J, Hu K, Jin Q, Wang J, Qian Z. 2020. Characterization of spike glycoprotein

of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. Nat

Commun 11:1620.

17. Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, Li F. 2020. Cell entry mechanisms

of SARS-CoV-2. Proc Natl Acad Sci U S A 117:11727-11734.

18. Bestle D, Heindl MR, Limburg H, Van Lam van T, Pilgram O, Moulton H, Stein DA,

Hardes K, Eickmann M, Dolnik O, Rohde C, Klenk HD, Garten W, Steinmetzer T,

Böttcher-Friebertshäuser E. 2020. TMPRSS2 and furin are both essential for proteolytic

activation of SARS-CoV-2 in human airway cells. Life Sci Alliance 3.

19.   Mykytyn AZ, Breugem TI, Riesebosch S, Schipper D, van den Doel PB, Rottier RJ,

Lamers MM, Haagmans BL. 2021. SARS-CoV-2 entry into human airway organoids is

serine protease-mediated and facilitated by the multibasic cleavage site. Elife 10.

20.   Seidah NG, Prat A. 2012. The biology and therapeutic targeting of the proprotein

convertases. Nat Rev Drug Discov 11:367-383.

21.   Van de Ven WJ, Creemers JW, Roebroek AJ. 1991. Furin: the prototype mammalian

subtilisin-like proprotein-processing enzyme. Endoproteolytic cleavage at paired basic

residues of proproteins of the eukaryotic secretory pathway. Enzyme 45:257-270.

22.   Moulard M, Decroly E. 2000. Maturation of HIV envelope glycoprotein precursors by

cellular endoproteases. Biochim Biophys Acta 1469:121-132.

23.   Millet JK, Whittaker GR. 2015. Host cell proteases: Critical determinants of coronavirus

tropism and pathogenesis. Virus Res 202:120-34.

24.   Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, Schiergens

TS, Herrler G, Wu NH, Nitsche A, Muller MA, Drosten C, Pohlmann S. 2020. SARS-

CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven

Protease Inhibitor. Cell 181:271-280.e8.

25.   Xia S, Lan Q, Su S, Wang X, Xu W, Liu Z, Zhu Y, Wang Q, Lu L, Jiang S. 2020. The role

of furin cleavage site in SARS-CoV-2 spike protein-mediated membrane fusion in the

presence or absence of trypsin. Signal Transduct Target Ther 5:92.

26.   Johnson BA, Xie X, Bailey AL, Kalveram B, Lokugamage KG, Muruato A, Zou J, Zhang

X, Juelich T, Smith JK, Zhang L, Bopp N, Schindewolf C, Vu M, Vanderheiden A, Winkler

ES, Swetnam D, Plante JA, Aguilar P, Plante KS, Popov V, Lee B, Weaver SC, Suthar

MS, Routh AL, Ren P, Ku Z, An Z, Debbink K, Diamond MS, Shi PY, Freiberg AN,

Menachery VD. 2021. Loss of furin cleavage site attenuates SARS-CoV-2 pathogenesis.

Nature doi:10.1038/s41586-021-03237-4.

27.   Peacock TP, Goldhill DH, Zhou J, Baillon L, Frise R, Swann OC, Kugathasan R, Penn R,

Brown JC, Sanchez-David RY, Braga L, Williamson MK, Hassard JA, Staller E, Hanley

B, Osborn M, Giacca M, Davidson AD, Matthews DA, Barclay WS. 2021. The furin

cleavage site in the SARS-CoV-2 spike protein is required for transmission in ferrets. Nat

Microbiol doi:10.1038/s41564-021-00908-w.

28.   Johnson BA, Xie X, Bailey AL, Kalveram B, Lokugamage KG, Muruato A, Zou J, Zhang

X, Juelich T, Smith JK, Zhang L, Bopp N, Schindewolf C, Vu M, Vanderheiden A, Winkler

ES, Swetnam D, Plante JA, Aguilar P, Plante KS, Popov V, Lee B, Weaver SC, Suthar

MS, Routh AL, Ren P, Ku Z, An Z, Debbink K, Diamond MS, Shi PY, Freiberg AN,

Menachery VD. 2021. Loss of furin cleavage site attenuates SARS-CoV-2 pathogenesis.

Nature 591:293-299.

29.   Afar DE, Vivanco I, Hubert RS, Kuo J, Chen E, Saffran DC, Raitano AB, Jakobovits A.

2001. Catalytic cleavage of the androgen-regulated TMPRSS2 protease results in its

secretion by prostate and prostate cancer epithelia. Cancer Res 61:1686-92.

30.   Shrimp JH, Kales SC, Sanderson PE, Simeonov A, Shen M, Hall MD. 2020. An Enzymatic

TMPRSS2 Assay for Assessment of Clinical Candidates and Discovery of Inhibitors as

Potential Treatment of COVID-19. bioRxiv doi:10.1101/2020.06.23.167544.

31.   Kawase M, Shirato K, van der Hoek L, Taguchi F, Matsuyama S. 2012. Simultaneous

treatment of human bronchial epithelial cells with serine and cysteine protease inhibitors

prevents severe acute respiratory syndrome coronavirus entry. J Virol 86:6537-45.

32. Nimishakavi S, Raymond WW, Gruenert DC, Caughey GH. 2015. Divergent Inhibitor

Susceptibility among Airway Lumen-Accessible Tryptic Proteases. PLoS One

10:e0141169.

33. Hörnich BF, Großkopf AK, Schlagowski S, Tenbusch M, Kleine-Weber H, Neipel F,

Stahl-Hennig C, Hahn AS. 2021. SARS-CoV-2 and SARS-CoV spike-mediated cell-cell

fusion differ in the requirements for receptor expression and proteolytic activation. J Virol

doi:10.1128/jvi.00002-21.

34. Cyranoski D. 2020. Profile of a killer: the complex biology powering the coronavirus

pandemic. Nature 581:22-26.

35. Matoba Y, Aoki Y, Tanaka S, Yahagi K, Katsushima Y, Katsushima F, Sugawara K,

Matsuzaki Y, Mizuta K. 2016. HeLa-ACE2-TMPRSS2 Cells Are Useful for the Isolation

of Human Coronavirus 229E. Jpn J Infect Dis 69:452-4.

36. Thomas G. 2002. Furin at the cutting edge: from protein traffic to embryogenesis and

disease. Nature Reviews Molecular Cell Biology 3:753-766.

37. Örd M, Faustova I, Loog M. 2020. Biochemical evidence of furin specificity and potential

for phospho-regulation at Spike protein S1/S2 cleavage site in SARS-CoV2 but not in

SARS-CoV1   or   MERS-CoV.   bioRxiv

doi:10.1101/2020.06.23.166900:2020.06.23.166900.

38. Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. 2020. The proximal origin

of SARS-CoV-2. Nat Med 26:450-452.

39. Schjoldager KT, Clausen H. 2012. Site-specific protein O-glycosylation modulates

proprotein processing - deciphering specific functions of the large polypeptide GalNAc-

transferase gene family. Biochim Biophys Acta 1820:2079-2094.

40.   Benton DJ, Wrobel AG, Xu P, Roustan C, Martin SR, Rosenthal PB, Skehel JJ, Gamblin

886   SJ. 2020. Receptor binding and priming of the spike protein of SARS-CoV-2 for membrane

887   fusion. Nature 588:327-330.

888   41.   Belouzard S, Chu VC, Whittaker GR. 2009. Activation of the SARS coronavirus spike

889   protein via sequential proteolytic cleavage at two distinct sites. Proc Natl Acad Sci U S A

890   106:5871-6.

891   42.   Hallenberger S, Bosch V, Angliker H, Shaw E, Klenk HD, Garten W. 1992. Inhibition of

892   furin-mediated cleavage activation of HIV-1 glycoprotein gp160. Nature 360:358-361.

893   43.   Susan-Resiga D, Essalmani R, Hamelin J, Asselin MC, Benjannet S, Chamberland A, Day

894   R, Szumska D, Constam D, Bhattacharya S, Prat A, Seidah NG. 2011. Furin Is the Major

895   Processing Enzyme of the Cardiac-specific Growth Factor Bone Morphogenetic Protein

896   10. J Biol Chem 286:22785-22794.

897   44.   Cheng YW, Chao TL, Li CL, Chiu MF, Kao HC, Wang SH, Pang YH, Lin CH, Tsai YM,

898   Lee WH, Tao MH, Ho TC, Wu PY, Jang LT, Chen PJ, Chang SY, Yeh SH. 2020. Furin

899   Inhibitors Block SARS-CoV-2 Spike Protein Cleavage to Suppress Virus Production and

900   Cytopathic Effects. Cell Rep 33:108254.

901   45.   Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. 2020. Structure,

902   Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell 181:281-292.e6.

903   46.   Hoffmann M, Mösbauer K, Hofmann-Winkler H, Kaul A, Kleine-Weber H, Krüger N,

904   Gassen NC, Müller MA, Drosten C, Pöhlmann S. 2020. Chloroquine does not inhibit

905   infection of human lung cells with SARS-CoV-2. Nature 585:588-590.

906   47.   Vincent MJ, Bergeron E, Benjannet S, Erickson BR, Rollin PE, Ksiazek TG, Seidah NG,

907   Nichol ST. 2005. Chloroquine is a potent inhibitor of SARS coronavirus infection and

908   spread. Virol J 2:69.

909   48.   Bayati A, Kumar R, Francis V, McPherson PS. 2021. SARS-CoV-2 infects cells following

910   viral entry via clathrin-mediated endocytosis. J Biol Chem 296:100306.

911   49.   Platt EJ, Wehrly K, Kuhmann SE, Chesebro B, Kabat D. 1998. Effects of CCR5 and CD4

912   cell surface concentrations on infections by macrophagetropic isolates of human

913   immunodeficiency virus type 1. J Virol 72:2855-64.

914   50.   Plante JA, Mitchell BM, Plante KS, Debbink K, Weaver SC, Menachery VD. 2021. The

915   variant gambit: COVID-19's next move. Cell Host Microbe 29:508-515.

916   51.   Peacock TP, Penrice-Randal R, Hiscox JA, Barclay WS. 2021. SARS-CoV-2 one year on:

917   evidence for ongoing viral adaptation. J Gen Virol 102.

918   52.   Rajah MM, Hubert M, Bishop E, Saunders N, Robinot R, Grzelak L, Planas D, Dufloo J,

919   Gellenoncourt S, Bongers A, Zivaljic M, Planchais C, Guivel-Benhassine F, Porrot F,

920   Mouquet H, Chakrabarti LA, Buchrieser J, Schwartz O. 2021. SARS-CoV-2 Alpha, Beta,

921   and Delta variants display enhanced Spike-mediated syncytia formation. EMBO J

922   doi:10.15252/embj.2021108944:e108944.

923   53.   Matsuyama S, Nagata N, Shirato K, Kawase M, Takeda M, Taguchi F. 2010. Efficient

924   activation of the severe acute respiratory syndrome coronavirus spike protein by the

925   transmembrane protease TMPRSS2. J Virol 84:12658-64.

926   54.   Heurich A, Hofmann-Winkler H, Gierer S, Liepold T, Jahn O, Pöhlmann S. 2014.

927   TMPRSS2 and ADAM17 cleave ACE2 differentially and only proteolysis by TMPRSS2

928   augments entry driven by the severe acute respiratory syndrome coronavirus spike protein.

929   J Virol 88:1293-307.

930   55.   Monteil V, Kwon H, Prado P, Hagelkrüys A, Wimmer RA, Stahl M, Leopoldi A, Garreta

931   E, Hurtado Del Pozo C, Prosper F, Romero JP, Wirnsberger G, Zhang H, Slutsky AS,

932   Conder R, Montserrat N, Mirazimi A, Penninger JM. 2020. Inhibition of SARS-CoV-2

933  Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. Cell

934  181:905-913.e7.

935 56. Hoffmann M, Mösbauer K, Hofmann-Winkler H, Kaul A, Kleine-Weber H, Krüger N,

936  Gassen NC, Müller MA, Drosten C, Pöhlmann S. 2020. Chloroquine does not inhibit

937  infection of human lung cells with SARS-CoV-2. Nature doi:10.1038/s41586-020-2575-3.

938 57. Bellamine A, Pham TNQ, Jain J, Wilson J, Sahin K, Dallaire F, Seidah NG, Durkee S,

939  Radošević K, Cohen É A. 2021. L-Carnitine Tartrate Downregulates the ACE2 Receptor

940  and Limits SARS-CoV-2 Infection. Nutrients 13.

941 58. Padmanabhan P, Desikan R, Dixit NM. 2020. Targeting TMPRSS2 and Cathepsin B/L

942  together may be synergistic against SARS-CoV-2 infection. PLoS Comput Biol

943  16:e1008461.

944 59. Papa G, Mallery DL, Albecka A, Welch LG, Cattin-Ortolá J, Luptak J, Paul D, McMahon

945  HT, Goodfellow IG, Carter A, Munro S, James LC. 2021. Furin cleavage of SARS-CoV-

946  2 Spike promotes but is not essential for infection and cell-cell fusion. PLoS Pathog

947  17:e1009246.

948 60. Altulea D, Maassen S, Baranov MV, van den Bogaart G. 2021. What makes

949  (hydroxy)chloroquine ineffective against COVID-19: insights from cell biology. J Mol

950  Cell Biol doi:10.1093/jmcb/mjab016.

951 61. Self WH, Semler MW, Leither LM, Casey JD, Angus DC, Brower RG, Chang SY, Collins

952  SP, Eppensteiner JC, Filbin MR, Files DC, Gibbs KW, Ginde AA, Gong MN, Harrell FE,

953  Jr, Hayden DL, Hough CL, Johnson NJ, Khan A, Lindsell CJ, Matthay MA, Moss M, Park

954  PK, Rice TW, Robinson BRH, Schoenfeld DA, Shapiro NI, Steingrub JS, Ulysse CA,

955  Weissman A, Yealy DM, Thompson BT, Brown SM, National Heart L, Network BIPCT.

2020. Effect of Hydroxychloroquine on Clinical Status at 14 Days in Hospitalized Patients

With COVID-19: A Randomized Clinical Trial. JAMA 324:2165-2176.

62.   Rosenke K, Jarvis MA, Feldmann F, Schwarz B, Okumura A, Lovaglio J, Saturday G,

Hanley PW, Meade-White K, Williamson BN, Hansen F, Perez-Perez L, Leventhal S,

Tang-Huau TL, Callison J, Haddock E, Stromberg KA, Scott D, Sewell G, Bosio CM,

Hawman D, de Wit E, Feldmann H. 2020. Hydroxychloroquine prophylaxis and treatment

is ineffective in macaque and hamster SARS-CoV-2 disease models. JCI Insight 5.

63.   Glowacka I, Bertram S, Müller MA, Allen P, Soilleux E, Pfefferle S, Steffen I, Tsegaye

TS, He Y, Gnirss K, Niemeyer D, Schneider H, Drosten C, Pöhlmann S. 2011. Evidence

that TMPRSS2 activates the severe acute respiratory syndrome coronavirus spike protein

for membrane fusion and reduces viral control by the humoral immune response. J Virol

85:4122-34.

64.   Raghuvamsi PV, Tulsian NK, Samsudin F, Qian X, Purushotorman K, Yue G, Kozma MM,

Hwa WY, Lescar J, Bond PJ, MacAry PA, Anand GS. 2021. SARS-CoV-2 S protein:ACE2

interaction reveals novel allosteric targets. Elife 10.

65.   Zhang H, Wada J, Hida K, Tsuchiyama Y, Hiragushi K, Shikata K, Wang H, Lin S, Kanwar

YS, Makino H. 2001. Collectrin, a collecting duct-specific transmembrane glycoprotein, is

a novel homolog of ACE2 and is developmentally regulated in embryonic kidneys. J Biol

Chem 276:17132-9.

66.   Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. 2020. Structural basis for the recognition

of SARS-CoV-2 by full-length human ACE2. Science 367:1444-1448.

67.   Yamamoto M, Kiso M, Sakai-Tagawa Y, Iwatsuki-Horimoto K, Imai M, Takeda M,

Kinoshita N, Ohmagari N, Gohda J, Semba K, Matsuda Z, Kawaguchi Y, Kawaoka Y,

Inoue JI. 2020. The Anticoagulant Nafamostat Potently Inhibits SARS-CoV-2 S Protein-

Mediated Fusion in a Cell Fusion Assay System and Viral Infection In Vitro in a Cell-

Type-Dependent Manner. Viruses 12.

68. Koch J, Uckeley ZM, Doldan P, Stanifer M, Boulant S, Lozach PY. 2021. TMPRSS2

expression dictates the entry route used by SARS-CoV-2 to infect host cells. EMBO J

40:e107821.

69. Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, Niemeyer D,

Jones TC, Vollmar P, Rothe C, Hoelscher M, Bleicker T, Brünink S, Schneider J, Ehmann

R, Zwirglmaier K, Drosten C, Wendtner C. 2020. Virological assessment of hospitalized

patients with COVID-2019. Nature 581:465-469.

70. Zhou L, Niu Z, Jiang X, Zhang Z, Zheng Y, Wang Z, Zhu Y, Gao L, Huang H, Wang X,

Sun Q. 2020. The SARS-CoV-2 targets by the pscRNA profiling of ACE2, TMPRSS2 and

Furin proteases. iScience doi:10.1016/j.isci.2020.101744:101744.

71. Mayer G, Boileau G, Bendayan M. 2004. Sorting of furin in polarized epithelial and

endothelial cells: expression beyond the Golgi apparatus. J Histochem Cytochem 52:567-

580.

72. Antalis TM, Bugge TH, Wu Q. 2011. Membrane-anchored serine proteases in health and

disease. Prog Mol Biol Transl Sci 99:1-50.

73. Chu H, Chan JF, Yuen TT, Shuai H, Yuan S, Wang Y, Hu B, Yip CC, Tsang JO, Huang

X, Chai Y, Yang D, Hou Y, Chik KK, Zhang X, Fung AY, Tsoi HW, Cai JP, Chan WM,

Ip JD, Chu AW, Zhou J, Lung DC, Kok KH, To KK, Tsang OT, Chan KH, Yuen KY.

2020. Comparative tropism, replication kinetics, and cell damage profiling of SARS-CoV-

2 and SARS-CoV with implications for clinical manifestations, transmissibility, and

laboratory studies of COVID-19: an observational study. Lancet Microbe 1:e14-e23.

74. Gunst JD, Staerke NB, Pahus MH, Kristensen LH, Bodilsen J, Lohse N, Dalgaard LS,

Brønnum D, Fröbert O, Hønge B, Johansen IS, Monrad I, Erikstrup C, Rosendal R, Vilstrup

E, Mariager T, Bove DG, Offersen R, Shakar S, Cajander S, Jørgensen NP, Sritharan SS,

Breining P, Jespersen S, Mortensen KL, Jensen ML, Kolte L, Frattari GS, Larsen CS,

Storgaard M, Nielsen LP, Tolstrup M, Sædder EA, Østergaard LJ, Ngo HTT, Jensen MH,

Højen JF, Kjolby M, Søgaard OS. 2021. Efficacy of the TMPRSS2 inhibitor camostat

mesilate in patients hospitalized with Covid-19-a double-blind randomized controlled trial.

EClinicalMedicine 35:100849.

75. Shapira T, Monreal IA, Dion SP, Jager M, Désilets A, Olmstead AD, Vandal T, Buchholz

DW, Imbiakha B, Gao G, Chin A, Rees WD, Steiner T, Nabi IR, Marsault E, Sahler J,

August A, Van de Walle G, Whittaker GR, Boudreault PL, Aguilar HC, Leduc R, Jean F.

2021. A novel highly potent inhibitor of TMPRSS2-like proteases blocks SARS-CoV-2

variants of concern and is broadly protective against infection and mortality in mice.

bioRxiv doi:10.1101/2021.05.03.442520.

76. Ou T, Mou H, Zhang L, Ojha A, Choe H, Farzan M. 2021. Hydroxychloroquine-mediated

inhibition of SARS-CoV-2 entry is attenuated by TMPRSS2. PLoS Pathog 17:e1009212.

77. Morens DM, Fauci AS. 2020. Emerging Pandemic Diseases: How We Got to COVID-19.

Cell 183:837.

78. Lodge R, Lalonde JP, Lemay G, Cohen EA. 1997. The membrane-proximal

intracytoplasmic tyrosine residue of HIV-1 envelope glycoprotein is critical for basolateral

targeting of viral budding in MDCK cells. EMBO J 16:695-705.

79. Forget J, Yao XJ, Mercier J, Cohen EA. 1998. Human immunodeficiency virus type 1 vpr

protein transactivation function: mechanism and identification of domains involved. J Mol

Biol 284:915-23.

80.   Schmidt F, Weisblum Y, Muecksch F, Hoffmann HH, Michailidis E, Lorenzi JCC,

1028   Mendoza P, Rutkowska M, Bednarski E, Gaebler C, Agudelo M, Cho A, Wang Z,

1029   Gazumyan A, Cipolla M, Caskey M, Robbiani DF, Nussenzweig MC, Rice CM,

1030   Hatziioannou T, Bieniasz PD. 2020. Measuring SARS-CoV-2 neutralizing antibody

1031   activity using pseudotyped and chimeric viruses. J Exp Med 217.

1032

1033

 FIGURE LEGENDS

Figure 1: **Processing of S-peptides by Furin and TMPRSS2. (A)** Schematic representation of

the primary structure of preproS and its domains and the predicted Furin-like S1/S2 site generating

the S1- and S2-subunits, as well as the S2' site preceding the fusion peptide (FP). The signal

peptide (SP), N-terminal domain (NTD), receptor binding domain (RBD) to ACE2, the two heptad

repeats HR1 and HR2, the transmembrane domain (TM), the cytosolic tail (CT) and the C-terminal

V5-tag are indicated. **(B)** *In vitro* Furin activity against peptides mimicking the S1/S2 (and its

mutants) and S2' cleavage site sequence of the spike protein from SARS-CoV-2 and SARS-CoV-

1, as described in SI-Table 1. Each substrate was tested at a final protease concentration of 2 and

100 nM. **(C)** *In vitro* TMPRSS2 activity (at 50 nM) against peptides mimicking the S1/S2 and S2'

cleavage site sequence of the spike protein from SARS-CoV-2 (and its mutants) described in SI-

Table 1.

Figure 2: **Processing of spike-glycoprotein in HeLa cells. (A)** Western blot analyses of the

processing of WT proS into V5-tagged S2 and S2' by the proprotein convertases Furin, PC5A,

PACE4 and PC7 following co-transfection of their cDNAs in HeLa cells. The migration positions

of immature proS$_{im}$, S2 and S2' as well as the actin loading control are emphasized. V = empty

pIRES-EGFP-V5 vector. **(B)** Western blot analyses of HeLa cells following co-transfection with

cDNAs coding for either WT-S-protein or its double Ala-mutant [R685A + R682A] (μS1/S2) in the

absence or presence of Furin cDNA at a ratio S:protease = 1:2. *Inconsistently observed oligomeric

forms of proS. **(C)** Identification of S2' cleavage site by MS/MS. WT-spike-glycoprotein was

immunoprecipitated from HeLa cells using V5 agarose beads then resolved by SDS

electrophoresis SDS/PAGE and subjected to silver staining (left panel); the positions of the slices

are indicated (1 to 10). The MS/MS analysis of peptides generated by a Lys-specific protease

$(K_{814}\downarrow)$ are indicated; the data represent the ratio of SFIEDLLFNK825 to $\underline{R}_{815}$SFIEDLLFN$\underline{K}_{825}$

(right panel). (**D**) Western blot analyses of HeLa cells co-transfected with V5-tagged spike protein,

WT (**S**) or its Furin-optimized S2' (KRRKR$_{815}\downarrow$SF) mutant (μS2'), and empty vector (**V**) or Furin.

(**A, B**) The estimated % cleavages into S1/S2 and S2' are shown and were calculated as the ratio

of the V5-immunoreactivity of the cleaved form to the sum of all forms. The data are representative

of at least three independent experiments.

Figure 3: **Processing of spike-glycoprotein at S2' is enhanced in presence of ACE2.** Western blot

showing the impact of ACE2 on the processing of WT and μS1/S2 spike-glycoproteins by Furin.

HeLa cells expressing empty vector (**V**), WT proS (**A**) or its μS1/S2 mutant (**B**) without or with

Furin, ACE2 or both were analysed by Western blotting using anti-V5 antibody. The ratio of cDNAs

used was S:ACE2:Furin = 1:1:1. The data are representative of at least three independent

experiments.

Figure 4: **Inhibition of PCs by BOS compounds.** (**A**) Chemical motif of BOS-inhibitors and

representative structure of BOS-318. (**B**) *In vitro* BOS-inhibition of the cleavage of the fluorogenic

dibasic substrate FAM-QRVRRAVGIDK-TAMRA by each of the proprotein convertases Furin,

PC5 (PCSK5), PACE4 (PCSK6) and PC7 (PCSK7). All experiments were performed in 10

different wells and the average pIC$_{50}$ (in nM) was calculated. Shown for comparison is the

inhibitory pIC$_{50}$ of the Furin-like inhibitor RVKR-cmk performed >100 times. (**C**) *In vitro*

inhibition of Furin by the BOS compounds. Furin (2 nM) was incubated with increasing

concentration of BOS-inhibitors, and its enzymatic activity against the synthetic peptides

DABSYL/Glu-TNSP**RRAR**$\downarrow$SVAS-EDANS (5 μM) was measured at pH 7.5 (n=3). (**D**) Golgi

assay: table representing the effects of BOS-inhibitors on U2OS cells expressing each of Furin,

PC5A, PACE4 and PC7 simultaneously transduced with a BacMam-delivered construct

containing a Golgi-targeting sequence followed by a 12-amino acid Furin/PCSK cleavage site from Bone Morphogenic Protein 10 (BMP10) and GFP at the C terminus (GalNAc-T2-GGGGS-DSTARIRR↓NAKG-GGGGS-GFP). Dibasic cleavage releases NAKG-GGGGS-GFP thereby reducing the Golgi-associated fluorescence estimated by imaging. (**E**) Furin-inhibitors (BOS) abrogate endogenous processing of the spike-glycoprotein. Hela cells were transiently transfected with a cDNA encoding an empty vector (V) or with one expressing the V5-tagged spike (S) glycoprotein (spike-V5). At 5h pre-transfection, cells were treated with vehicle DMSO (NT, duplicate) or with the Furin-inhibitors at indicated concentrations, or RVKR-cmk at 50 μM. At 24h post-transfection media were replaced with fresh ones lacking (NT) or containing the inhibitors for an additional 24h. Cell extracts were analyzed by Western blotting using a mAb-V5. All data are representative of at least three independent experiments.

Figure 5: **Furin-like inhibitors and Camostat treatment decrease SARS-CoV-2 infection in Calu-3 Cells.** (**A**) Replication kinetics was studied at 12, 24 and 48h post-infection by plaque assay to determine PFUs of SARS-CoV-2 virus in the supernatant of infected Calu-3 cells treated or not with 1μM BOS-318, BOS-857 and BOS-981. A line graph represents results of the triplicate plaque assay results (mean ± SD). (**B**) The virus titers (PFU per milliliter) released in the supernatant (24h post-infection) of infected Calu-3 cells treated with indicated concentrations of BOS-318 were determined by plaque assay (mean ± SD of triplicates, *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$) (left panel). The selectivity index (SI) of BOS-318 in Calu-3 cells as shown in top right panel was determined by CC_{50}/IC_{50}. The left y axis indicates the inhibition of virus titer (percent) relative to that of the untreated control group (red). The right y axis indicates the cell viability (percent) relative to that of the untreated control group (green). The CC_{50} (50% cytotoxic concentration), IC_{50} (half maximal inhibitory concentration), and SI (selectivity index) values for

each inhibitor are as shown. Representative plaque images of infected Calu-3 cells treated with

indicated doses of BOS-inhibitors are shown in the bottom right panel. (**C**) Immunoblots for the

infected Calu-3 cells (right panel) and viral particles secreted in the supernatant (left panel) with

and without treatment with BOS-inhibitors indicate reduced viral protein levels. Immunoblots

were probed for the full-length (proSm) and cleaved (S2) fragments of viral S protein and

nucleocapsid (N) protein as indicated; β-Actin was included as the loading control for the cells.

(**D**) The virus titers (PFU per milliliter) released in the supernatant (24h post-infection) of infected

Calu-3 cells treated with BOS-318 and/or Camostat (Camo) were determined by plaque assay

(mean ± SD of duplicates, *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$) (top panel). Representative

plaque images of infected Calu-3 cells are shown in the bottom panel. Color plaques differentiate

the lawn (one color gray per well) from individual plaques (independent colors).

Figure 6: **Processing of SARS-CoV-2 S by Furin-like convertases and TMPRSS2 is critical**
for viral entry in human lung epithelial cells but not in model HEK293 cells stably expressing
ACE2. (**A**) Furin cleavage of proS at the S1/S2 site is required for SARS-CoV-2 pseudoviral entry

in Calu-3 cells but not HEK293T-ACE2 cells. Cells were inoculated with nanoluciferase-

expressing HIV particles pseudotyped with SARS-CoV-2: wild-type spike (WT), double Ala-

mutant spike (μS1/S2) or Furin-optimized spike (μS2'). Inhibition of proS processing at S1/S2 by

a novel Furin-like inhibitor (BOS-318) during pseudovirion packaging prevents viral entry in

Calu-3 cells but not in HEK293T-ACE2 cells. (**B**) Western blot analyses show inhibition by BOS-

318 of proS processing at S1/S2 site. Purified pseudovirions and cellular extracts of producing

HEK293-T17 cells treated or not with BOS-318 inhibitor were separated on SDS-PAGE gel and

analyzed for HIV-1 p24 and V5-tagged S-protein (proSm or cleaved, S2) as indicated. (**C**) Pre-

treatment of Calu-3 cells with (B) 1 μM BOS-318, (C) 100 μM Camostat or both (B+C) markedly

reduces viral entry. In Panels A and C, Calu-3 cells were transduced with nanoluciferase-expressing HIV particles pseudotyped with SARS-CoV-2 S WT, µS1/S2 or µS2' for 72h and analyzed for nano-luciferase expression. Viral entry was expressed as fold increase over that given by bald particles (pseudovirions made in the absence of S). In panels **A** and **C** each dot represents a different experiment with median luciferase activity calculated from three biological replicates. Two to four experiments were performed for each cell type. Error bars indicate standard deviation (SD) from the mean.

Figure 7: **Spike-induced cell-to-cell fusion relies on Furin cleavage at S1/S2.** (**A**) Cell-to-cell fusion between donor cells (HeLa) expressing the fusogenic SARS-CoV-2 spike protein along with the HIV trans-activator Tat, and acceptor cells (TZM-bl) that express ACE2. Upon fusion, Tat is transferred from donor to acceptor cells, thereby inducing luciferase expression. (**B**) Cell-to-cell fusion was evaluated using confocal microscopy. A representative immunocytochemistry of Hela cells transfected with a vector expressing SARS-CoV-2 spike co-cultured with TZM-bl cells for 18h. The number of syncytia (multiple nuclei) was examined using CellMask™ to probe for the plasma membrane and Dapi to stain the nuclei. (**C**) Donor cells were transfected with vectors expressing either no protein (empty vector, V), µS1/S2, or WT-spike (S) in the absence (NT) or presence of vehicle (DMSO) or with the Furin-inhibitors BOS-318, BOS-981, BOS-857 (300 nM) or RVKR (10 µM). Acceptor cells were transfected with a vector expressing ACE2. After 48h, donor and acceptor cells were co-cultured for 18h. Relative luminescence units (RLU) were normalized to the V value arbitrarily set to 1. Data are presented as mean values ± SD (n=3), One-Way ANOVA, Dunn-Sidàk multiple comparison test. (**D**) Donor HeLa cells expressing WT-S or its indicated mutants and variants were co-cultured with acceptor TZM-bl cells expressing ACE2. The extent of fusion is represented as a ratio between the RLU measured for each condition

and that of donor cells expressing empty vector. The bar graph represents the average of 3

experiments performed in triplicates. Data are presented as mean values ± SEM (n=3), One-way

Anova Bonferroni multiple comparison test. Two-Way ANOVA, Dunn-Sidàk multiple

comparison test.

Figure 8: **Spike-glycoprotein processing by Furin and TMPRSS2 in a co-culture system and**

their role in cell-to-cell fusion. Donor HeLa cells expressing empty vector (V), WT-S-HA or

µS1/S2-HA were co-cultured with acceptor TZM-bl cells expressing V, ACE2, or ACE2 +

TMPRSS2 (TMP). From the same experiment, cell-to-cell fusion (**A**) was assessed in parallel by

Western blotting of spike-glycoproteins in cells and media using an anti-V5 mAb (**B**). Secreted

forms of spike protein (S1) in the media were detected with anti HA-HRP upon

immunoprecipitation with anti-HA agarose. The bar graph shows one representative fusion assay

done un duplicate. The corresponding Western-blot is representative of three independent

experiments. Values of S2 and S2' relative to β-actin are shown (nd = too low or not detected).

Figure 9: **Exogenous TMPRSS2-generated shedding of ACE2 differentially regulates S-induced**

fusion at the plasma membrane of WT-S *versus* µAS1/S2. Donor HeLa cells expressing double

tagged (N-terminal HA-tag; C-terminal V5-tag) spike-glycoprotein WT (S) or its S1/S2 mutant

(µAS1/S2) were co-cultured with acceptor TZM-bl cells expressing (empty vector, V), ACE2, or

ACE2 + TMPRSS2 (TMP) and treated with DMSO (vehicle control) or Camostat (120 µM).

Within the same experiment, cell-to-cell fusion (**A**) was assessed in parallel with spike processing

in cells and media by Western-blot (**B, C**). (**A**) The extent of fusion is represented as a ratio

between the RLU measured for each condition and that of donor cells expressing V. The bar graph

represents the average of 2 experiments performed in triplicates. (**B, C**). Western blot analyses of

media and cell extracts of the co-cultured cells with donor cells overexpressing double tagged (N-

terminal HA-tag; C-terminal V5-tag) spike-glycoprotein, WT (S) (**B**) or μAS1/S2 (**C**). The arrow

points to a putative degradation product of S2', that is absent in presence of Camostat. Media were

subjected to immunoprecipitation with anti-HA agarose for the secreted forms of spike protein (S1,

S1$_L$) followed by Western blotting with anti HA-HRP. In the cell extracts Spike-glycoproteins and

ACE2 were immunoblotted with anti-V5 mAb and a polyclonal ACE2 antibody, respectively. The

data are representative of three independent experiments.

Figure 10: **The C-terminal collectrin-like domain of ACE2 may be critical for the regulation**
of cell-to-cell fusion of spike-glycoprotein when exogenous TMPRSS2 is present. (**A**)
Schematic representation of the primary structure of human ACE2 with emphasis on the C-
terminal collectrin-like domain (aa 616-768, light gray), TMPRSS2 cleavage region (aa 697-716,
black) and the polybasic amino-acid segments in which K/R were mutated to A (C0 and C4)
(amino acids underlined and in bold). Also shown are the peptidase domain (aa 19-615, white)
containing the regions involved in the interaction with the spike SARS-CoV protein (hatched) and
transmembrane domain (TM). (**B**) HeLa cells were co-transfected with ACE2, WT (ACE2) or its
mutant ACE2 (C0+C4), and TMPRSS2, WT (TMPRSS2) or its S441A active-site mutant
(μTMPRSS2), or empty vector (V). Media and cell extracts were analyzed by western blotting for
shed ACE2 (sACE2) and ACE2, respectively. The migration positions of the ~95 kDa and ~80
kDa sACE2 are emphasized. (**C**) Donor HeLa cells expressing WT-S-HA or μAS1/S2-HA were
co-cultured with acceptor TZM-bl cells expressing ACE2, WT (ACE2) or its mutant ACE2
(C0+C4) in presence or absence of TMPRSS2. From the same experiment, cell-to-cell fusion was
assessed (**C**), in parallel with WB analyses of cells and media (**D**). The extent of fusion is
represented as a ratio between the RLU measured for each condition and that of donor cells
expressing an empty vector. The bar graph represents the average of 3 experiments performed in
triplicates. Data are presented as mean values ± SD (n=3), One-way Anova Turkey's multiple

comparison test. (**D**) Co-culture media were subjected to immunoprecipitation with anti-HA

agarose for the secreted forms of spike protein (S1) followed by western blot with anti HA-HRP.

Spike-glycoproteins in the cell extracts were immunoblotted with anti-V5 mAb. The Western blot

data are representative of three independent experiments.

Figure 11: **Proposed model for the processing of S-protein and its blockade by Furin and**

TMPRSS2 inhibitors. <u>Boxed left panel</u>: schematic representation of the S-glycoprotein domains

of SARS-CoV-2, including the N-terminal (NTD) and C-terminal (CTD) domains of S1, the Furin-

S1/S2 and the Furin/TMPRSS2-S2' processing sites as well as the fusogenic α-helix that follows

S2'. Binding of the RBD domain of S1 to the membrane associated ACE2 in target cells, and the

cell surface expression of TMPRSS2 and Furin are also schematized. <u>Right panels</u>: (**1**) BOS-

inhibitors (or μS1/S2 mutant) completely prevent cell-to-cell fusion of donor Hela cells expressing

S-glycoprotein with acceptor HeLa-ACE2 cells, which lack endogenous TMPRSS2. In this

context, this reveals that Furin is a major processing enzyme cleaving at S1/S2 and generating S2'.

(**2**) In acceptor HeLa cells expressing TMPRSS2 (+), maximal prevention of cell-to-cell fusion

can be achieved by a combination of Furin (BOS, phenocopying the μS1/S2 or μAS1/S2 mutants)

and TMPRSS2 (Camostat) inhibitors blocks S2' production, ACE2-shedding (sACE2) and the

separation of sACE2-S1$_L$ complex from S2. (**3**) Optimal blockade of SARS-CoV-2 infection of

Calu-3 cells, which express endogenously both Furin and TMPRSS2, is also achieved by a

combination of Furin (BOS) and TMPRSS2 (Camostat) inhibitors.

SUPPORTING INFORMATION (SI): TABLE 1; FIGURES 1-7

Table 1: **Sequences of the different peptides mimicking the Cov spike cleavage sites that have been tested in the enzymatic assay.** The arrow indicates the expected cleavage site.

SI-Figure 1: **Importance of Furin in the processing of the Spike-glycoprotein.** (**A**) HeLa cells were first transfected with control non-targeting siRNA (siCTL) or siRNA Furin (siFur) at final concentrations of 20 nM, or mock transfected (N) and 24h later, transfected with empty vector (V) or with that coding for a V5-tagged spike-glycoprotein for an additional 48h. Following lysis, proteins were resolved on SDS-PAGE followed by WB with anti-V5 or anti-Furin antibodies. (**B**) HeLa cells transfected with empty vector (V), V5-tagged wild type spike-protein (WT) or its S1/S2 site mutant (μS1/S2) were treated with Endo-F and Endo-H or mock treated (NT) and analyzed as described in panel A. (**C, D**) HeLa cells transfected with V5-tagged wild type spike-protein (WT) or S1/S2 single mutants (**C**) or S2' single or double mutants (**D**) in the absence (V) or presence of overexpressed Furin were lysed and analyzed by WB.

SI-Figure 2: **Immunocytochemistry of the co-localization of ACE2 and S-protein or μS1/S2-S in HeLa cells.** Immunofluorescence of S-protein (green), WT (S) or μS1/S2, and ACE2 (red) were revealed using the spike S2-antibody GTX632604 in non-permeabilized (NP) conditions or anti-V5 in permeabilized (P) conditions, and ACE2 antibody AF933. The confocal co-localizations are shown in the merged Figures. Scale bar = 10 μm.

SI-Figure 3: **Furin-like inhibitors strongly reduce SARS-CoV-2 infection in Calu-3 cells.** Calu-3 cells were treated with indicated concentrations of (**A**) BOS-857 and (**B**) BOS-981 and infected with SARS-CoV-2 for 24h. Virus titers in the supernatant were determined by plaque assay on VeroE6 cells (mean plaque forming units [PFU] per ml) ± SD of triplicates, $^*p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$). The selectivity index (SI) of (**A**) BOS-857, and (**B**) BOS-981 in

Calu-3 cells as shown in top right panel was determined by CC_{50}/IC_{50}. The left y axis indicates

the inhibition of virus titer (percent) relative to that of the untreated control group (red). The

right y axis indicates the cell viability (percent) relative to that of the untreated control group

(green). Representative plaque images of infected Calu-3 cells treated with indicated doses of

BOS-inhibitors are shown in the bottom right panel. Color plaques differentiate the lawn (one

color gray per well) from individual plaques (independent colors).

SI-Figure 4: **Furin-like inhibitors modestly reduce virus production in SARS-CoV-2-infected**

Vero E6 cells in a concentration-dependent manner. (A) Vero E6 cells treated or not with 1µM

BOS-318, BOS-857 or BOS-981 were infected with SARS-CoV-2 for up to 45h. Virus titers in

the supernatant obtained at 12, 24 and 48 h post infection were determined by plaque assay on

Vero E6. A line graph represents results of the triplicate plaque assay (mean PFU/ml ± SD). **(B,**

C, and D) Virus released in the supernatant (48 hr post infection) of infected Vero E6 cells treated

with indicated concentrations of (B) BOS-318, (C) BOS-857, or (D) BOS-981 were determined

by plaque assay (mean ± SD of triplicates, $*p < 0.05$; $**p < 0.01$; $***p < 0.001$).

SI-Figure 5: **Combination of BOS-981 and Camostat reduces SARS-CoV-2 replication.** Calu-

3 cells were treated with BOS-981 and/or Camostat (Camo) and infected with SARS-CoV-2 for

24h. Virus titers in the supernatant were determined by plaque assay on VeroE6 (mean PFU/ml ±

SD of duplicates, $*p < 0.05$. Representative plaque images of infected Calu-3 cells are shown in the

bottom panel. Color plaques differentiate the lawn (one color gray per well) from individual

plaques (independent colors).

SI-Figure 6: **Cell-to-cell fusion assay: correlation between syncytia formation and luciferase**

activity. (**A**) Cell-to-cell fusion between donor cells (HeLa) and acceptor cells (TZM-bl) was

evaluated using confocal microscopy. Hela cells transfected with: (**a**) an empty vector (V), or

expressing (**b**) HIV-gp160 and Tat, (**c**) SARS-CoV-2 spike, or (**d**) μS1/S2 were co-cultured with

TZM-bl cells for 18h and the number of syncytia was examined using CellMask™ to probe for

the plasma membrane and Dapi to stain the nuclei. (**B**) Donor cells were transfected with vectors

expressing either no protein (V), Tat, WT-spike (S), Tat and WT-spike (Tat + S) or Tat and HIV-

gp160 (Tat + gp160). Acceptor cells were transfected with a vector expressing no protein (V), with

ACE2 or directly with Tat as a positive control (hatched bar). After 48h, cells were co-cultured for

18h. Luminescence was normalized to the V value arbitrarily set to 1. Data are presented as mean

values ± SD (n=3) and a representative experiment is shown. (**C**) Donor cells were transfected

with increasing amount of plasmid expressing WT-spike and acceptor cells were transfected with

a vector expressing ACE2. After 48h, cells were co-cultured for 18h, and prepared for

luminescence or microscopy. Correlation between the number of syncytia counted by microscopy

(n=10 per condition) and the luciferase activity was determined, and the calculated correlation

coefficient is R^2=0.87.

SI-Figure 7: **Secretion of S1.** HeLa cells were transiently co-expressed with double-tagged spike

protein (N-terminal HA-tag; C-terminal V5-tag), WT (S) or its mutants, μS1/S2 or μAS1/S2, and

ACE2 alone or in combination with TMPRSS2, WT (TMPRSS2) or its S441A active-mutant

(μTMPRSS2), at a ratio S:ACE2:TMPRSS2 = 1:0.5:0.5. Immunoblot of the 24h conditioned

media was first probed for secreted S1, S1' and $S1_L$ (HA-HRP antibody), stripped and next probed

for shed ACE2 (sACE2).

Figure 1

Figure 1: **Processing of S-peptides by Furin and TMPRSS2.** (**A**) Schematic representation of the primary structure of preproS

and its domains and the predicted Furin-like S1/S2 site generating the S1- and S2-subunits, as well as the S2' site preceding the

fusion peptide (FP). The signal peptide (SP), N-terminal domain (NTD), receptor binding domain (RBD) to ACE2, the two heptad

repeats HR1 and HR2, the transmembrane domain (TM), the cytosolic tail (CT) and the C-terminal V5-tag are indicated. (**B**) *In*

vitro Furin activity against peptides mimicking the S1/S2 (and its mutants) and S2' cleavage site sequence of the spike protein

from SARS-CoV-2 and SARS-CoV-1, as described in SI-Table 1. Each substrate was tested at a final protease concentration of 2

and 100 nM. (**C**) *In vitro* TMPRSS2 activity (at 50 nM) against peptides mimicking the S1/S2 and S2' cleavage site sequence of

the spike protein from SARS-CoV-2 (and its mutants) described in SI-Table 1.

Figure 2

Figure 2: **Processing of spike-glycoprotein in HeLa cells.** (**A**) Western blot analyses of the processing of WT proS into V5-tagged S2 and S2' by the proprotein convertases Furin, PC5A, PACE4 and PC7 following co-transfection of their cDNAs in HeLa cells. The migration positions of immature proSim, S2 and S2' as well as the β-Actin loading control are emphasized. V = empty pIRES-EGFP-V5 vector. (**B**) Western blot analyses of HeLa cells following co-transfection with cDNAs coding for either WT-S-protein or its double Ala-mutant [R685A + R682A] (μS1/S2) in the absence or presence of Furin cDNA at a ratio S:protease = 1:2. *Inconsistently observed oligomeric forms of proS. (**C**) Identification of S2' cleavage site by MS/MS. WT-spike-glycoprotein was immunoprecipitated from HeLa cells using V5 agarose beads then resolved by SDS electrophoresis SDS/PAGE and subjected to silver staining (left panel); the positions of the slices are indicated (1 to 10). The MS/MS analysis of peptides generated by a Lys-specific protease (K814↓) are indicated; the data represent the ratio of $SFIEDLLFNK_{825}$ to $R_{815}SFIEDLLFNK_{825}$ (right panel). (**D**) Western blot analyses of HeLa cells co-transfected with V5-tagged spike protein, WT (S) or its Furin-optimized S2' ($KRRKR_{815}$↓SF) mutant (μS2'), and empty vector (V) or Furin. (**A, B**) The estimated % cleavages into S1/S2 and S2' are shown and were calculated as the ratio of the V5-immunoreactivity of the cleaved form to the sum of all forms. The data are representative of at least three independent experiments.

A

B

Figure 3: **Processing of spike-glycoprotein at S2' is enhanced in presence of ACE2.** Western blot showing the impact of ACE2 on the processing of WT and µS1/S2 spike-glycoproteins by Furin. HeLa cells expressing empty vector (V), WT proS (**A**) or its µS1/S2 mutant (**B**) without or with Furin, ACE2 or both were analysed by Western blotting using anti-V5 antibody. The ratio of cDNAs used was S:ACE2:Furin = 1:1:1. The data are representative of at least three independent experiments.

A

Structure of Furin inhibitors

BOS-318

B

	Furin pIC50	PC5 pIC50	PACE4 pIC50	PC7 pIC50
BOS-318	8.8±0.4 (n=10)	6.7±0.15 (n=8)	6.7±0.15 (n=6)	7.4±0.22 (n=8)
BOS-981	9.3±0.5 (n=10)	7.5±0.22 (n=10)	6.9±0.1 (n=4)	6.9±0.2 (n=8)
BOS-857	9.4±0.3 (n=10)	7.6±0.2 (n=10)	6.7±0.24 (n=5)	6.9±0.3 (n=8)
decanoyl-RVKR-cmk	9.1±0.43 (n=447)	9.9±0.38 (n=162)	9.2±0.24 (n=214)	9.6±0.63 (n=198)

Figure 4

C

S1/S2:
TNSPRRAR↓SVAS

IC_{50}

BOS-981: 4 nM ± 0.7
BOS-857: 32 nM ± 3.9
BOS-318: 35 nM ± 4.9

X-axis: [BOS-inhibitor] nM
Y-axis: Furin activity (%)

D

	Golgi pIC50 (U20S)
BOS-318	7.7±0.24 (n=22)
BOS-981	8.3±0.25 (n=12)
BOS-857	7.6±0.2 (n=12)
decanoyl-RVKR-cmk	5.1±0.33 (n=8)

E

HeLa-ACE2

Figure 4: **Inhibition of PCs by BOS compounds.** (**A**) Chemical motif of BOS-inhibitors and representative structure of BOS-318. (**B**) *In vitro* BOS-inhibition of the cleavage of the fluorogenic dibasic substrate FAM-QRVRRAVGIDK-TAMRA by each of the proprotein convertases Furin, PC5 (PCSK5), PACE4 (PCSK6) and PC7 (PCSK7). All experiments were performed in 10 different wells and the average pIC_{50} (in nM) was calculated. Shown for comparison is the inhibitory pIC_{50} of the Furin-like inhibitor dec-RVKR-cmk performed >100 times. (**C**) *In vitro* inhibition of Furin by the BOS compounds. Furin (2 nM) was incubated with increasing concentration of BOS-inhibitors, and its enzymatic activity against the synthetic peptides DABSYL/Glu-TNSP**RRAR**↓SVAS-EDANS (5 μM) was measured at pH 7.5 (n=3). (**D**) Golgi assay: table representing the effects of BOS-inhibitors on U2OS cells expressing each of Furin, PC5A, PACE4 and PC7 simultaneously transduced with a BacMam-delivered construct containing a Golgi-targeting sequence followed by a 12-amino acid Furin/PCSK cleavage site from Bone Morphogenic Protein 10 (BMP10) and GFP at the C terminus (GalNAc-T2-GGGGS-DSTARIRR↓NAKG-GGGGS-GFP). Dibasic cleavage releases NAKG-GGGGS-GFP thereby reducing the Golgi-associated fluorescence estimated by imaging. (**E**) Furin-inhibitors (BOS) abrogate endogenous processing of the spike-glycoprotein. Hela cells were transiently transfected with a cDNA encoding an empty vector (V) or with one expressing the V5-tagged spike (S) glycoprotein (spike-V5). At 5h pre-transfection, cells were treated with vehicle DMSO (NT, duplicate) or with the Furin-inhibitors at indicated concentrations, or RVKR-cmk at 50 μM. At 24h post-transfection media were replaced with fresh ones lacking (NT) or containing the inhibitors for an additional 24h. Cell extracts were analyzed by Western blotting using a mAb-V5. All data are representative of at least three independent experiments.

Figure 5: **Furin-like inhibitors and Camostat treatment decrease SARS-CoV-2 infection in Calu-3 Cells.** (**A**) Replication kinetics was studied at 12, 24 and 48h post-infection by plaque assay to determine PFUs of SARS-CoV-2 virus in the supernatant of infected Calu-3 cells treated or not with 1µM BOS-318, BOS-857 and BOS-981. A line graph represents results of the triplicate plaque assay results (mean ± SD). (**B**) The virus titers (PFU per milliliter) released in the supernatant (24h post-infection) of infected Calu-3 cells treated with indicated concentrations of BOS-318 were determined by plaque assay (mean ± SD of triplicates, *p < 0.05; **p < 0.01; ***p < 0.001) (left panel). The selectivity index (SI) of BOS-318 in Calu-3 cells as shown in top right panel was determined by CC_{50}/IC_{50}. The left y axis indicates the inhibition of virus titer (percent) relative to that of the untreated control group (red). The right y axis indicates the cell viability (percent) relative to that of the untreated control group (green). The CC_{50} (50% cytotoxic concentration), IC_{50} (half maximal inhibitory concentration), and SI (selectivity index) values for each inhibitor are as shown. Representative plaque images of infected Calu-3 cells treated with indicated doses of BOS-inhibitors are shown in the bottom right panel. (**C**) Immunoblots for the infected Calu-3 cells (right panel) and viral particles secreted in the supernatant (left panel) with and without treatment with BOS-inhibitors indicate reduced viral protein levels. Immunoblots were probed for the full-length (proSm) and cleaved (S2) fragments of viral S protein and nucleocapsid (N) protein as indicated; β-Actin was included as the loading control for the cells. (**D**) The virus titers (PFU per milliliter) released in the supernatant (24h post-infection) of infected Calu-3 cells treated with BOS-318 and/or Camostat (Camo) were determined by plaque assay (mean ± SD of duplicates, *, p < 0.05; **, p < 0.01; ***, p < 0.001) (top panel). Representative plaque images of infected Calu-3 cells are shown in the bottom panel. Color plaques differentiate the lawn (one color gray per well) from individual plaques (independent colors).

Figure 6: **Processing of SARS-CoV-2 S by Furin-like convertases and TMPRSS2 is critical for viral entry in human lung epithelial cells but not in model HEK293 cells stably expressing ACE2**. (**A**) Furin cleavage of proS at the S1/S2 site is required for SARS-CoV-2 pseudoviral entry in Calu-3 cells but not HEK293T-ACE2 cells. Cells were inoculated with nanoluciferase-expressing HIV particles pseudotyped with SARS-CoV-2: wild-type spike (WT), double Ala-mutant spike (μS1/S2) or Furin-optimized spike (μS2'). Inhibition of proS processing at S1/S2 by a novel Furin-like inhibitor (BOS-318) during pseudovirion packaging prevents viral entry in Calu-3 cells but not in HEK293T-ACE2 cells. (**B**) Western blot analyses show inhibition by BOS-318 of proS processing at S1/S2 site. Purified pseudovirions and cellular extracts of producing HEK293-T17 cells treated or not with BOS-318 inhibitor were separated on SDS-PAGE gel and analyzed for HIV-1 p24 and V5-tagged S-protein (proSm or cleaved, S2) as indicated. (**C**) Pre-treatment of Calu-3 cells with 1 μM BOS-318 (B), 100 μM Camostat (C) or both (B+C) markedly reduces viral entry. In Panels A and C, Calu-3 cells were transduced with nanoluciferase-expressing HIV particles pseudotyped with SARS-CoV-2 S WT, μS1/S2 or μS2' for 72h and analyzed for nano-luciferase expression. Viral entry was expressed as fold increase over that given by bald particles (pseudovirions made in the absence of S). Each dot represents a different experiment with median luciferase activity calculated from three biological replicates. Two to four experiments were performed for each cell type. Error bars indicate standard deviation (SD) from the mean.

A

Cell-to cell fusion assay

Donor

Hela

Tat   Tat

S

ACE2

TZM-bl

Acceptor

ACE2

S

Tat

Tat

Luc

Syncytium

B

S in TZM-bl-ACE2

C   ACE2 in acceptor cell

Fold RLU

p<0.0001

p=0.0001

p=0.0005

p<0.0001

V   μS1/S2   NT   DMSO   RVKR   BOS 318   BOS 981   BOS 857

Donor cells

S-WT

D

Fold RLU

p=0.0258

p=0.0330

S   μS1/S2   μS2'   P681H   P681R

Figure 7: **Spike-induced cell-to-cell fusion relies on Furin cleavage at S1/S2.** (**A**) Cell-to-cell fusion between donor cells (HeLa) expressing the fusogenic SARS-CoV-2 spike protein along with the HIV trans-activator Tat, and acceptor cells (TZM-bl) that express ACE2. Upon fusion, Tat is transferred from donor to acceptor cells, thereby inducing luciferase expression. (**B**) Cell-to-cell fusion was evaluated using confocal microscopy. A representative immunocytochemistry of Hela cells transfected with a vector expressing SARS-CoV-2 spike co-cultured with TZM-bl cells for 18h. The number of syncytia (multiple nuclei) was examined using CellMask™ to probe for the plasma membrane and Dapi to stain the nuclei. (**C**) Donor cells were transfected with vectors expressing either no protein (empty vector, V), μS1/S2, or WT-spike (S) in the absence (NT) or presence of vehicle (DMSO) or with the Furin-inhibitors BOS-318, BOS-981, BOS-857 (300 nM) or RVKR (10 μM). Acceptor cells were transfected with a vector expressing ACE2. After 48h, donor and acceptor cells were co-cultured for 18h. Relative luminescence units (RLU) were normalized to the V value arbitrarily set to 1. Data are presented as mean values ± SD (n=3), One-Way ANOVA, Dunn-Sidàk multiple comparison test. (**D**) Donor HeLa cells expressing WT-S or its indicated mutants and variants were co-cultured with acceptor TZM-bl cells expressing ACE2. The extent of fusion is represented as a ratio between the RLU measured for each condition and that of donor cells expressing empty vector. The bar graph represents the average of 4 experiments performed in triplicates. Data are presented as mean values ± SEM (n=3 or 4), One-way Anova Dunnett's multiple comparison test.

Figure 8

Figure 8: **Spike-glycoprotein processing by Furin and TMPRSS2 in a co-culture system and their role in cell-to-cell fusion.** Donor HeLa cells expressing empty vector (V), WT-S-HA or μS1/S2-HA were co-cultured with acceptor TZM-bl cells expressing V, ACE2 + V, or ACE2 + TMPRSS2. From the same experiment, cell-to-cell fusion (**A**) was assessed in parallel with spike processing in cells and media by Western blotting (**B**). (**B**) spike-glycoproteins in the cell extracts were analyzed by Western blotting using an anti-V5 mAb. Secreted forms of spike protein (S1) in the media were detected with anti HA-HRP upon immunoprecipitation with anti-HA agarose. The bar graph represents the average of 3 experiments performed in triplicates. Data are presented as mean values ± SD Two-way Anova Turkey's multiple comparison test. The corresponding Western-blot is representative of three independent experiments. Values of S2 and S2' relative to β-actin are shown (nd = too low or not detected).

A

B

Media — IP: HA; WB: HA

Cells

C

Media — IP: HA; WB: HA

Cells

Figure 9: **Exogenous TMPRSS2-generated shedding of ACE2 differentially regulates S-induced fusion at the plasma membrane of WT-S *versus* μAS1/S2.** Donor HeLa cells expressing double tagged (N-terminal HA-tag; C-terminal V5-tag) spike-glycoprotein WT (S) or its S1/S2 mutant (μAS1/S2) were co-cultured with acceptor TZM-bl cells expressing (empty vector, V), ACE2 + V, or ACE2 + TMPRSS2 and treated with DMSO (vehicle control) or Camostat (120μM). Within the same experiment, cell-to-cell fusion (**A**) was assessed in parallel with spike processing in cells and media by Western-blot (**B, C**). (**A**) The extent of fusion is represented as a ratio between the RLU measured for each condition and that of donor cells expressing V. The bar graph represents the average of 2 experiments performed in triplicates. (**B, C**). Western blot analyses of media and cell extracts of the co-cultured cells with donor cells overexpressing double tagged (N-terminal HA-tag; C-terminal V5-tag) spike-glycoprotein, WT (S) (**B**) or μAS1/S2 (**C**). The arrow points to a putative degradation product of S2', that is absent in presence of Camostat. Media were subjected to immunoprecipitation with anti-HA agarose for the secreted forms of spike protein (S1, $S1_L$) followed by Western blotting with anti HA-HRP. spike-glycoproteins in the cell extracts were immunoblotted with anti-V5 mAb. The data are representative of three independent experiments.

Figure 10

Figure 10: **The C-terminal collectrin-like domain of ACE2 may be critical for the regulation of cell-to-cell fusion of spike-glycoprotein when exogenous TMPRSS2 is present.** (**A**) Schematic representation of the primary structure of human ACE2 with emphasis on the C-terminal collectrin-like domain (aa 616-768, light grey), TMPRSS2 cleavage region (aa 697-716, black) and the polybasic amino-acid segments in which K/R were mutated to A (C0 and C4) (amino acids underlined and in bold). Also shown are the peptidase domain (aa 19-615, white) containing the regions involved in the interaction with the spike SARS-CoV protein (hatched) and transmembrane domain (TM). (**B**) HeLa cells were co-transfected with ACE2, WT (ACE2) or its mutant ACE2 (C0+C4), and TMPRSS2, WT (TMPRSS2) or its S441A active-site mutant (μTMPRSS2), or empty vector (V). Media and cell extracts were analyzed by western blotting for shed ACE2 (sACE2) and ACE2, respectively. The migration positions of the ~95 kDa and ~80 kDa sACE2 are emphasized. (**C**) Donor HeLa cells expressing WT-S-HA or μAS1/S2-HA were co-cultured with acceptor TZM-bl cells expressing ACE2, WT (ACE2) or its mutant ACE2 (C0+C4) in presence or absence of TMPRSS2. From the same experiment, cell-to-cell fusion was assessed (**C**), in parallel with WB analyses of cells and media (**D**). The extent of fusion is represented as a ratio between the RLU measured for each condition and that of donor cells expressing an empty vector. The bar graph represents the average of 3 experiments performed in triplicates. Data are presented as mean values ± SD (n=3), One-way Anova Turkey's multiple comparison test. (**D**) Co-culture media were subjected to immunoprecipitation with anti-HA agarose for the secreted forms of spike protein (S1) followed by western blot with anti HA-HRP. spike-glycoproteins in the cell extracts were immunoblotted with anti-V5 mAb. The Western blot data are representative of three independent experiments.

Figure 11: **Proposed model for the processing of S-protein and its blockade by Furin and TMPRSS2 inhibitors.**

Boxed left panel: Schematic representation of the S-glycoprotein domains of SARS-CoV-2, including the N-terminal (NTD) and C-terminal (CTD) domains of S1, the Furin-S1/S2 and the Furin/TMPRSS2-S2' processing sites as well as the fusogenic α-helix that follows S2'. Binding of the RBD domain of S1 to the membrane associated ACE2 in target cells, and the cell surface expression of TMPRSS2 and Furin are also schematized. Right panels: **(1)** BOS-inhibitors (or μS1/S2 mutant) completely prevent cell-to-cell fusion of donor Hela cells expressing S-glycoprotein with acceptor HeLa-ACE2 cells, which lack endogenous TMPRSS2. In this context, this reveals that Furin is a major processing enzyme cleaving at S1/S2 and generating S2'. **(2)** In acceptor HeLa cells expressing TMPRSS2 (+), maximal prevention of cell-to-cell fusion can be achieved by a combination of Furin (BOS, phenocopying the μS1/S2 or μAS1/S2 mutants) and TMPRSS2 (Camostat) inhibitors blocks S2' production, ACE2-shedding (sACE2) and the separation of sACE2-S1L complex from S2. **(3)** Optimal blockade of SARS-CoV-2 infection of Calu-3 cells, which express endogenously both Furin and TMPRSS2, is also achieved by a combination of Furin (BOS) and TMPRSS2 (Camostat) inhibitors.