Memory precision of object-location binding is unimpaired in *APOE* ε4-carriers with spatial navigation deficits

Helena M. Gellersen[1], Gillian Coughlan[2], Michael Hornberger[3], Jon S. Simons[1*]

[1]Department of Psychology, University of Cambridge, United Kingdom
[2]Rotman Research Institute, Baycrest, Toronto, Canada
[3]Norwich Medical School, University of East Anglia, Norwich, United Kingdom

[*]Corresponding author

Abstract

Research suggests that tests of memory fidelity, feature binding and spatial navigation are promising for early detection of subtle behavioural changes related to Alzheimer's disease (AD). In the absence of longitudinal data, one way of testing the early detection potential of cognitive tasks is through the comparison of individuals at different genetic risk for AD. Most studies have done so using samples aged 70 years or older. Here, we tested whether memory fidelity of long-term object-location binding may be a sensitive marker even among cognitively healthy individuals in their mid-60s by comparing participants at low and higher risk based on presence of the ε4-allele of the apolipoprotein gene (*n*=26 ε3ε3 and *n*=20 ε3ε4 carriers). We used a continuous report paradigm in a visual memory task that required participants to recreate the spatial position of objects in a scene. We employed mixture modelling to estimate the two distinct memory processes that underpin the trial-by-trial variation in localisation errors: retrieval success which indexes the proportion of trials where participants recalled any information about an object's position and the precision with which participants retrieved this information. Prior work has shown that these memory paradigms that separate retrieval success from precision are capable of detecting subtle differences in mnemonic fidelity even when retrieval success could not. Nonetheless, a Bayesian analysis found good evidence that ε3ε4 carriers did not remember fewer object locations ($F(1, 42)=.450$, $p=.506$, $BF_{01}=3.02$), nor was their precision for the spatial position of objects reduced compared to ε3ε3 carriers ($F(1, 42)=.12$, $p=.726$, $BF_{01}=3.19$). Importantly, ε3ε4-carriers from the same sample have previously been reported to exhibit wayfinding deficits in a spatial navigation task (Coughlan et al. 2019, *PNAS*, 116(19)). The sensitivity of memory fidelity tasks may therefore not extend to individuals with one ε4-allele in their early to mid-60s. These results provide further support to prior proposals that spatial navigation may be a sensitive marker for the earliest AD-dependent cognitive changes, even before episodic memory. More research in preclinical AD is needed to confirm this hypothesis by direct comparison of memory fidelity and spatial navigation tasks.

Keywords: Alzheimer's disease, APOE, early detection, memory, spatial navigation

Introduction

Alzheimer's disease (AD) has a long preclinical phase during which pathological neural changes occur without overt, detrimental effects on behaviour.[1–4] This long preclinical phase offers the possibility of interventions that may target further pathological changes and prevent irreversible cell death.[5,6] Cognitive tests are the most cost-effective and simple way to screen for cognitive impairment related to dementia, yet standard neuropsychological tests typically fail to detect these subtle preclinical symptoms of AD pathology.[7,8] In the absence of longitudinal data, carriers of the major genetic risk factor for late-onset Alzheimer's disease are a good model to test the diagnostic sensitivity of cognitive tests.[7,9–11] Heterozygotes who carry one ε4-allele of the apolipoprotein (*APOE*) gene are three times more likely to develop AD compared to the "wildtype" ε3ε3 carriers, while those with two alleles (homozygotes) have a 15-fold higher likelihood.[12] People with an ε4-allele tend to exhibit AD pathology and other biomarkers at an earlier age,[13] and show a faster rate of cognitive decline than other genotypes.[14,15]

Current efforts in early detection research deem tests of feature binding, mnemonic discrimination and spatial navigation as particularly promising markers of incipient cognitive decline as they all crucially depend on the integrity of regions affected in the earliest stages of AD.[1,16–19] For example, posterior parietal areas are among the first to exhibit amyloid beta pathology,[1,16] while the transentorhinal region, the transition zone between lateral and medial aspects of the entorhinal cortex (ERC), is the earliest site of tau neurofibrillary tangles, which then spread to the hippocampus.[17–19] Memory for high fidelity representations crucially relies on hippocampus, ERC, perirhinal cortex and posterior parietal regions. The ERC supports high fidelity representations of objects, spatial features of objects and the environment,[20–26] which are bound by the hippocampus into holistic episodes[27–31] that can be faithfully reinstated upon retrieval with the concerted action of the recollection network.[32,33] Importantly, tau spreads from transentorhinal to parietal regions of the recollection network that are crucial for memory fidelity,[21] and the communication of these brain regions is already altered in preclinical AD and genetic-risk carriers.[34–36] Based on these preclinical changes, tasks that tap into the functional integrity of AD vulnerable brain regions may prove clinically meaningful, which has prompted researchers to investigate the utility of mnemonic fidelity tests in preclinical and prodromal dementia cases.

In studies across the AD continuum, a number of tests of memory fidelity and relational binding is in use: the mnemonic similarity task, which requires the discrimination between previously studied target items and highly similar lures;[22] memory binding tasks to test for intact associations between the identity of an object and its features (e.g. location, colour);[23] and continuous report paradigms in which feature binding is not assessed by categorical responding but requires participants to precisely recreate studied object features.[24,25] Older adults, individuals with mild cognitive impairment (MCI) and preclinical individuals with positive AD biomarkers exhibit significant deficits in differentiating novel stimuli from previously studied targets under conditions of high feature overlap.[22,27–34,37,38] Similarly, MCI and AD patients, and individuals without apparent cognitive deficits but very high AD risk based on genotype or biomarkers of AD pathology, also perform significantly worse in tests of feature binding, showing a marked decline in representational fidelity.[39–5960]

Despite evidence of reduced memory fidelity in aging and prodromal AD, its sensitivity as a marker of preclinical genetic vulnerability in otherwise cognitively normal adults has rarely been tested. Moreover, mean ages in these studies are typically around 70 or older.[61,62] Although prior studies have shown that AD pathology begins to accrue in midlife,[63] most studies of memory fidelity do not include middle-aged individuals (exceptions are [32,64]). Here we include both middle-aged and older adults to determine whether memory fidelity

differs by *APOE* status (ε3ε3 vs ε4-carriers). We focus on a version of a feature binding task that uses study-test delays that preclude the use of working memory. Studies on short-term feature binding suggest these metrics may be sensitive to cognitive decline in familial and prodromal AD cases.[45,59,60] However, carriers of the *APOE* ε4-allele may have an advantage in short study-test delays but may be predisposed to accelerated rate of forgetting thereafter.[52,60] A longer study-test delay may be able to index such faster forgetting.

We capitalise on current evidence for subtle deficits in preclinical AD by using a memory precision task with demands on memory binding, fidelity of mnemonic representations and visuospatial construction, all abilities that are particularly affected by AD pathology even from preclinical stages onwards.[6,31,65] Moreover, we use a computational modelling approach to isolate the probability of recalling any task-relevant content and the precision of the retrieved memory representations.[24,66] This is in contrast to previous studies that rely on categorical definitions of correctly vs. incorrectly recalled information. Our precision metric therefore adds a potentially more sensitive qualitative dimension to the memory assessment by indexing the trial-by-trial variability in memory fidelity, which may be capable of identifying even subtle differences between individuals that categorical responding may fail to unmask.[50,67]

Only one study has tested the fidelity of relational binding with longer memory retention intervals using continuous report paradigms in an *APOE* genotyped cohort.[52] Zokaei and colleagues show a reduction in the fidelity of object-location binding for preclinical ε4 homozygote older adults, but no such effect for ε3ε4 heterozygotes when using the mean error between target location and response as a performance metric. The presence of an effect of the ε4-allele on the fidelity of long-term object-memory binding is promising as it suggests that this task is potentially sensitive to preclinical AD-related changes even in individuals in their 60s. There may be ways in which such object-location tasks could be made more sensitive, such that performance could index even subtler changes in memory, such as those that might exist in a medium risk group such as ε4-heterozygotes. Other studies typically use mean target-response distance to index memory fidelity,[52,68] yet this metric includes both successfully retrieved and forgotten locations, thereby resulting in a potentially strong influence of guess trials on performance. It is therefore possible that performance reductions might be observed not just in ε4 homozygotes but also heterozygotes for object-location binding when using a more specific index of precision that controls for accessibility of any information from memory. Another option may be to increase the number of objects tested per study display. Increasing the amount of interfering information may place further demands on transentorhinal and hippocampal processes,[69–71] and may result in a higher number of misbinding errors among individuals with poorer mnemonic representations, such as in cases with preclinical or prodromal AD.[59,68] Here, we use both approaches to investigate whether continuous report paradigms can be made even more sensitive than previously achieved.

We examine the utility of a novel test of memory fidelity of relational binding that engages regions vulnerable to early AD, supplemented with a mixture modelling approach that specifically indexes precision, to test the effect of the ε4-allele on the precision of object-location binding beyond short-term memory retention. We compare model-derived metrics with those used in prior studies with continuous report paradigms such as those by Zokaei et al.[52] to determine if the separation of precision and retrieval success may be able to tease apart subtle differences in memory abilities between *APOE* genotypes.[50] We also increase the degree of interference between exemplars for a given study episode to place more demands on high fidelity representations and pattern separation. Moreover, we apply our test to a sample that has previously been characterised in terms of spatial navigation abilities.[9] An added benefit of our study is therefore to test whether a fidelity metric for spatial memory

will be similarly sensitive to AD risk as spatial navigation measures. To our knowledge no other study to date provides data on the effect of the ε4-allele on spatial memory fidelity and spatial navigation in the same preclinical sample.

Methods and methods

Participants

The study was carried out at the University of East Anglia, Norwich with ethical approval from the Faculty of Medicine and Health Sciences Ethics Committee at UEA (Reference FMH/2016/2017-11). Participants provided written informed consent before participation. Exclusion criteria included psychiatric or neurological disease, substance dependence disorder and clinical depression and/or anxiety. Individuals with medicated cholesterol or blood pressure were included in the study, but medication intake was recorded to ensure no medical differences between genetic groups. The sample presented here was previously described by Coughlan and colleagues.[9] The sample size was based on that of prior studies that investigated the effect of the ε4-allele on spatial navigation.[72]

Forty-nine participants completed the spatial precision memory task. We included *n*=26 individuals with the ε3ε3 genotype aged 53 to 74 (*M*=63.38, *SD*=5.99; 13 females) and *n*=20 individuals with the ε3ε4 genotype aged 54 to 80 (*M*=64.80, *SD*=6.83; 5 females) for our main analysis. Three volunteers with the ε4ε4 genotype aged 63 or 64 years also completed the test battery (*M*=63.33, *SD*=.58; 1 female). Given the small number of ε4 homozygotes and the differences between ε3ε4 and ε4ε4 carriers in general, in this study our main analysis focused on a comparison of ε3ε3 carriers and ε4 heterozygotes to avoid the admixture of different genotypes. In a sensitivity analysis, we determined whether the addition of the high risk ε4 homozygotes influenced the results.

The genotype groups did not differ significantly in terms of age or scores on the Addenbrookes Cognitive Examination (ACE) regardless of whether the total score or the memory subscore was used (all $t<1$, $p>.4$). In all analyses, we controlled for age and sex. Sample demographics and standard neuropsychological scores are shown in Table 1.

Table 1. Demographics and standard neuropsychological test scores by *APOE* genotype group.

	ε3ε3 (n=26) Mean (SD)	ε3ε4 (n=20) Mean (SD)	ε4ε4 (n=3) Mean (SD)
Age	63.4 (6.07)	64.8 (6.83)	63.3 (.58)
Sex			
Female	13 (50%)	5 (25%)	1 (33%)
Male	13 (50%)	15 (75%)	2 (67%)
Education	14.25 (2.31)	13.80 (2.26)	14.67 (.58)
ACE Total	93.9 (4.91)	94.6 (2.42)	93.0 (3.61)
ACE Memory	25.0 (1.50)	25.1 (1.00)	24.7 (1.53)
ROCF immediate	33.3 (2.59)	31.8 (2.69)	32.0 (2.65)
ROCF delayed	21.4 (5.73)	21.9 (4.67)	22.2 (9.78)

Note: ACE: Addenbrookes Cognitive Examination; ROCF: Rey-Osterrieth Complex Figure. Delayed copy was three minutes after presentation.

Precision memory task

Details for the precision memory task can be found in the Supplementary Material. Briefly, participants were asked to remember the identity and locations of objects in a scene. Each encoding display consisted of a trial-unique background image with three objects pseudorandomly arranged around an invisible circle centred at the midpoint of the image. Object positions were constrained to maintain a minimum of 62.04° between objects to avoid spatial overlap.

Participants undertook five practice trials before beginning the actual task. They were told that memorising the details of object and their locations during the study phase was crucial to support good performance in the test phase. Participants received feedback during training and only moved on to the main task after a minimum of five practice displays and after the experimenter had verified good comprehension of the task. The main task comprised five study-test blocks. In each of the five blocks, participants first viewed five displays during the study phase. After encoding, an interference task required participants to count backwards from a random number between 50 and 100 for 12 seconds to prevent active rehearsal of memorised displays. Each test trial began with an identification question where participants were asked to determine which of two presented objects had previously been shown. If they chose correctly, the associated background image appeared, and participants were asked to move the object around the screen to recreate its studied location as precisely as possible.

In total, participants viewed 25 displays and completed 75 test trials, each containing an identification and a localisation question.

Figure 1. Schematic of the precision memory task.

APOE genotyping

DNA was collected using a Darcon tip buccal swab (LE11 5RG; Fisher Scientific). Buccal swabs were refrigerated at 2–4 °C until DNA was extracted using the QIAGEN QIAamp DNA Mini Kit (M15 6SH; QIAGEN). DNA was quantified by analyzing 2-μL aliquots of each extraction on a QUBIT 3.0 fluorometer (LE11 5RG; Fisher Scientific). DNA extractions were confirmed by the presence of a DNA concentration of 1.5 μg or higher per 100 μg of AE buffer as indicated on the QUBIT reading. PCR amplification and plate read analysis was performed using Applied Biosystems 7500 Fast Real-Time PCR System (TN23 4FD; Thermo Fisher Scientific). TaqMan Genotyping Master Mix was mixed with two single-nucleotide polymorphisms of *APOE* (rs429358 at codon 112 and rs7412 at codon 158). These two single-nucleotide polymorphisms determine the genotype of *APOE*2, E3, and E4 (2007; Applied Biosystems).

Mixture modelling

Models fitted to the data and distribution of responses across all participants by genotype are shown in Fig. 2A, 2B. We fit probabilistic mixture models to the location placement errors expressed as the degrees separating the response from the target. This is a

well-established method to probe memory fidelity that has been widely used by both our work and others across short-term and long-term memory studies.[24,66,73–75] The approach aims to determine the distribution of trial responses in order to examine which retrieval mechanisms best explain the observed responses: i) correctly recalled locations, ii) random guesses or iii) a misbinding error in which the location of the target item is confused with that of another object also presented on the same display. Guess trials can be modelled using a uniform distribution. The proportion of trials within the uniform distribution represents the guess rate *pU* and 1-*pU* therefore expresses retrieval success *pT*. Correctly remembered items can be modelled by a circular Gaussian (von Mises) distribution centred at the target location with its standard deviation reflecting the precision with which locations are recalled. Larger standard deviations (*SD*) correspond to lower fidelity of the recalled responses. Finally, misbinding errors can be modelled by von Mises distributions centred around the two distractor items.

To maximise comparability with the only other study on the effect of the *APOE* genotype on location memory precision, we also used Bayesian modelling implemented with the MemToolbox in MATLAB 2016a (code available at: http://visionlab.github.io/ MemToolbox/).[73 75] Of note, that study employed a short- rather than a long-term memory task. We fit three models to the error data collapsed across all participants by *APOE* genotype group to test whether the same model components could explain responding in each group. The models contained the following components (Fig. 2A): Model 1 (von Mises distribution only) assumes that no guessing occurred; Model 2 (uniform + von Mises distribution) assumes that responses reflect a mixture of guessed trials and correctly recalled locations with response-to-target distance varying across trials; Model 3 (uniform + von Mises + von Mises for non-targets) extends Model 2 by additionally assuming that some incorrect responses were due to object-location misbinding. For more details on modelling and comparison with an alternative model fitting procedure based on work by Bays and colleagues[24,25,50] refer to the Supplementary Material.

This approach allowed us to test if the *APOE* ε4-allele affects the probability of correctly retrieving information from memory and/or the fidelity (i.e. the precision with which this information is recalled). Mixture modelling is superior to other approaches that distinguish between retrieval success and fidelity of retrieved information because the estimation of the uniform distribution accounts for guess responses placed near the target item.

A model selection procedure using Deviance Information Criterion favoured Model 2 which describes the standard mixture model with a uniform distribution for guessed responses and a von Mises distribution for the precision of correctly recalled responses as the best model. All further analyses are conducted using Model 2. Details on model selection and model metrics for the final model are shown in the Supplementary Material.

Finally, we calculated retrieval success and precision for each subject. When using individual subjects for model fitting, fewer trials are available for parameter estimation. Moreover, the estimation procedure may produce more robust and precise estimates for some subjects compared to others. To improve robustness of estimates for precision and retrieval success, we calculated a cut-off for guessing from the mixture modelling approach across the full sample following the examples of prior studies (for details see Supplementary

Material).[24,50] Localisation errors exceeding 63° response-to-target distance were deemed as failure to retrieve an object's location. Consequently, for each subject we calculated retrieval success as the proportion of trials with errors ≤63°. A measure of imprecision was derived from the standard deviation across all responses with localisation errors ≤63°.

Statistical analysis: Testing APOE group differences in retrieval success and memory precision.

We employed a combination of frequentist (two-tailed tests with a statistical significance level of $p<.05$) and Bayesian methods to test for APOE genotype effects.

Group differences based on permutations. We first tested for differences in guessing (*g*) and imprecision (*SD*) estimates for the standard mixture models fit to all responses across subjects in the ε3ε3-carrier and ε3ε4-carrier group, respectively. To obtain a *p*-value, true group differences were compared to the distribution of standardised differences obtained from random group assignments over 1000 permutations (sample 1 with *n*=26 to match the number of participants in the ε3ε3 group; sample 2 with *n*=20, as in the ε3ε4 group). This approach has the advantage of operating on more robust model parameters due to reduced noise resulting from larger number of trials available for mixture modelling.

Group differences while controlling for age and sex. Next, we carried out analyses on individual subject data while controlling for nuisance variables using a linear model with sex and age as covariates and APOE genotype as between-subjects factor of interest. Dependent variables were the proportion of correctly identified items, and the measures of retrieval success and precision based on a 63° cut-off for guessing (see Supplementary Material). We also calculated the Bayes Factor for the contrast of the model with covariates (age, sex) and the full model with covariates and ε4-status as between-subjects factor of interest using the R package *BayesFactor* (https://CRAN.R-project.org/package=BayesFactor). For the model comparison, the numerator was the control model with covariates, and the denominator was the model of interest (including both covariates and APOE genotype) such that a factor >1 was indicative of more support for the null as opposed to the alternative hypothesis. A Bayes Factor of >3 was deemed as strong evidence in support of the null hypothesis.[50,76,77]

Sensitivity analyses. Although our modelling approach suggested that misbinding errors did not account for a significant proportion of location memory responses across all participants, this does not take into account potential differences in misbinding frequency between participants. We therefore conducted a control analysis termed 'nearest neighbour analysis' as used in prior work.[52,78] This analysis allowed us to test whether there was a difference in the nature of incorrect responses between genetic groups by taking into account the occurrence of misbinding errors. A significant *APOE* effect on the distance to the nearest neighbour would suggest that error responses in the two groups are not caused by the same mechanisms. The group with significantly smaller nearest neighbour difference is likely to commit more misbinding errors.

In order to make our results more comparable with prior studies that used a similar object-location binding paradigm without mixture modelling to separate retrieval success from precision,[52,79] we also calculated the mean absolute error between targets and responses (see Supplementary Material). This would allow us to determine whether a modelling approach to separate retrieval success and memory precision may be more sensitive to detect

APOE ε4 effects on memory. To compare performance of the ε4ε4 carriers to that of the ε3ε3 group, we calculated their performance in terms of *Z*-scores based on the mean and standard deviation of test scores in the ε3ε3 group.

a.

Figure 2. Tested models and results from the mixture modelling approach.

(a) Proposed models to capture location memory performance. In Model 1, all object locations are assumed to be correctly recalled without any guess responses (probability of guessing: pU=0). The mean distance of responses from the target can be represented by the width of the von Mises (circular Gaussian) distribution, expressing the precision of memory recall (expressed as the standard deviation (*SD*) of the von Mises distribution where higher values reflect less precision; for a more intuitive interpretation where higher values reflect better performance, the *SD* value can be converted to the von Mises distribution concentration parameter *K*; see Supplementary Material). Model 2 assumes a mixture of guessed and correctly remembered responses, where the proportion of responses that fall within the uniform distribution is denoted by the parameter pU that captures the proportion of guessed responses. For a more intuitive understanding where higher values reflect

higher performance this parameter can also be expressed as retrieval success denoted by *pT*, the proportion of trials within the von Mises distribution, i.e. trials in which the target location was correctly recalled. Model 3 assumes that responses reflect a combination of guessing, correctly remembered responses with variable degree of precision, and swaps of target and distractor locations, represented as von Mises distributions centered at the locations of distractor objects. **(b) Distribution of location errors by ε4-status in native circular space (left hand side) and the Standard Mixture Model (von Mises + uniform) fit to responses.** Model 2 was identified as the best fitting model in a model comparison procedure detailed in the Supplementary Material.

Results

Effects of genotype on memory performance.

A summary of the memory performance metrics as a function of *APOE* group is shown in Table 2. Fig. 3 shows the comparison of retrieval success and precision estimates as a function of ε4-status.

Group differences based on memory metrics derived from modelling across subjects by APOE group. The results of the permutation analysis are shown in Fig. 3A. The error distributions derived from across subjects in each *APOE* group exhibited considerable overlap. The distribution of permutation-based group differences derived from random assignments to groups confirmed that the differences in guessing and imprecision between the *APOE* groups were not significant (*guessing*: $z=.31, p=.704=$; *imprecision*: $z=-.59, p=.555$).

Group differences based on memory metrics derived for each subject. We investigated the effect of *APOE* genotype on object identification, retrieval success for location and precision for correctly retrieved items at the individual subject level (Fig. 3C). The linear models controlling for age and sex found no significant effect of *APOE* on identification of objects ($F(1, 42)=1.14, p=.292, BF_{01}=2.17$), retrieval success for object locations ($F(1, 42)=.450, p=.506, BF_{01}=3.02$) or the precision of recreating locations of correctly retrieved items ($F(1, 42)=.12, p=.726, BF_{01}=3.19$).

To facilitate comparison with a previous study on *APOE* genotype effects on localisation performance in a continuous report paradigm we also compared mean absolute angular disparity between targets and responses across all trials.[52] There were no significant differences between *APOE* ε4-status groups ($F(1, 42)=1.14, p=.292, BF_{01}=1.99$).

Control analysis accounting for misbinding errors. In line with the results from the overall group comparisons based on permutations, there were no significant differences in the occurrence of swap errors. The Bayes Factor further provided good evidence in support of the absence of an *APOE* ε4 effect ($F(1, 42)=.83, p=.367; BF_{01}=2.54$).

Sensitivity analysis: effect of inclusion of ε4 homozygotes
Even with the addition of the high risk ε4 homozygotes the comparison between ε4-carriers and non-carriers showed no effect of AD genetic risk on object identification ($F(3, 45)=.25,$

p=.861, BF_{01}=3.16), retrieval success ($F(3, 45)$=1.36 p=.268, BF_{01}=3.33), precision of object location memory ($F(3, 45)$=.87, p=.462, BF_{01}=3.35), mean absolute localisation error ($F(3, 45)$=1.43, p=.246, BF_{01}=3.36) or the occurrence of swap errors ($F(3, 45)$=1.45, p=.242; BF_{01}=3.09). Looking at the performance of each ε4ε4 carrier individually, none of the *Z*-scores of the ε4ε4 carriers computed on the basis of the ε3ε3 group were larger than ±1.65 in any of item identification, retrieval success or precision, indicating no significant reductions in any of the participants at the highest genetic risk.

Table 2. Summary of memory performance across subjects by *APOE* genotype.

Metric	ε3ε3 (*n*=26)	ε3ε4 (*n*=23)	All ε4 carriers (*n*=26)
Identification accuracy	.83 (.06)	.81 (.08)	.83 (.08)
Location retrieved success	.80 (.13)	.80 (.16)	.81 (.15)
Localisation Precision	22.4 (5.31)	22.1 (3.97)	22.16 (4.09)
Mean target-response distance	35.5 (16.1)	35.1 (16.2)	34.10 (15.7)
Mean distance to nearest item	20.7 (5.60)	21.0 (5.00)	20.67 (4.84)
Model-derived estimates calculated across all subjects per group			
pU [95% CI]	.31 [.28; .34]	.29 [.26; .33]	.30 [.28; .33]
SD [95% CI]	17.90 [16.82; 19.32]	18.84 [17.54; 20.49]	18.35 [17.43; 19.30]

Note: Retrieval success and precision were calculated based on a model derived cut-off score for guessing at a response-to-target distance of 63° (see Supplementary Material for details). *CI:* credibility interval of the posterior distribution derived from the Bayesian estimation procedure.

Figure 3. Precision memory performance by *APOE* genotype.

(a) Distribution of standardised group differences derived from 1000 permutations where $n=26$ subjects were randomly assigned to one sample and $n=20$ subjects to another (to match the actual group sizes in our sample). Retrieval success and precision were obtained using mixture modelling on all trials across subjects for the ε3ε44 and the ε3ε4 group, respectively. The red dots represent the standardised true differences in model metrics calculated by subtracting the scores of the ε3ε4 from those of the ε3ε3 group (for *guessing*: $z=.31$; for *SD*: $z=-.5$). (b) Mean and standard deviation of identification accuracy, retrieval success and precision for each *APOE* group. Retrieval success refers to the proportion of trials falling within 63° of the target object. Precision reflects the standard deviation in response-to-target distance for all trials within 63° of the target object.

Discussion

In this study we aimed to determine whether the precision of long-term memory for object-location bindings is affected in healthy middle- and older-aged *APOE* ε4-carriers who do not exhibit impairments on standard neuropsychological tests. We used a continuous report paradigm in which participants were asked to recreate object locations as precisely as possible[24] and employed Bayesian mixture modelling to separate memory retrieval success from the precision of retrieved locations.[66,73] We hypothesised that the precision task may be capable of identifying subtle changes in memory fidelity in preclinical *APOE* ε4-carriers. Prior evidence suggests that such tasks have potential for early detection of AD-related cognitive changes because 1) the task involves relational binding of objects and locations, which is impaired in prodromal AD and taps into the integrity of the entorhinal cortex and hippocampus,[34,68,80–82] 2) a continuous metric may be a more sensitive index than categorical measures of retrieval,[50,67] and 3) memory fidelity relies on communication between hippocampus and cortical regions, which exhibit altered connectivity in the early course of AD.[1,16,24,83–90] Previously, preclinical ε4 homozygotes at high risk of AD were impaired on a similar memory fidelity task, while heterozygotes were not.[52] Here, we aimed to increase sensitivity of such continuous report paradigms by increasing the to-be-recalled information per test display and by separating memory precision from retrieval success. We then tested if these adjustments may be capable of picking up subtle differences between controls and a genetic risk group, even if the risk group was comprised of individuals with moderate genetic risk of AD (ε4 heterozygotes), who represent 47% of the general population.[91]

However, we found robust evidence for the absence of an effect of the ε4-allele on object-location long-term memory performance in preclinical middle-aged and older adults, regardless of whether the risk groups included only ε4 heterozygotes or additionally added the ε4 homozygotes. Carriers of the ε4 allele did not recall fewer locations of objects, nor was the precision of their retrieved object-location associations affected. E4-carriers also did not commit more misbinding errors of item identity and location. To our knowledge, this is the first study comparing cognitively healthy *APOE* genotype groups, using a mixture modelling approach to separate retrieval success from retrieval precision, with study-test delays that prevented the involvement of short-term memory. Intriguingly, despite this absence of spatial memory deficits, the ε4-carriers in this sample did exhibit altered wayfinding trajectories in real-time while navigating a virtual environment in the SeaHero Quest game, as previously shown by Coughlan and colleagues.[9]

Few studies have previously investigated memory fidelity in individuals at higher risk for AD during preclinical stages of the disease. Preclinical individuals with higher AD risk based on biomarkers or the *APOE* genotype have been reported to show poorer performance in mnemonic discrimination (combined group of heterozygotes and homozygotes) and continuous report tasks of feature binding in long-term memory (homozygotes).[31,59,92] Specifically, they exhibit a greater tendency to falsely label as old novel lures that are similar to studied stimuli.[31,62,92] They also have higher rates of misbinding and larger placement errors.[52] In the case of continuous report tasks it was shown that individuals with the highest genetic risk for late-onset AD (ε4 homozygotes) and familial AD do indeed have poorer representational quality for object-location binding, as measured using the mean error between target and response.[52,59] This metric encompasses both the success of retrieving any information from memory and the precision with which this information is recalled. In this study, we therefore tested whether the mixture modelling approach we applied might render this task more sensitive by isolating precision from retrieval success, and potentially identifying subtle reductions even in a genetic risk cohort consisting mainly of ε4 heterozygotes as in the present study. However, our data show that this was not the case, with Bayes factors providing good evidence for the null hypothesis that ε4-carriers are unimpaired on object recall or object localisation precision.

Given that *APOE* ε4 homozygotes and those with autosomal dominant AD show pathological markers at an earlier age than ε3ε4 carriers,[12,93,94] it is possible that impairments in the fidelity of object-location binding may be more apparent in preclinical ε4ε4 and early-onset AD cases, compared to the ε3ε4 carriers investigated in this study whose mean age was significantly below that of most other studies testing the use of memory fidelity tasks for early detection.[61,62,64,95] Our sensitivity analysis showed that the null effect was still confirmed after inclusion of a smaller group of ε4 homozygotes in the AD risk group.

Similar to results for continuous report paradigms, mnemonic similarity tasks have also found reduced memory fidelity in ε4-carriers.[61,62] These studies included participants who were on average five years older than those in our sample (mean ages of 70 vs. 65 years), which may have been a reason for the differences between their findings and ours. Notably, their ε4-carriers had deficits on the standard neuropsychological assessments, while no differences in such tasks were found in our sample. Those two studies[61,62] could therefore not establish the relative sensitivity of standard and memory fidelity tasks for early preclinical AD signs, although they do suggest that fidelity metrics may be a promising early detection tool in prodromal MCI.

Another strategy to test the sensitivity of early detection tasks is to classify cognitively normal preclinical older adults based on tau and amyloid AD biomarker status. To date, this has been done to test for the sensitivity of mnemonic discrimination tests, which shown a correlation between both tau and amyloid beta loads with mnemonic discrimination performance.[31,32,64,96,97] Again, mean ages in these samples (70+) tend to be significantly older than the participants in the present study (~65), although in one study the association between tau levels and object mnemonic discrimination could still be observed in individuals aged below 70 years.[31] Interestingly, the association of AD biomarker concentration and mnemonic discrimination deficits remained even after accounting for *APOE* status.[32] Unfortunately no such biomarker data exist for memory precision binding tasks.

Nevertheless, findings suggest that risk classification based on biomarkers, as opposed to ε4-genotype, may be a better strategy to test the sensitivity of memory precision for early detection of AD in preclinical samples without cognitive signs on standard tests.[98]

Currently, it is difficult to compare the sensitivity of precision binding and mnemonic similarity tasks to early AD-related decline given the considerable difference in mean ages of study samples and the different approaches to index preclinical AD cases. Future studies are needed to better understand the relationship between continuous and categorical metrics of memory fidelity as measured using our precision task and mnemonic discrimination tests, respectively. Based on moderate correlations between test scores, mnemonic similarity judgments and object-location binding may involve similar but also somewhat dissociable mechanisms.[99] Pattern separation likely contributes to good performance in both tasks.[99–101] Both mnemonic similarity and precision rely on the integrity of, and effective communication between, perirhinal cortex, ERC and hippocampus, particularly subregions CA3 and dentate gyrus.[28,84,102–105] Functional MRI data suggest that the precision task further engages posterior parietal cortex regions upon retrieval to allow for the faithful reinstatement and visuospatial reconstruction of object features[24,84] and that hippocampal activity at encoding scales with precision.[20] Importantly, MTL-parietal connectivity tracks memory precision, which is of note given that these connections may already be altered in the early stages of AD, and even among relatively young ε4-carriers.[83,84,106–108]

Despite the absence of a precision memory deficit in the present sample, Coughlan and colleagues recently described suboptimal navigation patterns in these same ε3ε4 carriers (a subset of whom were enrolled in this study).[9] This subtle navigational deficit was attributed to a bias toward navigating close to environmental boundaries. This very specific impairment is thought to be a result of early tau pathology in the ERC thought to alter the integrity of grid cell representations, which are essential for updating self-motion during navigation.[9,72,109–111] This interpretation is in line with recent evidence suggesting that preclinical ε4-carriers only exhibit spatial navigation deficits in the absence of nearby landmarks or environmental boundaries that normally correct for accumulating temporal error in the grid-cell code.[110,112]

Although grid cells are most famously involved in spatial navigation, they also support visual memory.[113] Research suggests that both visual and navigational processes are supported by the entorhinal cortex via common mechanisms that include the formation of spatial or visual maps via grid cells.[114,115] Specifically, grid cells code for spatial locations in a visual scene much in the same way in which they code for space during exploration of a 3D environment.[113,114] Based on these findings it has been proposed that grid cells support both spatial navigation and relational memory.[115] It may therefore be surprising that we did not find any effect in our relational memory precision task, while impairments were observed in the navigation task by Coughlan et al.[9] However, the border bias is a very specific behaviour in ε4-carriers that appears when arenas have larger open spaces where anchoring spatial maps to nearby landmarks cannot be used as a corrective strategy.[9,72,110] Therefore, it has no direct equivalent in 2D visual scene memory in our precision task. This may explain why there is an effect of the ε4-allele on virtual reality spatial navigation but not in object-location memory precision in our sample. A preference for environmental borders may indeed be the very first

sign of AD risk dependent behavioural changes, whereas impairment in relational memory may arise at a later stage.[62,92,116]

Despite our relatively small sample size, our power analysis suggested that our study had moderate power to detect a difference in precision memory between ε3ε3 and ε3ε4 carriers similar in magnitude to that of the spatial navigation test in the same individuals (Supplementary Material).[9] Even though power was moderate, our Bayesian analysis provided good evidence in favour of a null effect, suggesting that the absence of a genotype effect was not due to an inadequate sample size. If a genotype effect on object-location precision does indeed exist in this sample, it is likely to be rather small and may be less meaningful for early detection efforts. This small effect may in part be due to the high heterogeneity of ε3ε4 carriers, given that only a subgroup will move on to actually develop AD.[12] It would be informative to follow up the present sample longitudinally to test whether any differences in memory precision are apparent in the current data when comparing participants who do or do not subsequently exhibit cognitive decline associated with AD. Additionally, as discussed above, a promising strategy to test the sensitivity of the precision task in preclinical cases in future studies may be to use biomarkers for classifying individuals into risk groups. This would not only allow studies to determine the sensitivity of memory fidelity metrics but to also assess the specificity of these tasks to AD-related cognitive decline. This is particularly important given the high heterogeneity of ε4-carriers and MCI patients. To date there is still a lack of studies on memory fidelity that stratify MCI patient groups based on AD biomarkers.[40,45,46]

To our knowledge this is the first study to employ a modelling approach to separate episodic memory retrieval success and precision and test the sensitivity of mnemonic fidelity metrics to preclinical AD risk as measured in a contrast of ε3ε3 and ε3ε4 carriers. Prior work in high risk AD individuals (familial, ε3ε4/ε4ε4, tau and amyloid positive cases) has suggested that object-location memory fidelity may be a sensitive marker for preclinical AD cases and that this effect can be detected in samples aged 70 and older.[32,52,59 62,92] We provide robust evidence that this may not be the case for middle-aged ε3ε4 carriers who were, on average, five years younger than individuals in prior studies. The sensitivity of memory fidelity tasks may therefore not extend to ε4 heterozygotes in their early to mid-60s. Despite no APOE genotype effect on object-location memory precision, prior evidence has shown that ε3ε4 carriers from the same sample do exhibit subtle behavioural deficits in spatial navigation.[9] These results provide further support to prior proposals that spatial navigation may be a sensitive marker for the earliest AD-dependent cognitive changes, even before episodic memory.[72,117] More research in preclinical AD is needed to confirm this hypothesis by direct comparisons of memory fidelity and spatial navigation tasks.

Acknowledgements: The authors would like to thank their funders and all volunteers who have participated in this study.

Funding: This work was supported by an ARUK grant (ARUK-SHQ2018-001). HMG is funded by a Medical Research Council doctoral training grant (#RG86932) and a Pinsent Darwin Award. GC was funded by a Foundation Grant from CIHR (#143311), MH by BBSRC, NIHR, Wellcome Trust and the UK Department for Transport and JSS by a James S. McDonnell Foundation Scholar award #220020333. The funders had no role in the conceptualisation, analysis or publication of this data.

Competing interests: The authors have no competing interests to declare.

Data availability statement: Data are available at https://osf.io/42sp9/

References

1. Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement*. 2011;7(3):280-292. doi:10.1016/j.jalz.2011.03.003

2. Jack CR, Knopman DS, Jagust WJ, et al. Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol*. 2010;9(1):119-128. doi:10.1016/S1474-4422(09)70299-6

3. Jack CR, Holtzman DM. Biomarker modeling of alzheimer's disease. *Neuron*. 2013;80(6):1347-1358. doi:10.1016/j.neuron.2013.12.003

4. Sutphen CL, Jasielec MS, Shah AR, et al. Longitudinal cerebrospinal fluid biomarker changes in preclinical Alzheimer disease during middle age. *JAMA Neurol*. 2015;72(9):1029-1042. doi:10.1001/jamaneurol.2015.1285

5. Chetelat G, Villemagne VL, Bourgeat P, et al. Relationship between atrophy and beta-amyloid deposition in Alzheimer disease. *Ann Neurol*. 2010;67(3):317-324. doi:10.1002/ana.21955

6. Rentz DM, Parra Rodriguez MA, Amariglio R, Stern Y, Sperling R, Ferris S. Promising developments in neuropsychological approaches for the detection of preclinical Alzheimer's disease: A selective review. *Alzheimer's Res Ther*. 2013;5(6):1. doi:10.1186/alzrt222

7. O'Donoghue MC, Murphy SE, Zamboni G, Nobre AC, Mackay CE. APOE genotype and cognition in healthy individuals at risk of Alzheimer's disease: A review. *Cortex*. 2018;104:103-123. doi:10.1016/j.cortex.2018.03.025

8. Salmon DP. Neuropsychological Features of Mild Cognitive Impairment and Preclinical Alzheimer's Disease. In: *Behavioral Neurobiology of Aging*. Springer, Berlin, Heidelberg; 2011:187-212. doi:10.1007/7854_2011_171

9. Coughlan G, Coutrot A, Khondoker M, Minihane AM, Spiers H, Hornberger M. Toward personalized cognitive diagnostics of at-genetic-risk Alzheimer's disease. *Proc Natl Acad Sci*. 2019;116(19):201901600. doi:10.1073/pnas.1901600116

10. Grilli MD, Wank AA, Bercel JJ, Ryan L. Evidence for Reduced Autobiographical Memory Episodic Specificity in Cognitively Normal Middle-Aged and Older Individuals at Increased Risk for Alzheimer's Disease Dementia. *J Int Neuropsychol Soc*. 2018;24(10):1073-1083. doi:10.1017/S1355617718000577

11. Flowers SA, Rebeck GW. APOE in the normal brain. *Neurobiol Dis*. 2020;136(October). doi:10.1016/j.nbd.2019.104724

12. Raber J, Huang Y, Ashford JW. ApoE genotype accounts for the vast majority of AD risk and AD pathology. *Neurobiol Aging*. 2004;25(5):641-650. doi:10.1016/J.NEUROBIOLAGING.2003.12.023

13. Risacher SL, Kim S, Shen L, et al. The role of apolipoprotein E (APOE) genotype in early mild cognitive impairment (E-MCI). *Front Aging Neurosci*. 2013;5. doi:10.3389/fnagi.2013.00011

14. Caselli RJ, Dueck AC, Locke DEC, et al. Longitudinal modeling of frontal cognition in APOE ε4 homozygotes, heterozygotes, and noncarriers. *Neurology*. 2011;76:1383-1388.

15. Caselli RJ, Reiman EM. Characterizing the preclinical stages of Alzheimer's disease and the prospect of presymptomatic intervention. *J Alzheimer's Dis*. 2013;33(SUPPL. 1). doi:10.3233/JAD-2012-129026

16. Buckner RL, Snyder AZ, Shannon BJ, et al. Molecular, Structural, and Functional Characterization of Alzheimer's Disease: Evidence for a Relationship between Default

Activity, Amyloid, and Memory. *J Neurosci*. 2005;25(34):7709-7717. doi:10.1523/jneurosci.2177-05.2005

17. Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol*. 1991;82(4):239-259. doi:10.1007/BF00308809

18. Sperling R. The Challenges of Disambiguating Preclinical Alzheimer's Disease from Cognitive Aging. In: Cabeza R, Nyberg L, Park DC, eds. *Cognitive Neuroscience of Aging: Linking Cognitive and Cerebral Aging.* ; 2016. doi:10.1093/acprof:oso/9780199372935.001.0001

19. Sepulcre J, Schultz AP, Sabuncu M, et al. In vivo tau, amyloid, and gray matter profiles in the aging brain. *J Neurosci*. 2016;36(28):7364-7374. doi:10.1523/JNEUROSCI.0639-16.2016

20. Korkki SM, Richter FR, Simons JS. Hippocampal-cortical encoding activity predicts the precision of episodic memory. *bioRxiv*. Published online 2020:1-38. doi:10.1101/2020.11.10.376814

21. Khan UA, Liu L, Provenzano FA, et al. Molecular drivers and cortical spread of lateral entorhinal cortex dysfunction in preclinical Alzheimer's disease. *Nat Neurosci*. 2014;17(2):304-311. doi:10.1038/mp.2011.182.doi

22. Stark SM, Yassa MA, Lacy JW, Stark CEL. A task to assess behavioral pattern separation (BPS) in humans: Data from healthy aging and mild cognitive impairment. *Neuropsychologia*. 2013;51(12):2442-2449. doi:10.1016/j.neuropsychologia.2012.12.014

23. Parra MA, Abrahams S, Fabi K, Logie R, Luzzi S, Sala S Della. Short-term memory binding deficits in Alzheimers disease. *Brain*. 2009;132(4):1057-1066. doi:10.1093/brain/awp036

24. Richter FR, Cooper RA, Bays PM, Simons JS. Distinct neural mechanisms underlie the success , precision , and vividness of episodic memory. Published online 2016:1-18. doi:10.7554/eLife.18260

25. Peich MC, Husain M, Bays PM. Age-related decline of precision and binding in visual working memory. *Psychol Aging*. 2013;28(3):729-743. doi:10.1037/a0033236

26. Reagh ZM, Noche JA, Tustison NJ, Delisle D, Murray EA, Yassa MA. Functional Imbalance of Anterolateral Entorhinal Cortex and Hippocampal Dentate/CA3 Underlies Age-Related Object Pattern Separation Deficits. *Neuron*. 2018;97(5):1187-1198.e4. doi:10.1016/j.neuron.2018.01.039

27. Berron D, Neumann K, Maass A, et al. Age-related functional changes in domain-specific medial temporal lobe pathways. *Neurobiol Aging*. 2018;65:86-97. doi:10.1016/j.neurobiolaging.2017.12.030

28. Yassa MA, Stark SM, Bakker A, Albert MS, Gallagher M, Stark CEL. High-resolution structural and functional MRI of hippocampal CA3 and dentate gyrus in patients with amnestic Mild Cognitive Impairment. *Neuroimage*. 2010;51(3):1242-1252. doi:10.1016/j.neuroimage.2010.03.040

29. Ally BA, Hussey EP, Ko PC, Molitor RJ. Pattern separation and pattern completion in Alzheimer's disease: Evidence of rapid forgetting in amnestic mild cognitive impairment. *Hippocampus*. 2013;23(12):1246-1258. doi:10.1002/hipo.22162

30. Leal SL, Yassa MA. Integrating new findings and examining clinical applications of pattern separation. *Nat Neurosci*. 2018;21(2):163-173. doi:10.1038/s41593-017-0065-1

31. Berron D, Cardenas-blanco A, Bittner D, et al. Higher CSF Tau Levels Are Related to Hippocampal Hyperactivity and Object Mnemonic Discrimination in Older Adults. *J Neurosci*. 2019;39(44):8788-8797. doi:10.1523/JNEUROSCI.1279-19.2019

32. Webb CE, Foster CM, Horn MM, Kennedy KM, Rodrigue KM. NeuroImage Beta-

amyloid burden predicts poorer mnemonic discrimination in cognitively normal older adults. *Neuroimage*. 2020;221(April):117199. doi:10.1016/j.neuroimage.2020.117199

33. Stark SM, Stark CEL. Age-related deficits in the mnemonic similarity task for objects and scenes. *Behav Brain Res*. 2017;333(April):109-117. doi:10.1016/j.bbr.2017.06.049

34. Reagh ZM, Roberts JM, Ly M, Diprospero N, Murray E, Yassa MA. Spatial discrimination deficits as a function of mnemonic interference in aged adults with and without memory impairment. *Hippocampus*. Published online 2014. doi:10.1002/hipo.22224

35. Reagh ZM, Ho HD, Leal SL, et al. Greater loss of object than spatial mnemonic discrimination in aged adults. *Hippocampus*. 2016;26(4):417-422. doi:10.1002/hipo.22562

36. Gellersen HM, Trelle AN, Henson RN, Simons JS. Executive function and high ambiguity perceptual discrimination contribute to individual differences in mnemonic discrimination in older adults. *PsyArXiv*. Published online 2020. doi:10.31234/osf.io/84vzc

37. Trelle AN, Carr VA, Wilson EN, et al. Association of CSF biomarkers with hippocampal-dependent memory in preclinical Alzheimer disease. *Neurology*.

38. Yassa MA, Mattfeld AT, Stark SM, Stark CEL. Age-related memory deficits linked to circuit-specific disruptions in the hippocampus. *Proc Natl Acad Sci*. 2011;108(21):8873-8878. doi:10.1073/pnas.1101567108

39. Parra MA, Saarimäki H, Bastin ME, et al. Memory binding and white matter integrity in familial Alzheimer's disease. *Brain*. 2015;138(5):1355-1369. doi:10.1093/brain/awv048

40. Mowrey WB, Lipton RB, Katz MJ, et al. Memory Binding Test Predicts Incident Amnestic Mild Cognitive Impairment. *J Alzheimer's Dis*. 2016;53(4):1585-1595. doi:10.3233/JAD-160291

41. Della Sala S, Parra MA, Fabi K, Luzzi S, Abrahams S. Short-term memory binding is impaired in AD but not in non-AD dementias. *Neuropsychologia*. 2012;50(5):833-840. doi:10.1016/j.neuropsychologia.2012.01.018

42. Parra MA, Calia C, García AF, et al. Refining memory assessment of elderly people with cognitive impairment: Insights from the short-term memory binding test. *Arch Gerontol Geriatr*. 2019;83(February 2018):114-120. doi:10.1016/j.archger.2019.03.025

43. Pietto M, Parra MA, Trujillo N, et al. Behavioral and Electrophysiological Correlates of Memory Binding Deficits in Patients at Different Risk Levels for Alzheimer's Disease. *J Alzheimer's Dis*. 2016;53(4):1325-1340. doi:10.3233/JAD-160056

44. Valdés MC, Clark R, Wang S, et al. The striatum , the hippocampus , and short-term memory binding : Volumetric analysis of the subcortical grey matter ' s role in mild cognitive impairment. *NeuroImage Clin*. 2020;25(September 2019):102158. doi:10.1016/j.nicl.2019.102158

45. Koppara A, Frommann I, Polcher A, et al. Feature Binding Deficits in Subjective Cognitive Decline and in Mild Cognitive Impairment. *J Alzheimer's Dis*. 2015;48(S1):S161-S170. doi:10.3233/JAD-150105

46. Troyer AK, Murphy KJ, Anderson ND, et al. Associative recognition in mild cognitive impairment: Relationship to hippocampal volume and apolipoprotein E. *Neuropsychologia*. 2012;50(14):3721-3728. doi:10.1016/j.neuropsychologia.2012.10.018

47. Konijnenberg E, den Braber A, ten Kate M, et al. Association of amyloid pathology with memory performance and cognitive complaints in cognitively normal older adults: a monozygotic twin study. *Neurobiol Aging*. 2019;77:58-65.

doi:10.1016/j.neurobiolaging.2019.01.006

48. Hampel H. Amyloid-β and cognition in aging and Alzheimer's disease: Molecular and neurophysiological mechanisms. *J Alzheimer's Dis*. 2013;33(SUPPL. 1). doi:10.3233/JAD-2012-129003

49. Rentz DM, Amariglio RE, Becker JA, et al. Face-name associative memory performance is related to amyloid burden in normal elderly. *Neuropsychologia*. 2011;49(9):2776-2783. doi:10.1016/j.neuropsychologia.2011.06.006

50. Korkki SM, Richter FR, Jeyarathnarajah P, Simons JS. Healthy Ageing Reduces the Precision of Episodic Memory Retrieval. *Psychol Aging*. 2020;35(1):124-142. doi:10.1037/pag0000432

51. Chen P-C, Chang Y-L. Associative memory and underlying brain correlates in older adults with mild cognitive impairment. *Neuropsychologia*. 2016;85:216-225. doi:10.1016/j.neuropsychologia.2016.03.032

52. Zokaei N, Čepukaitytė G, Board AG, Mackay CE, Husain M, Nobre AC. Dissociable effects of the apolipoprotein-E (APOE) gene on short- and long-term memories. *Neurobiol Aging*. 2019;73:115-122. doi:10.1016/j.neurobiolaging.2018.09.017

53. Polcher A, Frommann I, Koppara A, Wolfsgruber S, Jessen F, Wagner M. Face-Name Associative Recognition Deficits in Subjective Cognitive Decline and Mild Cognitive Impairment. *J Alzheimer's Dis*. 2017;56(3):1185-1196. doi:10.3233/JAD-160637

54. Delhaye E, Bahri MA, Salmon E, Bastin C. Impaired perceptual integration and memory for unitized representations are associated with perirhinal cortex atrophy in Alzheimer's disease. *Neurobiol Aging*. 2019;73:135-144. doi:10.1016/j.neurobiolaging.2018.09.021

55. Oedekoven CSH, Jansen A, Keidel JL, Kircher T, Leube D. The influence of age and mild cognitive impairment on associative memory performance and underlying brain networks. *Brain Imaging Behav*. 2015;9(4):776-789. doi:10.1007/s11682-014-9335-7

56. Van Geldorp B, Heringa SM, Van Den Berg E, Olde Rikkert MGM, Biessels GJ, Kessels RPC. Working memory binding and episodic memory formation in aging, mild cognitive impairment, and Alzheimers dementia. *J Clin Exp Neuropsychol*. 2015;37(5):538-548. doi:10.1080/13803395.2015.1037722

57. Bastin C, Bahri MA, Miévis F, et al. Associative memory and its cerebral correlates in Alzheimer's disease: Evidence for distinct deficits of relational and conjunctive memory. *Neuropsychologia*. 2014;63(1):99-106. doi:10.1016/j.neuropsychologia.2014.08.023

58. Atienza M, Atalaia-Silva KC, Gonzalez-Escamilla G, Gil-Neciga E, Suarez-Gonzalez A, Cantero JL. Associative memory deficits in mild cognitive impairment: The role of hippocampal formation. *Neuroimage*. 2011;57(4):1331-1342. doi:10.1016/j.neuroimage.2011.05.047

59. Liang Y, Pertzov Y, Nicholas JM, et al. Visual short-term memory binding deficit in familial Alzheimer's disease. *Cortex*. 2016;78(February):150-164. doi:10.1016/j.cortex.2016.01.015

60. Pavisic IM, Nicholas JM, Pertzov Y, et al. Visual Short-term Memory Impairments in Presymptomatic Familial Alzheimer ' s Disease : a Longitudinal Observational Study. *ResearchSquare*. Published online 2020. doi:10.21203/rs.3.rs-67640/v1

61. Sinha N, Berg CN, Tustison NJ, et al. APOE ε4 status in healthy older African Americans is associated with deficits in pattern separation and hippocampal hyperactivation. *Neurobiol Aging*. 2018;69:221-229. doi:10.1016/j.neurobiolaging.2018.05.023

62. Sheppard DP, Graves L V., Holden HM, Delano-Wood L, Bondi MW, Gilbert PE. Spatial pattern separation differences in older adult carriers and non-carriers for the

apolipoprotein E epsilon 4 allele. *Neurobiol Learn Mem*. 2016;129:113-119. doi:10.1016/j.nlm.2015.04.011

63. Ritchie K, Ritchie CW, Jaffe K, Skoog I, Scarmeas N. Is late-onset Alzheimer's disease really a disease of midlife? *Alzheimer's Dement Transl Res Clin Interv*. 2015;1(2):122-130. doi:10.1016/j.trci.2015.06.004

64. Maass A, Berron D, Harrison TM, et al. Alzheimer's pathology targets distinct memory networks in the ageing brain. *Brain*. 2019;142(8):2492-2509. doi:10.1093/brain/awz154

65. Ritchie K, Carrière I, Su L, et al. The midlife cognitive profiles of adults at high risk of late-onset Alzheimer's disease: The PREVENT study. *Alzheimer's Dement*. 2017;13(10):1089-1097. doi:10.1016/j.jalz.2017.02.008

66. Bays PM, Catalao RFG, Husain M. The precision of visual working memory is set by allocation of a shared resource. *J Vis*. 2011;9(10). doi:10.1167/9.10.7.The

67. Zokaei N, Burnett Heyes S, Gorgoraptis N, Budhdeo S, Husain M. Working memory recall precision is a more sensitive index than span. *J Neuropsychol*. 2015;9(2):319-329. doi:10.1111/jnp.12052

68. Hampstead BM, Towler S, Stringer AY, Sathian K. Continuous measurement of object location memory is sensitive to effects of age and mild cognitive impairment and related to medial temporal lobe volume. *Alzheimer's Dement Diagnosis, Assess Dis Monit*. 2018;10:76-85. doi:10.1016/j.dadm.2017.10.007

69. Newsome RN, Duarte A, Barense MD. Reducing perceptual interference improves visual discrimination in mild cognitive impairment: Implications for a model of perirhinal cortex function. *Hippocampus*. 2012;22(10):1990-1999. doi:10.1002/hipo.22071

70. Kirwan CB, Stark CEL. Overcoming interference: An fMRI investigation of pattern separation in the medial temporal lobe. *Learn Mem*. 2007;14(9):625-633. doi:10.1101/lm.663507

71. Reagh ZM, Yassa MA. Object and spatial mnemonic interference differentially engage lateral and medial entorhinal cortex in humans. *Proc Natl Acad Sci U S A*. 2014;111(40):E4264-E4273. doi:10.1073/pnas.1411250111

72. Kunz L, Schröder TN, Lee H, et al. Reduced grid-cell-like representations in adults at genetic risk for Alzheimer's disease. *Science (80-)*. 2015;350(6259):430-433. doi:10.1126/science.aac8128

73. Suchow JW, Brady TF, Fougnie D, Alvarez GA. Modeling visual working memory with the MemToolbox. *J Vis*. 2013;13(10):1-8. doi:10.1167/13.10.9

74. Zhang W, Luck SJ. Discrete fixed-resolution representations in visual working memory. 2008;453(May):233-236. doi:10.1038/nature06860

75. Zokaei N, Grogan J, Fallon SJ, et al. Short-term memory advantage for brief durations in human APOE ε 4 carriers. *Sci Rep*. Published online 2020:1-10. doi:10.1038/s41598-020-66114-6

76. Keysers C, Gazzola V, Wagenmakers EJ. Using Bayes factor hypothesis testing in neuroscience to establish evidence of absence. *Nat Neurosci*. 2020;23(7):788-799. doi:10.1038/s41593-020-0660-4

77. Jeffreys H. *Theory of Probability*. 3rd ed. Oxford University Press; 1961.

78. Pertzov Y, Miller TD, Gorgoraptis N, et al. Binding deficits in memory following medial temporal lobe damage in patients with voltage-gated potassium channel complex antibody-associated limbic encephalitis. *Brain*. 2013;136(8):2474-2485. doi:10.1093/brain/awt129

79. Zokaei N, Giehl K, Sillence A, et al. Sex and APOE: A memory advantage in male APOE ε4 carriers in midlife. *Cortex*. 2017;88:98-105.

doi:10.1016/j.cortex.2016.12.016

80.   Weigard AS, Sathian K, Hampstead BM. Model-based assessment and neural correlates of spatial memory deficits in mild cognitive impairment. *Neuropsychologia.* 2020;136(October 2019):107251. doi:10.1016/j.neuropsychologia.2019.107251

81.   Charles DP, Browning PGF, Gaffan D. Entorhinal cortex contributes to object-in-place scene memory. *Eur J Neurosci.* 2004;20(11):3157-3164. doi:10.1111/j.1460-9568.2004.03777.x

82.   McIlvain G, Schwarb H, Cohen NJ, Telzer EH, Johnson CL. Mechanical properties of the in vivo adolescent human brain. *Dev Cogn Neurosci.* 2018;34:27-33. doi:10.1016/j.dcn.2018.06.001

83.   Foo H, Mather KA, Jiang J, Thalamuthu A, Wen W, Sachdev PS. Genetic influence on ageing-related changes in resting-state brain functional networks in healthy adults: A systematic review. *Neurosci Biobehav Rev.* 2020;113(March):98-110. doi:10.1016/j.neubiorev.2020.03.011

84.   Cooper RA, Ritchey M. Cortico-hippocampal network connections support the multidimensional quality of episodic memory. *Elife.* 2019;8:1-37. doi:10.7554/elife.45591

85.   Jack CR, Wiste HJ, Weigand SD, et al. Age, Sex, and *APOE* ε4 Effects on Memory, Brain Structure, and β-Amyloid Across the Adult Life Span. *JAMA Neurol.* 2015;72(5):511. doi:10.1001/jamaneurol.2014.4821

86.   Xie W. A neurocognitive mechanism for precision of visual working memory representations. Published online 2018. doi:10.1017/CBO9781107415324.004

87.   Stevenson RF, Zheng J, Mnatsakanyan L, et al. Hippocampal CA1 gamma power predicts the precision of spatial memory judgments. *Proc Natl Acad Sci.* 2018;115(40):10148-10153. doi:10.1073/pnas.1805724115

88.   Sullivan MD, Anderson JAE, Turner GR, Spreng RN. Intrinsic neurocognitive network connectivity differences between normal aging and mild cognitive impairment are associated with cognitive status and age. *Neurobiol Aging.* 2019;73:219-228. doi:10.1016/j.neurobiolaging.2018.10.001

89.   Harrison TM, Maass A, Adams JN, Du R, Baker SL, Jagust WJ. Tau deposition is associated with functional isolation of the hippocampus in aging. *Nat Commun.* 2019;10(1). doi:10.1038/s41467-019-12921-z

90.   Berron D, van Westen D, Ossenkoppele R, Strandberg O, Hansson O. Medial temporal lobe connectivity and its associations with cognition in early Alzheimer's disease. *Brain.* 2020;143(4):1233-1248. doi:10.1093/brain/awaa068

91.   Liu CC, Kanekiyo T, Xu H BG. Nat Rev Neurol. *Nat Rev Neurosci.* 2013;9(2):106-118. doi:10.1038/nrneurol.2012.263.Apolipoprotein

92.   Sinha N, Berg CN, Tustison NJ, et al. APOE ε4 status in healthy older African Americans is associated with deficits in pattern separation and hippocampal hyperactivation. *Neurobiol Aging.* 2018;69:221-229. doi:10.1016/j.neurobiolaging.2018.05.023

93.   Dai MH, Zheng H, Zeng LD, Zhang Y. The genes associated with early-onset Alzheimer's disease. *Oncotarget.* 2018;9(19):15132-15143. doi:10.18632/oncotarget.23738

94.   Corder EH, Saunders AM, Strittmatter WJ, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science (80-).* 1993;261(5123):921-923.

95.   Leal SL, Ferguson LA, Harrison TM, Jagust WJ. Development of a mnemonic discrimination task using naturalistic stimuli with applications to aging and preclinical Alzheimer's disease. *Learn Mem.* 2019;26(7):219-228. doi:10.1101/lm.048967.118

96. Leal SL, Ferguson LA, Harrison TM, Jagust WJ. Development of a mnemonic discrimination task using naturalistic stimuli with applications to aging and preclinical Alzheimer ' s disease. Published online 2019:219-229. doi:10.1101/lm.048967.118.26

97. Marks SM, Lockhart SN, Baker SL, Jagust WJ. Tau and β-amyloid are associated with medial temporal lobe structure, function, and memory encoding in normal aging. *J Neurosci*. 2017;37(12):3192-3201. doi:10.1523/JNEUROSCI.3769-16.2017

98. Sperling RA, Donohue MC, Raman R, et al. Association of Factors With Elevated Amyloid Burden in Clinically Normal Older Individuals. 2020;02115(6):735-745. doi:10.1001/jamaneurol.2020.0387

99. Clark R, Tahan AC, Watson PD, Severson J, Cohen NJ, Voss M. Aging affects spatial reconstruction more than spatial pattern separation performance even after extended practice. *Hippocampus*. 2017;27(6):716-725. doi:10.1002/hipo.22727

100. Ngo CT, Newcombe NS, Olson IR. The ontogeny of relational memory and pattern separation. *Dev Sci*. 2018;21(2):1-11. doi:10.1111/desc.12556

101. Hassevoort KM, Khan NA, Hillman CH, Cohen NJ. Differential development of relational memory and pattern separation. *Hippocampus*. 2020;30(3):210-219. doi:10.1002/hipo.23146

102. Berron D, Schütze H, Maass A, et al. Strong Evidence for Pattern Separation in Human Dentate Gyrus. *J Neurosci*. 2016;36(29):7569-7579. doi:10.1523/JNEUROSCI.0518-16.2016

103. Reagh ZM, Yassa MA. Object and spatial mnemonic interference differentially engage lateral and medial entorhinal cortex in humans. *Proc Natl Acad Sci U S A*. 2014;111(40):E4264-E4273. doi:10.1073/pnas.1411250111

104. Bennett IJ, Stark CEL. Mnemonic discrimination relates to perforant path integrity: An ultra-high resolution diffusion tensor imaging study. *Neurobiol Learn Mem*. 2016;129:107-112. doi:10.1016/j.nlm.2015.06.014

105. Nilakantan AS, Bridge DJ, VanHaerents S, Voss JL. Distinguishing the precision of spatial recollection from its success: Evidence from healthy aging and unilateral mesial temporal lobe resection. *Neuropsychologia*. 2018;119(August):101-106. doi:10.1016/j.neuropsychologia.2018.07.035

106. Hodgetts CJ, Shine JP, Williams H, et al. Increased posterior default mode network activity and structural connectivity in young adult APOE-ε4 carriers: a multimodal imaging investigation. *Neurobiol Aging*. 2019;73:82-91. doi:10.1016/j.neurobiolaging.2018.08.026

107. Adams JN, Maass A, Harrison TM, Baker SL, Jagust WJ. Cortical tau deposition follows patterns of entorhinal functional connectivity in aging. *Elife*. 2019;8:1-22. doi:10.7554/eLife.49132

108. Xue C, Yuan B, Yue Y, et al. Distinct Disruptive Patterns of Default Mode Subnetwork Connectivity Across the Spectrum of Preclinical Alzheimer ' s Disease. 2019;11(November):1-11. doi:10.3389/fnagi.2019.00307

109. Lithfous S, Dufour A, Després O. Spatial navigation in normal aging and the prodromal stage of Alzheimer's disease: Insights from imaging and behavioral studies. *Ageing Res Rev*. 2013;12(1):201-213. doi:10.1016/j.arr.2012.04.007

110. Bierbrauer A, Kunz L, Gomes CA, et al. Unmasking selective path integration deficits in Alzheimer's disease risk carriers. *Sci Adv*. 2020;6(35):1-22. doi:10.1126/sciadv.aba1394

111. Levine TF, Allison SL, Stojanovic M, Fagan AM, Morris JC, Head D. Spatial navigation ability predicts progression of dementia symptomatology. *Alzheimer's Dement*. 2020;16(3):491-500. doi:10.1002/alz.12031

112. Hardcastle K, Ganguli S, Giocomo LM. Environmental Boundaries as an Error

Correction Mechanism for Grid Cells. *Neuron*. 2015;86(3):827-839. doi:10.1016/j.neuron.2015.03.039

113. Killian NJ, Jutras MJ, Buffalo EA. A map of visual space in the primate entorhinal cortex. *Nature*. 2012;491(7426):761-764. doi:10.1038/nature11587

114. Nau M, Julian JB, Doeller CF. How the Brain's Navigation System Shapes Our Visual Experience. *Trends Cogn Sci*. 2018;22(9):810-825. doi:10.1016/j.tics.2018.06.008

115. Bicanski A, Burgess N. A Computational Model of Visual Recognition Memory via Grid Cells. *Curr Biol*. 2019;29(6):979-990.e4. doi:10.1016/j.cub.2019.01.077

116. Berteau-Pavy F, Park B, Raber J. Effects of sex and APOE ε4 on object recognition and spatial navigation in the elderly. *Neuroscience*. 2007;147(1):6-17. doi:10.1016/j.neuroscience.2007.03.005

117. Coughlan G, Laczó J, Hort J, Minihane AM, Hornberger M. Spatial navigation deficits — overlooked cognitive marker for preclinical Alzheimer disease? *Nat Rev Neurol*. 2018;14(July):1-11. doi:10.1038/s41582-018-0031-x