Important declarations

Please remove this info from manuscript text if it is also present there.

Associated Data

Required Statements

Competing Interest statement:
The authors declare that they have no competing interests.

Funding statement:
This study was supported by the Czech Science Foundation (Grant no. GJ17-24795Y).

Water stress and nitrogen supply affect floral traits and pollination in the white mustard, *Sinapis alba* (Brassicaceae)

Asma Akter [Corresp., 1, 2], Jan Klečka [2]

[1] Department of Zoology, Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic

[2] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, České Budějovice, Czech Republic

Corresponding Author: Asma Akter
Email address: asma.akter@entu.cas.cz

Changes in environmental conditions are likely to have a complex effect on the growth of plants, their phenology, plant-pollinator interactions, and reproductive success. The current world is facing an ongoing climate change along with other human-induced environmental changes. Most research has focused on the impact of increasing temperature as a major driving force for climate change, but other factors may have important impacts on plant traits and pollination too and these effects may vary between different seasons. In addition, it is likely that the effects of multiple environmental factors, such as increasing temperature, water stress, and nitrogen enrichment are not independent. Therefore, we tested the impact of three key factors – season, water, and nitrogen supply – on plant traits, pollination, and seed production in *Sinapis alba* (Brassicaceae). Plants were grown in three seasons with two levels of water and nitrogen supply in a factorial design. We collected data on multiple vegetative and floral traits and assessed the response of pollinators in the field. Additionally, we evaluated the effect of growing conditions on seed set in plants exposed to pollinators and in hand-pollinated plants. Our results show that water stress impaired vegetative growth, decreased flower production, reduced visitation by pollinators and seed set, while nitrogen availability played an important role in nectar production. Seasonality modulated the effect of water and nitrogen availability on vegetative and floral traits and strongly affected flowering phenology and flower production. We demonstrated that changes in water and nitrogen availability alter plant vegetative and floral traits, which impacts flower visitation and consequently plant reproduction. We conclude that ongoing environmental changes such as altered precipitation regimes and nitrogen enrichment may thus affect plant-pollinator interactions with negative consequences for the reproduction of wild plants and insect-pollinated crops.

Water stress and nitrogen supply affect floral traits and pollination in the

white mustard, *Sinapis alba* (Brassicaceae)

Asma Akter[1,2] & Jan Klečka[1]

[1] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, České Budějovice,

Czech Republic

[2] Department of Zoology, Faculty of Science, University of South Bohemia, České Budějovice,

Czech Republic

Corresponding Author:

Asma Akter

Email address: asma.akter84@gmail.com

Abstract

Changes in environmental conditions are likely to have a complex effect on the growth of plants,

their phenology, plant-pollinator interactions, and reproductive success. The current world is

facing an ongoing climate change along with other human-induced environmental changes. Most

research has focused on the impact of increasing temperature as a major driving force for climate

change, but other factors may have important impacts on plant traits and pollination too and

these effects may vary between different seasons. In addition, it is likely that the effects of

multiple environmental factors, such as increasing temperature, water stress, and nitrogen

enrichment are not independent. Therefore, we tested the impact of three key factors – season,

water, and nitrogen supply – on plant traits, pollination, and seed production in *Sinapis alba*

(Brassicaceae). Plants were grown in three seasons with two levels of water and nitrogen supply

in a factorial design. We collected data on multiple vegetative and floral traits and assessed the

response of pollinators in the field. Additionally, we evaluated the effect of growing conditions

on seed set in plants exposed to pollinators and in hand-pollinated plants. Our results show that

water stress impaired vegetative growth, decreased flower production, reduced visitation by

pollinators and seed set, while nitrogen availability played an important role in nectar production.

Seasonality modulated the effect of water and nitrogen availability on vegetative and floral traits

and strongly affected flowering phenology and flower production. We demonstrated that changes

in water and nitrogen availability alter plant vegetative and floral traits, which impacts flower

visitation and consequently plant reproduction. We conclude that ongoing environmental

changes such as altered precipitation regimes and nitrogen enrichment may thus affect plant-

pollinator interactions with negative consequences for the reproduction of wild plants and insect-

pollinated crops.

Introduction

Ecosystems worldwide are facing accelerating global change characterised by increasing

temperature and changing levels of precipitations, coupled with increasing supply of nitrogen

and other nutrients, biological invasions, and habitat loss (Hoover *et al.,* 2012). Testing the

effects of these environmental changes on plant growth and reproductive fitness are necessary to

understand potential impacts of climate change on the productivity and functioning of natural

and agricultural ecosystems (Rustad, 2008). For pollinator-dependent plants, changes of these

factors may impact their relationship with pollinators and consequently the success of pollination

and plant reproduction (Scaven and Rafferty, 2013; Gérard *et al.,* 2020). Reproductive success of

animal-pollinated plants generally depends on floral traits, which act as an advertisement of

rewards to their pollinators (Hegland and Totland, 2005; Basnett *et al*., 2019). Despite strong

selection from pollinators, plant populations naturally show significant variation in their

morphological, phenological, and floral traits. A part of this variation results from heritable

genetic differences among individuals, while the rest (phenotypic plasticity) is caused by local

environmental factors (Holtsford and Ellstrand, 1992; Gray and Brady, 2016). Changing

environmental factors may thus alter plant-pollinator interactions as a consequence of changing

plant traits (Carroll *et al.,* 2001; Scaven and Rafferty, 2013).

Increasing temperature and water stress can have a major effect on the physiological and

phenological development of plants (Schweiger *et al*., 2010). The global average annual

temperature is rising gradually, with higher increases in the average and minimum temperatures

reported during the winter than the summer months (NOAA, 2021). Consequently, phenological

shifts are visible in many early-flowering plants (Kehrberger and Holzschuh, 2019). Increased

average temperatures may allow them to initiate growth and flowering earlier because of earlier

snowmelt and higher spring temperatures (Fitter and Fitter, 2002; Güsewell *et al.,* 2017).

Water availability is changing in a complex way as many regions of the world are facing

water scarcity and other regions are facing increased precipitation (Christensen *et al.,* 2007).

Although water availability is an important determinant of plant growth, its effect on floral traits

is less clear. Water availability can directly influence the flowering time and duration (Bernal *et*

al., 2011; Lasky *et al.,* 2016) and plants with adequate water supply may achieve greater height

and floral abundance (Galen, 2000; Carromero and Hamrick, 2005), increased nectar production

(Zimmermann and Pyke, 1988, Carroll *et al.,* 2001), and higher nectar sucrose content (Wyatt *et*

al., 1992). Water stress may lead to reduced floral resources (Rering *et al.,* 2020) including

impaired pollen and seed development (Barnabás *et al.,* 2008; Hedhly, 2011; Snider and

Oosterhuis, 2011). Changes in water availability can also affect various stages of phenological

growth differently in the same species, for example, severe water deficiency in maize delays

silking but not anthesis (Blum, 1996). The consequences of water stress for flower visitation by

insect pollinators are still poorly known, but there is evidence that alteration of floral rewards by

water stress may lead to decreased flower visitation (Descamps *et al.,* 2018).

Variation in nitrogen supply is another key driver of local plant diversity (Bobbink *et al.,*

2010). At the global scale, anthropogenic nitrogen deposition increased more than 10 times over

the last century (Galloway *et al.,* 2004; Fowler *et al.,* 2013) and is now around 200 Tg N per year

with a wide range of negative environmental consequences (Battye *et al.,* 2017). In a plant

community, small-scale heterogeneity of soil nitrogen content at the scale of a few meters can

lead to variation in plant size and reproductive success (Scott-Wendt *et al.,* 1988), possibly

including their mutualistic relationship with pollinators. Under the conditions of nitrogen

limitation, increasing nitrogen supply can enhance plant growth and enable plants to produce

floral rewards of higher quality (Gardener and Gillman, 2001; Burkle and Irwin, 2009; Burkle

and Irwin, 2010). In particular, nitrogen enrichment can increase the amount of nectar produced

per flower and alter the concentration and composition of amino acids in nectar, which may

affect pollinator preferences and foraging behaviour (Pyke, 1981; Baker and Baker, 1983; Inouye

and Waller 1984; Gardener and Gillman, 2002; Hoover *et al.,* 2012). For example, several

studies on fruit and vegetable plant species showed that pollinator preferences for their flowers

strongly depended on nectar volume or sugar concentration (Jabłoński *et al.,* 1984; Schmidt *et al.,*

2015; Roldán-Serrano and Guerra-Sanz, 2005). A study on *Citrus* plants also showed that

honeybee visitation rate was positively correlated with nectar volume (Albrigo *et al.,* 2012).

These investigations indicated that both the quantity and quality of nectar influences flower

preferences of pollinators.

Changing environmental conditions can alter plant traits and disrupt interactions of plants

with pollinators, but the consequences for pollination and seed production remain poorly known.

Given the pace of ongoing climate change, which alters not only the temperature but also water

availability for plants, and still increasing anthropogenic nitrogen deposition, it is important to

investigate how these factors act interactively. Such interactive effects have been rarely

considered in experimental studies on plant-pollinator interactions (Hoover *at al.*, 2012). To fill

this gap, we examined the interactive impact of water and nitrogen supply on vegetative and

floral traits, pollination, and seed production in *Sinapis alba* across three different seasons. It is

an economically important crop, cultivated over a wide geographic range for oil and fodder and

is partly self-incompatible with seed production strongly dependent on pollination by insects. In

our study we aimed to answer the following questions: 1) What are the interactive effects of

water and nitrogen supply on vegetative and floral traits of *S. alba?* 2) Are these effects

consistent in different seasons? 3) How does intraspecific trait variation caused by growing

conditions affect flower visitation by pollinators, pollination efficacy, and seed production?

Materials & Methods

The experimental plant, *Sinapis alba*

Sinapis alba (white mustard) is a rapidly growing annual plant from the Brassicaceae family with

a short vegetation period. This crop is widely cultivated for seeds, oil, fodder, or as a catch crop.

Flowers are yellow, produced in an elongated raceme, have four petals, four sepals, and 6

stamens, of which four are long and two are short. Fruit is a pod with usually four seeds but can

have up to eight seeds (Jauzein, 2011). A wide range of pollinating insects visit this plant but the

European honey bee (*Apis mellifera*), bumble bees and solitary bees are the main pollinators in

Europe (Flacher *et al.,* 2020).

Growing *S. alba* under variable conditions in the greenhouse

This experiment was conducted in the greenhouse, where *S. alba* seedlings were grown in the

winter (30 plants per treatment), spring (60 plants per treatment), and summer (45 plants per

treatment) in 2017-2018. Growing conditions varied among the three seasons in temperature and

sunlight. The temperature in the greenhouse fluctuated in a near-natural way and was on average

21°C in the winter 2017-2018, 25°C in the spring 2017, and 29°C in the summer 2018. A

minimum 10 hours of daylight (natural daylight + artificial light) was maintained for all seasons.

Plants grown in the winter received 10-12 hours of daylight, plants grown in the spring received

12-16 hours, and plants grown in the summer received 16 hours of daylight. The plants were

grown in a combination of garden soil: compost soil: sand = 2:2:1 by volume. Seeds were

germinated in germination trays and seedlings were transferred to individual pots (11 x 11 x 11

cm) four days after germination.

We divided the seedlings from each season into four treatment groups which received a

different combination of two watering regimes and two levels of nitrogen supply. Based on the

average ca. 700 mm annual precipitation in the Czech Republic (source: Czech

Hydrometeorological Institute, https://www.chmi.cz/), we determined the lower level of water

for one pot as 20 ml and the higher level of water as 40 ml per day. As the minimum N

application recommended for the *S. alba* is 12 kg N ha^{-1} with a $\leq$ 330 mm annual precipitation

(Brown *et al.,* 2005) and the N deposition is 100 kg N ha^{-1} for Europe for two years, we applied

NPK fertiliser, corresponding to 0.242 g N pot^{-1} ($\sim$ 100kg N ha^{-1}) as higher level of application

and 0.121 g N pot-1 ($\sim$50 kg N pot^{-1}) as lower level, once a week for 8 weeks according to

Vitousek *et al.,* 1997. Before conducting the main experiment, we performed a preliminary trial

to determine the described water level, N$_2$ application, and soil mixture to optimize the growth

conditions for *S. alba*. The position of plants in the greenhouse was altered regularly to avoid any

possible impact of environmental gradients, e.g. the light level, within the greenhouse. Plants

infected with diseases or attacked by aphids were discarded from the experiment.

Assessment of plant morphology, phenology and nectar production

We measured several vegetative and flora traits of individual plants in each treatment group.

Plant height and leaf number were measured several times throughout the growing period and

final height was taken after the end of flowering. The number of leaves was counted for the main

shoot only. Stem diameter of each plant was measured 20 cm above ground. The number of days

until the onset of flowering was counted from the day of the seedling transfer to the opening of

first flower and the total number of flowers bloomed were counted until the end of flowering.

Nectar was collected from 4 flowers per plant in each treatment group after one day of flowering

by using calibrated 0.5 µl capillary tubes (Drummond Microcaps®), which allowed us to

measure the volume of nectar. Additional data were collected on the plant fresh weight and dry

weight of plants grown in the spring to determine the effect of different combinations of water

and nitrogen availability on plant biomass. Overall, we collected data on plant height, the

number of leaves, and stem diameter from 11-20 plants, sampled nectar from 15-25 plants,

measured the onset of flowering in 15-30 plants, and counted the total number of flowers in 12-

15 plants per treatment and season. All data collected during the experiments are deposited in

Figshare (https://doi.org/10.6084/m9.figshare.13317686).

Pollination efficacy treatment and field pollination observations

To determine the impact of water and nitrogen supply on the pollination efficacy in *S. alba*, we

carried out self- and cross-pollination in 6 plants per treatment grown in the winter. We marked

flowers selected for the experiment, performed cross-pollination manually using pollen from a

different plant, and recorded fruit and seed production after the completion of seed development.

Plants grown in the spring and the summer were brought outside and placed in a sunny location

nearby the Institute to assess the pollinator response and natural pollination efficacy under field

conditions. First part of the pollinator observations was carried out in the spring 2017 (May 17 –

May 31) and a second part in the summer 2018 (July 26 – July 31). We always exposed four

plants at the same time (one plant from each of the four water x nitrogen combinations), placed 1

meter apart in a square configuration. We observed their visitation by naturally occurring

pollinators for 30 minutes, from 9:00 to 14:00 each day. Altogether, we carried out observations

of forty-four groups of four plants, which resulted into a total 22 hours of observation.

Pollinators were observed, collected, and identified in the field. After the end of each observation,

the plants were brought back to the greenhouse, their open flowers were marked, and seed

production through the natural pollination was measured after fruiting (data:

https://doi.org/10.6084/m9.figshare.13317686).

Statistical analyses

We used generalised linear models (GLM) to assess the individual and interactive impact of
water, N_2, and season on the plant vegetative and floral traits. The availability of water, nitrogen,
and the season were used as factors in the analyses. Depending on the type of the response
variable, we specified the GLM with either Gaussian error distribution, overdispersed Poisson
("quasipossion", the number of flowers), or Gamma distribution with a log link function (onset
of flowering and nectar volume). We analysed data from the outdoor flower visitation
experiment using GLM with water, nitrogen level, and season as factors, using the overdispersed
Poisson ("quasipossion") error distribution. We analogously analysed also data on seed set of
plants depending on growing conditions. We always examined the distribution of residuals to
verify that the models fitted the data well. We conducted all analyses in R Version 3.6.3 (R Core
Team 2020). Most plots were created using GraphPad Prism (Version 6.01, for Windows,
GraphPad Software, San Diego, California USA, www.graphpad.com).

Results

Vegetative traits

We observed a complex response of the selected vegetative traits of *S. alba* to differences in the
growing conditions (Table 1, Fig.1). Plant height was affected by the three-way interaction of
water availability, nitrogen supply, and season, i.e. the effect of each variable was dependent on
the values of the other two variables ($F = 4.94$, $P = 0.008$, Fig. 1A). While higher water
availability made the plants taller, increasing nitrogen availability made the plants shorter. The
magnitude of these effects varied across the three seasons (Fig. 1A). In addition, plants grown in
the summer were almost 50% shorter than those grown in the winter. Stem diameter was larger
in plants grown under high water availability ($F = 40.84$, $P < 0.001$, Fig. 1B) and varied also
depending on nitrogen supply in interaction with the season ($F = 9.51$, $P < 0.001$) – higher
nitrogen supplies increased stem diameter in plants grown in the winter and in the summer, but
not in the spring (Fig. 1B). On the other hand, the number of leaves was affected only to a
limited degree by water availability ($F = 16.20$, $P < 0.001$, Fig. 1C) and the season ($F = 7.48$, $P <$

0.001), while nitrogen supply had no measurable effect ($F = 0.09$, $P = 0.768$). Finally, dry weight
of the plants grown under high water availability was 2.6 times higher compared to those grown
under low water availability, with a positive effect of nitrogen supply only at high water
availability (water x nitrogen interaction: $F = 5.52$, $P = 0.023$, Fig. 1D).

Phenology and flower resource variation

The onset of flowering was delayed by ca. 21 days in the winter compared to the spring and
summer ($F = 69.45$, $P < 0.001$) and slightly delayed also by high nitrogen supply ($F = 6.21$, $P =$
0.013, Table 2, Fig. 2A). The number of flowers produced over the plants' flowering period was
affected positively by water ($F = 24.24$, $P < 0.001$) and slightly negatively by nitrogen supply (F
$= 6.83$, $P = 0.010$), with a week evidence for their interactive effect ($F = 3.60$, $P = 0.060$), and
varied among the three seasons (Table 2, Fig. 2B). Finally, nectar volume showed a complex
dependence on the interaction of water, nitrogen supply, and season ($F = 3.56$, $P = 0.030$).
Higher water availability increased nectar volume in the winter, but not in the spring and
summer, while higher nitrogen availability increased nectar volume under low water availability
in both spring and summer (Fig. 2C).

The dependence of flower visitation on growing conditions
We observed flower visitation by eight major types of flower-visiting insects in the spring and
summer which we distinguished as: the honeybee (*Apis mellifera*), solitary bees, wasps,
bumblebees, rapeseed beetles (*Brassicogethes* (=*Meligethes*) sp.), other beetles, hoverflies, and
other flies. Rapeseed beetles were the most abundant flower visitors in the spring, followed by
honeybees, while solitary bees were dominant in the summer, followed by hoverflies (data:
https://doi.org/10.6084/m9.figshare.13317686).
Plants grown with high amount of water were visited more frequently than the plants
grown with low amount of water ($F = 23.57$, $P < 0.001$) and the total number of flower visitors
was higher in the spring than in the summer ($F = 14.19$, $P < 0.001$) (Fig. 3A, 3B). Nitrogen
supply under which the plants were grown did not consistently affect their flower visitation ($F =$
0.26, $P = 0.612$). Flower visitation was also affected by the number of open flowers ($F = 18.92$,
$P < 0.001$, Fig. 3C) and by plant height ($F = 8.89$, $P = 0.003$, Fig. 3D), but the effect of water

availability and season remained significant even after accounting for the variation in flower
number and height (GLM, $F = 5.46$, $P = 0.021$ for the effect of water and $F = 6.58$, $P = 0.011$ for
the effect of the season), i.e. the differences in flower visitation between plants grown under
different conditions could not be explained simply by differences in plant height and flower
number. In addition to differences in total flower visitation, we detected changes in the
composition of the flower visitors observed on plants grown under different water availability
according to a redundancy analysis (RDA) performed separately for observations from the spring
($F = 4.0$, $P = 0.004$) and summer ($F = 3.1$, $P = 0.028$), while nitrogen supply did not affect the
composition of flower visitors ($F = 0.37$, $P = 0.869$ for the spring data and $F = 0.57$, $P = 0.669$
for the summer data). Some flower visitors visited plants grown under high water availability
more frequently, particularly solitary bees and rapeseed beetles, while other flower visitors did
not show a clear preference (Figs. 4 and 5).
258

Pollination efficacy and seed production
Our hand pollination experiment confirmed that *S. alba* is partially self-incompatible. Plants
cross-pollinated by hand using a brush produced ca. 3.9 times higher number of seeds per flower
than the self-pollinated ones; on average 2.9 compared to 0.7 seeds per flower (Fig. 6A).
However, the seed set depended not only on the mode of pollination (self-pollinated compared to
cross-pollinated) but on its interaction with nitrogen availability ($F = 10.64$, $P = 0.002$).
Specifically, higher nitrogen availability increased seed set in self-pollinated plants, but
decreased seed set in cross-pollinated plants. In addition, higher water availability increased seed
set in both self-pollinated and cross-pollinated plants irrespective of the nitrogen level ($F = 5.24$,
$P = 0.028$) (Fig. 6B).
Plants exposed to natural pollination in the spring and summer produced a variable
number of seeds per flower depending on the interaction of water availability and season ($F =$
8.93, $P = 0.004$) and on the nitrogen availability ($F = 5.08$, $P = 0.028$). We observed a slightly
higher number of seeds per flower in plants grown under high water availability in the spring
2017, but lower in the summer 2018. In addition, plants grown under higher nitrogen availability
produced a lower number of seeds per flower (Fig. 6C). As we showed above, plants grown
under different combinations of water and nitrogen availability varied in their total production of
flowers. Combined with the variation in the number of seeds produced per flower, this led to

differences in the total seed set per plant (Fig. 6D). Specifically, total seed set was higher in
plants grown under high water availability, but the effect was stronger in the summer than in the
spring (the interaction between water availability and season: $F = 5.03$, $P = 0.029$).
280
281

Discussion
282

The effect of environmental changes on plant traits
283

Our results highlight that water stress is a key factor for both vegetative and floral traits
(Descamps *et al.,* 2018), while changes of nitrogen supply had a more limited impact in our case.
The effects of these factors were mostly interactive and dependent on the season. Higher amount
of water positively affected plant growth, especially in the spring and summer. Nitrogen
enrichment played a more complex role in the vegetative growth of *S. alba* and its effect was
modulated by the season and water availability. Interestingly, previous research has shown that
while water deficiency may lead to reduced biomass production and diminished nitrogen uptake
in plants (Cossani *et al*., 2012), sufficient nitrogen supply may enhance their drought tolerance
and increase water use efficiency (Quemada and Gabriel, 2016).

Flowering phenology, the number of flowers, and nectar production of *S. alba* were also
affected by growing conditions. In particular, nectar production was affected by a complex
interaction of all three variables, where nectar production increased with higher nitrogen supply
to a variable degree depending on water availability and season. This is a complex relationship,
not reported by previous studies on other plant species. Contrary to our results, several studies
showed that nectar production may decline in response to water reduction and increased
temperature (Keasar *et al*., 2008; Scaven and Rafferty, 2013; Takkis *et al*., 2015). In our case,
nectar production did not decrease under these conditions and also showed an opposite result
compared to Hoover *et al.* (2012), where nectar production of *Cucurbita maxima* decreased with
higher nitrogen supply and increased with increasing temperature. However, our results are in
line with a previous study by of Villarreal and Freeman (1990), who showed that nectar
production of *Ipomopsis longiflora* was positively affected by water supplementation. It seems
that the effects of varying environmental conditions on nectar production are highly species-
specific (Lu *et al*., 2015).

Both vegetative and floral traits displayed significant differences among plants grown in
different seasons under the same water and nitrogen supply levels. In particular, plants grown in
the winter were generally smaller and their onset of flowering was significantly delayed
compared to spring and summer. Generally, the growth rate and reproductive success of plants is
the highest within a certain range of optimal temperatures and decreases rapidly beyond this
optimal range (Vasseur *et al*., 2014, Hatfield and Prueger, 2015). Phenological shifts in many
plants are also closely related to temperature (Jagadish *et al*., 2016; Kehrberger and Holzschuh,
2019). Previous studies on *Borago officinalis* also showed that increasing temperature may
diminish flower production or lead to flower bud abortion, which may reduce the total number of
flowers produced during the plant's flowering period (Saavedra *et al*., 2003; Descamps *et al.,*
2018). Similarly, the total number of flowers of *S. alba* significantly dropped in our experiment
in the summer when the temperature in the greenhouse averaged ca. 29°C. However, the fact that
plants produced more flowers with high amount of water, even during the summer, shows that
the impact of thermal stress on flower production can be reduced by water supplementation (see
also Mahan *et al*., 1995; Li *et al*., 2020). An optimal temperature is also required for the
maximum nectar secretion (Pacini and Nepi, 2007, Lu *et al*., 2015). In our case, *S. alba* produced
comparatively higher amount of nectar in the spring when the temperature reached intermediate
values (average ca. 25°C). Apart from the temperature, seasonal differences in vegetative and
floral traits in our plants could have been driven also by differences in day length and light
intensity.
327

Impact on pollination and seed production
Our experiments showed that differences in traits among plants grown under different
environmental conditions had a cascading effect on the number and identity of flower visitors of
S. alba and on its reproduction. Flower visitation rate of insect-pollinated plants depends on
visual cues indicating high floral reward such as the number of open flowers (Conner and Rush,
1996; Akter *et al.,* 2017) and the size of floral display (Grindeland *et al*., 2005; Parachnowitsch
et al., 2010; Biella *et al*., 2019), and on the amount and quality of nectar and pollen (Cresswell,
1999; Grindeland *et al*., 2005). Other morphological features can also influence plant detection
by potential pollinators, such as plant height (Junker *et al*., 2013; Klecka *et al.,* 2018a;
Hernández-Villa *et al*., 2020), local plant clustering (Elliot and Irwin, 2009; Akter *et al.,* 2017),

and flower colour (Reverté *et al.,* 2016). Measurements of flower visitation with plants grown in

the spring 2017 and plants grown in the summer 2018 revealed that and in both cases plants

grown with higher amount of water had a significantly higher number of flower visitors

compared to plants grown under low amount of water irrespective of nitrogen supply. This is

likely a consequence of differences in vegetative and floral traits induced by differences in water

availability. As discussed above, plants grown with high amount of water were taller and

produced more flowers and these characteristics had a positive effect on the visitation of

individual plants as reported in other plant species (e.g. Mitchell *et al.* 2004; Akter *et al.,* 2017;

Klecka *et al.,* 2018a). However, other modifications of plant traits induced by water stress also

apparently decreased the visitation of plants grown with low amount of water, because the effect

of water availability on the number of flower visitors per plant persisted even after accounting

for differences in the number of open flowers and plant height in our analysis. We believe that

the remaining unexplained variation could be related to nectar chemistry (Petanidou *et al.,* 2006;

Hoover *et al.,* 2012) or flower scent (Farré-Armengol *et al.*, 2020).

Besides having lower flower visitation, plants grown under low amount of water had

different composition of flower visitors compared to plants grown with higher amount of water.

In the spring 2017, plants grown with high amount of water received more frequent visits from

rapeseed beetles, solitary bees, and hoverflies than plants grown under low amount of water,

while the other flower visitors, including honeybees and bumblebees, did not discriminate among

the plants. The results were similar in the summer 2018, although rapeseed beetles were almost

absent. The observed differences in the flower visitation in the spring 2017 and the summer 2018

may be influenced by the differences in overall insect abundance or weather, but not necessarily

by the growing conditions of the plants. For instance, increasing temperature may affect flower

visitation by a number of mechanisms, from differences in plant traits caused by high

temperature stress (Descamps *et al.,* 2018), through phenological shifts of plant flowering and

pollinator emergence (Hegland *et al.,* 2009; Bartomeus *et al.,* 2011), to changes in pollinator

foraging activity caused by their responses to temperature (Corbet *et al.,* 1993; Slamova *et al.*

2011), and direct and indirect effects of temperature on the fitness and mortality of pollinating

insects (Scaven and Rafferty, 2013).

Finally, seed production of *S. alba* was also affected by water and nitrogen availability,

apparently both directly through physiological mechanisms and indirectly through changes in

insect pollination. Our hand pollination assessment confirmed that *S. alba* is a partially self-

incompatible plant (Fan *et al.*, 2007). Low water availability reduced seed production per flower

in both self-pollinated and cross-pollinated plants, which is consistent with previous studies

suggesting that water stress may lead to seed or pod abortion (e.g. New *et al.*, 1994; Behboudian

et al., 2001). However, we also observed an intriguing effect of nitrogen availability on seed set:

increased nitrogen availability increased seed set in self-pollinated plants, but decreased seed set

in cross-pollinated plants. We are not aware of any studies which would show that high nitrogen

supply can cause seed abortion.

Seed count per flower from the naturally pollinated plants in the spring 2017 also showed

a similar trend as in plants cross-pollinated by hand, where the number of seeds per flower

increased in plants grown with high water availability, but decreased with high nitrogen

availability. In contrast, in experiments done in the summer 2018, the number of seeds per flower

was not affected by nitrogen availability and decreased in plants grown with high amount of

water. Total seed set per plant was unaffected by nitrogen availability and increased in plants

grown under high water availability – moderately in the spring 2017 but much more in the

summer 2018. This may stem from differences in the composition of the flower visitor

community between plants grown under low and high water availability and from the higher total

visitation rate in the spring 2017 compared to summer 2018. The level of pollen limitation

(Knight *et al.*, 2005) was thus higher in the summer 2018, which likely explains why the number

of seeds per flower was lower and was more strongly reduced in plants grown with low amount

of water.

Conclusions

We have shown that multiple drivers of environmental change have a complex and interactive

impact on plant traits, visitation by pollinators, and seed production. Our model species, *S. alba*,

is an important crop and a close relative to many other economically important crops and

vegetables from the Brassicaceae family, hence our experiment shows how different climatic

drivers may affect both vegetative growth and crop yield in plants form this family in the future

extreme climatic events. We conclude that not only increasing temperature, but also reduced

precipitation and nitrogen enrichment, may impact plant-pollinator interactions with negative

consequences for the reproduction of wild plants as well as insect-pollinated crops.

Acknowledgements

We are grateful to Pavla Koloušková for her valuable help during the field observations and Jana

Jersáková for her help with the laboratory analyses.

References

Akter A, Biella P, Klecka J. 2017. Effects of small-scale clustering of flowers on pollinator
  foraging behaviour and flower visitation rate. *PLoS ONE* 12(11): e0187976. doi:
  01.1371/journal.pone.0187976

Albrigo LG, Russ R, Rouseff R, Bazemore RA. 2012. Some citrus flower characteristics and
  honeybee preference. *Proceedings of the Florida State Horticultural Society* 125: 112–118.

Baker HG, Baker I. 1983. Floral nectar sugar constituents in relation to pollinator type. In: Jones
  C. E., Little R. J. (eds.) Handbook of experimental pollination biology. Van Nostrand
  Reinhold, New York, 117–141.

Barnabás B, Jäger K, Fehér A. 2008. The effect of drought and heat stress on reproductive
  processes in cereals. *Plant Cell and Environment* 31(1):11-38.

Bartomeus I, Ascher JS, Wagner D, Danforth BN, Colla S, Kornbluth S, Winfree R. 2011.
  Climate-associated phenological advances in bee pollinators and bee-pollinated plants.
  Proceedings of the National Academy of Sciences 108 (51): 20645-20649.

Basnett S, Ganesan R, Devy SM. 2019. Floral traits determine pollinator visitation in
  Rhododendron species across an elevation gradient in the Sikkim Himalaya. *Alp Botany* 129:
  81–94.

Battye, W., Aneja, V. P., & Schlesinger, W. H. (2017). Is nitrogen the next carbon? Earth's
  Future, 5(9), 894-904.

Behboudian MH, Ma Q, Turner NC, Palta JA. 2001. Reactions of chickpea to water stress: yield
  and seed composition. *Journal of the Science of Food and Agriculture* 81(13): 1288-1291.

Bernal M, Estiarte M, Peñuelas J. 2011. Drought advances spring growth phenology of the
  Mediterranean shrub *Erica multiflora*. *Plant Biology* 13: 252-257.

Biella P, Akter A, Ollerton J, Tarrant S, Janeček Š, Jersáková J, Klecka J. 2019. Experimental
loss of generalist plants reveals alterations in plant-pollinator interactions and a constrained
flexibility of foraging. *Scientific Reports* 9: 7376.

Brown, J., J.B. Davis, and A. Esser. 2005. Pacific Northwest condiment yellow mustard (Sinapis
alba L.) grower guide. Subcontract report - National Renewable Energy Laboratory.
University of Idaho. [Online]. Available at http://www.nrel.gov/docs/fy05osti/36307.pdf.

Blum A. 1996. Crop responses to drought and the interpretation of adaptation. *Plant Growth*
Regul 20: 135–148.

Bobbink R, Hicks K, Galloway J. et al. 2010. Global assessment of nitrogen deposition effects
on terrestrial plant diversity: a synthesis. *Ecological Applications* 20(1): 30-59.

Burkle L, Irwin RE. 2009. The importance of interannual variation and bottom–up nitrogen
enrichment for plant–pollinator networks. *Oikos* 118: 1816-1829.

Burkle LA, Irwin RE. 2010. Beyond biomass: measuring the effects of community-level nitrogen
enrichment on floral traits, pollinator visitation and plant reproduction. *Journal of*
Ecology 98: 705– 717.

Carroll AB, Pallardy SG, Galen C. 2001. Drought stress, plant water status, and floral trait
expression in fireweed, *Epilobium angustifolium* (Onagraceae). *American Journal of*
Botany 88: 438– 446.

Carromero W, Hamrick JL. 2005. The mating system of *Verbascum thapsus* (Scrophulariaceae):
the effect of plant height. *International Journal of Plant Sciences* 166: 979– 983.

Christensen JH, Carter TR, Rummukainen M, Amanatidis G. 2007. Evaluating the performance
and utility of regional climate models: the PRUDENCE project. *Climatic Change* 81: 1–6.

Conner JK, Rush S. 1996. Effects of flower size and number on pollinator visitation to wild
radish, *Raphanus raphanistrum*. *Oecologia, 105(4), 509-516.*

Corbet SA, Fussell M, Ake R, Fraser A, Gunson C, Savage A, Smith K. 1993. Temperature and
the pollinating activity of social bees. *Ecological Entomology* 18: 17-30.

Cossani CM, Slafer GA, Savin R. 2012. Nitrogen and water use efficiencies of wheat and barley
under a Mediterranean environment in Catalonia. *Field Crops Research* 128: 109-118.

Cresswell JE. 1999. The influence of nectar and pollen availability on pollen transfer by
individual flowers of oil seed rape (*Brassica napus*) when pollinated by bumblebees
(*Bombus lapidarius*). *Journal of Ecology* 87(4): 670–677.

Descamps C, Quinet M, Baijot A, Jacquemart AL. 2018. Temperature and water stress affect
plant-pollinator interactions in *Borago officinalis* (Boraginaceae). *Ecology and Evolution* 8:
3443-3456.

Elliott SE, Irwin RE. 2009. Effects of flowering plant density on pollinator visitation, pollen
receipt, and seed production in *Delphinium barbeyi* (Ranunculaceae). *American Journal of*
Botany 96: 912-919.

Farré-Armengol G, Fernández-Martínez M, Filella I, Junker RR, Peñuelas J. 2020. Deciphering
the biotic and climatic factors that influence floral scents: A systematic review of floral
volatile emissions. *Frontiers in Plant Science* 11: 1154.

Fitter AH, Fitter RSR. 2002. Rapid changes in flowering time in British plants. *Science* 296:
1689–1691.

Flacher F, Raynaud X, Hansart A, Geslin B, Motard B, Verstraet S, Bataille M, Dajoz I. 2020.
Below-ground competition alters attractiveness of an insect-pollinated plant to
pollinators. *AoB PLANTS* 12(4): plaa022. doi:10.1093/aobpla/plaa022

Fowler, D. *et al.* 2013The global nitrogen cycle in the twenty-first century. *Philos. Trans. R. Soc.*
B 368, 20130164.

Galen C. 2000. High and dry: drought stress, sex-allocation trade-offs, and selection on flower
size in the alpine wildflower *Polemonium viscosum* (Polemoniaceae). *American*
Naturalist 156: 72– 83.

Galloway JN *et al.* 2004. Nitrogen cycles: past, present, and future. *Biogeochemistry* 70: 153–
226.

Gardener MC, Gillman MP. 2001. The effects of soil fertilizer on amino acids in the floral nectar
of corncockle, *Agrostemma githago* (Caryophyllaceae). *Oikos* 92:101–106.

Gardener MC, Gillman MP. 2002. The taste of nectar-a neglected area of pollination ecology.
Oikos 98: 552-557.

Gérard M, Vanderplanck M, Wood T, Michez D. 2020. Global warming and plant–pollinator
mismatches. *Emerging Topics in Life Sciences* 4 (1): 77–86.

Gray SB, Brady SM. 2016. Plant developmental responses to climate change. *Arid Soil Research*
and Rehabilitation 2: 49-58.

Grindeland JM, Sletvold N, Ims, RA. 2005. Effects of floral display size and plant density on
pollinator visitation rate in a natural population of *Digitalis purpurea*. *Functional Ecology*
19: 383-390.

Güsewell S, Furrer R, Gehrig R, Pietragalla B. 2017. Changes in temperature sensitivity of
spring phenology with recent climate warming in Switzerland are related to shifts of the
preseason. *Global Change Biol*ogy 23: 5189– 5202.

Hatfield JL, Prueger JH. 2015. Temperature extremes: Effect on plant growth and development.
Weather and Climate Extremes 10 (A): 4-10.

Hedhly A. 2011. Sensitivity of flowering plant gametophytes to temperature
fluctuations. *Environmental and Experimental Botany* 74: 9– 16.

Hegland SJ, Totland Ø. 2005. Relationships between species' floral traits and pollinator
visitation in a temperate grassland. *Oecologia* 145: 586–594.

Hegland SJ, Nielsen A, Lázaro A, Bjerknes AL, Totland Ø. 2009. How does climate warming
affect plant-pollinator interactions? *Ecology Letters* 12(2): 184– 195.

Hernández-Villa V, Vibrans H, Uscanga-Mortera E, Aguirre-Jaimes A. 2020. Floral visitors and
pollinator dependence are related to floral display size and plant height in native weeds of
central Mexico. *Flora* 262: 151505.

Hoover SER, Ladley JJ, Shchepetkina AA, Tisch M, Gieseg SP, Tylianakis JM. 2012. Warming,
CO_2, and nitrogen deposition interactively affect a plant-pollinator mutualism. *Ecology*
Letters 15: 227-234.

Holtsford TP, Ellstrand NC. 1992. Genetic and environmental variation in floral traits affecting
outcrossing rate in *Clarkia tembloriensis* (Onagraceae). *Evolution* 46: 216–225.

Fan HL, Sun WC, Yan N, Zhu HX, Wu JY, Zhang YH, Zeng J, Ye J, Liu YL. 2007. Analysis of
self-compatibility in *Sinapis alba* (L.) Boiss. In Proceedings of the 12th International
Rapeseed Congress, Wuhan, China, 26–30 March 2007, 368–369.

Inouye DW, Waller GD. 1984. Responses of honeybees (*Apis mellifera*) to amino acid solutions
mimicking nectars. *Ecology* 65: 618– 625.

Jabłoński B, Król S, Pliszka K, Żurowska Z. 1984. Nectar secretion and pollination of the
blueberry (*Vaccinium corymbosum* L.). *Acta Horticulture* 165: 133–144.

Jagadish SK, Bahuguna RN, Djanaguiraman M, Gamuyao R, Prasad PV, Craufurd
PQ. 2016. Implications of high temperature and elevated CO_2 on flowering time in
plants. *Frontiers in Plant Science* 7: 913.
Jauzein P. 2011. *Flore des champs cultivés*. 2nd edn. Versailles, France: Editions Quae.
Junker RR, Blüthgen N, Brehm T, Binkenstein J, Paulus J, Schaefer MH, Stang M. 2013.
Specialization on traits as basis for the niche-breadth of flower visitors and as structuring
mechanism of ecological networks. *Functional Ecology* 27(2): 329-341.
Keasar T, Sadeh A, Shmida A. 2008. Variability in nectar production and standing crop, and
their relation to pollinator visits in a Mediterranean shrub. *Arthropod-Plant Interactions*
2: 117–123.
Kehrberger S, Holzschuh A. 2019. Warmer temperatures advance flowering in a spring plant
more strongly than emergence of two solitary spring bee species. *PLoS ONE* 14(6):
e0218824.
Klecka J, Hadrava J, Koloušková P. 2018a. Vertical stratification of plant–pollinator interactions
in a temperate grassland. *PeerJ* 6: e4998. doi:10.7717peerj.4998
Knight, T. M., Steets, J. A., Vamosi, J. C., Mazer, S. J., Burd, M., Campbell, D. R., ... & Ashman,
T. L. (2005). Pollen limitation of plant reproduction: pattern and process. *Annual Review of*
Ecology, Evolution and Systematics, 36, 467-497.
Lasky JR, Uriarte M, Muscarella R. 2016. Synchrony, compensatory dynamics, and the
functional trait basis of phenological diversity in a tropical dry forest tree community:
effects of rainfall seasonality. *Environmental Research Letters* 11: 115003.
Li Y, Guan K, Peng B, Franz TE, Wardlow B, Pan M. 2020. Quantifying irrigation cooling
benefits to maize yield in the US Midwest. *Glob Change Biology* 26: 3065– 3078.
Lu NN, Li XH, Li L, Zhao ZG. 2015. Variation of nectar production in relation to plant
characteristics in protandrous *Aconitum gymnandrum*. *Journal of Plant Ecology* 8(2): 122–
129.
Mahan JR, McMicheal BL, Wanjura DF. 1995. Methods for reducing the adverse effects of
temperature stress on plants: A review. *Environmental and Experimental Botany* 35: 251–
258.
Mitchell RJ, Karron JD, Holmquist KG, Bell JM. 2004. The influence of *Mimulus ringens* floral
display size on pollinator visitation patterns. *Functional Ecology* 18: 116-124.

New B, Duthion C, Turc O. 1994. Phenological response of pea to water stress during
reproductive development. *Crop science* 34(1): 141-146.

NOAA National Centres for Environmental Information, State of the Climate: Global Climate
report for the Annual 2020, published online January 2021, retrieved on September 13, 2021
from https://www.ncdc.noaa.gov/sotc/global/202013

Pacini E, Nepi M. 2007. Nectar production and presentation. In: Nicolson S.W., Nepi M.,
Pacini E. (eds) Nectaries and Nectar. Springer, Dordrecht.

Parachnowitsch AL, Kessler A. 2010. Pollinators exert natural selection on flower size and floral
display in *Penstemon digitalis*. *New Phytologist* 188(2): 393-402.

Petanidou T, Van Laere A, Ellis WN, Smets E. 2006. What shapes amino acid and sugar
composition in Mediterranean floral nectars? *Oikos* 115(1): 155–169.

Pyke GH. 1984. Optimal Foraging Theory: A Critical Review. *Annual Review of Ecology and*
Systematics 15(1): 523-575.

Quemada M, Gabriel JL. 2016. Approaches for increasing nitrogen and water use efficiency
simultaneously. *Global Food Security* 9: 29-35.

R Core Team 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/.

Rering, CC, Franco, JG, Yeater, KM, Mallinger, RE. 2020. Drought stress alters floral volatiles
and reduces floral rewards, pollinator activity, and seed set in a global
plant. *Ecosphere* 11(9): e03254.

Reverté S, Retana J, Gómez JM, Bosch J. 2016. Pollinators show flower colour preferences but
flowers with similar colours do not attract similar pollinators. *Annals of Botany* 118(2): 249-
257.

Roldán-Serrano AS, Guerra-Sanz JM. 2005. Reward attractions of zucchini flowers (*Cucurbita*
pepo L.) to bumblebees (*Bombus terrestris* L.). *European Journal of Horticultural*
Science 70: 23–28.

Rustad LE. 2008. The response of terrestrial ecosystems to global climate change: towards an
integrated approach. *Science of the Total Environment* 404: 222–235.

Saavedra F, Inouye DW, Price MV, Harte J. 2003. Changes in flowering and abundance
of *Delphinium nuttallianum* (Ranunculaceae) in response to a subalpine climate warming
experiment. *Global Change Biology* 9: 885-894.

Scaven VL, Rafferty NE. 2013. Physiological effects of climate warming on flowering plants

and insect pollinators and potential consequences for their interactions. *Current Zoology*

59(3): 418-426.

Schmidt K, Filep R, Orosz-Kovács Z, Farkas A. 2015. Patterns of nectar and pollen presentation

influence the attractiveness of four raspberry and blackberry cultivars to pollinators. *The*

Journal of Horticultural Science and Biotechnology 90: 47–56.

Schweiger O, Biesmeijer JC, Bommarco R, Hickler T, Hulme PE. 2010. Multiple stressors on

biotic interactions: How climate change and alien species interact to affect

pollination. *Biological Reviews* 85: 777–795.

Scott-Wendt J, Chase RG, Hossner LR. 1988. Soil chemical variability in sandy Ustalfs in semi-

arid Niger, West Africa. *Soil Science* (June): 414.

Slamova I, Klecka J, Konvicka M. 2011. Diurnal behavior and habitat preferences of *Erebia*

aethiops, an aberrant lowland species of a mountain butterfly clade. *Journal of Insect*

Behavior, *24*(3), 230-246.

Snider JL, Oosterhuis DM. 2011. How does timing, duration, and severity of heat stress

influence pollen-pistil interactions in angiosperms? *Plant Signaling & Behavior* 6(7): 930–

933.

Takkis K, Tscheulin T, Tsalkatis P, Petanidou T. 2015. Climate change reduces nectar secretion

in two common Mediterranean plants. *AoB Plants* 7: 111.

Vasseur DA, DeLong JP, Gilbert B, Greig HS, Harley CDG, McCann KS, Savage V, Tunney TD,

O'Connor MI. 2014. Increased temperature variation poses a greater risk to species than

climate warming. *Proceedings of the Royal Society B: Biological Sciences* 281(1779): 2013-

2612.

Villarreal AG, Freeman CE. 1990. Effects of temperature and water stress on some floral nectar

characteristics in *Ipomopsis longiflora* (Polemoniaceae) under controlled conditions.

*Botanical Gazette*151: 5-9.

Vitousek, P.M., Aber, J.D., Howarth, R.W., Likens, G.E., Matson, P.A., Schindler,

W. *et al.* (1997). Human alteration of the global nitrogen cycle- sources and

consequences. *Ecol. Appl.*, **7**, 737– 750.

Wyatt R, Broyles SD, Derda GS. 1992. Environmental influences on nectar production in

milkweeds (Asclepias syriaca and A. exaltata). *American Journal of Botany* 79: 636-642.

612 Zimmermann M, Pyke GH. 1988. Reproduction in *Polemonium*: assessing the factors limiting

613   seed set. *American Naturalist* 131: 723-738.

Figure 1

The impact of water and nitrogen supply on vegetative traits of *S. alba* across three seasons.

Four vegetative traits were measured: A) plant height, B) plant diameter, C) the number of leaves, and D) dry weight. The boxplots show the median and interquartile range. The results of statistical tests are summarised in Table 1.

Figure 2

The impact of water and nitrogen supply on floral traits of *S. alba* across three seasons.

A) the onset of flowering (the day of the opening of the first flower), B) the number of flowers produced per plant, and C) nectar production (median and interquartile range is shown). The results of statistical tests are summarised in Table 2.

Figure 3

Flower visitation of plants grown under varying water and nitrogen availability.

The number of flower visitors per plant per 30 minutes during two observation periods are shown: A) spring 2017 and B) summer 2018 (median and interquartile range is shown). Flower visitation also varied depending on the number of open flowers (C) and plant height (D).

Figure 4

The composition of the flower visitor community varied depending on water availability under which the plants were grown.

The results of RDA show the effect of water availability and the lack of an effect of nitrogen supply during the plant growth period on the composition of the flower visitor community in the spring 2017 (A) and the summer 2018 (B).

Figure 5

Flower visitation by major flower visitor groups.

A) plants grown in the spring 2017, and B) in summer 2018. The number of visitors per 30
min. is shown (mean ± SE).

Figure 6

Seed production of *S. alba* grown under different growing conditions.

Seed production in plants subjected to self-pollination and cross-pollination by hand: A) seed production per flower and B) seed production per pod (median and interquartile range is shown). Seed production of plants subjected to natural pollination in the spring 2017 and the summer 2018: C) the number of seeds per flower, and D) the total seed set per plant (median and interquartile range is shown).

Table 1(on next page)

The effects of water availability, nitrogen supply, and season on selected vegetative traits of *S. alba*. *F* and *P* values for individual variables and their interactions estimated by generalised linear models (see Methods) are shown.

Variable	Plant height		Stem diameter		Number of leaves		Dry weight	
	F	*P*	*F*	*P*	*F*	*P*	*F*	*P*
Water	**23.2**	**<0.001**	**32.9**	**<0.001**	**8.8**	**0.004**	**105.72**	**<0.001**
Nitrogen	**22.3**	**<0.001**	**22.3**	**<0.001**	0.1	0.78768	0.22	0.643
Season	**60.7**	**<0.001**	**41.9**	**<0.001**	**9.0**	**<0.001**	-	-
Water x Nitrogen	2.1	0.147	2.1	0.154	1.9	0.167	**5.52**	**0.023**
Water x Season	2.2	0.113	**3.1**	**0.047**	**3.3**	**0.039**	-	-
Nitrogen x Season	0.457	0.636	**14.4**	**<0.001**	0.2	0.817	-	-
Water x Nitrogen x Season	**4.94**	**0.008**	1.2	0.306	0.1	0.951	-	-

Table 2(on next page)

The effects of water availability, nitrogen supply, and season on floral traits of *S. alba*. *F* and *P* values for individual variables and their interactions estimated by generalised linear models (see Methods) are shown.

Variable	Day of first flower		Number of flowers		Nectar volume	
	F	*P*	*F*	*P*	*F*	*P*
Water	**11.6**	**<0.001**	**11.6**	**<0.001**	**4.9**	**0.028**
Nitrogen	**8.4**	**0.004**	**8.4**	**0.004**	0.2	0.622
Season	**25.1**	**<0.001**	**25.1**	**<0.001**	**10.74**	**<0.001**
Water x Nitrogen	**6.1**	**0.015**	**6.1**	**0.015**	**21.95**	**<0.001**
Water x Season	**3.5**	**0.033**	**3.5**	**0.033**	1.2	0.318
Nitrogen x Season	1.3	0.265	1.3	0.265	**8.8**	**<0.001**
Water x Nitrogen x Season	0.01	0.950	1.3	0.288	**3.6**	**0.030**