**Title:** Urban fox squirrels exhibit tolerance to humans but respond to stimuli from natural

predators

**Running title:** Fox squirrel behavior in urban areas

**Authors:** Anna Kittendorf[1], Ben Dantzer[1,2]

[1]Department of Psychology, University of Michigan, Ann Arbor, MI, 48109, USA

[2]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI,

48109, USA

**Correspondence:** Ben Dantzer (dantzer@umich.edu)

**Acknowledgements**

This research was a part of a senior honor's thesis conducted by AK in the Department of

Psychology at the University of Michigan. It was funded by the University of Michigan. Thanks

to Dantzer Lab members (especially Amy-Charlotte Devitz) for input on this project. We

acknowledge that the University of Michigan and some of the other properties that we worked on

during this project reside on the traditional Territories of the Three Fire Peoples – the Ojibwe,

Odawa, and Bodewadmi.

**Abstract**

Animals in urban areas that experience frequent exposure to humans often behave differently

than those in less urban areas, such as less vigilance or anti-predator behavior. These behavioral

shifts may be an adaptive response to urbanization but it may be costly if animals in urban areas

also exhibit reduced anti-predator behavior in the presence of natural predators. We tested the

hypothesis that increased tolerance to humans in urban populations of fox squirrels (*Sciurus*

*niger*) reduces their responses to predators (hawks and domestic dogs). In trials with only a

human observer as the stimulus, urban squirrels exhibited reduced vigilance and anti-predator

behavior compared to those in less urban areas but there was little evidence that urbanization

altered the correlations among the different behaviors. Next, we exposed squirrels in multiple

urban and less urban sites to acoustic playbacks of a control stimulus (non-predatory bird calls),

a natural predator (hawk), and dogs and recorded their vigilance and three different anti-predator

behaviors when a human approached them while either broadcasting one of these three

playbacks or no playbacks at all. Urban squirrels exhibited increased vigilance and anti-predator

behavior when exposed to a human paired with hawk playbacks compared to the control

playbacks. This indicates that urban squirrels did perceive and assess risk to the natural predator

appropriately despite exhibiting increased tolerance to humans. These results provide little

support for the hypothesis that increased tolerance to humans causes animals to lose their fear of

natural predators.

**Keywords**: Anti-predator behavior, cross-habituation, first alert distance, flight initiation

distance, stimulus generalization, urbanization

**Introduction**

Behavior plays an important role in enabling animals to persist through environmental

change (Baldwin, 1896; Bartholomew, 1964; West Eberhard, 1989; Price et al., 2003; Snell-

Rood, 2013). Accordingly, it seems to play a major role in facilitating the ability of animals to

cope with new challenges that they face in urban environments (Ditchkoff et al., 2006;

Tuomainen and Candolin, 2011; Lowry et al., 2013; Ryan and Partan, 2014). One of the most

common behavioral adjustments of animals in urban environments is reduced expression of anti-

predator behavior in the presence of humans. This is often reflected in measures of flight

initiation distance (FID), which is the distance at which an animal flees from an approaching

human and is thought to be an approximation of their sensitivity to risk of an approaching

predator (Cooke, 1980; Ydenberg & Dill, 1986; Lima and Dill, 1990). Individuals with shorter

FIDs are considered to be bolder than individuals with longer FIDs since they demonstrate

reduced fear of the "predator".

Substantial evidence supports the hypothesis that animals in more urban environments

exhibit less anti-predator behavior, as reflected by a lower FID. For example, a meta-analysis of

180 bird species, 16 lizard species, and 16 mammal species, Samia et al. (2015) showed that

populations of these species that experienced elevated levels of human disturbance exhibited

lower FID. This could be because anti-predator behaviors carry energetic or time costs as

vigilance and escape behaviors take time away for resource acquisition (Ydenberg and Dill,

1986; Cooper and Frederick, 2007; Møller, 2012) and animals in urban environments may

optimize resource acquisition by exhibiting lower levels of anti-predator behavior.

Although reductions in the expression of anti-predator in urban environments is generally

thought to be adaptive (Møller, 2008; Carrette et al., 2016), there may be potential costs for

urban animals if they reduce their overall expression of anti-predator behavior to not only

humans but also natural predators. This could be due to the phenomenon of cross-habituation or

stimulus-generalization. For example, birds that are habituated to a threatening stimulus that are

then presented with a second simulated predator exhibit an attenuated response to this second

stimulus compared to a group of naïve birds (Hinde, 1954; see also Curio, 1993). This type of

stimulus generalization can occur where an animal habituated to one stimulus exhibits an

attenuated response to a second stimulus from the same *or* different sensory modality (Guttman

and Kalish, 1956; Thompson and Spencer, 1966; Rankin et al., 2009). Related concepts occur in

the context of "behavioral spillover" where individuals that exhibit high levels of a behavior in

one context also exhibit it in another context even though it may not be adaptive, such as animals

exhibiting higher levels of boldness in a courtship context also exhibiting higher boldness in the

presence of a predator (Arnqvist and Henriksson, 1997; Sih et al., 2004).

If urban animals also exhibit reductions in anti-predator behavior to natural predators, it

could conceivably have important impacts on wildlife populations by increasing their

vulnerability to predators (Geffroy et al., 2015). To date, there is little consensus about whether

animals in urban areas or those exposed to increased human presence exhibit a reduced response

to threats from natural predators (Fitzgerald and Stronza, 2016). For example, some studies show

that individuals in areas with higher human activity exhibit less of a behavioral response when

natural predators were observed visiting the area (Olson and Acevedo-Gutiérrez, 2017) or due to

acoustic playbacks of a natural predator (McCleery, 2009). The latter suggests animals

experiencing frequent exposure to human activity exhibit reduced responses to other stimuli from

natural predators. Other studies show the response of animals in more urban areas to a stimulus

from a natural predator is not attenuated compared to those in more rural locations (Labra and

Leonard, 1999; Coleman et al., 2008; Seress et al., 2011; Bokony et al., 2012; Cavalli et al.,

2016; Weaver et al., 2018; Vincze et al., 2019).

In this study, we characterized the vigilance and anti-predator of fox squirrels (*Sciurus*

*niger*) in urban and less urban areas to achieve the following three objectives. First, we

conducted standard FID trials (with only stimuli from a human observer) to examine if squirrels

in urban areas showed reduced vigilance and anti-predator behavior towards a human observer

compared to those in less urban areas. If squirrels in urban areas did exhibit reduced vigilance

and anti-predator behavior, this would support the hypothesis that squirrels in our urban study

populations were more tolerant to human presence, which would be consistent with numerous

other studies (Samia et al., 2015). Squirrels were located in their natural habitat and we recorded

the following four aspects of their vigilance and anti-predator behavior. First, we recorded the

distance to which the observer could get to before they exhibited vigilance behavior towards the

observer ("first alert distance" or FAD, similar to Fernández-Juricic and Schroeder, 2003;

Blumstein et al., 2005). Second, we recorded how close the observer could get to them before

they ran away (FID). Third, was the probability that the squirrel escape by running up a tree.

Fourth, the latency following the trial it took them to resume their typical behavior (foraging or

traveling off tree).

Our second objective was to assess the degree of phenotypic correlations among the

different behaviors we recorded from these trials. We did this because previous studies suggest

that phenotypic correlations among different vigilance or anti-predator behaviors (i.e., the

behavioral syndrome) may differ between urban and rural populations (Evans et al., 2010; Scales

et  al., 2011; Carrette and Tella, 2017; Lapiedra et al., 2017). This could be because selection

against anti-predator behavior in urban environments produces a correlated response in other

risk-taking behaviors. If there was a strong phenotypic correlation between two behaviors within

the same squirrel recorded from less urban sites compared to those in urban sites, this would

support the hypothesis that squirrels in urban areas exhibit greater behavioral flexibility.

Our final objective was to examine if urban animals exhibit reduced behavioral responses

to stimuli from natural predators when they are in the presence of humans. To do so, we

quantified the four behaviors described above when fox squirrels in urban or less urban areas

were presented with a human observer with a control acoustic playback (common non-

threatening bird), a human observer paired with the playback of a natural predator (hawk), or a

human observer paired with a playback of an invasive predator (dog). We predicted that squirrels

in the urban areas but not those in the less urban areas would exhibit no change in vigilance and

anti-predator behavior when they were exposed to the human+dog or human+hawk stimuli

compared to the human+control playback. This would support the hypothesis that animals in

urban environments exhibit less vigilance and anti-predator not only to humans but also when

faced with natural predators.

127                                        **Materials and Methods**

*Study species and sites*

Fox squirrels are ubiquitous in urban and suburban environments in the midwestern

United States (McCleery, 2008, 2009). Although arboreal tree squirrel species like fox squirrels

are common in urban areas worldwide, they continue to experience predation from natural

predators, although it may be rare (McCleery et al., 2008), and from domestic cats and dogs

(Koprowski, 1994; Wauters et al., 1997; Tumlison, 2012; Loss et al., 2013; Jokimäki et al.,

2017).

We studied natural populations of adult fox squirrels from six sites in and around Ann

Arbor, Michigan (Table 1). Sites were chosen based upon estimates of human population density

(Center for International Earth Science Information Network, 2018) with urban sites having

higher human density than less urban sites (see below and Table 1). Urban sites included

Prospect Park as well as two locations on the University of Michigan's (UM) main campus

(North and Central Campus) that are ~3-4 km away from one another. Prospect Park is near

downtown Ypsilanti, Michigan and about 13 km away from UM main campus. Less urban sites

included Nichols Arboretum, County Farm Park, and Saginaw Forest. Nichols Arboretum is

located ~1 km away UM main campus, County Farm Park is about ~4.5 km away, and Saginaw

Forest is ~7 km away. At all research sites, dogs are allowed but hunting is not. Approval to

conduct this research at each site was obtained from UM (Central Campus, Nichols Arboretum,

North Campus, Saginaw Forest), Washtenaw County Parks & Recreation (County Farm Park),

and the City of Ypsilanti (Prospect Park). All procedures were non-invasive and involved

behavioral observation or short-term exposure to playbacks of acoustic stimuli. All experiments

followed the guidelines set by the Animal Behavior Society/Association for the Study of Animal

Behaviour (Anonymous, 2012) and the US National Research Council and were approved by the

UM Institutional Animal Care and Use Committee (protocol # PRO00009076). We note that

squirrels used in the study may be STRANGE (sensu Webster and Rutz, 2020) in the sense that

individual squirrels likely have different rearing histories (though they are unknown and none

should have been reared in captivity) and that compliance to take part in the study was likely

biased towards squirrels that did not immediately run away when approached by the human

observer.

Given that increased exposure to humans may cause animals in urban areas to exhibit less

anti-predator behavior towards them (McCleery, 2009; Rodriguez-Prieto et al., 2009; Vincze et

al., 2016; Uchida et al., 2019), we focused on human presence as the major factor difference

between our study sites (which should also reflect general urbanization). Sites were classified as

"urban" based upon having a human population density >1000 persons per km$^2$ whereas the less

urban sites had anywhere from 25-250 persons per km$^2$ (Saginaw) to 250-1000 persons per km$^2$

(County Farm Park, Nichols Arboretum). To support these classifications, we human and dog

presence while we were visiting sites conducting our behavioral observations. We counted the

total number of dogs (on or off leash) but only counted the number of humans up to 50. If human

presence exceeded 50 people, then a rough estimate of 50, 75, or 100 was recorded. Human

presence was recorded as 100 for all numbers estimated to be >100. Although human and dog

presence varied among the different sites (Table 1), the number of humans observed per hour of

observation averaged over all the urban sites (mean ± SE = 7.23 ± 3.33 humans/hr) was higher

than those observed averaged over all the less urban sites (0.51 ± 0.40: Mann-Whitney-Wilcoxon

Test, W = 1, $p = 0.1$). We observed fewer dogs per hour at the urban sites (0.065 ± 0.06 dogs/hr)

compared to the less urban sites (0.33 ± 0.10, Mann-Whitney-Wilcoxon Test, W = 9, $p = 0.1$).

These differences in humans or dogs observer per hour were not significant but in general

support our assumption that our urban sites likely experience greater exposure to humans.

Adult squirrels and sex were visually identified by size (small juvenile squirrels were

excluded) and anatomy (males were identified by presence of testes), respectively. Similar to

most studies that measure anti-predator using FID, trials were conducted on unmarked squirrels

at each site. Squirrels were located by walking around each site and trials were started when

squirrels were observed. Focal individuals were selected randomly, however, only squirrels that

were feeding or foraging on the ground were included in this experiment. Because we did not

mark squirrels individually, it is possible that the same squirrel was observed on different days,

although we visited different areas of each study site to try and reduce this possibility. It is

unlikely that the same squirrel was observed multiple times on the same day, because after each

trial was completed, the observer traveled approximately 20 meters away from the previous

location and started a trial with a different squirrel. Additionally, sites were only revisited after at

least three days since the previous visit to reduce the possibility of a squirrel becoming

habituated to the trials should it be sampled again. Although we cannot address habituation in

this study, if squirrels at these sites were habituating to our protocols, we would expect that their

behavioral responses would decline with trial number or date when the trial was conducted. The

fact that none of our behavioral variables were associated with date of when the trial was

conducted (Tables 2-4) supports our assumption that squirrels were not habituating to our

protocols.

*Quantifying behavioral responses of squirrels*

In total, we observed fox squirrels for 52.36 hours over 30 different days. A single

observer (AK) conducted all trials. Sites were visited between 800 and 1900 h EST and data

were collected from October 2019 through January 2020. Trials with no playbacks were

conducted from 25 October 2019 to 16 November 2019 (from 812 to 1810 h) whereas trials

using playbacks (playback trials) were conducted from 18 November 2019 to 27 January 2020

(from 802 to 1609 h). Both the order in which sites were visited and the order in which playback

versions across playback treatments (see below) were randomized. No two sites were visited on

the same day. All trials with no playbacks were conducted prior to the playback trials in this

study. This was due to personnel limitations and prevents us from directly comparing trials with

and without playbacks given that squirrel behavior likely changes seasonally from October to

January due to food caching behavior in autumn but not winter. Trials were not conducted when

it was raining or snowing. Results from two one-way ANOVAs showed that the time of day for

playback trials did not vary among the three different treatment groups ($F_{2,91} = 0.44$, $p = 0.65$)

and that the time of day when baseline trials were conducted did not vary among squirrels at the

urban or less urban sites (t-test: $t_{75.4} = 0.21$, $p = 0.83$). Air temperature varied during all the trials

varied from -6.1° to 10.5°C (mean = 2.7° C).

We measured the behavioral responses of squirrels to humans or humans plus different

playbacks using protocols developed for tree squirrels (Dill and Houtman, 1989; Gustafson and

VanDruff, 1990; McCleery, 2009). At the beginning of each trial, a marker was placed at the

starting position of the observer and trial data were recorded (GPS location, time of day,

temperature, general weather conditions, and squirrel sex). The squirrel was approached by a

single observer (AK) at a slow and steady pace in a direct line to the squirrel (see Fig. S1 in

Appendix I). Additional markers were placed when a squirrel displayed the first alert and when

they fled. The FAD was defined as the distance between the observer and the squirrel when it

first stopped moving (froze) and looked at the observer with one or both eyes. FID was recorded

as the distance between the squirrel's initial position and the observer location when it actively

fled (stopped feeding and foraging and moved rapidly away from observer). A marker was also

placed at an estimate of the squirrel's initial position to the place where they fled to if refuge was

not taken in a tree, and the distance between this marker and the squirrel's initial position was

recorded as "flight distance". We recorded this because some studies have noted that FID is

variable depending on intruder starting distances and distance to a refuge (Dill & Houtman,

1989; Blumstein, 2003). Consequently, the distance between the observer and the focal squirrel

at the start of the trial (starting positions, hereafter referred to as "initial distance") and the

distance between the focal squirrel's initial position and the nearest tree were also measured

("distance to nearest tree").

If the focal animal took refuge in a tree other than the one nearest, distance between the

squirrel's initial position and its refuge tree of choice ("distance to the chosen tree") was also

recorded. If an individual took refuge in a tree, a laser rangefinder was used to measure how high

they climbed, and this distance was recorded (same as vertical escape distance in Uchida et al.,

2017). Lastly, latency to resume behavior was recorded ("latency"). This was measured with a

stopwatch to determine how long it took for the animal to cease alert/vigilance behavior and

resume typical activity (foraging or traveling off tree). When the human observer was recording

latency, they maintained as large a distance as possible from the tree to reduce the influence on

the squirrel's behavior. Out of all the trials, nearly all squirrels ceased alert behavior and

resumed typical behavior within a couple minutes (n = 171 trials, mean ± SE = 130.5 ± 10.3 s),

but there was one individual squirrel that remained alert for longer than ten minutes and we

recorded its latency as ten minutes. Distances were measured with a tape measure and presented

in meters.

*Playback trials*

Playback trials were conducted using the same protocol shown above, with the addition

of an acoustic stimulus being broadcasted while the observer approached the squirrels. The

control stimulus consisted of recordings of black-capped chickadee calls (*Poecile atricapillus*).

Black-capped chickadees are not known to be predators of fox squirrels (Korschgen, 1981;

Koprowski, 1994) and a previous study in another tree squirrel species showed that individuals

exhibited a significantly reduced response to black-capped chickadee playbacks compared to

calls of other anthropogenic noises (car alarm, buzzer) and playbacks of red-tailed hawks (Bohls

and Koehnle, 2017). We therefore expected that black-capped chickadee recordings would

represent a neutral vocalization for fox squirrels and they can act as control to ensure that any

differences in squirrel behavior across playback treatments are attributable to the vocalization

information of the playback rather than an added exposure to noise. To simulate the threat of a

terrestrial predator, recordings of domestic dogs barking were broadcasted. Domestic dogs are

terrestrial predators of fox squirrels (Koprowski, 1994; Wauters et al., 1997) and other species of

tree squirrels that live in the same habitats as fox squirrels also adjust their risk-taking behavior

according to the abundance of domestic dogs (Bowers and Breland, 1996; Cooper et al., 2008).

For the avian predator, recordings of red-tailed hawk (*Buteo jamaicensis*) calls were broadcasted

to the focal individual. Red-tailed hawks were chosen since they are year-round predators of fox

squirrels in Michigan (Koprowski, 1994; personal observations) and other studies illustrate that

tree squirrels respond to hawk playbacks with increased anti-predator behavior (McCleery, 2009;

Lilly et al., 2019). No post-processing of sound files was performed.

Playbacks of the recordings were broadcast to individuals at the start of the trial and

when the observer began the approach and suspended when the squirrel took flight.

Vocalizations were broadcast through a JAMBOX speaker (Jawbone, San Francisco, CA)

connected to an Apple iPhone 6s (Mountain View, CA) with a constant volume set for the

speaker and phone. The amplitude of the playbacks measured from 1 m away from the speaker

was variable among the chickadee (67-80 dB), dog (69-77 dB), and hawk (78-86 dB) playbacks

(measured using a BAFX Sound Level Meter, BAFX3370). We note that the initial starting

distance of the playbacks was inherently variable as we could not standardize the distance

between the observer and squirrel when the trials were started. Consequently, the actual realized

sound level of the playbacks experienced by a squirrel varied. Given how the trials were

conducted in real time (not video recorded), the single observer (AK) was not blind to the

playback treatments or locations of where the experiments took place. All vocalization

recordings were found online (Control A: Place, 2015; Control B: Floyd, 2017a; Control C:

Floyd, 2017b; Dog A: Simion, 2016; Dog B: Simion, 2018; Dog C: Simion, 2017; Hawk A:

Chartier, 2008; Hawk B: Addison, 2017; Hawk C: Wilson, 2010). Each playback treatment

(control, dog, or hawk) had three separate recordings/exemplars (A, B, or C). We tested whether

there was any exemplar effects in separate ANOVAs that included playback exemplar (A, B, C),

playback treatment (chickadee, dog, hawk), and an interaction between the two. We did this for

each of our four behavioral response variables and did not find any significant interactions

between playback exemplar and treatment ($p = 0.12$-$0.99$), suggesting that the version of the

playback treatment did not influence the behavioral response.

*Statistical analyses*

We analyzed the data from trials with and without playbacks separately because the two

experiments were not conducted synchronously and seasonal changes from Fall to Winter in

Michigan may alter squirrel behavior. In trials without playbacks, we used three separate linear

mixed-effects models (LMMs) to examine the effects of urbanization on FAD, FID, and latency

to resume activity following the trial. Although the linear distance a squirrel climbed up a tree

(from base of tree to location of squirrel) has been used in other studies of tree squirrels (e.g.,

Uchida et al., 2017), the distance a squirrel climbed up a tree in our study exhibited a Poisson

distribution where many squirrels did not climb up a tree at all and a few climbed up very high.

Consequently, a generalized linear mixed-effect model (GLMM) with binomial errors was used

to examine the effects of urbanization on the probability that squirrels climbed a tree to escape

during the trial. Models included site category of data collection (urban, less urban), distance to

the nearest tree, sex, Julian date of the trial, and initial distance of the observer as fixed effects.

Distance to the nearest tree was included not only because previous studies show it can impact

anti-predator behavior (measured using FID: Dill & Houtman, 1989; Blumstein, 2003) but also

because it helps control for any differences in vegetation among the different study sites, which

could impact their behavior. The same model structure was used in separate LMMs or the

GLMM for data from the playback trials to examine the effects of the acoustic playback

manipulations on the four squirrel behaviors described above but the models included an

interaction between playback treatment (control, dog, hawk) and site category (urban, less

urban). We assessed the statistical significance of the latter interaction between playback

treatment and site using post-hoc Tukey's Honest Significant Differences that were corrected for

multiple comparisons.

To assess the degree of association between FAD or FID and the two other behavioral

variables (probability of climbing a tree, latency to resume typical behavior), we ran two separate

LMMs (one for FAD and the other for FID). Each model included fixed effects for site category,

probability of climbing a tree, latency to resume typical behavior, and interactions between site

category and each of the two behavioral variables. We also report the Pearson' or point-biserial

(escaping up tree was binary variable) correlations between these variables. We did not run a

similar model looking at the association between FAD and FID given that they are clearly not

independent of one another (for all squirrels [n = 171] across all treatments in this study: Pearson

$r = 0.88$, 95% CI = 0.84-0.91).

Continuous predictor variables were standardized to a mean of 0 and SD of 1. We

confirmed model diagnostics visually and all models met the appropriate assumptions (normality

of residuals, constant variance, no high leverage observations). Latency to resume behavior was

log+1 transformed (base e) to improve homoscedasticity and normality. There were also no

predictor variables that were found to be colinear as all variance inflation factors (VIFs) were

less than 3.68 (Zuur et al., 2010), though the higher VIF were due to interaction terms and VIF of

variables not in interactions were <1.5. Because we had repeated samples from the same site, we

also included a random intercept for site in all of our models. All analyses were conducted in R

version 4.02 (R Core Team, 2020) with lme4 (version 1.1.23, Bates et al., 2015) and p-values

estimated using lmerTest (version 3.1.2, Kuznetsova et al., 2017). Tukey's post-hoc tests were

used to evaluate if the responses to the playback treatments differed between squirrels in urban

and less urban areas using emmeans (1.5.2-1: Lenth, 2020) and *p* values from these analyses

were adjusted for multiple comparisons. Mean and SE are presented below.

330                                            **Results**

331    *Behavioral responses to human-stimuli only*

332            Overall, urban squirrels (n = 38, 20 females and 18 males) exhibited greater tolerance to

humans as they allowed a human observer to get closer to them before they exhibited vigilance

(FAD) or fled (FID) and tended to be less likely to climb a tree during the trial and more quickly

return to typical behavior following the trial compared to those in less urban sites (n = 39, 19

females and 20 males; Table 2, Fig. 1). FID in the squirrels at the urban site (6.36 ± 0.52 m) was

97.2% shorter compared to those at the less urban sites (12.54 ± 0.62 m, *p* = 0.043, Table 2, Fig.

1B). Although the average FAD for squirrels observed at the urban site (10.09 ± 0.74 m) was

43.8% shorter than for those at the less urban sites (14.51 ± 0.69 m), this difference were not

significant (*p* = 0.10, Table 2, Fig. 1A). Squirrels at the urban sites were less likely to climb a

tree while the observer approached (34.2% of trials) compared to those at the less urban sites

(48.6%), although this difference was not significant ($p = 0.093$, Table 2, Fig. 1C). Latency to

resume behavior following the trial was shorter for urban squirrels ($43.9 \pm 14.4$ s) compared to

those at the less urban sites ($157.51 \pm 25.35$ s), but this difference was not significant ($p = 0.11$,

Table 2, Fig. 1D). Trials where the observer started the trial at a longer initial distance to the

squirrel had significantly higher FAD and FID but not probability of climbing a tree or latency to

resume behavior following the trial (Table 2). There were no significant effects of sex, Julian

date, or distance to the nearest tree on FAD, FID, probability of climbing a tree, or latency

(Table 2).

*Patterns of associations among behaviors*

351            Squirrels at urban sites ($n = 38$) that exhibited a higher FAD also exhibited a significantly

higher latency to resume typical behavior following the trial ($r = 0.59$, 95% CI $= 0.33$-$0.77$, $t_{22.9}$

$= 3.07$, $p = 0.005$) but this association between these two variables was significantly reduced in

squirrels at the less urban sites ($n = 39$, $r = 0.09$, 95% CI $= -0.23$-$0.39$), as indicated by the nearly

significant interaction between site category and latency ($t_{33.1} = -2.01$, $p = 0.053$, Table 3, Fig.

2A). Squirrels at both urban sites that exhibited a higher FID also tended to exhibit a higher

latency to resume typical behavior following the trial ($r = 0.53$, 95% CI $= 0.26$-$0.73$, $t_{68.2} = 1.86$,

$p = 0.066$), but the association between these two behaviors was slightly lessened in squirrels at

less urban sites ($r = 0.18$, 95% CI $= -0.14$-$0.47$), although this was not significant ($t_{70.5} = -1.17$, $p$

$= 0.24$, Table 3, Fig. 2B). In squirrels at both urban ($r = 0.46$, 95% CI $= 0.17$-$0.68$) and less

urban sites ($r = 0.39$, 95% CI $= 0.09$-$0.63$), squirrels that were more likely to escape the human

observer by climbing a tree also had a longer latency to resume typical behavior following the

trial ($t_{72.9} = 2.08$, $p = 0.041$) and this did not depend upon whether the squirrels were at urban or

less urban sites ($t_{72.3} = 0.05$, $p = 0.96$).

*Behavioral responses to stimuli from natural predators*

366            The effects of the playbacks on FAD depended upon whether the squirrels were located

at the urban or less urban sites (site x playback treatment: $F_{2, 81.7} = 6.07$, $p = 0.003$), with urban

squirrels exhibiting the highest FAD when exposed to hawk playbacks and squirrels at less urban

sites exhibiting higher overall FAD and no response to the playback treatments (Fig. 3A).

Average FAD for urban squirrels exposed to the hawk vocalizations (n = 12 trials, 17.17 ± 1.33

m) was 37% longer than squirrels who were exposed to the control playback (n = 11, 12.56 ±

0.94 m, Tukey's $p = 0.0016$) and 20% longer than those exposed to the dog playbacks (n = 20,

14.30 ± 0.83 m, Tukey's $p = 0.0025$, Fig. 3A). By contrast, the FAD of squirrels at the less urban

sites were just higher overall (Fig. 3A) and the FAD of those who were exposed to the hawk

playbacks (n = 16, 14.62 ± 1.14 m) did not differ from those exposed to the control playback (n

= 15, 14.46 ± 0.88 m, Tukey's $p = 0.95$) or dog playbacks (n = 20, 14.85 ± 1.05 m, Tukey's $p =$

0.64, Fig. 3A). There were no significant differences in FAD for squirrels exposed to dog

playbacks and those exposed to the control playback for squirrels at urban sites (Tukey's $p =$

0.83) or those at less urban sites (Tukey's $p = 0.84$).

380            Similar to FAD, the effects of the playbacks on FID also depended upon whether the

squirrels were located at the urban or less urban sites (site x playback treatment: $F_{2, 81.2} = 6.28$, $p$

= 0.003), with urban squirrels exhibiting the highest FID when exposed to hawk playbacks and

squirrels at less urban sites exhibiting the highest FID in response to the dog playbacks (Fig. 3B).

Average FID for urban squirrels exposed to the hawk vocalizations (n = 12 trials, 14.95 ± 0.98

m) was 44.3% longer than squirrels who were exposed to the control playbacks (n = 11, 10.37 ±

3.13 m, Tukey's $p = 0.0006$) and 29.9% longer than those exposed to the dog playbacks (n = 20,

11.86 ± 2.65 m, Tukey's $p = 0.0005$, Fig. 3A). In squirrels at urban sites, there was no difference

in FID between those exposed to the control playback and those exposed to the dog vocalizations

(Tukey's $p = 0.90$). In squirrels at less urban sites, FID for those exposed to the dog playbacks

(13.60 ± 0.98 m,) was 12.2% higher than those exposed to the control playback (12.12 ± 0.88 m,

Tukey's $p = 0.049$), but was similar to those exposed to the FID of those exposed to hawk

playbacks (13.11 ± 0.96 m, Tukey's $p = 0.66$, Fig. 3B). Unlike urban squirrels, the FID of those

at the less urban sites who were exposed to hawk playbacks was similar compared to those

exposed to the control playback (Tukey's $p = 0.34$).

The playback treatments had no significant effects on the probability that squirrels in

urban or less urban sites exhibited an escape response by climbing a tree, as indicated by the lack

of interactions between playback treatment and site (Table 4, Fig. 3C). Squirrels at both the

urban and less urban sites exhibited the same response when it comes to their latency to resume

their typical behavior following the trial (hawk > dog > control, Table 4, Fig. 3D), as indicated

by the lack of interaction between site and playback treatment (site x playback treatment: $F_{2, 80.5}$

$= 0.20, p = 0.82$). Squirrels at urban sites that were exposed to the hawk playbacks took 436%

longer to resume their pre-trial behavior (213.33 ± 28.9 s) compared to those who were exposed

to the control playback (39.82 ± 7.70 s, Tukey's $p < 0.0001$) and 147% longer than those

exposed to dog playbacks (86.25 ± 13.30 s, Tukey's $p = 0.011$). Squirrels at the urban sites also

took 114% longer to resume typical behavior if they were exposed to dog playbacks compared to

those exposed to the control playback (Tukey's $p < 0.001$). Similarly, squirrels at the less urban

sites that were exposed to the hawk playbacks took 218% longer to resume their pre-trial

behavior (291.25 ± 35.45 s) compared to those who were exposed to the control playback (91.47

± 25.45 s, Tukey's $p < 0.0001$) and 55.7% longer than those exposed to dog playbacks (187.05 ±

21.38 s, Tukey's $p = 0.045$, Fig. 3D). Squirrels at the less urban sites also took 95.6% longer to

resume typical behavior if they were exposed to dog playbacks compared to those exposed to the

control playback (Tukey's $p < 0.001$).

There was no effect of sex or Julian date of trial, on any of the behaviors (Table 4). There

was no effect of the initial distance that a squirrel was from a tree when the trial started on FAD,

FID, or probability to climb a tree, but, squirrels were less likely to climb a tree if they were

closer to one when the trial started (Table 4). Trials that started with the human observer a

greater distance away from the squirrel had higher FAD, FID, and latency to resume typical

behavior, but not the probability to climb a tree (Table 4).

420                                        **Discussion**

Overall, squirrels at urban sites in the no playback trials exhibited a significantly shorter

FID compared to those at the less urban sites but there minimal differences in the associations

between the behaviors in squirrels at urban versus less urban sites. In the trials with playbacks,

squirrels at the urban sites exhibited higher FAD (hawk > control > dog) and FID (hawk >

control = dog) when exposed to hawk playbacks compared to control or dog playbacks. By

contrast, squirrels at the less urban sites had higher overall FAD and FID than those at urban

sites regardless of playback treatment, there was no effect of hawk or dog playbacks on FAD

(hawk = control = dog) and they exhibited the highest FID when exposed to dog playbacks (dog

> control = hawk). Squirrels at both urban and less urban sites were not more likely to climb a

tree following playbacks from possible predators (hawk = dog = control) but both urban and less

urban squirrels exhibited a longer latency to resume typical behavior following the hawk or dog

playbacks compared to the control (hawk > dog > control). In terms of the STRANGEness of our

results (Webster and Rutz, 2020), our results may be generalizable to other squirrel populations

or different species but we note that our results are biased towards squirrels that voluntarily

participated in the trials (i.e., did not run away when approached).

Similar to most other studies in terrestrial animals (Samia et al., 2015) and in studies in

tree squirrels (McCleery, 2009; Engelhardt and Weladji, 2011; Sarno et al., 2015; Uchida et al.,

2020), our results from trials with no playbacks suggest that squirrels in urban sites were more

tolerant of humans. Specifically, squirrels in urban areas exhibited a shorter FAD and FID, lower

probability to climb a tree to escape the human observer, and a shorter latency to resume typical

behavior following the trial, although only FID was significantly different. The congruency of

our results with previous studies strongly supports this assumption that squirrels at our urban

sites were more tolerant to humans. This is surprising given that these are presumably sympatric

populations with large amount of gene flow among them as the linear distance between some

urban and less urban sites is ~1 km. Unless selection favoring reductions in anti-predator

behavior is extremely strong in urban areas or features of urban landscapes strongly impede gene

flow (Johnson and Munshi-South, 2017), it seems likely that these behavioral differences are

driven by plasticity given that the likely exchange of individuals between suburban and urban

sites prevents local genetic adaptation to these different sites (see discussion in Sol et al., 2013).

It is also possible that these behavioral differences are due to personality-dependent colonization

of urban habitats (Carrete and Tella, 2010; Sprau and Dingemanse, 2017), but we cannot

distinguish among these possibilities at this time.

Urban environments may favor individuals with reduced anti-predator behavior (as this

may increase resource acquisition) and this could cause some correlated shift in other behaviors.

There was little evidence that the correlations among the different behaviors we recorded

differed between squirrels in urban and less urban sites. We found that the association between

latency to resume typical behavior following a trial and FAD differed between urban and less

urban populations. Squirrels in urban but not less urban sites that allowed humans to get closer to

them before exhibiting first vigilance (FAD) also took a significantly shorter amount of time to

resume typical behavior following the end of the trial. However, there was no significant

deviance in the association between FID and latency or probability to escape by climbing a tree

and latency in squirrels from urban and less urban sites. As such, our results somewhat differ

from previous studies (Evans et al., 2010; Scales et al., 2011; Bokony et al., 2012; Myers and

Hyman, 2016; Carrete and Tella, 2017) that indicate that urban environments may enhance or

lessen the associations between different behaviors within the same individual. At this point we

cannot infer whether the shift in phenotypic correlations we documented here (for FAD and

latency) is adaptive, evidence of a constraint or correlated response to selection, or merely due to

environmental co-variance.

Although squirrels at our urban sites were more tolerant of humans, they still exhibited a

robust behavioral response to acoustic stimuli from natural predators. Specifically, they exhibited

increased vigilance (FAD) and anti-predator behavior (FID), latency to resume typical behavior

after the trial) when exposed to the playbacks of a natural predator (hawk) compared to the

control playback or the dog playbacks. These results indicate that urban squirrels do still pay

attention to predation risk and can discriminate and respond accordingly by becoming vigilant

and fleeing when the human is at a greater distance if the human is also paired with hawk

playbacks. By contrast, squirrels at less urban sites did not exhibit differences in FAD when

exposed to the different playbacks, perhaps due to some ceiling effect given that FAD of

squirrels at less urban sites was much higher than FAD of squirrels at urban sites. Our results

therefore reject the hypothesis that urban squirrels are less responsive to natural predators due to

increased tolerance to humans. Previous studies (see also Labra and Leonard, 1999; Seress et al.,

2011; Cooper et al., 2008; Bokony et al., 2012; Cavalli et al., 2016; Weaver et al., 2018; Vincze

et al. 2019) together with our results support that animals in urban habitats or those frequently

exposed to humans, even if more tolerant to human presence, still exhibit increases in anti-

predator behavior in response to a non-human predator.

485            We expected that squirrels would respond to the dog playbacks in similar way to how

they responded to the hawk playbacks as both are stimuli from potential predators. Previous

studies in tree squirrels also show that they exhibit increased anti-predator behavior (vigilance or

FID) to the physical presence of a dog with a human handler (Gustafson and VanDruff, 1990;

Cooper et al., 2008) or behave in such a way in areas with high levels of dogs and cats that

suggests that they perceive a higher predation risk in such areas (i.e., giving up density was

higher in study areas where cats and dogs are present: Bowers and Breland, 1996). Instead, we

found that squirrels at both sites did not differ in their FAD when exposed to dog playbacks

compared to the control playbacks. Additionally, only squirrels in less urban sites had a

significantly higher FID when exposed to dog playbacks compared to the control playback.

These results differ from previous studies in tree squirrels and owls showing that FAD and/or

FID were increased in squirrels or owls in urban areas when they were presented with a human

plus dog compared to just a human (Gustafson and VanDruff, 1990; Cooper et al., 2008; Cavalli

et al., 2016). This suggests that squirrels in urban sites were more tolerant to dog vocalizations

when paired with a human observer, whereas squirrels in less urban areas viewed the sounds of

dogs paired with humans as threatening. We predict that squirrels in urban environments exhibit

selective tolerance where the response to humans or stimuli from their commensals (dogs) is

attenuated but the increased response to natural predators is maintained despite this tolerance to

humans and their dogs.

Together, our results provide insight into how urbanization may shape the behavioral

characteristics animals. As most studies on this topic are in birds (Samia et al., 2015), which can

escape from humans using flight, it is important to consider if the same patterns are found in

terrestrial animals. Our results show that non-volant animals in urban environments exhibit less

vigilance and anti-predator behavior. We also show that squirrels in urban environments were

more tolerant to humans but still exhibited a strong behavioral response to acoustic stimuli from

a natural predator (hawks). Although we do not wish to imply that tolerance or habituation to

humans is cost-free, most studies fail to find evidence that populations where individuals are

more tolerant of humans (or in some cases habituated to their presence) also exhibit reduced anti-

predator behavior to stimuli from natural predators. Given that urbanization is unlikely to slow,

increased effort is needed to determine if increased tolerance and/or habituation to humans

carries costs (Geffroy et al,. 2015). Some studies suggest the costs of human tolerance may be

more nuanced, such as tolerance to humans reducing the latency to return to the nest following a

disturbance in nesting shorebirds, but potentially causing increased chick mortality due to the

presence of dogs that often are paired with human stimuli (Baudains and Lloyd,  2007). Other

studies that increased tolerance of humans could even be beneficial for populations that cannot

avoid anthropogenic stimuli due to seasonal food pulses coinciding with a large influx of tourists

(Wheat and Wilmers, 2016). Clearly more work is needed on this subject, especially on a greater

number of species including species that are not "urban exploiters" like tree squirrels, but the

existing evidence rejects hypothesis that there is a cost to human tolerance in terms of lowering

the anti-predator behavior of animals to other natural predators.

**Data Availability Statement**

All data will be archived at Dryad before final acceptance and all regulations set by the journal

will be followed.

**References**

Addison, A., (2017, January 27). *Red-tailed Hawk (Buteo jamaicensis)* [XC401311].

https://www.xeno-canto.org/401311

Anonymous. 2012. Guidelines for the treatment of animals in behavioural research and teaching.

Anim Behav 83:301-309.

Arnqvist G, Henriksson S. 1997. Sexual cannibalism in the fishing spider and a model for the

evolution of sexual cannibalism based on genetic constraints. Evol Ecol 11:255-273.

Baldwin JM. 1896. A new factor in evolution. The American Naturalist 30:441-451.

Bartholomew GA. 1964. The roles of physiology and behaviour in the maintenance of

homeostasis in the desert environment. Symp Soc Exp Biol 18:7-29.

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4.

J Statistical Software, 67:1-48. doi:10.18637/jss.v067.i01.

Baudains TP, Lloyd P. 2007. Habituation and habitat changes can moderate the impacts of

human disturbance on shorebird breeding disturbance. Animal Conservation 10: 400-407.

Blumstein DT. 2003. Flight-initiation distance in birds is dependent on intruder starting distance

J Wildlife Management 67: 852-857.

Blumstein DT, Fernández-Juricic E, Zollner PA, Garity SC. 2005. Inter-specific variation in

avian responses to human disturbance. J Applied Ecology 42:943-953.

Bohls P, Koehnle TJ. 2017. Responses of eastern gray squirrels (*Sciurus carolinensis*) to

predator calls and their modulation by coat color. The American Midland Naturalist 178:226-

236.

Bókony V, Kulcsár A, Tóth Z, Liker A. 2012. Personality traits and behavioral syndromes in

differently urbanized populations of house sparrows (*Passer domesticus*). PLoS One 7:e36639.

Bowers MA, Breland B. 1996. Foraging of gray squirrels on an urban-rural gradient: use of the

GUD to assess anthropogenic impact. Ecological Applications 6:1135-1142.

Carrete M, Tella JL. 2010. Individual consistency in flight initiation distances in burrowing owls:

a new hypothesis on disturbance-induced habitat selection. Biol Lett 6:167-170.

Carrete M, Tella JL. 2017. Behavioral correlations associated with fear of humans differ between

rural and urban burrowing owls. Frontiers in Ecology and Evolution 5:54.

Cavalli M, Baladrón AJ, Isacch JP, Biondi LM, Bó MS. 2016. Differential risk perception of

rural and urban Burrowing Owls exposed to humans and dogs. Behavioural Processes 124:60-65.

Center for International Earth Science Information Network - CIESIN - Columbia University.

(2018). *Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11*.

NASA Socioeconomic Data and Applications Center (SEDAC).

https://doi.org/https://doi.org/10.7927/H49C6VHW

Chartier A (2008, June 24). *Red-tailed Hawk (Buteo jamaicensis borealis)*. [XC31161].

https://www.xeno-canto.org/31161

Coleman A, Richardson D, Schechter R, Blumstein DT. 2008. Does habituation to humans

influence predator discrimination in Gunther's dik-diks (*Madoqua guentheri*)? Biol Lett 4:250-

252.

Cooke AS. 1980. Observations on how close certain passerine species will tolerate an

approaching human in rural and suburban areas. Biol Cons 18:85-88.

Cooper CA, Neff AJ, Poon DP, Smith GR. 2008. Behavioral responses of eastern gray squirrels

in suburban habitats differing in human activity levels. Northeastern Naturalist 15:619-625.

Cooper WE, Frederick WG. 2007. Optimal flight initiation distance. J Theoretical Biology

244:59–67. https://doi.org/10.1016/j.jtbi.2006.07.011

Crooks KR. 2002 Relative sensitivities of mammalian carnivores to habitat fragmentation. Cons

Biol 16:488-502.

Dill LM, Houtman R. 1989. The influence of distance to refuge on flight initiation distance in the

gray squirrel (*Sciurus carolinensis*). Can J Zool 67:233–235. https://doi.org/10.1139/z89-033

Ditchkoff SS, Saalfeld ST, Gibson CJ. 2006. Animal behavior in urban ecosystems:

modifications due to human-induced stress. Urban Ecosyst 9:5-12.

Dochtermann NA. 2011. Testing Cheverud's conjecture for behavioral correlations and

behavioral syndromes. Evolution 65:1814-1820.

Engelhardt SC, Weladji RB. 2011. Effects of levels of human exposure on flight initiation

distance and distance to refuge in foraging eastern gray squirrels (*Sciurus carolinensis*). Can J

Zool 89:823-830.

Evans J, Boudreau K, Hyman J. 2010. Behavioural syndromes in urban and rural populations of

song sparrows. Ethology 116:588-595.

Fernández-Juricic E, Schroeder  N. 2003. Do variations in scanning behavior affect tolerance to

human disturbance? Applied Animal Behaviour Science 84:219-234.

Fitzgerald LA, Stronza AL. 2016. In defense of the ecotourism shield: a response to Geffroy et

al. Trends Ecol Evol 31:94.

Floyd, T. (2017a, January 2). *Black-capped Chickadee (Poecile atricapillus)*. [XC348981].

https://www.xeno-canto.org/348981

Floyd T. (2017b, January 25). *Black-capped Chickadee (Poecile atricapillus)*. [XC352827].

https://www.xeno-canto.org/352827

Geffroy B, Samia DSM, Bessa E, Blumstein DT. 2015. How nature-based tourism might

increase prey vulnerability to predators. Trends Ecol Evol 30:755-765.

Gustafson EJ, VanDruff LW. 1990. Behavior of black and gray morphs of *Sciurus carolinensis*

in an urban environment. The American Midland Naturalist 123:186-192.

Guttman N, Kalish HI. 1956. Discriminability and stimulus generalization. J Exp Psych 51:79-

88.

Hinde RA. 1954. Factors governing the changes in strength of a partially inborn response, as

shown by the mobbing behaviour of the chaffinch (*Fringilla coelebs*). II. The waning of the

response. Proc Roy Soc B 142:331-358.

Johnson MTJ, Munshi-South J. 2017. Evolution of life in urban environments. Science 358:607.

https://doi.org/10.1126/science.aam8327

Jokimäki J, Selonen V, Lehikoinen A, Kaisanlahti-Jokimäki M-L. 2017. The role of urban

habitats in the abundance of red squirrels (*Sciurus vulgaris*, L.) in Flnland. Urban Forestry &

Urban Greening 27:100-108.

Koprowski JL. 1994. *Sciurus niger*. Mammalian Species 479:1–9.

https://doi.org/10.2307/3504263

Korschgen LJ. 1981. Foods of fox and gray squirrels in Missouri. J Wildlife Management

45:260-266.

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest package: tests in linear mixed

effects models. Journal of Statistical Software 82:1-26. doi: 10.18637/jss.v082.i13

Labra A, Leonard R. 1999. Intraspecific variation in antipredator responses of three species of

lizards (*Liolaemus*): possible effects of human presence. J Herpetol. 33:441-448.

Lapiedra O, Chejanovski Z, Kolbe JJ. 2017. Urbanization and biological invasion shape animal

personalities. Global Change Biology 23:592-603.

Lenth R. 2020. emmeans: estimated marginal means, aka least-squares means. R package version

1.5.2-1. https://CRAN.R-project.org/package=emmeans

Lilly MV, Lucore EC, Tarvin KA. 2019. Eavesdropping grey squirrels infer safety from bird

chatter. PLoS One 14:e0221279.

Lima SL, Dill L. 1990. Behavioral decisions made under the risk of predation: a review and

prospectus. Can J Zool 68:610-640.

Loss SR, Will T, Marra PP. 2013. The impact of free-ranging domestic cats on wildlife of the

United States. Nature Communications 4:1396.

Lowry H, Lill A, Wong BBM. 2013. Behavioural responses of wildlife to urban environments.

Biological Reviews 88:537–549. https://doi.org/10.1111/brv.12012

McCleery RA. 2009. Changes in fox squirrel anti-predator behaviors across the urban-rural

gradient. Landscape Ecology 24:483–493. https://doi.org/10.1007/s10980-009-9323-2

McCleery RA, Lopez RR, Silvy NJ, Gallant DL. 2008. Fox squirrel survival in urban and rural

environments. Journal of Wildlife Management 72:133–137. https://doi.org/10.2193/2007-138

Møller AP. 2008. Flight distance of urban birds, predation, and selection for urban life. Behav

Ecol Sociobiol 63:63-75.

Møller AP. 2012. Urban areas as refuges from predators and flight distance of prey. Behav Ecol

23:1030-1035.

Myers RE, Hyman J. 2016. Differences in measures of boldness even when underlying

behavioral syndromes are present in two populations of the song sparrow (*Melospiza melodia*). J

Ethol 34:197-206.

Olson JK, Acevedo-Gutiérrez A. 2017. Influence of human exposure on the anti-predator

response of harbor seals (*Phoca vitulina*). Aquatic Mammals 43:673-681.

Place A. (2015, September). *Black-capped Chickadee (Poecile atricapillus)*. [XC289220].

https://www.xeno-canto.org/289220

Price TD, Qvarnström A, Irwin DE. 2003. The role of phenotypic plasticity in driving genetic

evolution. Proc Roy Soc B 270:1433-1440.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton D, Colombo J, Coppola G, Geyer MA,

Glanzman DL, Marsland S, McSweeney F, Wilson DA, Wu C-F, Thompson RF. 2009.

Habituation revisited: an updated and revised description of the behavioral characteristics of

habituation. Neurobiol Learn Mem 92:135-138.

Rodriguez-Prieto I, Ferndández-Juricic E, Martin J, Regis Y. 2009. Antipredator behavior in

blackbirds: habituation complements risk allocation. Behav Ecol 20:371-377.

Ryan AM, Partan SR. 2014. Urban wildlife behavior. In: McCleery RA, Moorman CE, Peterson

MN, editors. Urban Wildlife Conservation. New York, USA: Springer p. 149-173.

Samia DSM, Nakagawa S, Nomura F, Rangel TF, Blumstein DT. 2015. Increased tolerance to

humans among disturbed wildlife. Nat Comm 6:8877.

Sarno RJ, Parsons M, Ferris A. 2015. Differing vigilance among gray squirrels (*Sciuridae

carolinensis*) along an urban-rural gradient on Long Island. Urban Ecosyst 18:517-523.

Scales J, Hyman J, Hughes M. 2011. Behavioral syndromes break down in urban song sparrow

populations. Ethology 117:887-895.

Seress G, Bókony V, Heszberger J, Liker A. 2011. Response to predation risk in urban and rural

house sparrows. Ethology 117:896-907.

Sih A, Bell A, Johnson JC. 2004. Behavioral syndromes: An ecological and evolutionary

overview. Trends Ecol Evol 19:372–378. https://doi.org/10.1016/j.tree.2004.04.009

Simion D. (2016, August 1). *German Shepard Sound* [Audio]. http://soundbible.com/2146-

German-Shepard.html

Simion D. (2017, May 12). *Doberman Pinscher Sound* [Audio]. http://soundbible.com/2194-

Doberman-Pinscher.html

Simion D. (2018, May 4). *Labrador Barking Dog Sound* [Audio]. http://soundbible.com/2215-

Labrador-Barking-Dog.html

Snell-Rood EC. 2013. An overview of the evolutionary causes and consequences of behavioural

plasticity. Anim Behav 85:1004-1011.

Sol D, Lapiedra O, González-Lagos C. 2013. Behavioural adjustments for a life in the city. Anim

Behav 85:1101-1112.

Sprau P, Dingemanse NJ. 2017. An approach to distinguish between plasticity and non-random

distributions of behavioral types along urban gradients in a wild passerine bird. Front Ecol Evol

5:92.

Thompson RF, Spencer WA. 1966. Habituation: a model phenomenon for the study of neuronal

substrates of behavior. Psychological Review 73:16-43.

Tumlison R. 2012. Unusual foraging tactics by a red-tailed hawk in an urban environment.

Wilson J Ornithol 124:818-820.

Tuomainen U, Candolin U. 2011. Behavioural responses to human-induced environmental

change. Biol Rev 86:640-657.

Uchida K, Suzuki KK, Shimamoto T, Yanagawa H, Koizumi I. 2017. Escaping height in a tree

represents a potential indicator of fearfulness in arboreal squirrels. Mammal Study 42:39-43.

Uchida K, Suzuki KK, Shimamoto T, Yanagawa H, Koizumi I. 2019. Decreased vigilance or

habituation to humans? Mechanisms on increased boldness in urban animals. Behav Ecol,

30:1583–1590. https://doi.org/10.1093/beheco/arz117

Uchida K, Shimamoto T, Yanagawa H, Kolzumi I. 2020. Comparison of multiple behavioral

trials between urban and rural squirrels. Urban Ecosyst 23:745-754.

Vincze E, Papp S, Preiszner B, Seress G, Bókony V, Liker A. 2016. Habituation to human

disturbance is faster in urban than rural house sparrows. Behav Ecol 27:1304-1313.

Vincze E, Pipoly I, Seress G, Preiszner B, Papp S, Németh B, Liker A, Bókony V. 2019. Great

tits take greater risk toward humans and sparrowhawks in urban habitats than in forests.

Ethology 125:686-701.

Wauters LA, Somers L, Dhondt AA. 1997. Settlement behaviour and population dynamics of

reintroduced red squirrels *Sciurus vulgaris* in a park in Antwerp, Belgium. Biological

Conservation 82:101-107.

Weaver M, Ligon RA, Mousel M, McGraw KJ. 2018. Avian anthrophobia? Behavioral and

physiological responses of house finches (*Haemorhous mexicanus*) to human and predator

threats across an urban gradient. Landscape and Urban Planning 179:46-54.

Webster, M.M. & Rutz, C. (2020) How STRANGE are your study animals? Nature 582, 337-

340.

West-Eberhard MJ. 1989. Phenotypic plasticity and the origins of diversity. Ann Rev Ecol Syst

20:249-278.

Wheat RE, Wilmers CC. 2016. Habituation reverses fear-based ecological effects in brown bears

(*Ursus arctos*). Ecosphere 7:e01408.

Wilson T. (2010, May 29). *Red-tailed Hawk (Buteo jamaicensis)* [XC53797]. https://www.xeno-

canto.org/53797

Ydenberg RC, Dill LM. 1986. The economics of fleeing from predators. Advances in the Study

of Behavior, 16:229–249.

Zuur AF, Ieno EN, Elphick CS. 2010. A protocol for data exploration to avoid common

statistical problems. Methods Ecol Evol 1:3-14. https://doi.org/10.1111/j.2041-

210X.2009.00001.x

**Table 1.** Behavioral data from fox squirrels were collected at six sites around Ann Arbor, Michigan, USA located in areas that were classified as urban or less urban. Urban sites were in areas with human population densities of greater than 1000 persons/km$^2$ whereas less urban sites were in areas with population densities of 250 – 1000 persons/km$^2$ (County Farm Park, Nichols Arboretum) or fewer than 250 persons per square km (Saginaw Forest). Population density estimates obtained from the Center for International Earth Information Network of Columbia University (2018). Total days of observation and hours over these days are provided. When we visited the sites, human and dog presence was quantified (see methods) with mean and 1 standard deviation (SD) shown below.

| Site Name | GPS Location of Site | Site Category | Population Density (persons/ sq km) | Human Presence | | Dog Presence | | Total Trials | Total Observation Days | Total Observation Hours |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | Mean Count | SD | Mean Count | SD | | | |
| Central Campus | N 42°16.622' W083°44.293 ' | Urban | > 1000 | 97.7 | 7.3 | 0.39 | 0.61 | 33 | 6 | 9.68 |
| North Campus | N 42°17.753' W083°42.620 ' | Urban | > 1000 | 68.0 | 11.4 | 0 | 0 | 25 | 4 | 6.18 |
| Prospect Park | N 42°14.825' W083°36.093 ' | Urban | > 1000 | 3.6 | 10.2 | 0.69 | 1.55 | 23 | 4 | 6.08 |
| County Farm Park | N 42°15.430' W083°42.518 ' | Less urban | 250 - 1000 | 1.6 | 1.3 | 0.41 | 0.50 | 32 | 5 | 8.73 |
| Nichols Arboretum | N 42°16.940' W083°43.410 ' | Less urban | 250 - 1000 | 11.9 | 2.3 | 2.32 | 1.19 | 28 | 5 | 9.07 |
| Saginaw Forest | N 42°16.235' W083°48.465 ' | Less urban | 25 - 250 | 0.37 | 0.50 | 0.20 | 0.41 | 30 | 6 | 12.62 |

**Table 2.** Differences between fox squirrels at urban and less urban sites that were not exposed to any playbacks for first alert distance (FAD), flight initiation distance (FID), probability of escaping the observer by climbing a tree during the trial, and latency to resume behavior following the trial. A random effect for site identity was included in the model or FAD ($\sigma^2$ = 6.9), FID ($\sigma^2$ = 6.24), probability of climbing a tree ($\sigma^2$ = 1.23), and latency ($\sigma^2$ = 1.22). Latency was ln+1 transformed. Results are from 77 trials from six sites.

| Response Variable | Variable | *b* | SE | *t* or *z* | *P* |
|---|---|---|---|---|---|
| First alert distance (FAD) | Intercept (Less urban, Female) | 14.11 | 1.63 | 8.67 | 0.0007 |
| | Site (Urban) | -4.79 | 2.26 | -2.11 | 0.10 |
| | Distance to tree | -0.37 | 0.38 | -0.98 | 0.33 |
| | Sex (Male) | 1.17 | 0.66 | 1.77 | 0.08 |
| | Julian Date | 0.15 | 0.40 | 0.39 | 0.70 |
| | Initial Distance | 2.63 | 0.35 | 7.51 | <0.0001 |
| Flight initiation distance (FID) | Intercept (Less urban, Female) | 12.16 | 1.54 | 7.89 | 0.0009 |
| | Site (Urban) | -6.22 | 2.14 | -2.90 | 0.043 |
| | Distance to tree | -0.56 | 0.35 | -1.59 | 0.12 |
| | Sex (Male) | 0.85 | 0.61 | 1.38 | 0.17 |
| | Julian Date | 0.01 | 0.37 | 0.03 | 0.97 |
| | Initial Distance | 1.38 | 0.32 | 4.24 | <0.0001 |
| Probability of climbing a tree | Intercept (Less urban, Female) | 0.71 | 0.79 | 0.89 | 0.37 |
| | Site (Urban) | -1.89 | 1.12 | -1.68 | 0.09 |
| | Distance to tree | -0.09 | 0.32 | -0.27 | 0.78 |
| | Sex (Male) | 0.17 | 0.55 | 0.31 | 0.76 |
| | Julian Date | 0.47 | 0.35 | 1.34 | 0.18 |
| | Initial Distance | 0.23 | 0.30 | 0.76 | 0.45 |
| Latency to resume behavior | Intercept (Less urban, Female) | 4.48 | 0.67 | 6.70 | 0.002 |
| | Site (Urban) | -1.87 | 0.93 | 1.99 | 0.11 |

| | | | | |
|---|---|---|---|---|
| Distance to tree | -0.07 | 0.13 | -0.57 | 0.57 |
| Sex (Male) | 0.06 | 0.23 | 0.25 | 0.80 |
| Julian Date | 0.07 | 0.14 | 0.52 | 0.61 |
| Initial Distance | 0.002 | 0.12 | 0.02 | 0.98 |

**Table 4.** Associations between latency to resume behavior following the trial ("Latency") and first alert distance (FAD), flight initiation distance (FID), and probability of escaping the human observer by climbing a tree during the trial ("Went up tree?") in fox squirrels at urban and less urban sites that were not exposed to any playbacks. A random effect for site identity was included in the model or FAD ($\sigma^2 = 0.02$), FID ($\sigma^2 = 0.08$), and probability of climbing a tree ($\sigma^2 = 1.03$). FAD, FID, and latency were ln+1 transformed. Results are from 77 trials from six sites.

| Response Variable | Variable | *b* | SE | *t* or *z* | *P* |
|---|---|---|---|---|---|
| First alert distance (FAD) | Intercept (Urban, Female) | 2.42 | 0.10 | 24.2 | 0.0004 |
| | Site (Less urban) | 0.27 | 0.14 | 1.91 | 0.16 |
| | Latency | 0.25 | 0.08 | 3.07 | 0.005 |
| | Went up tree? | -0.10 | 0.07 | -1.39 | 0.17 |
| | Site (Less urban) x Latency | -0.23 | 0.12 | -2.01 | 0.053 |
| | Site (Less urban) x Went up tree? | 0.13 | 0.09 | 1.41 | 0.16 |
| Flight initiation distance (FID) | Intercept (Urban, Female) | 1.94 | 0.18 | 11.0 | 0.0005 |
| | Site (Less urban) | 0.63 | 0.25 | 2.54 | 0.067 |
| | Latency | 0.16 | 0.08 | 1.86 | 0.067 |
| | Went up tree? | -0.08 | 0.07 | -1.18 | 0.24 |
| | Site (Less urban) x Latency | -0.14 | 0.12 | -1.17 | 0.24 |
| | Site (Less urban) x Went up tree? | 0.06 | 0.09 | 0.70 | 0.48 |
| Latency to resume behavior | Intercept (Urban, Female) | 2.77 | 0.61 | 4.56 | 0.01 |
| | Site (Less urban) | 1.61 | 0.86 | 1.88 | 0.13 |
| | Went up tree? | 0.43 | 0.21 | 2.08 | 0.041 |
| | Site (Less urban) x Went up tree? | 0.01 | 0.26 | 0.05 | 0.96 |

**Table 4.** Effects of acoustic playbacks (control, hawk, dog) on first alert distance (FAD), flight initiation distance (FID), probability of escaping human observer by climbing a tree, and latency to resume behavior following the trial for fox squirrels observed at urban or less urban sites. Reference (intercept) was "less urban" for site, "control playback" for treatment, and "female" for sex. A random effect for site identity was included in the model or FAD ($\sigma^2 = 6.5$), FID ($\sigma^2 = 4.9$), probability of climbing a tree ($\sigma^2 = 0.07$), and latency ($\sigma^2 = 0.37$). Latency was ln+1 transformed. Results are from 94 trials from six study sites.

| Response Variable | Variable | *b* | SE | *t* or *z* | *P* |
|---|---|---|---|---|---|
| First alert distance (FAD) | Intercept (Less urban, Female, Control) | 14.94 | 0.89 | 16.71 | <0.0001 |
| | Site (Urban) | -2.32 | 1.33 | -1.74 | 0.11 |
| | Dog playbacks | 0.51 | 0.90 | 0.57 | 0.57 |
| | Hawk playbacks | -0.29 | 0.98 | -0.30 | 0.76 |
| | Distance to tree | 0.36 | 0.29 | 1.23 | 0.22 |
| | Sex (Male) | 0.29 | 0.58 | 0.51 | 0.61 |
| | Julian Date | -0.10 | 0.28 | -0.34 | 0.73 |
| | Initial Distance | 3.20 | 0.29 | 10.84 | <0.0001 |
| | Site (Urban) x Dog playbacks | 0.07 | 1.35 | 0.05 | 0.96 |
| | Site (Urban) x Hawk playbacks | 4.29 | 1.49 | 2.86 | 0.005 |
| Flight initiation distance (FID) | Intercept (Less urban, Female, Control) | 12.3 | 0.95 | 7.31 | <0.0001 |
| | Site (Urban) | -1.64 | 1.39 | -1.18 | 0.27 |
| | Dog playbacks | 1.89 | 0.79 | 2.40 | 0.018 |
| | Hawk playbacks | 1.21 | 0.85 | 1.42 | 0.16 |
| | Distance to tree | 0.08 | 0.26 | 0.30 | 0.76 |
| | Sex (Male) | 0.01 | 0.50 | 0.01 | 0.99 |
| | Julian Date | -0.39 | 0.25 | -1.59 | 0.11 |
| | Initial Distance | 2.99 | 0.26 | 11.51 | <0.0001 |
| | Site (Urban) x Dog playbacks | -1.52 | 1.18 | -1.29 | 0.20 |

| | | | | | |
|---|---|---|---|---|---|
| | Site (Urban) x Hawk playbacks | 2.53 | 1.30 | 1.94 | 0.056 |
| Probability of climbing a tree | Intercept (Less urban, Female, Control) | -0.20 | 0.62 | -0.33 | 0.74 |
| | Site (Urban) | -0.61 | 0.92 | -0.66 | 0.51 |
| | Dog playbacks | 1.14 | 0.82 | 1.39 | 0.16 |
| | Hawk playbacks | 1.26 | 0.90 | 1.40 | 0.16 |
| | Distance to tree | -0.64 | 0.27 | -2.34 | 0.02 |
| | Sex (Male) | 0.61 | 0.55 | 1.10 | 0.27 |
| | Julian Date | -0.26 | 0.27 | -0.97 | 0.33 |
| | Initial Distance | 0.01 | 0.27 | 0.04 | 0.96 |
| | Site (Urban) x Dog playbacks | -1.52 | 1.20 | -1.27 | 0.20 |
| | Site (Urban) x Hawk playbacks | 0.18 | 1.34 | 0.14 | 0.89 |
| Latency to resume behavior | Intercept (Less urban, Female, Control) | 3.83 | 0.39 | 9.69 | 0.0001 |
| | Site (Urban) | -0.60 | 0.57 | -1.05 | 0.33 |
| | Dog playbacks | 1.14 | 0.23 | 4.91 | <0.0001 |
| | Hawk playbacks | 1.71 | 0.25 | 6.76 | <0.0001 |
| | Distance to tree | 0.03 | 0.07 | 0.39 | 0.70 |
| | Sex (Male) | 0.24 | 0.15 | 1.61 | 0.11 |
| | Julian Date | 0.07 | 0.07 | -0.91 | 0.36 |
| | Initial Distance | 0.06 | 0.08 | 0.82 | 0.41 |
| | Site (Urban) x Dog playbacks | 0.03 | 0.35 | 0.10 | 0.92 |
| | Site (Urban) x Hawk playbacks | 0.22 | 0.39 | 0.57 | 0.57 |

**Figures and Figure Legends**

**Figure 1.** Variation in A) vigilance (first alert distance: FAD), B) anti-predator behavior (flight initiation distance: FID), C) probability of escaping up a tree during the trial, and D) latency to resume behavior following the trial among fox squirrels at urban (n = 38 trials) and less urban (n = 39) sites in trials where squirrels were not exposed to any playbacks. Squirrels in urban areas had significantly shorter FID compared to those in the less urban sites, but there were no other significant differences (Table 2). Upper and lower hinges correspond to first and third quartile, respectively. Upper/lower whiskers extend from the hinge to the highest/lowest value that is within 1.5x the interquartile range. Solid horizontal line shows median.

**Figure 2.** Associations between the latency of fox squirrels in urban and less urban study areas to resume typical behavior after a disturbance trial with A) first alert distance (FAD), B) flight initiation distance (FID), and C) probability of escaping the human observer by climbing up a tree. FAD and latency to resume typical behavior were significantly positively associated in urban squirrels but not those in less urban areas. In squirrels in both urban and less urban areas, those that had a higher FID or climbed a tree also took longer to resume typical behavior. differed between urban and less urban populations of squirrels. Note that latency is ln+1 transformed.

**Figure 3.** Effects of human observer approaching a squirrel while broadcasting one of three playback treatments (control playback, hawk or dog vocalizations) on A) first alert distance (FAD), B) flight initiation distance (FID), C) probability of escaping up a tree during the trial, and D) latency to resume typical behavior following the trial. Trials were conducted at less urban (n = 15 control, 20 dog, 16 hawk) and urban (n = 11 control, 20 dog, 12 hawk) sites. Results shown in Table 4. Upper and lower hinges correspond to first and third quartile, respectively. Upper/lower whiskers extend from the hinge to the highest/lowest value that is within 1.5x the interquartile range. Solid horizontal line shows median.

**Appendix I**

**Figure S1.** Overview of trial protocol. A marker was placed at the initial position of the observer when they initially identified a squirrel for a trial. The distance between observer initial position and the squirrel's initial position is the initial distance. From this position, the observer walked at a slow, steady, and constant  pace in a direct line towards the squirrel. The observer then dropped a marker to represent FAD when the squirrel first displayed "alert" behavior (freezing with eye contact). An additional marker was dropped when the squirrel fled from its initial position to represent FID. Another marker was placed at an estimate of the squirrel's initial position. In playback manipulation trials, acoustic stimuli were broadcast when the observer started approaching the squirrel and stopped playing the stimuli when the squirrel fled. Distance from the squirrel's initial position to the nearest tree was recorded along with how high in the tree the squirrel took refuge. If the squirrel took refuge in a tree other than the one nearest, then distance to the chosen tree was recorded. Once the squirrel took refuge latency to resume normal activity (foraging or traveling off tree) was recorded. Trials ended when squirrels resumed normal activity or if latency exceeded 10 min.