Cell patterning by secretion-induced plasma membrane flows

Veneta Gerganova[1,&], Iker Lamas[1,&], David M. Rutkowski[2,&], Aleksandar Vještica[1], Daniela Gallo Castro[1], Vincent Vincenzetti[1], Dimitrios Vavylonis[2,*] and Sophie G Martin[1,*]

[1]Department of Fundamental Microbiology, University of Lausanne, CH-1015 Switzerland.
[2]Department of Physics, Lehigh University, Bethlehem PA 18015 USA.
*Correspondence to: sophie.martin@unil.ch; vavylonis@lehigh.edu
[&]These authors contributed equally to this work.

Abstract

Cells self-organize using reaction-diffusion and fluid-flow principles. Whether bulk membrane flows contribute to cell patterning has not been established. Here, using mathematical modelling, optogenetics and synthetic probes, we show that polarized exocytosis causes lateral membrane flows away from regions of membrane insertion. Plasma membrane-associated proteins with sufficiently low diffusion and/or detachment rates couple to the flows and deplete from areas of exocytosis. In rod-shaped fission yeast cells, zones of Cdc42 GTPase activity driving polarized exocytosis are limited by GTPase activating proteins (GAPs). We show that membrane flows pattern the GAP Rga4 distribution and that coupling of a synthetic GAP to membrane flows is sufficient to establish the rod shape. Thus, membrane flows induced by Cdc42-dependent exocytosis form a negative feedback restricting the zone of Cdc42 activity.

Introduction

Cells are highly polarized entities that exhibit one (or several) fronts with distinct lipid and protein plasma membrane composition. Reaction-diffusion systems have come a long way in explaining polarity establishment through positive and negative feedback regulations. For instance, in yeast cells, scaffold-mediated positive amplification of Cdc42 GTPase activity underlies the formation of one (or two) cell fronts (*1, 2*), with negative feedback promoting oscillations (*3, 4*). Biochemical reactions are also modulated by physical processes such as fluid flows, which may happen in the cytosol and at the membrane. In large metazoan cells, cytoskeletal-driven cortical and cytosolic fluid flows, causing advective protein movements, play important roles in cell polarization (*5*), and can in principle promote polarity establishment (*6*). Membrane flows are predicted to occur and equilibrate membrane tension between spatially segregated zones of exo- and endocytosis, such as in tip-growth of walled pollen tubes (*7, 8*) or in cell migration, as initially proposed in (*9*). However, even though membrane flows have been observed in migrating amoebae, apicomplexans, amoeboid macrophages and neuronal growth cones (*10-13*), their role in cell patterning has not been explored.

In contrast to metazoan cells, fungal cells are not thought to harbour an actin cortex that may contribute to cortical flow forces. In fission yeast cells, which are rod-shaped and grow by tip extension at one or both cell poles, Cdc42 GTPase activity promotes local exocytosis for polarized cell growth. Membrane is retrieved through endocytosis, which also occurs mainly

at cell poles. The width of the cell correlates with the size of the Cdc42-GTP zones at cell poles, which is regulated by GTPase-activating proteins (GAPs) inhibiting Cdc42 activity (*14*). In particular, Cdc42 GAPs that localize to the sides of rod-shaped fission yeast cells through unknown mechanisms are thought to control cell width by restricting the size of the Cdc42 active zone (*14-16*).

We recently described the use of the CRY2-CIB1 optogenetic system in fission yeast cells (*2*). In our CRY2-CIB1 setup, blue light activates the cytosolic photosensor domain CRY2PHR (CRY2) to bind its ligand CIBN, which is linked to the plasma membrane (PM)-interacting domain of the mammalian protein Rit1 (RitC). RitC, which is a disordered peptide that reversibly associates with the membrane through hydrophobic and charge interactions (*17*), specifically associates with the PM but not with secretory vesicles or other endomembranes (*18, 19*). Blue light also promotes CRY2 oligomerization (*20*). Here, we describe our serendipitous discovery of membrane flows away from sites of secretion, revealed by the behaviour of the CRY2-CIB1 system. We show that plasma membrane-associated proteins with sufficiently low diffusion and/or detachment rates, including the Cdc42 GAP Rga4, are displaced from sites of exocytosis. We further reveal that coupling of a synthetic GAP to membrane flows is sufficient to establish the rod cell shape. Thus, secretion-induced membrane flows underlie a negative feedback that shapes the cell by restricting the size of the secretion zone

Results

Whereas CIBN-RitC is nearly uniform at the PM in the dark and blue light initially caused uniform CRY2 recruitment, we unexpectedly observed that both proteins cleared from most interphase cell poles within 5 min (Fig 1A, Movie 1). Similarly, in pre-divisional cells, CRY2 and CIBN cleared from mid-cell, where exocytosis is targeted, but re-appeared at non-growing cell poles (Fig S1A, Movie 2). CRY2 depletion at cell poles after 5 min correlated with local levels of Cdc42-GTP, marked by CRIB-3GFP (Fig 1A-B), the secretory vesicle associated Rab11-like GTPase Ypt3 (Fig 1B, S1B), exocyst subunits Exo70 and Exo84 (Fig S1C-D), and with fast depletion rates (Fig S1E). Thus, light-induced changes in CRY2-CIBN interactions provoke their clearance from zones of exocytosis.

CRY2 was not depleted from cell poles upon Brefeldin A (BFA) treatment, which completely stops secretion (*21, 22*), with some cell poles even gaining in CRY2 signal (Fig 1C, S1F-G). CRY2 was also not depleted upon secretion inactivation in the *ypt3-i5* mutant (Fig 1D, S1H). We note that both conditions also caused reduction in CRIB-GFP, perhaps due to loss of positive feedback between growth and polarity (*23*). GFP-Ypt3 intensity was also slightly reduced in BFA-treated cells (Fig S1G). Nevertheless, comparison of cell poles with similar GFP intensity clearly showed that CRY2 does not deplete upon secretion block. Thus, polarized secretion is essential for CRY2 depletion from cell poles.

Cell poles are also sites of endocytosis, which is strictly dependent on actin patch assembly by the Arp2/3 complex. To test the role of endocytosis, we disrupted all actin structures with Latrunculin A (LatA). Because LatA treatment elicits activation of the MAPK stress signalling pathway, which leads to loss of cell polarity, we performed these experiments in cells lacking the MAPK Sty1, in which this stress response is abrogated, and cell polarity is retained (*24*). In LatA-treated *sty1Δ* cells, actin patches were lost, but CRY2 still depleted from cell poles (Fig 1E, S1I). As LatA also depolymerizes actin cables, we can conclude that cytoplasmic actin cable

retrograde flow is not driving depletion. We note however that the extent of depletion was reduced relative to untreated samples. We hypothesize that this reduction is due to reduced rates of exocytosis in LatA-treated cells both due to absence of actin cable-mediated vesicle transport and to accommodate the absence of endocytosis, but we cannot exclude a possible direct contribution of endocytosis in CRY2 depletion from cell poles. We conclude that endocytosis may partially contribute but is not essential for CRY2 depletion from cell poles.

Simulations of membrane flow-dependent depletion

How may RitC-CIBN and CRY2 become depleted from sites of polarized growth? Because they are not native to *S. pombe* cells, their localization is unlikely to be directly regulated by endogenous yeast proteins. A reaction-diffusion mechanism would require a source of energy to maintain the pattern and potentially non-linear interactions through activation/deactivation, which is not likely for CIBN/CRY2. We have also shown above that endocytosis could not solely explain the depletion. We thus hypothesized that deposition and retrieval of membrane material through exo- and endocytic events at sites of growth locally perturbs the distribution of peripherally associated membrane proteins, such as CIBN-RitC, through bulk membrane flows causing associated protein movement towards cell sides.

We performed computer simulations to test the hypothesis that exo- and endocytic events generate flows strong enough to deplete membrane-associated proteins at growth sites. We modelled proteins as particles with association/dissociation rate constants k_{on} and k_{off}, diffusing on a sphere approximating the cell tip, and displaced by exo/endocytotic events over zones of width σ_{exo} and σ_{endo} (Fig 2A, Movie 3; see Methods section for details). Particles are displaced radially by up to a distance of R_{cutoff} around the site of each vesicle delivery/internalization (Fig 2B, Eq M2, Eq M3), as an approximation for how proteins in a fluid membrane move, adjusting their position to accommodate the change of membrane area on a background of membrane tension. A minimum value R_{cutoff} ~1 μm was required to observe significant depletion in subsequent analysis (see Fig S9E). To calibrate the model, we measured rates of fission yeast endocytosis (Table 1). We also measured cell growth to estimate the net rate of new membrane addition (Table 1). Rates of exocytosis were derived from the sum of new membrane required to both compensate endocytosis and provide net growth.

We first considered conditions where overall membrane delivery by exocytosis balances internalization by endocytosis, and assumed that neither exocytic nor endocytic vesicles carry membrane-associated proteins. This assumption is based on prior work showing that RitC does not associate with endomembranes (*18, 19*), and on our findings that endocytosis is not necessary for depletion. In these conditions, we found tip depletion when exocytosis is narrower than endocytosis, $\sigma_{exo} < \sigma_{endo}$ (Fig 2C, Fig S2A). This is a condition which we verified in cells (Fig S3 and Methods). When $\sigma_{exo} = \sigma_{endo}$, the active diffusion generated by the randomness of endo- and exocytic events was not by itself sufficient to drive significant concentration gradients (Fig S2A, S2C). Under net growth conditions, where exocytosis exceeds endocytosis by the amount of extra membrane needed for cell growth, depletion was stronger and faster, and occurred even when $\sigma_{exo} = \sigma_{endo}$ (Fig S2A-C). Net growth conditions in absence of endocytosis also led to depletion, even if the rate of exocytosis was concomitantly reduced to retain the same net membrane area addition (Fig S2E). This latter situation, which may best mimic growth upon LatA treatment, showed somewhat reduced

depletion, similar to our experimental observations in LatA-treated cells. Thus, an area-change model of membrane flow can reproduce the observed depletion from cell poles.

Using the measured endocytosis and exocytosis profiles in our model, we plotted the tip occupation ratio, a measure of depletion, as a function of k_{off} and diffusion coefficient D (Fig 2D). The model makes two key, testable predictions: First, significant tip depletion requires both parameters to be sufficiently small – this is the limit where particle movement and exchange are too slow to counteract flow away from the tip (Movies 3, 4). Second, membrane material flowing away from the cell tip accumulates in a region of enhanced density near the end of the depletion area (Fig 2E). Lateral peak formation requires slow diffusion rates (Fig S2D). Both tip depletion and lateral build-up were stronger in conditions of net growth (Fig S2B).

Evidence of membrane flow in vivo

The finding that depletion is only observed for proteins with slow dynamics (low k_{off} or D) provides an explanation for the acute change in CIBN localization upon light-induced CRY2 interaction. Indeed, light activation promotes not only CIBN-CRY2 binding but also formation of CRY2 oligomers through a distinct binding interface (*20, 25*). CRY2 oligomers form bright foci (see Fig 1A), predicted to reduce dynamics through clustering of membrane anchors. Using FRAP to estimate D and k_{off} (Fig S4 and supplemental text), we found that CRY2 interaction causes a >10-fold decrease in CIBN-RitC D and k_{off}, to values consistent with tip depletion in the simulations (Fig 3A). CRY2-CIBN tip depletion was more marked in experiments than predicted by the measured D and k_{off} values, which we reasoned is due to CRY2 oligomerization not only decreasing membrane-unbinding dynamics but also biasing CRY2 binding from the cytosol. This can be accounted for in the model by increasing the rate of particle dimerization on the membrane, which leads to exacerbated tip depletion (Fig S5). We propose that tip depletion events revealed by optogenetics are due to acute clustering-mediated changes in membrane-binding properties.

Simulations of protein removal by endocytic vesicles neglecting membrane flows, which were previously shown to maintain the polarization of slow-diffusing transmembrane proteins delivered by secretory vesicles (*26*), generate tip depletion for similar, low values of k_{off} and D. However, in contrast to simulations with flow displacement, they do not produce a lateral accumulation peak (Fig S2F-G). Depletion and lateral accumulation peaks occur in a model with combined endocytic removal and displacement by flow (Fig. S2H-I). Thus, protein accumulation on the edges of the depletion zone is a characteristic feature of the model with protein displacement by membrane flow. As for tip depletion, protein accumulation on the edges of the depletion zone was dependent upon k_{off} and D, and was strongly enhanced by oligomerization (Fig S5D). Concordant with the flow displacement model, a lateral CRY2 peak corresponding to the edge of the CRIB-labelled depletion zone increased over time (Fig 3B, Fig 1A white arrowhead). The rates of CRY2 levels decrease from cell pole and increase at edges were indistinguishable, suggesting they reflect the same molecular process (Fig 3B). Kymographs of CRY2 oligomers at cell tips also directly showed events of flow away from the cell pole (Fig 3C). We observed similar lateral CRY2 peaks on the edges of the depletion zone at the division site (Fig 3D). These observations support the existence of bulk membrane flows carrying associated proteins away from sites of polarized secretion.

Protein dynamics condition tip depletion

To test the role of membrane binding in flow-mediated protein patterning without the amplifying effect of oligomerization, we constructed a series of one, two or three tandem copies of the membrane anchor RitC tagged with sfGFP (Fig 4A). As RitC binds the membrane peripherally and can exchange between membrane and cytosolic pools, our hypothesis was that tandem copies would lead to increased residence time at the PM. Indeed, estimated k_{off} and D rates extracted from FRAP experiments were higher for 1xRitC than 2xRitC and 3xRitC (Fig 4B). A control 3xsfGFP-1xRitC construct showed only slightly reduced k_{off} compared to sfGFP-1xRitC, but more reduced lateral diffusion, suggesting a possible effect of protein size or perhaps weak sfGFP-mediated oligomerization. The measured dynamic parameters of 2xRitC and 3xRitC, but not 1xRitC and 3xsfGFP-1xRitC, fall within the range predicted in the model to cause local depletion (Fig 4B). Indeed, cell pole depletion was observed for 2xRitC and more prominently 3xRitC, but not 1xRitC constructs (Fig 4C). This is particularly visible in comparison with the uniform localization mCherry-1xRitC co-expressed in these cells (Fig 4A). Signal depletion at cell poles correlated with Cdc42-GTP levels (Fig S6A). In time-lapse experiments, 3xRitC was observed to dynamically deplete from sites marked by CRIB at both growing cell poles and division sites, and repopulate cell poles during cell division when growth stops and CRIB disappears (Fig 4D, Movie S5). Thus, slowing down the dynamics of a peripherally membrane-associated protein suffices to displace it from sites of active secretion.

We further confirmed that 3xRitC depletion from cell poles was independent of endocytosis by probing its distribution in LatA-treated cells, again using the *sty1Δ* mutant background to alleviate the LatA-triggered stress response. Depletion from cell poles is reflected in a < 1 fluorescence ratio between cell tips and cell sides: 0.8 for 2xRitC and 0.5 for 3xRitC (Fig 4E). In *sty1Δ* cells, the tip-to-side ratio of 3xRitC was 0.5 as in wildtype cells and was unchanged upon LatA treatment, indicating that 3xRitC depletion from cell poles does not depend on endocytosis.

Collectively, these data demonstrate that the membrane residence time of peripheral membrane-associated proteins, which is defined by both membrane dissociation and lateral diffusion rates, is a strong predictor of protein distribution around sites of polarized exocytosis induced membrane flows (Fig 4F). The dependence of protein distribution on membrane affinity should invite careful consideration of this parameter when designing probes to evaluate lipid distribution.

Rga4 GAP localization is consistent with membrane flow coupling

To probe the role of membrane flows in cell patterning, we focused on the Cdc42 GAP Rga4, which is excluded from cell poles and forms membrane-associated clusters previously described to form a 'corset' pattern around the site of growth (*15*). Quantitative profiles around the cell cortex indeed showed that Rga4 is depleted from the CRIB-labelled cell poles and its distribution is skewed towards the cell tip with higher CRIB intensity (Fig 5A) (*15, 27*).

Through structure-function analysis of GFP-tagged Rga4 fragments, we defined determinants of membrane association and cluster formation. Consistent with previous observations (*27*), a cortex-binding domain (CBD; aa 623-685) was both necessary and sufficient for membrane

association (Fig 5B-C). This region could be replaced by RitC to target Rga4 N-terminus to the cortex. CBD sequence analysis revealed two conserved sequences: a weak amphipathic helix prediction (membrane-binding (MB) motif #1) and a region rich in basic residues, predicted by a high BH score (28), (MB motif #2) (Fig S7A). Mutations of key residues in these two motifs did not individually block cortical binding, but together resulted in cytosolic localization. This was the case whether the mutations were introduced in a short Rga4 fragment containing coiled-coils (cc) and the CBD (cc-CBD), or whether they were introduced in the full-length Rga4 expressed from the native genomic locus (Fig 5D). Full-length Rga4 containing mutations in both membrane-association motifs, which was cytosolic, was non-functional, as shown by the round shape of cells also lacking the other two Cdc42 GAPs Rga3 and Rga6 (Fig S7B) (29). Thus, these data identify two membrane-binding motifs in the CBD, which are together essential for Rga4 localization at the plasma membrane.

The N-terminal fragment lacking the CBD and full-length Rga4 with a non-functional CBD formed cytosolic condensates, as previously observed upon overexpression of a cytosolic form of Rga4 (14), (Fig 5C-D). These condensates disappeared upon exposure to high temperature (37°C for 6h) or 5% 1,6-hexanediol, thought to interfere with weak hydrophobic interactions (Fig S7C-D), suggesting phase-separation properties. Importantly, in the N-terminal fragment, condensates were absent upon further truncation of the coiled coils (Fig 5C; compare fragments 2 and 3). The Rga4 N-terminus containing the LIM domains likely also contributes to condensate formation, as we did not observe condensates in cytosolic fragments lacking this region (Fig 5D, fragment 9). We conclude that Rga4 forms larger molecular assemblies for which the coiled coils and perhaps other determinants are necessary.

A minimal Rga4 fragment containing both coiled coils and CBD (cc-CBD) recapitulated the main features of Rga4 localization, as it formed membrane-associated clusters that were partly depleted from cell poles (Fig 5E-F). We used this fragment to probe how a simplified endogenous protein may be affected by membrane flows. Truncating the coiled coils led to faster membrane-binding dynamics and re-population of the cell poles (CBD fragment, Fig 5E-G). Tandem multimerization of the CBD (4xCBD) increased membrane affinity and restored depletion from cell poles (Fig 5E-G), as did artificial CBD oligomerization by fusion with CRY2 (Fig S8A). Truncating one of the membrane-binding motifs (cc-CBD-Δ2) to reduce membrane affinity led to faster protein dynamics, as confirmed by FRAP analysis (Fig 5G) and TIRF microscopy (Fig S8B-C, Movie 6-7). This truncation caused enrichment at cell poles (Fig 5E-G), indicating affinity of the coiled coil region for cell tips. Similar enrichment at cell poles was observed upon mutations in either MB motifs #1 or #2 to reduce membrane binding affinity (Fig 5D; fragments 7 and 8). Conversely, increasing membrane-affinity by multimerization (4x cc-CBD-Δ2) restored tip depletion and enrichment to cell sides (Fig 5E-F). The estimated k_{off} and D of all protein fragments largely co-varied, falling in parameter space regions consistent with their observed distribution. We conclude that, even with a preferential association at cell poles, the oligomerization and two membrane-binding motifs of Rga4 ensure a sufficiently long membrane residence time that couples Rga4 to membrane flows towards cell sides.

Flow-driven negative feedback shapes the cell

To directly test the importance of the GAP lateral localization, we constructed a synthetic GAP (optoGAP) consisting of CRY2, the membrane-binding RitC, and the Rga4 GAP domain. RitC targets optoGAP to the plasma membrane. In the dark, the distribution is predicted to be uniform including at cell poles because RitC unbinding and lateral diffusion dynamics are fast enough (as measured in Fig 4B). In the light, oligomerization of CRY2 is predicted to increase

membrane residence time by clustering several RitC molecules, and thus couple the optoGAP probe to membrane flows leading to depletion from cell poles. Indeed, a control CRY2-RitC construct lacking the GAP domain was uniformly distributed in wild-type cells in the dark, but was depleted from cell tips within 7 min of light exposure (Fig 6A). In contrast, in cells lacking all endogenous Cdc42 GAPs, which display very large zones of Cdc42 activity leading to absence of polarized exocytosis, round morphology and poor growth (*29*), CRY2-RitC distribution was uniform in both dark and light conditions. Remarkably, expression of optoGAP in these mutant cells in the light led to a strong reduction in the size of Cdc42-GTP zones, with concomitant local depletion of optoGAP (Fig 6B-C), restoring viability and the rod cell shape (Fig 6D, 6F). By contrast, upon growth in the dark, optoGAP was present around the cell periphery (Fig 6B) and was toxic to the cells (Fig 6F), which retained a round morphology (Fig 6D). The round shape and toxicity are likely due to uniform GAP activity suppressing Cdc42-GTP, as the majority of dark-grown cells showed no cortical CRIB (Fig 6E). These data agree with the conclusion that lateral localization of the Cdc42 optoGAP is critical to constrain the zone of Cdc42-GTP activity and build the rod cell shape.

To further probe the mode of optoGAP exclusion from sites of growth, we imaged cells during the dark-to-light transition, which acutely increases the optoGAP membrane residence time. Light exposure provoked strong local optoGAP depletion coinciding with establishment of a zone of Cdc42 activity, followed by outgrowth of a tube of restricted width (Fig 6G, Movies 8 and 9). During depletion, which occurred before tube outgrowth, clusters of optoGAP exhibited lateral flow movements away from the CRIB-labelled zone, captured in kymographs of the cell periphery. These movements are consistent with the view that optoGAP couples to membrane flows. We note that upon longer growth in the light, optoGAP did not form visible lateral peaks on the edges of the CRIB zone and instead appeared depleted from proximal regions of the growing cell pole, though small amounts were present along the shaft of the tube. We suspect that the large pools of optoGAP present in the back of the cell titrate the majority of optoGAP through strong CRY2-mediated oligomerization. We conclude that coupling of Cdc42 inhibition to membrane flows, which are themselves induced by Cdc42 activity, forms a spatial negative feedback sufficient to restrict zones of Cdc42 activity and define cell width (Fig 6I).

We also observed strong, directional movements of optoGAP away from the division site in cells grown continuously in the light. Most optoGAP cells are monopolar, growing at only one cell pole. At cell division, the optoGAP cleared from the division site, concurrent with the arrival of CRIB marking Cdc42-GTP. Remarkably, optoGAP clusters displayed long-distance (several μm) unidirectional movement towards the cell pole that was growing in the preceding interphase (Fig 6H, Movie 10). While further work will be required to establish the cause of these movements with certainty, these observations agree with the idea of long-distance membrane flows at the PM caused by targeted secretion at the division site and endocytosis still present at the previously growing half of the cell.

Discussion

Eukaryotic cell membranes are in constant flux, with exocytosis and endocytosis universally controlling plasma membrane homeostasis. Serendipitous observations of an optogenetic system in fission yeast cells led to our discovery that membrane-associated proteins can be depleted from sites of exocytosis. Our complementary simulations and experiments indicate

that this depletion is best explained by bulk membrane flow away from the secretion site, where membrane-associated proteins are displaced in a fountain-flow manner away from zones of polarized secretion, due to membrane area addition in the centre of a wider subtraction area. Importantly, not all membrane-associated proteins are depleted from secretion sites: only those with sufficiently long residence time at the membrane are. Because membrane flows are a consequence of polarized secretion, they are likely to be prevalent in cells and pattern membrane-associated proteins, coordinating with the cytoskeleton that can independently tune membrane rheology (*30, 31*).

The observation that non-native proteins (CRY2, RitC multimers) are displaced from sites of secretion raises the question of whether secretion site-derived bulk membrane flows also pattern the localization of native proteins. We have focused here on the Cdc42 GAP Rga4, showing that a short fragment containing elements of oligomerization (cc) and membrane binding (CBD) displays localization consistent with flow coupling. Thus, membrane flows are likely to be an important factor to pattern the native Rga4 protein. Additional factors may influence Rga4 localization. For instance, previous work proposed that the lateral localization of Rga4 is due to an exclusion mechanism controlled by the Pom1 kinase (*27*). The documented interaction of Rga4 with Pom1, which accumulates at cell poles, may indeed provide an explanation for the cell pole enrichment of cc-CBD fragments with low membrane affinity (*27, 32*). However, we note that membrane flows can be sufficient to explain Rga4 exclusion from growing cell ends, without invoking a Pom1-dependent exclusion mechanism.

Our experiments with the artificial Cdc42 optoGAP demonstrate that bulk membrane flows can be harnessed to pattern and shape the cell. These experiments provide a striking example of cell shape regulation by a simple change in protein localization. Indeed, the optoGAP is constitutively active whether cells are grown in dark or light conditions, as shown by its effect on the localization of the Cdc42-GTP CRIB reporter, but its localization is dramatically different. In the dark, optoGAP localizes indiscriminately to the entire plasma membrane, preventing Cdc42 activity, leading to sick, round cells. In the light, CRY2 oligomerization leads to increased membrane residence time, coupling optoGAP to membrane flows, which displace it from sites of secretion. This creates a local depletion of Cdc42 inhibition, allowing enhancement of activity and establishment of polarized growth. We note that strong oligomerization of immobile optoGAP in the non-growing part of the cell likely titrates a large fraction of optoGAP, explaining the very low signals along the newly grown cell portion. Because Cdc42 activity promotes secretion, which creates flows that displace the Cdc42 inhibitor to lateral regions, this results in the formation of Cdc42 activity zones of defined dimension (Fig 6I). Thus, membrane flows underlie a negative feedback that shapes the cell.

Membrane flows may pattern a variety of membrane-associated factors. Besides Rga4, membrane flows may promote the lateral localization of other proteins that organize the yeast cell shape and division, and explain their exclusion from both cell poles in wildtype cells, as well as from the only growing pole in monopolar mutants (for instance Rga6, Mid1, Blt1, Cdr2, Skb1 (*16, 27, 33-37*)). Establishment of daughter cell identity in *S. cerevisiae* through exocytosis-mediated septin ring formation is likely to belong to the same class of morphogenetic processes (*38*). Membrane flows may also help couple endocytosis around sites of exocytosis, which also promote polarity establishment (*39, 40*). Other proteins localize to secretion sites despite membrane flows, thanks to fast membrane unbinding, directed cytoskeletal transport or preferred affinity to static components. For instance, pole-enriched Pom1 shows association to microtubule-transported factors and very fast membrane

unbinding dynamics (*32, 41*). The Cdc42 GTPase also has to resist dilution from secretion-derived membrane addition, through rapid turnover at the membrane and likely other mechanisms (*42*).

Protein patterning via membrane lipid flow is likely to take place in all eukaryotic cells, in parallel to other patterning processes, such as actin cortex-dependent transport in animal cells. In yeast cells, membrane flow away from cell tips occurs because the membrane secreted by polarized exocytosis associated with net growth far exceeds that needed for growth and is recovered by endocytosis over a broader zone. The excess membrane delivery is about 4 to 5-fold according to our estimates (Table 1) and other measurements made in a recent study (*43*). This is likely also the case in tip-growing fungi and pollen tubes (*44*). More broadly, long range membrane flow should be a general phenomenon occurring even in cells maintaining their size, whenever regions of exocytosis and endocytosis are spatially segregated or when they overlap but have unequal widths. They may thus accompany cellular processes ranging from the formation of neuronal connections, cilia and immunological synapses (*45, 46*) to cell migration and cytokinesis (*47, 48*).

Our simulations and experiments clearly demonstrate how the dynamic properties of membrane-associated proteins are the determining factor for membrane-flow patterning. Membrane proteins that couple to lipid flow will spatially segregate if they remain associated on the membrane for a sufficiently long time and their diffusion is sufficiently slow relative to the flow rate and length scale involved, such that transport dominates diffusive spreading. We thus found that protein depletion at tips of yeast cells only occurs for proteins with slow enough membrane detachment and diffusion rates. Animal and plant cells, like yeast cells, likely harbour membrane-associated proteins that couple to flows and are displaced from sites of secretion. While transmembrane proteins in yeast cells are well known to exhibit particularly slow diffusion rates (perhaps due to interaction with the static cell wall (*26*), which may also hinder their movement along with lipid flow), the lateral diffusion of proteins associated with the inner leaflet (for instance Rho/Ras-family proteins) is only about 2 to 4-fold slower in yeast than animal cells (*18, 49-51*). Some membrane proteins in animal cells can have lateral diffusion coefficients significantly reduced upon tandem linkage or oligomerization (*52, 53*), compatible with flow coupling. Because protein localization can be altered by a simple change in membrane unbinding rates, as shown for instance by the acute localization change of CRY2 complexes, similar changes may occur in physiological conditions through post-translational modifications that modulate membrane affinity and/or oligomerization.

Materials and Methods

<u>Model methods</u>

Simulations of effects of exocytosis and endocytosis on protein distributions.

To investigate conditions that lead to depletion of proteins from a growing cell tip, we simulate the motion of non-interacting particles moving along a surface representing the plasma membrane under the effects of membrane binding/unbinding, diffusion, and membrane delivery/removal by exocytosis/endocytosis. The biophysical assumptions of our model are presented together with the model implementation below. For simplicity, the

particles in these simulations are restricted to the surface of a sphere with radius 1.8 μm with a fixed point on the sphere surface specified to be the growing tip. All arc lengths are measured with respect to this fixed point.

Time Step. The simulation was evolved through time primarily by using the Gillespie algorithm (*54*). An event is chosen from the list of possible events that include particle association and dissociation from the membrane, endocytosis, and exocytosis (excluding diffusion). The probability of a specific event occurring is given by the ratio of the event rate to the sum of the rates of all possible events (i.e. $P_i = r_i / \sum r_j$ where r_i is the rate of event i). The time until the selected event occurs is calculated as $\Delta t = -\ln(u') / \sum r_j$ where u' is a uniform random number between 0 and 1.

Diffusion. Particle diffusion, with diffusion coefficient *D*, is implemented with Brownian dynamics. The magnitude of the displacement due to diffusion over time Δt is calculated as $\sqrt{2D\Delta t}\xi(t)$ independently in both directions in the plane tangent to the sphere, and $\xi(t)$ is a random Gaussian distributed variable with zero mean and unit variance. After calculating the displacement in the tangent plane, the positions of each particle are then projected down to the sphere surface while keeping the distance conserved to correctly describe diffusion on a spherical surface (*55*). Since particles are assumed to diffuse independently of each other, we only calculated the displacement due to diffusion prior to endocytosis or exocytosis events, or before saving the data (except for a model with particle dimerization described below). We ensured that this implementation of diffusion on the sphere surface reproduced the correct scaling behaviour for the angular diffusion (*55*) by measuring the change in the polar angle, $\langle \theta(t)^2 \rangle$, as a function of time up to a time step of 1000 s as shown in Fig S9A.

Membrane binding and unbinding. Particles on the sphere undergo an unbinding event with rate k_{off} where they are removed from the sphere surface and placed in an internal pool representing the cytoplasm. Particles in this internal pool are placed back on the surface anywhere with uniform probability with rate k_{on}. This process assumes a sufficiently fast cytoplasmic diffusion to ensure uniform cytoplasmic concentration. We fix $k_{on} = 0.01\ \text{s}^{-1}$ since within our independent particle approximation this rate should only affect the average number of particles on the surface but not the relative level of tip depletion at steady state. To calculate smooth concentration profiles, there are on average approximately 50,000 particles on the surface at steady state in the simulations.

Particle movement by exocytosis and endocytosis. We assume that exocytosis and endocytosis near cell tips create a local flow that displaces lipids and proteins around the site of the vesicle event (*42, 56*). Our assumption is supported by the appearance of a flat plasma membrane in electron micrographs of fission yeast cell at cell sides and tips, except at sites of vesicle traffic or around eisosomes (*57-59*), similar to *S. cerevisiae* (*59, 60*). This observation indicates the presence of a membrane tension, which, combined with turgor pressure, can flatten the vesicle membrane delivered by exocytosis (*61, 62*), causing lateral flow. Similarly, flow of lipids towards endocytic vesicles must occur over a sufficiently wide area, preventing membrane strains larger than a few % that would otherwise lead to formation of membrane pores (*63*).

We simulate the movement associated with exocytosis as occurring instantaneously (i.e. no diffusion or membrane binding/unbinding occurs during the movement of the particles for either exocytosis or endocytosis). Our simulations are technically an area-change model as they do not explicitly include hydrodynamics, considered in many prior membrane models,

such as (*64*). Our rationale for this choice is that unlike single-component membranes, cellular membrane hydrodynamics remain unresolved (*65*), likely resembling those of a two dimensional multiphase fluid in the presence of obstacles, such as immobile transmembrane proteins connected to the cell wall (*66*). We thus judged these complexities would not justify the assumptions and additional computational complexity of a full hydrodynamic model.

We do however consider a few key hydrodynamic parameters. Firstly, similar to animal cells, we anticipate a fraction of the membrane, as high as 50%, to be immobile, with the fluid component undergoing Darcy-type flow (*64, 65, 67*). The fraction α of the fluid component of the membrane (that could be heterogeneous, containing lipids and proteins with different diffusion coefficients), is a parameter that we vary in the model. Peripherally-associated membrane proteins whose viscous drag is determined by the lipid membrane are assumed to be physically and hydrodynamically coupled to this lipid flow. To account for proteins whose drag and diffusion coefficient is influenced by interactions with structures that remain stationary or less mobile with respect to lipid flow, such as in the cell wall or in the cytoplasm, we introduce a parameter γ measuring the ratio of lipid to protein velocity. Unless otherwise indicated, we assume $\gamma = 1$. To simulate the effect of slow equilibration of membrane tension over long distances (*31*), we also introduce a cutoff distance to the flow induced by individual exocytosis and endocytosis events. As we show below, our results on tip depletion are robust with respect to the precise values of the mobile fraction α, velocity ratio γ, and cutoff distance.

In the simulation, exocytosis and endocytosis events take place on the surface of the sphere and affect the positions of nearby particles. The location of these events is selected randomly from an experimentally determined distribution (see below). Lipids on the surface at arc length distance s from the event are assumed to be pushed away radially from the center of exocytosis as shown in Fig 2B by arc length distance $\Delta r(s)$. It is assumed that the difference between the spherical cap surface area corresponding to mobile lipids, A_{cap}, measured at the final and initial arc length positions of the particle from the center of exocytosis equals the surface area of the exocytotic vesicle A_{exo}:

$$A_{cap}(s + \Delta r(s)) - A_{cap}(s) = A_{exo}, \qquad A_{cap}(s) = \alpha \, 2\pi R^2 \left[1 - \cos\left(\frac{s}{R}\right)\right], \qquad (M1)$$

where s is the arc length distance, R is the radius of the sphere, and parameter α describes the fraction of mobile proteins and lipids between 0 (no mobility) and 1 (all mobile). This gives for the displacement of membrane proteins:

$$\Delta r_{exo}(s) = \frac{1}{\gamma}\left\{R' \, acos\left[1 - \left(\frac{R}{R'}\right)^2 \left[1 - \cos\left(\frac{s}{R}\right)\right] - \frac{A_{exo}}{2\pi R'^2 \alpha}\right] - s\right\}, \quad R' = \sqrt{R^2 + A_{exo}/4\pi}, \qquad (M2)$$

where R' is the radius the sphere would have to accommodate the membrane area contributed by the exocytotic vesicle. The parameter γ describes the coupling between the proteins and the flowing membrane with higher values correlating with weaker coupling. The displacement is only applied to particles up to an upper arc length distance, $s < R_{cutoff}$, beyond which we assume the effects of membrane delivery are not felt immediately. The small difference between R and R' was introduced to avoid artificial accumulation of particles at the opposite sphere end in the absence of a cutoff R_{cutoff} but is otherwise insignificant for the simulations with a cutoff distance smaller than the arc length corresponding to the half-sphere. For an endocytosis event, particles within the cutoff distance R_{cutoff} are moved radially towards the endocytic point, similar to the calculation for exocytosis, with displacement magnitude:

$$\Delta r_{endo}(s) = \frac{1}{\gamma}\left\{-R'\ acos\left[1-\left(\frac{R}{R'}\right)^2\left[1-cos\left(\frac{s}{R}\right)\right]+\frac{A_{endo}}{2\pi R'^2\alpha}\right]+s\right\}\ ,\ \ R'=\sqrt{R^2-A_{endo}/4\pi},\quad \text{(M3)}$$

where A_{endo} is the surface area of the endocytic vesicle. When the displacement magnitude $\Delta r_{endo}(s)$ calculated by Equation (M3) is larger than s, particles are moved to the endocytosis event point, for the simulations where endocytosis does not contribute to internalization; for the model with endocytic removal they are internalized into the internal pool of particles.

The dependence of the model on parameters α and γ occurs approximately through their product, $\alpha\gamma$, as the distance proteins are displaced by exocytosis is $\Delta r_{exo} \approx A_{exo}/[\alpha\gamma 2\pi R|sin(s/R)|]$, and similarly for endocytosis, as long as $s > 0.25$ µm and $\alpha > 0.1$ for the reference parameters of Table 1.

Distribution of exocytosis and endocytosis. The distribution for exocytosis/endocytosis is based on the experimental distribution as function of arc-length distance from the cell tip, $P_{exp}(s)$, measured by imaging the profile along a medial confocal plane. The distribution $P_{exp}(s)$ can be well described by a Gaussian of standard deviation σ_{exo} and σ_{endo} for exocytosis and endocytosis, respectively. To convert this two dimensional distribution to a three dimensional one, we assume that the experimentally observed profiles account for events within a slab of width on the order of the microscope's resolution along the axis perpendicular to the glass slide ($b = 0.71$ µm) (Fig S9B). Thus, assuming cylindrical symmetry, the distribution of an event at arc length s from the position defined as cell tip used in simulations is calculated as

$$P(s) = C\,P_{exp}(s)f(s),\ \ \text{where}\ \ f(s)=\begin{cases}1, & s\leq R\,asin\left(\frac{b}{2R}\right)\\ \pi/\left[2\,asin\left(\frac{b}{2\,Rsin(s/R)}\right)\right], & R\,asin\left(\frac{b}{2R}\right)<s\leq\frac{\pi R}{2}\\ \pi\sin\left(\frac{s}{R}\right)/\left[2\,asin\left(\frac{b}{2R}\right)\right], & s>\frac{\pi R}{2}\end{cases}\quad\text{(M4)}$$

where $P_{exp}(s)$ is the experimental 2D probability distribution, $P(s)$ is the adjusted 3D probability distribution used in simulations, C is a normalization constant and $f(s)$ is the fraction of the circumference of the circle at a given value of s outside the slab region (i.e. thin line region of the circle circumference divided by the thick line region in Fig S9B). The expression for $f(s)$ initially assumes that for $s > \pi R/2$ the shape is that of a spherocylinder (as this is the shape of the actual cell) which is then mapped back onto a sphere by keeping the density of events constant by multiplying by $\sin(s/R)$. Agreement between experimental and simulated $P(s)$ considering only the slab region is shown in Fig S9C. The azimuthal angle for the exocytotic or endocytic event is picked from a uniform distribution.

Rates of membrane area delivery by endocytosis and exocytosis. We calibrated the model based on measurements performed as described in the supplemental text of Experimental Methods.

Firstly, the measured average rate of old end tip growth was 0.0305 ± 0.0151 µm/min (n = 75 cells; average $\pm$ stdev), which, multiplying by $2\pi R$ corresponds to net membrane area added of 0.345 µm^2/min = 20.7 µm^2/hr.

Secondly, we measured the rates of endocytosis at growing old end tips by counting the number of Fim1 patches in a medial section and extrapolating to the whole cell tip area. Please see experimental details in the supplemental file. Since the Gaussian profile of the endocytosis distribution extends beyond the hemispherical tip region, we considered separately the medial section rates at the hemi-spherical region (arclength from tip 0 to

$\pi R/2$, with 24.35 ± 10.56 events/min) and cylindrical segment (arclength $\pi R/2$ to πR, with 5.26 ± 3.85 events/min counting both sides). The fraction of events in a medial section of the hemi-spherical region would be $\int_0^{\pi R/2} P_{exp}(s)ds / \int_0^{\pi R/2} f(s)P_{exp}(s)ds$. This ratio is 0.233 using the measured σ_{endo}, leading to 104.53 endocytosis events/min for the whole hemi-spherical region. The fraction of cylindrical area imaged in a medial section is $b/(\pi R) = 0.126$, leading to 41.62 endocytosis events/min in the cylindrical region around the cell tip. Summing the numbers for hemi-spherical and cylindrical regions provides a rate of $r_{endo} = 146.2$ /min = 2.44 /s as listed on Table 1. This corresponds to endocytosed membrane area of 0.939 μm^2/min, using the estimate of endocytic vesicle radius in Table 1.

Thirdly, we estimated the rates of exocytosis by assuming that the rate of membrane area delivered by exocytosis is equal to the sum of net area added due to growth plus the membrane area endocytosed (sum = 1.28 μm^2/min). Using the estimate of exocytic vesicle radius in Table 1 leads to r_{exo} = 40.9 /min = 0.68 /s as listed on Table 1.

Measurement of tip occupancy ratio and reference parameters. The tip occupancy ratio is calculated by dividing the density of particles at an arc length of $\pi R/4$ = 1.41 μm (a quarter of half sphere total tip arc length) away from the top tip of the sphere by the density of particles in the remaining area of the sphere. The reference parameter values are listed in Table 1, at the bottom of this section.

Effect of parameters k_{on}, cut-off distance R_{cutoff}, mobile fraction α, and velocity ratio γ. Results presented in the main figures and Fig S2 used values of $k_{on} = 0.01 \text{ s}^{-1}, \alpha = 0.5, R_{cutoff} = 2$ μm. We also examined the effect of varying these parameters. As expected, the steady state reached is independent of k_{on} (Fig S9D). Decreasing the value of R_{cutoff} to 1.0 μm decreases overall depletion and lateral peak magnitude, shifting the peak towards the left, while increasing R_{cutoff} to 2.5 μm reverses the trend (Fig S9E). We find that decreasing α to a value of 0.1 (low mobile fraction) results in larger depletion and higher lateral peak, while increasing α to a value of 1.0 (highest mobile fraction) reverses the trend (Fig S9F). Increasing the value of γ to 10.0 (10-fold lower protein flow compared to lipid flow) results in decreased lateral peak and a decreased tip exclusion compared to same conditions with γ =1.0 (Fig S9F).

Dimerization simulations. To investigate the effect that oligomerization has on the extent of tip depletion and lateral peak height, we performed additional simulations in which we allow surface particles representing proteins to bind together to form dimers (Fig S5A). We did not explicitly include the effects of oligomerization in our main simulations (Fig 2) where the values of D and k_{off} measured from FRAP can be thought to reflect average behavior from individual particle to aggregate. Here we show that the binding of particles together into multimers further enhances the depletion and lateral peak, even without a change in the diffusion coefficient or membrane dissociation rate from the multimer. To avoid artificial concentration gradients as a result of implicit energy input to the system, we implemented a scheme designed to satisfy detailed balance in equilibrium, in the absence of exocytosis and endocytosis. In addition to association and dissociation of particles between cytoplasm and surface representing the plasma membrane (reaction 1 in Fig S5A with rate constants k_{on}, k_{off}), dimers can be formed in two ways: either two surface particles can bind reversibly to form a dimer (reaction 2 in Fig S5A with rate constants k_+, k_-) or a particle from the bulk cytoplasm and a surface particle can bind reversibly to form a dimer (reaction 3 in Fig S5A with rate constants $k_{on,2}, k_{off,2}$). While two particles are bound in the dimer state

they have the same position and they move together under diffusion, exocytosis, and endocytosis. To distinguish between the effects of oligomerization versus diffusion, as well as for simplicity, we used the same diffusion coefficient for surface monomers and dimers.

With rate constants in units of inverse time, detailed balance in absence of exocytosis and endocytosis requires $k_{on}N_b = k_{off}N_{sm}$, $k_+N_{sm}^2 = k_-N_{sd}$, $k_{on,2}N_bN_{sm} = k_{off,2}N_{sd}$, where N_b is the number of bulk particles in the cytoplasm, N_{sm} is the number of surface bound particles as monomers, and N_{sd} is the number of surface particles in dimers. We also have $N_b + N_{sm} + N_{sd} = N_{tot}$, where N_{tot} is the total number of particles which is constant throughout the simulation. Simultaneously solving these four equations results in expressions for the number of particles in each state in equilibrium, $N_{b,eq}$, $N_{sm,eq}$, and $N_{sd,eq}$ and a fourth equation, the detailed balance condition, for the reaction rate constants which can be written as $k_+ = k_{on,2} k_- k_{off}/(k_{off,2} k_{on})$. In these simulations we kept $k_{on} = 0.01 \text{ s}^{-1}$.

To investigate the effect of dimerization on tip depletion, we then turned on exocytosis and endocytosis. We define a parameter describing the effect of cooperativity, which is the ratio of equilibrium rate of particle association to the surface through dimer formation (reaction 3) versus the equilibrium rate of bulk cytoplasm particle attaching to the surface as monomer (reaction 1): $p = k_{on,2}N_{b,eq}N_{sm,eq}/(k_{on}N_{b,eq}) = k_{on,2}N_{sm,eq}/k_{on}$. The value of p thus implicitly defines the value of rate constant $k_{on,2}$, as well as that of k_+ through detailed balance when $k_{off}, k_{off,2}$, and k_- are given.

All reactions, except dimer formation through surface monomers, were handled with the Gillespie algorithm, similar to the case without dimerization described earlier. When a cytoplasmic particle binds to the surface to form a dimer, it is placed at the same position as a randomly-picked surface monomer particle. In the reverse reaction, one of the two particles in the dimer is selected to become a cytoplasmic particle while the other maintains its original position.

To handle the reaction of two surface particles binding together to form a dimer we used the method implemented in Smoldyn (*68*), where particles associate when separated by a distance less than a binding radius, σ_b, and are placed at an unbinding radius, σ_u, upon unbinding into two surface monomers, using $\sigma_b = \sigma_u$. To determine σ_b we followed the 3D reaction scheme of Smoldyn as a guide (where σ_b is determined based on the timestep, diffusion coefficient, and k_+), however as reactions in 2D involve additional complexities (*69, 70*), we adjusted σ_b by explicitly checking that detailed balance is satisfied in simulations without endocytosis or exocytosis. Values of σ_b ranged from $5\cdot10^{-5}$ to 0.05 µm as p was varied. The association reactions were checked over a fixed time interval $\Delta t_{fixed} = 0.01 \text{ } s$. Diffusion of surface particles was implemented every Δt_{fixed} and prior to any endocytosis or exocytosis event. Dimerization between a pair of particles occurs if they are at a distance less than or equal to σ_b at the end of such a timestep. The unbinding reaction of a dimer into two surface particles is handled by the Gillespie algorithm and the rate constant k_- where the particles are placed a distance of σ_u away from each other with random orientation. Due to the additional computational costs associated with calculating distances between particle pairs, we reduced the total number of particles to $N_{tot} = 5000$ and simulations were run for 10,000 s.

We find that as p increases (keeping $k_{off} = k_- = k_{off,2}$), the number of dimers at steady state increases at the expense of monomers on the surface and cytoplasm particles (Fig S5B). The tip depletion and height of lateral peak also increases as a function of p for fixed values

of D and k_{off}, when these were taken to be close to those of CRY2+CIBN-RitC (Fig S5C-E). This result shows that dimerization, and more generally, oligomerization, can lead to an enhancement of tip depletion and lateral peak. A decrease of the dimer dissociation rates $k_- = k_{off,2}$ compared to the monomer k_{off} further enhances this effect (Fig S5F-H).

Determination of membrane diffusion and dissociation rates by FRAP

Diffusion coefficients and membrane dissociation constants were determined by fitting FRAP recovery profiles to a 1D model (Fig S4). The method relies on the different effects of diffusion and association/dissociation on the recovery, with diffusion providing smoothing of profile over time (*71*) and association/dissociation contributing to uniform recovery. Detailed methods are described in the supplementary text.

Experimental materials and methods

Media and growth conditions

For microscopy experiments, cells were grown to exponential phase in Edinburgh minimal medium (EMM) supplemented with amino acids as required, unless indicated otherwise. For experiments in Fig 1, Fig 3, Fig S1 and Fig S3, cells were first precultured in 3 ml of EMM in dark conditions at 30 °C for 6–8 h, exponentially growing precultures were diluted (optical density [O.D.]$_{600nm}$ = 0.02) in 10 ml of EMM and incubated in dark conditions overnight at 30 °C, except for *ypt3-i5* and control strains that were cultured at 25 °C in YE. To allow proper aeration of the culture, 50 ml Erlenmeyer flasks were used. For experiments in Fig 4-6, Fig S4 and Fig S6-8, *S. pombe* cells were grown at 25°C. All live-cell imaging was performed on EMM-ALU agarose pads, except for the experiment shown in Fig 1E and 4E, for which lectin coated 96-well plates were used.

The control CRY2-mCherry-RitC was transformed in the triple mutant background (*rga3Δ rga4Δ rga6Δ*) and in wild type cells (Fig 6A). For the triple mutant, the strain was cultured either continuously in light or in dark conditions for 72hrs prior imaging. In the wild type background, the strain was cultured under dark conditions and light exposure was performed at the microscope using 488nm laser over 3 Z-stack planes to trigger CRY2 oligomerization over 15min time period. The optoGAP strain (Fig 6) was generated and grown continuously in light conditions (details on strain generation below). To compare cell morphologies in light and dark conditions (Fig 6B-E), the strains were grown on YE plates over 2 days at 30°C in the presence of light. Cultures in EMM-ALU were then prepared and split in incubators with or without light and diluted over 4 days to maintain exponential growth until imaged. The strains grown in the dark were not exposed to any light source until the moment of imaging. From these cultures, half of the cell material was used for AiryScan imaging and half was used for Calcofluor imaging. To study the dark-to-light transition (Fig 6G), cultures grown in the dark as above were shifted to light conditions at the microscope during live-cell imaging performed on EMM-ALU 2%-agar pads using blue light exposure at 5 focal planes through the cell depth every 2 minutes. For the drop-test assay (Fig 6F), cultures were grown in the light as above, adjusted to OD$_{600}$ = 0.3, subjected to 5x serial dilutions, and 5µl of each dilution was spotted on YE plates and grown at 30°C under light or dark conditions.

Strain construction

Strains used in this study are listed in Table S1. Standard genetic manipulation methods for *S. pombe* transformation and tetrad dissection were used. Gene tagging was performed at endogenous genomic locus at the 3' end, yielding C-terminally tagged proteins, as described (*72*). GFP-Ypt3 protein was ectopically expressed in addition to the endogenous gene by transforming a WT strain with PmeI-linearized pBSII(KS⁺)-based single integration vector (pSM2366-3'UTR*ade6*-GFP-Ypt3-hphMX-5'UTR*ade6*) targeting the *ade6* locus (*73*). Gene tagging, gene deletion, and plasmid integration were confirmed by diagnostic PCR for both sides of the insertion. The construction of plasmids and strains expressing the CRY2-CIB1 optogenetic system was done as described in (*2*).

To generate the RitC strain series, a synthetic gBlock DNA fragment was ordered (Integrated DNA Technologies, Inc.) containing four RitC repeats separated by linker sequences with restriction sites for the generation of 1x, 2x, and 3xRitC constructs. For all ordered synthetic gBlocks with repeat sequences codon usage was modified between repeats to allow successful synthesis. The RitC constructs are N-terminally tagged with sfGFP through sub-cloning in the stable integration vector (S.I.V.) backbone pAV0327 for *ura4* integration under expression from *act1* promoter and with *tdh1* terminator ((*73*); Addgene ID: 133488). Plasmid numbers and descriptions are in Table S2. Plasmids maps are available upon request. 1xRitC was additionally tagged N-terminally with mCherry in the S.I.V. pAV607 for *ade6* locus integration ((*73*); Addgene ID: 133504). The 3xsfGFP-1xRitC tagging was performed on pAV327 backbone carrying one sfGFP sequence and subcloning of a synthetic gBlock with 2 additional sfGFP sequences and the RitC sequence. Plasmids were integrated as described (*73*). Strains with either mCherry-1xRitC or CRIB-3xmCherry along with the respective sfGFP tagged RitC construct were generated through mating and tetrad dissection.

For the structure-function analysis of Rga4, fragments were amplified from the full-length gene originally amplified from cDNA using primers listed in Table S3, and first cloned in pSM621 (pREP41-GFP) for C-terminal tagging with GFP. All tagged fragments were subsequently amplified and introduced in S.I.V. for *ura4* single integration pAV327 in *S. pombe* with *act1* promoter driven expression and *tdh1* terminator (Fig 5C-E, S7C). The cc-CBD and cc-CBD-Δ2 fragments were additionally also expressed under a *pom1* promoter at the *ura4* locus to allow resolution of single clusters by TIRF imaging (Fig S8B-C). All integrations of AfeI linearized plasmids were performed in a *rga4Δ::kanMX* strain (YSM3826). Synthetic geneBlock DNA fragments were ordered for the generation of the multimerized 4xCBD and 4xcc-CBD-Δ2 constructs and introduced in S.I.V.s by In-Fusion cloning (Takara Bio Inc.) (Fig 5E). The CRY2-mCherry-CBD construct was also generated on a S.I.V. backbone for *ura4* integration and driven by *act1* promoter pAV0328 ((*73*); Addgene ID: 133489) (Fig S8A). Immediately after transformation, this strain was grown and stored only in dark conditions in the absence of any light source until the moment of imaging. Point-mutations in Rga4 membrane binding domains 1 and 2 were generated through several rounds of side-directed mutagenesis with primers listed in Table S3 for both fragments and full length Rga4. The mutagenized fragments were introduced in S.I.V.s and transformed at *ura4* locus as described above. The mutagenesis of the full length Rga4 was performed on a plasmid carrying the open reading frame and 112 base pairs of up- and downstream sequence. The entire sequence was excised (BamHI/SalI) and transformed in *rga4Δ::ura4+* background for integration at the endogenous locus, selected by growth on 5-FOA and verified by absence of growth on EMM-AL medium. The strains were genotyped and verified by sequencing for the presence of

mutations. Each mutagenized allele was subsequently tagged with GFP-natMX via standard PCR-based WACH approach (*72*) and additionally introduced in *rga3Δ::hphMX rga6Δ::kanMX* background through the standard *S. pombe* mating protocol and tetrad dissection.

The optoGAP was generated on a S.I.V. for *ade6* locus integration (pAV0356, ((*73*); Addgene ID: 133468)) with *act1* promoter driving the expression of CRY2-mCherry-RitC-rga4(aa665-933). Fragments were assembled by standard cloning. CRY2-mCherry-RitC control was generated by AscI/SacI digestion of the optoGAP backbone vector and re-introduction of RitC through standard cloning. The PmeI linearized plasmids were transformed in a *rga3Δ::hphMX rga4Δ::ura4+ rga6Δ::kanMX* strain (YSM3136; (*29*)) and in wild type background for the CRY2-mCherry-RitC control. The optoGAP strain was continuously grown in the presence of light at 30°C, unless otherwise indicated. We note that attempted transformation of these cells in the dark resulted in absence of colonies, suggesting toxicity as subsequently shown in the serial dilution assay (Fig 6F). This growth defect was absent for the CRY2-mcherry-RitC control. Both CRY2-RitC and the optoGAP strain (*rga3Δ::hphMX rga4Δ::ura4+ rga6Δ::kanMX ade6::pact1-CRY2-mCherry-RitC-rga4(aa665-933) ::ade6+*) were then transformed with pAV046 for CRIB-3xGFP-kanMX integration at *leu32* locus in light conditions.

BH-search prediction performed at https://hpcwebapps.cit.nih.gov/bhsearch/ with window size for residue averaging of 15 amino acids and values for amino acids set to standard BH parameters.

Microscopy

Spinning-disk microscopy

Images in Fig 1, Fig 3, Fig S1 and Fig S3 were acquired using a spinning disk confocal microscope, essentially as described (*74, 75*). Image acquisition was performed on a Leica DMI6000SD inverted microscope equipped with an HCX PL APO 100X/1.46 numerical aperture oil objective and a PerkinElmer Confocal system. This system uses a Yokogawa CSU22 real-time confocal scanning head, solid-state laser lines, and a cooled 14-bit frame transfer EMCCD C9100-50 camera (Hamamatsu) and is run by Volocity (PerkinElmer). When imaging strains expressing the CRY2-CIB1 optogenetic system, an additional long-pass color filter (550 nm, Thorlabs, Newton, New Jersey, USA) was used for bright-field (BF) image acquisition, in order to avoid precocious photoactivation caused by the white light. In *ypt3-i5* experiments, an objective heater (Bioptechs, Butler, PA, USA) was used for temperature control. Experiments in Fig 1, Fig 3 and Fig S1 were carried out using cell mixtures. Cell mixtures were composed of 1 sample strain of interest (the sample optogenetic strain co-expressing the indicated GFP-tagged protein) and up to 2 control strains namely:

1.   RFP control: An RFP bleaching correction strain, expressing cytosolic CRY2PHR-mCherry.
2.   Drug performance control: A strain expressing specific GFP or RFP markers tagging molecular components targeted by the drug employed in the specific experiment (see Fig S1F and S1I).

Strains were handled in dark conditions throughout. Red LED light was used in the room in order to manipulate strains, to prepare the agarose pads and lectin-coated 96-well plates. Strains were cultured separately. For experiments using agarose pads, exponentially growing cells (O.D.$_{600nm}$ = 0.4–0.6) were mixed in 2:1:1 (strain of interest, RFP control, and drug

performance control) ratio and harvested by soft centrifugation (2 min at 1,600 rpm). The cell mixture slurry (1 µl) was placed on a 2% EMM-ALU agarose pad, covered with a #1.5-thick coverslip, and sealed with vaseline, lanolin, and paraffin (VALAP). Samples were imaged after 5–10 min of rest in dark conditions. For experiments using lectin-coated 96-well plates, wells were first coated using 100 µl of soy bean lectin (L1395, Sigma-Aldrich) over night. Coated wells were washed 3 times using 200 µl of H_2O and dried for 4 h at room temperature. Exponentially growing cells (YE liquid media, O.D.$_{600nm}$ = 0.4–0.6) were mixed in 2:1:1 (strain of interest, RFP control, and drug performance control) ratio to a final O.D.$_{600nm}$ = 0.1. The cell mixture slurry (200 µl) was placed in lectin-coated wells and left for 45 min in the dark to allow cell-lectin attachment. Finally, cells were washed 3 times with 200 µl YE and 200 µl fresh YE liquid media was added for imaging.

CRY2PHR signal depletion at the cell tips and cell middle.

Spinning disk time-lapse images shown in Fig 1A, Fig S1A, Movie 1 and Movie 2 correspond to middle sections focal planes. Images were acquired every 30 s for GFP, RFP and bright-field channels. UV channel was acquired every 1.5 min in order to avoid photo-toxicity caused by UV light. Lasers were set to 100%; shutters were set to sample protection; and in all instances, the GFP channel was imaged first (for CRY2 photo-activation) and then the additional channels. RFP exposure time was set to 200 ms, whereas the GFP exposure was set to 1 s. Cells were monitored in these conditions for 20 min.

CRY2PHR tip depletion dynamic and cell-side accumulation dynamic analyses.

Strains co-expressing CRIB, Ypt3, Exo84 or Exo70 GFP-tagged proteins and the CRY2-CIB1 system were monitored in order to assess tip depletion dynamics. In these experiments an RFP control strain was added to the cell mixtures. Results of these experiments are shown in Fig 1B, Fig 3B-C, Fig S1B-E. Lasers were set to 100%; shutters were set to sample protection; and in all instances, the GFP channel was imaged first and then the RFP channel. RFP exposure time was always set to 200 ms, whereas the GFP exposure time varied depending on the monitored protein. Cells were monitored in these conditions for 285 s and images were acquired every 15 s.

Tip depletion dynamics upon disrupting exocytosis using Brefeldin A.

For Brefeldin A (BFA; B6542, Sigma-Aldrich) experiments shown in Fig 1C and Fig S1F-G cells co-expressing CRIB-3GFP or GFP-Ypt3 and the CRY2-CIB1 system were imaged. An RFP control strain and a strain expressing Anp1-GFP were mixed in this experiment. Anp1-GFP is a Golgi marker that in normal conditions forms prominent puncta (Fig S1F, left). Upon blocking of endomembrane trafficking due to BFA treatment, Anp1 puncta disappear and the protein re-localizes to the ER (Fig S1F, right). BFA (diluted in ethanol, final concentration 300 µM) was added to the cell mixture slurry. Cells were placed on top of 2% EMM-ALU + BFA 300 µM agarose pads to be imaged. Lasers were set to 100%; shutters were set to sample protection; and in all instances, the GFP channel was imaged first and then the RFP channel. RFP exposure time was always set to 200 ms, whereas the GFP exposure time varied depending on the monitored protein. Cells were monitored in these conditions for 285 s and images were acquired every 15 s.

Tip depletion dynamics upon disrupting exocytosis using *ypt3-i5* ts allele.

For *ypt3-i5* experiments shown in Fig 1D and Fig S1H, *ypt3-i5* and *ypt3+* cells co-expressing CRIB-3GFP and the CRY2-CIB1 system were imaged. In these experiments an RFP control

strain was added to the cell mixtures. Cells were cultured at 25 °C throughout the experiment and placed on 2% EMM-ALU agarose pad sealed with VALAP. 2 sets of pads were prepared, one to be imaged at 25 °C and another one to be imaged at 36 °C. Prior to imaging, pads were incubated for 30 mins either at 25 °C or 36 °C and imaged at the respective temperatures by setting the objective heater accordingly. Lasers were set to 100%; shutters were set to sample protection; and in all instances, the GFP channel was imaged first and then the RFP channel. RFP exposure time was always set to 200 ms, whereas the GFP exposure was set to 1 s. Cells were monitored in these conditions for 285 s and images were acquired every 15 s.

Tip depletion dynamics upon disrupting endocytosis using Latrunculin A.

For Latrunculin A (LatA; BML-T119-0500; Enzo Life Sciences) experiments shown in Fig 1E and Fig S1H-I *sty1Δ* cells co-expressing CRIB-3GFP and the CRY2-CIB1 system were imaged on lectin-coated 96-well plates. An RFP control strain and a strain co-expressing Exo70-GFP and Fim1-mCherry were mixed in this experiment. The Fim1-mCherry actin patch marker serves to verify actin depolymerization due to LatA treatment (Fig S1I). LatA (diluted in DMSO, final concentration 50 μM) was added to 200 μl YE liquid media. Lasers were set to 100%; shutters were set to sample protection; and in all instances, the GFP channel was imaged first and then the RFP channel. RFP exposure time was always set to 200 ms, whereas the GFP exposure was set to 1 s. Cells were monitored in these conditions for 285 s and images were acquired every 15 s.

Cortical distribution profiles of exocytosis (Exo70-GFP) and endocytosis (Fim1-mCherry) and counting endocytic events at the cell cortex.

A strain expressing endogenous tagged Exo70-GFP and Fim1-mCherry was monitored in order to assess their cortical distribution at the cell tips. Results of these experiments are shown in Fig S3. Lasers were set to 100% and shutters were set to sample protection. Cells were monitored in these conditions for 300 s and images were acquired every 15 s (21 timepoints).

Airyscan imaging

All images in Fig 4-6, S4 and S6-8 (except S7B and S8B) were acquired on a Zeiss LSM 980 system with Plan-Apochromat 63x/1.40 Oil DIC objective and acquired by the ZEN Blue software (Zeiss). Imaging was set in super-resolution mode with frame bidirectional maximum speed scanning. Laser power was kept at < 0.4%, with pixel time around 6.6 μs and frame acquisition time <2s per channel. All other settings were optimized as recommended by the ZEN Blue software.

For the Latrunculin A (LatA; BML-T119-0500; Enzo Life Sciences) experiment shown in Fig 4E *sty1Δ* cells co-expressing sfGFP-3xRitC and CRIB-3xmCherry were imaged using lectin-coated 96-well plates. Cells were grown in YE at 25°C, LatA (diluted in DMSO, final concentration 50 μM) was added to 200 μl YE liquid media. Cells were imaged after 30min of treatment. LifeAct-GFP expressing strain was used as a control for LatA depolymerization.

In Fig 5A, z-stacks were set at optimal interval with 0.16 μm spacing over a total of 4μm from the medial plane of the cell. All other imaging shows medial focal planes. Frames for movie S5 and Fig 4D were acquired with 488nm and 561nm lasers every 5 min; frames for movies S8, S9 and Fig 6G were acquired with 488nm and 561nm lasers every 1min30sec and frames for movie S10 and Fig 6H every 2min.

FRAP was performed on cells on a microscope slide (maximum imaging time of 5 min) by finding the focal plane closest to the cover slip and defining a rectangular ROI in the middle

of the visible signal (see Fig S4A). 2 snapshots were taken prior to bleaching, which was done using 20% laser power and a 5 time repeat of the bleach zone. To determine the depth of the bleached signal immediately post-bleaching, z-stacks optimally spaced at 0.16µm interval were acquired across the entire depth of the cell, which allowed to calculate that bleaching occurs over a depth of 2.0µm-2.5µm. For 2xRitC and 3xRitC, recovery was imaged for 5 min, every 10s; 1xRitC and Rga4 fragments were imaged at maximum speed every 729ms for 1 min. For FRAP of CRY2-CIBN in the light-activated state, cells were first found in focus using the 561nm laser after which they were illuminated with transmitted white light for 45s and a snapshot of CRY2 to CIBN recruitment at the mid-plane of the cell was taken. The FRAP was then performed as described above at the focal plane closest to the cover slip over a 5 min recovery time, acquired every 10s, at each time point acquisition was performed first with 561nm laser, followed by 488nm laser for a total frame acquisition time of 900ms. The recovery for the FRAP of CIBN-GFP alone was recorded over 1 min, every 700ms maximum speed imaging. All AiryScan images were processed in Zeiss Zen Blue software.

Other imaging

For cell length measurements (Fig 6D) as well as cell morphology imaging Fig S7D, cells were stained with 2µg/ml final concentration calcofluor (Fluorescent brightener 28, F3543, Sigma) and imaged with a Leica epifluorescence microscope (63X magnification). TIRF microscopy (Fig S8B-C) was performed on a DeltaVision OMX SR imaging system, equipped with a 60x 1.49 NA TIRF oil objective (oil 1.514), an illumination pathway for ring-TIRF and a front illuminated sCMOS camera size 2560x2160 pxl (manufacturer PCO). Imaging settings were: 512x512 pxl field of view, 100ms exposure time, laser power of 20% with TIRF angle 488nm at 86.7°. Samples were placed on a 0.17 +/- 0.01 mm thick glass slide and imaged within 15 minutes. Imaging was performed every second over a 60s imaging period.

Image analysis of concentration profile

All image-processing analyses were performed with ImageJ software (http://rsb.info.nih.gov/ij/). Image and time-lapse recordings were imported to the software using the Bio-Formats plugin (http://loci.wisc.edu/software/bio-formats). Time-lapse recordings were aligned using the StackReg plugin (http://bigwww.epfl.ch/thevenaz/stackreg/) using the rigid body method. All optogenetic data analyses were performed using MATLAB (R2019a), with scripts developed in-house. Figures were assembled with Adobe Photoshop CC2019 and Adobe Illustrator 2020. Details of image quantification procedures are available in the supplemental text.

Statistical analysis

The Kruskal-Wallis test was used for comparison of CRY2 depletion for identical values of CRIB intensity in Fig 1 and Fig S1. The student t-test with unequal variance was used in Fig 4E to compare tip to side ratios of GFP signal and in Fig S8C. In Fig 6D, the student t-test with equal variance was used. Experiments were conducted in triplicates, except where indicated. For the exact number of cells in each experiment, please refer to the experimental details.

References

1. L. Kozubowski, K. Saito, J. M. Johnson, A. S. Howell, T. R. Zyla, D. J. Lew, Symmetry-breaking polarization driven by a Cdc42p GEF-PAK complex. *Curr Biol* **18**, 1719-1726 (2008); published online EpubNov 25 (

2. I. Lamas, L. Merlini, A. Vjestica, V. Vincenzetti, S. G. Martin, Optogenetics reveals Cdc42 local activation by scaffold-mediated positive feedback and Ras GTPase. *PLoS Biol* **18**, e3000600 (2020); published online EpubJan (10.1371/journal.pbio.3000600).

3. M. Das, T. Drake, D. J. Wiley, P. Buchwald, D. Vavylonis, F. Verde, Oscillatory Dynamics of Cdc42 GTPase in the Control of Polarized Growth. *Science*, (2012); published online EpubMay 17 (10.1126/science.1218377).

4. A. S. Howell, M. Jin, C. F. Wu, T. R. Zyla, T. C. Elston, D. J. Lew, Negative feedback enhances robustness in the yeast polarity establishment circuit. *Cell* **149**, 322-333 (2012); published online EpubApr 13 (10.1016/j.cell.2012.03.012).

5. R. Illukkumbura, T. Bland, N. W. Goehring, Patterning and polarization of cells by intracellular flows. *Curr Opin Cell Biol* **62**, 123-134 (2020); published online EpubFeb (10.1016/j.ceb.2019.10.005).

6. M. C. Wigbers, F. Brauns, C. Y. Leung, E. Frey, Flow Induced Symmetry Breaking in a Conceptual Polarity Model. *Cells* **9**, (2020); published online EpubJun 23 (10.3390/cells9061524).

7. G. Grebnev, M. Ntefidou, B. Kost, Secretion and Endocytosis in Pollen Tubes: Models of Tip Growth in the Spot Light. *Frontiers in plant science* **8**, 154 (2017)10.3389/fpls.2017.00154).

8. P. Rowghanian, O. Campas, Non-equilibrium Membrane Homeostasis in Expanding Cellular Domains. *Biophys J* **113**, 132-137 (2017); published online EpubJul 11 (10.1016/j.bpj.2017.06.001).

9. M. S. Bretscher, Endocytosis: relation to capping and cell locomotion. *Science* **224**, 681-686 (1984); published online EpubMay 18 (10.1126/science.6719108).

10. M. Tanaka, T. Kikuchi, H. Uno, K. Okita, T. Kitanishi-Yumura, S. Yumura, Turnover and flow of the cell membrane for cell migration. *Scientific reports* **7**, 12970 (2017); published online EpubOct 11 (10.1038/s41598-017-13438-5).

11. P. R. O'Neill, J. A. Castillo-Badillo, X. Meshik, V. Kalyanaraman, K. Melgarejo, N. Gautam, Membrane Flow Drives an Adhesion-Independent Amoeboid Cell Migration Mode. *Dev Cell* **46**, 9-22 e24 (2018); published online EpubJul 2 (10.1016/j.devcel.2018.05.029).

12. S. Gras, E. Jimenez-Ruiz, C. M. Klinger, K. Schneider, A. Klingl, L. Lemgruber, M. Meissner, An endocytic-secretory cycle participates in Toxoplasma gondii in motility. *PLoS Biol* **17**, e3000060 (2019); published online EpubJun (10.1371/journal.pbio.3000060).

13. J. Dai, M. P. Sheetz, Axon membrane flows from the growth cone to the cell body. *Cell* **83**, 693-701 (1995); published online EpubDec 1 (10.1016/0092-8674(95)90182-5).

14. F. D. Kelly, P. Nurse, Spatial control of Cdc42 activation determines cell width in fission yeast. *Mol Biol Cell* **22**, 3801-3811 (2011); published online EpubOct (mbc.E11-01-0057 [pii]10.1091/mbc.E11-01-0057).

15. M. Das, D. J. Wiley, S. Medina, H. A. Vincent, M. Larrea, A. Oriolo, F. Verde, Regulation of cell diameter, For3p localization, and cell symmetry by fission yeast Rho-GAP Rga4p. *Mol Biol Cell* **18**, 2090-2101 (2007); published online EpubJun (

16. M. T. Revilla-Guarinos, R. Martin-Garcia, M. A. Villar-Tajadura, M. Estravis, P. M. Coll, P. Perez, Rga6 is a Fission Yeast Rho GAP Involved in Cdc42 Regulation of Polarized Growth. *Mol Biol Cell*, (2016); published online EpubMar 9 (10.1091/mbc.E15-12-0818).

17. A. D. Migliori, L. A. Patel, C. Neale, The RIT1 C-terminus associates with lipid bilayers via charge complementarity. *Comput Biol Chem* **91**, 107437 (2021); published online EpubApr (10.1016/j.compbiolchem.2021.107437).

18. F. O. Bendezu, V. Vincenzetti, D. Vavylonis, R. Wyss, H. Vogel, S. G. Martin, Spontaneous Cdc42 polarization independent of GDI-mediated extraction and actin-based trafficking. *PLoS Biol* **13**, e1002097 (2015); published online EpubApr (10.1371/journal.pbio.1002097).

19. B. Onken, H. Wiener, M. R. Philips, E. C. Chang, Compartmentalized signaling of Ras in fission yeast. *Proc Natl Acad Sci U S A* **103**, 9045-9050 (2006); published online EpubJun 13

20. L. J. Bugaj, A. T. Choksi, C. K. Mesuda, R. S. Kane, D. V. Schaffer, Optogenetic protein clustering and signaling activation in mammalian cells. *Nat Methods* **10**, 249-252 (2013); published online EpubMar (10.1038/nmeth.2360).

21. T. G. Turi, P. Webster, J. K. Rose, Brefeldin A sensitivity and resistance in Schizosaccharomyces pombe. Isolation of multiple genes conferring resistance. *J Biol Chem* **269**, 24229-24236 (1994); published online EpubSep 30 (

22. M. Marek, V. Vincenzetti, S. G. Martin, Sterol biosensor reveals LAM-family Ltc1-dependent sterol flow to endosomes upon Arp2/3 inhibition. *J Cell Biol* **219**, (2020); published online EpubJun 1 (10.1083/jcb.202001147).

23. D. Bonazzi, J. D. Julien, M. Romao, R. Seddiki, M. Piel, A. Boudaoud, N. Minc, Symmetry Breaking in Spore Germination Relies on an Interplay between Polar Cap Stability and Spore Wall Mechanics. *Dev Cell* **28**, 534-546 (2014).

24. D. R. Mutavchiev, M. Leda, K. E. Sawin, Remodeling of the Fission Yeast Cdc42 Cell-Polarity Module via the Sty1 p38 Stress-Activated Protein Kinase Pathway. *Curr Biol* **26**, 2921-2928 (2016); published online EpubNov 7 (10.1016/j.cub.2016.08.048).

25. L. Duan, J. Hope, Q. Ong, H. Y. Lou, N. Kim, C. McCarthy, V. Acero, M. Z. Lin, B. Cui, Understanding CRY2 interactions for optical control of intracellular signaling. *Nat Commun* **8**, 547 (2017); published online EpubSep 15 (10.1038/s41467-017-00648-8).

26. J. Valdez-Taubas, H. R. Pelham, Slow diffusion of proteins in the yeast plasma membrane allows polarity to be maintained by endocytic cycling. *Curr Biol* **13**, 1636-1640 (2003); published online EpubSep 16 (S0960982203006584 [pii]).

27. H. Tatebe, K. Nakano, R. Maximo, K. Shiozaki, Pom1 DYRK regulates localization of the Rga4 GAP to ensure bipolar activation of Cdc42 in fission yeast. *Curr Biol* **18**, 322-330 (2008); published online EpubMar 11 (

28. H. Brzeska, J. Guag, K. Remmert, S. Chacko, E. D. Korn, An experimentally based computer search identifies unstructured membrane-binding sites in proteins: application to class I myosins, PAKS, and CARMIL. *J Biol Chem* **285**, 5738-5747 (2010); published online EpubFeb 19 (10.1074/jbc.M109.066910).

29. D. Gallo Castro, S. G. Martin, Differential GAP requirement for Cdc42-GTP polarization during proliferation and sexual reproduction. *J Cell Biol* **217**, 4215-4229 (2018); published online EpubDec 3 (10.1083/jcb.201806016).

30. A. Kusumi, T. K. Fujiwara, R. Chadda, M. Xie, T. A. Tsunoyama, Z. Kalay, R. S. Kasai, K. G. Suzuki, Dynamic organizing principles of the plasma membrane that regulate signal transduction: commemorating the fortieth anniversary of Singer and Nicolson's fluid-mosaic model. *Annu Rev Cell Dev Biol* **28**, 215-250 (2012)10.1146/annurev-cellbio-100809-151736).

31. Z. Shi, Z. T. Graber, T. Baumgart, H. A. Stone, A. E. Cohen, Cell Membranes Resist Flow. *Cell* **175**, 1769-1779 e1713 (2018); published online EpubDec 13 (10.1016/j.cell.2018.09.054).

32. O. Hachet, M. Berthelot-Grosjean, K. Kokkoris, V. Vincenzetti, J. Moosbrugger, S. G. Martin, A phosphorylation cycle shapes gradients of the DYRK family kinase Pom1 at the plasma membrane. *Cell* **145**, 1116-1128 (2011); published online EpubJun 24 (S0092-8674(11)00540-X [pii] 10.1016/j.cell.2011.05.014).

33. N. N. Padte, S. G. Martin, M. Howard, F. Chang, The cell-end factor pom1p inhibits mid1p in specification of the cell division plane in fission yeast. *Curr Biol* **16**, 2480-2487 (2006); published online EpubDec 19

34. S. G. Martin, M. Berthelot-Grosjean, Polar gradients of the DYRK-family kinase Pom1 couple cell length with the cell cycle. *Nature* **459**, 852-856 (2009); published online EpubJun 11 (

35. J. B. Moseley, A. Mayeux, A. Paoletti, P. Nurse, A spatial gradient coordinates cell size and mitotic entry in fission yeast. *Nature* **459**, 857-860 (2009); published online EpubJun 11

36. L. Deng, J. B. Moseley, Compartmentalized nodes control mitotic entry signaling in fission yeast. *Mol Biol Cell* **24**, 1872-1881 (2013); published online EpubJun (10.1091/mbc.E13-02-0104).

37. M. Guzman-Vendrell, S. Baldissard, M. Almonacid, A. Mayeux, A. Paoletti, J. B. Moseley, Blt1 and Mid1 provide overlapping membrane anchors to position the division plane in fission yeast. *Mol Cell Biol* **33**, 418-428 (2013); published online EpubJan (10.1128/MCB.01286-12).

38. S. Okada, M. Leda, J. Hanna, N. S. Savage, E. Bi, A. B. Goryachev, Daughter cell identity emerges from the interplay of Cdc42, septins, and exocytosis. *Dev Cell* **26**, 148-161 (2013); published online EpubJul 29 (10.1016/j.devcel.2013.06.015).

39. R. T. A. Pedersen, J. E. Hassinger, P. Marchando, D. G. Drubin, Spatial regulation of clathrin-mediated endocytosis through position-dependent site maturation. *J Cell Biol* **219**, (2020); published online EpubNov 2 (10.1083/jcb.202002160).

40. M. Jose, S. Tollis, D. Nair, J. B. Sibarita, D. McCusker, Robust polarity establishment occurs via an endocytosis-based cortical corralling mechanism. *J Cell Biol* **200**, 407-418 (2013); published online EpubFeb 18 (10.1083/jcb.201206081).

41. V. Gerganova, C. Floderer, A. Archetti, L. Michon, L. Carlini, T. Reichler, S. Manley, S. G. Martin, Multi-phosphorylation reaction and clustering tune Pom1 gradient mid-cell levels according to cell size. *eLife* **8**, (2019); published online EpubMay 3 (10.7554/eLife.45983).

42. A. T. Layton, N. S. Savage, A. S. Howell, S. Y. Carroll, D. G. Drubin, D. J. Lew, Modeling vesicle traffic reveals unexpected consequences for Cdc42p-mediated polarity establishment. *Curr Biol* **21**, 184-194 (2011); published online EpubFeb 8 (10.1016/j.cub.2011.01.012).

43. J. Lemière, Y. Ren, J. Berro, Rapid adaptation of endocytosis, exocytosis and eisosomes after an acute increase in membrane tension in yeast cells. *bioRxiv*, 342030 (2021)10.1101/342030).

44. G. Steinberg, M. A. Penalva, M. Riquelme, H. A. Wosten, S. D. Harris, Cell Biology of Hyphal Growth. *Microbiol Spectr* **5**, (2017); published online EpubApr (10.1128/microbiolspec.FUNK-0034-2016).

45. G. M. Griffiths, A. Tsun, J. C. Stinchcombe, The immunological synapse: a focal point for endocytosis and exocytosis. *J Cell Biol* **189**, 399-406 (2010); published online EpubMay 3 (10.1083/jcb.201002027).

46. J. Roos, R. B. Kelly, The endocytic machinery in nerve terminals surrounds sites of exocytosis. *Curr Biol* **9**, 1411-1414 (1999); published online EpubDec 2 (10.1016/s0960-9822(00)80087-1).

47. J. A. Castillo-Badillo, A. C. Bandi, S. Harlalka, N. Gautam, SRRF-Stream Imaging of Optogenetically Controlled Furrow Formation Shows Localized and Coordinated Endocytosis and Exocytosis Mediating Membrane Remodeling. *ACS Synth Biol* **9**, 902-919 (2020); published online EpubApr 17 (10.1021/acssynbio.9b00521).

48. K. S. Gerien, J. Q. Wu, Molecular mechanisms of contractile-ring constriction and membrane trafficking in cytokinesis. *Biophys Rev* **10**, 1649-1666 (2018); published online EpubDec (10.1007/s12551-018-0479-3).

49. V. Y. Kiselev, D. Marenduzzo, A. B. Goryachev, Lateral dynamics of proteins with polybasic domain on anionic membranes: a dynamic Monte-Carlo study. *Biophys J* **100**, 1261-1270 (2011); published online EpubMar 2 (10.1016/j.bpj.2011.01.025).

50. K. C. Vinnakota, D. A. Mitchell, R. J. Deschenes, T. Wakatsuki, D. A. Beard, Analysis of the diffusion of Ras2 in Saccharomyces cerevisiae using fluorescence recovery after photobleaching. *Phys Biol* **7**, 026011 (2010); published online EpubJun 4 (10.1088/1478-3975/7/2/026011).

51. B. Khalili, L. Merlini, V. Vincenzetti, S. G. Martin, D. Vavylonis, Exploration and stabilization of Ras1 mating zone: A mechanism with positive and negative feedbacks. *PLoS Comput Biol* **14**, e1006317 (2018); published online EpubJul (10.1371/journal.pcbi.1006317).

52. G. R. Hammond, Y. Sim, L. Lagnado, R. F. Irvine, Reversible binding and rapid diffusion of proteins in complex with inositol lipids serves to coordinate free movement with spatial information. *J Cell Biol* **184**, 297-308 (2009); published online EpubJan 26 (10.1083/jcb.200809073).

53. R. Iino, I. Koyama, A. Kusumi, Single molecule imaging of green fluorescent proteins in living cells: E-cadherin forms oligomers on the free cell surface. *Biophys J* **80**, 2667-2677 (2001); published online EpubJun (10.1016/S0006-3495(01)76236-4).

54. D. T. Gillespie, Exact stochastic simulation of coupled chemical reactions. *J. Phys. Chem.* **81**, 2340–2361 (1977)https://doi.org/10.1021/j100540a008).

55. P. Castro-Villarreal, A. Villada-Balbuena, J. M. Mendez-Alcaraz, R. Castaneda-Priego, S. Estrada-Jimenez, A Brownian dynamics algorithm for colloids in curved manifolds. *J Chem Phys* **140**, 214115 (2014); published online EpubJun 7 (10.1063/1.4881060).

56. A. W. McClure, M. Minakova, J. M. Dyer, T. R. Zyla, T. C. Elston, D. J. Lew, Role of Polarized G Protein Signaling in Tracking Pheromone Gradients. *Dev Cell* **35**, 471-482 (2015); published online EpubNov 23 (10.1016/j.devcel.2015.10.024).

57. H. Cheng, R. Sugiura, W. Wu, M. Fujita, Y. Lu, S. O. Sio, R. Kawai, K. Takegawa, H. Shuntoh, T. Kuno, Role of the Rab GTP-binding protein Ypt3 in the fission yeast exocytic pathway and its connection to calcineurin function. *Mol Biol Cell* **13**, 2963-2976 (2002); published online EpubAug (10.1091/mbc.01-09-0463).

58. N. Wang, I. J. Lee, G. Rask, J. Q. Wu, Roles of the TRAPP-II Complex and the Exocyst in Membrane Deposition during Fission Yeast Cytokinesis. *PLoS Biol* **14**, e1002437 (2016); published online EpubApr (10.1371/journal.pbio.1002437).

59. J. H. Lee, J. E. Heuser, R. Roth, U. Goodenough, Eisosome Ultrastructure and Evolution in Fungi, Microalgae, and Lichens. *Eukaryot Cell* **14**, 1017-1042 (2015); published online EpubOct (10.1128/EC.00106-15).

60. W. Kukulski, M. Schorb, M. Kaksonen, J. A. Briggs, Plasma membrane reshaping during endocytosis is revealed by time-resolved electron tomography. *Cell* **150**, 508-520 (2012); published online EpubAug 3 (10.1016/j.cell.2012.05.046).

61. L. V. Chernomordik, M. M. Kozlov, Mechanics of membrane fusion. *Nature structural & molecular biology* **15**, 675-683 (2008); published online EpubJul (10.1038/nsmb.1455).

62. W. Shin, G. Arpino, S. Thiyagarajan, R. Su, L. Ge, Z. McDargh, X. Guo, L. Wei, O. Shupliakov, A. Jin, B. O'Shaughnessy, L. G. Wu, Vesicle Shrinking and Enlargement Play

Opposing Roles in the Release of Exocytotic Contents. *Cell reports* **30**, 421-431 e427 (2020); published online EpubJan 14 (10.1016/j.celrep.2019.12.044).

63. M. Chabanon, J. C. S. Ho, B. Liedberg, A. N. Parikh, P. Rangamani, Pulsatile Lipid Vesicles under Osmotic Stress. *Biophys J* **112**, 1682-1691 (2017); published online EpubApr 25 (10.1016/j.bpj.2017.03.018).

64. B. Fogelson, A. Mogilner, Computational Estimates of Membrane Flow and Tension Gradient in Motile Cells. *Plos One* **9**, (2014); published online EpubJan 17 (ARTN e84524 10.1371/journal.pone.0084524).

65. A. E. Cohen, Z. Shi, Do Cell Membranes Flow Like Honey or Jiggle Like Jello? *Bioessays* **42**, (2020); published online EpubJan (ARTN 1900142 10.1002/bies.201900142).

66. K. D. Moran, D. J. Lew, How Diffusion Impacts Cortical Protein Distribution in Yeasts. *Cells* **9**, (2020); published online EpubMay (ARTN 1113 10.3390/cells9051113).

67. Z. Kalay, T. K. Fujiwara, A. Otaka, A. Kusumi, Lateral diffusion in a discrete fluid membrane with immobile particles. *Phys Rev E* **89**, (2014); published online EpubFeb 26 (ARTN 022724 10.1103/PhysRevE.89.022724).

68. S. S. Andrews, D. Bray, Stochastic simulation of chemical reactions with spatial resolution and single molecule detail. *Phys Biol* **1**, 137-151 (2004); published online EpubSep (Pii S1478-3967(04)83249-2 10.1088/1478-3967/1/3/001).

69. O. N. Yogurtcu, M. E. Johnson, Theory of bi-molecular association dynamics in 2D for accurate model and experimental parameterization of binding rates. *Journal of Chemical Physics* **143**, (2015); published online EpubAug 28 (Artn 084117 10.1063/1.4929390).

70. B. O'Shaughnessy, D. Vavylonis, Kinetic regimes and cross-over times in many-particle reacting systems. *Europhys Lett* **45**, 653-658 (1999); published online EpubMar 15 (DOI 10.1209/epl/i1999-00217-9).

71. J. Ries, C. Klose, C. Walch-Solimena, P. Schwille, How to measure slow diffusion in yeast cell membranes. *Proc Spie* **6991**, (2008)Artn 69910w 10.1117/12.787043).

72. J. Bahler, J. Q. Wu, M. S. Longtine, N. G. Shah, A. McKenzie, 3rd, A. B. Steever, A. Wach, P. Philippsen, J. R. Pringle, Heterologous modules for efficient and versatile PCR-based gene targeting in Schizosaccharomyces pombe. *Yeast* **14**, 943-951 (1998); published online EpubJul (10.1002/(SICI)1097-0061(199807)14:10<943::AID-YEA292>3.0.CO;2-Y).

73. A. Vjestica, M. Marek, P. J. Nkosi, L. Merlini, G. Liu, M. Berard, I. Billault-Chaumartin, S. G. Martin, A toolbox of Stable Integration Vectors (SIV) in the fission yeast Schizosaccharomyces pombe. *J Cell Sci*, (2019); published online EpubDec 4 (10.1242/jcs.240754).

74. F. O. Bendezu, S. G. Martin, Actin cables and the exocyst form two independent morphogenesis pathways in the fission yeast. *Mol Biol Cell* **22**, 44-53 (2011); published online EpubJan 1 (mbc.E10-08-0720 [pii] 10.1091/mbc.E10-08-0720).

75. O. Dudin, F. O. Bendezu, R. Groux, T. Laroche, A. Seitz, S. G. Martin, A formin-nucleated actin aster concentrates cell wall hydrolases for cell fusion in fission yeast. *J Cell Biol* **208**, 897-911 (2015); published online EpubMar 30 (10.1083/jcb.201411124).

76. B. P. Ziemba, J. D. Knight, J. J. Falke, Assembly of Membrane-Bound Protein Complexes: Detection and Analysis by Single Molecule Diffusion. *Biochemistry* **51**, 1638-1647 (2012); published online EpubFeb 28 (10.1021/bi201743a).

77. J. D. Knight, M. G. Lerner, J. G. Marcano-Velazquez, R. W. Pastor, J. J. Falke, Single Molecule Diffusion of Membrane-Bound Proteins: Window into Lipid Contacts and Bilayer Dynamics. *Biophysical Journal* **99**, 2879-2887 (2010); published online EpubNov 3 (10.1016/j.bpj.2010.08.046).

78. B. A. Camley, F. L. H. Brown, Motion of objects embedded in lipid bilayer membranes: Advection and effective viscosity. *Journal of Chemical Physics* **151**, (2019); published online EpubSep 28 (Artn 124104 10.1063/1.5121418).

79. J. R. Houser, D. J. Busch, D. R. Bell, B. Li, P. Y. Ren, J. C. Stachowiak, The impact of physiological crowding on the diffusivity of membrane bound proteins. *Soft Matter* **12**, 2127-2134 (2016)10.1039/c5sm02572a).

80. D. F. Kucik, E. L. Elson, M. P. Sheetz, Weak dependence of mobility of membrane protein aggregates on aggregate size supports a viscous model of retardation of diffusion (vol 76, pg 314, 1999). *Biophysical Journal* **76**, 1720-1720 (1999); published online EpubMar (

81. J. Chow, Correlation Matrix Scatterplot (https://www.mathworks.com/matlabcentral/fileexchange/53043-correlation-matrix-scatterplot). *MATLAB Central File Exchange*, (2020); published online EpubRetrieved August 12, 2020 (

82. H. Wang, X. Tang, J. Liu, S. Trautmann, D. Balasundaram, D. McCollum, M. K. Balasubramanian, The multiprotein exocyst complex is essential for cell separation in Schizosaccharomyces pombe. *Mol Biol Cell* **13**, 515-529 (2002); published online EpubFeb (

83. F. O. Bendezu, V. Vincenzetti, S. G. Martin, Fission yeast Sec3 and Exo70 are transported on actin cables and localize the exocyst complex to cell poles. *PLoS One* **7**, e40248 (2012)10.1371/journal.pone.0040248).

84. J. Q. Wu, J. Bahler, J. R. Pringle, Roles of a fimbrin and an alpha-actinin-like protein in fission yeast cell polarization and cytokinesis. *Mol Biol Cell* **12**, 1061-1077 (2001); published online EpubApr (

85. V. Sirotkin, J. Berro, K. Macmillan, L. Zhao, T. D. Pollard, Quantitative analysis of the mechanism of endocytic actin patch assembly and disassembly in fission yeast. *Mol Biol Cell* **21**, 2894-2904 (2010); published online EpubAug 15 (10.1091/mbc.E10-02-0157).

86. M. Kohli, V. Galati, K. Boudier, R. W. Roberson, P. Philippsen, Growth-speed-correlated localization of exocyst and polarisome components in growth zones of Ashbya gossypii hyphal tips. *J Cell Sci* **121**, 3878-3889 (2008); published online EpubDec 1 (10.1242/jcs.033852).

87. N. Taheri-Talesh, T. Horio, L. Araujo-Bazan, X. Dou, E. A. Espeso, M. A. Penalva, S. A. Osmani, B. R. Oakley, The tip growth apparatus of Aspergillus nidulans. *Mol Biol Cell* **19**, 1439-1449 (2008); published online EpubApr (10.1091/mbc.E07-05-0464).

88. D. L. Delgado-Alvarez, O. A. Callejas-Negrete, N. Gomez, M. Freitag, R. W. Roberson, L. G. Smith, R. R. Mourino-Perez, Visualization of F-actin localization and dynamics with live cell markers in Neurospora crassa. *Fungal Genet Biol* **47**, 573-586 (2010); published online EpubJul (10.1016/j.fgb.2010.03.004).

89. D. McCusker, A. Royou, C. Velours, D. Kellogg, Cdk1-dependent control of membrane-trafficking dynamics. *Mol Biol Cell* **23**, 3336-3347 (2012); published online EpubSep (10.1091/mbc.E11-10-0834).

90. J. Y. Tinevez, N. Perry, J. Schindelin, G. M. Hoopes, G. D. Reynolds, E. Laplantine, S. Y. Bednarek, S. L. Shorte, K. W. Eliceiri, TrackMate: An open and extensible platform for single-particle tracking. *Methods* **115**, 80-90 (2017); published online EpubFeb 15 (10.1016/j.ymeth.2016.09.016).

91. J. Huang, Y. Huang, H. Yu, D. Subramanian, A. Padmanabhan, R. Thadani, Y. Tao, X. Tang, R. Wedlich-Soldner, M. K. Balasubramanian, Nonmedially assembled F-actin cables incorporate into the actomyosin ring in fission yeast. *J Cell Biol* **199**, 831-847 (2012); published online EpubNov 26 (10.1083/jcb.201209044).

Acknowledgments

We thank Stephan Gruber (Unil), Jan-Willem Veening (Unil), Serge Pelet (Unil), Laura Merlini (Unil), and Olivia Muriel-Lopez (Unil) for careful reading of the manuscript.

Funding:

ERC consolidator grant CellFusion (SGM)

Swiss National Science Foundation grant 310030B_176396 (SGM)

National Institutes of Health Grant R01GM114201 (DV)

National Institutes of Health Grant R35GM136372 (DV)

Author contributions:

IL, VG, AV and SGM conceived the project. IL performed all experiments in Fig 1, 3, S1 and S3. VG performed all other experiments in Fig 4-6 and S6-8. DMR and DV conceptualized the simulations and FRAP analysis methods. DMR performed all simulations and FRAP analyses. AV performed the initial RitC multimerization experiments. DG performed initial Rga4 structure-function analysis. VV provided technical support. SGM and DV coordinated the project and acquired funding. SGM and DV wrote the first draft of the manuscript, which was revised by all authors.

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:
All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

Figure 1. Depletion of membrane-associated CRY2 around sites of secretion

A. Spinning disk imaging time-lapse of CRY2-mCherry, CIBN-mTag-BFP2-RitC and CRIB-3GFP cells grown in the dark. Time 0 is the first timepoint after illumination. Yellow arrowheads indicate depletion zones. White arrowheads point to lateral peaks. **B.** Correlation plot of CRY2 depletion after 5 min and pole GFP intensity of CRIB-3GFP and GFP-Ypt3. Depletion cut-off is indicated by red to green colour change. **C.** Cells as in (A) treated with 300μM brefeldin A (BFA) or solvent (EtOH). **D.** Cells as in (A) carrying or not the *ypt3-i5* temperature-sensitive allele and grown at 36°C for 30 min to inactivate secretion. The control at the permissive

temperature of 25°C is shown in Fig S1H. **E.** Cells as in (A) carrying a *sty1* deletion (*sty1Δ*) treated or not with 50µM Latrunculin A (LatA) to prevent endocytosis. In (C-E), the GFP channel is a 5-min temporal sum projection. mCherry shows individual time points. Yellow arrowheads indicate depletion zones. Bottom graphs show CRIB-3GFP fluorescence plotted against CRY2 depletion at cell tips, as in (B). Data points are coloured with different shades of green according to CRIB-3GFP fluorescence. The indicated Kruskal-Wallis test *p*-value compares CRY2 depletion values between cells of equivalent CRIB-3GFP intensities. Scale bars are 3µm in all figures.

Figure 2. Simulations of membrane-flow-induced concentration gradients

A. Particles representing proteins have diffusion coefficient *D* on a spherical cell and exchange with a uniform internal pool with rates k_{off}, k_{on}. Exocytosis (endocytosis) events occur with a distribution of width σ_{exo} (σ_{endo}) around a point representing cell tip. The green particle represents any protein that reversibly associates with the PM. **B.** Particle displacement away/towards site of exocytosis/endocytosis by a distance Δr up to a maximum distance of R_{cutoff}. **C.** Depletion of particles starting from uniform distribution occurs for $\sigma_{exo} < \sigma_{endo}$, under static conditions (exocytosis rate = measured endocytosis rate). **D.** Tip occupation ratio vs k_{off} and *D*, same conditions as (C). Points are an average taken from t = 15,000-20,000 s after the start of the simulation. **E.** Scaled number density vs arclength shows depletion and lateral peak near sphere equator at low k_{off} values, same conditions as (C). Dashed vertical grey line represents equator; dashed horizontal black line represents isotropic density value. Lines are time averaged from 10,000 to 20,000 s.

Figure 3. Experimental validation of membrane flows with the optogenetic CRY2-CIBN system

A. D and k_{off} of CIBN-RitC and CRY2 with or without CRY2 co-expression, derived from FRAP measurements. Points are best D and k_{off} fits for individual cells. Boxes are data point lower and upper quartiles. Lines show D and k_{off} combinations resulting in same overall FRAP recovery (from ridge of averaged R^2 data fits, Fig S4). Curved corner region corresponds to comparable contributions of D and k_{off}; outer limits to recovery dominated by diffusion or exchange. Grey shaded regions have predicted tip occupation below indicated levels (from growth condition simulations). **B.** CRY2 cortical profiles around cell poles in time-lapse imaging as in Fig 1A. Time 0 is the first post-illumination image. Please compare to Fig 2E. The green dashed line shows the average CRIB distribution. The inset shows the average fluorescence value over time in the shaded areas. Blue shaded area is the standard deviation for time point t=270sec. **C.** Kymographs of the cell pole in cells as in Fig 1A. Arrowheads mark cluster of CRY2 that move away from the center of the CRIB-labelled region, suggestive of flow-driven displacement. **D.** CRY2 cortical profiles around division sites in time-lapse imaging as in Fig S1A. Traces represent average values of cells at different times of the cell division process. Time 0 is the first post-illumination image. The green dashed line shows the average CRIB distribution.

Figure 4. The lateral diffusion and detachment rates of peripheral membrane proteins conditions their cortical distribution by coupling to bulk membrane flows

A. Airyscan imaging localization of indicated sfGFP-tagged RitC constructs in cells co-expressing mCherry-1xRitC. Colour coded boxes match colour coding in panels B, C and E. **B.** D and k_{off} of indicated RitC constructs derived from FRAP measurements. Points are best D and k_{off} fits for individual cells. Boxes are data point lower and upper quartiles. Lines show D and k_{off} combinations resulting in same overall FRAP recovery (from ridge of averaged R^2 data fits, Fig S4). Curved corner region corresponds to comparable contributions of D and k_{off}; outer limits to recovery dominated by diffusion or exchange. Grey shaded regions have predicted tip occupation below indicated levels (from growth condition simulations). **C.** Distribution of sfGFP-RitC constructs at the cortex and CRIB signal distribution (from cells as in Fig S6). **D.** Time-lapse of 3xRitC co-expressed with CRIB-3xmCherry with 5min interval between frames. White arrowheads point to tip depletion events at sites of growth, the blue arrowhead indicates tip repopulation event when growth stops during cell division, yellow arrowheads point to local depletion events at the future division site also marked by CRIB. **E.** Ratios of tip-to-side signal measured for the indicated constructs in the indicated growth conditions. LatA treatment experiments were performed in *sty1Δ* cells on cells grown on lectin coated plates in rich media (YE) as previously reported (*24*). The bottom panel shows corresponding images. *p*-values are from student t-test with unequal variance. **F.** Scheme of secretion-induced membrane flow and influence on the distribution of peripheral membrane-associated proteins. Local secretion associated with endocytosis over a wider zone leads to bulk membrane flows away from the secretion site. Membrane-associated proteins (green) with sufficiently long membrane residence time (high affinity) couple to the flows and are depleted from the secretion zone.

Figure 5. Structure-function analysis of the Cdc42 GAP Rga4 defining membrane-binding and oligomerization motifs coupling to membrane flows

A. Maximum projection image of Rga4-RFP and CRIB-3GFP over a medial 4μm cell section. The graph shows average Rga4 distribution in sum projections along the cell cortex in 70 cells

from 3 experiments. Profiles were aligned from the cell tip with higher CRIB intensity (green line). Shaded areas indicate stdev. **B.** Scheme of Rga4 and truncation fragments. The sequence alignment shows the CBD across *Schizosaccharomyces* species (*S.p. = S. pombe*; *S.o. = S. octosporus*; *S.c. = S. cryophilus*; *S. j. = S. japonicus*). Fragments are numbered to help guidance through panels (C-E). **C.** Localization of indicated Rga4 truncations, showing role of CBD in membrane localization and role of coiled coils (cc) in cluster formation. **D.** Localization of full-length Rga4 and the cc-CBD minimal Rga4 with or without mutations in MB #1 and #2. For full-length Rga4, max projections of 10 time points at 2s interval are shown. **E.** Localization of the minimal cc-CBD Rga4 fragment, as well as single or 4x tandem CBD and cc-CBD-Δ2 constructs. Colour-coded boxes match colour coding in panels F and G. **F.** Average cortical distribution of indicated constructs. Shaded areas show standard deviation. **G.** Diffusion and membrane detachment rates of Rga4 fragments as in (B) derived from FRAP measurements. Points are best D and k_{off} fits for individual cells. Boxes are data point lower and upper quartiles. Lines show D and k_{off} combinations resulting in same overall FRAP recovery (from ridge of averaged R^2 data fits, Fig S4). Curved corner region corresponds to comparable contributions of D and k_{off}; outer limits to recovery dominated by diffusion or exchange. Grey shaded regions have predicted tip occupation below indicated levels (from growth condition simulations).

Figure 6. Bulk membrane flow-induced Cdc42 GAP depletion forms a negative feedback restricting the secretion zone.

A. Localization of CRY2-mCherry-RitC probe in light versus dark conditions in wild type and *rga3Δ rga4Δ rga6Δ* cells. Arrowheads point to depletion zones in light conditions. **B.** *rga3Δ rga4Δ rga6Δ CRIB-3GFP* cells expressing or not optoGAP continuously grown in the light or dark for 96h. **C.** CRIB-GFP cortical profiles in light-grown optoGAP cells vs. cells not expressing optoGAP as in (B). **D.** Aspect ratio of cells as in (B). p-values calculated from student t-test with equal variances. **E.** Percentage of cells as in (B) with CRIB cortical signal. **F.** Serial dilution of indicated cells on rich medium. **G.** Cells indicated as a, b, c in (B, dark conditions) grown in the dark and shifted to light at time 0. Kymographs at the bottom were constructed over segmented lines of invariant length marked by the yellow arrows (i to iv). OptoGAP clusters show directional movement away from the newly forming polarity site but

are almost static at lateral positions. **H.** Cells indicated as a, b, c in (B, light conditions) grown in the light. Time 0 is set at the first time point with CRIB-3GFP signal at the division site. Kymographs at the bottom were constructed over segmented lines of invariant length marked by the yellow arrows (i, ii). During cell division, OptoGAP clusters display extensive directional movement away from the division site towards the previously growing cell end. Note that in the 12-min time-lapse shown, the cells are not growing. **I.** A double negative feedback regulates the size of the Cdc42-GTP zone. Cdc42-GTP promotes secretion, which induces membrane-associated flow of Rga4 GAP to cell sides where it restricts Cdc42 activation, thus constraining the Cdc42-GTP zone.

Table 1. Reference parameter values used in simulations.

Variable	Value	Description
R	1.8 µm	Radius of model sphere, approximation to *S. pombe* radius
R_{cutoff}	2 µm	Cutoff range for exocytosis/endocytosis effect on sphere surface
α	0.5	Mobile membrane fraction
γ	1	Ratio of lipid to protein velocity
R_{exo}	50 nm	Radius of individual exocytotic vesicles (*57, 58, 82*)
A_{exo}	3.14×10^{-2} µm^2	Surface area of individual exocytotic vesicles
R_{endo}	22.6 nm	Radius of individual endocytotic vesicles based on measurements in *S. cerevisiae* (*60*)
A_{endo}	6.42×10^{-3} µm^2	Surface area of individual endocytotic vesicles
r_{endo}	2.44/s	Rate of endocytosis events [This work]
$r_{exo,growth}$	0.68/s	Rate of exocytosis events under growth conditions (20.7 µm^2/hr) [This work]
$r_{exo,static}$	0.50/s	Static rate of exocytosis events (no net added area)
σ_{exo}	0.848 µm	Standard deviation of exocytosis distribution [This work]
σ_{endo}	1.838 µm	Standard deviation of endocytosis distribution [This work]

Supplementary Materials for

Cell patterning by secretion-induced plasma membrane flows

Veneta Gerganova[1,&], Iker Lamas[1,&], David M. Rutkowski[2,&], Aleksandar Vještica[1], Daniela Gallo Castro[1], Vincent Vincenzetti[1], Dimitrios Vavylonis[2,*] and Sophie G Martin[1,*]

*Corresponding authors: sophie.martin@unil.ch; vavylonis@lehigh.edu

This PDF file includes:

Other Supplementary Materials for this manuscript:

Movies S1 to S10

SUPPLEMENTAL METHODS

Determination of membrane diffusion and dissociation rates by FRAP

Diffusion coefficients and membrane dissociation constants were determined by fitting FRAP recovery profiles to a 1D model (Fig S4). The method relies on the different effects of diffusion and association/dissociation on the recovery, with diffusion providing smoothing of profile over time (*71*) and association/dissociation contributing to uniform recovery.

Experimental FRAP profiles. Experimental fluorescence intensity profiles were acquired after photobleaching of a rectangular region and imaging, on the top surface of *S. pombe* cells (Fig S4A; see experimental details in the experimental methods). We corrected for photobleaching after the initial FRAP event by multiplying all intensities by e^{kt} after background subtraction, where k is the exponential decay constant of the average intensity of a neighboring non-bleached cell. From these photobleach-corrected images we determined the intensity profile $I(x,t)$ by defining a rectangular region of interest along the long axis of a bleached cell with width ~2 μm, excluding the region close to the cell tips, and calculated the average intensity along this line at points spaced 0.0425 μm (1 pixel) apart through the bleaching recovery using ImageJ's getProfile function (Fig S4B).

Model with diffusion and association/dissociation used for fitting. To match the experimental bleach recovery, we modelled the evolution of the concentration profile of a membrane-associated protein, $c_m(x,t)$, along a 1D line with reflecting boundaries at positions x_a, x_b, defining the size of the finite membrane reservoir (approximately the location of cell poles and beyond, to include the back side of the cell). The x-dimension in the model corresponds to the long axis of the cell. We assumed $c_m(x,t)$ obeys the following equations, which include both the effects of diffusion and membrane binding/unbinding:

$$\frac{\partial c_m(x,t)}{\partial t} = D\frac{\partial^2 c_m(x,t)}{\partial x^2} - k_{off}c_m(x,t) + k_{on}C(t)$$

$$C(t) = C_{tot} - \int_{x_a}^{x_b} c_m(x,t)dx, \tag{M5}$$

where c_m is concentration on the membrane , C is the number of proteins in the cytoplasm, C_{tot} is the total number of proteins, and k_{off} , k_{on} are the membrane dissociation and association rate constants. We do not write a separate diffusion equation for the cytoplasm, assuming that cytoplasmic diffusion is sufficiently fast. We also specifically consider the limit where most of the proteins are on the membrane: $C(t) \ll C_{tot}$, (equivalently, the limit of sufficiently large k_{on}). The Green's function describing the probability for a single protein being at position x at time t, given that it was at position x_0 at time $t = 0$ is

$$p(x,t|x_0,0) = e^{-k_{off}t}p_D(x,t) + \frac{1-e^{-k_{off}t}}{x_b-x_a}, \tag{M6}$$

where $p_D(x,t)$ accounts for the probability of the particle position due to diffusion alone:

$$p_D(x,t|x_0,0) = \frac{1}{x_b-x_a} + \frac{2}{x_b-x_a}\sum_{n=1}^{\infty} e^{-Dt\lambda_n^2}\cos(\lambda_n(x-x_a))\cos(\lambda_n(x_0-x_a)), \text{ where } \lambda_n = \frac{n\pi}{x_b-x_a} \tag{M7}$$

General procedure of fitting experimental I(x,t) profiles. Assuming $c_m(x,t)$ is proportional to the experimental $I(x,t)$, we solve for $c_m(x,t)$ given an initial condition $c_m(x,0)$ that is proportional to the intensity profile imaged immediately after the photobleaching, $I(x,0)$. We then fit the model

to both x and t after photobleaching simultaneously, using SciPy's optimization curve fitting to get a single best fit value for D and k_{off} as described below. The sum in Eq. (M7) was carried up to $n = 500$. We checked that replacing this by 250 did not have a significant effect on the values determined for D and k_{off} for our experimental data.

Evaluation of initial intensity distribution and simulation boundary positions x_a and x_b. The intensity profile is divided into three regions: left, middle, and right (Fig S4B). The intensity in the middle region, defined to be between x_d and x_e where $x_a \leq x_d \leq x_e \leq x_b$, is determined by the intensity profile I_i^0 immediately after photobleaching, evaluated at pixels i of the projected profile, corresponding to positions x_i (Fig S4B). The middle region is therefore treated as a sum of $n = \Delta x/(x_e - x_d)$ separate integrals, where Δx is the pixel size. The lengths of the left and right flanking regions are both equal to l_F, with $l_F = x_d - x_a = x_b - x_e$. We average the intensity of the pre-bleached cell over the middle region and set this as the initial intensity of the two flanking regions, I_F^0. The value of l_F determines the uniform intensity at long times, I_∞, according to mass conservation:

$$(2l_F + x_e - x_d)I_\infty = \sum_{i=0}^{n} I_i^0 \Delta x + 2l_F \, I_F^0. \tag{M8}$$

The long time intensity I_∞, or equivalently l_F, is determined via the fitting procedure. In mutants where the l_F value drifted towards 0 or large numbers, it was restricted to be in the range 2-10 μm.

Evolution of concentration on a discrete lattice starting from a given initial condition. The equation describing the concentration or intensity as a function of position x_i corresponding to pixel i, and time is found by integration, $I_{model}(x_i, t) = \int_{x_a}^{x_b} I(x', 0)p(x_i, t|x', 0)dx'$, which, using Equations (M6) and (M7) gives:

$$I_{model}(x_i, t) = \frac{1}{x_b - x_a}\left[2l_F I_F^0 + \frac{1}{2}\sum_{i=0}^{n}(I_i^0 + I_{i+1}^0)\Delta x\right]$$

$$+ \frac{2\exp(-k_{off}t)}{x_b - x_a}\left[I_F^0 S_{x_a,x_d}(x_i, t) + I_F^0 S_{x_e,x_b}(x_i, t) + \frac{1}{2}\sum_{j=0}^{n}(I_j^0 + I_{j+1}^0)S_{x_j,x_{j+1}}(x_i, t)\right]$$

$$\tag{M9}$$

where $S_{x_1,x_2}(x, t) = \sum_{m=1}^{\infty}\frac{e^{-Dt\lambda_m^2}}{\lambda_m}\cos(\lambda_m(x - x_a))\left[\sin(\lambda_m(x_2 - x_a)) - \sin(\lambda_m(x_1 - x_a))\right]$ and λ_n is given in Equation (M7). Here, the initial intensity $I(x, 0)$ at $x_i < x < x_{i+1}$ was assumed to be given by $0.5(I_i^0 + I_{i+1}^0)$. An example of a fit over time to the experimental FRAP data for a cell expressing 1xRitC-GFP using this expression is shown in Fig S4C.

Accuracy of determining D and k_{off}. The fit to each FRAP profile determines the single best fit values for D and k_{off} for each cell, corresponding to the largest R^2 value that is a measure of the goodness of the fit. These two values are not independent of each other: since both contribute to recovery, increasing the value of one parameter can be partly compensated by a decrease in the other (even though they influence the evolution of the profile differently, depending on the size of the bleached region). Additionally, one of the two parameters may be too small and thus dominated by the other. These complications are reflected in a large scatter of values for D and k_{off} among cells (Fig 3A, 4B and 5G). To check that despite this scatter, averaging over cells is meaningful, we developed a global procedure to determine the best D and k_{off} simultaneously for all cells. For

each cell of a given cell type, we varied D and k_{off} over six orders of magnitude and calculated the R^2 value. We then plotted the R^2 value as function of D and k_{off}, averaged over all cells of the same cell type (Fig S4D-F). The best fitting regions indicated by high values of R^2 typically form a pair of branches: one horizontal (constant k_{off}) and the other vertical (constant D) which intersect to form a curved bend. The median of the best fit value for D and k_{off} as determined by the 1D model is indicated by a red circle. For most photobleached proteins this highest average R^2 (indicated by a blue circle) is close to the median of the best fit values from the 1D model. One case does not show particularly good consistency (1x cc-CBD-Δ2), where the highest average R^2 is at very low values of k_{off}), suggesting that the values determined for D and k_{off} for these proteins is not reliable. Vertical or horizontal regions of high R^2 that extend beyond the boundaries of the graphs also indicate that the parameter varied along such a region may only be determined up to an upper limit. They typically correspond to scattered D and k_{off} values in Fig 3A, 4B and 5G. Finally, the line defining the ridge of the pair of branches that intersect to form a curved bend in Fig S4D,E is plotted as continuous line in Fig 3A,D and 4D to indicate the different combinations of D and k_{off} that provide similar overall recovery.

Simulations to validate FRAP fitting procedure. One limitation of our FRAP fit method to extract D and k_{off} is that we neglect the diffusion along the radial cylindrical direction. To examine the implications of this limitation, as well as another check of our method, we simulated FRAP experiments accounting for diffusion and membrane association/dissociation of particles on the surface of a cylinder with reflecting boundary conditions at the ends. We used a similar method as used for the sphere model with 50,000 particles on the surface of a cylinder with radius 2 μm and length 12 μm. As with the discrete sphere model, we fix $k_{on} = 0.01 \ s^{-1}$. The displacement is calculated as $\sqrt{2D\Delta t}\xi(t)$ independently in both directions in the plane tangent to the cylinder surface, where Δt is the timestep determined from the Gillespie algorithm and $\xi(t)$ is a random Gaussian distributed variable with zero mean and unit variance. The displacement along the long axis of the cylinder needs no modification, but the displacement in the radial direction is projected down onto the cylinder surface while maintaining the distance as with the sphere model. To ensure that the radial diffusion is properly implemented, we measure the change in the azimuthal angle, $\langle\phi(t)^2\rangle$, as a function of time up to a time step of 1000 s, similar to Fig S9A.

The bleaching recovery was measured over a region of width 2.36 μm which is similar to the average experimental width of experimental cells. We took snapshots of particle positions every 1 s for a total of 500 s. To simulate initial photobleaching we label particles on the surface which are within a certain region at a given frame number as being bleached. This simulated bleach region is rectangular on the surface of the cylinder with a long axis of 3.315 μm and an arc length of half the circumference. To calculate the simulated intensity profile, $I_{sim}(x,t)$, we counted the number of particles in this region over time that were not labelled as being bleached in the initial frame and further assume that the intensity is proportional to the number of particles. To get better averages for the bleaching recovery, we simulate box photobleaching at each snapshot and follow the k independent recovery trajectories from their respective bleached frame, where k is the number of snapshots. We then average the intensity curves for each bleaching recovery at the same amount of time passed since the bleaching frame to obtain $\langle I_{sim}(x,t)\rangle$. We then use $\langle I_{sim}(x,t)\rangle$ up to a time of 250 s as the input for the 1D model to predict D and k_{off}. When using $D = 10^{-3}$ μm²/s and $k_{off} = 10^{-2}$ /s for input parameter values of the discrete simulations, which are approximately in the range of 2xRitC-GFP, there is good agreement between the values inputted

into the simulations i.e. the values determined by the R^2 plot (red circle), and the values determined by the 1D model (Fig S4D). This result indicates that the 1D model can correctly predict D and k_{off} values. For input parameters with zero diffusion ($D = 0$, $k_{off} = 10^{-2}$ /s) and zero membrane unbinding ($D = 10^{-3}$ µm²/s, $k_{off} = 0$) shown in Fig S4G the red dashed line indicates the non-zero input parameters are accurately determined, however a small nonzero value is assigned to the other parameter. '

Comments on measured D and k_{off} values. Even though the estimated values of D or k_{off} can be partly compensated by an inverse change in the other, the FRAP recovery fits to the 1xRitC, 2xRitC, 3xRitC constructs do indicate that the main parameter responsible for slowing dynamics with increasing number of RitC domains is k_{off} (Fig S4E). This trend most likely indicates the cooperative nature of the dissociation process for constructs with multiple binding sites (*76*). Our results also point to a potential strong reduction of the diffusion coefficient with increasing number of RitC domains, however more precise measurements are needed to resolve if such a reduction is consistent with a weak size dependence of *D* expected for dilute systems (*77, 78*) or else if this is an indication of crowding mechanisms and microdomain formation in the fluid membrane component (*79, 80*). Interestingly, our measurements also indicate slower diffusion of 3xsfGP-1xRitC compared to sfGP-1xRitC. This observation could indicate that cytoplasmic drag might dominate diffusion of the 3xsfGP construct; further studies could explore if this diffusion reflects size-dependent frictional interaction with cytoplasm adjacent to inner membrane leaflet or else if it is due to oligomerization and related processes.

Image analysis of concentration profiles

All image-processing analyses were performed with Image J software (http://rsb.info.nih.gov/ij/). Image and time-lapse recordings were imported to the software using the Bio-Formats plugin (http://loci.wisc.edu/software/bio-formats). Time-lapse recordings were aligned using the StackReg plugin (http://bigwww.epfl.ch/thevenaz/stackreg/) using the rigid body method. All optogenetic data analyses were performed using MATLAB (R2019a), with scripts developed in-house. Figures were assembled with Adobe Photoshop CC2019 and Adobe Illustrator 2020.

Correlating CRY2PHR-mCherry tip depletion dynamics with endogenous GFP-tagged protein signal intensities at cell tips.

CRY2PHR-mCherry tip depletion dynamics was monitored in strains co-expressing CRIB (N = 3, n = 275 cells), Ypt3 (N = 3, n = 184 cells), Exo84 (N = 3, n = 290 cells) and Exo70 (N = 3, n = 281 cells) GFP-tagged proteins (sample strains) in normal conditions (Fig 1B and S1C-E); in strains co-expressing CRIB-3GFP and GFP-Ypt3 treated with BFA (CRIB: N = 3, n = 219 cells; Ypt3: N = 3, n = 230 cells) or ethanol as control (CRIB: N = 3, n = 127 cells; Ypt3: N = 3, n = 136 cells); in *ypt3-i5* CRIB-3GFP and *ypt3+* CRIB-3GFP strains at 25 ° C (*ypt3+* CRIB-3GFP: N = 3, n = 191 cells; *ypt3-i5* CRIB-3GFP: N = 3, n = 132 cells) and 36 °C (*ypt3+* CRIB-3GFP: N = 3, n = 229 cells; *ypt3-i5* CRIB-3GFP: N = 3, n = 204 cells); and in *sty1Δ* CRIB-3GFP strain treated with LatA (N = 3, n = 276 cells) or left untreated (N = 3, n = 293 cells). Tip depletion was assessed upon photoactivation of the CRY2-CIB1 system by recording the RFP fluorescence intensity over an ROI that was 12 pixels wide by 25 pixels long (≈ 1 µm by 2.075 µm) and drawn perpendicular to the cell tip cortex over 20 timepoints, as shown

by the blue arrow in the scheme on the right. The RFP and GFP fluoresce profiles across the plasma membrane were recorded over time. The GFP signal was used to correlate the presence of endogenous proteins with the CRY2PHR-mCherry tip depletion dynamics. Note that in these sets of experiments the initial timepoint was not included in the analysis, since CRY2PHR does not yet achieve its maximal plasma membrane recruitment (see Fig 3B). Thus, the second time point was set as the initial timepoint (t_0) in these experiments (19 timepoints in total). This corresponds to t = 0 s in Fig 3B.

To derive photobleaching correction coefficients, the average camera background signals (*Bckg*) from 6 cell-free regions were measured; and fluorophore bleaching was measured from entire cells, using RFP controls for RFP and sample cells for GFP.

RFP bleaching correction coefficient =
$$\left(RFP\ Intensity\ of\ cell_{tn} - Bckg_{tn} \right) \Big/ \left(RFP\ Intensity\ of\ cell_{t0} - Bckg_{t0} \right) \quad \text{(E1)}$$

GFP bleaching correction coefficient =
$$\left(GFP\ Intensity\ of\ cell_{tn} - Bckg_{tn} \right) \Big/ \left(GFP\ Intensity\ of\ cell_{t0} - Bckg_{t0} \right) \quad \text{(E2)}$$

where *RFP Intensity* and *GFP Intensity* of cell stands for the signal measured from entire RFP control and sample cells, respectively; *Bckg* stands for the average fluorescence intensity of 6 independent cell-free regions; t_n represents a given time point along the time course of the experiment; and t_0 represents the initial time point (n = 19 time points). These coefficients were corrected by a moving average smoothing method (moving averaged values = 5). GFP and RFP profiles of sample cells were independently analysed as follows. First, GFP and RFP signals were background and bleaching corrected, using Eq E1 and E2:

$$\textit{Single} - \textit{Cell GFP/RFP profile}_{tn} = \left((Profile\ signal_{tn} - Bckg_{tn}) / bleaching\ correction\ coefficient_{tn} \right) \quad \text{(E3)}$$

where *Profile signal* intensity represents the GFP or RFP raw profile values across the plasma membrane, *Bckg* stands for the average fluorescence intensity of 6 independent cell-free regions, and t_n represents a given time point along the time course of the experiment.

Corrected single-cell GFP and RFP profiles were then aligned to their peaks in order to correct for cell growth and/or cell movement. The RFP peak at t_0 was used as a reference point to align single-cell RFP and GFP profiles over time. The profiles resulting from the peak alignment were used to calculate the cortical GFP and RFP fluorescence values over time:

Cortical RFP Trace $_{tn}$ =
$$\left(RFP\ intensity_{peak-1pixel\ tn} + RFP\ intensity_{peak\ tn} + RFP\ intensity_{peak+1pixel\ tn} \right) \Big/ 3 \quad \text{(E4)}$$

Cortical GFP Trace $_{tn}$ =
$$\left(GFP\ intensity_{peak-1pixel\ tn} + GFP\ intensity_{peak\ tn} + GFP\ intensity_{peak+1pixel\ tn} \right) \Big/ 3 \quad \text{(E5)}$$

The GFP and RFP fluorescence intensities at RFP$_{peak\ pixel\ position\ at\ t0}$ $\pm$ 1 pixel were averaged in order to extract CRY2PHR-mCherry tip depletion (RFP channel) and GFP-tagged traces across the duration of the time-lapse. From here on, GFP traces were saved and further analyses were performed only for RFP traces. First, the amplitude of the RFP tip depletion was calculated as follows:

$Amplitude\ RFP\ depletion\ =$
$$((Cortical\ RFP\ Trace_{t0} - average(Cortical\ RFP\ Trace_{t13-t18}) \Big/ Cortical\ RFP\ Trace_{t1}) * 100 \qquad (E6)$$

where *Cortical RFP trace t_0* represents the initial RFP signal at the plasma membrane and *average (Cortical RFP trace t_{13-18})* represents the average signal at the plasma of the last 5 timepoints of the trace. In addition, single-cell RFP traces derived from Eq E4 were also normalized to their own maximum RFP intensity:

$$Max.\ Normalized\ RFP\ trace\ = (Cortical\ RFP\ Trace_{tn} \Big/ Cortical\ RFP\ Trace_{tmax}) \qquad (E7)$$

where *Cortical RFP trace t_n* represents the RFP value of the trace at the timepoint *n* and *Cortical RFP trace t_{max}* represent the maximum RFP value of the single-cell trace.

From here on, normalized RFP traces from Eq E7 were sorted according to the amplitude of depletion derived from Eq E6. When the *Amplitude RFP depletion* was lower than 20 %, then the normalized RFP trace from Eq E7 was labelled as "Non-depleted". These traces are represented in red-coloured dots in Fig 1B and similar cluster plots. Cumulative GFP values of "Non-depleted" RFP traces were calculated as follows:

$$GFP\ value_{Non-depleted\ trace} = Cortical\ GFP\ Trace_{t1} + Cortical\ GFP\ Trace_{t2} ... + Cortical\ GFP\ Trace_{tn}$$
$$(E8)$$

Where *Cortical GFP trace t_n* represents the GFP value at timepoint n from the GFP trace calculated in Eq E5. Thus, GFP value for "Non-depleted" RFP traces correspond to the sum of GFP values along the 19 timepoints.

In contrast, when the *Amplitude RFP depletion* was greater than 20 %, then the normalized RFP trace from Eq E7 were labelled as "Depleted". These "Depleted" traces were used to calculate the tip depletion half-times:

$$Max.\ Normalized\ RFP\ trace\ t_{max}: t_n\ (y) = a * e^{(-b*tmax:tn)} \qquad (E9)$$

$$Tip\ Depletion\ t_{1/2} = \frac{ln(0.5)}{-b} \qquad (E10)$$

where *Max. Normalized RFP trace $t_{max}:t_n$* represent the normalized RFP trace from Eq E7 starting from the timepoint of its maximal RFP value (t_{max}) to the last timepoint of the trace (t_n) and t_n represents the last timepoint of the trace. If t_{max} corresponds to t_0, then it means that the tip was depleting from the very beginning of the recording. In contrast, if t_{max} does not corresponds to t_0, then it means that the tip depletion started later on. Therefore, a *Depletion delay* variable was also calculated.

$$Tip\ Depletion\ delay = t_{max} - t_0 \qquad (E11)$$

where *t_{max}* represents the timepoint at which the maximal RFP value was detected along the single-cell trace and *t_0* represents the first timepoint of the trace. 67% of traces started depleting from t_0, 19% from t_1 and a few % thereafter.

Finally, cumulative GFP values for "Depleted" RFP traces were calculated as follows:

$$GFP\ value_{Depleted\ trace} = Cortical\ GFP\ Trace_{tmax} + Cortical\ GFP\ Trace_{tmax+1} ... +$$
$$Cortical\ GFP\ Trace_{tn} \qquad (E12)$$

where *Cortical trace t_{max}* represents the GFP value of the GFP trace calculated in Eq E5 at timepoint t_{max}. Thus, GFP value for "Depleted" RFP traces correspond to the sum of GFP values from t_{max} to t_n (*n = 19 timepoints*).

After these data analyses, RFP traces were further sorted based on the R^2 value of the fitting from Eq E9 and the *Tip depletion delay* value from Eq E11. "Depleted" RFP traces matching any of these criteria were discarded from the regression analysis:

1) The depletion delay from Eq E11 is more than 90 s (starts depleting after t7).
2) R^2 of curve fitting from Eq E9 is lower than 0.7, except in *ypt3-i5* and LatA experiments, where $R^2 > 0.5$ was accepted.

Additionally, "non-depleted" RFP traces, which depleted less than 20% but initiated depletion after t14 (delay from Eq E11 is more than 200 s) were discarded, as these may represent tips that may in fact eventually deplete, after the time of the time-lapse.

Finally, the cumulative GFP values from Eq E8 and Eq E12 and the amplitude of RFP depletion from Eq E6 of non-depleted and depleted traces were used to perform a linear regression analysis. The linear regression was performed using the correlation matrix scatterplot function for MatLab developed by John Chow* (*81*).

Statistical analyses shown in Fig 1C-D-E and Fig S1G-H were performed by comparing the 2nd quintiles (20% - 40% of GFP signal) of control and treated samples using Kruskal-Wallis test. Quintiles were defined based on GFP intensities of control and treated samples. The range of GFP values within an experiment used to define quintiles was set by the lowest and highest GFP values amongst control and treated samples.

CRY2PHR-mCherry lateral peak formation.

The lateral accumulation of CRY2PHR-mCherry at the edges of the depletion zone at cell tips in a strain co-expressing CRIB-3GFP and the CRY2-CIB1 system (N = 3, n = 268 traces in 134 cells) was monitored by recording the RFP and GFP fluorescence profiles over an ROI that was 3-pixels wide by more than 171-pixels long (≈ 0.25 µm by 14.2 µm) drawn along the cortex of sample cells, as shown by the blue arrow on the scheme on the right. The ROI was drawn from one cell side towards the other cell side, passing across the tip of the cell. The lateral accumulation of CRY2PHR-mCherry at the edges of the depletion zone at cell division site was monitored in the same strain (N = 3, n = 36 traces in 18 ROIs) by recording the RFP and GFP fluorescence profiles over an ROI that was 3-pixels wide by more than 60-pixels long (≈ 0.25 µm by 5 µm) drawn along the cortex of

 sample cells. The RFP and GFP fluoresce profiles were recorded over time from sample strains. The GFP signal was used to align single-cell RFP and GFP profiles. Because CRY2PHR only achieves its maximal plasma membrane recruitment at the second image acquisition, this was set as t = 0 s. (see Fig 3B).

Photobleaching correction coefficients were derived as in Eq E1-E2 and single-cell RFP and GFP cortical profiles were corrected using Eq E3. Single-cell cortical GFP profiles from all the timepoints (*n = 20 timepoints*) were averaged in order to generate an *average Cortical GFP profile*. The peak of the *average Cortical GFP profile* was used as a reference for the center of the cell tip. The peak pixel position of the *average Cortical GFP profile* was fitted to a Gaussian distribution, where the parameter *b* corresponds to the peak position:

$$\textit{GFP Gaussian fit } (\boldsymbol{y}) = a * e^{-(x-b)^2\big/2\sigma^2} \quad \text{(E13)}$$

Finally, all single-cell GFP and RFP profiles were aligned to the center of the cell tip, divided symmetrically in two traces and averaged per timepoint (Fig 3B, bottom).

Tip depletion and cell-side accumulation traces were calculated from *single-cell RFP profiles* by averaging the RFP intensities at pixel positions shown in Fig 3B (bottom) for every timepoint. For each cell the average fluorescence value at the cell tip (sampled over 15-pixel length = 1.25 μm), showing depletion over time, and at the edge of the depletion zone (sampled over 10-pixel length = 0.83 μm), showing accumulation over time, was calculated. Average traces are shown in Fig 3B (top).

Cortical distribution profiles of exocytosis (Exo70-GFP) and endocytosis (Fim1-mCherry) at the cell tips and cell middle.

To experimentally address the relative distribution of endo- and exocytosis, we compared the distribution of the exocyst subunit Exo70 and the actin-binding Fimbrin Fim1, as markers of exo- and endocytic events, respectively (*82-85*). Both markers are enriched at cell poles and division sites, and we examined their distribution at both locations (Fig S3). The distribution profiles were generated from the sum projection images of entire time-lapses (n = 21 timepoints). Exo70-GFP and Fim1-mCherry fluorescence profiles were recorded over ROIs drawn from one cortical cell side towards the other cell side (N = 3, n = 149 profiles) and at the cell middle cortex (N = 3, n = 70 profiles) (see Fig S3A and S3B). ROIs were 6-pixel wide by more than 171-pixel long (≈ 0.5 μm by 14.2 μm) for cell tips and 6-pixel wide by 92-pixel long (≈ 0.5 μm by 7.64 μm) for the cell middle.

Single cell profiles were first corrected for the background signal using Eq E14 and E15.

$$\textit{RFP corrected profile} = (RFP\ raw\ profile\ - Bckg_{RFP}) \quad \text{(E14)}$$

$$\textit{GFP corrected profile} = (GFP\ raw\ profile\ - Bckg_{GFP}) \quad \text{(E15)}$$

where *RFP/GFP raw profile* stands for Exo70-GFP and Fim1-mCherry profiles generated from ROIs drawn at cell cortex; and $Bckg_{RFP/GFP}$ stand for the average fluorescence intensity of 6 independent cytosolic regions. $Bckg_{RFP/GFP}$ serve to set fluorescence threshold present in the cytosol of Exo70-GFP and Fim1-mCherry cells. Due to the ubiquitous Fim1-mCherry cortical localization and GFP auto-fluorescence, *Bckg* ROIs were placed in the interception between the nucleus and the cytosol in the centre of cells co-expressing both markers. Upon background correction, RFP and GFP profile values below the threshold (i.e. < 0 A.U.) were set to 0 A.U. in order to prevent numerical aberrations during the analysis.

Single-cell *corrected GFP/RFP profiles* were then normalized to their maximum and minimum values and fitted to a Gaussian distribution using Eq E13. Finally, single-cell *corrected GFP/RFP profiles* were aligned to their geometrical center, averaged and normalized to the maximum and 0 values (Fig S3).

We note that Exo70-GFP exhibits a narrower distribution than Fim1-mCherry at growing cell poles (Fig S3A). We observed similar, even more extensive distribution differences in the middle of pre-divisional cells (Fig S3B). These observations show that endocytosis occurs in a region centred over, but wider than, the zone of polarized exocytosis. These findings are in line with previous observations in several fungi (*86-89*). The measured distributions at cell poles were used in the simulations.

Estimating endocytic and exocytic rates.

To estimate the rate of endocytosis, we counted the number of Fim1-mCherry patches at the cell side of cells co-expressing Exo70-GFP and Fim1-mCherry markers across the duration of the whole time-lapse. We also measured the average fluorescence intensity over the same cell-side regions in sum intensity projection images. These values were used to calculate an average Fim1-mCherry patch fluorescence. We performed this analysis at cell sides rather than cell tips to avoid overlap of endocytic signal at cell tips due to high density, which would lead to an underestimation of the number of events. By dividing the Fim1-mCherry distribution profile described above by the average patch fluorescence value, the average endocytic rate per pixel position across the Fim1-mCherry distribution profile was calculated as follows:

$$\textbf{\textit{Endocytic rate}}_n = ((^{(Mean\ Fim1-mCherry\ fluo._n/mean\ patch\ fluo.)}/_{21}) * 4) * 0.75 \qquad (E16)$$

where *Mean Fim1-mCherry fluo.* $_n$ stands for the average RFP fluorescence value at pixel position n; *mean patch fluo.* stands for the average Fim1-mCherry patch fluorescence. The value 21 represents the number of time points that contribute to the average Fim1-mCherry distribution profile. This division aims to estimate the number of endocytic event per image. This is multiplied by 4 to obtain the number of events per minute (imaging interval = 15s). Finally, the result of this operation was corrected by 0.75 in order to correct for the slight over sampling in our images, as the average lifetime of a Fim1-mCherry patch is about 20 s (*85*).

The rate of exocytosis events under growth conditions was calculated from the addition of the above rates of endocytosis and rates membrane addition contributing to cell growth. Cell growth was measured in time-lapse imaging as the difference in length over time between the invariant birth scar location and the pole of the cell, and converted to amount of required membrane to accommodate this growth (see model methods for further details).

The obtained values were used in the simulations and are presented in Table 1.

Cortical distribution profiles of GFP-1RitC, GFP-2RitC, GFP-3RitC and 3GFP-1RitC.

Strains co-expressing CRIB-3mCherry and any of the GFP-1RitC (N = 3, n = 266 profiles), GFP-2RitC (N = 3, n = 194 profiles), GFP-3RitC (N = 3, n = 246 profiles) or 3GFP-1RitC (N = 3, n = 274 profiles) constructs were imaged. Cortical distribution profiles were generated from single timepoint middle section images. The RFP and GFP fluorescence profiles were recorded over an ROI that was 3-pixels wide by more than 171-pixels long (≈ 0.25 μm by 14.2 μm) drawn along the cortex of sample cells. The ROI was drawn from one cell side towards the other cell side, passing across the tip of the cell. To correct single-cell profiles for the average camera background signals (*Bckg*), the average signal from 6 cell-free regions was subtracted using Eq E14 and E15. Single-cell *corrected GFP/RFP profiles* were then aligned to their geometrical center and divided symmetrically in two traces starting from the cell pole. To normalize for the differential plasma membrane binding affinity of the GFP-1RitC, GFP-2RitC, GFP-3RitC or 3GFP-1RitC constructs, and compare fluorescence distribution, the area under the curve (AUC) of single-cell *corrected GFP traces* were normalized to 1:

$$\textbf{\textit{Normalized GFP trace}} = Corrected\ GFP\ trace/AUC_{trace} \qquad (E17)$$

where *Corrected GFP trace* represents the fluorescence profile from the cell pole after background correction and *AUC_trace* stands for its area under the curve. Finally, *normalized GFP traces* were averaged (Fig 3E).

For the quantification of tip to side fluorescence intensities of the RitC series (Fig 4E) and correlation to CRIB-3xmCherry (Fig S6), 7 px-wide lines were drawn, one 12 px-long at the tip and two covering the cell sides to measure an average fluorescence signal at cell tip and at cell sides, respectively. Background corrections from the average value of 6 ROIs outside any cell from each individual field of view were performed for each channel. Tip/side ratios were calculated and plotted against CRIB intensity as scatter plots, with R^2 value calculated from the trend line generated in Excel. Data was collected from three replicate experiments for a total of 100-180 cells per strain.

Distribution profile of Rga4 and Rga4 fragments

For the corset profiles of Rga4 in Fig 5A, a 7 px-wide line was drawn along the cell cortex starting from the cell tip with higher CRIB intensity on a sum projection image obtained from z-stack AiryScan acquisition. Background correction from the average value of 6 ROIs in regions outside cells from each individual field of view was performed.

Cortical profiles for Rga4 fragments (Fig 5F) were manually drawn as 7px-wide cortical lines from cell tip to cell tip along each side of the cell and data was collected using the ImageJ multi-plot function. The middle of each profile was identified, the profiles were split in half and aligned to cell tips using MatLab. Background correction from the average value of 6 ROIs outside of cells from each individual field of view was performed. The plots represent data from three replicate experiments with a total of 150-180 cells per strain, shaded area is standard deviation between cells.

TIRF movies were analysed using the TrackMate Fiji plug-in (*90*) and tracking was performed on individual cells. The particle diameter was set to 0.3 μm with maximum gap linking of two frames, and linking range of 0.15 μm. Quality control for spot identification was set at cut-off of 19 and was visually confirmed for each movie.

All boxplots and violin plots were generated using the BoxPlotR web-tool (http://shiny.chemgrid.org/boxplotr/) with definition of whisker extend by Tukey.

Analysis of optoGAP

The quantification of optoGAP and no GAP in Fig 6E shows the number of cells with a clear CRIB zone divided by the total number of cells in both light and dark condition. Quantifications were performed from 3 replicate experiments with a total number of counted cells indicated in the figure. Error bars represent standard deviation between experiments. The CRIB fluorescence intensity profile in Fig 6C was quantified by drawing, for each cell with a CRIB zone, two 7 px-wide profiles along the cell cortex starting at the middle of the cell tip. Data was collected using the multi-plot function of ImageJ and corrected for background from ROIs outside of cells for each individual field of view. For the *rga3Δ rga4Δ rga6Δ* cells without transgene, only cells with a single CRIB zone were considered. Data was collected from three replicate experiments for a total of 150 cells per strain. Shaded area represents standard deviation between cells. The aspect ratio in Fig 6D was calculated from the measured cell length to cell width of dividing cells in three replicate calcofluor experiments for 500-600 cells per strain. Statistics were performed on the averages of the three experiments using a student t-test with equal variances in Microsoft Excel.

Cortical kymographs

Kymographs shown in Fig 3C and Fig S3 were generated with the MultipleKymograph (https://www.embl.de/eamnet/html/body_kymograph.html) ImageJ plugin using a 1-pixel-wide and 6-pixel-wide ROI drawn along the cell cortex, respectively. Kymographs in Fig 6G-H were generated with the FIJI stack > reslice tool without interpolation using a static 15-pixel-wide segmented line drawn along the cell cortex.

Table S1: *S. pombe* strains used in this study

Number	Genotype	Source
Fig 1, Fig S1, Fig S3		
YSM3568	*h- leu1-32::Ppak1-CRIB-3xGFP-KanMX+:leu1+ ura4-D18::Ptdh1-CIBN-mTagBFP2-Ritc-ADH1term-Pact1-CRY2PHR-mCherry:ura4+ ade6+ leu1-32*	(*2*)
YSM3564	*ura4-294::Pact1-CRY2PHR-mCherry::ura4+ ade6+ leu1+ ura4-294*	(*2*)
YSM3803	*h- ade6-D19::pnmt41-GFP-ypt3-hph+::Ade6+ ura4-D18::Ptdh1-CIBN-mTagBFP2-Ritc-ADH1term-Pact1-CRY2PHR-mCherry:ura4+ ade6- leu1+ ura4-D18*	This study.
YSM3801	*exo70-GFP-Kan+ ura4-::Ptdh1-CIBN-mTagBFP2-RitC-Adh1term-Pact1-CRY2PHR-mCherry::ura4+ ade6+ leu1+ ura4-*	This study.
YSM3802	*exo84-GFP-Kan+ ura4-::Ptdh1-CIBN-mTagBFP2-RitC-Adh1term-Pact1-CRY2PHR-mCherry:ura4+ ade6+ leu1+ ura4-*	This study.
YSM3807	*h- fim1-mCherry-Nat+ exo70-GFP-Kan+ ade6+ leu1+ ura4+*	This study.
YSM3809	*ypt3-I5(ts) leu1-32::CRIB-3GFP-Kan+::leu1+ ura4-::Ptdh1-CIBN-mTagBFP2-RitC-Adh1term-Pact1-CRY2PHR-mCherry::ura4+ ade6+ leu1-32 ura4-*	This study.
YSM3808	*sty1::bsd leu1-32::CRIB-3GFP-kanMX-leu+ ura4-::Ptdh1-CIBN-mTagBFP2-RitC-Adh1term-Pact1-CRY2PHR-mCherry::ura4 ade6+ leu1-32 ura4-*	This study.
Fig 3, Fig S6		
YSM3568	*h- leu1-32::ppak1-CRIB-3xGFP-KanMX+:leu1+ ura4-D18::Ptdh1-CIBN-mTagBFP2-Ritc-ADH1term-pact1-CRY2PHR-mCherry:ura4+ ade6+ leu1-32*	(*2*)
YSM3564	*ura4-294::pact1-CRY2PHR-mCherry::ura4+ ade6+ leu1+ ura4-294*	(*2*)
Fig 4, Fig S6, S7, Fig S8		
YSM3811	*h- ura4+::pact1:sfGFP-3RitCb:terminatortdh1 ade6-M210 leu+*	This study.
YSM3812	*h- ura4+::pact1:sfGFP-2RitC:terminatortdh1 ade6-M210 leu+*	This study.
YSM3813	*h- ura4+::pact1:sfGFP-1RitC:terminatortdh1 ade6-M210 leu+*	This study.
YSM3814	*h- ura4::pact1-3sfGFP-1RitC-tdhterminator::ura4+ ade6-M210 leu1+*	This study.
YSM3815	*h+ ura4+::pact1:sfGFP-1RitC:terminatortdh1 ade6+::pact1:mCherry-1RitC:termScADH1 leu+*	This study.
YSM3816	*h- ura4+::pact1:sfGFP-2RitC:terminatortdh1 ade6+::pact1:mCherry-1RitC:termScADH1 leu+*	This study.
YSM3817	*h+ ura4+::pact1:sfGFP-3RitCa:terminatortdh1 ade6+::pact1:mCherry-1RitC:termScADH1 leu+*	This study.
YSM3818	*ura4::pact1-3sfGFP-1RitC-tdhterminator ade6+::pact1:mCherry-1RitC:termScADH1 leu1+*	This study.
YSM3819	*ura4+::pact1:sfGFP-1RitC:terminatortdh1 his5+::pact1:CRIB[gic2aa1-181]-3mCherry::bsdMX ade6+ leu1+*	This study.
YSM3820	*ura4+::pact1:sfGFP-2RitC:terminatortdh1 his5+::pact1:CRIB[gic2aa1-181]-3mCherry::bsdMX ade6+ leu1+*	This study.
YSM3821	*ura4+::pact1:sfGFP-3RitCb:terminatortdh1 his5+::pact1:CRIB[gic2aa1-181]-3mCherry::bsdMX ade6+ leu1+*	This study.
YSM3822	*ura4::pact1-3sfGFP-1RitC-tdhterminator::ura4+ his5+::pact1:CRIB[gic2aa1-181]-3mCherry::bsdMX ade6+ leu1+*	This study.
YSM3905	*sty1Δ::kanMX ura4+::pact1:sfGFP-3RitCb:terminatortdh1 his5+::pact1:CRIB[gic2aa1-181]-3mCherry::bsdMX ade6+ leu1+*	This study.
YSM1916	*h+ leu1-32:pact1:LifeAct-GFP:leu1+ leu1- ura4-D18*	(*91*)
Fig 5, Fig S7, S8		
YSM3823	*ura4-294:pshk1:CRIB-3GFP:ura4+ rga4-RFP:kanMX leu1-32*	This study.
YSM3826	*h+ rga4Δ::kanMX ade6-M210 leu1-32 ura4-D18*	This study.
YSM3827	*h+ rga4Δ::kanMX ura4::pact1-rga4(1-685)-GFP::ura4+ ade6-M210 leu1-32*	This study.
YSM3828	*h+ rga4Δ::kanMX ura4::pact1-rga4(1-623)-GFP:ura4+ ade6-M210 leu1-32*	This study.
YSM3829	*h+ rga4Δ::kanMX ura4::pact1-rga4(1-470)-GFP::ura4+ ade6-M210 leu1-32*	This study.
YSM3830	*h+ rga4Δ::kanMX ura4::pact1-rga4(1-623)-RitC-GFP::ura4+ ade6-M210 leu1-32*	
YSM3831	*h+ rga4Δ::kanMX ura4::pact1-rga4(607-685)-GFP:ura4+ ade6-M210 leu1-32*	This study.
YSM3832	*h+ rga4Δ::kanMX ura4::pact1-rga4(465-685)-GFP::ura4+ ade6-M210 leu1-32*	This study.
YSM3833	*h+ rga4Δ::KanMx ura4-pact1-rga4(465-685-F630A-L633A-L640A)-GFP::ura4+ ade6-M210 leu1-32*	This study.

YSM3834	*h+ rga4Δ::kanMX ura4::pact1-rga4(465-685KR670-1AA; KR675-6AA)-GFP::ura4+ ade6-M210 leu1-32*	This study.
YSM3835	*h+ rga4Δ::kanMX ura4::pact1-rga4(465-685-F630A-L633A-L640A- KR670-1AA; KR675-6AA)-GFP::ura4+ ade6-M210 leu1-32*	This study.
YSM3836	*h+ rga4::kanMX ura4::pact1-rga4(465-666)-GFP::ura4+ ade6-M210 leu1-32*	This study.
YSM1408	*h- rga4-GFP:kanMX ade6+ leu1-32 ura4-D18*	(*29*)
YSM3837	*h- rga4-F630A,-L633A,-L640A-GFP-natMX ade6- leu1- ura4-*	This study.
YSM3838	*h- rga4-KR670/671AA-KR675/676AA-R783G-GFP-NAT ade6- leu1- ura4-*	This study.
YSM3839	*h- rga4-F630A,-L633A,-L640A-KR670/671AA,-KR675/676AA-GFP-natMX ade6- leu1- ura4-*	This study.
YSM3840	*rga3Δ::hphMX rga6Δ::kanMX rga4-GFP:kanMX ade6- leu1- ura4-*	This study.
YSM3841	*rga3Δ::hphMX rga6Δ::kanMX rga4-FLL-GFP-kanMX ade6- leu1- ura4-*	This study.
YSM3842	*rga6Δ::kanMX rga3Δ::hphMX rga4-KR670/671AA-KR675/676AA-GFP-natMX ade6- leu1- ura4-*	This study.
YSM3843	*rga6Δ::kanMX rga3Δ::hphMX rga4-F630A,-L633A,-L640A-KR670/671AA,-KR675/676AA-GFP-natMX ade6- leu1- ura4-*	This study.
YSM3844	*h+ rga4Δ::kanMX ura4::pact1-rga4(4x607-685)-GFP:ura4+ ade6-M210 leu1-32*	This study.
YSM3845	*h+ rga4Δ::kanMX ura4::pact1-rga4 (4x465-666)-GFP-ura4+ ade6-M210 leu1-32*	This study.
YSM3846	*h+ rga4Δ::kanMX ura4::pact1-CRY2-mcherry-rga4(607-685)::ura4+ ade6-M210 leu1-32*	This study.
YSM3847	*h+ rga4Δ::kanMX ura4::ppom1-rga4(465-685)-GFP::ura4+ ade6-M210 leu1-32*	This study.
YSM3848	*h+ rga4Δ::kanMX ura4::ppom1-rga4(465-666)-GFP::ura4+ ade6-M210 leu1-32*	This study.

	Fig 6	
YSM3136	*h- rga6Δ::kanMX rga3Δ::hphMX rga4Δ::ura4+ ade6- leu1-*	(*29*)
YSM3907	*h- ade6::pact-CRY2-mCherry-RitC::ade6+ leu1-32::CRIB-3xGFP-kanMX::leu1+ ura4-D18*	This study.
YSM3906	*h- rga3Δ::hphMX rga4Δ::ura4+ rga6Δ::kanMX ade6::pact1-CRY2-mcherry-RitC::ade6+ leu1-32::CRIB-3xGFP-kanMX::leu1+*	This study.
YSM3824	*h- rga3Δ::hphMX rga4Δ::ura4+ rga6Δ::kanMX ade6::pact1-CRY2-mCherry-RitC-rga4GAP::ade6+ leu1-32*	This study.
YSM3825	*h- rga3Δ::hphMX rga4Δ::ura4+ rga6Δ::KanMX ade6::CRY2-mCherry-Ritc-rga4GAP::ade6+ leu1::CRIB-3GFP-KanMX:leu1+*	This study.

Table S2: Plasmids used in this study

Number	Description	Purpose
pAV0327	pUra4AfeI-pact1-sfGFP-terminatortdh1	*ura4* integration
pAV0328	pUra4AfeI-pact1-mCherry-terminatortdh1	*ura4* integration
pAV0356	pAde6$_{PmeI}$	*ade6* integration
pAV0605	pUra4$_{AfeI}$-pact1-sfGFP-1RitC-terminatortdh1	*ura4* integration
pAV0602	pUra4$_{AfeI}$-pact1-sfGFP-2RitC-terminatortdh1	*ura4* integration
pAV0601	pUra4$_{AfeI}$-pact1-sfGFP-3RitC-terminatortdh1	*ura4* integration
pAV0607	pAde6$_{PmeI}$-pact1-mCherry-1RitC-termScAdh1	*ade6* integration
pSM2652	pUra4$_{AfeI}$-pact1-3sfGFP-1RitC-tdhterminator	*ura4* integration
pSM0621	pREP41-GFP	cloning backbone
pSM2672	pUra4$_{AfeI}$-pact1-rga4(aa1-685)-GFP-terminatortdh1	*ura4* integration
pSM2641	pUra4$_{AfeI}$-pact1-rga4(aa1-623)-GFP-terminatortdh1	*ura4* integration
pSM2673	pUra4$_{AfeI}$-pact1-rga4(aa1-470)-GFP-terminatortdh1	*ura4* integration
pSM2642	pUra4$_{AfeI}$-pact1-rga4(aa1-623)-RitC-GFP-terminatortdh1	*ura4* integration
pSM2617	pUra4$_{AfeI}$-pact1-rga4(aa607-685)-GFP-terminatortdh1	*ura4* integration
pSM2381	pUra4$_{AfeI}$-pact1-rga4(aa465-685)-GFP-terminatortdh1	*ura4* integration
pSM2424	pUra4$_{AfeI}$-pact1-rga4(aa465-685-F630A, L633A, L640A)-GFP-terminatortdh1	*ura4* integration
pSM2383	pUra4$_{AfeI}$-pact1-rga4 (aa465-685-KR670-1AA; KR675-6AA)-GFP-terminatortdh1	*ura4* integration
pSM2480	pUra4$_{AfeI}$-pact1-rga4 (aa465-685-F630A, L633A, L640A ;KR670-1AA; KR675-6AA)-GFP-terminatortdh1	*ura4* integration
pSM2382	pUra4$_{AfeI}$-pact1-rga4 (aa465-666)-GFP-terminatortdh1	*ura4* integration
pSM2635	pUra4$_{AfeI}$-pact1-4xrga4(aa607-685)-GFP-terminatortdh1	*ura4* integration
pSM2479	pUra4$_{AfeI}$-pact1-4xrga4(aa465-666)-GFP-terminatortdh1	*ura4* integration
pSM2384	pUra4$_{AfeI}$-ppom1-rga4(aa465-685)-GFP-terminatortdh1	*ura4* integration
pSM2385	pUra4$_{AfeI}$ -ppom1-rga4(aa465-666)-GFP-terminatortdh1	*ura4* integration
pSM2637	pUra4$_{AfeI}$-pact1-CRY2-mcherry-rga4(aa607-685)	*ura4* integration
pSM2919	pAde6$_{PmeI}$-pact1-CRY2-mCherry-RitC	*ade6* integration
pSM2787	pAde6$_{PmeI}$-pact1-CRY2-mCherry-RitC-rga4(aa665-933)	*ade6* integration
pSM1692	pSp72-112bp$_{upstream}$-Rga4$_{ORF}$(F630A, L633A, L640A)-+112bp$_{downstream}$	mutagenesis; target endogenous locus
pSM1421	pSp72-112bp$_{upstream}$-Rga4$_{ORF}$(KR670/671AA : KR675/676AA)-+112bp$_{downstream}$	mutagenesis; target endogenous locus
pSM1694	pSp72-112bp$_{upstream}$-Rga4$_{ORF}$(F630A, L633A, L640A, KR670/671AA : KR675/676AA)-+112bp$_{downstream}$	mutagenesis; target endogenous locus
pAV0046	pJK148-ppak1-Ntermgic2aa2-181-3GFP-KanMX	*leu32* integration
pSM2366	pAde6-Pnmt41-GFP-Ypt3	*ade6* integration

Table S3: Primers used for mutagenesis. Bold residues indicate changes from the wildtype sequence

Number	Mutation	5'-3' sequence
osm2347	rga4 (F630A;L633A)_for	CTCCGCGAGAATCG**GC**TTCGCGT**G**CACAAATGGTTGCATC
osm2348	rga4 (F630A;L633A)_rev	GATGCAACCATTTGT**G**CACGCGAA**GC**CGATTCTCGCGGAG
osm2349	rga4 (L640A)_for	AATGGTTGCATCTAGC**GC**TGGATTTAGACCCAAAG
osm2350	rga4 (L640A)_rev	CTTTGGGTCTAAATCCA**GC**GCTAGATGCAACCATT
osm2351	rga4 (KR670/671AA:KR675/676AA)_for	AGAAGTCATTTTCT**GCAGCG**TTTCATTGG**GCAGCA**GATTCTCCTCACAT
osm2352	rga4 (KR670/671AA:KR675/676AA)_rev	ATGTGAGGAGAA**TCTGCT**GCCCAATGAAA**CGCTGC**AGAAAATGACTTCT

Figure S1. Controls and further evidence of depletion of membrane-associated CRY2 around sites of secretion

A. Time-lapse of CRY2-mCherry, CIBN-mTag-BFP2-RitC and CRIB-3GFP in a pre-divisional cell grown in the dark. Time 0 is the first timepoint after illumination. **B.** Time-lapse of CRY2-mCherry redistribution in cells co-expressing CIBN-mTag-BFP2-RitC and GFP-Ypt3 (see Fig 1B for quantification). **C-D.** Time-lapse of CRY2-mCherry redistribution in cells co-expressing CIBN-mTag-BFP2-RitC and Exo84-GFP (C) or Exo70-GFP (D). The bottom graph shows correlation between extent of CRY2 depletion after 5 min and pole GFP intensity. In (B-D), cells were grown in the dark. Time 0 is the first timepoint after illumination. The GFP channel is shown as sum projection over the 5 min time-lapse. The mCherry channel shows individual time points. **E.** Correlation between CRY2 depletion halftime and extent of depletion after 5 min. **F.** Example of Anp1-GFP-expressing cells mixed with cells shown in Fig 1C and panel (G), showing the effect of brefeldin A (BFA) on collapse of Golgi to the ER. **G.** Cells as in (B) treated with 50nM brefeldin A (BFA) or solvent (EtOH). **H.** Cells as in Fig 1D, but at 25°C. In (G-H), the GFP channel is shown as sum projection over the 5 min time-lapse. The mCherry channel shows individual time points. Graphs on the right are as in Fig1C-E with depletion cut-off shown in red and green backgrounds. Data points are coloured with different shades of green according to GFP fluorescence levels. The indicated Kruskal-Wallis test *p*-value compares CRY2 depletion values between cells of equivalent CRIB-3GFP intensities. **I.** Example of cell expressing both Fim1-mCherry and Exo70-GFP mixed with cells shown in Fig 1E, showing the effect of Latrunculin A (LatA) on loss of actin patches (but not exocyst localization). Throughout, yellow arrowheads indicate depletion zones. Scale bars are 3μm.

Figure S2. Simulation results and conditions for tip depletion and lateral peak.

A. Time evolution of particles (green) on the surface of the sphere with $k_{off} = 10^{-5}/s$ and $D = 10^{-4}\ \mu m^2/s$ under static conditions (top row; net area added is zero with 0.5 endocytosis events/s and 2.44 endocytosis events/s) and under net growth conditions (bottom row; net area added 20.7 $\mu m^2/hr$ with 0.68 exocytosis events/s and 2.44 endocytosis events/s). The spherical shell representing the cell remains at a constant size; exocytosis and endocytosis only affect the positions of particles on the surface. Left: both rows have $\sigma_{exo} = 0.85\ \mu m < \sigma_{endo} = 1.84\ \mu m$ derived from experimental measurements; depletion near the top tip is evident in both cases but occurs faster under growth conditions. Right: both rows have $\sigma_{exo} = \sigma_{endo} = 1.84\ \mu m$; slight depletion near the tip is only evident under growth conditions. **B.** Scaled number density as a function of arclength for the same simulations as Fig 2E but under growth conditions. Compared to Fig 2E, the lateral peak is shifted further towards the back of the sphere and tip depletion is stronger. **C.** Tip occupation ratio as a function of k_{off} for static and growth conditions for $\sigma_{exo} = \sigma_{endo}$ and $\sigma_{exo} < \sigma_{endo}$ as in panel A, for $D = 10^{-3}\ \mu m^2/s$. Switching from $\sigma_{exo} = \sigma_{endo}$ to $\sigma_{exo} < \sigma_{endo}$ or from static to growth conditions both result in increased tip depletion. **D.** Increasing diffusion coefficient while keeping the tip occupancy ratio nearly constant reduces the lateral peak height (net growth, $\sigma_{exo} < \sigma_{endo}$ as in A). **E.** Scaled number density as a function of arclength, for simulations where exocytosis is turned off/endocytosis remains (left); endocytosis is turned off/exocytosis remains the same as under growth conditions (middle); and endocytosis is turned off/exocytosis is reduced so that the net area addition rate is 20.7 $\mu m^2/hr$ (right). Reference parameters are for net growth, $\sigma_{exo} < \sigma_{endo}$ as in (A). Stopping exocytosis leads to tip enhancement for low enough k_{off}. As these simulations do not account for anticipated slowing down of endocytosis due to reduction of plasma membrane lipid reservoir, we consider them consistent with the results of experiments in Fig 1C-D. Stopping endocytosis maintains tip depletion, in line with experiments of Fig 1E. **F.** Scaled number density as a function of arclength at $D = 10^{-4}\ \mu m^2/s$ under no flow displacement conditions where exocytosis/endocytosis does not move the particles on the sphere surface but endocytosis pulls particles into the internal pool if they are within R_{endo} of the endocytosis event (using 2.44 endocytosis events/s and $\sigma_{endo} = 1.84\ \mu m$ as in (A)). Depletion is still evident but the lateral peak near the equator of the sphere disappears. Compare to Fig 2E. **G.** Tip occupation ratio versus k_{off} under no flow displacement conditions as in (F). Tip depletion is less pronounced compared to simulations with displacement in Fig 2D, for the same conditions. **H.** Scaled number density as a function of arclength at $D = 10^{-4}\ \mu m^2/s$ under both flow displacement conditions and endocytosis pulling particles into the internal pool if they are within R_{endo} of the endocytosis event (using 2.44 endocytosis events/s and $\sigma_{endo} = 1.84\ \mu m$ as in (A)). **I.** Tip occupation ratio versus k_{off} under flow displacement conditions and endocytic removal as in (H).

Figure S3. Distribution of exo- and endocytosis in fission yeast cells

A-B. Time-lapse, sum projection and kymographs of the cell pole (A) and the early, pre-constriction division site (B) of cells expressing Exo70-GFP and Fim1-mCherry, marking exo- and endocytic sites, respectively. The graph shows the average distribution of these markers. Shaded areas show standard deviation.

Figure S4. Fitting for diffusion coefficient and membrane dissociation rate with FRAP.

A. A rectangular region along the cylindrical section on one side of the cell was photobleached and the recovery imaged over time. **B.** Schematic of 1D fluorescence recovery model for an experimental profile, extracted by vertical projection along a rectangular region shown in yellow, for the frame after photobleaching ($t = 0$, black circles). The boundaries between the experimentally-sampled middle region and the left and right flanking regions are indicated by dashed vertical lines. **C.** Example recovery curves. Single cell expressing 1xRitC-GFP imaged every 0.729 s and fit using the 1D model over 30 frames. Experimental data shown as lines with filled points while model fits shown as lines without points ($D = 6.85 \times 10^{-3} \ \mu m^2/s$, $k_{off} = 0.108 \ /s$ and l_F as shown by the end points of the model curves). The black line of the model is

initialized to go through the experimental data points at $t = 0$. **D-F.** Plots of averaged R^2 calculated from FRAP fits for all protein constructs investigated as function of D and k_{off}. Red circles indicate the median of the D and k_{off} values for each protein from individual fits of the 1D model while blue dashed circles indicate the best fit using the average of individual R^2 maps. The number of cells averaged in each plot is indicated by the value of n. **G.** Averaged R^2 plots for simulated FRAP due to both diffusion and membrane binding/unbinding on the surface of a cylinder. Good agreement is found between the input parameters (red circle or dashed line) and the highest value of R^2 (blue circle).

Figure S5. Modified model to include dimer formation on spherical surface, diffusing in the presence of membrane flows.

A. Reactions used in dimerization simulations. The diffusion coefficient of dimers is the same as that of monomers. Reactions obey detailed balance at steady state in absence of endocytic or exocytic flow. **B.** Plots of the number of average particles at steady state as a function of parameter p. Simulations in this and panels D and E have 5000 particles and rate constants as described in methods. **C.** Snapshots of spheres with increasing p value show that tip depletion and lateral concentration increase with increasing p. Color scale shows local density. **D.** Scaled number density of particles as a function of arclength show a similar trend where depletion and lateral peak increase with increasing p. **E.** Tip occupation ratio also decreases as a function of p. **F.** Plots of the number of average particles at steady state as a function of parameter p. Simulations in this and panels G and H have 5000 particles and dimer off rates ($k_{off,2}$ and k_{-}) equal to a quarter of the single particle off rate (k_{off}). **G.** Scaled number density of particles as a function of arclength have an enhanced depletion and lateral peak compared to panel D. **H.** Tip occupation ratio is slightly decreased compared to panel E.

Figure S6. Membrane affinity-dependent depletion of membrane-associated RitC at sites of polarized secretion.

A. Localization of indicated sfGFP-tagged RitC constructs in cells co-expressing CRIB-mCherry. The graphs on the right show correlation plots between the ratio of the tip to side GFP signal and the CRIB-mCherry tip intensity. Note the progressive shift towards the left of the data points upon RitC multimerization, as well as the more marked negative correlation. **B.** sfGFP-3xRitC and CRIB-3xmCherry in *sty1Δ* cells non-treated or treated with LatA for 30min. The cells were mixed with LifeAct-GFP expressing cells as control, showing actin depolymerization upon LatA treatment.

Figure S7. Structure-function analysis of the Cdc42 GAP Rga4 defining membrane-binding and oligomerization motifs

A. Scheme of Rga4 and truncation fragments. The sequence alignment shows the CBD across *Schizosaccharomyces* species (*S.p.* = *S. pombe*; *S.o.* = *S. octosporus*; *S.c.* = *S. cryophilus*; *S. j.* = *S. japonicus*). Membrane-binding (MB) motif 1 shows a weak amphipathic helix prediction, of which three hydrophobic residues (FLL, red circles and asterisks) were mutated to alanine. MB motif 2 was identified to have a high basic and hydrophobic (BH) scale, predictive of membrane association (*28*). Asterisks highlight four basic residues (KRKR) that were mutated to alanine. **B.** Shape of *rga3Δ rga6Δ* double mutant cells with indicated *rga4* alleles shown by calcofluor imaging. **C.** Dissolution of cytosolic condensates of Rga4 N-terminus lacking the CBD (N-ΔCBD) upon treatment with 10% 1,6-hexanediol or 37°C. **D.** Cytosolic condensates of full-length Rga4 with mutations in MB #1 and #2 are dissolved upon treatment with 5% 1,6-hexanediol or 37°C.

Figure S8. Further characterization of minimal Rga4 fragments

A. Localization of CRY2-CBD in cells grown in the dark and illuminated at time 0. **B.** Localization of cc-CBD and cc-CBD-Δ2 Rga4 fragments by TIRF-microscopy. See Movies 6-7 for time-lapse imaging and cluster tracking. **C.** Quantification of track duration, speed and fluorescence intensity for clusters of Rga4 fragment as in (B). The track duration of cc-CBD-Δ2 clusters was significantly reduced, while their speed was significantly increased, consistent with shorter membrane residence time and faster lateral diffusion. The track duration times and diffusion coefficient implied by the measurements (displacement per time step $\approx (4Dt)^{1/2}$) are in the same range as the median k_{off} and D of the FRAP fits in Fig S4F. p-values calculated from student t-test with unequal variances. The fluorescence intensity per cluster was not altered, suggesting unchanged oligomerization properties, as predicted from truncation of the membrane-binding region.

Figure S9. Model calibration and dependence on parameters k_{on}, α, R_{cutoff}, γ.

A. Test of implementation of diffusion on a sphere. Change in polar angle due to diffusion maintains the correct scaling of particle displacement over time, regardless of the timestep. Colored curves show results of averaged mean squared polar angle displacement (in radians squared) for particles diffusing on a sphere with $D = 10^{-3}$ μm²/s, $R = 2$ μm. Dashed lines show asymptotic limits at short and long times from (55). The colored curves overlap, each starting from the smallest time, $t = dt$. **B.** Limited vertical resolution of microscope, b, leads to varying fractions of the cell surface being observed at a given arc length, s. **C.** Test of Equation (M4): if only events in a slab of height b are counted, sampling in 3D from $P(s)$ recovers the experimentally observed probability distribution, $P_{exp}(s)$, for both the endocytosis and exocytosis profiles. **D.** Varying k_{on} has no effect on the tip occupation ratio over the range of D and k_{off} investigated, as expected. **E.** Lower (higher) values of the cutoff range of exocytosis/endocytosis, R_{cutoff}, leads to a reduction (increase) in tip depletion, along with reduction and leftward (increase and rightward) shift of the lateral peak. Other results in this paper used $R_{cutoff} = 2$ μm. **F.** Parameter α (fraction of flowing membrane) affects tip depletion with lower (higher) values of α lead to enhanced (reduced) tip depletion and lateral peak. Other results in this paper used $α = 0.5$. Increasing parameter γ leads to a reduced tip depletion and lateral peak as proteins are less influenced by the flowing lipids.

Movie legends

Movie 1. Depletion of membrane-associated CRY2 at cell poles

Spinning disk microscopy time-lapse of CRY2-mCherry, CIBN-mTag-BFP2-RitC and CRIB-3GFP cells grown in the dark. Time 0 is the first timepoint after illumination. Images are at 30s interval. The UV channel (CIBN-mTagBFP2-RitC) was acquired every 4 timepoints. Depletion is observed at both poles of the three medial cells. Note that cells containing only CRY2-mCherry, which were used for normalization purposes, are also visible. Scale bar is 5μm.

Movie 2. Depletion of membrane-associated CRY2 at division sites

Spinning disk microscopy time-lapse of CRY2-mCherry, CIBN-mTag-BFP2-RitC and CRIB-3GFP cells grown in the dark. Time 0 is the first timepoint after illumination. Images are at 30s interval. The UV channel (CIBN-mTagBFP2-RitC) was acquired every 4 timepoints. Depletion is observed initially at cell poles and then, as cells enter mitosis and prepare for cytokinesis, at mid-cell, concomitant with CRIB-3GFP accumulation. Scale bar is 5μm.

Movie 3. Simulation of membrane-associated protein depletion by membrane flows

Simulation under static conditions (equal rates of membrane delivery by exo- and endocytosis), with $k_{off} = 10^{-5}$/s and $D = 10^{-4}$ μm^2/s showing depletion from the zone of exocytosis on the top of the sphere.

Movie 4. Simulation showing lack of membrane flow-dependent depletion for a protein with fast membrane unbinding rate

Simulation under static conditions (equal rates of membrane delivery by exo- and endocytosis), with $k_{off} = 10^{-2}$/s and $D = 10^{-4}$ μm^2/s showing absence of depletion from the zone of exocytosis on the top of the sphere.

Movie 5. Localization of 3xRitC during the cell growth cycle

Airyscan microscopy time-lapse of sfGFP-3xRitC and CRIB-3xmCherry in cells during mitotic growth. 3xRitC depletes from zones of growth labelled by CRIB at cell poles during interphase and repopulates the cell poles during mitosis when polarized cell growth stops. 3xRitC also transiently depletes from mid-cell, labelled by CRIB in pre-divisional cells until the initiation of septum invagination. White arrowheads show depletion zones. Scalebar is 5μm.

Movie 6. Dynamics of Rga4 cc-CBD fragment in TIRF microscopy

Tracking of cortical clusters of the Rga4 cc-CBD fragment in cells imaged by TIRF microscopy detected by TrackMate ImageJ plugin. Detected clusters are circles. All trajectories are shown.

Movie 7. Dynamics of Rga4 cc-CBD-Δ2 fragment in TIRF

Tracking of cortical clusters of the Rga4 cc-CBD–Δ2 fragment in cells imaged by TIRF microscopy detected by TrackMate ImageJ plugin. Detected clusters are circles. All trajectories are shown.

Movie 8. Light-induced polarity establishment in *rga3Δ rga4Δ rga6Δ* optoGAP cells – example of monopolar growth

Airyscan microscopy time-lapse of *rga3Δ rga4Δ rga6Δ* cells expressing CRIB-3GFP and optoGAP grown in the dark and illuminated with blue light from t = 0. OptoGAP, shown in magenta, initially decorates the entire cell periphery, with CRIB absent from the cell cortex. The cells initiate polarized growth, with depletion of optoGAP, and accumulation of CRIB in a restricted zone.

Movie 9. Light-induced polarity establishment in *rga3Δ rga4Δ rga6Δ* optoGAP cells – example of bipolar growth

Airyscan microscopy time-lapse of *rga3Δ rga4Δ rga6Δ* cells expressing CRIB-3GFP and optoGAP grown in the dark and illuminated with the blue light from t = 0. OptoGAP, shown in magenta, initially decorates the entire cell periphery, with CRIB absent from the cell cortex. The cells initiate 2 zones of polarized growth, with depletion of optoGAP, accumulation of CRIB in a restricted zone and outgrowth of a tube.

Movie 10. Long-range movement of optoGAP in dividing in *rga3Δ rga4Δ rga6Δ* cells

Airyscan microscopy time-lapse of *rga3Δ rga4Δ rga6Δ* cells expressing CRIB-3GFP and optoGAP grown in the light. OptoGAP, shown in magenta, strongly decorates the back of these monopolar cells, with the site of growth indicated by CRIB localization. As cells enter division, growth ceases, CRIB disappears from the cell pole and reappears at the division site. During cell division, optoGAP exhibits long-range movements away from the division site towards the previously growing cell end.