QTL Mapping in Outbred Tetraploid (and Diploid) Diallel Populations

Rodrigo R. Amadeu*, Patricio R. Munoz*, Chaozhi Zheng[†], Jeffrey B. Endelman[‡]

*Horticultural Sciences Department, University of Florida, Gainesville, FL, USA

[†]Biometris, Wageningen University and Research, Wageningen, The Netherlands

[‡]Department of Horticulture, University of Wisconsin, Madison, WI, USA

Running title: QTL mapping in tetraploid diallel

Key words: multiparental, polyploidy, dominance, haplotypes

Corresponding author:

Jeffrey B. Endelman, University of Wisconsin–Madison

1575 Linden Dr., Madison, WI 53706

Phone: 608-250-0754

E-mail: endelman@wisc.edu

ABSTRACT

Over the last decade, multiparental populations have become a mainstay of genetics

research in diploid species. Our goal was to extend this paradigm to autotetraploids by

developing software for quantitative trait locus (QTL) mapping in connected F1

populations derived from a set of shared parents. For QTL discovery, phenotypes are

regressed on the dosage of parental haplotypes to estimate additive effects. Statistical

properties of the model were explored by simulating half-diallel diploid and tetraploid

populations with different population sizes and numbers of parents. Across scenarios,

the number of progeny per parental haplotype (pph) largely determined the statistical

power for QTL detection and accuracy of the estimated haplotype effects. Multi-allelic

QTL with heritability 0.2 were detected with 90% probability at 25 pph and genome-wide

significance level 0.05, and the additive haplotype effects were estimated with over 90%

accuracy. Following QTL discovery, the software enables a comparison of models with

multiple QTL and non-additive effects. To illustrate, we analyzed potato tuber shape in a

half-diallel population with 3 tetraploid parents. A well-known QTL on chromosome 10

was detected, for which the inclusion of digenic dominance lowered the Deviance

Information Criterion (DIC) by 17 points compared to the additive model. The final

model also contained a minor QTL on chromosome 1, but higher order dominance and

epistatic effects were excluded based on the DIC. In terms of practical impacts, the

software is already being used to select offspring based on the effect and dosage of

particular haplotypes in breeding programs.

INTRODUCTION

For over three decades, the genetic mapping of quantitative trait loci (QTL) using DNA markers has been essential to basic and applied research. Early studies in plants focused on experimental diploid populations created from two parents (Lander and Botstein 1989; Knapp *et al.* 1990). For inbred parents, genetic marker alleles are easily coded to uniquely identify the two parental haplotypes at each locus (Young and Tanksley 1989). For two outbred parents of ploidy ϕ, there are 2ϕ parental haplotypes (or homologs), but in the case of biallelic SNPs there are only two marker alleles. This difficulty was initially circumvented by using single-dose markers to create separate maternal and paternal maps in diploid species (Grattapaglia and Sederoff 1994) and separate maps for each homologous chromosome in polyploids (Wu *et al.* 1992). Advances in estimation theory (Ritter and Salamini 1996; Wu *et al.* 2002) and software (Van Ooijen and Voorrips 2001; Margarido *et al.* 2007) eventually made it possible to reconstruct F1 diploid offspring in terms of their parental haplotypes, and QTL mapping soon followed (Van Ooijen 2004). Inference of parental haplotypes in F1 polyploid offspring based on bi-allelic SNPs is even more challenging, but several software packages are now available for linkage and QTL analysis (Hackett *et al.* 2017; Bourke *et al.* 2018; Mollinari and Garcia 2019; da Silva Pereira *et al.* 2020).

Not long after QTL mapping in biparental populations became a reality, several groups explored the use of multiple, connected biparental populations via theory and simulation (Rebai and Goffinet 1993; Muranty 1996; Liu and Zeng 2000). We use the term diallel to represent a general class of mating designs in which groups of offspring are derived from no more than two founders (Fig. 1). The transition from theory to

practice for diallel mapping began in maize with 25 diverse inbred lines mated to a

common parent (Yu *et al.* 2008; Buckler *et al.* 2009), and similar efforts have been

published for other crops (Nice *et al.* 2017; Song *et al.* 2017). Examples of QTL

mapping in connected F1 populations derived from outbred diploids have also been

published (Rosyara *et al.* 2013; Bink *et al.* 2014). Compared with MAGIC (multiparent

advanced generation intercross) designs, in which offspring contain haplotypes from

more than two founders (Zhang *et al.* 2014; Huang *et al.* 2015), diallel designs have the

advantage of mimicking how plant breeding programs typically work. This is particularly

important for breeding programs with limited resources to create specialized populations

for discovery research.

The objective of this research was to develop and apply software for QTL mapping

in autotetraploid diallel populations (i.e., connected F1 populations), for which no other

software is available. Our software, named diaQTL and available as a package for the R

Computing Environment (R Core Team 2020), estimates additive effects by regression

of phenotypes on the dosage of parental haplotypes. For a diallel with p tetraploid

parents, there are $4p$ parental haplotypes (some of which may contain identical alleles

at a given QTL). diaQTL computes parental haplotype dosage based on output from the

companion software PolyOrigin, which estimates parental genotype probabilities using

hidden Markov models to perform multi-point linkage analysis (Zheng *et al.* 2021).

diaQTL can also analyze connected F1 populations from outbred diploid parents, using

parental genotype probabilities from the software RABBIT (Zheng *et al.* 2015).

Our software is also unique for its ability to model dominance effects in tetraploid

linkage mapping populations. The dominance deviation for one locus is the residual

genetic effect with the additive model. In polyploids, a hierarchy of dominance effects

can be defined by regression of the dominance deviation on combinations of parental

haplotypes (Kempthorne 1957). Digenic dominance effects are the regression

coefficients for a pair of haplotypes, or diplotype; trigenic dominance effects are the

regression coefficients for triplotypes; etc.

After exploring the statistical properties of QTL mapping in outbred diallel

populations via simulation, we illustrate the use of diaQTL to analyze tuber shape in a

tetraploid potato diallel with 3 parents.

MATERIALS AND METHODS

QTL Model

A mixed model is used to relate phenotypes to the effects of parental haplotypes

and potential covariates. For the t^{th} measurement (e.g., plot) of clone i, the following

equation specifies the additive model for m QTL in a diallel with p parents of ploidy ϕ

(bold font designates a vector):

$$y_{it} = \mu + \boldsymbol{x}_{it}\boldsymbol{c} + \sum_{j=1}^{m}\sum_{k=1}^{\phi p} W_{ijk}\alpha_{jk} + \varepsilon_{it} \tag{1}$$

The response variable y_{it} denotes the phenotype; μ is the intercept; $\boldsymbol{c}$ is an optional

column-vector of fixed effects with corresponding row-vector of covariates $\boldsymbol{x}_{it}$; the

additive QTL effects α_{jk} are random regression coefficients corresponding to the

dosage of parental haplotype k at locus j, denoted W_{ijk}; and ε_{it} is the residual.

The dosage of parental haplotype k at locus j, denoted W_{ijk}, is calculated from an

input file containing the parental genotype probabilities for every offspring. For

haplotypes originating in founders that are not the parents of clone i, W_{ijk} is trivially zero.

For the other haplotypes, W_{ijk} is an expectation over the 100 unique parental genotype

states in a biparental tetraploid F1 or 35 unique states in a uniparental (selfed) tetraploid

S1 (Zheng *et al.* 2021). If $w_k(abcd)$ denotes the dosage of parental haplotype k in

parental genotype $abcd$ (each letter represents a parental haplotype, not necessarily

unique), and $P_{ij}(abcd)$ is the genotype probability for clone i at locus j, then

$$W_{ijk} = \sum_{abcd} w_k(abcd)P_{ij}(abcd) \tag{2}$$

For diploids, the sum in Eq. 2 is over the 4 parental genotype states for a biparental F1

or 3 states for a uniparental S1.

The additive model can be extended to include dominance effects. Denoting the

digenic dominance effect for parental haplotypes k and k' at locus j as $\beta_{jkk'}$, which is a

random regression coefficient corresponding to diplotype dosage $W_{ijkk'}$, the additive +

digenic model is

$$y_{it} = \mu + \mathbf{x}_{it}\mathbf{c} + \sum_{j=1}^{m}\left(\sum_{k=1}^{p\phi} W_{ijk}\alpha_{jk} + \sum_{k=1}^{p\phi}\sum_{k'=k}^{p\phi} W_{ijkk'}\beta_{jkk'}\right) + \varepsilon_{it} \tag{3}$$

Because the order of the parental haplotypes in diplotype kk' does not matter, the sum

over k' in Eq. 3 is restricted to values $k' \geq k$. For partial diallel designs, not all

diplotypes are present, and the sum over k' must be restricted accordingly.

Furthermore, digenic dominance may only be included for a subset of the m loci based

on model selection procedures (see below). The dosage $W_{ijkk'}$ is computed

analogously to Eq. 2, based on the expectation over parental genotypes. For

tetraploids, the formula is

$$W_{ijkk'} = \sum_{abcd} w_{kk'}(abcd) P_{ij}(abcd) \qquad (4)$$

The symbol $w_{kk'}(abcd)$ denotes the dosage of diplotype kk' in genotype $abcd$, which

equals the number of times kk' occurs in the $\binom{4}{2} = 6$ possible combinations of length 2

(sampling from $abcd$ without replacement). The definition of higher order dominance

effects is analogous to the digenic case and therefore not explicitly shown.

Additive x additive epistatic effects can also be modeled. Denoting the epistatic

effect for parental haplotypes k and k' at loci j and j', respectively, as $(\alpha\alpha)_{jkj'k'}$, the

extension of Eq. 1 is

$$y_{it} = \mu + \boldsymbol{x}_{it}\boldsymbol{c} + \sum_{j=1}^{m}\left(\sum_{k=1}^{p\phi} W_{ijk}\alpha_{jk} + \sum_{j'>j}^{p\phi}\sum_{k=1}^{p\phi}\sum_{k'=k}^{p\phi} W_{ijkj'k'}(\alpha\alpha)_{jkj'k'}\right) + \varepsilon_{it} \qquad (5)$$

132 Typically, epistasis is not modeled for all pairs of loci (model selection procedures are

133 used), so limits for the sum over j' are not explicit. The dosage $W_{ijkj'k'}$ of the two-locus

134 diplotype is based on the expectation over the parental genotypes at both loci

135 (designated $abcd$ and $a'b'c'd'$, respectively):

$$W_{ijkj'k'} = \sum_{abcd} \sum_{a'b'c'd'} w_k(abcd)w_{k'}(a'b'c'd')P_{ij}(abcd)P_{ij'}(a'b'c'd') \qquad (6)$$

136   The meaning of the symbols in Eq. 6 is the same as in Eq. 2.

Polygenic Effects

Polygenic effects can be added to the QTL model for improved understanding of

trait genetic architecture and genomic prediction (latter is beyond the scope of this

manuscript). Each hierarchy of QTL effects discussed above has an analogous

polygenic effect with covariance based on identity-by-descent (IBD) relationships, which

can be computed using diaQTL function *IBDmat*.

The additive relationship $A_{ii'}$ equals ploidy times kinship, which is the probability

that a randomly chosen homolog in clone i is IBD at a particular locus to a randomly

chosen homolog in clone i' (Gallais 2003). For locus j, the probability that both

randomly chosen homologs are parental haplotype k is $\left(\frac{W_{ijk}}{\phi}\right)\left(\frac{W_{i'jk}}{\phi}\right)$, and thus the total

probability of IBD (for any parental haplotype) is $\sum_{k=1}^{\phi p}\left(\frac{W_{ijk}}{\phi}\right)\left(\frac{W_{i'jk}}{\phi}\right)$. Averaged over m

loci, the additive relationship becomes

$$A_{ii'} = \frac{\phi}{m}\sum_{j=1}^{m}\sum_{k=1}^{\phi p}\left(\frac{W_{ijk}}{\phi}\right)\left(\frac{W_{i'jk}}{\phi}\right) = \frac{1}{\phi m}\sum_{j=1}^{m}\sum_{k=1}^{\phi p}W_{ijk}W_{i'jk} \qquad (7)$$

Eq. 7 is computed for each chromosome, and the value used for the polygenic effect is

based on averaging all chromosomes that do not contain a QTL; in other words, by

extending the leave-one-chromosome-out concept of Yang *et al.* (2014).

IBD relationships for dominance and epistatic effects can be defined analogously to

additive relationship (Gallais 2003). The digenic dominance relationship $D_{ii'}$ equals the

binomial coefficient $\binom{\phi}{2}$ times the probability that a randomly chosen pair of homologs

in clone i are IBD at a particular locus to a randomly chosen pair in clone i'. For locus j,

the probability that both randomly chosen homologs are parental diplotype kk' is

$\left[\dfrac{W_{ijkk'}}{\binom{\phi}{2}}\right]\left[\dfrac{W_{i'jkk'}}{\binom{\phi}{2}}\right]$, which leads to the following expression:

$$D_{ii'} = \binom{\phi}{2}\frac{1}{m}\sum_{j=1}^{m}\sum_{k=1}^{\phi p}\sum_{k'=k}^{\phi p}\left[\frac{W_{ijkk'}}{\binom{\phi}{2}}\right]\left[\frac{W_{i'jkk'}}{\binom{\phi}{2}}\right] = \binom{\phi}{2}^{-1}\frac{1}{m}\sum_{j=1}^{m}\sum_{k=1}^{\phi p}\sum_{k'=k}^{\phi p}W_{ijkk'}W_{i'jkk'} \qquad (8)$$

Trigenic and quadrigenic dominance relationships are defined analogously. The additive

x additive epistasis relationship $E_{ii'}$ is also based on IBD probabilities for diplotypes, but

in this case the two haplotypes are at different loci (j and j') , so the probability that both

diplotypes are $jkj'k'$ is $\left[\dfrac{W_{ijkj'k'}}{\phi^2}\right]\left[\dfrac{W_{i'jkj'k'}}{\phi^2}\right]$ (using the same notation as Eq. 5). The

epistatic relationship is based on the average over v pairs of loci from different

chromosomes:

$$E_{ii'} = \frac{1}{\phi^2 v}\sum_{j}\sum_{j'>j}\sum_{k=1}^{\phi p}\sum_{k'=1}^{\phi p}W_{ijkj'k'}W_{i'jkj'k'} \qquad (9)$$

Bayesian Implementation

R package BGLR (Pérez and de los Campos 2014) was used to implement the

regression model due to its flexibility for modeling random effects and its ability to

handle binary phenotypes with a generalized linear model (in which case the left-hand

side of Eq. 1 is the linear predictor, not the phenotype). BGLR uses a Bayesian

framework and Markov Chain Monte Carlo (MCMC) to generate samples from the

posterior density. The additive haplotype effects for each QTL (Eq. 1) are independent

and identically distributed (i.i.d.) with a "BayesC" prior (Habier *et al.* 2011), which is a

two-component mixture of a normal and Dirac distribution (i.e., point mass). This prior

was chosen because it accommodates a greater range of complexity for the QTL allelic

series than a single normal. The mixing probability and variance of the normal

distribution are random (hyper)parameters with their own prior densities (beta and

inverse chi-squared, respectively), chosen according to the default rules implemented in

the BGLR software. The digenic dominance effects for each locus are i.i.d. with a

BayesC prior (separate from the prior for the additive effects), and the same holds for

higher order dominance or epistatic effects. The prior density for the additive polygenic

effect is multivariate normal with covariance matrix $\mathbf{A} \times V_{poly}$ ("RKHS" model in BGLR),

and the variance component V_{poly} has an inverse chi-squared prior. The residuals are

i.i.d. normal with variance V_{ε}, which has an inverse chi-squared prior.

If the QTL effects were modeled as fixed, Eq. 1 would be overparameterized, and

constraints would be needed to ensure estimability (Hackett *et al.* 2014). Although

constraints are not theoretically needed for random effects, in practice we observed

large variation in the sum of the additive effects during the Markov chain, even as the

differences between the additive effects remained fairly stable. Because ultimately it is

these differences that are biologically meaningful, and to reduce the Bayesian credible

interval (described below) for the genetic effects, a constant was added to all additive

effects (separately for each locus) at each iteration of the Markov chain to impose the

constraint of zero sum, and the model intercept was adjusted accordingly. Each group

of non-additive effects was similarly constrained to have zero sum.

QTL genetic variances were calculated for each iteration of the Markov chain from

the corresponding genetic effects. If $u_{ij} = \sum_{k=1}^{\phi p} W_{ijk} \alpha_{jk}$ denotes the additive value for

clone i due to locus j, the additive variance for that locus is $V_{a,j} = \mathrm{var}_i[u_{ij}] = \overline{u_{.j}^2} - (\overline{u_{.j}})^2$,

and non-additive variances were computed analogously. The total genetic variance for

locus j, denoted $V_{Q,j}$, is the sum of additive and non-additive variances. The proportion

of variance due to the additive effects for locus j is

$$h_j^2 = \frac{V_{a,j}}{V_\varepsilon + \lambda V_{poly} + \sum_{j'} V_{Q,j'}} \tag{10}$$

The parameter λ in Eq. 10 equals the mean diagonal of the additive relationship matrix,

to account for inbreeding in selfed populations (Endelman and Jannink 2012). The

proportion of variance due to non-additive or polygenic effects is calculated analogously

to Eq. 10.

Two kinds of Bayesian credible interval (CI) are computed in diaQTL. The CI for

model parameters is based on the quantiles of the Markov chain (after discarding the

burn-in iterations). The CI for QTL location uses a profile likelihood based on the

methodology in the R/qtl package (Broman and Sen 2009). Let $f_k = b(e^{LL_k}) d_k$ be the

probability of QTL location being in marker-bin k, where LL_k is the posterior mean of the

log-likelihood, d_k is the bin-width in cM, and the normalization constant b is chosen such

that $1 = \sum_k f_k$. The cumulative distribution at bin k is $F_k = \sum_{i=1}^{k} f_i$. The lower bin for the

CI with probability $1 - \alpha$ is one less than the largest k satisfying $F_k \leq \alpha/2$, and the upper

bin is one more than the smallest k satisfying $F_k \geq 1 - \alpha/2$.

The burn-in and total number of iterations for MCMC were determined using the

Raftery and Lewis (1992) diagnostic implemented in R package coda (Plummer *et al.*

2006). This diagnostic is based on estimating quantile q of a parameter within the

interval (*q-r, q+r*) with probability *s* (= 0.95 in our analysis). diaQTL function *set_params*

returns the value of the diagnostic for each of the genetic and residual variances, and

the number of iterations was chosen based on the largest value. For QTL discovery with

a one-dimensional scan, which relies on the posterior mean (see below), we used q =

0.5, *r* = 0.1. For estimating model parameters and the 90% CI at discovered QTL, we

used *q* = 0.05, *r* = 0.025.

Model Selection

In accordance with the statistical principle of parsimony, we used the Deviance

Information Criterion (DIC) to guide the selection of genetic models (Lenarcic *et al.*

2012; Pérez and de los Campos 2014). DIC uses the posterior mean deviance ($-2 \times$

log likelihood) to measure model fit and then adds a penalty for model complexity, which

equals the difference between the posterior mean deviance and the deviance evaluated

at the posterior mean (Spiegelhalter *et al.* 2002). The diaQTL package utilizes DIC for

both QTL discovery (with function *scan1*) and post-discovery exploration of multiple

QTL and non-additive models (with function *fitQTL*).

For QTL discovery, we used the conventional Neyman-Pearson hypothesis testing

framework, with the additive QTL model (Eq. 1) as the alternative hypothesis. The null

hypothesis follows Eq. 1, but instead of additive QTL effects, there is one effect for each

parent to model general combining ability (GCA). The GCA effects have i.i.d. normal

priors, with an inverse chi-squared prior for the variance ("BRR" model in BGLR).

Although GCA effects are not explicit in Eq. 1, they are implicitly present because the

average of the additive haplotype effects for each parent equals its GCA.

The test statistic (i.e., "score") for each marker bin during the genome scan is

$-\Delta\mathrm{DIC}$, which equals the DIC of the null hypothesis minus the DIC of the alternative

hypothesis. A typical convention for accepting a more complex model is that the DIC

should decrease by at least 5, preferably 10 (Lunn *et al.* 2012), but this is not adequate

to control the genome-wide Type I error rate during QTL discovery. To determine a

proper threshold, one can use diaQTL function *scan1_permute* to run a stratified

permutation test (Churchill and Doerge 1994), randomly permuting the phenotypes

within each F1 population. The largest $-\Delta$DIC for each permutation is recorded, and

the $1 - \alpha$ quantile of this distribution is used for QTL discovery with significance level α.

Alternatively, due to the generic nature of linkage mapping populations for a given

mating design, number of parents, ploidy, and genome size, simulation can be used to

determine the appropriate threshold. Half-diallel mating designs (without selfing) were

simulated using the software PedigreeSim V2.0 (Voorrips and Maliepaard 2012) and

auxiliary R package PedigreeSimR (https://www.github.com/rramadeu/PedigreeSimR).

The simulated genomes contained 1 to 12 linkage groups, each with a length of 100 cM,

loci evenly spaced at 0.1 cM, and recombination based on Haldane's map function. For

tetraploids, only bivalents were allowed with no preferential pairing of homologous

chromosomes. The number of parents was varied from 2 to 10. A genome-wide scan

with *scan1* was run for 1000 simulations, using standard normal deviates as

phenotypes, and the largest $-\Delta$DIC for each simulation was recorded. For each ploidy

(2x and 4x) and significance level ($\alpha = 0.01, 0.05, 0.1, 0.2$), a bivariate monotone

regression spline was fit for genome size and number of parents using R/scam (Pya

and Wood 2015). Predicted values from the spline are returned by diaQTL function

DIC_thresh.

Power Simulation

A factorial numerical experiment with $3 \times 3 \times 3 \times 2 = 54$ treatment scenarios, and

1000 simulations per scenario, was used to estimate statistical power and accuracy as

a function of the total population size (N = 200, 400, 800), number of parents (3, 4, 6)

with a half-diallel mating design, heritability (h^2 = 0.1, 0.2, 0.4), and ploidy (2, 4),

assuming a genome size of 12 Morgans (simulated as described above). The $-\Delta$DIC

threshold was chosen to maintain significance level $\alpha = 0.05$. Simulations were

performed using a high-performance computing facility at the University of Florida

(HiPerGator 2.0), allocating 2GB of RAM for each thread.

For each simulation, a single additive QTL was randomly assigned to one locus,

and all other loci were used as markers for mapping. Unreplicated phenotypes were

simulated by $y_i = \sum_{k=1}^{\phi p} W_{ik} \alpha_k + \varepsilon_i$ (same symbols as Eq. 1), using i.i.d. standard normal

deviates for the additive haplotype effects and i.i.d. normal residuals with variance

chosen to achieve the target heritability. If the realized heritability, computed from the

simulated QTL effects and residuals (Eq. 10), deviated from the prescribed value by

more than 0.01 (due to finite sample size), the simulation was discarded before QTL

mapping. Perfect knowledge about the haplotype dosage W_{ik} was assumed, which is

asymptotically achievable with PolyOrigin as marker density increases (Zheng *et al.*

2021). Genotyping error would reduce power and accuracy (Zhang *et al.* 2014; Bourke

et al. 2019).

Statistical power was the proportion of simulations in which both flanking markers of

the simulated QTL were declared significant. The accuracy of the inferred QTL position

was the distance between the simulated position and most significant marker. Accuracy

for the posterior mean estimates of the additive effects was evaluated based on the

Pearson correlation with simulated values. Monotone splines for power and accuracy

were fit using R/scam (Pya and Wood 2015).

To ensure the results were robust to polygenic effects, we simulated traits with an

additive QTL and polygenic effect for a half-diallel design with 4 parents and total

population size N = 200. The multivariate normal deviate was simulated using *mvrnorm*

from R/MASS (Venables and Ripley 2002), with covariance equal to the additive

relationship matrix (Eq. 7) times a polygenic variance component. Increasing the

proportion of variance due to the polygenic effect up to 0.3, while keeping the QTL h^2 at

0.2, had negligible effect on statistical power (Fig. S1).

Potato Data

The potato dataset is a half-diallel population (without selfing) for three parents from

the University of Wisconsin (UW) breeding program: Villetta Rose, W6511-1R, and

W9914-1R. The population was genotyped using version 3 of the potato SNP array, and

allele dosage was called using R package *fitPoly* (Voorrips *et al.* 2011; Zych *et al.*

2019). There were 5334 markers distributed across 12 chromosome groups, and their

physical positions were based on the potato DMv4.03 reference genome (Potato

Genome Sequencing Consortium 2011; Sharma *et al.* 2013). Parental genotype

probabilities and a genetic map were calculated using the software PolyOrigin (Zheng *et*

al. 2021), which identified 19 of the 434 progeny as outliers, and phenotype data were

unavailable for 2 more. The remaining 413 clones were distributed as follows: Villetta

Rose × W6511-1R (154 individuals), Villetta Rose × W9914-1R (113 individuals), and

W6511-1R × W9914-1R (146 individuals). The genetic map produced by PolyOrigin

spanned 12.1 Morgans and contained 2781 marker-bins.

The half-diallel population was evaluated as part of a larger field trial in 2018 at the

UW Hancock Agricultural Research Station. The trial used an augmented design with

four incomplete blocks and seven repeated checks per block. Each plot was a single

row sown with 15 seeds (tuber pieces) at 30 cm in-row spacing and 90 cm between

rows. The trial was harvested mechanically, and tubers were passed through an optical

sizer after washing to measure tuber length and width. Tuber shape was calculated as

the average length/width (L/W) ratio of all tubers weighing 170–285g. To improve the

normality of the residuals, the trait was transformed to $\log\left[(L/W) - 1\right]$. Initial statistical

analysis using ASReml-R v4 (Butler *et al.* 2018) was based on the linear model $y_{ij} =$

$\mu + block_j + clone_i + \varepsilon_{ij}$. Broad-sense heritability on a plot basis was estimated at 0.91

by treating $clone_i$ and $block_j$ as random effects (i.i.d. normal). BLUEs for each clone

were estimated by treating $clone_i$ as fixed and then used as the response variable for

analysis with diaQTL.

To give an idea about computational requirements, a variable of size 430 MB was

created by the diaQTL function *read_data* from the potato input CSV files. Based on the

output from *set_params*, 500 iterations were used with *scan1* for QTL discovery, which

required 3 minutes using 2 cores of a 3.1 GHz Intel Core i5 processor. Several functions

in the diaQTL package (*read_data*, *scan1*, *IBDmat*) can utilize multiple cores for parallel

execution.

Software and Data Availability

Version 1.00 of diaQTL and version 0.2 of PedigreeSimR, which were current at the

time of manuscript publication, are available as supplemental files through FigShare

under the GPL-3 license. For the most up-to-date version of diaQTL, download the

package from https://github.com/jendelman/diaQTL. The potato dataset is distributed

with the diaQTL package and used in the tutorial vignette.

RESULTS

Statistical power and accuracy

Simulated half-diallel populations were generated to study the influence of ploidy, number of parents, and genome size on the threshold needed to control the genome-wide Type I error rate at $\alpha = 0.05$. The test statistic is $-\Delta$DIC, which equals the Deviance Information Criterion for the null hypothesis (GCA but no QTL effects) relative to the alternative hypothesis of an additive QTL. The threshold was higher for tetraploids compared to diploids and increased approximately linearly with the number of parents (Fig. 2). There was also an approximately linear relationship between the threshold and logarithm of genome size (Fig. S2). As expected from previous research in diploids (Lander and Botstein 1989), the threshold was not influenced by population size. QTL discovery can also be conducted using markers as covariates or with digenic dominance, but higher $-\Delta$DIC thresholds are needed to maintain the same significance level (Fig. S3).

The influence of population size, number of parents, ploidy, and QTL heritability on statistical power was investigated for a genome of 12 Morgans (the size of the potato genome). Power increased with population size and QTL heritability but decreased with the number of parents and ploidy (Fig. S4). For a given h^2, the interplay between these factors could be largely summarized by the number of progeny per parental haplotype, abbreviated pph, but the diploid (tetraploid) results were consistently below (above) the regression spline (Fig. 3 and S5). At $h^2 = 0.1$ and $\alpha = 0.05$, power reached 0.5 and 0.9 with 30 and 70 pph, respectively, while only 10 and 25 pph (respectively) were needed for $h^2 = 0.2$. At $h^2 = 0.4$, the power was 1 for all scenarios (Fig. S4). As power increased,

so did the accuracy of the inferred QTL position, as measured by the distance between

the most significant marker and QTL (Fig. S6). The positional accuracy was 6 cM when

the power was 0.5 and decreased to 3 cM when power was 0.9.

The accuracy of the predicted haplotype effects, as measured by the correlation

with simulated values, was also largely explained by the pph metric (Fig. 4). For $h^2 = 0.1$

and $\alpha = 0.05$, 40 pph was sufficient to achieve an accuracy of 0.9, while for $h^2 = 0.2$,

only 20 pph was needed. Across all scenarios, the heritability of the QTL could be

estimated with essentially no error (Fig. S4), even when the accuracy of the haplotype

effects was only 0.7.

Potato diallel

To demonstrate diaQTL with a real dataset, we analyzed potato tuber shape,

measured by the length to width (L/W) ratio, in a half diallel population with three

parents. The ideal range of values for tuber shape is determined by the end-product

(e.g., long tubers for French fries, round tubers for potato chips) and to some extent

cultural preferences. The three parents were developed for the red-skin fresh market in

the US, for which a round to slightly oblong shape is expected. The distribution of L/W

values for the population ranged from 1.0 to 2.0 (Fig. S7), while the commercial checks

used in the field experiment had values between 1.1 and 1.2.

A genome scan with the additive model identified a large peak on chromosome 10

at 63 cM (Fig. 5A), designated 10@63. This region coincides with the location of the

classical *Ro* (round) QTL in potato (Van Eck *et al.* 1994), and the causal gene has been

identified as *StOVP20*, an *OVATE Family Protein* with orthologs that affect fruit shape in

tomato, melon, and cucumber (Wu *et al.* 2018). *StOVP20* is not present in the DM

reference genome, but based on synteny analysis (Wu *et al.* 2018), it is flanked by the

DM gene models (PGSC0003DMG400006678, PGSC0003DMG400006679). This

corresponds to the DM genome interval (48,982,521 - 49,022,687), which lies within the

95% CI for the 10@63 QTL (48,203,284 - 50,782,097).

The next highest peak in the genome scan was on chromosome 1 at 133 cM. The

$-\Delta$DIC score was at the threshold for $\alpha = 0.05$ but clearly above the threshold for $\alpha = $

0.1 (Fig. 5A). The model with both QTL reduced the DIC by 23 points compared to the

model with only 10@63, so it was accepted over the single QTL model (Table 1).

More complex models involving dominance and/or additive x additive epistatic

effects were evaluated next (Table 1). Epistasis lowered the DIC by 6 points compared

to the two-QTL additive model, whereas digenic dominance at 10@63 lowered the DIC

by 17 points. Including both of these non-additive effects raised the DIC, as did higher

order dominance at 10@63 and dominance at 1@133 (Table 1). We therefore selected

the model with additive and digenic effects at 10@63 and additive effects at 1@133,

which accounted for 26%, 8%, and 5% of the total variance, respectively (Table 2). To

quantify the importance of smaller undetected QTL, an additive polygenic effect was

included, which lowered the DIC by 78 points and accounted for 29% of the total

variance (Table 2).

Of the 12 parental haplotypes at 10@63, the largest additive effect (in magnitude)

was estimated for haplotype 2 of parent W6511-1R (Fig. 5B), and the negative sign

means it reduced the L/W ratio. Haplotypes 2 and 4 of parent Villetta Rose also reduced

the L/W ratio. For parent W9914-1R, all four haplotypes had similar additive effects but

showed some differentiation with respect to non-additive effects (Fig. S8 and S9).

DISCUSSION

Although this is the first publication on statistical power and accuracy for QTL

mapping in tetraploid diallel populations, some comparisons with previous studies on

diploid diallel or biparental tetraploid populations are possible. Among previous

simulation studies, Muranty (1996) is unique for explicitly showing the loss of statistical

power as the number of outbred diploid parents increased at a fixed total population

size. We observed the same trend (Fig. 3), but the exact values for power are not

comparable because of differences in the significance level (higher values produce

higher power). Muranty (1996) also reported that the specific form of the mating design

(e.g., cyclic, factorial, half-diallel) had little impact on power, which was confirmed in our

simulation (data not shown). Bourke *et al.* (2019) reported statistical power in a

biparental tetraploid population of size 200, which corresponds to 25 pph, was 0.8 for a

bi-allelic, additive QTL with $h^2 = 0.1$ and high genotypic information content. This is

higher than the value of 0.4 for the regression spline in Fig. 3, which may be the result

of the simpler genetic architecture (bi-allelic vs. multi-allelic in our simulation).

Compared to existing software, R/diaQTL has unique features for modeling

dominance and epistasis with multi-allelic QTL in tetraploids. The inner product of the

genetic dosages used for QTL mapping of non-additive effects, when averaged across

many loci, generates non-additive relationships (see Methods). Due to Mendelian

segregation, these realized relationships exhibit a distribution around the expected

value based on pedigree (Fig. S10; Amadeu *et al.* 2020). Non-additive relationship

matrices probably have limited utility for QTL mapping, but we envision potential

applications with genomic selection. Breeding values include 1/2 of the additive x

additive epistasis and 1/3 of the digenic dominance in tetraploids (Gallais 2003),

although this result is only exact when genotype frequencies equal the product of

haplotype frequencies (Kempthorne 1957), which requires a complete diallel with selfs.

Our statistical model allows each parental haplotype to have a different effect, but

we expect some haplotypes will carry the same QTL allele. Although the use of random

effects for the regression coefficients helps offset the statistical complexity of the

haplotype model, from a genetic point of view, we are interested in inferring identity-by-

state relationships. At present, a user of our software is limited to hypothesizing that

haplotypes with similar effect carry the same allele. One direction for future research

would be to incorporate methods to formally test such hypotheses (Jannink and Wu

2003; Crouse *et al.* 2020).

Large plant breeding programs often maintain distinct research enterprises devoted

to genetic discovery vs. variety selection. Genetic discovery projects typically

characterize novel populations, such as a panel of distantly related germplasm or

larger-than-normal, unselected biparental populations, but this additional effort is not

always feasible for small programs. There is also the potential disconnect between the

development of a genetic marker in the discovery population and its successful

deployment in the breeding populations; such markers often fail because they are not

haplotype-specific. With a diallel QTL analysis of the breeding population, this additional

step would be unnecessary. The diaQTL function *haplo_plot* allows users to visualize

which parental haplotypes are present at a particular locus (Fig. S11), and it is

straightforward to select progeny-based on the dosage of particular haplotypes using

diaQTL function *haplo_get* (see package tutorial).

The University of Wisconsin potato breeding program provides an example to

discuss the feasibility of using breeding populations for discovery research. A typical

mating scheme (for each market type) involves 10 females and four males (male fertility

is scarce), from which 40 F1 populations with 500 progeny each are created and

evaluated as single plants in the field. Using visual selection for plant maturity and tuber

type, 800 individuals (= 4%) advance to the first clonal stage, for genotyping and further

phenotyping. Under uniform selection, this population size corresponds to 20 pph for the

female parents (more for the male parents), which translates into 80% power at $h^2 = 0.2$

based on our simulation results.

CONCLUSION

The R/diaQTL software is a user-friendly tool for QTL mapping and haplotype

analysis in connected F1 populations derived from outbred parents. The number of

progeny (or gametes, if selfed populations are used) per parental haplotype is a critical

design parameter to consider when developing populations. By enabling selection on

the effect and dosage of haplotypes without having to develop haplotype-specific

markers, diaQTL can improve the efficiency of breeding.

Acknowledgments: Financial support provided by USDA NIFA Award No. 2019-67013-

29166. We thank Maria Caraza-Harter for testing the diaQTL software.

Author contributions: RRA and JBE developed the software, analyzed the data, and

drafted the manuscript. All authors contributed to the methodology and edited the

manuscript.

LITERATURE CITED

Amadeu, R. R., L. A. C. Lara, P. Munoz, and A. A. F. Garcia, 2020 Estimation of molecular pairwise relatedness in autopolyploid crops. G3: Genes, Genomes, Genetics 10: 4579-4589.

Bink, M. C., J. Jansen, M. Madduri, R. E. Voorrips, C. E. Durel, *et al.*, 2014 Bayesian QTL analyses using pedigreed families of an outcrossing species, with application to fruit firmness in apple. Theoretical and Applied Genetics 127: 1073{1090.

Bourke, P. M., C. A. Hackett, R. E. Voorrips, R. G. F. Visser, and C. Maliepaard, 2019 Quantifying the power and precision of QTL analysis in autopolyploids under bivalent and multivalent genetic models. G3: Genes, Genomes, Genetics 9: 2107-2122.

Bourke, P. M., G. van Geest, R. E. Voorrips, J. Jansen, T. Kranenburg, *et al.*, 2018 polymapR: linkage analysis and genetic map construction from F1 populations of outcrossing polyploids. Bioinformatics 34:3496-3502.

Broman, K. W. and S. Sen, 2009 A Guide to QTL Mapping with R/qtl, volume 46. Springer.

Buckler, E. S., J. B. Holland, P. J. Bradbury, C. B. Acharya, P. J. Brown, *et al.*, 2009 The genetic architecture of maize flowering time. Science 325: 714-718.

Butler, D., B. Cullis, A. Gilmour, B. Gogel, and R. Thompson, 2018 ASReml-R Reference Manual Version 4. VSN International Ltd, Hemel Hempstead, HP1 1ES, UK.

Churchill, G. A. and R. W. Doerge, 1994 Empirical threshold values for quantitative trait mapping. Genetics 138: 963-971.

Crouse, W. L., S. N. P. Kelada, and W. Valdar, 2020, Inferring the allelic series at QTL in multiparental populations. Genetics 216: 957-983.

da Silva Pereira, G., D. C. Gemenet, M. Mollinari, B. A. Olukolu, J. C. Wood, *et al.*, 2020 Multiple QTL mapping in autopolyploids: A random-effect model approach with application in a hexaploid sweetpotato fullsib population. Genetics 215: 579-595.

Endelman, J. B. and J.-L. Jannink, 2012 Shrinkage estimation of the realized relationship matrix. G3: Genes, Genomes, Genetics 2: 1405-1413.

Gallais, A., 2003 Quantitative genetics and breeding methods in autopolyploids plants. INRA, Paris, First edition.

Grattapaglia, D. and R. Sederoff, 1994 Genetic linkage maps of Eucalyptus grandis and Eucalyptus urophylla using a pseudo-testcross: mapping strategy and RAPD markers. Genetics 137: 1121-1137.

Habier, D., R. L. Fernando, K. Kizilkaya, and D. J. Garrick, 2011 Extension of the Bayesian alphabet for genomic selection. BMC Bioinformatics 12: 186.

Hackett, C. A., B. Boskamp, A. Vogogias, K. F. Preedy, and I. Milne, 2017 TetraploidSNPMap: software for linkage analysis and QTL mapping in autotetraploid populations using SNP dosage data. Journal of Heredity 108: 438-442.

Hackett, C. A., J. E. Bradshaw, and G. J. Bryan, 2014 QTL mapping in autotetraploids using SNP dosage information. Theoretical and Applied Genetics 127:1885-1904.

Huang, B. E., K. L. Verbyla, A. P. Verbyla, C. Raghavan, V. K. Singh, *et al.*, 2015 MAGIC populations in crops: current status and future prospects. Theoretical and Applied Genetics 128: 999-1017.

Jannink, J.-L. and X.-L. Wu, 2003 Estimating allelic number and identity in state of QTLs in interconnected families. Genetical Research 81: 133{144.

Kempthorne, O., 1957 One-Locus Polyploid Population. *In* An introduction to genetic statistics, chapter 18, pp. 400-409, John Wiley & Sons, New York.

Knapp, S., W. Bridges, and D. Birkes, 1990 Mapping quantitative trait loci using molecular marker linkage maps. Theoretical and applied genetics 79: 583-592.

Lander, E. S. and D. Botstein, 1989 Mapping Mendelian factors underlying quantitative traits using RFLP linkage maps. Genetics 121: 185-199.

Lenarcic, A. B., K. L. Svenson, G. A. Churchill, and W. Valdar, 2012 A general Bayesian approach to analyzing diallel crosses of inbred strains. Genetics 190: 413-435.

Liu, Y. and Z.-B. Zeng, 2000 A general mixture model approach for mapping quantitative trait loci from diverse cross designs involving multiple inbred lines. Genetics Research 75: 345-355.

Lunn, D., C. Jackson, N. Best, A. Thomas, and D. Spiegelhalter 2012 The BUGS Book: A practical introduction to Bayesian analysis, CRC Press, Boca Raton.

Margarido, G. R., A. P. Souza, and A. A. Garcia, 2007 OneMap: software for genetic mapping in outcrossing species. Hereditas 144: 78-79.

Matias, F. I., K. G. Xavier Meireles, S. T. Nagamatsu, S. C. Lima Barrios, C. Borges do Valle, *et al.*, 2019 Expected genotype quality and diploidized marker data from genotyping-by-sequencing of Urochloa spp. tetraploids. The Plant Genome 12: 190002.

Mollinari, M. and A. A. F. Garcia, 2019 Linkage analysis and haplotype phasing in experimental autopolyploid populations with high ploidy level using hidden Markov models. G3: Genes, Genomes, Genetics 9: 3297-3314.

Muranty, H., 1996 Power of tests for quantitative trait loci detection using fullsib families in different schemes. Heredity 76: 156-165.

Nice, L. M., B. J. Steffenson, T. K. Blake, R. D. Horsley, K. P. Smith, *et al.*, 2017 Mapping agronomic traits in a wild barley advanced backcross-nested association mapping population. Crop Science 57: 1199-1210.

Pérez, P. and G. de Los Campos, 2014 Genome-wide regression and prediction with the BGLR statistical package. Genetics 198: 483-495.

Potato Genome Sequencing Consortium, 2011 Genome sequence and analysis of the tuber crop potato. Nature 475: 189-195.

Plummer M., N. Best, K. Cowles, and K. Vines, 2006 CODA: convergence diagnosis and output analysis for MCMC. R News 6: 7–11.

Pya, N. and S. N. Wood, 2015 Shape constrained additive models. Statistics and Computing 25: 543-559.

R Core Team, 2020 R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Raftery, A. E. and S. M. Lewis, 1992 One Long Run with Diagnostics: Implementation Strategies for Markov chain Monte Carlo. Statistical Science 7: 493-497.

Rebai, A. and B. Goffinet, 1993 Power of tests for QTL detection using replicated progenies derived from a diallel cross. Theoretical and Applied Genetics 86: 1014-1022.

Ritter, E. and F. Salamini, 1996 The calculation of recombination frequencies in crosses of allogamous plant species with applications to linkage mapping. Genetics Research 67: 55-65.

Rosyara, U. R., M. C. Bink, E. van de Weg, G. Zhang, D. Wang, *et al.*, 2013 Fruit size QTL identification and the prediction of parental QTL genotypes and breeding values in multiple pedigreed populations of sweet cherry. Molecular Breeding 32: 875-887.

Sharma, S. K., D. Bolser, J. de Boer, M. Sønderkær, W. Amoros, *et al.*, 2013 Construction of reference chromosome-scale pseudomolecules for potato: integrating the potato genome with genetic and physical maps. G3: Genes, Genomes, Genetics 3: 2031-2047.

Song, Q., L. Yan, C. Quigley, B. D. Jordan, E. Fickus, *et al.*, 2017 Genetic characterization of the soybean nested association mapping population. The Plant Genome 10: 1-14.

Spiegelhalter, D. J., N. G. Best, B. P. Carlin, and A. Van Der Linde, 2002 Bayesian measures of model complexity and fit. Journal of the royal statistical society: Series b (statistical methodology) 64: 583-639.

Uitdewilligen, J. G. A. M. L., A.-M. A. Wolters, B. B. D'hoop, T. J. A. Borm, R. G. F. Visser, *et al.*, 2013 A next-generation sequencing method for genotyping-by-sequencing of highly heterozygous autotetraploid potato. PLoS ONE 8: e62355.

Van Eck, H. J., J. M. Jacobs, P. Stam, J. Ton, W. J. Stiekema, *et al.*, 1994 Multiple alleles for tuber shape in diploid potato detected by qualitative and quantitative genetic analysis using RFLPs. Genetics 137: 303-309.

Van Ooijen, J., 2004 MapQTL® 5, software for the mapping of quantitative trait loci in experimental populations of diploid species. Kyazma B.V., Wageningen, Netherlands.

Van Ooijen, J. and R. Voorrips, 2001 Joinmap® 3, software for the calculation of genetic linkage maps. Kyazma B.V., Wageningen, Netherlands.

Verhoeven, K. J., J. L. Jannink, and L. M. McIntyre, 2006 Using mating designs to uncover QTL and the genetic architecture of complex traits. Heredity 96: 139-149.

Venables, W. N. and B. D. Ripley, 2002 Modern applied statistics with S. Fourth edition. Springer, New York.

Voorrips, R. E., G. Gort, and B. Vosman, 2011 Genotype calling in tetraploid species from bi-allelic marker data using mixture models. BMC Bioinformatics 12: 172.

Voorrips, R. E. and C. A. Maliepaard, 2012 The simulation of meiosis in diploid and tetraploid organisms using various genetic models. BMC Bioinformatics 13: 248.

Wu, K., W. Burnquist, M. Sorrells, T. Tew, P. Moore, *et al.*, 1992 The detection and
estimation of linkage in polyploids using single-dose restriction fragments. Theoretical
and Applied Genetics 83: 294-300.

Wu, R., C.-X. Ma, I. Painter, and Z.-B. Zeng, 2002 Simultaneous maximum likelihood
estimation of linkage and linkage phases in outcrossing species. Theoretical population
biology 61: 349-363.

Wu, S., B. Zhang, N. Keyhaninejad, G. Rodríguez, H. Kim, *et al.*, 2018 A common
genetic mechanism underlies morphological diversity in fruits and other plant organs.
Nature communications 9: 1-12.

Yang, J., N. A. Zaitlen, M. E. Goddard, P. M. Visscher, and A. L. Price, 2014
Advantages and pitfalls in the application of mixed-model association methods. Nature
genetics 46: 100-106.

Young, N. and S. Tanksley, 1989 Restriction fragment length polymorphism maps and
the concept of graphical genotypes. Theoretical and Applied Genetics 77: 95-101.

Yu, J., J. B. Holland, M. D. McMullen, and E. S. Buckler, 2008 Genetic design and
statistical power of nested association mapping in maize. Genetics 178: 539-551.

Zhang, Z., W.Wang, and W. Valdar, 2014 Bayesian modeling of haplotype effects in
multiparent populations. Genetics 198: 139-156.

Zheng, C., R. Amadeu, P. Munoz, and J. Endelman, 2021 Haplotype reconstruction in
connected tetraploid F1 populations. Genetics (accepted) doi:10.1101/2020.12.18.423519

Zheng, C., M. P. Boer, and F. A. van Eeuwijk, 2015 Reconstruction of genome ancestry
blocks in multiparental populations. Genetics 200: 1073-1087.

Zych, K., G. Gort, C. A. Maliepaard, R. C. Jansen, and R. E. Voorrips, 2019 FitTetra
2.0: improved genotype calling for tetraploids with multiple population and parental data
support. BMC Bioinformatics 20: 148.

 FIGURE CAPTIONS

Figure 1. Examples of diallel populations with four parents (labeled a,b,c,d); each square

represents a family of full-sibs. (A) full diallel; (B) half diallel with selfing; (C) half diallel without

selfing; (D) circular; (E) linear; (F) factorial; (G) testcross or nested design; (H) arbitrary partial

diallel. Figure adapted from Verhoeven *et al.* (2006).

Figure 2. Threshold to control the genome-wide Type 1 error rate at $\alpha = 0.05$, for a half-diallel

design without selfing. The statistic $-\Delta$DIC is the Deviance Information Criterion for the null

hypothesis (no QTL) relative to the alternative hypothesis of an additive QTL. Each point is

based on 1,000 simulations, and the best-fit linear regression is shown as a solid line.

Figure 3. Statistical power to detect a multi-allelic QTL at significance level $\alpha = 0.05$, as a

function of the number of progeny per parental haplotype, number of parents, ploidy, and QTL

heritability (h^2). Each point is the average of 1,000 simulations with an additive model. The

dashed line is a monotone increasing, concave spline for $h^2 = 0.1$, and the solid line is the spline

for $h^2 = 0.2$.

Figure 4. Accuracy of the estimated haplotype effects, as measured by Pearson's correlation

with the simulated values. Each point is the average of 1,000 simulations with an additive

model. The dashed line is a monotone increasing, concave spline for $h^2 = 0.1$, and the solid line

is the spline for $h^2 = 0.2$.

Figure 5. R/diaQTL results for potato tuber shape using a half-diallel population with three

parents: VillettaRose, W6511-1R, and W9914-1R. (A) Genome scan with *scan1*. The dashed

horizontal lines correspond to $\alpha = 0.05$ (gold) and $\alpha = 0.1$ (red). The most significant marker was

solcap_snp_c2_25522 on potato chromosome 10. (B) Additive effect estimates for the 12

parental haplotypes using *fitQTL*. Error bars are the 90% CI.

TABLES

Table 1. Model comparison for potato tuber shape based on ΔDIC, which is the

Deviance Information Criterion relative to a null model with parental GCA effects. QTL

are designated as chromosome@cM, with A = additive, D = digenic, T = trigenic, and

AA = additive x additive epistasis.

Model	Effects	ΔDIC
1	10@63(A)	-160
2	10@63(A) + 1@133(A)	-183
3	10@63(A) + 1@133(A) + 10@63 x 1@133 (AA)	-189
4	10@63(A+D) + 1@133(A)	**-200**
5	10@63(A+D) + 1@133(A) + 10@63 x 1@133 (AA)	-196
6	10@63(A+D+T) + 1@133(A)	-197
7	10@63(A+D) + 1@133(A+D)	-196

657 **Table 2**. Proportion of variance for potato tuber shape based on Model 4 from Table 1,

658 plus a polygenic effect. QTL are designated as chromosome@cM.

Effect	Proportion of Variance
10@63 additive	0.26
10@63 digenic dominance	0.08
1@133 additive	0.05
polygenic	0.29
residual	0.31

Supplemental Figures

QTL Mapping in Outbred Tetraploid (and Diploid) Diallel Populations

Rodrigo R. Amadeu*, Patricio R. Munoz*, Chaozhi Zheng[†], Jeffrey B. Endelman[‡]

*Horticultural Sciences Department, University of Florida, Gainesville, FL, USA

[†]Biometris, Wageningen University and Research, Wageningen, The Netherlands

[‡]Department of Horticulture, University of Wisconsin, Madison, WI, USA

Figure S1. The influence of simulating polygenic effects on statistical power for detecting an additive QTL with h^2 = 0.2 in a half-diallel with 4 parents and total population size N = 200. The height of each bar is the proportion of 1000 simulations in which the QTL was detected, and the error bars are $\pm 1.96\,\text{SE}$ for the standard error (SE) of a binomial proportion.

Figure S2. The threshold for QTL discovery, −ΔDIC, increased approximately linearly with the logarithm of genome size. Results are shown for significance level $\alpha = 0.05$, using simulated half-diallel populations, and the number of parents (4 and 10) is shown at the top of each panel. Each point is based on 1000 simulations.

Figure S3. The threshold for QTL discovery, −ΔDIC, was higher for a QTL model with additive and dominance effects (A+D) compared to only additive (A) effects. Results are shown for significance level $\alpha = 0.05$, using 1000 simulated half-diallel populations. The number of parental haplotypes equals ploidy (2 or 4) times the number parents (2 to 10).

Figure S4. QTL mapping results from simulated half-diallel populations with varying population size, QTL heritability, number of parents, and ploidy. The genome-wide Type I error rate was controlled at $\alpha = 0.05$. A) Average distance in cM between the peak LOD score position and simulated QTL position. B) Average power to detect the QTL. C) Average estimated QTL heritability based on the diaQTL linear model. Error bars are standard errors based on 1,000 simulations.

Figure S5. Statistical power to detect QTL for different significance levels α, as a function of the number of progeny per parental haplotype, number of parents, ploidy, and QTL heritability (h^2). Each point is the average of 1,000 simulations with an additive model. The dashed line is a monotone increasing, concave spline for h^2 = 0.1, and the solid line is the spline for h^2 = 0.2.

Figure S6. Average distance between the position of the most significant marker and simulated QTL position, as function of power, ploidy and number of parents. Each point represents an average over the multiple population sizes and heritability values shown in Figure S4.

Figure S7. Distribution of the length to width (L/W) ratio for tubers of three full-sib families of autotetraploid potato. The images illustrate the extreme phenotypes of this population with L/W = 1.0 as a round tuber shape, L/W = 2.0 as an elongated tuber shape.

Trait: tuber_shape
Marker: solcap_snp_c2_25522

Figure S8. Estimated QTL effects for the 10@63 QTL with the digenic dominance model (#4 in Table 1). The digenic dominance effects are above the diagonal, and below the diagonal are the sum of the additive and digenic effects. The four haplotypes for each parent are labeled .1 to .4. The figure is part of the graphical output from diaQTL function *fitQTL*.

Figure S9. Estimated additive x additive epistatic effects between the 10@63 and 1@133 QTL (model #3 in Table 1). The four haplotypes for each parent are labeled .1 to .4. The figure is part of the graphical output from diaQTL function *fitQTL*

Figure S10. The distribution of realized relationships between full-sibs (FS) and half-sibs (HS) in the potato diallel population, computed with diaQTL function *IBDmat*. The four panels correspond to additive (A), digenic dominant (D), trigenic dominant (T), and additive x additive epistatic (E) effects. The expected values based on pedigree are A = 1/2, D = 2/9, T = 1/12, E = 1/4 for full-sibs and A = 1/4, D = 1/36, T = 0, E = 1/16 for half-sibs.

Figure S11. Visualizing the inheritance of parental haplotypes from VillettaRose and W6511-1R in one F1 offspring (W15268-53R), using diaQTL function *haplo_plot*. This clone was selected as one of three offspring containing two copies of the W6511-1R.2 haplotype, which was estimated to have the largest negative effect (Fig. 5) and is therefore desirable for breeding potato varieties with round tubers. The dashed line marks the position of the 10@63 QTL for tuber shape. The presence of two copies of a parental haplotype is possible in tetraploid offspring due to double reduction, a phenomenon in which the diploid gamete contains portions of both sister chromatids from one homologous chromosome.