Title:

Sensitive detection of DNA contamination in tumor samples via microhaplotypes

Short Title:

Detection of DNA contamination in tumor samples via microhaplotypes

Authors:

Brett Whitty and John F. Thompson[1*]

Research and Development, PGDx, 2809 Boston Street, Baltimore, MD 21224

[1]Current address: Homology Medicines Inc., 1 Patriots Park, Bedford, MA 01730

[*]Corresponding Author: Thompson.john.f@gmail.com

Abstract

Background: Low levels of sample contamination can have disastrous effects on the accurate

identification of somatic variation in tumor samples. Detection of sample contamination in DNA is

generally based on observation of low frequency variants that suggest more than a single source of DNA

is present. This strategy works with standard DNA samples but is especially problematic in solid tumor

FFPE samples because there can be huge variations in allele frequency (AF) due to massive copy number

changes arising from large gains and losses across the genome. The tremendously variable allele

frequencies make detection of contamination challenging. A method not based on individual AF is

needed for accurate determination of whether a sample is contaminated and to what degree.

Methods: We used microhaplotypes to determine whether sample contamination is present.

Microhaplotypes are sets of variants on the same sequencing read that can be unambiguously phased.

Instead of measuring AF, the number and frequency of microhaplotypes is determined. Contamination

detection becomes based on fundamental genomic properties, linkage disequilibrium (LD) and the

diploid nature of human DNA, rather than variant frequencies. We optimized microhaplotype content

based on 164 single nucleotide variant sets located in genes already sequenced within a cancer panel.

Thus, contamination detection uses existing sequence data and does not require sequencing of any

extraneous regions. The content is chosen based on LD data from the 1000 Genomes Project to be

ancestry agnostic, providing the same sensitivity for contamination detection with samples from

individuals of African, East Asian, and European ancestry.

Results: Detection of contamination at 1% and below is possible using this design. The methods

described here can also be extended to other DNA mixtures such as forensic and non-invasive prenatal

testing samples where DNA mixes of 1% or less can be similarly detected.

Conclusions: The microhaplotype method allows sensitive detection of DNA contamination in FFPE

tumor samples. These methods provide a foundation for examining DNA mixtures in a variety of

contexts. With the appropriate panels and high sequencing depth, low levels of secondary DNA can be

detected and this can be valuable in a variety of applications.

Background

In many genetics applications, contamination is not a significant issue because small amounts of

extraneous DNA do not affect the experimental outcome. For example, variant frequency in germ-line

DNA is approximately 0%, 50% or 100% non-reference so it requires a substantial fraction of DNA from

other sources to affect variant calls. Contamination of up to 5% has been shown to have virtually no

impact on Mendelian variant calling (1). However, even with low-level DNA contamination, there can be

a significant impact on some applications like detecting low frequency somatic variants in tumor DNA

samples or deconvoluting complex mixtures in forensic samples. Clinical diagnostic samples are

presumed to be pure but there is always a concern that unexpected contamination could affect results

and escape standard detection methods. Knowledge of whether a sample is contaminated with another

human sample and, if so, the extent of that low-level contamination, is critical for assessing whether

variant calls can be trusted.

Variant detection is often carried out by measuring the variable lengths of short tandem repeats (STRs)

or by genotyping/sequencing single-nucleotide variants (SNVs). When unusual frequencies are observed,

the degree of contamination is estimated by quantitating the frequency of the extra variants (2-4).

However, accurate estimation becomes difficult at low variant frequency because the true signal can be

obscured by the competing noise arising from technical issues like STR stutter or genotyping/sequencing

errors. This is especially problematic with analog measurement systems that are often used with STR,

genotyping, and classical sequencing methods. In contrast, next-generation sequencing (NGS) methods

produce a digital signal with both STRs and SNVs, allowing greater sensitivity at low contamination

levels.

There are a number of other factors that can further complicate the analysis of DNA contamination.

With clinical samples, the contaminating DNA can be of different quality than the primary DNA sample.

This can add to the noise in the system because the DNAs may respond differently depending on the

assay. When tumor samples are being studied, there is the added complication of varying copy numbers

as large regions of the tumor genome may be highly amplified or deleted and samples typically contain

variable amounts of normal and tumor tissue. When large fractions of the genome have altered copy

number, the allelic frequencies (AFs) of SNVs can vary substantially and change from the typical 50-50

AF to much more extreme values. These altered allele ratios can overlap the range of AFs for what is

typically seen with contamination and thus a sample that is 100% pure can mistakenly appear to be

significantly contaminated. Thus, pure samples can be incorrectly reported as contaminated, resulting in

a false positive (FP) signal with perfectly suitable samples being discarded. In order to avoid these

problems, a method that is independent of genotype AF must be used for determining contamination

status with tumor DNA. Analysis based on microhaplotype (MH) counting can be used to achieve this

goal.

For individuals with a normal set of chromosomes, each autosomal region of any size has two sets of

alleles. Those alleles may be the same (homozygous) or different (heterozygous); but, under normal

circumstances with a few specific exceptions, there should never be more than two haplotypes in a

genomic region in any individual. More than two haplotype versions in a given region indicates

contamination. Each haplotype appears with a characteristic frequency in different populations and

there may be several MHs at the same genomic positions in any population. When DNA from two

individuals is mixed, accidentally or intentionally, there is a possibility for 1-4 MHs in each region.

Microhaplotypes can be defined as multiple variants on the same sequence read such that the phasing

of the variants is known and the haplotype can be unambiguously determined. MHs have been used for

forensic purposes in order to detect contamination and identify individuals (5-8), to detect transplant

DNA in a host background (9), and to determine ancestry (10-13). Use of MHs for routine detection and

estimation of contamination in clinical samples has not been described but is a powerful application of

these methods.

Methods

Sequencing Samples

De-identified DNA and tumor samples were obtained from several commercial and academic providers.

These included BioIVT, Conversant Bio, Cureline, Duke, Folio Bio, Fundacio Institute, Indivumed,

iSpecimen, NCI, and Proteogenex. Samples were received either as FFPE material or extracted DNA.

Extraction of DNA from FFPE specimens was carried out using standard techniques. DNA was prepared

for sequencing using hybridization capture with panels directed primarily at exons in >500 genes

implicated in cancer. Depending on the gene, some intronic and promoter DNA was also targeted. DNA

was sequenced using an Illumina NextSeq500. The PGDx elio™ tissue complete assay was used for

sample preparation and analysis (14, 15).

Variant calling

Low frequency variants were called using a machine learning approach (16) with replicate reads

compressed to a single sequence. Even with this approach, technology limitations can induce false

positives at frequencies of 1% or less. Because the false positive variant calls could artificially inflate the

number of 3rd/4th MHs, MH calculations included 3rd MHs only if they surpassed a minimum fraction of

variant calls (typically >0.2%). Each read covering the positions of all component SNVs for the MH set

was tagged and counted.

***In silico* read mixing**

Artificially contaminated samples were made by *in silico* mixing at multiple artificial contamination

levels. Contamination levels of x% were generated by combining randomly selected 100-x% reads from

the sample with x% reads from the contaminant and running the resultant set of reads through the

standard analysis pipeline. It was found that the same results were achieved if only the reads mapping

to the MH regions being examined were included in the mixing. These analyses ran on the pipeline much

faster. Thus, later experiments used only the truncated data set to speed analysis and allow more

conditions to be tested.

Results

Choice of initial variant sets

There are many MHs that have been characterized in the literature, and their usefulness for

distinguishing individuals and ancestries is well documented (7. 11, 12). However, most of the well-

characterized MHs are not in regions chosen for commercial panel sequencing. In addition, the criteria

for selecting MHs for discriminating individuals and ancestry are not the same as for detecting

contamination where the goal is not distinguishing individuals but maximizing the number of MHs with

>2 versions. Since high depth of sequencing is required for sensitive contamination detection, regions

already targeted for analysis were examined so that additional, non-productive sequencing in known

MH regions would not be necessary. For gene panel sequencing, this means that literature MH data is of

limited value. Each novel MH set we chose thus requires testing for behavior across samples and

ancestries. We have used a >500 gene cancer panel directed against FFPE tumor DNA. The panel

included regions outside of the target exons and these regions were also inspected for potentially

informative SNV sets.

The criteria for inclusion in the final set of variants included maximizing the frequency of 3rd MHs while

minimizing the likelihood of sequencing errors in those variants. This was achieved by avoiding regions

with higher inherent sequencing error rates such as insertions, deletions, and variants within long

homopolymeric regions and other repetitive elements. The regions targeted for panel sequencing were

examined in the gnomAD database (gnomad.broadinstitute.org/; 17) for multiple SNVs within 180 bp of

each other and with 5-95% AF. Variants not in a segmental duplication or a low confidence region were

examined for LD in the 1000 Genomes Project samples (ldlink.nci.nih.gov/?tab=ldhap; 18). SNV sets in

complete or near complete LD were discarded. Pairs and triplets with at least three haplotypes and the

3rd and greater haplotypes having a total frequency of >1% were selected for evaluation in real samples.

Variant sets with median coverage >100x for the relevant variants on the read were used to generate a

phased haplotype. Variant sets prone to higher frequency 3rd/4th microhaplotypes in purportedly pure

samples were eliminated because they would be likely to produce high noise relative to signal. When

nearby SNV sets provided duplicate information because of high LD between SNV sets, those with the

highest coverage and fewest 3rd/4th microhaplotypes in pure samples were included in the panel and the

alternatives omitted to minimize over-counting particular genomic regions. The 106 SNV sets that

passed these initial evaluations are shown in Supplemental Table 1. In addition to the genomic

coordinates and dbSNP identifiers, the frequency of MHs is shown for the five major ancestry groupings

defined by 1000 Genomes as well as for the African American sub-grouping. In addition, the median

coverage for the relevant SNVs across >700 samples, the number of times the SNV set generated more

than 2 MHs in "pure" samples, and the DNA length containing the SNVs are also listed.

Assessing and estimating contamination levels

In order to determine the criteria with which to detect contamination, *in silico* mixing of reads from

individual pure samples was carried out so that the precise level of contamination would be known and

the sensitivity of the MH assay could be determined. The initial samples used for establishing

contamination methods need to be as close to 100% pure as possible for maximizing sensitivity. After

potentially artifactual MH sets were eliminated as described above, samples were examined using the

initial MH sets to ensure that they were as pure as possible by including only those samples with the

fewest 3rd/4th MH sites.

The gnomAD database was used as a starting point for selecting SNVs and, at the time it was initially

queried, had a higher proportion of individuals of European ancestry (~60%) than individuals with other

geographic ancestries. Because of this, variation in European ancestry individuals had a larger impact on

the initial choice of SNVs than variation in other ancestries. Thus, there is a possibility that samples from

different ancestral backgrounds could respond differently when examined independently using SNV sets

selected from gnomAD and not filtered based on ancestry. The initial testing of MH sets for establishing

contamination criteria was carried out using three individuals of African American ancestry, three

individuals of East Asian ancestry, and three individuals of European ancestry. These ancestries were

self-declared by the individuals and confirmed using Principle Component Analysis (PCA) of the MH sets

based on 1000 Genome MH frequencies. The self-declared and PCA-determined ancestries of these

samples were in agreement. However, with other samples, the self-declared Asian ancestries were

sometimes South Asian rather than East Asian. Only individuals with PCA-confirmed East Asian ancestry

were used in this and subsequent analyses. In addition to diverse ancestral histories, samples were

selected for mixing experiments based on having two or fewer SNV sets with >2 MHs.

Precise levels of read mixing can be generated *in silico* but the functional level of contamination was

found to often differ significantly from the known mixing proportions. This issue was particularly obvious

when samples with divergent DNA qualities were mixed. The longer, higher quality DNA appeared to

have a higher concentration than what had been added to the mix. This arises because longer reads map

better and are more likely to contain all the variants being examined and thus this DNA appears to be

relatively more frequent than the shorter, degraded DNA. This effect is seen with all types of DNA but is

especially noticeable with FFPE DNA where DNA quality is highly dependent on the methods used to fix

and store the sample. This is consistent with previous results that showed DNA with reduced extraction

yields tend to be lower quality and more susceptible to contamination (19). To minimize the impact of

this effect, all DNA mixes were examined in a pairwise fashion where one sample was the primary

sample and the second was the contaminant. These roles were then switched and the results averaged.

This allowed an over-contamination with one sample set to be partially mitigated by the paired under-

contamination of the opposite sample set.

As shown in Fig 1, the maximum number of MHs with 3rd haplotypes ranged from 15-30% with this set of

MHs. Most mixes had 20-25% 3rd haplotypes with significant overlap of individuals of European and

African American ancestry while those with East Asian ancestry had fewer 3rd MHs, in the range of 15-

20%. These results mirror the 1000 Genomes MH frequencies where the European and African American

frequencies for 3rd MHs are expected to be about 50% higher than East Asian frequencies. When mixes

were made with DNA from individuals of different ancestry, the fraction of MHs with 3rd

microhaplotypes increased with some combinations having >30% MHs with >2 haplotypes.

Figure 1 Percent of SNV sets with >2 MHs for individual samples with 106 panel

NGS reads from samples from different individuals of European (Euro), African American (AfAm), and

East Asian (EaAs) ancestry were mixed at frequencies of 0-10%. The number of MH sets with >2

haplotypes were counted and plotted for each individual pair.

While the number of 3rd MHs is useful for assessing whether a sample is contaminated, it is less useful

for estimating the level of contamination as this value reaches a maximum at around 2% contamination

with these coverage levels. The 3rd MH frequency is more useful in this regard so we have used its

median frequency to estimate contamination level. As shown in Fig 2, there is a direct correlation

between the level of contamination and the 3rd MH median frequency. While this comparison does

provide an estimate of contamination levels, this panel and set of analysis parameters yields a nearly

two-fold difference in estimates of contamination for different sample pairs.

Figure 2 Median Frequency for 3rd MHs in individual samples

NGS reads from samples from different individuals of European (Euro), African American (AfAm), and

East Asian (EaAs) ancestry were mixed at frequencies of 0-10%. For all SNV sets with a 3rd MH, the

median frequency was calculated for each pair.

To solve the problem of reduced sensitivity with East Asian samples as well as the high variability of

contamination estimation for different individuals, new MH sets were identified and evaluated for

inclusion in a larger panel. Both old and new MHs were examined for optimal properties. 11 MHs from

the 106 panel with >5 SNV sets with >2 MHs were eliminated to reduce noise. Another 7 SNV sets with

median coverage <250x were eliminated so that more precise 3rd MH frequencies could be generated.

No new SNV sets with those properties were added to the revised panel.

In addition to basic quality metrics, the contamination assay should also be ancestry agnostic so that

contamination can be detected regardless of the DNA source. The population frequencies of 3rd and 4th

MHs were compared based on 1000 Genome data. SNV sets without such data were eliminated from

consideration on the revised panel. While it would have been desirable to use all of the population

groupings from 1000 Genomes in this analysis, there were not enough available SNV sets to allow

balancing of all five major ancestry groupings so only data from African, East Asian, and European

ancestry were applied to achieve balanced variation.

28 of the 106 MHs in the original panel had no 3rd MHs among East Asians. 23 of these were eliminated.

Other MHs that adversely affected ancestry balance were also eliminated. The average frequency of

3rd/4th MHs in the original panel ranges from a low of 8.2% for East Asians to a high of 12.7% in

Europeans. In the revised 164 MH panel, the average frequency of 3rd MHs was 13.9% for all three

ancestries. These results are shown in Table 1 for the complete 106 and 164 panels as well as the SNV

subsets that were shared or unique to each panel. The number of SNV sets with no 3rd MH frequency is

also shown for each ancestry.

Table 1 Panel characteristics

Properties of the 106 and 164 SNV set panels are compared. In addition, the subsets of SNVs in only one

or both panels were also compared. Comparisons include the median coverage, the mean value of SNV

sets with >2 MHs, the total frequency of 3rd and 4th MHs among individuals of African (Afri), East Asian

(EaAs), European (Euro). Admixed American (AdAm), and African Americans (AfAm) as defined by the

1000 Genomes project. The number of SNV sets with no 3rd MH is also shown for each group.

Panel	MHs with LD data	Med Cov	Mean Pure MH>2	Med Len	Mean Fraction of Population with 3rd and 4th MHs						Number of MHs with no examples of 3rd/4th MHs					
					Afri	EaAs	Euro	AdAm	SoAs	AfAm	Afri	EaAs	Euro	AdAm	SoAs	AfAm
106	102	716	2.83	71	0.110	0.081	0.130	0.125	0.129	0.127	9	28	7	3	5	1
164	164	768	1.78	59	0.139	0.139	0.139	0.156	0.153	0.155	9	10	12	1	6	2
Both 106/164	50	776	1.66	46	0.122	0.120	0.127	0.142	0.143	0.140	4	5	3	1	2	1
Only 106	52	695	3.88	92	0.098	0.044	0.132	0.108	0.115	0.114	5	23	4	2	3	0
Only 164	114	768	1.83	61	0.146	0.147	0.144	0.161	0.157	0.161	5	5	9	0	4	1

To assess the impact of the revised SNV sets, *in silico* contamination pairs were examined as before but

with added constraints to minimize noise. To be considered contaminated, we used a threshold of a

minimum of 25 SNV sets with >2 MHs with frequency >0.2%. 3[rd] MHs with frequency <0.2% are usually

technical artifacts due to sequencing and thus artificially lower the median frequency if not eliminated.

15 sets of DNA from each ancestry were mixed as sample/contaminant pairs with averaged results

shown in Fig 3. As expected, the East Asian fraction of >2 MHs was nearly identical to the European

results. The African American samples that we examined had a higher frequency of >2 MHs than the

other groupings but nearly identical to the samples mixed across ancestries. This is predicted by 1000

Genomes data. Their subgroupings of African ancestry included seven different sub-populations, only

one of which was US-based. When the African American subgrouping is compared to the other African

populations, it has a higher 3[rd]/4[th] MH frequency (Table 1) and thus the higher frequency in our

population reflects the fact our African ancestry samples are primarily from African American

individuals.

Figure 3 Percent of SNV sets with >2 MHs for individual samples with 164 panel

The number of 3rd MHs for pairs of individuals of East Asian, African American, and European ancestry is

shown for mixes of 0.1 – 40%. Results from pairs of individuals with different ancestry is included as

"Mixed", all on a log scale.

With the ability to confidently detect whether a sample is contaminated via the number of SNV sets with

>2 MHs, the next goal was to better estimate the level of contamination. The *in silico* mixing studies

provide an empirical calibration, but it is also useful to understand the theoretical basis for those

findings. As shown in Fig 4, the expected frequency of the contaminating 3rd MH depends on the nature

of the starting and incoming genotypes. Some combinations will generate no 3rd MH, others will

generate a 3rd MH with a frequency half the level of incoming contaminant, and others will generate a

genotype that is the same frequency as the contaminant. Since most variant combinations will be at the

50% level, examination of the median value for the 3rd MH should yield a value that is half of the true

contamination level.

Figure 4 Expected frequencies for 3rd MHs based on genotype

The primary limitations on the sensitivity of contamination detection are the practical sequencing

depths for the SNV sets in question and the inherent error rates for sequencing. The higher the

sensitivity desired, the greater sequencing depth required. The usefulness of added depth is limited by

sequencing error rates that are typically on the order of 0.1% for Illumina. It is possible to employ

various error correction methods to add to sensitivity, but the need for that is dependent on the

sensitivity required for the application and its usefulness limited by the available sequencing depth.

Based on these limitations, we have aimed for detection of contamination at least as low as 1% in these

samples.

For determining the 3rd allele frequency, only those MHs with >0.2% frequency are used to avoid over-

inclusion of artifactual MHs that arise from sequencing errors. Additionally, if there are only a few 3rd

MHs, the median frequency could be heavily skewed by having so few values. To avoid this, only

samples with >25 of the potential 164 3rd MHs are considered contaminated. As seen in Fig 3, the

average value for the number of 3rd MHs among all ancestries with 1% contamination is >45 with this

panel so this threshold should not eliminate any truly contaminated samples. All 3rd MHs with frequency

>0.2% are then used for determining the median frequency.

As shown in Fig 5, contamination level using this panel is independent of ancestry. All three ancestries

and mixed samples behave nearly identically. A more detailed listing of results is listed in Table 2 where

the individual ancestries and mixed samples are shown with various thresholds for detection. 15

sample/contaminant pairs were tested for each ancestry with the same number of each mixed ancestry

pairs. The observed frequencies of the median 3rd MH (e.g. 0.005) are multiplied by 2 based on the

discussion above to indicate the actual contamination levels within the sample. When all of the samples

are aggregated, 40% of samples with 0.5% contamination were detected. If the threshold is set at 1%, all

contaminated samples were detected. The detection threshold can be set wherever required based on

the detection needs and the likely impact of contamination on final results.

Table 2 Detection efficiency versus contamination level

Observed 3rd MH Frequency (x2)

East Asian		1	1.5	2	2.5	3	4	5	7	9
	0.5	8	0	0	0	0	0	0	0	0
	1	15	2	0	0	0	0	0	0	0
	1.5	15	12	0	0	0	0	0	0	0
In silico	2	15	14	10	0	0	0	0	0	0
Mixing	2.5	15	15	15	8	0	0	0	0	0
Levels	3	15	15	15	15	6	0	0	0	0
	4	15	15	15	15	15	3	0	0	0
	5	15	15	15	15	15	15	1	0	0
	10	15	15	15	15	15	15	15	15	9
African American		1	1.5	2	2.5	3	4	5	7	9
	0.5	3	0	0	0	0	0	0	0	0
	1	15	0	0	0	0	0	0	0	0
	1.5	15	10	0	0	0	0	0	0	0
In silico	2	15	14	5	0	0	0	0	0	0
Mixing	2.5	15	15	15	4	0	0	0	0	0
Levels	3	15	15	15	14	5	0	0	0	0
	4	15	15	15	15	13	1	0	0	0
	5	15	15	15	15	15	12	2	0	0
	10	15	15	15	15	15	15	15	14	7
European		1	1.5	2	2.5	3	4	5	7	9

In silico Mixing Levels	1	1.5	2	2.5	3	4	5	7	9
0.5	8	0	0	0	0	0	0	0	0
1	15	4	0	0	0	0	0	0	0
1.5	15	13	4	0	0	0	0	0	0
2	15	15	12	0	0	0	0	0	0
2.5	15	15	15	8	0	0	0	0	0
3	15	15	15	13	4	0	0	0	0
4	15	15	15	14	14	3	0	0	0
5	15	15	15	15	15	12	1	0	0
10	15	15	15	15	15	15	15	13	7

Mixed	1	1.5	2	2.5	3	4	5	7	9

In silico Mixing Levels	1	1.5	2	2.5	3	4	5	7	9
0.5	5	0	0	0	0	0	0	0	0
1	15	3	0	0	0	0	0	0	0
1.5	15	14	0	0	0	0	0	0	0
2	15	15	11	0	0	0	0	0	0
2.5	15	15	15	7	1	0	0	0	0
3	15	15	15	15	6	0	0	0	0
4	15	15	15	15	15	2	0	0	0
5	15	15	15	15	15	14	0	0	0
10	15	15	15	15	15	15	15	14	9

All (%)	1	1.5	2	2.5	3	4	5	7	9

In silico Mixing Levels	1	1.5	2	2.5	3	4	5	7	9
0.5	40	0	0	0	0	0	0	0	0
1	100	15	0	0	0	0	0	0	0
1.5	100	82	7	0	0	0	0	0	0
2	100	97	63	0	0	0	0	0	0
2.5	100	100	100	45	2	0	0	0	0
3	100	100	100	95	35	0	0	0	0
4	100	100	100	98	95	15	0	0	0
5	100	100	100	100	100	88	7	0	0
10	100	100	100	100	100	100	100	93	53

Figure 5 Contamination Detection as a function of ancestry

The median frequency of 3[rd] MHs for pairs of individuals of East Asian, African American, and European

ancestry is shown. Results from pairs of individuals with different ancestry is included as "Mixed", all on

a log scale for mixing levels of 0.1 – 40%.

In addition to detecting contamination, it is also possible to use this method to identify sources of

contamination. A partial set of genotypes for the contaminating DNA can be generated from the MH

data and then matched with samples that have been tested previously or contemporaneously to

determine if they are the source of contamination. In a contaminated sample, MHs with 1 or 4 different

haplotypes can be assigned their correct genotypes directly. MHs with 2 or 3 different haplotypes can

generally be assigned a partial or complete genotype based on frequency. The frequency of the $2^{nd}/3^{rd}$

haplotypes will provide at least one and sometimes two genotypes corresponding to the contaminant.

Comparison of the known and partial genotypes with other samples allows them to be ruled in or out as

potential sources of contamination. This procedure has been applied to multiple contaminated samples

to successfully determine the source in real clinical laboratory situations. Often, the contamination

source can be shown not to arise from any samples known to have been handled in the laboratory so

the contamination must have originated prior to sample receipt.

Discussion

The level of DNA contamination that is an issue varies across applications. In some cases, contamination

levels that impact results are obvious. With other applications, signals arising from contamination can be

easily confused with real signals. With diagnosis of somatic variants in cancer, this is especially a

problem because variant identification is attempted at and below the very limits of variant detection.

Being able to separate the tumor signal from the contamination noise is critical for the proper diagnosis

of tumor etiology. Because tumor samples often have highly unusual copy number patterns, use of

simple genotype frequencies can cause errors in contaminant detection. Multiple samples that we

assessed as contaminated based on standard methods using raw genotype frequencies were found to

be pure by MH analysis. In contrast to the FFPE tumor samples, circulating free DNA (cfDNA) purified

from the plasma of cancer patients is not as prone to the same issues because most of the cfDNA

purified from plasma is from normal cells so typical Mendelian ratios are observed. MH analysis in such

samples is still valuable for sensitive contamination detection, but such samples are not as likely to

generate false positives as DNA from FFPE tumor samples where copy number variation is much more

significant.

For most purposes, the approximation that the percent contamination is twice the median frequency for

the 3rd MH is sufficient. This relationship arises from the likelihood that a 3rd MH is most likely

heterozygous so is present on only one of the two incoming alleles. The uncertainty around this

probability introduces the potential for multiple sources of error and those should be considered if a

more exact measurement is necessary. At low coverage, two confounding factors arise. First, low

coverage leads to high AF variability due to the stochastic nature of read accumulation in poorly covered

SNV sets. Only SNV sets with 3rd MHs are included in the median calculation rather than all SNV sets so,

at low coverage, AF inflation occurs among the variants that are counted. The observed AF is higher than

reality because there are not enough reads to generate the proper AF value. By requiring a large number

of 3rd MHs prior to calling a sample contaminated, this effect is mitigated. Superimposed on this effect is

the appearance of homozygous 3rd MHs before heterozygous 3rd MHs as coverage increases. If coverage

is deep, this is not an issue but, at low coverage, homozygous variants will appear first because they are

being double counted. Further complicating the relationship is the presence of low frequency technical

artifacts and the copy number issues that can occur with cancer samples. Many cancer samples have

large copy number variations so different SNV sets will appear at different fractional rates depending on

whether the 3rd MH is under- or over-represented in copy number changes. The more MH sets used in

the calculations, the more likely it is that these effects can be averaged out.

The MH approach is perfectly suited for use when somatic variants are sought because such studies use

very high coverage in order to detect rare variants. The sensitivity of contamination detection is

dependent on the coverage of the regions used in the analysis. Thus, other panel-based and exome NGS

approaches are also immediately extendable to similar studies. A typical whole genome approach with

less than 50x coverage would not be amenable to sensitive detection using individual MH sets because

the low coverage would prevent the observation of infrequent 3rd MHs. To overcome this, combinations

of MH sets could be used. The simplest approach would be to combine MH sets that are in high LD and

combine the appropriate SNV calls in order to achieve the needed depth of coverage across multiple

high LD regions. Other methods for pooling MH sets could also be used. With the entire genome as a

source, MH sets that behave as needed and are subject to the lowest error rates could be employed.

The same techniques used for these cancer samples can also be used with forensic samples to detect

major and minor contributors to a sample. Unlike tumor samples, copy number should not be an issue,

which makes interpretation easier. Like the tumor samples we have examined, the major and minor

contributors to a sample may have different DNA qualities that can affect the accuracy of MH

frequencies. Some MHs used in these panels have more than 4 potential haplotypes so could indicate

more than one contaminant. Panels designed with a larger number of >4 potential MHs would be

appropriate when forensic samples are being examined in order to clarify the number of contributors to

a sample. As with contamination assignment described above, MH assignment can be accomplished at

high enough efficiency to identify possible sources of the contamination as well as ancestry.

Another useful application of the MH methods is for more sensitive detection of fetal DNA in Non-

invasive prenatal testing (NIPT). Panels can be designed to focus on genomic regions with the most

likely potential copy number variation (e. g. chr 5, chr13, chr18, or chr21) and the frequency of 3[rd] MHs

in targeted regions compared to the rest of the genome. In all regions, fetal DNA will share one

genotype with the mother (main contributor of DNA) and have one from the father. In genomic regions

where the mother is heterozygous and the father's contribution to the fetal DNA is a different allele,

there will be a low frequency 3[rd] MH. If the frequency of that 3[rd] MH varies across genomic regions, it is

indicative of a copy number variation. Because the fetal "contamination" signal will always be just a

single copy of non-matching genotype for a single birth, the frequency distributions will be less variable

than with a typical contaminant. For fraternal twin births, MH frequencies should be the sum of the

individual fetal contributors. For SNV sets where both parents are heterozygous with four different MHs,

it would be possible to see 4 MHs rather than just 3. With the potential for high copy depth and lower

detection levels, NIPT could be done earlier in a pregnancy using the MH approach rather than standard

methods.

The methods described here provide a foundation for examining DNA mixtures in a variety of contexts.

With the appropriate choice of SNV panels and high sequencing depth, very low levels of secondary DNA

can be detected and this information can be valuable in a variety of applications.

Abbreviations

AdAm: Admixed American

AF: Allele Frequency

AfAm: African American

Afri: African

cfDNA: circulating free DNA

EaAs: East Asian

Euro: European

FFPE: Formalin-fixed Paraffin-embedded

FP: False Positive

LD: Linkage Disequilibrium

MH: Microhaplotype

NGS: Next-generation Sequencing

NIPT: Non-invasive Prenatal Testing

PCA: Principle Component Analysis

SNV: Single Nucleotide Variant

SoAs: South Asian

STR: Short Tandem Repeat

References

1. Flickinger M, Jun G, Abecasis GR, Boehnke M, Kang HM. Correcting for Sample Contamination in

Genotype Calling of DNA Sequence Data. Am J Hum Genet. 2015 Aug 6;97(2):284-90.

2. Cibulskis K, McKenna A, Fennell T, Banks E, DePristo M, Getz G. ContEst: estimating cross-

contamination of human samples in next-generation sequencing data. Bioinformatics. 2011 Sep

15;27(18):2601-2.

3. Sakarya, 2017, https://grail.com/wp-

content/uploads/2018/05/HiTSeq_2017_Sakarya_Conta_POS_Final.pdf

4. Fiévet A, Bernard V, Tenreiro H, Dehainault C, Girard E, Deshaies V et al. ART-DeCo: easy tool for

detection and characterization of cross-contamination of DNA samples in diagnostic next-generation

sequencing analysis. Eur J Hum Genet. 2019 May;27(5):792-800.

5. Kidd KK, Pakstis AJ, Speed WC, Lagace R, Chang J, Wootton S, et al. Microhaplotype loci are a powerful

new type of forensic marker. FSI Supplement Series. 2013. 4 (1):e123-e124.

6. Kidd KK, Pakstis AJ, Speed WC, Lagacé R, Chang J, Wootton S, et al. Current sequencing technology

makes microhaplotypes a powerful new type of genetic marker for forensics. Forensic Sci Int Genet.

2014 Sep;12:215-24.

7. Kidd KK, Speed WC. Criteria for selecting microhaplotypes: mixture detection and deconvolution.

Investig Genet. 2015 Jan 28;6:1.

8. Oldoni F, Kidd KK, Podini D. Microhaplotypes in forensic genetics. Forensic Sci Int Genet. 2019 Jan;

38:54-69.

9. Debeljak M, Freed DN, Welch JA, Haley L, Beierl K, Iglehart BS, et al. Haplotype counting by next-

generation sequencing for ultrasensitive human DNA detection. J Mol Diagn. 2014 Sep;16(5):495-503.

10. Bulbul O, Pakstis AJ, Soundararajan U, Gurkan C, Brissenden JE, Roscoe JM, et al. Ancestry inference

of 96 population samples using microhaplotypes. Int J Legal Med. 2018 May;132(3):703-711.

11. Kidd KK, Speed WC, Pakstis AJ, Podini DS, Lagacé R, Chang J, et al. Evaluating 130 microhaplotypes

across a global set of 83 populations. Forensic Sci Int Genet. 2017 Jul;29:29-37.

12. Oldoni F, Yoon L, Wootton SC, Lagacé R, Kidd KK, Podini D. Population genetic data of 74

microhaplotypes in four major U.S. population groups. Forensic Sci Int Genet. 2020; 49:102398.

13. Phillips C, McNevin D, Kidd KK, Lagacé R, Wootton S, de la Puente M, et al. MAPlex - A massively

parallel sequencing ancestry analysis multiplex for Asia-Pacific populations. Forensic Sci Int Genet. 2019

Sep;42:213-226.

14. Phallen J, Sausen M, Adleff V, Leal A, Hruban C, White J, et al. Direct detection of early-stage cancers

using circulating tumor DNA. Sci Transl Med. 2017 9:2415.

15. Georgiadis A, Durham JN, Keefer LA, Bartlett BR, Zielonka M, Murphy D, et al. Noninvasive Detection

of Microsatellite Instability and High Tumor Mutation Burden in Cancer Patients Treated with PD-1

Blockade. Clin Cancer Res. 2019 25:7024-7034.

16. Wood DE, White JR, Georgiadis A, Van Emburgh B, Parpart-Li S, Mitchell J, et al. A machine learning

approach for somatic mutation discovery. Sci Transl Med. 2018 10:7939. PMID: 3018562

17. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational constraint

spectrum quantified from variation in 141,456 humans. Nature. 2020 581: 434-443.

18. Machiela MJ, Chanock SJ. LDlink: a web-based application for exploring population-specific

haplotype structure and linking correlated alleles of possible functional variants. Bioinformatics. 2015 Jul

2

19. Sehn JK, Spencer DH, Pfeifer JD, Bredemeyer AJ, Cottrell CE, Abel HJ, et al. Occult Specimen

Contamination in Routine Clinical Next-Generation Sequencing Testing. Am J Clin Pathol. 2015

Oct;144(4):667-74.