Long term survival of *Dehalococcoides mccartyi* strains in mixed cultures under electron

acceptor and ammonium limitation

Nadia Morson[1], Olivia Molenda[2], Katherine J. Picott[2], Ruth E. Richardson[3] and Elizabeth A.

Edwards[1,2]

[1]Department of Cell & Systems Biology, University of Toronto, Toronto, ON, Canada

[2]Department of Chemical Engineering & Applied Chemistry, University of Toronto, Toronto,

ON, Canada

[3]School of Civil and Environmental Engineering, Cornell University, Ithaca, NY, United States

of America

Corresponding Author: Elizabeth A. Edwards, Department of Chemical Engineering and Applied

Chemistry, University of Toronto, 200 College Street, Toronto, ON, Canada M5S 3E5; Tel.: +1

416 946 3506; fax: 416 978 8605; e-mail: elizabeth.edwards@utoronto.ca

Keywords: bioremediation, reductive dechlorination, quantitative PCR, *Dehalococcoides*, vinyl

chloride

TOC Art

Abstract

Few strains of *Dehalococcoides mccartyi* harbour and express the vinyl chloride reductase

(VcrA) that catalyzes the dechlorination of vinyl chloride (VC), a carcinogenic soil and

groundwater contaminant. The *vcrA* operon is found on a Genomic Island (GI) and therefore

believed to participate in horizontal gene transfer. To try to induce horizontal gene transfer of the

vcrA-GI, we blended two enrichment cultures in medium without ammonium while providing

VC. We hypothesized that these conditions would select for a mutant strain of *D. mccartyi* that

could both fix nitrogen and respire VC. However, after more than 4 years of incubation, we

found no evidence for horizontal gene transfer of the *vcrA*-GI. Rather, we observed VC-

dechlorinating activity attributed to the trichloroethene reductase TceA. Sequencing and protein

modelling revealed a mutation in the predicted active site of TceA which may have influenced

substrate specificity. We also identified two nitrogen-fixing *D. mccartyi* strains in the KB-1

culture. The presence of multiple strains of *D. mccartyi* with distinct phenotypes is a feature of

natural environments and certain enrichment cultures (such as KB-1) and may enhance

bioaugmentation success. The fact that multiple distinct strains persist in the culture for decades

and that we could not induce horizontal gene transfer of the *vcrA*-GI suggests that it is not as

mobile as predicted, or that mobility is restricted in ways yet to be discovered to specific sub-

clades of *Dehalococcoides*.

Introduction

Chlorinated solvents are among the most prevalent and persistent soil and groundwater contaminants in industrialized countries (Moran, Zogorski and Squillace 2007). Tetra- or perchloroethene (PCE) and trichloroethene (TCE) contamination originates from the on-going use of dry-cleaning solvents and metal degreasing agents, respectively (Doherty 2000). These compounds, and their transformation intermediates such as vinyl chloride (VC), are known to have toxic or carcinogenic effects so their widespread soil and groundwater contamination poses a great concern to human health (Müller *et al.* 2004). In the late 1980s, following the discovery that microorganisms could completely dechlorinate these solvents under anaerobic conditions (Freedman and Gossett 1989), bioremediation and bioaugmentation emerged as highly successful treatment options for these problematic pollutants (Löffler and Edwards 2006). A microbial culture enriched from soil contaminated with chlorinated solvents in Southern Ontario, Canada, resulted in the KB-1™ mixed microbial consortium (Duhamel *et al.* 2002). KB-1™ has been used commercially for bioaugmentation for more than 20 years to remediate chlorinated compounds (Major *et al.* 2002; Löffler and Edwards 2006). In the KB-1™ culture, chlorinated ethenes, such as PCE and TCE, are sequentially dechlorinated via *cis*-dichloroethene (cDCE) and VC to ethene (Duhamel *et al.* 2002; Perez-de-Mora *et al.* 2017; Molenda *et al.* 2020). Reductive dechlorination in KB-1™ is primarily performed by multiple *Dehalococcoides mccartyi* strains in a growth-linked process called organohalide respiration (Duhamel *et al.* 2002; Perez-de-Mora *et al.* 2017; Molenda *et al.* 2020). Based on 16S rRNA sequence similarity, all *D. mccartyi* strains are categorized into three phylogenetic clades or sub-groups: Pinellas, Victoria, and Cornell. The Pinellas clade is represented by strain CBDB1, the Victoria clade is represented by strain VS, and the Cornell clade is represented by the type strain, strain 195 (Hendrickson *et al.*

2002; Löffler *et al.* 2013). Lab-grown KB-1 cultures enriched on different chlorinated substrates

select for different strains of *D. mccartyi* that express different reductive dehalogenases (Perez-

de-Mora *et al.* 2017). Reductive dehalogenases belong to a broad protein family (PF13486) and

have been classified into Ortholog Groups (OGs) on the basis of >90% amino acid pairwise

identity (Hug *et al.* 2013; Molenda *et al.* 2020). In the TCE-enriched KB-1 culture, strains

belonging to the Pinellas clade are dominant, expressing the VC reductases VcrA (OG 8) and

BvcA (OG 28) (Molenda *et al.* 2020). Recently, the TCE-enriched KB-1 culture was also found

to contain low abundance Cornell clade strains, which express the TCE reductase TceA (OG 5)

(Molenda *et al.* 2020). VC reductases are the most critical since VC is the most toxic

dechlorination intermediate and following VC dechlorination, complete dechlorination is

achieved.

Interestingly, the operon which encodes VcrA was found on a mobile genetic element,

called the *vcrA*-Genomic Island (GI) (McMurdie *et al.* 2011). In the KB-1 TCE-enrichment

culture, the *vcrA*-GI was identified in a circularized and extrachromosomal state within the cell

which could theoretically be transferred between *D. mccartyi* strains through horizontal gene

transfer (HGT) (Regeard *et al.* 2005; McMurdie *et al.* 2011; Molenda *et al.* 2019). However,

there is no direct evidence of HGT of the *vcrA*-GI, and the mechanism of transfer between *D.*

mccartyi remains unknown. To try to induce HGT of the *vcrA*-GI, we blended the KB-1 TCE-

enrichment culture with another mixed culture, called Donna II (Fennell, Gossett and Zinder

1997). We called the blend of the two enrichment cultures DKB (Donna + KB-1). The KB-1

TCE-enriched consortium is the ideal *vcrA*-GI donor because the *vcrA*-containing *D. mccartyi*

strain is highly abundant in this culture. The Donna II culture is a mixed microbial consortium

that contains only one *D. mccartyi* strain, strain 195, that dechlorinates PCE to VC via

organohalide respiration utilizing the PCE dehalogenase PceA (OG 30) and TceA (Fennell,

Gossett and Zinder 1997). Subsequently, VC is dechlorinated to ethene slowly via co-

metabolism since *D. mccartyi* strain 195 does not contain the *vcrA*-GI or any other VC

reductases and therefore cannot grow on VC (Regeard *et al.* 2005). Of particular interest for

experimental design, *D. mccartyi* strain 195 is capable of nitrogen fixation in the absence of

available nitrogen sources via the *nif* operon (Lee *et al.* 2009). Whereas the *vcrA*-GI-containing

D. mccartyi strain from TCE-enriched KB-1 does not have a nitrogen fixation operon and thus an

incapability for nitrogen fixation. In a theoretical HGT event, the KB-1 TCE-enrichment culture

would contain the donor strain of the *vcrA*-GI and *D. mccartyi* strain 195 in Donna II would be

the recipient strain. By applying the selective pressure of providing only VC as the sole energy

source (electron acceptor) in medium without ammonium, we thought that a hybrid *D. mccartyi*

strain which could fix nitrogen and respire VC would emerge.

After 4 years of observation, we found no evidence to support horizontal gene transfer of

the *vcrA*-GI. Instead, in two of three replicates of the DKB culture, we observed VC

dechlorination activity even though *vcrA* gene copies were low and *tceA* and *D. mccartyi* 16S

rRNA copies were high. We then sequenced and modelled the TceA of the DKB culture and

found a mutation in the predicted active site that is hypothesized to influence substrate

specificity. Additionally, over the course of the study, two previously unknown Cornell strains of

D. mccartyi were identified in KB-1 that contain nitrogen-fixing genes. We determined that these

genes were expressed and active at low ammonium concentrations, a beneficial feature of the

KB-1 culture for bioremediation.

Materials and Methods

Enrichment cultures

The KB-1 culture originated from microcosms prepared with aquifer materials from a TCE-

contaminated site in southern Ontario in 1996 (Duhamel *et al.* 2002). The KB-1 TCE-enrichment

culture is maintained biweekly with 0.76 mM TCE as the electron acceptor and $5 \times$ electron

equivalents of methanol (MeOH) as the electron donor, referred to as KB-1/TCE-MeOH, as

previously described (Duhamel, Mo and Edwards 2004; Duhamel and Edwards 2006, 2007). The

Donna II culture originated from an enrichment culture seeded with digester sludge from a

wastewater treatment plant in Ithaca, NY, USA. The Donna II culture was maintained batch-style

at Cornell University with 110 µM of PCE as the electron acceptor and butyrate as the electron

donor, as previously described (Smatlak, Gossett and Zinder 1996; Fennell, Gossett and Zinder

1997). The "DKB" culture was formed at Cornell University in 2012 when 1000 mL of KB-

1/TCE-MeOH and 700 mL of Donna II were combined. The DKB culture was maintained with

PCE and butyrate, similar to the Donna II culture. In 2015, the DKB culture was shipped to the

University of Toronto and was subsequently used as inoculum to create DKB sub-cultures grown

under different conditions of varying selective pressures, as described below.

Experimental Set-up and Monitoring

DKB sub-cultures were fed either PCE or VC as electron acceptor, and butyrate as electron

donor. Additionally, DKB sub-cultures were either cultured in medium with ammonium (▲), or

without ammonium (△) requiring nitrogen fixation. Therefore, four experimental conditions

were prepared: (i) DKB PCE ▲, (ii) DKB PCE △, (iii) DKB VC ▲, and (iv) DKB VC △

(Figure S1). Additionally, a KB-1 control culture, KB-1/TCE-MeOH, was used as inoculum for

sub-culture triplicates into medium without ammonium and amended with VC, referred to as

KB-1 VC ∆; this condition was expected to be a negative control for KB-1 growth under VC-

degrading, ammonium-limiting conditions (Figure S1). To create each sub-culture, each 2 mL

sample of culture (KB-1/TCE-MeOH or DKB) was centrifuged at 13,000 × g for 15 minutes at

room temperature, the supernatant was discarded, and the pellet was resuspended by flicking and

pipetting in 2 mL of anaerobic, autoclaved distilled H_2O. This process was repeated three times,

to wash ammonium from the pellet. Following, the 2 mL volume of washed inoculum was

transferred into 198 mL of anaerobic mineral medium (Duhamel and Edwards 2007) with (10

mM) or without NH_4Cl in 250 mL serum bottles sealed with butyl rubber stoppers (Geo-

Microbial Technologies Inc.). An additional 500 µL of vitamin stock (Edwards and Grbic-Galic

1994) was added to each bottle during set-up, and 50 µL was added for each 20 mL of medium

added during medium amendments. The culture was purged with N_2:CO_2 (80:20) gas, and the

headspace was over-pressurized with N_2:CO_2 for 3 seconds. The DKB sub-cultures were

amended periodically with 310 µelectron equivalents (µeeq) of gaseous VC (5 mL) or neat PCE

(4 µL) as electron acceptor, and 4 × eeq sodium butyrate stock as electron donor. Electron

equivalents were used to calculate the mass of electron donor or acceptor fed to the cultures as a

way to establish a ratio of electrons required for organohalide respiration given that each

hydrogenolysis reaction requires 2 electrons for the removal of a chlorine atom. Therefore, the

KB-1/TCE-MeOH sub-cultures, KB-1 VC ∆, were maintained with 310 µeeq VC as electron

acceptor, and 5 × eeq of methanol and ethanol mixture (50:50 on an eeq basis) as electron donor.

Cumulative electron acceptor consumed was monitored over time. Acceptor and donor were re-

amended as needed when depleted in each sub-culture, and medium was amended following

large volume sampling for nucleic acid extraction in batch style maintenance. Triplicate bottles

were prepared for each of the four test conditions. In 2016, to reduce number of bottles

monitored, only one of the triplicate cultures from the DKB sub-cultures with ammonium was

maintained; prior to this time, these triplicates were behaving similarly. Therefore, from 2016

onward, 8 DKB sub-cultures and three KB-1 VC Δ cultures were maintained and analyzed.

DNA Extraction

Samples were taken from KB-1/TCE-MeOH, Donna II, DKB, and DKB sub-cultures at various

times throughout the experiment for DNA extraction. For each DNA sample, a volume of 20 mL

of liquid culture was removed anaerobically. The 20 mL aliquot was centrifuged in a sealed

Falcon conical centrifuge tube at $6870 \times g$ for 20 minutes at room temperature, using a swinging-

bucket rotor. Aerobically, the supernatant was decanted, and the pellet was resuspended

immediately using 60 µL of Solution C1 and 500 µL of liquid from the PowerBead tube, from

the DNeasy PowerSoil Kit (Qiagen). All subsequent kit instructions were followed. Final DNA

elution volume was 50 µL of Solution C6 (elution buffer). DNA was quantified by NanoDrop or

using the High-Sensitivity DNA kit with the Qubit® 3.0 Fluorometer (ThermoFisher Scientific).

All DNA extracts were stored at -80°C.

RNA Extraction and cDNA synthesis

RNA was extracted from 10 mL of the KB-1 VC Δ cultures and the KB-1/TCE-MeOH culture.

Cells were pelleted by centrifuging at $5000 \times g$ at 4°C for 15 minutes. Pellets were then

resuspended in 500 µL of supernatant and stabilized with 1 mL of RNAprotect Bacteria Reagent

(Qiagen). The RNeasy® Protect Bacteria Mini Kit (Qiagen) was used to extract RNA from the

cells by physical disruption via bead-beating. The kit procedure was followed, and RNA was

eluted in 30 µL of RNase-free water. Following, RNA was DNase-treated using 10 µL of DNase

A and 70 µL of DNase buffer for 30 seconds. To purify, the RNA was cleaned up using the

RNeasy® MinElute® Cleanup kit (Qiagen) and eluted in 15 µL of RNase-free water. Following,

cDNA was synthesized from the RNA using the Invitrogen Superscript® VILO cDNA synthesis

kit (Invitrogen), and procedure was followed according to kit instructions. cDNA was quantified

using the High-Sensitivity DNA kit with the Qubit® 3.0 Fluorometer. All cDNA extracts were

stored at -80°C.

***D. mccartyi* strain biomarker selection for Quantitative PCR of 16S rRNA and functional**

genes

To track the growth of the different *D. mccartyi* strains in the DKB sub-cultures, we designed

strain-specific biomarkers for quantitative PCR (qPCR). The abundance of all *D. mccartyi* strains

was quantified by the 16S rRNA gene, using primers Dhc1f and Dhc264r (Hendrickson *et al.*

2002), since there is a single 16S rRNA gene copy per genome. To quantify the *vcrA*-containing

strain from KB-1/TCE-MeOH and all extrachromosomal *vcrA*-GIs, we used primers vcrA670f

and vcrA440r targeting the *vcrA* gene (Molenda, Quaile and Edwards 2016). The tceA500f and

tceA795r (Fung *et al.* 2007) primers were used for tracking *tceA*-containing *D. mccartyi* strains

in KB-1/TCE-MeOH and *D. mccartyi* strain 195. To quantify strain 195, we designed primers

for a unique *rdhA* (DET_RS00960), that is truncated (truncated *rdhA*, "t*rdhA"*) and predicted to

encode a non-functional reductive dehalogenase. It is only 1200 bp long compared to functional

rdhA with lengths of approximately 1500 bp. As well, this gene contains two iron-sulfur cluster

binding domains $(CX_2CX_2CX_3CP)_2$ but does not contain commonly conserved motifs: a twin-

arginine TAT membrane export sequence (RRXFXK) nor a cobalamin binding domain. Primers

for t*rdhA* were designed for the *D. mccartyi* strain 195 genome, using Primer2 (Untergasser *et al.*

2012) in Geneious 8.1.9 (Kearse *et al.* 2012). To track nitrogen fixation (*nif)* genes, we

quantified the *nifD* gene which encodes the nitrogenase molybdenum-iron protein ▢-chain. The

nifD primers used in this study were used to characterize nitrogen fixation in *D. mccartyi* strain

195 (Lee *et al.* 2009). These *nifD* primers were also used for transcription analysis of nitrogen

fixation genes in KB-1 VC Δ and KB-1/TCE-MeOH, by reverse transcription (RT)-qPCR.

Lastly, we went back to archived DNA samples to quantify the vinyl chloride reductase, *bvcA*

(Figure S2), using primers bvcA318f and bvcA555r (Waller *et al.* 2012). More information on

primers can be found in Supplementary Information (Table S1).

As a qPCR standard for absolute quantification, a plasmid with concatenated sequences

corresponding to the *D. mccartyi* 16S rRNA gene, *vcrA*, *tceA,* and *bvcA* was used, as previously

described (Molenda *et al.* 2019). This concatenated plasmid allowed us to calculate accurate

ratios of these *rdhA* genes to 16S rRNA gene copies. For *nifD* and t*rdhA* qPCR standards,

regions were PCR amplified, purified and cloned into *Escherichia coli* (TOPO™ TA Cloning™

Kit for Sequencing, Invitrogen) (Additional methods in Supplemental Information).

All qPCRs were prepared in a UV PCR cabinet and each qPCR was run in duplicate or

triplicate, using a CFX96 real-time PCR detection system. Each 20 µL qPCR reaction was

prepared in UV-treated UltraPure nuclease-free water containing 10 µL of EvaGreen® Supermix

(Bio-Rad Laboratories, Hercules, CA), 0.5 µL of each primer (forward and reverse, from 10 µM

stock solutions), and 2 μL of DNA template or standard plasmid dilution series, from 10^1 to 10^7

copies of plasmid per μL. Thermocycler program: initial denaturation at 95°C for 2 min,

followed by 40 cycles of denaturation at 98°C for 5 s, varied annealing temperatures (Table S1),

followed by extension for 10 s at 72°C. All qPCR standards and quality metrics (efficiency,

standard curve details, etc.) can be found in Supplementary Information (Table S2), with

calculations of absolute gene abundances per mL of culture (Table S3). The quantification limit

(corresponding to lowest calibration standard) was $\sim 1 \times 10^3$ copies/mL, but values as low as

1×10^1 copies/mL could be detected. To compare *nifD* expression among different experimental

conditions, the absolute *nifD* gene abundances of genomic DNA and cDNA were determined.

Using this data, we calculated *nifD* transcript per gene (TPG) ratios, as the ratio of transcript

copies per mL of culture to gene copies per mL of culture.

Clone Library Preparation and Sequencing of *tceA*

A clone library of *tceA* was generated from one of the DKB VC ∆ cultures. As control, a *tceA*

clone library was generated in parallel using the KB-1/TCE-MeOH culture. The two *tceA* clone

libraries were generated using the TOPO™ XL-2 Complete PCR Cloning Kit (Invitrogen). PCR

primers were designed to amplify the 2200 bp *tceAB* region (Table S1). Blunt end *tceA* PCR

products were produced using the Platinum SuperFi polymerase, bands were extracted from an

agarose gel, and cloned into One Shot™ OmniMAX™ 2 T1R Chemically Competent *E. coli*

cells. 50 µL or 100 µL of transformed *E. coli* were spread on 50 µg/mL kanamycin LB plates,

with 40 µL of 40 mg/mL X-gal in dimethylformamide (DMF) solution and 40 µL of filter-

sterilized 0.1 M IPTG for blue-white selection. Following, 10 colonies were selected each from

the DKB sub-culture and from the KB-1/TCE-MeOH culture and transferred cultures onto a

patch plate for further analysis. From the patch plate, colonies were incubated in to 10 mL of 50

µg/mL kanamycin LB broth, and plasmids were extracted using a QIAprep Spin Miniprep Kit

(Qiagen). Each plasmid was PCR amplified using the T3 and T7 cloning primers to confirm

successful transformation. For the first round of Sanger sequencing, plasmids were sequenced

using the T3 and T7 cloning primers, at the SickKids Center for Applied Genomics (TCAG)

sequencing/synthesis facility (Toronto, Canada). Ten sequences each from the forward and

reverse were then aligned, and two consensus sequences were generated at a 99% nucleotide

identity threshold. Using the consensus sequences, primers were designed (Table S1) for the

remaining 845 bp. For the second round of sequencing, the final 99% nucleotide identical

consensus sequences were generated by aligning these 20 sequences. The cloned *tceA* sequences

from this study, the *tceA* published on NCBI (AAW39060), and the *tceA* from the Donna II

metagenome (IMG-M taxon ID: 2032320001) were aligned using the MUSCLE aligner (Edgar

2004) in Geneious 8.1.9.

TceA protein modelling

Protein models were independently produced for TceA from the *Dehalococcoides mccartyi* 195

isolate and the TceA from one of the DKB VC ∆ cultures. Each sequence was first trimmed to

remove the TAT signal peptide sequence predicted by SignalP-5.0 (Almagro Armenteros *et al.*

2019). The models were produced from four different predictive modelling servers, AlphaFold2,

Robetta, I-TASSER, and Phyre2, and assessed based on quality scores determined by the

MolProbity assessment tool (Chen *et al.* 2010; Roy, Kucukural and Zhang 2010; Song *et al.*

2013; Kelley *et al.* 2015; Studer *et al.* 2020; Jumper *et al.* 2021). The AlphaFold2 and Robetta

models had the highest quality scores and were used for further analysis; however, all models

were used to validate the predicted spatial locations of the mutated residues. The cobalamin

cofactor was modelled into the active site using AutoDock Vina which was consistent with the

docking position of known crystal structures (Trott and Olson 2009). The iron-sulfur clusters

were superimposed from the *Sulfurospirillum multivorans* PceA crystal structure (PDB ID:

4UR2) (Bommer *et al.* 2014). The substrate access channels were predicted using the CAVER

3.0 plugin in PyMol v2.3.4, only the channels predicted on the catalytic face of the cobalamin

cofactor were kept (Chovancova *et al.* 2012). Models were imaged and the molecular

interactions were predicted using PyMol v2.3.4 (The PyMOL Molecular Graphics System,

Version 2.3.4 Schrödinger).

Nitrogenase operon gene and protein alignments

The *D. mccartyi* strain 195 nitrogen fixation operon (DET_RS05950-5990), encoding the

nitrogenase enzyme, was aligned with the nitrogenase operons of the two KB-1/TCE-MeOH

strains (KBTCE2: B1773_05150-05185, KBTCE3: B1774_04800-04835). An alignment of these

operons was performed using the built-in MUSCLE aligner (Edgar 2004) in Geneious 8.1.9.

Results and Discussion

Long term growth of DKB sub-cultures under varying degrees of selective pressure

To analyze the effects of experimental conditions on growth and dechlorination, we monitored

the cumulative electron acceptor consumed over 4 years (Table S3). Overall, cultures grown

without ammonium (△) did not consume as much electron acceptor as cultures with ammonium

(▲) in each series (Figure 1A). The DKB PCE ▲ culture consumed twice as much PCE as the

DKB PCE △ cultures, while the DKB VC ▲ culture consumed nearly six times more VC than

the DKB VC △ cultures over the same time. The DKB VC △ cultures, grown under the highest

selective pressure conditions of VC without ammonium, dechlorinated the least amount of

electron acceptor of all DKB sub-cultures (Figure 1A). These findings confirmed that the

selective pressures on the DKB sub-cultures impacted the metabolism of *D. mccartyi* to varying

degrees.

In each DKB and KB-1 culture, we tracked the abundance of key genes of *D. mccartyi*

strains and related that to strain abundance (Figure 1B-H). The *D. mccartyi*-specific 16S rRNA

gene target tracked the abundance of all *D. mccartyi* strains ("total *Dhc*"). The *vcrA* gene target

provided an indication of *D. mccartyi* strain KBTCE1 from KB-1 that contained the *vcrA*-GI

("KB-1/*vcrA*") and captured any extrachromosomal copies of the *vcrA*-GI. Similarly, the *bvcA*

gene target measured the *bvcA*-containing strain from KB-1 ("KB-1/*bvcA*"). *D. mccartyi* strain

195 and two KB-1 strains, KBTCE2 and KBTCE3, contained both *tceA* and *nifD* gene targets, so

the strain 195 biomarker (truncated *rdhA*, or t*rdhA*) was used to distinguish between *D. mccartyi*

from KB-1 ("KB-1/*tceA*") and strain 195 (195/t*rdhA*").

Trends in the abundance of key strain biomarkers helped decipher what occurred in the

DKB (blend of KB-1 and Donna II) and KB-1 control cultures over nearly 4 years of incubation.

The KB-1/*vcrA* strain was highly abundant (10^7 to 10^9 copies/mL of culture) in all cultures

except the DKB VC △ (no ammonium) replicate cultures (Figure 1D-F). In DKB VC △1 and △2

cultures (Figure 1D-E), the KB-1/*vcrA* strain decreased in abundance over time, whereas in the

DKB VC △3 culture, KB-1/*vcrA* abundance was maintained (Figure 1F). Conversely, an

increase in the 195/t*rdhA* strain was observed in DKB VC △1 and △2 cultures, whereas it was

below or near the quantification limit of 10^3 copies/mL in the DKB VC △3 culture throughout

the experiment (Figure 1D-F). All together, these results suggest that the dominant strain in DKB

VC △1 and △2 cultures was *D. mccartyi* strain 195 from the Donna II culture, and the dominant

strain in the DKB VC △3 culture was *D. mccartyi* strain KBTCE1 from the KB-1 culture (Figure

S3). An interesting and complementary result is that the non-dominant *D. mccartyi* strains, such

as KB-1/*bvcA* and KB-1/*tceA*, persisted in low abundances throughout the 4 years of incubation.

If the *vcrA* island were to be transferred to strain 195, the "hybrid" *D. mccartyi* strain would have

both *vcrA* and t*rdhA* and thus would appear in Figures 1D-F as an equal abundance of KB-1/*vcrA*

and 195/t*rdhA,* yet this result was never observed. Therefore, these results refute the existence of

a predicted hybrid strain that acquired the *vcrA*-GI.

However, early in the experiment, we noticed that KB-1 VC Δ cultures were able to

dechlorinate VC even without ammonium provided in the growth medium (Figure 1A, Table

S4). This condition was intended to be a control where dechlorination and growth were not

anticipated. As dechlorination continued, it was suspected that the KB-1 VC Δ culture may

contain nitrogen-fixing bacteria, later justified by the high abundance of *D. mccartyi nifD*

copies/mL of culture (Figure 1F). We therefore analyzed nitrogen fixation activity in the KB-

1/TCE-MeOH and KB-1 VC Δ cultures.

Complete nitrogenase operons identified in two KB-1 strains

In 2012, when this experiment was conceived, the only *D. mccartyi* strain that was known to fix

nitrogen was the *D. mccartyi* strain 195 isolate (Lee *et al.* 2009). After the DKB experiment had

begun, metagenomic sequencing of the KB-1/TCE-MeOH culture revealed the presence of more

than three strains of *D. mccartyi* (Molenda *et al.* 2019). The *vcrA*-GI containing strain, KBTCE1,

was the most abundant strain and belonged to the Pinellas clade. However, the other two strains,

KBTCE2 and KBTCE3, were both *tceA*-containing Cornell strains that were not previously

detected in the culture. Genome analysis revealed that strains KBTCE2 and KTBCE3 contained

complete nitrogenase operons, encoded by *nif* genes, and were therefore predicted to fix

nitrogen. The nitrogenase operons in strains KBTCE2 and KBTCE3 share 99.0% nucleotide

pairwise identity with the Cornell strain 195 isolate nitrogenase operon (DET_RS05950-

RS05990) (Figure 2A, Table S8). The KBTCE2 and KBTCE3 nitrogenase operons were almost

identical to each other, with one single nucleotide polymorphism (SNP) in the *nifK* gene (Figure

2A), which is the ▯-subunit of the iron-molybdenum protein in the nitrogenase complex

(Raymond *et al.* 2004).

To determine if KBTCE2 and KBTCE3 strains could utilize the *nif* operon to fix

nitrogen, we quantified the *nifD* gene copies and transcripts in the KB-1 VC △ cultures and the

KB-1/TCE-MeOH culture, referred to here as KB-1 TCE ▲. In the KB-1 VC △ cultures, the

absolute abundance of *nifD* transcripts was 2.3 (± 1.7) times more than the KB-1 TCE ▲ culture,

even though the absolute abundance of the *nifD* gene was 24 (± 9.6) times less (Figure 2B).

Therefore, *D. mccartyi* strains KBTCE2 and KBTCE3 in the KB-1 VC △ cultures were actively

transcribing the *nif* operon indicative of nitrogen fixation. However, the KBTCE2 and KBTCE3

strains are *tceA*-containing strains and do not contain *vcrA*, so these strains are not able to obtain

energy from VC in these experimental conditions. This is different from the KB-1 TCE ▲

culture in which strains KBTCE2 and KBTCE3 can obtain energy through the dechlorination of

TCE. Therefore, it is a conundrum as to how these strains obtain energy when grown in the

presence of VC as the sole electron acceptor in the KB-1 VC △ cultures, especially since

nitrogen fixation is an energetically expensive process (Leigh and Dodsworth 2007). This

finding might suggest syntrophic relationships between *D. mccartyi* strains, in the exchange of

energy for available nitrogen sources. Alternatively, this activity may point towards the function

of the constitutively expressed reductive dehalogenase (OG 15) observed under "starvation"

conditions in *D. mccartyi* strain 195 (DET_RS07915) (Johnson *et al.* 2008; Rahm and

Richardson 2008) and in KB-1 (DQ177510) (Waller *et al.* 2012; Liang *et al.* 2015). It may be

that RdhA from OG 15 can sustain the population with an unknown electron acceptor in what is

perceived as "starvation". Further work is required to determine if a reductive dehalogenase from

OG 15 is expressed in the KB-1 VC Δ cultures and its role in providing energy to *D. mccartyi*

strains.

The *nifD* transcripts-per-gene (TPG) ratio was calculated for the KB-1 VC Δ cultures and

for KB-1 TCE ▲ (Figure 2C). The TPG ratios of KB-1 cultures were comparable to approximate

TPG ratios of the *D. mccartyi* strain 195 isolate (Lee *et al.* 2009) and the PW4 groundwater

aquifer-derived enrichment culture dominated by Cornell strains (Kaya *et al.* 2019) (Figure 2C,

Table S6). Therefore, we identified two more strains of *D. mccartyi* that can fix nitrogen, both

originating from the KB-1/TCE-MeOH culture. *D. mccartyi* strains KBTCE2, KBTCE3, and 195

are the only strains currently known to fix nitrogen, all belonging to the Cornell clade (Molenda

et al. 2020). Therefore, the nitrogen fixation characteristic of *D. mccartyi* appears to be specific

to the Cornell clade.

***D. mccartyi* strain 195 TceA predicted VC-dechlorinating activity**

D. mccartyi strain 195 in the Donna II culture contains *tceA* and *D. mccartyi* strains KBTCE2

and KBTCE3 in the KB-1/TCE-MeOH also contain *tceA* (Molenda *et al.* 2020). The *tceA* gene

encodes the TCE reductive dehalogenase, TceA, which catalyzes the reductive dechlorination of

TCE to cDCE and cDCE to VC (Tang *et al.* 2013). TceA can also dechlorinate 1,2-

dichloroethane (1,2-DCA) to ethene and trace amounts of VC (Duhamel and Edwards 2007).

However, it was recently discovered that TceA can dechlorinate VC to ethene coupled to

organohalide respiration in the presence of sufficient vitamin B_{12} (Yan *et al.* 2021). Ethene

formation occurred when vitamin B_{12} concentrations were 10 µg/L or greater and dechlorination

rates were positively correlated to B_{12} concentrations (Yan *et al.* 2021). In the DKB experiment,

vitamin B_{12}, in the form of cyanocobalamin, was maintained at a concentration of 6 µg/L in the

culture medium, which would suggest that VC would not be consumed. However, our data

revealed slow VC dechlorination where ethene was formed at a rate of 0.88 ($\pm$ 0.58) μmol/day in

the DKB VC Δ1 culture (Figure S4) which was comparable to the rate of Cl$^-$ released when

vitamin B$_{12}$ concentrations were 10 μg/L (Yan *et al.* 2021). Furthermore, the observed increase

in *tceA* and *D. mccartyi* with the simultaneous decrease in *vcrA* (Figure 1D-F) suggested that *D.*

mccartyi were able to sustain growth on VC dechlorination with *tceA* with 6 μg/L of vitamin B$_{12}$

(Figure S4).

To determine if the active TceA was from the Donna II or KB-1 culture, we generated a

tceA clone library from the DKB VC Δ1 culture (Table S7) and the KB-1/TCE-MeOH culture as

a control. The cloned *tceA* sequence from the KB-1/TCE-MeOH culture was identical (100%

nucleotide pairwise identity) to the *tceA* in published KBTCE2 and KBTC3 genomes (Figure

S5A). The cloned *tceA* sequence from the DKB VC Δ1 culture was 100% identical to the *tceA* in

D. mccartyi strain 195 from the Donna II metagenome sequenced in year 2010 (Figure S5B), but

not the original sequence of isolated strain 195, sequenced in 2005. Therefore, the observed VC-

dechlorinating activity was not the result of a mutation during this experiment.　TceA from the

Donna II metagenome was aligned to TceA from the isolated *D. mccartyi* strain 195 and other

TceA from OG 5 (Figure S6). In this alignment, two non-synonymous mutations were identified

between the strain 195 TceA and Donna II TceA: V261A and I481T. Interestingly, the I481T

mutation was also observed in the recently sequenced TceA of strain FL2, which was shown to

have higher rates of VC respiratory activity compared to strain 195, even at high vitamin B$_{12}$

concentrations (Yan *et al.* 2021).

To explore whether these two mutations could have an impact on the substrate specificity

and activity of the protein, we generated models of the TceA structure using the sequence from

D. mccartyi strain 195 isolate and from the Donna II metagenome. Models were generated

independently through several prediction servers to increase confidence in the predicted

locations of each mutation (Table S9), with the highest scoring models produced by the Robetta

and AlphaFold2 servers (Song *et al.* 2013; Jumper *et al.* 2021). In each model the I481T

mutation was consistently near the active site in both the *D. mccartyi* strain 195 isolate and the

Donna II strain 195 TceA models (Figure 3B). Whereas the location of the V261A mutation had

some variability but was always predicted to be removed from the active site. It is unlikely that

the V261A mutation plays a role in altering the enzymatic activity (Figure 3A). Visualization of

the predicted I481T location led to two potential hypotheses that could contribute to enhanced

VC respiration: alteration of the substrate access channel, and hydrogen bonding between the

residue and an [4Fe-4S] cluster to shift its reduction potential. These are described below.

The crystal structure of PceA, a PCE reductive dehalogenase, from the organohalide-

respiring bacterium *Sulfurospirillum multivorans,* depicts a narrow substrate channel leading to

the buried hydrophobic binding pocket (Bommer et al., 2014). To visualize where the substrate

may enter the active site in our models, we predicted the structure all of the tunnels with access

to the catalytic face of the cobalamin cofactor using Caver 3.0 (Chovancova *et al.* 2012). The

models predicted that the I481T mutation is situated at the mouth of the major substrate access

channel in each model (Figure 3B). Access channels can have a drastic impact on the substrate

scope of an enzyme as they act as a filter for the active site. It has been established in directed

evolution research that mutations in the substrate channel can lead to a change in enzyme activity

(Kokkonen *et al.* 2019). Thus, it is speculated that the I481T mutation in TceA could alter

substrate preferences.

The I481T mutation is also situated in a position where it can form interactions with one

of the [4Fe-4S] clusters. While the level of influence that the [4Fe-4S] clusters and their redox

potential have on RDase activity is unstudied, it is well known that the environment surrounding

the clusters affects their redox potential (Langen *et al.* 1992; Birrell *et al.* 2016). The presence of

hydrogen-bonding residues has been suggested to have an impact on the redox potential though

there are other factors at play and a direct correlation with hydrogen bonding is not clear

(Stephens, Jollie and Warshel 1996). The I481T mutation introduces the potential for a hydrogen

bonding interaction and is predicted to be in close proximity (2.8 Å) of the [4Fe-4S] binding

position (Figure 3C). As the redox potential provides a driving force for respiration and

reduction of the electron acceptor, changes to the environment around the [4Fe-4S] clusters

should be considered as a possible mechanism of fine-tuning the RDase activity towards a

certain substrate.

Predictive modelling is a powerful tool in gaining insight as to the potential impact of

these mutations; however, biochemical characterization and mutagenesis experiments need to be

done to unequivocally determine the substrate range of the Donna II *D. mccartyi* strain 195

TceA. Furthermore, it would be beneficial to determine if mutations in TceA enhance VC

respiration when combined with vitamin B_{12} supplementation, similarly to strain FL2 (Yan *et al.*

2021), or if TceA mutants can completely respire VC with lower concentrations of vitamin B_{12}.

Implications for horizontal gene transfer of the *vcrA*-GI

One objective of this experiment was to try to promote HGT of the *vcrA*-GI from *D. mccartyi*

strain KBTCE1 into strain 195. As previously mentioned, HGT would be manifested by

high/equal abundance of both KB-1/*vcrA* and 195/*tceA* biomarkers with VC dechlorination

(Table 1). These criteria were not met in any DKB sub-cultures. To further confirm that the

vcrA-GI had not integrated into the *D. mccartyi* strain 195 genome, we PCR-amplified the

integration site at the *ssrA* gene locus (McMurdie *et al.* 2011). We did not observe *vcrA*-GI

integration in any DKB sub-cultures at the *ssrA* locus (Figure S7). As a result, we definitively

concluded that HGT of the *vcrA*-GI did not occur in under the experimental conditions of the

DKB experiment. In the family of KB-1 cultures enriched on different substrates, eight *D.*

mccartyi genomes were closed (Molenda *et al.* 2020). These genomes revealed that clade-

specific, and even strain-specific, characteristics provide *D. mccartyi* populations with ecological

advantages to persist within mixed communities and to prevent gene transfer of mobile genetic

elements between dissimilar strains. With the benefit of many more genome sequences, we now

have evidence that *D. mccartyi* clade speciation plays a role in strain compatibility for HGT. To

date, there are no Cornell strains that contain the *vcrA*-GI, only Pinellas and Victoria clades.

There are several other clade-specific characteristics, including nitrogen fixation, which is found

only in the Cornell clade, as previously described. HGT between clades may also be prevented

by cellular defense mechanisms such as CRISPR-Cas systems (Molenda *et al.* 2019, 2020).

With improved appreciation of the diversity of strains even within the KB-1/TCE-MeOH culture

itself, there is also no evidence of *vcrA*-GI HGT between the Pinellas and Cornell strains now

known to exist within this culture.

Starvation and long-term survival of *D. mccartyi* strains

These clade-specific and strain-specific characteristics may also play a role in the observed

persistence of *D. mccartyi* strains and their long-term survival under starvation conditions. As

previously mentioned, this finding may point toward RdhA of unknown function, such as the

starvation RdhA (OG 15) which has not been functionally characterized but commonly observed

in transcriptomic analyses (Johnson *et al.* 2008; Rahm and Richardson 2008; Waller *et al.* 2012;

Tang *et al.* 2013). Furthermore, the long-term survival of *D. mccartyi* strains may explain why at

least eight strains of *D. mccartyi* were identified in the KB-1 culture that has been maintained in

the laboratory for more than 20 years (Molenda *et al.* 2020). The observations of the DKB

experiment provide insights into the growth and survival of *D. mccartyi* in environments with

low flow rates or where cells are attached to a surface, such as in the natural environment, that

may also experience long-term starvation conditions. Organohalogens naturally exist in the

environment, typically at low concentrations, and were the natural substrates for organohalide

respiring bacteria before exposure to high concentrations of anthropogenic sources of

organohalogens (Field 2016). It is likely that *D. mccartyi* have always been capable of long-term

survival in the absence of organohalogens and these conditions have promoted strain variation

among *D. mccartyi* communities.

While this experiment started off as a simple attempt to promote HGT, the breadth and

depth of knowledge of *D. mccartyi* has developed immensely since the experiment began in

2012. In the time since the DKB culture was prepared at Cornell University, *D. mccartyi* was

renamed (Löffler *et al.* 2013), the number of closed genomes of *D. mccartyi* in NCBI quintupled

from 5 to 25 (Kube *et al.* 2005; Sung *et al.* 2006; Pöritz *et al.* 2013; Wang *et al.* 2014; Molenda,

Quaile and Edwards 2016; Molenda *et al.* 2020), prophages (Waller *et al.* 2012), mobile genetic

elements (Molenda *et al.* 2019), and CRISPR-Cas systems (Molenda *et al.* 2019) were identified

in KB-1, and the genomes in the KB-1/TCE-MeOH (Molenda *et al.* 2020) and Donna II cultures

(IMG-M taxon ID: 2032320001) were sequenced. Considering these advancements, the findings

of the DKB experiment contribute to this knowledge and will inform attempts to induce HGT in

the future.

Funding

This work was supported by the Government of Canada through Genome Canada and the

Ontario Genomics Institute (grant number 2009-OGI-ABC-1405 to E.A.E.), and the Natural

Sciences and Engineering Research Council (NSERC) of Canada (NSERC PGS-D to O.M. and

K.J.P. and Discovery grant funding to E.A.E). This work was also supported by the Government

of Ontario through the ORF-GL2 program and the Unites States Department of Defense through

the Strategic Environmental Research and Development Program (SERDP) under the contract

(grant number W912HQ-07-C-0036 project ER-1586).

Acknowledgements

We acknowledge Garrett Debs, a researcher in the Richardson lab at Cornell University, who

maintained the DKB in the early days, before it was sent to the University of Toronto.

Conflict of Interest

The authors have no conflicts of interest to declare.

References

Almagro Armenteros JJ, Tsirigos KD, Sønderby CK *et al.* SignalP 5.0 improves signal peptide

predictions using deep neural networks. *Nat Biotechnol* 2019;**37**:420–3.

Birrell JA, Laurich C, Reijerse EJ *et al.* Importance of Hydrogen Bonding in Fine Tuning the

[2Fe-2S] Cluster Redox Potential of HydC from Thermotoga maritima. *Biochemistry*

2016;**55**:4344–55.

Bommer M, Kunze C, Fesseler J *et al.* Structural basis for organohalide respiration. *Science (80-*

) 2014;**346**:455–8.

Chen VB, Arendall WB, Headd JJ *et al.* MolProbity: All-atom structure validation for

macromolecular crystallography. *Acta Crystallogr Sect D Biol Crystallogr* 2010;**66**:12–21.

Chovancova E, Pavelka A, Benes P *et al.* CAVER 3.0: A Tool for the Analysis of Transport

Pathways in Dynamic Protein Structures. *PLoS Comput Biol* 2012;**8**:23–30.

Doherty RE. A History of the Production and Use of Carbon Tetrachloride, Tetrachloroethylene,

Trichloroethylene and 1,1,1-Trichloroethane in the United States: Part 1 - Historical

Background; Carbon Tetrachloride and Tetrachloroethylene. *Environ Forensics* 2000;**1**:69–

81.

Duhamel M, Edwards EA. Microbial composition of chlorinated ethene-degrading cultures

dominated by Dehalococcoides. *FEMS Microbiol Ecol* 2006;**58**:538–49.

Duhamel M, Edwards EA. Growth and yields of dechlorinators, acetogens, and methanogens

during reductive dechlorination of chlorinated ethenes and dihaloelimination of 1,2-

dichloroethane. *Environ Sci Technol* 2007;**41**:2303–10.

Duhamel M, Mo K, Edwards EA. Characterization of a Highly Enriched Dehalococcoides -

Containing Culture That Grows on Vinyl Chloride and. *Appl Environ Microbiol*

2004;**70**:5538–45.

Duhamel M, Wehr SD, Yu L *et al.* Comparison of anaerobic dechlorinating enrichment cultures

maintained on tetrachloroethene, trichloroethene, cis-dichloroethene and vinyl chloride.

Water Res 2002;**36**:4193–202.

Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res 2004;**32**:1792–7.

Edwards EA, Grbic-Galic D. Anaerobic degradation of toluene and o-xylene by a methanogenic

consortium. *Appl Environ Microbiol* 1994;**60**:313–22.

Fennell DE, Gossett JM, Zinder SH. Comparison of butyric acid, ethanol, lactic acid, and

propionic acid as hydrogen donors for the reductive dechlorination of tetrachloroethene.

Environ Sci Technol 1997;**31**:918–26.

Field JA. Natural production of organohalide compounds in the environment. In: Loffler FE,

Adrian L (eds.). *Organohalide-Respiring Bacteria*. Berlin, Heidelberg: Springer Berlin

Heidelberg, 2016, 7–30.

Freedman DL, Gossett JM. Biological reductive dechlorination of tetrachloroethylene and

trichloroethylene to ethylene under methanogenic conditions. *Appl Environ Microbiol*

1989;**55**:2144–51.

Fung JM, Morris RM, Adrian L *et al.* Expression of reductive dehalogenase genes in

Dehalococcoides ethenogenes strain 195 growing on tetrachloroethene, trichloroethene, or

2,3-dichlorophenol. *Appl Environ Microbiol* 2007;**73**:4439–45.

Hendrickson ER, Payne JA, Young RM *et al.* Molecular Analysis of Dehalococcoides 16S

Ribosomal DNA from throughout North America and Europe. *Appl Environ Microbiol*

2002;**68**:485–95.

Hug LA, Maphosa F, Leys D *et al.* Overview of organohalide-respiring bacteria and a proposal

for a classification system for reductive dehalogenases. *Philos Trans R Soc B Biol Sci*

2013;**368**:20120322–20120322.

Johnson DR, Brodie EL, Hubbard AE *et al.* Temporal transcriptomic microarray analysis of

"Dehalococcoides ethenogenes" strain 195 during the transition into stationary phase. *Appl*

Environ Microbiol 2008;**74**:2864–72.

Jumper J, Evans R, Pritzel A *et al.* Highly accurate protein structure prediction with AlphaFold.

Nature 2021;**596**:583–9.

Kaya D, Kjellerup B V., Chourey K *et al.* Impact of Fixed Nitrogen Availability on

Dehalococcoides mccartyi Reductive Dechlorination Activity. *Environ Sci Technol*

2019;**53**:14548–58.

Kearse M, Moir R, Wilson A *et al.* Geneious Basic: An integrated and extendable desktop

software platform for the organization and analysis of sequence data. *Bioinformatics*

2012;**28**:1647–9.

Kelley LA, Mezulis S, Yates CM *et al.* The Phyre2 web portal for protein modeling, prediction

and analysis. *Nat Protoc* 2015;**10**:845–58.

Kokkonen P, Bednar D, Pinto G *et al.* Engineering enzyme access tunnels. *Biotechnol Adv*

2019;**37**:107386.

Kube M, Beck A, Zinder SH *et al.* Genome sequence of the chlorinated compound-respiring

bacterium Dehalococcoides species strain CBDB1. *Nat Biotechnol* 2005;**23**:1269–73.

Langen R, Jensen GM, Jacob U *et al.* Protein control of iron-sulfur cluster redox potentials. *J*

Biol Chem 1992;**267**:25625–7.

Lee PKH, He J, Zinder SH *et al.* Evidence for nitrogen fixation by "Dehalococcoides

ethenogenes" strain 195. *Appl Environ Microbiol* 2009;**75**:7551–5.

Leigh JA, Dodsworth JA. Nitrogen Regulation in Bacteria and Archaea. *Annu Rev Microbiol*

2007;**61**:349–77.

Liang X, Molenda O, Tang S *et al.* Identity and substrate specificity of reductive dehalogenases

expressed in Dehalococcoides-containing enrichment cultures maintained on different

chlorinated ethenes. *Appl Environ Microbiol* 2015;**81**:4626–33.

Löffler FE, Edwards EA. Harnessing microbial activities for environmental cleanup. *Curr Opin*

Biotechnol 2006;**17**:274–84.

Löffler FE, Yan J, Ritalahti KM *et al.* Dehalococcoides mccartyi gen. nov., sp. nov., obligately

organohalide-respiring anaerobic bacteria relevant to halogen cycling and bioremediation,

belong to a novel bacterial class, Dehalococcoidia classis nov., order Dehalococcoidales

ord. nov. and famil. *Int J Syst Evol Microbiol* 2013;**63**:625–35.

Major DW, Cox EE, Dworatzek SM *et al.* Field Demonstration of Successful Bioaugmentation

To Achieve Dechlorination of Tetrachloroethene To Ethene. *Environ Sci Technol*

2002;**36**:5106–16.

McMurdie PJ, Hug LA, Edwards EA *et al.* Site-specific mobilization of vinyl chloride

respiration islands by a mechanism common in Dehalococcoides. *BMC Genomics*

2011;**12**:287.

Molenda O, Puentes Jácome LA, Cao X *et al.* Insights into origins and function of the

unexplored majority of the reductive dehalogenase gene family as a result of genome

assembly and ortholog group classification. *Environ Sci Process Impacts* 2020;**22**:663–78.

Molenda O, Quaile AT, Edwards EA. Dehalogenimonas sp. Strain WBC-2 genome and

identification of its trans-dichloroethene reductive dehalogenase, TdrA. *Appl Environ*

Microbiol 2016;**82**:40–50.

Molenda O, Tang S, Lomheim L *et al.* Extrachromosomal circular elements targeted by

CRISPR-Cas in Dehalococcoides mccartyi are linked to mobilization of reductive

dehalogenase genes. *ISME J* 2019;**13**:24–38.

Moran MJ, Zogorski JS, Squillace PJ. Chlorinated solvents in groundwater of the United States.

Environ Sci Technol 2007;**41**:74–81.

Müller JA, Rosner BM, Von Abendroth G *et al.* Molecular identification of the catabolic vinyl

chloride reductase from Dehalococcoides sp. strain VS and its environmental distribution.

Appl Environ Microbiol 2004;**70**:4880–8.

Perez-de-Mora A, Lacourt A, McMaster ML *et al.* Chlorinated electron acceptor availability

selects for specific Dehalococcoides populations in dechlorinating enrichment cultures and

in groundwater. 2017:44.

Pöritz M, Goris T, Wubet T *et al.* Genome sequences of two dehalogenation specialists -

Dehalococcoides mccartyi strains BTF08 and DCMB5 enriched from the highly polluted

Bitterfeld region. *FEMS Microbiol Lett* 2013;**343**:101–4.

Rahm BG, Richardson RE. Dehalococcoides' gene transcripts as quantitative bioindicators of

tetrachloroethene, trichloroethene, and cis-1,2-dichloroethene dehalorespiration rates.

Environ Sci Technol 2008;**42**:5099–105.

Raymond J, Siefert JL, Staples CR *et al.* The Natural History of Nitrogen Fixation. *Mol Biol*

Evol 2004;**21**:541–54.

Regeard C, Maillard J, Dufraigne C *et al.* Indications for acquisition of reductive dehalogenase

genes through horizontal gene transfer by Dehalococcoides ethenogenes strain 195. *Appl*

Environ Microbiol 2005;**71**:2955–61.

Roy A, Kucukural A, Zhang Y. I-TASSER: A unified platform for automated protein structure

and function prediction. *Nat Protoc* 2010;**5**:725–38.

Smatlak CR, Gossett JM, Zinder SH. Comparative kinetics of hydrogen utilization for reductive

dechlorination of tetrachloroethene and methanogenesis in an anaerobic enrichment culture.

Environ Sci Technol 1996;**30**:2850–8.

Song Y, Dimaio F, Wang RYR *et al.* High-resolution comparative modeling with RosettaCM.

Structure 2013;**21**:1735–42.

Stephens PJ, Jollie DR, Warshel A. Protein control of redox potentials of iron-sulfur proteins.

Chem Rev 1996;**96**:2491–513.

Studer G, Rempfer C, Waterhouse AM *et al.* QMEANDisCo-distance constraints applied on

model quality estimation. *Bioinformatics* 2020;**36**:1765–71.

Sung Y, Ritalahti KM, Apkarian RP *et al.* Quantitative PCR confirms purity of strain GT, a

novel trichloroethene-to-ethene-respiring Dehalococcoides isolate. *Appl Environ Microbiol*

2006;**72**:1980–7.

Tang S, Chan WWMM, Fletcher KE *et al.* Functional characterization of reductive

dehalogenases by using blue native polyacrylamide gel electrophoresis. *Appl Environ*

Microbiol 2013;**79**:974–81.

The PyMOL Molecular Graphics System, Version 2.3.4 Schrödinger L. No Title.

Trott O, Olson AJ. AutoDock Vina: Improving the Speed and Accuracy of Docking with a New

Scoring Function, Efficient Optimization, and Multithreading. *J Comput Chem*

2009;**32**:174–82.

Untergasser A, Cutcutache I, Koressaar T *et al.* Primer3-new capabilities and interfaces. *Nucleic*

Acids Res 2012;**40**:1–12.

Waller AS, Hug LA, Mo K *et al.* Transcriptional analysis of a Dehalococcoides-containing

microbial consortium reveals prophage activation. *Appl Environ Microbiol* 2012;**78**:1178–

86.

Wang S, Chng KR, Wilm A *et al.* Genomic characterization of three unique Dehalococcoides

that respire on persistent polychlorinated biphenyls. *Proc Natl Acad Sci* 2014;**111**:12103–8.

Yan J, Wang J, Villalobos Solis MI *et al.* Respiratory Vinyl Chloride Reductive Dechlorination

to Ethene in TceA-Expressing Dehalococcoides mccartyi. *Environ Sci Technol*

2021;**55**:4831–41.

Figure 1: Cumulative electron acceptor consumed (meeq) and strain-specific biomarker tracking (gene copies/mL culture). (A)

Cumulative electron acceptor consumed, given that 8 eeq are required per mol of PCE consumed, and 2 eeq per mol VC consumed.

(B) *D. mccartyi* abundance, by 16S rRNA gene copies per mL of culture. (C) *vcrA* copies per mL of culture, quantifying both genomic

and extrachromosomal *vcrA*-GI. (D) *D. mccartyi* strain 195-specific biomarker for a "truncated" *rdhA* (t*rdhA*). (E) *tceA* and (F) *nifD*

gene copies per mL of culture. Closed triangles represent ammonium provided in the medium (▲), and open triangles represent

ammonium not provided (△) in growth medium. The quantification limit was 1×10^3 copies/mL; values below this are unreliable. Error

bars represent the standard deviation (n=3), otherwise unaveraged triplicates are numbered (1,2,3) or a single bottle if not numbered

(n=1).

Figure 2: Nitrogen fixation in *D. mccartyi* (*Dhc*) in KB-1. (A) Alignment of nitrogenase genes (*nif*) in *D. mccartyi* strains 195

(DET_RS05950-5990), KBTCE2 and KBTCE3, where each black bar represents a misaligned nucleotide position. The top scale

indicates nucleotide length in kilobase pairs (kb). (B) *nifD* transcripts and gene copies per mL of KB-1 grown with (▲) and without

(△) ammonium provided. Error bars represent standard deviation (n=3). (C) *nifD* transcripts per gene (TPG) in KB-1, compared to

approximate TPG ratios from Cornell-dominant *Dhc* cultures: pure strain 195 culture (Lee *et al.* 2009) and mixed PW4 culture (Kaya

et al. 2019), grown with (▲) and without (△) ammonium.

Figure 3: Protein models of the Donna II metagenome *D. mccartyi* strain 195 TceA (blue) and *D. mccartyi* strain 195 isolate TceA

(yellow) with mutation sites highlighted in red. (A) The Donna II TceA mutant structure shows spatial locations of V261A and I481T

mutations. (B) View of the substrate access channel (depicted in white mesh) with the I481T mutation (red) in the Donna II TceA and

strain 195 isolate TceA. (C) Close up of the two residues next to a superimposed [4Fe-4S] cluster with the measured distance (2.8 Å)

indicating the distance of the T481 hydroxyl group to the cluster. Models were produced using the Robetta and AlphaFold2 online

prediction servers, [4Fe-4S] clusters were superimposed from the PceA crystal structure (PDB ID: 4UR2) (Song *et al.* 2013).

Table 1: Key features of *D. mccartyi* strains combined to try to promote horizontal gene transfer.

	D. mccartyi strain		
Feature	KBTCE1[a]	195	Predicted "hybrid"
Role in HGT	Donor strain	Recipient strain	HGT product
Source Enrichment Culture	KB-1	Donna II	DKB
Clade	Pinellas	Cornell	Cornell
vcrA-GI	*vcrA*-GI	--	*vcrA*-GI
Nitrogen Fixation Operon	0	1	1
Distinguishing Biomarkers[b]	*vcrA*	t*rdhA*	*vcrA* & t*rdhA*
Common Biomarkers[c]	16S	16S, *tceA, nifD*	16S, *tceA, nifD*

[a]When the study was initiated, a *vcrA*-containing Pinellas strain was known to dominate the KB-

1 TCE-enrichment culture. Subsequently, genomes corresponding to three strains (KBTCE1,

KBTCE2, and KBTCE3) were closed from the KB-1/TCE-MeOH culture (Molenda *et al.* 2020).

The KBTCE1 strain was most abundant when the culture was grown on TCE or VC.

[b] Key quantitative (q)PCR biomarkers used to distinguish strains.

[c] Biomarkers common to more than one strain, where 16S represents the *D. mccartyi* 16S rRNA

gene.