Genome evolution in bacteria isolated from million-year-old subseafloor sediments

William D. Orsi[1,2*], Tobias Magritsch[1], Sergio Vargas[1], Ömer K. Coskun[1], Aurele Vuillemin[1], Sebastian Höhna[1,2], Gert Wörheide[1,2,3], Steven D'Hondt[4], B. Jesse Shapiro[5,6,7], Paul Carini[8*]

[1]Department of Earth and Environmental Sciences, Paleontology & Geobiology, Ludwig-Maximilians-Universität München, Richard-Wagner-Strasse 10, 80333 Munich, Germany.

[2]GeoBio-CenterLMU, Ludwig-Maximilians-Universität München, Richard-Wagner-Strasse 10, 80333 Munich, Germany.

[3]SNSB- Bayerische Staatssammlung für Paläontologie und Geologie, Richard-Wagner-Strasse 10, 80333 Munich, Germany.

[4]Graduate School of Oceanography, University of Rhode Island, 215 South Ferry Road, 02882 Narragansett, USA.

[5]Department of Biological Sciences, University of Montreal, QC, Canada

[6]Department of Microbiology and Immunology, McGill University, QC, Canada

[7]McGill Genome Centre, Canada

[8]Department of Environmental Science, the BIO5 Institute, School of Plant Sciences, and the School of Comparative Animal & Biomedical Science, University of Arizona, Tucson, Arizona USA

Corresponding authors*:
Prof. Dr. William D. Orsi
Ludwig-Maximilians-Universität München, Department of Earth and Environmental Sciences,
Paleontology & Geobiology, Richard-Wagner-Strasse 10, 80333 Munich, Germany.

Corresponding authors*:
Dr. Paul Carini
Address: University of Arizona, Department of Environmental Science, School of Plant Science, BIO5
Institute. Tucson, Arizona 85721
E-Mail: paulcarini@arizona.edu
Phone/Fax: Phone: 520-621-1646

Running title: Genome evolution in subseafloor bacteria

Keywords: genome evolution; deep biosphere

Significance statement: In microbial populations that subsist in isolation from the surface world

in deep subseafloor sediment over millions of years, ultra-slow metabolic rates caused by long

term energy limitation are hypothesized to restrict the spread of newly evolved traits. It remains

unknown whether genomic evolution occurs under these extreme conditions. Our findings

demonstrate that genomes of cultivated bacterial strains from the genus *Thalassospira* isolated

from million-year-old abyssal sediment exhibit greatly reduced levels of homologous

recombination, elevated numbers of pseudogenes, and widespread evidence of relaxed purifying

selection. Our findings show that the genome evolution of these anciently buried bacteria has

proceeded in a manner dominated by genetic drift, whereby in small population sizes, and in the

absence of homologous recombination, mutations became fixed into the population which has

led to the emergence of new genotypes.

Abstract: Deep below the seafloor, microbial life subsists in isolation from the surface world

under perpetual energy limitation. The extent to which subsurface microbes evolve and adapt to

their subseafloor habitat is unclear, given their ultra-slow metabolic rates. Here we show that

genomes of *Thalassospira* bacterial populations cultured from million-year-old subseafloor

sediments evolve by point mutation, with a relatively low rate of homologous recombination and

a high frequency of pseudogenes. Ratios of synonymous to non-synonymous mutation rates

correlate with the accumulation of pseudogenes, consistent with a dominant role for genetic drift

in the subseafloor genomes, but not in type strains of *Thalassospira* isolated from surface world

habitats. The genome evolution of these anciently buried bacteria has apparently proceeded in a

genetic drift-like manner, whereby under long-term isolation with reduced access to novel

genetic material from neighbors, new mutations became fixed into the populations leading to the

emergence of new genotypes.

Main text.

The subseafloor biosphere contains one-third of all bacterial cells on Earth totaling $>10^{29}$ cells,

which subsist over geological timescales under perpetual energy limitation (*1*). Whether evolution and

ecological differentiation occurs in microbial populations below the seafloor has remained controversial. It

is generally agreed that extreme energy limitation restricts metabolic activity and growth (*2*), which are

necessary for new mutations to propagate through populations to foster ecological differentiation and
speciation (*3*). A metagenomic analysis showed that energy limitation and reduced growth restricted the
spread of new mutations through microbial communities over 5,000 years in the upper 10 meters of anoxic
continental shelf sediments (*4*). Yet, energy-starved cells under experimental laboratory conditions may
undergo adaptations that confer fitness under long-term energy limitation (*5, 6*). Moreover, bacteria
surviving for up to half a million years in deep subsurface permafrost have been shown to actively repair
their DNA (*7*)—a process that can tamper the rate of molecular evolution in the subsurface (*4*). Here, we
used the genomes of bacteria isolated from million-year-old subseafloor abyssal clay sediments to
investigate the nature of genome evolution in subseafloor bacteria that persist under extreme energy
limitation over long timescales.

Newly acquired mutations of functional significance can sweep through relatively fast-growing
bacteria in surface world habitats and influence ecological differentiation (*8*), but it is unclear whether this
occurs in ancient subseafloor sediment given the comparably slow subseafloor bacterial biomass turnover
rates that are estimated to be on thousand-year timescales (*9*). The metabolic rates of microbes persisting
in deep-sea abyssal clay are amongst the lowest observed in the subseafloor biosphere, such that these
sediments are often oxic through the entire sediment column to the underlying oceanic crust (*10*). Microbes
inhabiting such abyssal clay subseafloor settings are characterized by ultra-slow respiration rates and live
near the low-energy limit to life (*11*). We cored a 15 m sedimentary sequence of oxygenated abyssal clay
at a water depth of 6,000 m in the North Atlantic where the average sedimentation rate is an estimated 1
meter per million years (*12*). A relatively slow drawdown of O_2 with increasing depth at this site (Fig. 1A)
reflects oxidation of organic matter by aerobic microbes.

We isolated colony-forming bacteria on petri dishes following an 18-month incubation of sediment
and sterile ^{18}O-labeled artificial seawater (Fig S1) from 3 and 6 meters below the seafloor (mbsf)(see
Methods). Because the mean sedimentation rate is estimated to be on average 1 m million yr^{-1}, the age of
the sediments from which these bacteria were enriched and isolated are estimated to be 3 and 6 million
years old, respectively. The full length 16S rRNA gene sequences from the isolates had closest similarity
(90-99% sequence identity) to *Thalassospira xiamenensis and Thalassospira lohafexi* previously isolated
from marine sediments (*13,14*) and oligotrophic marine waters (*15*) (these previously isolated microbes
and their related cultured relatives are referred to as 'type strains' herein).

Several lines of evidence indicate the *Thalassospira* isolated from the 3 and 6 mbsf sediment
enrichments are endemic to the subseafloor abyssal clays and are not a contaminant from the water column
or other sources. First, the V4 hypervariable region of the 16S rRNA gene sequences from the sediment
slurry enriched *Thalassospira* cultures share >99% sequence identity with a phylotype (OTU6) previously
identified from the *in-situ* community determined from the frozen samples (Figure 1C). This OTU became

^{18}O-labeled during the 18-month incubation (atomic ^{18}O labeling of DNA: 59%) in the presence of sterile

^{18}O-labeled seawater (Fig S1), a proxy for growing microbes (*16*), and had an estimated doubling time in

the incubation of 36 ± 1.5 (mean ± SD) days. This phylotype consists of two amplicon sequence variants

(ASVs) that cluster with the *Thalassospira* subseafloor isolates (Otu6_ASV1, Otu6_ASV2), respectively,

and are distinguished by a single nucleotide polymorphism (SNP) that is conserved between the ASVs and

the subseafloor isolates (Figure 1C). The *in-situ* concentrations of both ASVs have highest abundance (ca.

1,000 16S rRNA gene copies g^{-1} sediment) between 4 - 6 mbsf, and both *Thalassospira* ASVs were detected

in the 3 and 6 mbsf sediment (Figure 1B). This shows that the *Thalassospira* strains isolated from the 3 and

6 mbsf sediment enrichments are derived from the same distinct 16S rRNA gene ASVs present within the

in-situ communities. The long-term physical isolation of these isolates in the subseafloor (see 'sediment

physical properties' in SI) subsisting under uninterrupted energy limitation within these ancient sediments

provides an opportunity to investigate how the relative effects of recombination, nucleotide substitution,

and gene decay have shaped the genomes of the cultivated subsurface *Thalassospira* strains since their

burial in the deep-sea clay millions of years ago.

Genome statistics

We sequenced the genomes of ten *Thalassospira* isolates each from the 3 and 6 mbsf sediment enrichments

using a hybrid assembly approach consisting of long-read Nanopore sequencing, corrected and polished via

short-read Illumina technologies at >100x coverage. The mean genome completeness of these hybrid

assemblies was estimated to be 99.7% ± 0.3%; (mean ± SD), with most being 100% complete and

representing the complete chromosome (Table S1). The mean length of these new *Thalassospira* genomes

was 4.71 ± 0.08 Mbp (mean ± SD, Fig. S2), with 4,567 ± 107 (mean ± SD) protein coding genes (Fig. S2),

and were assembled to an average of 12 ± 2 (mean ± SD) contigs (Table S1). The genome size and number

of protein encoding genes are similar to those observed within the existing *Thalassospira* type strains

isolated from the surface world (Fig. S2).

Core genome phylogenomic analysis

The core genome phylogeny of existing *Thalassospira* type species, and the newly isolated

subseafloor *Thalassospira* strains, consisted of 1,809 orthologous genes and revealed three clades of

subseafloor *Thalassospira*. One clade shares 96-97% genome-wide average nucleotide identity (ANI),

and 99.9% 16SrRNA gene sequence identity, with *T. xiamenensis* and *T. permensis* (Fig. S3). We named

isolates in this clade *T. xiamenensis* strain 'Neogene', after the Neogene eon (2.8 – 23 mya) which covers

both estimated ages of sediment (3 mya and 6 mya) from which the strains in this clade were isolated. The

subseafloor genomes in this clade correspond to the 16S rRNA gene ASV1 detected in the *in situ* frozen

sediment core samples (Fig. 1B, C). A second clade contained three isolates from 6 mbsf sediment shared

97% ANI with *T. xiamenensis* and *T. permensis.* Since all isolates in this clade were recovered from ca. 6

mya sediment deposited during the Miocene eon (5.33 – 23 mya), we report them as *T. xiamenensis* strain

'Miocene' (Fig. S3). However, despite sharing 97% ANI in the core genome with *T. xiamenensis* and *T.*

permensis (Fig. S3), *T. xiamenensis* strain 'Miocene' only shared 90% 16S rRNA gene sequence identity

with these closest related type strains. The subseafloor genomes in this clade also correspond to ASV1

detected in the *in situ* frozen sediment core samples (Fig. 1B, C). A third clade contained subseafloor

Thalassospira cultures isolated only from 3 mbsf sediment and shared 95-96% ANI with *T. lucentensis*

and *T. lohafexi* (Fig. S3). The subseafloor genomes in this clade correspond to the 16S rRNA gene ASV2

detected in the *in situ* frozen sediment core samples (Fig. 1B, C). Based on the genetic distinctness of

these isolates we consider them to be a new candidate species, according to recently provided criteria

based on genome-wide ANI (*17*). Because all isolates of this third clade were recovered from 3 mya

sediment, we propose the candidate name '*Candidatus* Thalassospira pliocenensis', named after the

Pliocene age (2.58 – 5.33 mya) of the deep-sea clay sediments from which they were isolated.

Pangenome analysis revealed that flexible genome content is conserved within each of the three

subseafloor clades, further evidence that each clade represents a genetically distinct population (Fig. S4).

Roles of mutation and recombination

The ratio of nucleotide diversity originating from mutations versus homologous recombination (r/m) can

be used to measure the relative effect of homologous recombination on the genetic diversification of

populations, (*18*). Due to the physical isolation of individual bacterial cells, reduced cell concentrations,

and the reduced availability of extracellular DNA for recombination in subseafloor sediments (*2*) we

hypothesized that rates of homologous recombination in the subseafloor *Thalassospira* populations would

be lower compared to the type strains. To test this, we used an established method (*19*) to calculate the

relative rate of recombination to mutation (R/θ), the mean length of recombined DNA (δ), and the mean

divergence of imported DNA (v) for branch tips (existent genomes) and internal nodes (ancestral states) in

the *Thalassospira* core genome phylogeny, which allows for a calculation of r/m (r/m = (R/θ) * δ * v). This

analysis showed that in the *Thalassospira* core genome, the r/m is approximately ten times lower in the

existent subseafloor core genomes (r/m = 0.078) compared to the type strains (r/m = 0.71) (Table 1),

indicating that homologous recombination plays a much lesser role in the diversification of the subseafloor

strains. The r/m values of the subseafloor *Thalassospira* are furthermore anomalously low compared to

free-living bacteria isolated from the surface world, which have r/m values that range from 0.1-64

(*18*). Concomitant with the ten-fold lower r/m values compared to the type strains (Table 1), the subseafloor

Thalassospira core genomes exhibit far fewer numbers of inferred imported DNA from recombination

events compared to the *Thalassospira* type strains and the ancestral states of the last common ancestors

(Fig. 2).

We considered the possibility that subseafloor *Thalassospira* populations may be those with the

necessary traits to survive at the time of burial (*2*), and little (or no) genome evolution may occur thereafter

(*4*). We looked for evidence of evolution by investigating pairwise substitution numbers in the subseafloor

Thalassospira genomes. We identified 10's to 1000's of nucleotide differences (single nucleotide

polymorphisms [SNPs]) within each subseafloor *Thalassospira* clade, suggesting that evolution by point

mutation has occurred since burial (Fig. S5). The SNPs in subseafloor *Thalassospira* strains were present

in a clade-specific manner (Fig. S6) and included genes with predicted annotations involved in flagellar

motility (FlhB, FliO), transcription (TetR and Fis family transcriptional regulators), cell wall biogenesis

(peptidase S41, peptidoglycan DD-metalloendoptidase M23), and transport and metabolism of amino acids

and carbohydrates (Fig. S6). Alternatively, the nucleotide diversity we observed in the distinct subsurface

Thalassospira clades may have been present at the time of burial. However, the subseafloor *Thalassospira*

genomes have widely different nucleotide diversity when compared to the predicted ancestral states of the

last common ancestor with the type strains (Fig. 2). Thus, it is likely the clade-specific pairwise substitutions

accrued during the time below the seafloor.

We considered the possibility that these substitutions occurred within the subseafloor populations

during the culture enrichment process. However, the doubling times of the *Thalassospira* (OTU6) in the

incubation measured with qPCR (36 ± 1.5 days; mean $\pm$ SD) indicate an estimated maximum of 15

doublings over the incubation (see SI). Given the number of generations, and the mutation rate for bacterial

genomes (*20*) we calculated less than a single mutation would be expected to arise in the subseafloor

Thalassospira genomes over the 15 generations that occurred during 18 month enrichment (4.7 Mbp in the

Thalassospira genome $\times 1 \times 10^{-9}$ mutations bp^{-1} generation$^{-1} \times$ 15 generations = 0.07 mutations). This is

insufficient to explain the observed interpopulation nucleotide diversity between the subseafloor genomes

which is 952 ± 177 nucleotide differences Mbp^{-1} (Fig. S5). Thus, the inter-population nucleotide diversity

of the subseafloor strains likely arose during their long-term subsistence in the ancient sediments and is not

the result of evolution during the laboratory incubation.

Substitutions and pseudogenes are fixed in subseafloor populations

The greatly reduced role homologous recombination in the subseafloor *Thalassospira* genomes

(Table 1, Fig. 2) coincided with higher numbers of pseudogenes (non-functional parts of the genome that

resemble functional genes), and non-synonymous substitutions (substitutions that alter the amnio acid

sequence of a protein). We identified 47.9 ± 8.57 (mean $\pm$ SD) pseudogenes in the genomes of subseafloor

Thalassospira isolates, which is significantly higher than the number of pseudogenes identified in the type

strains (22.1 ± 5.52 pseudogenes [mean ± SD]; Table 1) (two-sided T-test: P=1.5E-10). Similarly, we

observed a modest but significant elevation of genome-wide nonsynonymous to synonymous substitution

rates (dN/dS) in the genomes of the subseafloor *Thalassospira* cultures (0.035 ± 0.006; mean ± SD), relative

to the type strains (0.022 ±0.012; mean ± SD) (two-sided t-test P=0.0002; Table 1 and Fig. 3). Like the

SNPs, the composition of pseudogenes occurs in a clade-specific manner (ANOSIM: P = 0.001) (Fig. S6-

S8). Compared to the type strains, the predicted annotations of the pseudogenes in the subseafloor

Thalassospira genomes are skewed towards those involved in transcription, energy conservation, amino

acid and carbohydrate metabolism, and flagellar motility (Fig. S7). Subseafloor genomes have significantly

higher numbers of pseudogenes involved in flagellar biosynthesis (FliN, FliK, FlhO) compared to the type

strains (Fig. S8).

In microbial genomes, non-essential genes under relaxed purifying selection are more susceptible

to mutation-driven decay into pseudogenes (*21*). In small populations, mildly deleterious nonsynonymous

substitutions and pseudogenes are fixed into the population by chance through genetic drift (*22*). Thus, key

genomic signatures associated with drift-related evolutionary processes are elevated dN/dS ratios and a

high proportion of pseudogenes (*23,24*). In the absence of recombination, the effects of nonsynonymous or

slightly deleterious mutations compound with each generation because descendants carry the mutational

burden of the parent generation (*23,24*).

The elevated and correlated genome-wide pseudogene count and dN/dS ratios (Fig. 3) in the

subsurface *Thalassospira* isolates indicates that upon burial the population size became restricted relative

to surface-world *Thalassospira*. Thus, the burial in sediments likely had two significant effects. First, the

reduced energy availability—in the form of reduced quality and quantity of reactive organic carbon (*25*)—

limited the environmental carrying capacity, and thus the population size, in these ancient subseafloor clays.

Second, because the population size was limited, the chance for recombination events was also reduced due

to infrequent cell-cell contact stemming from lower cell abundances and subseafloor *Thalassospira* cells

were less likely to encounter genetically diverse recombination partners that might introduce genetic

diversity into the population. The reduction of homologous recombination resulted in mutation becoming

the dominant driver of evolution in these subseafloor *Thalassospira* strains as highlighted by the order of

magnitude lower r/m values in the genomes of subseafloor *Thalassospira* (Table 1). Thus, our observations

in subsurface *Thalassospira* populations are consistent with relaxed purifying selection in the absence of

recombination in a small microbial population. Although we see elevated dN/dS ratios and an accumulation

of pseudogenes across the core genomes of subsurface *Thalassospira*, some functions appear to be more

prone to gene decay than others. For example, genes predicted to be involved in flagellar motility were

present in both the SNP (*fliO*, *flhB*, *flgH*; Fig. S5) and pseudogene (*fliN*, *fliK*, *flhO*; Fig. S8) analysis,

suggesting physical restriction and energetic limitation has rendered motility as a superfluous function in
the highly compacted ancient deep-sea clay.

Lack of genome reduction despite energy limitation

The concerted action of mutation and decay into nonfunctional pseudogenes can culminate in the
loss of chromosomal DNA through a proposed deletional bias in microbial genomes (*26*). We did not
observe evidence of genome reduction in the subseafloor *Thalassospira* genomes that might be expected in
small, recombination-limited bacterial populations (*21*). We speculate that ultra-slow generation times of
subsurface *Thalassospira* due to perpetual energy limitation has prevented sufficient generations to pass
for the cells to lose superfluous DNA or genome reduction is balanced by the influx of new DNA by lateral
gene transfer into the flexible genome (*27*). Alternatively, our selection of genomes and the inferences
derived may be biased by our analysis of genomes from culturable microbes and genome reduction is often
associated with complex growth requirements or unculturability (*28*).

Outlook:

Free-living bacterial populations may engage in high rates of homologous recombination (*8*), which
may allow them to differentiate ecologically and purge deleterious mutations. However, the nearly
complete lack of recombination in the subseafloor *Thalassospira* genomes analyzed here shows that they
have evolved under a different regime akin to endosymbiotic bacteria, which lack homologous
recombination and are thus subject to genetic drift whereby deleterious mutations become fixed (*21*). In the
absence of recombination, the so-called 'Muller's Ratchet' eventually leads to the extinction of
endosymbiotic bacterial lineages (*21*). Our genomes show similar signs of evolution, namely greatly
reduced recombination and elevated dN/dS ratios and pseudogene numbers (Table 1). However, dN/dS
ratios observed in the subseafloor *Thalassospira* genomes are lower than those seen in endosymbiotic
bacteria (*24*) and genome reduction was absent. Thus, it appears that the subseafloor genomes are in a
middle state, perhaps "one click" in the Mullers Ratchet, represented by a single burial event followed by
a stable but low population size, in contrast to the repeated population bottlenecks experienced by
endosymbionts at each insect generation.

Subseafloor communities are relatively small and become smaller and more physically isolated
with increasing depth reaching the ultra-low concentrations of one cell per cubic centimeter of sediment
at 2,500 meters below seafloor (*29*). Because genetic drift has a stronger effect on populations with small
population sizes (*21,24*), microbes in the deep biosphere with small population sizes experiencing
reduced homologous recombination may be particularly prone to genetic drift-mediated evolution. For
example, a metagenomic analysis showed that an anaerobic subseafloor population displayed elevated

dN/dS ratios *(4)*. Because the subseafloor biosphere contains one-third of all bacterial cells on Earth (*1*),

our findings suggest drift-like evolutionary processes in the absence of homologous recombination may

be much more widely distributed in nature than previously thought. Future assessments of homologous

recombination and drift in single cell genomes from uncultured lineages of bacteria and archaea that

comprise most subsurface energy limited communities (*30*) could be used to assess how widespread this

evolutionary mechanism is within the subsurface biosphere.

Acknowledgements. This work was supported primarily by the Deutsche Forschungsgemeinschaft (DFG)

project OR 417/1-1 granted to W.D.O.   Publication of the manuscript was supported by the LMU

Mentoring Program. The expedition was funded by the US National Science Foundation through grant

NSF-OCE-1433150 to S.D., who led the expedition.   This is Center for Dark Energy Biosphere

Investigations (C-DEBI) publication number XXX. A portion of this work was performed as part of the

LMU Masters Program "Geobiology and Paleobiology" (MGAP). P.C. was supported by the University

of Arizona's Technology and Research Initiative Fund (the Water, Environmental, and Energy Solutions

initiative).

Author contributions. W.D.O., P.C., J.S., and G.W. conceived the work and experimental approach.

W.D.O., T.M., O.K.C., S.V., P.C., S.H., and A.V. contributed to the laboratory and bioinformatics analyses

and experimental work. S.D. obtained the samples during the KN223 R/V Knorr oceanographic expedition.

All authors discussed and wrote the manuscript and commented on the paper.

Competing interests. The authors declare that they have no competing interests.

Supplemental information. Supplemental Table 1, Supplemental Figures S1-S8, References 31-49.

Data and materials availability. Data are publicly available through NCBI BioProject PRJNA473406.

The 16S data are available in SRA BioSample accessions SAMN10929403 to SAMN10929517. Figures

and output files from the pangenomic analysis in Anvio are available online through FigShare

(https://figshare.com/s/06ba1287a00ab01a1ee). Additional data related to this paper may be requested

from the authors.

References

1.   J. Kallmeyer, R. Pockalny, R. R. Adhikari, D. C. Smith, S. D'Hondt, Global distribution of
microbial abundance and biomass in subseafloor sediment. *Proc Natl Acad Sci U S A* **109**,
16213-16216 (2012).
2.   M. A. Lever *et al.*, Life under extreme energy limitation: a synthesis of laboratory- and
field-based investigations. *FEMS Microbiol Rev* **39**, 688-728 (2015).
3.   O. X. Cordero, M. F. Polz, Explaining microbial genomic diversity in light of evolutionary
ecology. *Nat Rev Microbiol* **12**, 263-273 (2014).
4.   P. Starnawski *et al.*, Microbial community assembly and evolution in subseafloor
sediment. *Proc Natl Acad Sci U S A* **114**, 2940-2945 (2017).
5.   S. E. Finkel, Long-term survival during stationary phase: evolution and the GASP
phenotype. *Nat Rev Microbiol* **4**, 113-120 (2006).
6.   L. M. Wick, H. Weilenmann, T. Egli, The apparent clock-like evolution of Escherichia coli
in glucose-limited chemostats is reproducible at large but not at small population sizes
and can be explained with Monod kinetics. *Microbiology (Reading)* **148**, 2889-2902
(2002).
7.   S. S. Johnson *et al.*, Ancient bacteria show evidence of DNA repair. *Proc Natl Acad Sci U S
A* **104**, 14401-14405 (2007).
8.   B. J. Shapiro *et al.*, Population genomics of early events in the ecological differentiation
of bacteria. *Science* **336**, 48-51 (2012).
9.   B. A. Lomstein, A. T. Langerhuus, S. D'Hondt, B. B. Jorgensen, A. J. Spivack, Endospore
abundance, microbial growth and necromass turnover in deep sub-seafloor sediment.
NAture **484**, 101-104 (2012).
10.   H. Roy *et al.*, Aerobic microbial respiration in 86-million-year-old deep-sea red clay.
Science **336**, 922-925 (2012).
11.   S. D'Hondt *et al.*, Presence of oxygen and aerobic communities from sea floor to
basement in deep-sea sediments. *Nature Geoscience* **8**, 299-304 (2015).
12.   A. Vuillemin *et al.*, Archaea dominate oxic subseafloor communities over multimillion-
year time scales. *Sci Adv* **5**, eaaw4108 (2019).
13.   M. Li, S. Yang, Q. Lai, Z. Shao, Draft Genome Sequence of Thalassospira xiamenensis
Strain MCCC 1A03042(T). *Genome Announc* **5**, (2017).
14.   C. Liu, Y. Wu, L. Li, Y. Ma, Z. Shao, Thalassospira xiamenensis sp. nov. and Thalassospira
profundimaris sp. nov. *Int J Syst Evol Microbiol* **57**, 316-320 (2007).
15.   A. Hutz, K. Schubert, J. Overmann, Thalassospira sp. isolated from the oligotrophic
eastern Mediterranean Sea exhibits chemotaxis toward inorganic phosphate during
starvation. *Appl Environ Microbiol* **77**, 4412-4421 (2011).
16.   B. A. Hungate *et al.*, Quantitative microbial ecology through stable isotope probing. *Appl
Environ Microbiol* **81**, 7570-7581 (2015).

17. L. M. Rodriguez-R, K. Konstantinidis, Bypassing cultivation to identify bacterial species.
Microbe **9**, 111-118 (2014).
18. M. Vos, X. Didelot, A comparison of homologous recombination rates in bacteria and
archaea*. ISME J* **3**, 199-208 (2009).
19. X. Didelot, D. J. Wilson, ClonalFrameML: efficient inference of recombination in whole
bacterial genomes. *PLoS Comput Biol* **11**, e1004041 (2015).
20. J. W. Drake, B. Charlesworth, D. Charlesworth, J. F. Crow, Rates of spontaneous
mutation. *Genetics* **148**, 1667-1686 (1998).
21. J. P. McCutcheon, N. A. Moran, Extreme genome reduction in symbiotic bacteria. *Nat*
Rev Microbiol **10**, 13-26 (2011).
22. H. Ochman, L. M. Davalos, The nature and dynamics of bacterial genomes. *Science* **311**,
1730-1733 (2006).
23. N. A. Moran, Accelerated evolution and Muller's rachet in endosymbiotic bacteria. *Proc*
Natl Acad Sci U S A **93**, 2873-2878 (1996).
24. C. H. Kuo, N. A. Moran, H. Ochman, The consequences of genetic drift for bacterial
genome complexity. *Genome Res* **19**, 1450-1454 (2009).
25. E. R. Estes *et al.*, Persistent organic matter in oxic subseafloor sediment. *Nature*
Geoscience **12**, 126-131 (2019).
26. A. Mira, H. Ochman, N. A. Moran, Deletional bias and the evolution of bacterial
genomes. *Trends Genet* **17**, 589-596 (2001).
27. Nelson-Sathi *et al.*, Origins of major archaeal clades correspond to gene acquisitions
from bacteria. *Nature* **517**, 77-80 (2015).
28. P. Carini, A "Cultural" Renaissance: Genomics Breathes New Life into an Old Craft.
mSystems **4**, (2019).
29. F. Inagaki *et al.*, Exploring deep microbial life in coal-bearing sediment down to ~2.5 km
below the ocean floor. *Science* **349**, 420-424 (2015).
30. L. Solden, K. Lloyd, K. Wrighton, The bright side of microbial dark matter: lessons
learned from the uncultivated majority. *Curr Opin Microbiol* **31**, 217-226 (2016).
375
376

Figure legends

Figure 1: Isolated subsurface *Thalassospira* are most abundant at 3-7 mbsf and distinct from related type strains isolated from overlying water and sediments. (A) Vertical profile of total 16S rRNA gene concentrations determined via qPCR (squares), and oxygen concentrations (circles). 16S rRNA gene concentration points are the average abundances of three technical qPCR replicates with ranges shown with error bars. (B) qPCR-normalized average concentrations of the *Thalassospira* affiliated 'Otu6' ASVs. Error bars are ranges from three technical qPCR replicates. Asterisks mark the depths for the long term ^{18}O-water incubation experiments, enrichments, and cultivation. (C) Maximum likelihood (PhyML) phylogenetic analysis of the *Thalassospira* 16S rRNA gene ASVs (V4 hypervariable region), together with subseafloor and type strain *Thalassospira* 16S rRNA gene V4 regions. The presence of the SNP is displayed. Black, grey, and white dots at the nodes represent >90%, >70%, >50% bootstrap support, respectively.

Figure 2: Recombination in the conserved core genome is limited in subseafloor *Thalassospira* populations. The maximum likelihood (PhyML) phylogenetic tree is based on a concatenated alignment of 1,809 genes conserved across all *Thalassospira* genomes ('core genes'). Black circles on nodes represent bootstrap values >95%. The position of recombination events in the core genome are represented by dark blue dots. Positions of low nucleotide diversity and no recombination events in the core genome are shown in light blue. Nucleotide diversity at specific sites in the core genome are illustrated with a color gradient (white: less diversity, orange: more diversity). Histograms on the right display the total number of recombination events (imports) in each genome sequence, and ancestral state reconstructions (internal nodes), as detected by ClonalFrameML (*19*).

Figure 3: The number of pseudogenes and dN/dS ratios are elevated and correlated in Subseafloor *Thalassospira* populations. Subseafloor genomes accumulate more pseudogenes as a function of increasing dN/dS ratios compared to the type strains. Linear regressions for type strains, subseafloor strains, and all strains are displayed.

Table 1. The contribution of recombination and mutation to nucleotide diversity in subseafloor populations. The results from ClonalFrameML (*19)* analysis used to calculate the relative contributions of recombination and mutation in the core genome (r/m). *R/θ*: the relative rate of recombination compared to mutation, *δ*: the average length of recombined (imported) DNA, *v*: mean divergence of imported DNA. Also displayed are the average number of pseudogenes and dN/dS ratios (+/- standard deviation). ** two sided T-test: P = 0.005. *****two sided T-test: P = 0.000001.

422 **Figure 1**

435 **Table 1**

Group	# strains	*R/θ*	*δ*	*v*	r/m	# pseudogenes	dN/dS
All subseafloor and type strains	34	0.053	244	0.055	0.71	37 (+/- 8)	0.025 (+/- 0.011)
Type strains	13	0.04	333	0.053	0.71	21 (+/- 3)	0.022 (+/- 0.01)
Subseafloor	21	0.006	500	0.026	0.078	48 (+/- 4)****	0.038 (+/- 0.007)**

441 **Figure 2**

Figure 3

Supplemental Information

Genome evolution in bacteria isolated from million-year-old subseafloor sediments

William D. Orsi[1,2*], Tobias Magritsch[1], Sergio Vargas[1], Ömer K. Coskun[1], Aurele Vuillemin[1], Sebastian Höhna[1,2], Gert Wörheide[1,2,3], Steven D'Hondt[4], B. Jesse Shapiro[5,6,7], Paul Carini[8*]

[1]Department of Earth and Environmental Sciences, Paleontology & Geobiology, Ludwig-Maximilians-Universität München, Richard-Wagner-Strasse 10, 80333 Munich, Germany.

[2]GeoBio-CenterLMU, Ludwig-Maximilians-Universität München, Richard-Wagner-Strasse 10, 80333 Munich, Germany.

[3]SNSB- Bayerische Staatssammlung für Paläontologie und Geologie, Richard-Wagner-Strasse 10, 80333 Munich, Germany.

[4]Graduate School of Oceanography, University of Rhode Island, 215 South Ferry Road, 02882 Narragansett, USA.

[5]Department of Biological Sciences, University of Montreal, QC, Canada

[6]Department of Microbiology and Immunology, McGill University, QC, Canada

[7]McGill Genome Centre, Canada

[8]Department of Environmental Science, the BIO5 Institute, School of Plant Sciences, and the School of Comparative Animal & Biomedical Science, University of Arizona, Tucson, Arizona USA

Group	isolation depth	isolate	N50 contig size (bp)	size (Mb)	# contigs	GC content	completeness (CheckM)	Contamination (CheckM)	predicted genome size (Mb)	# PEGs	Predicted #f PEGs	% core genome
		3.4	649,568	4.7	13	53.5	100%	0%	4.7	4463	4463	40.5
		3.7	1,392,694	4.73	13	53.5	100%	0%	4.73	4463	4463	40.5
		3.F	1,049,738	4.64	13	53.5	100%	0%	4.64	4463	4463	40.5
		3.G	1,232,970	4.64	13	53.5	100%	0%	4.64	4463	4463	40.5
Ca. T. pliocenensis'	3 mbsf	3.H	994,209	4.64	13	53.5	100%	0%	4.64	4463	4463	40.5
		3.I	735,219	4.64	11	53.5	100%	0%	4.64	4466	4466	40.5
		3.J	1,232,974	4.64	11	53.5	100%	0%	4.64	4466	4466	40.5
		3.3	556,527	4.64	13	53.5	100%	0%	4.64	4463	4463	40.5
		3.6	1,047,050	4.63	13	53.5	100%	0%	4.63	4463	4463	40.5
		6.1	823,074	4.74	10	54.8	99.50%	0%	4.76	4651	4674	38.9
		6.2	671,859	4.59	9	54.8	97%	0%	4.73	4481	4620	40.4
		6.6	1,045,808	4.74	12	54.8	100%	0%	4.74	4622	4622	39.1
T. xiamenensis 'Neogene'	6 mbsf	6.18	1,220,195	4.75	14	54.8	100%	0%	4.75	4651	4651	38.9
		6.F	1,372,017	4.74	9	54.8	99.50%	0%	4.76	4637	4660	39.0
		6.G	1,372,000	4.73	7	54.8	99.50%	0%	4.75	4636	4659	39.0
		6.I	1,156,923	4.75	11	54.8	100%	0%	4.75	4651	4651	38.9
		6.J	1,392,823	4.64	8	54.8	98%	0%	4.76	4555	4672	39.7
	3 mbsf	3.1	1,245,526	4.75	12	54.8	100%	0%	4.75	4673	4673	38.7
		6.5	484,976	4.87	16	54.7	100%	0%	4.87	4730	4730	38.2
T. xiamenensis 'Miocene'	6 mbsf	6.9	484,974	4.87	15	54.7	100%	0%	4.87	4728	4728	38.3
		6.H	484,664	4.87	16	54.6	100%	0%	4.87	4732	4732	38.2

490 **Table S1. Genome summary statistics.**

Figure S1. Top panel: ^{18}O-labeling of 16S rRNA genes from the *Thalassospira* OTU6 (see Figure 1), after 7 and 18 months of incubation with ^{18}O-labeled water from the 3 mbsf sediment (data originally published in Vuillemin et al., 2019). **Middle panel:** Oxygen consumption over time in the 18 month slurry from the 3 mbsf incubation (filled circles), and slurries containing labeled water and autoclaved sediment (killed control). **Bottom photo**: cultivation of colony forming bacteria on solid media after the 18 month incubation of sediment slurries in sterile ^{18}O-labeled artificial sea water. No bacterial colonies formed on petri dishes that were inoculated with the killed control slurries.

Figure S2: Summary of genome properties for *Thalassospira* strains used in this study.

Points are derived from the analysis of existing genome sequences (for "Type" strains), and new
high-quality draft genomes sequenced as part of this study.

Figure S3. Average nucleotide identity (ANI) and the core genome phylogeny. The tree is
based on maximum likelihood and a concatenated alignment of 1,809 core genes

Figure S4: Pangenome analysis of all *Thalassospira* genomes included in the study. The internal
dendrogram is a UPGMA based on the presence/absence of shared gene orthologs. Black bars in the first
(inner) 34 circles show the occurrence of gene clusters in the genome of *Thalassospira* species. Grey
areas and light colors in the circles represent gene clusters that were detected in the corresponding
genome. The next eight bars show statistics for the pangenome analysis of each individual gene cluster
(inner circle to outer circle) # contributing genomes: # of genomes that has a hit in a gene cluster, (GC),
max # paralogs, single copy gene clusters (SCG), Functional Homogeneity index, Geometric
homogeneity index, combined homogeneity index, presence of a COG functional assignment. The
categories on the right side (below dendrogram) show the totals per genome for # of gene clusters found
in each genome, Num genes per kbp: Number of genes per kilobase pairs of genome.   Redundancy:
Multiple occurrence of single copy genes in a genome, Completeness: Calculated from the occurrences of
single copy gene set in a genome

569   **Figure S5.** The number of SNPs between pairs of subseafloor *Thalassospira* genomes.

 Figure S6. The number of interpopulation SNPs at different positions in the core genome
 alignment, for each of the three subseafloor populations. The gene annotations to the
 corresponding regions are shown.

Figure S7: The heatmap shows the presence/absence of pseudogenes in the subseafloor genomes
and the presence of these pseudogenes in type strains. Functional annotation (against eggNOG)
of pseudogenes found in the subseafloor genomes only are compared to functional annotations of
pseudogenes found in both the subseafloor and type strains.

 Figure S8: Histograms showing the average relative abundance of functional categories in
 pseudogenes found within each of the three subseafloor populations, compared to the type
 strains. The error bars represent standard deviations, and asterisks indicate functional categories
 of pseudogenes that were significantly higher in the subseafloor genomes compared to the type
 strain genomes (two sided T-Test, P<0.05).

Materials and Methods

Sampling, pore water chemistry, sediment properties. All samples were taken during Expedition KN223 of the *R/V Knorr* in the North Atlantic, from 26 October to 3 December 2014. At site 11 (22°47.0′ N, 56°31.0′ W, water depth ~5600 m) via a long core piston-coring device (~28 m). Additional details of sampling are published elsewhere (*11, 13*). Dissolved oxygen concentrations in the core sections were measured with optical O_2 sensors from the equilibrated core sections and measured with needle-shaped optical O_2 sensors (optodes) (PreSens, Regensburg, Germany) as described previously (*11, 13*). The dissolved O_2 data from Expedition KN223 are archived and available online in the Integrated Earth Data Applications (IEDA) database (http://www.iedadata.org/doi?id=100519).

Deep sea clay sediment particles have a grain size of <0.2 μm, and thus the pore space between the clay particles is smaller than that of a bacterial cell, limiting the movement of bacteria through pore space in the clays. Bioturbation can vertically redistribute cells within marine sediments, but bioturbation is restricted to the upper 0.5 meters of sediment (*31*), and thus cannot vertically transport sediment surface material and microbes to depths of 3 and 6 mbsf. Barring the possibility of vertical redistribution via bioturbation, and considering the mean sedimentation rate of 1 m per million years, it can be concluded that the bacterial cultures obtained from sediments collected at 3 and 6 mbsf have been physically isolated from the surface world for millions of years.

DNA extraction, qPCR, 16S rRNA gene sequencing. DNA extractions, qPCR, and 16S rRNA gene sequencing were performed previously and described in Vuillemin et al (*12*). In brief, subcores were sampled aseptically with sterile syringes were subsampled aseptically in an ultraviolet (UV)–sterilized DNA/RNA clean HEPA- filtered laminar flow hood. DNA extraction was extracted from 10 g of sediment transferred into 50 ml of Lysing Matrix E tubes (MP Biomedicals) containing silica glass beads and homogenized for 40 s at 6 m/s using a FastPrep-24 5G homogenizer (MP Biomedicals) in the presence of 15 ml of preheated (65°C) sterile- filtered extraction buffer [76 volume % 1 M NaPO4 (pH 8), 15 vol- ume % 200 proof ethanol, 8 volume % MoBio's lysis buffer solution C1, and 1 volume % SDS]. The samples were incubated at 99°C for 2 min and frozen overnight at −20°C, thawed, and frozen again at −20°C overnight, followed by additional incubation at 99°C for 2 min and a second homogenization using the settings described above. After the second homogenization, the samples were centrifuged for 15 min, and the supernatants were concentrated to a volume of 100 ml using 50-kDa Amicon centrifugal filters (Millipore). Coextracted PCR-inhibiting humic acids and other compounds

were removed from the concentrated extract using the PowerClean Pro DNA Cleanup Kit (MoBio).

Extraction blanks were performed alongside the samples to assess laboratory contamination during the

extraction process.

DNA was quantified fluorometrically using a Qubit with a double- stranded DNA high-sensitivity

kit (Life Technologies). qPCR was performed using the custom primer dual indexed approach that tar-

gets the V4 hypervariable region of the 16S rRNA gene using updated 16S rRNA gene primers

515F/806R (515F, 5′-GTGYCAG- CMGCCGCGGTAA-3′; 806R, GGACTACNVGGGTWTCTAAT)

(*32*). Barcoded V4 hypervariable regions of amplified 16S rRNA genes were sequenced on an Illumina

MiniSeq following an established protocol (*33*). Bioinformatic processing of these previously published

sequence data is described by Vuillemin et al (*12*) in detail.

Long term incubation set up. Prior to setting up the incubations, the subcores were sampled with sterile

syringes using the sample aseptic technique used for the DNA extraction. For each sample depth, seven

grams of abyssal clay was placed into sterile 20-mL glass flasks and incubated with 4 mL of sterile

artificial seawater composed of either $H_2^{18}O$ (97% atomic enrichment) or unlabeled artificial seawater.

Vials were crimp sealed, with an oxygenated headspace of approximately 10 mL, and incubated at 8 °C.

The artificial seawater was different from the porewater at depth because there was no added nitrate, but

there was also no added ammonia which should be similar to the *in situ* conditions where ammonia is

generally below detection (*12*). Oxygen was measured continuously throughout the incubations using

non-invasive fiberoptic measurements as described previously (*12*). Small fluctuations in the oxygen

measurements in the killed control, and experimental incubations, were likely due to temperature

fluctuations of the incubator itself (±1˚C), since the non-invasive fiber optic oxygen sensor spots are

temperature sensitive (*12*). Oxygen consumption was detectable over 18 months in slurries consisting of

sediment and sterile artificial seawater (Fig S1), suggesting the presence of actively respiring microbes.

We used qSIP to measure the atom % ^{18}O-enrichment of actively growing microbial taxa

as described previously (*12*). In brief, after 7 and 18 months incubations DNA was extracted and

subjected to Cesium Chloride (CsCl) density gradient centrifugation. The same 16S 515F/806R

primers (described above) were used in qPCR (described above) to determine density shifts in

the peak DNA of buoyant density (BD) for each incubation. 16S rRNA gene amplicons from

each fraction resulting from the density gradient fractionation were Illumina sequenced as

described previously (*12*). To identify contaminants that may have entered during the

fractionation process, we also included in the sequencing run extraction blanks from the SIP

fractionation. OTUs containing sequences from extraction blanks were removed. Excess atm%

^{18}O-enrichment was calculated for each OTU (including OTU6, corresponding to the subseafloor

Thalassospira) according to the equations for quantifying per OTU atomic enrichment.

The number of doublings for the *Thalassospira* OTU (OTU_6) detected at the 18 month

timepoint was calculated using qPCR normalized relative abundance of the 16S rRNA genes at

T0 and 18 months. The number of doublings was divided by the total number of days incubated

to calculate doubling times in days.

Enrichments, cultivation, and sub-cultivation. After the 18 months of incubation in sterile ^{18}O-labeled

artificial seawater, 25 µL of slurry was plated onto solid media (10 mg/mL yeast extract and 8 mg/mL

agar in artificial seawater), and after 2 days incubated in the dark at room temperature, abundant colonies

were observed growing on the surface of the petri dishes (Fig S1). No colonies were observed to grow on

control petri dishes that received 25 µL of ^{18}O-labeled artificial seawater slurry incubated for 18 months

using starting material from autoclaved sediment (killed controls). This indicated that the colony forming

bacteria were from the sediments themselves and not due to contamination that was introduced during the

experimental set up of the incubations. We attempted to culture chemoheterotrophic microbes directly

from the collected sediment samples using the same conditions, but no colony forming units were

observed on the petri dishes even after several months of incubation. Thus, a long term incubation of the

sediments at 8 °C simply in the presence of added water apparently stimulated the activity of many

subseafloor bacteria to a point at which they were then able to grow on the surface of a petri dish.

Ten colonies were picked from petri dishes containing colonies from the 3 and 6 mbsf slurries.

These colonies were streaked onto new petri dishes, and a single colony was picked from these newly

streaked bacteria and grown in sterile liquid media (10 mg/mL yeast extract in artificial seawater). Single

colonies were then grown up in liquid media, a portion used for DNA extraction and genome sequencing,

and the remaining volume frozen as glycerol stocks.

Assessing the possibility for genome evolution during the 18 month enrichment.

Since bacteria can evolve on lab experimental timescales (*5,6*), we considered the possibility that

all diversification and evolution happened during the 1.5 year enrichment. Using the qPCR-based estimate

for doubling time of the subseafloor Thalassospira OTU (OTU_6) in the incubation which was 36 (± 1.5)

days, the number of doublings with this rate over this time period would be approximately 15. According
to "Drake's rule" (*20*), bacteria experience on average one mutation per 300 genomes replicated, and thus
the amount of nucleotide diversity (hundreds to thousands of mutations : Fig S5) that could be accumulated
during the incubation is insufficient to explain the observed diversity between the three subseafloor
populations. We thus conclude that the inter-population nucleotide diversity is the result of new mutations
that were acquired after they were buried.

Genome sequencing, de novo assembly, and annotation. DNA was extracted from the isolates grown in
liquid culture until the end of exponential phase as described above. After reaching stationary phase,
cultures were pelleted via centrifugation and the supernatant was decanted. The cell pellets were
resuspended in a preheated (65°C) sterile filtered extraction buffer [76 volume % 1 M NaPO4 (pH 8), 15
volume %200 proof ethanol, 8 volume %MoBio lysis buffer solution C1,and 1 volume % SDS], and added
to lysing matrix E tubes (MP Biomedicals) containing silica glass beads and homogenized for 40 s at 6 m/s
using a FastPrep-24 5G homogenizer (MPBiomedicals). The samples were centrifuged for 15 min, and the
dissolved high molecular weight DNA in the supernatant was concentrated to a volume of 100 μL using
50-kDa Amicon centrifugal filters (Millipore). The concentrated extract was cleaned of proteins and other
non-genomic DNA organic matter using the PowerClean Pro DNA Cleanup Kit (MoBio). Extraction blanks
were performed alongside the samples to assess laboratory contamination during the extraction process.
Genomic libraries were prepared using the Nextera XT DNA Library Prep Kit (Illumina). Quality control
and quantification of the libraries were obtained on an Agilent 2100 Bioanalyzer System using the High
Sensitivity DNA reagents and DNA chips (Agilent Genomics). Metagenomic libraries were diluted to 1
nM using the Select-a-Size DNA Clean and Concentrator MagBead Kit (Zymo Research) and pooled for
further sequencing on the Illumina MiniSeq platform. Genomic libraries were sequenced to a depth of ca.
100x coverage using a high-output paired end 2 x 150 sequencing regent kit (Illumina).
In addition to Illumina sequencing, the high molecular weight genomic DNA was sequenced using the
NanoPore MinION. Sequencing libraries for the MinION were prepared using the Ligation Sequencing kit
(Oxford NanoPore Technologies), according to the manufacturers instructions. Barcoded libraries were
sequenced on the MinION using a Flongle R9 flow cell, base-called and demultiplexed using the MinIT
(Oxford NanoPore Technologies).
A hybrid assembly was performed using both the short (Illumina) and long (NanoPore) read
sequencing data using Unicycler (v.0.4.0), which uses *de novo* assembled Illumina data from SPADES to
polish the *de novo* assembled contigs obtained from NanoPore data using RACON (*34*). The combined
assemblies of Illumina and NanoPore data resulted in a relative low number of contigs (9-12 per genome),

and a predicted genome completeness of 100% of nearly all genomes (Table S1). Genome completeness
was determined using CheckM (*35*). Genomes were annotated using RASTk (*36*).

Core genome phylogenetic analyses. The core genome was defined as the set of orthologous
genes which were shared in all subseafloor and extant *Thalassospira* genomes. Orthologous genes were
defined as those sharing >30% amino acid similarity to the collective suite of genes encoded within the
type strain *Thalassospira xiamenensis* M-5. *T. xiamenensis* M-5 was chosen as the reference genome for
this purpose, because it is the only publicly available genome of a cultivated *Thalassospira* isolate that is
completely closed and represents a single chromosome and a 190 Kb plasmid (*14*). A total of 1,809
orthologous genes were identified that are encoded by all *Thalassospira* strains that had >30% sequence
similarity to genes encoded within the *T. xiamenensis* M-5 genome. Each of these 1,809 genes were
individually aligned between all *Thalassospira* strains using MUSCLE (*37*), and the individual 1,809
alignments were then concatenated into a single core genome alignment for the subsequent phylogenomic
analysis (ClonalFrameML, HyPhy, aBRSEL) using Geneious Prime (version 2019.2.1). After
concatenation of all core genes, the total size of the core genome alignment was 1,817,073 nucleotide
characters, and 34 taxa (21 subseafloor strains, and 13 type strain taxa). A Maximum-Likelihood phylogeny
was created using PhyML (*38*) with a GTR model of evolution and 100 bootstrap replicates, which was
implemented within SeaView (*39*). The resulting phylogenetic tree and the concatenated core genome
alignment were used as inputs for subsequent ClonalFrameML and dN/dS analyses.
The contributions of mutations and recombination to the genomic diversity in the concatenated core
genome alignment, the number of recombination events (imports) per genome, and the positions of
recombination hot spots, was investigated using ClonalFrameML (*19*). Nucleotides unaffected by
recombination are referred to as unimported and nucleotides subject to recombination are referred to as
imported (*19*). ClonalFrameML provides the relative rate of recombination to mutation (R/Theta), the mean
length of recombined DNA (Delta), and the mean divergence of imported DNA (Nu). These results were
used to calculate the relative contribution of recombination versus mutation to the overall genomic diversity
(r/m), using the following formula r/m = (R/Theta) * Delta * Nu. ClonalFrameML was performed in three
separate runs containing a core genome alignment that contained (1) all genomes, (2) only the subseafloor
genomes, and (3) only the type strains. The resulting r/m values from these three groups (presented in Table
1) were then used to interpret the relative importance of mutations compared to recombination, in the
separate groups (e.g., type strains versus subseafloor strains). We acknowledge that because the dataset
contains genomes covering the diversity of a single bacterial genus (*Thalassospira*), the only detectable
recombination most events are from donors from the species under study, so that the main source of
recombination is not external (*19*).

In addition to calculating site and rates of recombination in the core genome, ClonalFrameML also

estimates the ancestral sequences at internal nodes of the clonal genealogy, and any missing base calls in

the observed sequences. The reconstruction of ancestral sequence states is performed using maximum

likelihood and the ClonalFrame model can be thought of as a hidden Markov model (HMM) when the

ancestral and descendant genomes for each branch of the clonal genealogy have been observed or

reconstructed (*19*). The hidden state of the HMM records whether each nucleotide was subject to

recombination or not on the branch connecting the two genomes. We acknowledge that drawing inference

under the resulting ancestral recombination graph is a notoriously complex statistical problem (*19*). Instead,

here we use ClonalFrameML only to assess within-group recombination (e.g., between species within the

genus *Thalassospira*), and thus our analysis cannot assess the influence of external recombination (from

species outside the genus *Thalassospira*).

The ratio of non-synonymous (dN) to synonymous (dS) mutations in the core genome alignment

(global ω ratio) was estimated using HyPhy v2.2.4 (*40*), and applying the adaptive branch-site random

effects likelihood (aBSREL) approach (*41*) to all branches in all subfamilies. Because of the high similarity

of the subseafloor genomes, aBSREL was run multiple times using the core genome alignment with only

one representative of the nearly identical subseafloor genomes included in each separate run. For this, one

representative genome of *T. xiamenensis* strain 'Neogene', *T xiamenensis* strain 'Miocene, and 'Ca. T.

pliocenensis' were included together with all other *Thalassospira* type strain genomes in each aBSREL run.

Then, the aBSREL run was repeated with the same type strains but different subseafloor genomes from

those same three clades, until dN/dS estimates for all subseafloor genomes were obtained.

Pangenome analysis. All subseafloor and extant *Thalassospira* genomes were analyzed in Anvi'o v6.2

using pangenome workflow (*42*). Briefly, each genome was converted into an anvi'o contigs database.

Genes were functionally annotated using eggnog v5.0 (*43*) with eggNOG-mapper (*44*) and imported back

to each genome's anvi'o contig database. Genome storages were generated using 'anvi-gen-genomes-

storages' and 'anvi-pan-genome' was deployed with parameters '--min-bit 0.5' (*45*), '--mcl-inflation' 10

(*46*), and the flag '--use-ncbi-blast' (*47*). The anvi'o pan database and summary of gene clusters stored in

FIGshare (https://figshare.com/s/06ba1287a00ab01a1ee).

Identifying pseudogenes. We estimated the number of pseudogenes within the genomes using two

programs, Psi-Phi (*48*) and DFAST (*49*). Psi-Phi uses a conservative criterion considering a pseudogene

only when it lost >20% of its original length, and enhances pseudogene recognition among closely related

strains both in annotated regions by identifying incorrectly annotated open reading frames (ORFs) and in intergenic regions by detecting new pseudogenes (*48*). Psi-Phi classifies pseudogenes as either identified pseudogenes and those as being possible, but potentially not pseudogenes. To be conservative, we only considered genes identified as pseudogenes from Psi-Phi and did not consider those flagged as 'potential pseudogenes'. As a second check of pseudogene content, we searched genomes for pseudogenes using DFAST (*49*). The estimated number of pseudogenes per genome was then taken as an average of the numbers detected both using both methods (Psi-Phi and DFAST). On average, Psi-Phi identified a higher number of pseudogenes per genome (57 ±10) compared to DFAST (32 ±4), but the variation between methods for the same genome was consistent (average variation =27, standard deviation of averages = 7). This minimal variation between individual genomes indicates that biases inherent to the pseudogene prediction methods affected the different genomes equally, and thus allow for a pseudogene comparison between the genomes.

Supplemental references

31. L. R. Teal, M. T. Bulling, E. R. Parker, M. Solan, Global patterns of bioturbation intensity and mixed depth of marine soft sediments. *Aquatic Biology* **2**, 207-218 (2008)

32. A. E. Parada, D. M. Needham, J. A. Fuhrman, Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ Microbiol* **18**, 1403-1414 (2016).

33. M. Pichler *et al.*, A 16S rRNA gene sequencing and analysis protocol for the Illumina MiniSeq platform. *Microbiologyopen*, e00611 (2018).

34. R. R. Wick, L. M. Judd, C. L. Gorrie, K. E. Holt, Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**, e1005595 (2017).

35. D. H. Parks, M. Imelfort, C. T. Skennerton, P. Hugenholtz, G. W. Tyson, CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* **25**, 1043-1055 (2015).

36. R. Overbeek *et al.*, The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* **42**, D206-214 (2014).

37. R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nulceic Acids Res* **32**, 1792-1797 (2004).

38. S. Guindon, F. Lethiec, P. Duroux, O. Gascuel, PHYML Online--a web server for fast maximum likelihood-based phylogenetic inference. *Nucleic Acids Res* **33**, W557-559 (2005).

39. M. Gouy, S. Guindon, O. Gascuel, SeaView version 4: A multiplatform graphical user
interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* **27**, 221-
224.
40. S. L. Pond, S. D. Frost, S. V. Muse, HyPhy: hypothesis testing using phylogenies.
Bioinformatics **21**, 676-679 (2005).
41. M. D. Smith *et al.*, Less is more: an adaptive branch-site random effects model for
efficient detection of episodic diversifying selection. *Mol Biol Evol* **32**, 1342-1353 (2015).
42. A. M. Eren *et al*., Anvi'o: an advanced analysis and visualization platform for 'omics data.
PeerJ **3**, e1319 (2015).
43. J. Huerta-Cepas *et al*., eggNOG 5.0: a hierarchical, functionally and phylogenetically
annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids
Res* **47**, D209-D314 (2018).
44.   J. Huerta-Cepas *et al*., Fast Genome-Wide Functional Annotation through Orthology
Assignment by eggNOG-Mapper. *Mol Biol Evol* **34**, 2115-2122 (2017).
45.   M. N. Benedict et al., ITEP: and integrated toolkit for exploration of microbial pan-
genomes. *BMC Genomics* **15**, 12.1186/1471-2164-15-8 (2014).
46. S. van Dongen, C. Abreu-Goodger, Using MCL to extract clusters from networks. In
"Bacterial Molecular Networks: Methods and Protocols" Eds. J. van Helden, A.
Toussaint, D. Thieffry. Springer New York, New York, NY. Pages 281-295.
47.   S. F. Altschul *et al.,* Basic local alignment search tool*. J Mol Biol* **215**, 403-410 (1990).
48. E. Lerat, H. Ochman, Psi-Phi: exploring the outer limits of bacterial pseudogenes.
Genome Res **14**, 2273-2278 (2004).
49. Y. Tanizawa, T. Fujisawa, Y. Nakamura, DFAST: a flexible prokaryotic genome annotation
pipeline for faster genome publication. *Bioinformatics* **34**, 1037-1039 (2018).
843

844