Invasive garlic mustard demonstrates stronger relationships with pathotrophic than

mycorrhizal fungi

Joseph D. Edwards[a], Wendy H. Yang[a,b,c], Anthony C. Yannarell[a,d]

[a]Program in Ecology, Evolution, and Conservation Biology, University of Illinois at Urbana-

Champaign, Urbana, IL 61801 USA

[b]Department of Plant Biology, University of Illinois at Urbana-Champaign, Urbana, IL

61801 USA

[c]Department of Geology, University of Illinois at Urbana-Champaign, Urbana, IL 61801 USA

[d]Department of Natural Resources and Environmental Science, University of Illinois at Urbana-

Champaign, Urbana, IL 61801 USA

Corresponding author: Joseph Edwards. Email: jedwa9@illinois.edu. Tel: 217-244-7128, Fax:

217-244-7246

Abstract

Garlic mustard (*Alliaria petiolata*) has long been known to degrade mycorrhizal

mutualisms in soils it invades and may also promote the abundance of microbial pathogens

harmful to native plants or alter saprotrophic communities to disrupt nutrient cycling. Phenology

of other invasive species, like *Lepidium latifolium* and *Lonicera maackii*, plays a role in their

interactions with soil microbial communities, and so we may expect garlic mustard phenology to

influence its effects on native soil microbiomes as well. Here, we investigate differences in

fungal, bacterial, and archaeal community structure, as well as the abundance of key functional

groups, between garlic mustard present, absent, and removed treatments in central-Illinois forest

soils across different stages of the garlic mustard life cycle. Across its phenology, garlic mustard

present soils had different overall fungal community structure and greater abundance of

pathotrophic fungi than soils where garlic mustard was absent or removed. However, abundance

of ectomycorrhizal and saprotrophic fungi as well as bacterial and archaeal community structure

were similar between treatments and did not interact with garlic mustard phenology. The most

abundant overall fungal taxon was a plant pathogen, *Entorrhiza aschersoniana*, that was greatest

in garlic mustard present soils, particularly while the plants were flowering. These results

support the hypothesis that invasive plants form active relationships with microbial pathogens

that could contribute to their overall success in invading ecosystems.

Keywords

Garlic mustard, forest soil, microbial-mediated plant invasion, fungal pathogen, mycorrhizal

fungi, invasive plant phenology

Introduction

Novel interactions between exotic plants and native microbes can both facilitate the

spread of invasive species and contribute to harmful legacies of invasion on native plant

communities (Callaway et al. 2004; van der Putten 2010; Van der Putten et al. 2007). Some

invasive plants can actively disrupt or degrade mutualistic interactions between soil microbes

and native plants that depend on them (Pringle et al. 2009; Traveset and Richardson 2014),

giving invaders a relative fitness advantage over native plants. Other invaders have been shown

to accumulate local pathogens in soils (Eppinga et al. 2006; Mangla et al. 2008). Because

invasive plants are often less susceptible to native pathogens than are native plants (Agrawal et

al. 2005; Engelkes et al. 2008; Klironomos 2002; Mitchell and Power 2003), this "enemy

release" can provide another relative fitness benefit to invasive plants over natives (Kardol et al.

2007; Keane and Crawley 2002). Invasive plant effects on soil microbial communities and their

functions can vary with invasion age (Lankau 2011), plant phenology (Portier et al. 2019), and

soil properties (Li et al. 2006). Therefore, it is important to investigate these plant-microbe

interactions in different environmental contexts to avoid biases in the documented patterns

(Hulme et al. 2013).

Garlic mustard (*Alliaria petiolata,* Cavara and Grande), a prominent invasive forest herb

in North America (Rodgers et al. 2008), is often considered a classic example of invasion

success driven by mutualism suppression (Callaway et al. 2008). Garlic mustard produces

secondary metabolites that reduce inoculum potential, activity, and biomass of mycorrhizal fungi

(Anderson et al. 1996; Roberts and Anderson 2001). Because garlic mustard is a non-

mycorrhizal plant, this assault on the mycorrhizal mutualists of its competitors has been posited

to explain garlic mustard's high invasion success (Lankau et al. 2009; Stinson et al. 2006; Wolfe

et al. 2008). Concomitant with a decrease in the relative abundance of mycorrhizal fungi in garlic

mustard invaded soils is an increase in the relative abundance of pathotrophic and saprotrophic

fungi (Anthony et al. 2017; Anthony et al. 2019). Both of these groups could also contribute to

garlic mustard's invasion success. Pathotrophic and saprotrophic fungi could fill niches made

available by the suppression of mycorrhizal fungi (Anthony et al. 2017), with pathotroph

accumulation potentially also benefitting garlic mustard as its growth rate is unaffected when

inoculated with heterospecific pathogens (Klironomos 2002). Changes in saprotrophic bacterial

and archaeal communities can alter soil nutrient cycling (Hawkes et al. 2007; Hawkes et al.

2005), potentially conferring advantage over native plants (Fraterrigo et al. 2011; Lee et al.

2018); however, garlic mustard effects on these communities are more poorly characterized than

its effect on fungal communities (Cipollini and Cipollini 2016). Bacterial richness and diversity

has been found to be consistent in soils under garlic mustard and other native forest herbs (Burke

and Chan 2010). However, garlic mustard's impact on bacterial richness and abundance is

thought fluctuate with invasion age, increasing initially then eventually recovering in older garlic

mustard populations (Lankau 2011). Although garlic mustard is well-known for its suppression

of mycorrhizal fungi, it is not clear if its impacts on other functional groups within soil microbial

communities also play a role in its invasion success.

Our understanding of garlic mustard impacts on soil microbial communities derives

largely from studies on persistent invasions with high plant densities, mostly located in the north-

eastern United States (Anthony et al. 2017; Anthony et al. 2019; Burke et al. 2019; Stinson et al.

2006; Wolfe et al. 2008) but also in the midwestern United States (Lankau 2011; Lankau et al.

2014). After initial invasion, garlic mustard population densities increase rapidly then become

relatively stable (Anderson et al. 1996; Nuzzo 1999). Although garlic mustard population

densities do not appear to decrease over time, its influence on microbial communities changes

with invasion age, as garlic mustard produces less allelochemicals over time (Lankau et al. 2009)

thereby weakening its influence on soil microbes (Lankau et al. 2010). Soil bacterial and fungal

communities also evolve as invasions age, leading to a recovery in the diversity of these

communities in older garlic mustard stands (Lankau et al. 2011). Furthermore, landscape factors

like disturbance regime, soil types, and surrounding plant communities also influence how

quickly garlic mustard invasions spread and their population distribution patterns in invaded

forests (Haines et al. 2018; Urbanowicz et al. 2019). Because garlic mustard may interact with

soil microbiomes differently depending on ecoregions and invasion patterns (Urbanowicz et al.

2019), it is important to broaden our understanding of these interactions beyond the historically

studied settings.

Garlic mustard studies often focus on plant-microbe interactions during the active life

stage of garlic mustard earlier in the growing season (e.g., Anthony et al. 2017; Burke et al.

2019). However, invasive plant effects on soil microbial communities (Arthur et al. 2012) and

microbial processes (Portier et al. 2019) can vary with plant phenology. As garlic mustard

exhibits an early phenology relative to native competitors (Engelhardt and Anderson 2011), its

effects on soil microbial communities may depend on its own life stages or broader temporal

changes associated with surrounding plants and ecosystem conditions (Bennett et al. 2013;

Saikkonen 2007). Effects like mutualist suppression that confer a competitive advantage may be

more important for garlic mustard during its early phenology, but garlic mustard could also

produce allelochemical and soil ecological legacy effects that continue to indirectly influence

soil microbial communities in the non-growing season. The long-term persistence of indirect

garlic mustard effects is supported by observations that soil microbial communities can take

several years to return to pre-invasion condition after garlic mustard eradication (Anthony et al.

2019; Lankau et al. 2014). To our knowledge, no studies have explicitly investigated how garlic

mustard impacts on soil microbial communities change over the course of its life cycle.

Here, we investigated how garlic mustard affects the relative abundance of soil microbial

functional groups (ectomycorrhizal fungi, saprotrophic fungi, pathotrophic fungi, bacteria, and

archaea) across garlic mustard phenology in relatively recently invaded hardwood forests in the

midwestern United States. We hypothesized that the degree of garlic mustard effects on soil

microbial community composition varies with its phenological stage, reflecting changes in the

production of allelochemicals that directly affect soil microbes. We compared soil microbial

community composition among three garlic mustard treatments (present, absent, and

experimentally removed) established in multiple forests across central Illinois where most garlic

mustard populations were estimated to be 20-30 years old at the time of the study. We expected

to observe a decrease in ectomycorrhizal fungi relative abundance, an increase in pathotrophic

and saprotrophic fungi relative abundance, and a change in bacterial and archaeal community

composition associated with the presence of garlic mustard throughout the growing season and

persisting after senescence, but with the magnitude of garlic mustard effects declining at

flowering and even more so after senescence.

Methods

Study Sites

In 2017, we measured garlic mustard impacts on soil microbial communities across five

forests in central Illinois: Brownfield Woods, Collins Pond Woods, Hart Woods, Richter Woods,

and Trelease Woods. These sites are located within a 30 km north-south range and a 40 km east-

west range. Geographical coordinates, climate, and soil variables are reported in Table S1. All

sites are managed by the Illinois Natural History Survey (INHS;

https://www.inhs.illinois.edu/research/natural-areas-uiuc). According to documentation from

INHS managers, all of these sites began experiencing invasion between 1991 and 1995. Site

managers attempted to control garlic mustard populations through hand-pulling, herbicide

application, and "weed whacking", and as a result, garlic mustard densities at these sites were

generally maintained below the high densities often associated with a decline in garlic mustard

allelochemical production (Lankau 2011; Lankau et al. 2009).

In 2018, eight forests in central Illinois were sampled to expand the range of our

inference to a greater number of geographical sites: Allerton Park, Breens Woods, Bunny Forest,

Funk Woods, Richter Woods, Trelease Woods, Warbler Ridge, and the Vermillion River

Observatory. Two of the sites, Richter Woods and Trelease Woods, were sampled in both years;

garlic mustard populations in the other three 2017 sites were intensively managed via herbicide

application in early 2018 so those sites were excluded. Site descriptions can be found in Table

S1. These sites are located within a 130 km north-south range and a 140 km east-west range. As

the 2018 study sites are managed by multiple different agencies, records for invasion ages and

management were not available for some of the sites depending on the agency. Plant

communities from forests sampled in 2017 and 2018 are relatively homogenous across these

sites, consisting of a "prairie grove" species mix with *Quercus, Fraxinus, Carya, & Acer* species

dominating the canopies. The understory these forests are made up of a diverse mix of tree

seedlings, *Asimina triloba,* and other shrubs including *Sanicula, Laportea, Ageratina, Ribes,* and

Trillium species

Study Design

In mid-March of each year of this study, we established 10-m^2 experimental blocks in each forest. Each block contained 1-m^2 plots for each of the following three treatments: garlic mustard present (present), garlic mustard absent (absent), and garlic mustard removed via hand-pulling (removed). We established the garlic mustard removed treatment as a control to account for meter-scale differences in soils within each block that might promote the growth of garlic mustard and to understand if live garlic mustard plants are required to see their effects. In observational studies such as these it is often difficult to differentiate the effects of invasive plants on soil conditions from preexisting edaphic characteristics that may have promoted garlic mustard colonization. We accounted for this challenge by comparing soils with active garlic mustard plant present and those that were previously invaded but live plants were removed. This allowed us to better attribute the presence of live garlic mustard plants to changes in microbial communities, rather than simply draw association between invaded and uninvaded soils. In 2017, we established 4 replicate blocks in each of the five forests and collected soil samples in May and August, corresponding with the rosette and senescent life stages of garlic mustard, respectively. In 2018, only a single treatment block was established in each forest because site was considered the experimental unit for replication. This allowed us to increase the number of sites (from 5 to 8) and sampling dates per site (from 2 to 3) compared to 2017. In 2018, we also added a soil collection in June, during garlic mustard flowering, to account for potential changes in garlic mustard allelochemical production with changes in reproductive status (Cipollini and Cipollini 2016) or light availability (Smith 2015).

Soil samples were collected from 0-10 cm depth from the soil surface using a 2 cm

diameter hand corer. There was very little, if any, development of organic horizons in any of our

plots, so all samples consisted primarily of mineral soils. Four soil cores were collected from

each 1 m^2 treatment plot, composited, and homogenized by hand in the field. On the same day as

collection, the soil samples were transported to the University of Illinois at Urbana-Champaign

and frozen at -20□C for later analysis.

DNA extraction, sequencing, identification, and assignments

We used high-throughput DNA sequencing to determine the impact of garlic mustard on

microbial community structure as well as the abundance of key functional groups and individual

taxa. DNA was extracted from 500 mg of freeze-dried soil using the FastDNA SPIN Kit for Soils

(MP Biomedicals, Santa Ana, USA). The extracts were purified using cetyl trimethyl ammonium

bromide (CTAB). DNA extracts were submitted to the Roy J. Carver Biotechnology Center at

the University of Illinois at Urbana-Champaign for Fluidigm amplification (Fluidigm, San

Francisco, USA) and Illumina sequencing (Illumina, San Diego, USA). We assessed fungal

communities via the *ITS2* gene region using ITS3 and ITS4 (Ihrmark et al. 2012) primers to

amplify DNA and sequence amplicons via MiSeq bulk 2 x 250 bp V2. As we were interested in

the broad fungal community effect of garlic mustard invasion, we chose to use a universal ITS2

barcode region (Schoch et al. 2012). However, this barcode is known to discriminate against

early-diverging fungal lineages like arbuscular mycorrhizal fungi (Stockinger et al. 2010), so this

group was not included in our analysis. We assessed bacterial communities via the bacterial and

archaeal *16S* rRNA genes. The 2017 samples were amplified using V4_515 forward and V4_806

reverse primers, and amplicons were sequences via MiSeq bulk 2 x 250 bp V2. The 2018

samples were amplified using Arch519 forward and BAC785 reverse primers, and amplicons

were sequenced via NovaSeq 2 x 250 bp. Primer sequences and references can be found in Table

S2. We changed the *16S* primers from 2017 to a more permissive primer in 2018 to capture a

wider range of the bacterial and archaeal community in our analysis. Because different primer

sets could lead to differences in amplification efficiency between different bacterial and archaeal

taxa, we do not directly compare 2017 and 2018 *16S* datasets in our analysis.

A standard bioinformatics workflow was used to process sequences. Briefly, FLASH

(Magoč and Salzberg 2011) was used for merging paired-end reads before quality filtering via

the FLASHX-Toolkit (Gordon and Hannon 2010) to remove sequences with quality scores of 30

or less for 10% of bases or more. USEARCH (Edgar 2010) was used to cluster sequences with

97% similarity into operational taxonomic units (OTUs) as well as remove chimeric sequences

and singletons. QIIME (Caporaso et al. 2010) was used to convert OTU clusters into an OTU

table, assign taxonomy using the UNITE reference database (version 8, 2019 release), and then

to rarify all samples to a uniform read depth of 7172 (ITS-2017), 2776 (ITS-2018), 13263 (16S-

2017), and 153507 (16S-2018). The final fungal OTU tables were analyzed via the FUN-Guild

(Nguyen et al. 2016) database to assign OTUs with probable or highly probable matches to

ectomycorrhizal (ECM), pathotroph, or saprotroph functional groups.

Statistical Analysis

All statistical analysis was performed in the R statistical environment (Team 2013).

Overall patterns of beta diversity and community structure were analyzed with the VEGAN

package (Oksanen et al. 2010). To determine differences community composition based on

invasion status (treatment), garlic mustard phenological stage (sampling date), and the

interaction between treatment and phenological stage, we performed a permutational analysis of

variance (PERMANOVA) with the *adonis* function on Hellinger-transformed rarified OTU

count numbers. We added site (forest) as a term in the *adonis* model, rather than stratifying, to

account for the variance contributed by site to microbial communities. This allowed us to keep

the model consistent between sampling years, as there was no replication within sites in our 2018

sampling year. We also assessed differences in community composition and heterogeneity using

nonmetric multidimensional scaling (NMDS) ordinations on Bray-Curtis dissimilarity of rarified

OTU count numbers. Differences in the relative abundance of functional groups (i.e., pathotroph,

saprotroph, and ECM) and individual OTUs were determined using a linear-mixed model with

the LMER package (De Boeck et al. 2011). In this model, we used invasion status, phenological

stage, and their interaction as fixed effects and site as a random effect. Relative abundances were

log-transformed to meet assumptions of normality and homogeneity of variance. Post-hoc

pairwise differences between treatments and between time points were determined using a Tukey

test. Statistical significance was assessed as $P < 0.05$.

Results

Soil microbial community composition

In both 2017 and 2018, overall fungal (and to a lesser extent bacterial and archaeal)

community composition differed based on garlic mustard treatment and phenology, with no

interactions between those effects (Table 1). Garlic mustard treatment accounted for 1.8%

(p=0.04; Table 1; Figure 1A) and 3% (p=0.002; Table 1; Figure 1B) of the variation in the fungal

overall community composition in 2017 and 2018, respectively. Garlic mustard treatment

accounted for a smaller portion of bacterial and archaeal community variation relative to fungi,

with 1.1% (p=0.78; Figure 2A) and 2.8% (p=0.05; Table 1; Figure 2B) in 2017 and 2018,

respectively. Phenology was associated with a greater portion of community variation than garlic

mustard treatment: 3.3% (p=0.001; Table 1; Figure 1C) and 4.6% (p=0.001; Table 1; Figure 1D)

for fungal communities in 2017 and 2018, respectively, and 6.8% (p=0.001; Table 1; Figure 2C)

and 4.2% (p=0.001; Table 1; Figure 2D) for bacterial and archaeal communities in 2017 and

2018, respectively. There was no significant interaction between garlic mustard treatment and

phenology on fungal or bacterial and archaeal community composition in either year (Table 1).

Functional group relative abundance

Garlic mustard invasion did not have a consistent effect on soil microbes across all

functional groups (Figure 3). Garlic mustard did not significantly affect the relative abundance of

ECM or saprotrophs in soil fungal communities (Table 2, Figure 3AB). ECM relative abundance

increased as the growing season progressed in both 2017 ($F_{1,110}$=29.88, p<0.001) and 2018

($F_{2,56}$=11.85, p<0.001; Table 2; Figure 3CD), with no interaction between phenology and garlic

mustard treatment (Table 2). In contrast, the relative abundance of saprotrophs was not

significantly affected by sampling date in either year (Table 2, Figure 3). The relative abundance

of pathotrophic fungi within the fungal community was significantly increased by garlic mustard

presence compared to the absent or removed treatments in both 2017 ($F_{2,110}$=6.12, p<0.001) and

2018 ($F_{2,56}$=7.02, p=0.002; Table 2; Figure 3AB). In 2017, plant phenology did not significantly

affect pathotroph relative abundance (Figure 3C), but in 2018, pathotroph relative abundance

was greatest while plants were flowering and decreased significantly after senescence

($F_{2,56}$=2.73, p=0.03; Table 2; Figure 3D). The average relative abundances of the four most

abundant OTUs in each functional group (including the 4 most abundant OTUs overall) are listed

in Table S3.

OTU abundance

The most abundant fungal OTU was *Entorrhiza aschersoniana*, a pathogen which

accounted for an average of 3.4% of fungal communities across all treatments and sampling

dates in 2017 (with a maximum of 50.1%) and an average of 6.7% (with a maximum of 72.2%)

across all samples in 2018. This OTU was found in 85% of garlic mustard present samples

(averaging 10% of the fungal community in this treatment) and was the most abundant taxon in

30% of these samples in 2017. In 2018, this OTU was found in 80% of garlic mustard present

samples (averaging 8% of the fungal community in this treatment) and was the most abundant

taxon in 41% of these samples. In contrast, this taxon was found in only 13% of garlic mustard

absent samples (averaging 0.0003% of the fungal community in this treatment) in 2017 and 16%

of garlic mustard absent samples (averaging 0.008% of the fungal community in this treatment)

in 2018. *E. aschersoniana* relative abundance was consistently greater in garlic mustard present

plots than in absent or removed plots across sampling dates (2017: $F_{2,110}=21.07$, $p<0.001$; 2018:

$F_{2,56}=16.528$, $p<0.001$; Table 2; Figure 4). In garlic mustard present and removed soils, *E.*

aschersoniana relative abundance decreased after garlic mustard present plots senesced (2017:

$F_{2,110}=4.10$, $p=0.01$; 2018: $F_{4,56}=6.08$, $p<0.001$; Table 2; Figure 4). In 2018, *E. aschersoniana*

abundance was greatest in garlic mustard present treatments when the plants were flowering

(Figure 4). Garlic mustard did not significantly affect any of the other most abundant OTUs

based on the same statistical test used for *E. aschersoniana*.

Discussion

Garlic mustard's suppression of mycorrhizal fungi have long been known (Roberts and

Anderson 2001; Vaughn and Berhow 1999) while evidence for its interactions with other

microbial functional groups in more recent (Anthony et al. 2017). Here, we found to garlic

mustard to promote pathotrophic fungi, especially during its flowering stage, without

significantly affecting other microbial groups like ECM, saprotrophic fungi, or bacteria and

archaea. Though plant phenological stage is important for microbial community interactions with

other invasive species (Arthur et al. 2012; Portier et al. 2019), garlic mustard's phenology did not

significantly influence ECM and saprotrophic abundance or community structure. However,

phenological stage did appear to play a role in the abundance of a prominent soil pathogen, *E.*

aschersoniana, which was greatest when garlic mustard plants were active. The evidence we

found of garlic mustard promoting pathotrophic fungi, namely a specific pathotrophic taxon

whose abundance peaked at the plant's flowering stage, supports the broader hypothesis that

microbial pathogens are important for understanding plant invasions.

Although invasive plants can acquire a competitive advantage over native species via

passive immunity to native pathogens, referred to as "enemy release", we found soils beneath

garlic mustard to actively accumulate the pathotrophic fungi *E. aschersoniana.* The recruitment

of *E. aschersoniana* resulted from direct influence of live garlic mustard, as the relative

abundance of E. *aschersoniana* declined over time in the garlic mustard removed treatment and

after garlic mustard senescence in the present treatment. Pathogen accumulation by invasive

plants is often associated with generalist pathogens (Eppinga et al. 2006; Mangla et al. 2008).

However, this pathogen recruited by garlic mustard specifically targets the roots of other

common forest understory plants (sedges and rushes; Bauer et al. 2015) that might compete with

garlic mustard. *E. aschersoniana* may not infect garlic mustard due to significant differences in

root architecture between garlic mustard and this taxon's preferred hosts (Fineran 1980). We did

not observe any root galls characteristic of *E. aschersoniana* infection on garlic mustard plants

we removed when establishing our treatments. Furthermore, we could find no evidence in the

literature of this fungus, or any other members of its phylum, infecting Brassicaceae roots.

Without performing reciprocal transplant studies to determine whether the fungi cultivated by

garlic mustard would negatively affect native plants (or garlic mustard itself), we do not know if

this relationship confers garlic mustard a competitive advantage. *E. aschersoniana* has not been

documented in garlic mustard invasions in the northeast U.S. where mutualism suppression has

been demonstrated to facilitate garlic mustard invasion. Given that garlic mustard recruitment of

E. aschersoniana occurred across all 11 independent forest sites in our study, where we did not

observe evidence of mutualism suppression, the potential for this pathogen to play a role in garlic

mustard invasion is worthy of exploration.

Garlic mustard's inhibition of mycorrhizal fungi in its invasive range is one of its most

widely observed traits (Anderson et al. 2010; Anthony et al. 2017; Callaway et al. 2008; Pringle

et al. 2009; Rodgers et al. 2008; Stinson et al. 2006; Wolfe et al. 2008), though we did not find

support for mycorrhizal inhibition in this study. Evidence for garlic mustard mycorrhizal

suppression often comes from potting studies using extracts from garlic mustard tissues

(Callaway et al. 2008; Stinson et al. 2006). Though mycorrhizal suppression has been

demonstrated under field conditions (Anthony et al. 2017), it is difficult to quantify ecologically

relevant allelochemical concentrations from garlic mustard plants *in situ*, particularly as these

concentrations fluctuate with population age (Lankau et al. 2009). Further, many of the soils in

our study had higher clay contents than the characteristically sandy soil of more northern and

eastern North American forests invaded by garlic mustard (Table S1). Allelochemicals produced

by garlic mustard plants could have bound to these clay particles (Blair et al. 2005), potentially

ameliorating their effects on mycorrhizal fungi. As garlic mustard has been found to have

diminished effects on intact, rather than establishing, mycorrhizal networks (Barto et al. 2010),

this could help explain why we observed shifts in fungal community structure and pathotrophic

relative abundance without reductions in ECM abundance. The increases in pathotroph relative

abundance in garlic mustard invaded soils without evidence for mycorrhizal inhibition observed

here suggest a relationship between garlic mustard and pathotrophic fungi independent of

mycorrhizal allelopathy. Pathotroph accumulation can occur in the absence of mutualism

suppression for garlic mustard, but its role in promoting garlic mustard invasion will require

further investigation.

Our results demonstrate that plant-microbe interactions thought to facilitate garlic

mustard invasion, like mutualism suppression or enemy release, may vary with invasion age and

geography. Many studies investigating microbial community impacts of garlic mustard invasion

are conducted on long-term, high-density populations (Anthony et al. 2017; Anthony et al. 2019;

Burke et al. 2019; Stinson et al. 2006; Wolfe et al. 2008), demonstrating the eventual effects of

garlic mustard invasion. However, based on these studies it is difficult to predict microbial

community responses to more recent garlic mustard invasions (Corbin et al. 2017). Microbial

community suppression by garlic mustard peaks when invasive populations are around 40 years

old (Lankau 2011), which may represent a potential ecological filter on microbial communities

in garlic mustard invaded soils. Most garlic mustard populations in our study were younger than

40 years with low garlic mustard plant density characteristic of this age range, thus, the soils

surrounding these populations have not likely experienced garlic mustard's peak microbial

suppression capacity. Filtering of microbial communities in garlic mustard invaded soils caused

by peaks in allelopathy based on invasion age could explain differences in our results,

particularly mycorrhizal suppression, from studies conducted on older, higher density garlic

mustard populations. However, without direct comparisons we cannot definitively conclude

whether these differences are due to invasion age or other regional characterizes. Our results

indicate that garlic mustard invasions can occur without strong evidence for mycorrhizal

inhibition, challenging the paradigm that mutualism suppression is a critical driver of garlic

mustard invasiveness. It is possible that other microbially-mediated mechanisms, like pathogen

accumulation, play a larger role in garlic mustard invasion in some environmental contexts.

Conclusions

Garlic mustard is often cited as a key example of the mutualism degradation hypothesis

for plant invasion (Pringle et al. 2009; van der Putten 2010). However, we were unable to detect

this phenomenon for garlic mustard invasions in Illinois forests with relatively young invasions.

Instead, we demonstrated that garlic mustard actively promoted the relative abundance of fungal

pathotrophs, particularly *E. aschersoniana*, which could confer a competitive advantage over

native plants. While we did not observe garlic mustard phenological stage to influence its effect

on the abundance and structure of microbial functional groups, *E. aschersoniana* relative

abundance was affected by phenology. This observation together with a rapid decline in *E.*

aschersoniana relative abundance after garlic mustard removal suggests that live garlic mustard

directly recruits this pathogen. Overall, our study shows that in some contexts, either associated

with invasion age or regional environmental differences, garlic mustard interacts more strongly

with pathotrophic fungi than mycorrhizal fungi.

Acknowledgements

We would like to thank the Illinois Natural History Survey and Grand Prairie Friends,

particularly Jamie Ellis and Jeff Peyton, along with The Parklands Foundation and Dr. Rodger

Anderson for allowing access to their lands for conduction this research and their gracious help

in finding garlic mustard populations. We greatly appreciate the assistance from Ally Cook, Alex

Kent, Elle Lucadamo, Nikki Snyder, Xiangyu Zhang, Joseph Schlosser, Alonso Favela, and

Gabe Price for their help in establishing treatments and sample collection. This work was

supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture,

McIntire Stennis program under accession number 1011139 to ACY and a National Science

Foundation Dimensions of Biodiversity grant DEB 1831582 to WHY. JDE was also supported

by the cooperative State Research, Education, and Extension Service, US Department of

Agriculture, under project number ILLU 875-952. We are also grateful to Adam Davis and

James Dalling for comments of earlier versions of this manuscript.

References

Agrawal AA, Kotanen PM, Mitchell CE, Power AG, Godsoe W, Klironomos J (2005) Enemy
release? An experiment with congeneric plant pairs and diverse above and belowground
enemies Ecology 86:2979-2989

Anderson RC, Anderson MR, Bauer JT, Slater M, Herold J, Baumhardt P, Borowicz V (2010)
Effect of removal of garlic mustard (*Alliaria petiolata*, Brassicaeae) on arbuscular
mycorrhizal fungi inoculum potential in forest soils The Open Ecology Journal 3

Anderson RC, Dhillion SS, Kelley TM (1996) Aspects of the ecology of an invasive plant, garlic
mustard (*Alliaria petiolata*), in central Illinois Restoration Ecology 4:181-191 doi:DOI
10.1111/j.1526-100X.1996.tb00118.x

Anthony M, Frey S, Stinson K (2017) Fungal community homogenization, shift in dominant
trophic guild, and appearance of novel taxa with biotic invasion Ecosphere 8:e01951

Anthony M, Stinson K, Trautwig A, Coates-Connor E, Frey S (2019) Fungal communities do not
recover after removing invasive *Alliaria petiolata* (garlic mustard) Biological Invasions
21:3085-3099

Arthur MA, Bray SR, Kuchle CR, McEwan RW (2012) The influence of the invasive shrub,
Lonicera maackii, on leaf decomposition and microbial community dynamics Plant
Ecology 213:1571-1582

Barto K, Friese C, Cipollini D (2010) Arbuscular mycorrhizal fungi protect a native plant from
allelopathic effects of an invader Journal of chemical ecology 36:351-360

Bauer R, Garnica S, Oberwinkler F, Riess K, Weiß M, Begerow D (2015) Entorrhizomycota: a
new fungal phylum reveals new perspectives on the evolution of fungi PLoS One
10:e0128183

Bennett AE et al. (2013) Arbuscular mycorrhizal fungal networks vary throughout the growing
season and between successional stages PloS one 8:e83241

Blair AC, Hanson BD, Brunk GR, Marrs RA, Westra P, Nissen SJ, Hufbauer RA (2005) New
techniques and findings in the study of a candidate allelochemical implicated in invasion
success Ecology Letters 8:1039-1047

Burke DJ, Carrino-Kyker SR, Hoke A, Cassidy S, Bialic-Murphy L, Kalisz S (2019) Deer and
invasive plant removal alters mycorrhizal fungal communities and soil chemistry:
Evidence from a long-term field experiment Soil Biology and Biochemistry 128:13-21

Burke DJ, Chan CR (2010) Effects of the invasive plant garlic mustard (*Alliaria petiolata*) on
bacterial communities in a northern hardwood forest soil Canadian journal of
microbiology 56:81-86

Callaway RM et al. (2008) Novel weapons: Invasive plant suppresses fungal mutualists in
America but not in its native Europe Ecology 89:1043-1055 doi:Doi 10.1890/07-0370.1

Callaway RM, Thelen GC, Rodriguez A, Holben WE (2004) Soil biota and exotic plant invasion
Nature 427:731-733

Caporaso JG et al. (2010) QIIME allows analysis of high-throughput community sequencing
data Nature methods 7:335

Cipollini D, Cipollini K (2016) A review of garlic mustard (*Alliaria petiolata*, Brassicaceae) as
an allelopathic plant The Journal of the Torrey Botanical Society 143:339-348

Corbin JD, Wolford M, Zimmerman CL, Quirion B (2017) Assessing feasibility in invasive plant
management: a retrospective analysis of garlic mustard (*Alliaria petiolata*) control
Restoration Ecology 25:S170-S177

De Boeck P, Bakker M, Zwitser R, Nivard M, Hofman A, Tuerlinckx F, Partchev I (2011) The estimation of item response models with the lmer function from the lme4 package in R Journal of Statistical Software 39:1-28

Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST Bioinformatics 26:2460-2461

Engelhardt MJ, Anderson RC (2011) Phenological niche separation from native species increases reproductive success of an invasive species: *Alliaria petiolata* (Brassicaceae)–garlic mustard1 The Journal of the Torrey Botanical Society 138:418-433

Engelkes T et al. (2008) Successful range-expanding plants experience less above-ground and below-ground enemy impact Nature 456:946-948

Eppinga MB, Rietkerk M, Dekker SC, De Ruiter PC, Van der Putten WH, Van der Putten WH (2006) Accumulation of local pathogens: a new hypothesis to explain exotic plant invasions Oikos 114:168-176

Fineran JM (1980) The structure of galls induced by Entorrhiza C. Weber (Ustilaginales) on roots of the Cyperaceae and Juncaceae

Fraterrigo JM, Strickland MS, Keiser AD, Bradford MA (2011) Nitrogen uptake and preference in a forest understory following invasion by an exotic grass Oecologia 167:781

Gordon A, Hannon G (2010) Fastx-toolkit FASTQ/A short-reads preprocessing tools (unpublished) http://hannonlab cshl edu/fastx_toolkit 5

Haines DF, Aylward JA, Frey SD, Stinson KA (2018) Regional patterns of floristic diversity and composition in forests invaded by garlic mustard (*Alliaria petiolata*) Northeastern Naturalist 25:399-417

Hawkes CV, DeAngelis KM, Firestone MK (2007) Root interactions with soil microbial communities and processes. In: The Rhizosphere. Elsevier, pp 1-29

Hawkes CV, Wren IF, Herman DJ, Firestone MK (2005) Plant invasion alters nitrogen cycling by modifying the soil nitrifying community Ecology letters 8:976-985

Hulme PE, Pyšek P, Jarošík V, Pergl J, Schaffner U, Vila M (2013) Bias and error in understanding plant invasion impacts Trends in ecology & evolution 28:212-218

Ihrmark K et al. (2012) New primers to amplify the fungal ITS2 region–evaluation by 454-sequencing of artificial and natural communities FEMS microbiology ecology 82:666-677

Kardol P, Cornips NJ, van Kempen MM, Bakx-Schotman JT, van der Putten WH (2007) Microbe⬜mediated plant–soil feedback causes historical contingency effects in plant community assembly Ecological monographs 77:147-162

Keane RM, Crawley MJ (2002) Exotic plant invasions and the enemy release hypothesis Trends in ecology & evolution 17:164-170

Klironomos JN (2002) Feedback with soil biota contributes to plant rarity and invasiveness in communities Nature 417:67-70

Lankau RA (2011) Resistance and recovery of soil microbial communities in the face of *Alliaria petiolata* invasions New Phytologist 189:536-548 doi:10.1111/j.1469-8137.2010.03481.x

Lankau RA, Bauer JT, Anderson MR, Anderson RC (2014) Long-term legacies and partial recovery of mycorrhizal communities after invasive plant removal Biological invasions 16:1979-1990

Lankau RA, Nuzzo V, Spyreas G, Davis AS (2009) Evolutionary limits ameliorate the negative impact of an invasive plant Proceedings of the National Academy of Sciences 106:15362-15367

Lee MR, Flory SL, Phillips RP, Wright JP (2018) Site conditions are more important than
abundance for explaining plant invasion impacts on soil nitrogen cycling Ecosphere
9:e02454
Li W-h, Zhang C-b, Jiang H-b, Xin G-r, Yang Z-y (2006) Changes in soil microbial community
associated with invasion of the exotic weed, Mikania micrantha HBK Plant and soil
281:309-324
Magoč T, Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve genome
assemblies Bioinformatics 27:2957-2963
Mangla S, Inderjit, Callaway RM (2008) Exotic invasive plant accumulates native soil pathogens
which inhibit native plants Journal of Ecology 96:58-67
Mitchell CE, Power AG (2003) Release of invasive plants from fungal and viral pathogens
Nature 421:625-627
Nguyen NH et al. (2016) FUNGuild: an open annotation tool for parsing fungal community
datasets by ecological guild Fungal Ecology 20:241-248
Nuzzo V (1999) Invasion pattern of herb garlic mustard (*Alliaria petiolata*) in high quality
forests Biological Invasions 1:169-179
Oksanen J et al. (2010) Vegan: community ecology package. R package version 1.17-4 URL
http://CRAN R-project org/package= vegan
Portier E, Silver WL, Yang WH (2019) Invasive perennial forb effects on gross soil nitrogen
cycling and nitrous oxide fluxes depend on phenology Ecology 100:e02716
Pringle A, Bever JD, Gardes M, Parrent JL, Rillig MC, Klironomos JN (2009) Mycorrhizal
symbioses and plant invasions Annual Review of Ecology, Evolution, and Systematics
40:699-715
Roberts KJ, Anderson RC (2001) Effect of garlic mustard [*Alliaria petiolata* (Beib. Cavara &
Grande)] extracts on plants and arbuscular mycorrhizal (AM) fungi American Midland
Naturalist 146:146-152 doi:Doi 10.1674/0003-0031(2001)146[0146:Eogmap]2.0.Co;2
Rodgers VL, Stinson KA, Finzi AC (2008) Ready or not, garlic mustard is moving in: *Alliaria
petiolata* as a member of eastern North American forests Bioscience 58:426-436
doi:10.1641/B580510
Saikkonen K (2007) Forest structure and fungal endophytes Fungal Biology Reviews 21:67-74
Schoch CL et al. (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal
DNA barcode marker for Fungi Proceedings of the National Academy of Sciences
109:6241-6246
Smith LM (2015) Garlic mustard (*Alliaria petiolata*) glucosinolate content varies across a natural
light gradient Journal of chemical ecology 41:486-492
Stinson KA et al. (2006) Invasive plant suppresses the growth of native tree seedlings by
disrupting belowground mutualisms Plos Biology 4:727-731 doi:ARTN e140
10.1371/journal.pbio.0040140
Stockinger H, Krüger M, Schüßler A (2010) DNA barcoding of arbuscular mycorrhizal fungi
New Phytologist 187:461-474
Team RC (2013) R: A language and environment for statistical computing
Traveset A, Richardson DM (2014) Mutualistic interactions and biological invasions Annual
Review of Ecology, Evolution, and Systematics 45:89-113
Urbanowicz C, Pasquarella VJ, Stinson KA (2019) Differences in landscape drivers of garlic
mustard invasion within and across ecoregions Biological invasions 21:1249-1258

van der Putten WH (2010) Impacts of soil microbial communities on exotic plant invasions
Trends in Ecology & Evolution 25:512-519
Van der Putten WH, Klironomos JN, Wardle DA (2007) Microbial ecology of biological
invasions The ISME Journal 1:28-37
Vaughn SF, Berhow MA (1999) Allelochemicals isolated from tissues of the invasive weed
garlic mustard (*Alliaria petiolata*) Journal of Chemical Ecology 25:2495-2504 doi:Doi
10.1023/A:1020874124645
Wolfe BE, Rodgers VL, Stinson KA, Pringle A (2008) The invasive plant *Alliaria petiolata*
(garlic mustard) inhibits ectomycorrhizal fungi in its introduced range Journal of Ecology
96:777-783 doi:10.1111/j.1365-2745.2008.01389.x
559

Table 1. Results from PERMANOVA tests for differences in overall community composition for fungal (*ITS*) and bacterial (*16S*) communities based on garlic mustard treatments, phenology, and this interaction of treatment and phenology for 2017 and 2018 datasets.

Barcode Gene	Main Fixed Effect	2017			2018		
		F-value	R^2	p-value	F-value	R^2	p-value
ITS	Treatment	1.224	0.018	**0.042**	1.656	0.030	**0.002**
	Phenology	4.509	0.033	**0.001**	2.529	0.046	**0.001**
	Treatment*Phenology	0.893	0.012	0.977	0.938	0.034	0.719
	Site	4.668	0.137	**0.001**	4.943	0.361	**0.001**
16S	Treatment	0.781	0.011	0.781	1.175	0.028	**0.050**
	Phenology	9.889	0.068	**0.001**	2.093	0.025	**0.001**
	Treatment*Phenology	0.541	0.008	0.999	0.908	0.021	0.811
	Site	5.847	0.161	**0.001**	0.244	0.244	**0.001**

Table 2. ANOVA results for main effects on the relative abundance of fungal guilds (ECM, pathotroph, or saprotroph) and the fungal pathogen, *E. aschersoniana*. Site was included as a random effect in all models.

Fungal Functional Group or Taxon	Main fixed effect	*2017*			*2018*		
		df	*F-value*	*p-value*	*df*	*F-value*	*p-value*
ECM	Treatment	2	1.397	0.252	2	0.514	0.601
	Phenology	1	29.882	**<0.001**	2	11.855	**<0.001**
	Treatment*Phenology	2	0.953	0.388	4	0.908	0.466
PATHOTROPH	Treatment	2	6.129	**<0.001**	2	7.023	**0.002**
	Phenology	1	0.735	0.391	2	3.943	**0.026**
	Treatment*Phenology	2	0.108	0.897	4	2.737	**0.039**
SAPROTROPH	Treatment	2	0.168	0.845	2	1.234	0.300
	Phenology	1	0.002	0.965	2	0.846	0.435
	Treatment*Phenology	2	0.009	0.991	4	0.415	0.797
E. ASCHERSONIANA	Treatment	2	21.079	**<0.001**	2	16.528	**<0.001**
	Phenology	1	6.958	**0.009**	2	7.478	**0.001**
	Treatment*Phenology	2	4.108	**0.019**	4	6.088	**<0.001**

Figure 1. Fungal community composition expressed as nonmetric multidimensional scaling axes of Bray-Curtis dissimilarities of rarified count numbers for OTUs within each sample (circle), convex hulls (lines) depict the range of community composition across garlic mustard treatment (AB) and phenology (CD) in 2017 (AC; stress = 0.25) and 2018 (BD; stress = 0.23). Left panels include data across all sampling dates together for each treatment and right panels include data from all treatments together for a given sampling date.

Figure 2. Bacterial and archaeal community composition expressed as nonmetric
multidimensional scaling axes of Bray-Curtis dissimilarities of rarified count numbers for OTUs
within each sample (circle), convex hulls (lines) depict the range of community composition
across garlic mustard treatment (AB) and phenology (CD) in 2017 (AC; stress = 0.12) and 2018
(BD; stress = 0.17). Left panels include data across all sampling dates together for each treatment
and right panels include data from all treatments together for a given sampling date.

Figure 3. Relative abundance (%; mean ± SE) of ectomycorrhizal (ECM), pathotroph, and saprotroph functional groups based on garlic mustard treatment (AB) and phenology (CD) in 2017 (AC) and 2018 (BD). Different letters indicate statistically significant differences (p < 0.05) within each functional group in a given sampling year.

Figure 4. Relative abundance (%; mean ± SE) of the pathogenic fungi, *Entorrhiza aschersoniana,* based on garlic mustard treatment in 2017 (A) and 2018 (B). Different letters indicate statistically significant differences (p < 0.05) within a given sample year.