Genomic analysis provides novel insights into diversification and taxonomy of *Allorhizobium vitis* (i.e. *Agrobacterium vitis*)

Nemanja Kuzmanović[1,2*], Enrico Biondi[3], Jörg Overmann[4], Joanna Puławska[5], Susanne Verbarg[4], Kornelia Smalla[1], Florent Lassalle[6,7*]

[1]Julius Kühn Institute, Federal Research Centre for Cultivated Plants (JKI), Institute for Epidemiology and Pathogen Diagnostics, Messeweg 11-12, 38104 Braunschweig, Germany

[2]Julius Kühn Institute, Federal Research Centre for Cultivated Plants (JKI), Institute for Plant Protection in Horticulture and Forests, Messeweg 11-12, 38104 Braunschweig, Germany (current affiliation)

[3]Alma Mater Studiorum - University of Bologna, Viale G. Fanin, 42, 40127 Bologna, Italy

[4]Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, Inhoffenstrasse 7B, 38124 Braunschweig, Germany

[5]Research Institute of Horticulture, ul. Konstytucji 3 Maja 1/3, 96-100 Skierniewice, Poland

[6]Imperial College London, St-Mary's Hospital campus, Department of Infectious Disease Epidemiology, Praed Street, London W2 1NY, UK; Imperial College London, St-Mary's Hospital campus, MRC Centre for Global Infectious Disease Analysis, Praed Street, London W2 1NY, United Kingdom

[7]Wellcome Sanger Institute, Pathogens and Microbes Programme, Wellcome Genome Campus, Hinxton, Saffron Walden, CB10 1RQ, United Kingdom

*Correspondence: kuzmanovic1306@gmail.com, nemanja.kuzmanovic@julius-kuehn.de (N. Kuzmanović); f.lassalle@imperial.ac.uk (F. Lassalle)

Abstract

Background: *Allorhizobium vitis* (formerly named *Agrobacterium vitis* or *Agrobacterium* biovar 3) is the primary causative agent of crown gall disease of grapevine worldwide. We obtained and analyzed whole-genome sequences of diverse *All. vitis* strains to get insights into their diversification and taxonomy.

Results: Pairwise genome comparisons and phylogenomic analysis of various *All. vitis* strains clearly indicated that *All. vitis* is not a single species, but represents a species complex composed of several genomic species. Thus, we emended the description of *All. vitis*, which now refers to a restricted group of strains within the *All. vitis* species complex (i.e. *All. vitis sensu stricto*) and proposed a description of a novel species, *All. ampelinum* sp. nov. The type strain of *All. vitis sensu stricto* remains the current type strain of *All. vitis*, K309^T. The type strain of *All. ampelinum* sp. nov. is S4^T. We also identified sets of gene clusters specific to the *All. vitis* species complex, *All. vitis sensu stricto* and *All. ampelinum,* respectively, for which we predicted the biological function and infer the role in ecological diversification of these clades, including some we could experimentally validate. *All. vitis* species complex-specific genes confer tolerance to different stresses, including exposure to aromatic compounds. Similarly, *All. vitis sensu stricto*-specific genes confer the ability to degrade 4-hydroxyphenylacetate and a putative compound related to gentisic acid. *All. ampelinum*-specific genes have putative functions related to polyamine metabolism and nickel assimilation. Congruently with the genome-based classification, *All. vitis sensu stricto* and *All. ampelinum* were clearly delineated by MALDI-TOF MS analysis. Moreover, our genome-based analysis indicated that *Allorhizobium* is clearly separated from other genera of the family *Rhizobiaceae*.

Conclusions: Comparative genomics and phylogenomic analysis provided novel insights into

the diversification and taxonomy of *Allorhizobium vitis* species complex, supporting our

redefinition of *All. vitis sensu stricto* and description of *All. ampelinum.* Our pan-genome

analyses suggest that these species have differentiated ecologies, each relying on specialized

nutrient consumption or toxic compound degradation to adapt to their respective niche.

Keywords: *Allorhizobium vitis sensu stricto*, *Allorhizobium ampelinum*, *Rhizobiaceae*,

agrobacteria, grapevine crown gall, taxonomy, plant pathogenic bacteria, clade-specific

genes, ecological specialization, pan-genome analysis

Background

Allorhizobium vitis (formerly named *Agrobacterium vitis* or *Agrobacterium* biovar 3) is a

bacterium primarily known as a plant pathogen causing crown gall disease of grapevine (*Vitis*

vinifera) (1). This economically important plant disease may cause serious losses in nurseries

and vineyards. *All. vitis* is widely distributed pathogen, detected in almost all grapevine

growing regions throughout the world. This bacterium seems to be associated almost

exclusively with grapevine. It has been isolated from crown gall tumors, xylem sap, roots,

rhizosphere, non-rhizosphere soil of infected vineyards, decaying grape roots and canes in

soil, but also from the phyllosphere of grapevine plants (reviewed in (1)). In one exceptional

case, *All. vitis* was isolated from galls on the roots of kiwi in Japan (2).

All. vitis is an aerobic, non-spore-forming, Gram-negative, rod-shaped bacterium with

peritrichous flagella (3). It is a member of the alphaproteobacterial family *Rhizobiaceae*,

together with other genera hosting tumor-inducing plant pathogens, including

Agrobacterium and *Rhizobium*. With time, the taxonomy of *All. vitis* has undergone various

changes. Tumorigenic strains associated with crown gall of grapevine were initially defined

as an atypical group that could neither be classified as *Agrobacterium* biovar 1 (i.e.,

Agrobacterium tumefaciens species complex) nor as biovar 2 (i.e. *Rhizobium rhizogenes*) (4).

Afterwards, several studies classified these atypical strains as *Agrobacterium* biovar 3

(biotype 3), based on their biochemical and physiological characteristics (5-7). Serological

analysis using monoclonal antibodies also allowed differentiation of *Agrobacterium* biovar 3

strains (8). Polyphasic characterization involving DNA-DNA hybridization (DDH), phenotypic

and serological tests clearly showed that *Agrobacterium* biovar 3 strains represent a

separate species, for which the name *Agrobacterium vitis* was proposed (9). However, multi-

locus sequence analysis (MLSA) suggested that *A. vitis* is phylogenetically distinct from the

genus *Agrobacterium*, and prompted the transfer of this species to the revived genus

Allorhizobium (10, 11).

The genus *Allorhizobium* was created by de Lajudie et al. (12) and initially included single

species *Allorhizobium undicola*. Afterwards, Young et al. (13) proposed reclassification of *All.*

undicola and its inclusion into the genus *Rhizobium*, while Costechareyre et al. (14)

suggested that this species might belong to the genus *Agrobacterium*. However, these

studies employed single gene phylogenies, which were insufficient to support such

taxonomic revisions. The authenticity of the genus *Allorhizobium* and the clustering of *All.*

vitis within it was unequivocally confirmed by genome-wide phylogenies (15, 16). Moreover,

distinctiveness of *All. vitis* with respect to the genus *Agrobacterium* was further supported

by their different genome organization, with the genus *Agrobacterium* being characterized

by the presence of a circular chromosome and a secondary linear chromid (17, 18). Chromids

are defined as large non-dispensable plasmids carrying essential functions (19). In contrast

to *Agrobacterium*, the *All. vitis* strains carry two circular chromosomes (18, 20, 21).

However, the smaller circular chromosome (named chromosome II) was later classified as a

chromid in the fully sequenced strain *All. vitis* S4$^\mathrm{T}$ (19). Additionally, genomes of *All. vitis* and

other agrobacteria include a variable number of plasmids.

In recent years, genomics has significantly impacted the taxonomy of bacteria, leading to

the revisions in classification of different bacterial taxa. In particular, a novel genomics-based

taxonomy primarily relies on the calculation of various overall genome relatedness indices

(OGRIs) and estimation of genome-based phylogenies (22-24), largely replacing the

traditionally used methods of 16S rRNA gene phylogeny and DDH (25, 26). Genomic

information were also highly recommended as essential for the description of new rhizobial

and agrobacterial taxa (27). In addition, it has been recommended that some functions and

phenotypic characters may not be considered for taxonomic classification. This particularly

applies to the tumor-inducing ability of agrobacteria, which is mainly associated with the

dispensable tumor-inducing (Ti) plasmid.

Information on genetic diversity and relatedness of strains responsible for crown gall

disease outbreaks provide important insights into the epidemiology, ecology and evolution

of the pathogen. Numerous studies indicated that *All. vitis* strains are genetically very

diverse (reviewed in (1)). In our previous study, we analyzed a representative collection of

All. vitis strains originating from several European countries, Africa, North America, and

Australia using MLSA, which indicated a high genetic diversity between strains, clustered

into four main phylogenetic groups (28). These data suggested that *All. vitis* might not be a

homogenous species, but a species complex comprising several genomic species, warranting

further investigation of the diversification and evolution of *All. vitis* towards a more

complete elucidation of its taxonomy.

In this work, we selected representative strains belonging predominantly to the two most

frequent phylogenetic groups identified in our previous study (28) that included the well-

studied *All. vitis* type strain K309[T] and the fully sequenced strain S4[T], respectively. We

obtained draft genome sequences for 11 additional strains and performed comparative

genomic and phylogenetic analyses to reveal the diversification history and synapomorphies

of these groups. In parallel, we investigated phenotypic features of selected strains. The

combination of these approaches allows us to revise the taxonomy within this group,

notably by emending the description of *All. vitis* (*All. vitis sensu stricto*) and proposing the

new species *All. ampelinum*.

Results

Allorhizobium vitis **genome sequencing**

Draft genome sequences were obtained for 11 *All. vitis* strains (Table 1), with average

coverage depth ranging from 65- to 96- fold. The total size of draft genome assemblies

ranged from 5.67 to 6.52 Mb, with a GC content ranging 57.5-57.6% (Table 1), which was

similar to the genomes of other *All. vitis* strains sequenced so far (Table S1b).

Table 1

Core-genome phylogeny and overall genome relatedness indexes measurements

A core-genome phylogeny was inferred for 14 strains of *All. vitis* (Table 1) and 55 reference

Rhizobiaceae strains (Table S1a). A phylogenomic tree that was reconstructed from the

concatenation of 344 non-recombining core marker genes confirmed the grouping of

Allorhizobium species separately from other *Rhizobiaceae* genera (Figs. 1 and S1). The clade

comprising all members of the genus *Allorhizobium* was well separated from its sister clade,

which included members of the group provisionally named "*R. aggregatum* complex" (11),

as well as representatives of the genus *Ciceribacter*.

All. vitis strains formed a well-delineated clade within the *Allorhizobium* genus (Figs. 1 and

S1). Furthermore, *All. vitis* strains were clearly differentiated into two well-supported sub-

clades (clades A and B), while strain Av2 branched separately from each of these two clades

(Figs. 1 and S1). OGRIs values (Table S2) indicated that sub-clades A and B, as well as strain

Av2, represent separate genomic species. In other words, the core-genome phylogeny and

OGRI measurements showed that *All. vitis* is not a single species, but a species complex

composed of at least three separate genomic species.

The first genomic species, corresponding to sub-clade A, comprises the type strain of *All.*

vitis (strain K309^T) (Fig. 1). Although dDDH values suggested that the cluster containing

strains K309^T and KFB 253 might belong to a separate species compared to other strains

comprised in this sub-clade (Table S2e), this was not supported by the other four OGRIs

calculated here (Table S2a-d). Indeed, dDDH values for these strains (65.9-66.4 %) were

relatively close to the generally accepted threshold value of 70 %. A revised description of

the species *All. vitis*, hereafter referred to as *All. vitis sensu stricto*, is given below.

The second genomic species, corresponding to sub-clade B, included eight strains

originating from various geographic areas (Table 1; Fig. 1). It included the well-studied strain

S4^T, whose high-quality genome sequence was described previously (18). The dDDH value

obtained from the comparison of strain KFB 254 with strain IPV-BO 1861-5 was below, but

very close to the 70 % threshold value generally accepted for species delineation (Table S2e).

However, other OGRIs unanimously indicated that strains from this sub-clade belong to the

same species (Table S2a-d). A description of the novel species corresponding to sub-clade B,

for which the name *Allorhizobium ampelinum* sp. nov. is proposed, is given below.

The third genomic species comprised strain Av2 alone (Figs. 1 and S1, Table S2). To get a

more comprehensive insight into the diversity of the *All. vitis* species complex, we conducted

a second phylogenomic analysis where we included 34 additional genomes of *All. vitis* that

were available in GenBank but not yet published (Table S1b). Based on core-genome

phylogeny and ANIb calculations (Fig. S2, Table S3), additional strains were taxonomically

assigned as *All. vitis sensu stricto* (sub-clade A) and *All. ampelinum* (sub-clade B). Strain Av2

then grouped with three other strains originating from the USA (sub-clade D; Fig. S2). These

four strains comprised in the sub-clade D were genetically very similar and exhibited >99.8

ANI between each other (Table S3). Moreover, additional sub-clades C and E were apparent,

corresponding to two other new genomic species of *All. vitis* species complex (Fig. S2, Table

S3). Genomic species corresponding to sub-clade C and sub-clade D were closely related, as

their ANIb values were in the range 94.62-94.93 %, which is slightly below the threshold for

species delimitation (~95-96%) (29).

Pan-genome analyses

A ML pan-genome phylogeny of the 64 *Rhizobiaceae* genome dataset was estimated from a

matrix of the presence or absence of 33,396 orthologous gene clusters (Fig. 2; Fig. S3). The

pan-genome phylogeny (Fig. 2; Fig. S3) presented the same resolved sub-clades of the *All.*

vitis complex as the core-genome phylogeny (Fig. 1). Furthermore, *Rhizobiaceae* genera and

clades were generally differentiated based on the pan-genome tree (Fig. 2; Fig. S3).

Nevertheless, some inconsistencies were observed: tumorigenic strain *Neorhizobium* sp.

NCHU2750 was more closely related to the representatives of the genus *Agrobacterium*,

while nodulating *Pararhizobium giardinii* H152^{T} was grouped with *Ensifer* spp. (Fig. 2; Fig.

S3). These inconsistencies were also observed in another pan-genome phylogeny inferred

using parsimony (data not shown). Such limitations of gene content-based phylogenies have

previously been reported (30, 31).

Focusing on 14 *All. vitis* species complex strains, we identified 10,501 pan-genome gene

clusters. The core-genome ('strict core' and 'soft core' compartments) of the species

complex comprised 3,775 gene clusters (35.95% of total gene clusters), with 3,548 gene

clusters strictly present in all 14 strains (Fig. 3). The accessory genome contained 4,516 in the

cloud (43% of total gene clusters) and 2,210 gene clusters in the shell (21.05% of total gene

clusters) (Fig. 3).

Clade-specific gene clusters

Homologous gene families specific to particular clades of interest, i.e. with contrasted

presence pattern with respect to closely related clades, were identified using both

Pantagruel or GET_HOMOLOGUES software packages. Both sets of inferred clade-specific

genes were to a large extent congruent, although some differences were observed (Table

S4), owing to the distinct approaches employed by these software packages (32, 33). We

focused on clusters of contiguous clade-specific genes for which we could predict putative

molecular functions or association to a biological process. The results are summarized below

and in Table S4.

All. vitis species complex

Based on Pantagruel and GET_HOMOLOGUES analyses, we identified 206 and 236 genes,

respectively, that are specific to the *All. vitis* species complex (*Av*SC-specific genes), i.e.

present in all strains of *All. vitis sensu stricto*, *All. ampelinum* and *Allorhizobium* sp. Av2, and

in no other *Allorhizobium* strain. *Av*SC-specific genes are mostly located on the second

chromosome (chromid). While some *Av*SC-specific genes are found on the Ti plasmid and

include the type 4 secretion system, this likely only reflects a sampling bias whereby all *All.*

vitis species complex strains in our sample were tumorigenic and possessed a Ti plasmid. As

such, Ti plasmid-encoded genes directly associated with pathogenicity were not further

considered or discussed in this study.

Half of the *Av*SC-specific genes are gathered in contiguous clusters for most of which we

could predict putative function (Table S4); most of the other half are scattered on

chromosome 1 and have unknown function. Predicted functions of clustered genes revealed

that they are strikingly convergent: most are involved in either environmental signal

perception (4 clusters), stress response (2 clusters), aromatic compound and secondary metabolite biosynthesis (3 clusters) and/or aromatic compound degradation response (2 clusters). In addition, one cluster encodes a multicomponent K^+:H^+ antiporter, which is likely useful for adaptation to pH changes, and three clusters harbor several ABC transporter systems for sugar or nucleotide uptake. Finally, one cluster on chromosome 1 encodes a putative auto-transporter adhesin protein, which may have a role in plant commensalism and pathogenesis.

All studied *All. vitis* species complex strains carried a *pehA* gene encoding a polygalacturonase enzyme. Unlike other agrobacteria, *All. vitis* strains are known to produce a polygalacturonase, regardless of their tumorigenicity (34). However, this gene was present also in *All. taibaishanense* 14971[T], *All. terrae* CC-HIH110[T] and *All. oryziradicis* N19[T], but absent in *All. undicola* ORS 992[T] and in other studied members of the *Rhizobiaceae* family.

Furthermore, we detected the presence of gene encoding enzyme 1-aminocyclopropane-1-carboxylate deaminase (*acdS*) in all studied *All. vitis* species complex strains. This gene is considered to be important for plant-bacteria interaction through its involvement in lowering the level of ethylene produced by the plant (35). We found this gene in all other *Allorhizobium* spp., and in some other *Rhizobiaceae* (data not shown), including *R. rhizogenes* strains. However, *acdS* gene was not present in *Agrobacterium* spp., even when the similarity search (blastp) was extended to *Agrobacterium* spp. strains available in GenBank, consistent with previous findings (36).

Tartrate utilization ability was previously reported for most of the *All. vitis* strains (37-39). Therefore, we searched *All. vitis* species complex genomes for the presence of tartrate utilization (TAR) regions. All strains except IPV-BO 6186 and IPV-BO 7105 carried TAR gene clusters. Moreover, we could not find any *All. vitis*-like TAR regions in any other *Rhizobiaceae*

strain. Sequence comparison of TAR regions from *All. vitis* species complex strains using ANIb

algorithm (Table S5) showed they could be divided into four types (Fig. S4). The first type is

represented by a previously characterized TAR region called TAR-I, carried on the tartrate

utilization plasmid pTrAB3 of strain AB3 (39, 40). The second type included representatives

of TAR-II (carried on pTiAB3) and TAR-III (carried on pTrAB4) regions, which were previously

described to be related to each other (39, 41). A third TAR region type, which we designate

TAR-IV, was characterized by the absence of a second copy of *ttuC* gene (tartrate

dehydrogenase). The TAR-IV region type is found in *All. ampelinum* strain S4^T, in which the

tartrate utilization system is located on the large plasmid pAtS4c (initially named pTrS4) (40).

The TAR system of *Allorhizobium* sp. strain Av2 is a unique type (TAR-V), which is related to

region type TAR-I, but is characterized by the absence of the *ttuA* gene (a LysR-like

regulator). We compared the distribution of these TAR region types in strain genomes,

showing there is no TAR region type associated to any genomic species (Table S6). *All. vitis*

sensu stricto strains K309^T and KFB 253 carry a TAR-II/III region. In addition to TAR-II/III

region, strain KFB 239 carries a TAR-I region (Table S6), a combination similar to that found

in the well-characterized strain AB3 (39). *All. ampelinum* strains S4^T, IPV-BO 1861-5, KFB 264

and V80/94 contain a TAR-IV region, while the remaining *All. ampelinum* strains IPV-BO

5159, KFB 243, KFB 250 and KFB 254 additionally carry a TAR-II/III region (Table S6).

All. vitis sensu stricto

Using Pantagruel and GET_HOMOLOGUES pipelines, we identified 63 and 78 genes, that are

specific to *All. vitis sensu stricto* (Av-specific, present in all five strains and in none of *All.*

ampelinum), respectively. 32 of these *Av*-specific genes are clustered into four main loci in

the genome of strain K309^T, for which we could predict putative function (Table S4). One *Av*-

specific gene cluster (Av-GC1, Table S4) comprised genes functionally annotated to be

involved in the degradation process of salicylic acid and gentisic acid (2,5-dihydroxybenzoic

acid) (MetaCyc pathways PWY-6640 and PWY-6223). Av-GC1 was located on Contig 1

(LMVL02000001.1) of reference strain K309^T genome, which is likely part of the chromid,

based on its high ANI with the chromid (Chromosome 2) of strain S4^T, whose genome

sequence is complete. BLAST searches showed that this gene cluster is also present in some

representatives of *Agrobacterium deltaense*, i.e. *Agrobacterium* genomospecies G7 (data not

shown). Av-GC1 is predicted to encode the degradation of salicyl-CoA, an intermediate in

degradation of salicylic acid, to 3-fumarylpyruvate, via gentisic acid. Interestingly, strains KFB

239, IPV-BO 6186 and IPV-BO 7105 carried additional genes encoding the degradation of

salicylaldehyde to salicyl-CoA via salicylic acid and salicyl adenylate, as well as the gene

encoding the final step of gentisic acid degradation, the conversion of 3-fumarylpyruvate to

fumarate and pyruvate. The three strains encoding enzymes of the complete pathway for

degradation of salicylic acid and gentisic acid, and remaining strains K309^T and KFB 253

carrying a partial gene cluster, were phylogenetically separated and formed distinct sub-

clades within *All.* vitis *sensu stricto* (Fig. 1).

Another *Av*-specific gene cluster (Av-GC4, Table S4) was annotated to be involved in the

degradation of 4-hydroxyphenylacetate (MetaCyc pathway 3-HYDROXYPHENYLACETATE-

DEGRADATION-PWY). Gene content and comparative analysis of the contig carrying this

gene cluster suggested that Av-GC4 is carried on a putative plasmid of *All. vitis sensu stricto*

(data not shown).

In addition, *Av*-specific gene clusters Av-GC2 and Av-GC3 (Table S4) were both predicted

to be involved in amino-acid uptake and catabolism. However, we were not able to predict

the precise molecular function of proteins and substrates of enzymes encoded by these *Av*-

specific gene clusters. Both these gene clusters are likely located on a putative plasmid, as

suggested by the presence of plasmid-related genes (replication and/or conjugation

associated genes) on the same contigs.

All. ampelinum

Based on Pantagruel and GET_HOMOLOGUES analyses, we identified 97 genes and 128

genes, respectively, that are specific to *All. ampelinum* (*Aa*-specific, present in all 8 strains

and in none of *All. vitis sensu stricto*). Taking advantage of the finished status of strain S4^T

genome, we found that 52/97 specific genes identified by Pantagruel occur on plasmids

rather than chromosomes. This is a significant over-representation compared to the

distribution of all genes (21.4% on plasmids, Chi-squared test *p*-value < 10^{-6}) or core-genome

genes (5.8% on plasmids, Chi-squared test *p*-value < 10^{-16}). For eleven contiguous gene

clusters we could predict putative function (Table S4). The *Aa*-specific gene clusters encode a

variety of putative biological functions; an enrichment analysis of their functional

annotations revealed a set of high-level biological processes that were over-represented:

transport and metabolism of amino-acids or polyamines like putrescine (three separate

clusters), lysin biosynthesis (two separate clusters), and nickel assimilation. The latter

function is predicted for gene cluster Aa-GC10, which is located on the 631-kb megaplasmid

pAtS4e and encodes the NikABCDE Ni^{2+} import system and a nickel-responsive

transcriptional regulator NikR. Aa-GC10 additionally includes genes with predicted functions

such as cation-binding proteins and a chaperone/thioredoxin, which may be involved in the

biosynthesis of ion-associated cofactors.

Phenotypic and MALDI-TOF MS characterization

The phenotypic properties of the newly described species *All. ampelinum* are listed in Table

2. API 20NE and Biolog GEN III analyses did not reveal clear discriminative features between

All. vitis sensu stricto and *All. ampelinum*. However, a weak positive reaction for 4-

hydroxyphenylacetic (p-hydroxy-phenylacetic) acid for strains belonging to *All. vitis sensu*

stricto was recorded, unlike for those belonging to *All. ampelinum*, which were clearly

negative. As bioinformatic analyses suggested that *All. vitis sensu stricto* strains carry a gene

cluster encoding the degradation of 4-hydroxyphenylacetate, the metabolism of this

compound was assayed in a separate biochemical test. Our results indicated that all *All. vitis*

sensu stricto strains tested are able to metabolize 4-hydroxyphenylacetate, which was

recorded by a vigorous bacterial growth and a change of pH (~7.2 to ~6.5), indicating the

production of acid from the substrate oxidation. On the other hand, *All. ampelinum* strains

showed poor growth under culturing conditions, without change of pH.

Although *All. vitis sensu stricto* strains carry genes predicted to be involved in a

degradation process of gentisic acid, this biochemical property could not be demonstrated in

this study. Gentisic acid degradation genes could have lost their function or not be induced

under our test conditions. Alternatively, the predicted function might be incorrect and the

target substrate of these enzymes may be an unidentified compound more or less closely

related to gentisic acid.

We also tested the ability of *All. vitis* species complex strains to metabolize L-tartaric acid

and produce alkali from this compound. In the present study, we included only strains that

were not tested in our former work (42). Taken together, all tested *All. vitis* species complex

strains (Table 1) were able to produce alkali from tartrate. Interestingly, strains IPV-BO 6186

and IPV-BO 7105, for which we could not identify TAR gene clusters, were also positive for

this test.

As a broader way to characterize and phenotypically distinguish strains, we used MALDI-

TOF mass-spectrometry (MS) of pure bacterial cultures. MALDI-TOF MS revealed diversity

among the tested strains, while allowing to discriminate genomic species (Fig. S5).

Relationship of the genus *Allorhizobium* and related *Rhizobiaceae* genera

As indicated by the core-genome phylogeny, the genus *Allorhizobium* is clearly separated

from the other representatives of the family *Rhizobiaceae,* including the "*R. aggregatum*

complex", which, with the genus *Ciceribacter*, formed a well-delineated sister clade to

Allorhizobium clade (Figs. 1, S1 and S2). The genome-based comparisons showed a clear

divergence between these two clades. In particular, members of the genus *Allorhizobium*

shared >74.9% AAI among each other, and 70.79-72.63% AAI with members of the "*R.*

aggregatum complex"/*Ciceribacter* clade (Table S7). On the other hand, representatives of

the genera *Shinella*, *Ensifer* and *Pararhizobium* showed 71.46-75.85% AAI similarity between

genera. Similarly, representatives of genera *Neorhizobium* and *Pseudorhizobium* showed

72.24-76.18 % AAI similarity between genera. In other words, AAI values suggested that the

existing genera *Ensifer*, *Pararhizobium* and *Shinella*, or *Neorhizobium* and *Pseudorhizobium*

were more closely related than the genus *Allorhizobium* and the "*R. aggregatum*

complex"/*Ciceribacter* clade. gANI and POCP values similarly supported the divergence of

the members of *Allorhizobium* genus and the "*R. aggregatum* complex"/*Ciceribacter* clade

(Table S7). Members of the genus *Allorhizobium* exhibited gANI and POCP values ranging

73.55-76.86 and 55.27-66.17, respectively, when compared with members of the "*R.*

aggregatum complex"/*Ciceribacter* clade, values that were similar to these seen between

representatives of the genera *Agrobacterium* and *Neorhizobium* (gANI 74.66-77.45; POCP

59.96-65.58).

Discussion

***Allorhizobium vitis* is not a single species**

Genomic analyses allowed us to unravel the substantial taxonomic diversity within *All. vitis*.

In particular, whole-genome sequence comparisons and phylogenomic analyses clearly

showed that *Allorhizobium vitis* is not a single species, but represents a species complex

composed of several genomic species. Similarly, *Agrobacterium* biovar 1 (i.e. *A. tumefaciens*)

was initially considered a single species, but was later designated as a species complex

comprising closely related, but distinct genomic species. Several studies applying DDH

initially demonstrated this species diversity within *Agrobacterium* biovar 1 (43-45), which

was later supported by results obtained with AFLP (46, 47), housekeeping gene analysis (10,

11, 14) and whole-genome sequence analysis (30). Although Ophel and Kerr (9) also

performed DDH for several *All. vitis* strains, diversity within this species remained unknown

because these authors only studied strains that belonged to *All. vitis sensu stricto* as defined

here.

Our previous study based on the analysis of several housekeeping gene sequences

suggested the existence of several phylogenetic groups within *All. vitis* species complex (28).

The present study focused on two phylogenetic groups defined in our previous study: the

first comprises the type strain of *All. vitis* (strain K309[T]) (9, 48), whereas the second includes

the well-characterized and completely sequenced strain S4[T] (18). Consequently, we

amended the description of *All. vitis*, which now refers to the limited group within *All. vitis*

species complex strains (*All. vitis sensu stricto*) and proposed a description of a novel

species, *All. ampelinum* sp. nov. (see formal description below).

As indicated by the genome analysis of a larger set of strains available from the NCBI

GenBank database, the taxonomic diversity of *All. vitis* species complex is not limited to *All.*

vitis sensu stricto and *All. ampelinum* sp. nov. However, the description of sub-clades C, D

and E (Fig. S2) as separate species was considered outside the scope of this study, because

the sequencing of these strains was not conducted by our group and their draft genome

sequences are yet to be described in scientific publication(s). In addition, it is not clear

whether sub-clades C and D represent a single or separate species. Further comprehensive

genomic analysis of diverse members of these clades is required to elucidate relationships

between them.

Specific functions and ecologies suggested by clade-specific gene cluster analysis

The convergence of functions encoded by the *All. vitis* species complex-specific genes

suggests an ancient adaptation to different kind of stresses, including exposure to aromatic

compounds, competition with other rhizospheric bacteria and pH change. The occurrence of

multiple signal perception systems in the *All. vitis* species complex-specific gene set indicates

that adaptation to a changing environment is to be a key feature of their ecology.

We also searched genomes of *All. vitis* species complex strains for genes and gene

clusters that were previously reported as important for the ecology of this bacterium. In this

regard, polygalacturonase production, a trait associated with grapevine root necrosis (34,

49, 50), and tartrate degradation (37) were proposed to contribute to the specialization of

All. vitis to its grapevine host. In addition, polygalacturonase activity might be involved in the

process of the invasion of the host plant, as postulated previously for other rhizobia (51).

Although all *All. vitis* species complex strains carried the *pehA* gene encoding a

polygalacturonase enzyme, this gene was not restricted to this bacterial group, as it was also

present in all other *Allorhizobium* spp. strains included in our analysis, except for *All.*

undicola.

All *All. vitis* species complex strains included in this study, except for strains IPV-BO 6186

and IPV-BO 7105, carried TAR regions. However, all of them were able to metabolize tartrate

and produce alkali from this compound. Therefore, we speculate that strains IPV-BO 6186

and IPV-BO 7105 must carry another type of TAR system, distinct from those described so far

in other *All. vitis* strains. Furthermore, some diversity between TAR regions and variable

distribution patterns of different TAR regions among strains were observed, in line with

previously reported data (39). The existence of non-tartrate-utilizing strains was also

documented in the literature (39). Considering the fact that tartrate utilization in *All. vitis*

has only been observed as plasmid-borne (40, 41, 52), this suggests that tartrate utilization is

an accessory trait that can be readily gained via the acquisition of a plasmid encoding this

trait and selected for in tartrate-abundant environments. Because grapevine is rich in

tartrate (53), utilization of this substrate may enhance the competitiveness of *All. vitis*

species complex strains in colonizing this plant species (37).

We observed that an important fraction of the species-specific genes for *All. vitis sensu*

stricto and *All. ampelinum* occurred on chromids and plasmids, suggesting that these

replicons may be an important part of these species' adaptive core-genome, as previously

observed in the *A. tumefaciens* species complex (30). Ecological differentiation of the two

main species of the *All. vitis* species complex seems to rely on consumption of different

nutrient sources, including polyamines and nickel ion (potentially as a key cofactor of

ecologically important enzymes) for *All. ampelinum*, and phenolic compounds for *All. vitis*

sensu stricto.

Even though *All. vitis sensu stricto* strains carried a putative gene cluster which predicted

function was the degradation of gentisic acid, we could not experimentally demonstrate this

trait. Gentisic acid was detected in grapevine leaves (54) and is likely present in other parts

of this plant. This compound was reported as a plant defense signal that can accumulate in

some vegetable plants responding to compatible viral pathogens (55, 56). In addition, a sub-

clade within *All. vitis sensu stricto* composed of strains K309^{T} and KFB 253 carried a complete

pathway for degradation of salicylic acid through gentisic acid. Salicylic acid is recognized as

an important molecule for plant defense against certain pathogens (57). The role of salicylic

and gentisic acid in grapevine defense mechanism against pathogenic bacteria has not been

studied in detail, and further investigations are required to understand their effect against

tumorigenic agrobacteria. Furthermore, we predicted, and demonstrated that all studied *All.*

vitis sensu stricto strains have the specific ability to degrade 4-hydroxyphenylacetate, an

activity that may contribute to the detoxication of aromatic compounds and thus to the

survival of this bacterium in soil, notably in competition against bacteria lacking this

pathway.

Similarly, gene clusters putatively involved in polyamine metabolism or nickel assimilation

might confer to *All. ampelinum* the ability to persist in harsh environments. In this respect,

nickel import has been shown to be essential for hydrogenase function in *E. coli* (58).

Hydrogenase function has in turn been proposed as a potential mechanism for detoxication

of phenolic compounds in *A. vitis* (59) and may thus have an important role in survival in the

rhizosphere.

Delineation of the genus *Allorhizobium*

The genus *Allorhizobium* was clearly differentiated from other *Rhizobiaceae* genera based on

core- and pan-genome-based phylogenies, in line with previous studies employing genome-

wide phylogeny (15, 16). We included diverse *All. vitis* species complex strains into our

analysis, confirming that these bacteria, principally recognized as grapevine crown gall

causative agents, belong to the genus *Allorhizobium*.

On the other hand, the taxonomic status of the *"R. aggregatum* complex"/*Ciceribacter*

clade is still unresolved. Although MLSA suggested that *"R. aggregatum* complex" is a sister

clade of the genus *Agrobacterium*, the more thorough phylogenetic analyses performed in

this study rather showed that the *"R. aggregatum* complex" grouped with *Ciceribacter* spp.,

in a clade that is more closely related to the genus *Allorhizobium*. Presently, there is no

widely accepted criteria and scientific consensus regarding the delineation of new bacterial

genera (27). In this study, existing *Rhizobiaceae* genera were compared using several

delineation methods proposed in the literature, such as AAI (60, 61), POCP (62) and gANI/AF

(63), which we complemented with genome-based phylogenies. Taken together, our

genome-based analysis suggested that *Allorhizobium* represents a genus clearly separated

from other *Rhizobiaceae* genera, including closely related *"R. aggregatum*

complex"/*Ciceribacter* clade. A separate and more focused analysis is however required to

explore the taxonomic diversity and structure of the *"R. aggregatum* complex"/*Ciceribacter*

clade.

Conclusions

Whole-genome sequence comparisons and phylogenomic analyses placed *All. vitis* strains

within the genus *Allorhizobium*, which was clearly differentiated from other *Rhizobiaceae*

genera, including the closely related "*R. aggregatum* complex"/*Ciceribacter* clade. We

revealed an extensive and structured genomic diversity within *All. vitis*, which in fact

represents a species complex composed of several genomic species. Consequently, we

emended the description of *All. vitis*, now encompassing a restricted group of strains within

the *All. vitis* species complex (i.e. *All. vitis sensu stricto*) and proposed a description of a

novel species, *All. ampelinum* sp. nov. Further analyses including pan-genome reconstruction

and phylogeny-driven comparative genomics revealed loci of genomic differentiation

between these two species. Functional analysis of these species-specific loci suggested that

these species are ecologically differentiated as they can consume specific nutrient sources

(*All. ampelinum*), or degrade specific toxic compounds (*All. vitis sensu stricto*). We identified

another two potential genomic species within the *All. vitis* species complex, for which the

limited diversity of available isolates prevented further characterization. We also described

how accessory genomic regions associated with the colonization of grapevine host plant are

distributed across species, and how they combine to form diverse genotypes. However,

given the complete bias in sampling of *All. vitis* strains – all grapevine pathogens – the

ecological significance of this genetic diversity remains unclear. We encourage future studies

to integrate genomic data from new genomically diverse isolates, to further unravel the

ecological basis of *All. vitis* species complex diversification.

Emended description of *Allorhizobium vitis* (Ophel and Kerr 1990) Mousavi et al. 2016

emend. Hördt et al. 2020

The description of *Agrobacterium vitis* is provided by Ophel and Kerr (9). Young et al. (13)

proposed the transfer of *A. vitis* to the genus *Rhizobium*, but it was neither widely accepted

by the scientific community nor supported by further studies (14, 64, 65). Mousavi et al. (11)

reclassified this species to the genus *Allorhizobium*, which was included into the Validation

list no. 172 of the IJSEM (66). Hördt et al. (16) emended a description of *All. vitis* by including

genome sequence data for its type strain, which was published in the List of changes in

taxonomic opinion no. 32 (67).

As shown in this study, *All. vitis sensu stricto* includes a limited group of strains that can

be differentiated from other *All. vitis* genomic species and other *Allorhizobium* species based

on OGRIs, such as ANI, as well as by core-genome phylogeny. Moreover, *All. vitis sensu*

stricto can be differentiated from other species of *All. vitis* species complex by analysis of

sequences of housekeeping genes *dnaK*, *gyrB* and *recA* (28). Finally, this study demonstrated

that strains belonging to this species can be distinguished from *All. ampelinum* by MALDI-

TOF MS analysis. Unlike any *All. ampelinum*, all tested *All. vitis sensu stricto* strains are able

to produce acid in a medium containing 4-hydroxyphenylacetate. However, this apparently

species-specific trait is borne by a plasmid, and could possibly be transmitted to closely

related species.

The whole-genome sequence of type strain K309^T is available in GenBank under the

accessions LMVL00000000.2 and GCA_001541345.2 for the Nucleotide and Assembly

databases, respectively (48). The genomic G+C content of the type strain is 57.55%. Its

approximate genome size is 5.75 Mbp.

Basonym: *Agrobacterium vitis* Ophel and Kerr 1990.

The type strain, K309^T (= NCPPB 3554^T =HAMBI 1817^T = ATCC 49767^T = CIP 105853^T =

ICMP 10752^T = IFO 15140^T = JCM 21033^T = LMG 8750^T = NBRC 15140^T), was isolated from

grapevine in South Australia in 1977.

Description of *Allorhizobium ampelinum* sp. nov.

The description and properties of the new species are given in the protologue (Table 2).

All. ampelinum (am.pe.li'num. Gr. n. ampelos grapevine; Gr. adj. ampelinos and N.L. neut.

adj. ampelinum of the vine).

All. ampelinum strains were formerly classified in the species *All. vitis*. However, our

genomic data showed that they can be distinguished from *All. vitis sensu stricto* and other

All. vitis genomic species based on OGRIs (e.g. ANI and dDDH) and core-genome phylogeny,

as well as by analysis of sequences of housekeeping genes (28). Furthermore, *All. ampelinum*

can be differentiated from *All. vitis sensu stricto* by MALDI-TOF MS analysis.

The type strain, S4^T (= DSM 112012^T = ATCC BAA-846^T) was isolated from grapevine tumor

in Hungary in 1981.

Table 2

Methods

***Allorhizobium vitis* strains**

All. vitis strains used in this study were isolated from crown gall tumors on grapevine

originating from different geographical areas (Table 1). These strains were predominantly

representatives of the two main phylogenetic groups (C and D) delineated in our previous

study (28).

DNA extraction

For whole genome sequencing, genomic DNA was extracted from bacterial strains grown on

King's medium B (King et al. 1954) at 28°C for 24 h using NucleoSpin Microbial DNA kit

(Macherey-Nagel, Germany). The quality of the genomic DNA was assessed by

electrophoresis in 0.8% agarose gel.

Genome sequencing

Draft whole-genome sequences were obtained for 11 *All. vitis* strains (Table 1). DNA libraries

were obtained with Nextera XT DNA Library Prep Kit (Illumina, USA). Paired-end sequencing

(2 × 300 bp) was performed on an Illumina MiSeq platform generating 487,883- 2,309,377

paired reads per genome. Trimming and quality filtering of raw reads were conducted using

Trimmomatic (Galaxy Version 0.36.5) (68) implemented on the Galaxy Web server (69). The

read quality was assessed with FastQC (Galaxy Version 0.72+galaxy1)

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). In order to achieve higher

coverage for strains Av2, IPV-BO 1861-5, KFB239 and KFB 264, additional paired-end

sequencing (2 × 150 bp) was performed using an Illumina NextSeq 500 platform generating

1,037,619-1,443,575 paired reads. Demultiplexing and adapter clipping was done using the

bcl2fastq(2) conversion software (Illumina, USA).

Genome assembly and annotation

De novo genome assemblies were performed using the SPAdes genome assembler (Galaxy

Version 3.12.0+galaxy1) (70). For genomes sequenced on the MiSeq and NextSeq platforms,

both sets of reads were used for assembly. The genome sequences were deposited to

DDBJ/ENA/GenBank under the Whole Genome Shotgun projects accession numbers listed in

Table 1, under BioProject ID PRJNA557463.

The genome sequences were annotated using Prokka (Galaxy Version 1.13) (71) and NCBI

Prokaryotic Genomes Annotation Pipeline (PGAP) (72). Prokka Version 1.14.6 was used to

annotate genomes as a part of the Pantagruel pipeline (task 0; see below and

Supplementary Methods). Functional annotation of proteins encoded by each gene family

clustered by Pantagruel was conducted by the InterProScan software package Version 5.42-

78.0 (Jones et al., 2014) as implemented in the Pantagruel pipeline (Task 4). Additionally,

annotation of particular sequences of interest and metabolic pathway prediction were

performed using BlastKOALA and GhostKOALA (last accessed on December, 2020) (73).

Protein sequences analyzed were subjected to Pfam domain searches (database release

32.0, September 2018, 17929 entries) (74). Metabolic pathway prediction was performed

using KEGG (75) and MetaCyc (76) databases (last accessed on December, 2020).

The NCBI BLASTN and BLASTP (https://blast.ncbi.nlm.nih.gov/Blast.cgi), as well as

BLAST search tool of KEGG database (last accessed on December, 2020) (75), were used for

ad-hoc sequence comparisons at the nucleotide and amino acid levels, respectively.

Core- and pan-genome phylogenomic analyses

For phylogenomic analysis, whole genome sequences of 69 *Rhizobiaceae* strains were used,

including 14 strains of *All. vitis* (Table 1) and 55 reference *Rhizobiaceae* strains (Table S1a).

Additionally, in order to further explore the phylogenetic diversity of *All. vitis*, another core-

genome phylogeny was inferred from an extended dataset that also included 34 *All. vitis*

genomes available from GenBank but not yet published in peer-review journals by sequence

depositors (Table S1b). To build phylogenies based on the core-genome (supermatrix of

concatenated non-recombining core gene alignments) and on the pan-genome (homologous

gene cluster presence/absence matrix), we used the GET_HOMOLOGUES Version 10032020

(33) and GET_PHYLOMARKERS Version 2.2.8.1_16Jul2019 (77) software packages. Details of

the bioinformatic pipeline and used options are described in the Supplementary Methods.

Overall genome relatedness indices

To differentiate between the strains, different OGRIs were computed. For species

delimitation, we relied on the values of average nucleotide identity (ANI) (29, 78) and digital

DNA-DNA hybridization (dDDH) (79) among strain genomes. Because different

implementations of the ANI metric are known to give slightly different results (80), ANI was

calculated using several programs: PyANI Version 0.2.9 (for metrics ANIb and ANIm) (81)

(https://github.com/widdowquinn/pyani), OrthoANIu Version 1.2 (82) and FastANI Version

1.2 (83) tools. dDDH values were calculated using the Genome-to-Genome Distance

Calculator (GGDC) Version 2.1 (79).

For genus delimitation, we relied on average amino acid identity (AAI) (22, 60, 78),

genome-wide average nucleotide identity (gANI) and alignment fraction (AF) (84), and the

percentage of conserved proteins (POCP) (62). AAI values were calculated with CompareM

Version 0.0.23 (https://github.com/dparks1134/CompareM). gANI and AF values were

obtained by the ANIcalculator Version 1.0 (84). POCP values were calculated using

GET_HOMOLOGUES software package (33). Details of the used software and options are

given in the Supplementary Methods.

Genome gene content analyses and identification of clade-specific genes

To explore the distribution of genome gene contents, we conducted further pan-genome

analyses on more focused datasets, using two different bioinformatics pipelines, from which

we present a consensus. Firstly, a pan-genome database was constructed using the

Pantagruel pipeline Version 00aaac71f85a2afa164949b86fbc5b1613556f36 under the

default settings as described previously (31, 32) and in Supplementary Methods. Because of

computationally intensive tasks undertaken in this pipeline, the dataset was limited to the

Allorhizobium genus and its sister clade *"Rhizobium aggregatum* complex*"/Ciceribacter* (28

strains).

Secondly, we analyzed a more focused dataset comprised of the 14 *All. vitis* species

complex strains (Table 1) and four *Allorhizobium* spp. (*All. oryziradicis* N19[T], *All.*

taibaishanense 14971[T], *All. terrae* CC-HIH110[T] and *All. undicola* ORS 992[T]; Table S1a), using

the GET_HOMOLOGUES software package (33). Pan-genome gene clusters were classified

into core, soft core, cloud and shell compartments (85) and species-specific gene families

were identified from the pan-genome matrix. For details on the used scripts and options, see

Supplementary Methods.

Biochemical tests

All. vitis strains were phenotypically characterized using API and Biolog tests. The API 20NE

kit was used according to manufacturer's instructions (bioMérieux). Utilization of sole

carbon sources was tested with Biolog GEN III microplates using protocol A, according to the

instructions of the manufacturer (Biolog, Inc., Hayward, CA, USA).

The metabolism of 4-hydroxyphenylacetic acid (p-hydroxyphenylacetic acid; Acros

Organics, Product code: 121710250) and gentisic acid (2,5-dihydroxybenzoic acid; Merck,

Product Number: 841745) was performed in AT minimal medium (86, 87) supplemented

with yeast extract (0.1 g/L), bromthymol blue (2.5 ml/L of 1 % [w/v] solution made in 50%

ethanol), and the tested compound (1 g/L). Hydroxyphenylacetic and gentisic acid were

added as filter-sterilized 1% aqueous solutions. Bacterial growth and color change of the

medium were monitored during one week of incubation at 28°C and constant shaking (200

rpm/min). Metabolism of L(+)-tartaric acid, involving production of alkali from this

compound, was tested as described before (5).

MALDI-TOF Mass Spectrometry analysis

Sample preparation for MALDI-TOF mass spectrometry (MS) was carried out according to the

Protocol 3 described by Schumann and Maier (88). Instrument settings for the

measurements were as described previously by Tóth et al. (89). The dendrogram was

created using the MALDI Biotyper Compass Explorer software (Bruker, Version 4.1.90).

Abbreviations

AAI: Average amino acid identity; AF: Alignment fraction; ANI: Average nucleotide identity;

DDH: DNA-DNA hybridization; dDDH: Digital DDH; gANI: Genome-wide ANI; MS: Mass-

spectrometry; OGRI: Overall genome relatedness index; MLSA: Multi-locus sequence

analysis; PGAP: Prokaryotic genomes annotation pipeline; POCP: Percentage of conserved

proteins; TAR: Tartrate utilization; Ti: Tumor-inducing.

Acknowledgments

We would like to thank Cathrin Spröer and Boyke Bunk for conducting the Illumina

sequencing. We are grateful to Anja Frühling and Ulrike Steiner for support in phenotypic

tests.

Authors' contributions

NK and FL conceived and designed the study, and analyzed data. FL conceived and

implemented the bioinformatic analysis pipeline Pantagruel. SV performed phenotypic

characterization of the strains (API, Biolog and MALDI-TOF). EB provided bacterial strains. JP,

JO and KS were involved in interpreting data. NK and FL wrote the manuscript. All authors

read, discussed, edited and approved the final manuscript.

Funding

This research was supported by the Georg Forster Fellowship for postdoctoral research from

the Alexander von Humboldt-Foundation, Bonn, Germany. The work of FL was supported by

by a Medical Research Council (MRC) grant (MR/N010760/1) and the Wellcome Trust Grant

[206194]. Bioinformatic analyses were supported by the BMBF-funded de.NBI Cloud within

the German Network for Bioinformatics Infrastructure (de.NBI) (031A537B, 031A533A,

031A538A, 031A533B, 031A535A, 031A537C, 031A534A, 031A532B).

Availability of data and materials

The genome sequences generated in this study were deposited to DDBJ/ENA/GenBank

under the Whole Genome Shotgun projects accession numbers listed in Table 1, under

BioProject ID PRJNA557463. The versions described in this paper are first versions.

All other relevant data (including output of analyses) referring to this project have been

deposited on Figshare under the project accession 20894, available at figshare

(https://figshare.com/), with individual items accessible at DOIs:

10.6084/m9.figshare.17105267,   10.6084/m9.figshare.17125571,

10.6084/m9.figshare.16850071,   10.6084/m9.figshare.16849165,

10.6084/m9.figshare.13440218 and 10.6084/m9.figshare.17125568.

(DOIs to be published upon manuscript acceptance; private links for peer-review:

https://figshare.com/s/b6c7ccc50f0179a08ce6,

https://figshare.com/s/6aaae45f5eb5a81baeb5,

https://figshare.com/s/c88e7412cf58bc9b1b54,

https://figshare.com/s/77bb93622b1bf788bada,

https://figshare.com/s/f17bb1c00a960714e097,

https://figshare.com/s/eecc49ea166bbe061554).

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

[1]Julius Kühn Institute, Federal Research Centre for Cultivated Plants (JKI), Institute for

Epidemiology and Pathogen Diagnostics, Messeweg 11-12, 38104 Braunschweig, Germany.

[2]Julius Kühn Institute, Federal Research Centre for Cultivated Plants (JKI), Institute for Plant

Protection in Horticulture and Forests, Messeweg 11-12, 38104 Braunschweig, Germany

(current affiliation). [3]Alma Mater Studiorum - University of Bologna, Viale G. Fanin, 42,

40127 Bologna, Italy. [4]Leibniz Institute DSMZ-German Collection of Microorganisms and Cell

Cultures, Inhoffenstrasse 7B, 38124 Braunschweig, Germany. [5]Research Institute of

Horticulture, ul. Konstytucji 3 Maja 1/3, 96-100 Skierniewice, Poland. [6]Imperial College

London, St-Mary's Hospital campus, Department of Infectious Disease Epidemiology, Praed

Street, London W2 1NY, UK; Imperial College London, St-Mary's Hospital campus, MRC

Centre for Global Infectious Disease Analysis, Praed Street, London W2 1NY, United

Kingdom. [7]Wellcome Sanger Institute, Pathogens and Microbes Programme, Wellcome

Genome Campus, Hinxton, Saffron Walden, CB10 1RQ, United Kingdom

References

1. Kuzmanović N, Pulawska J, Hao L, Burr TJ. The ecology of *Agrobacterium vitis* and management of
crown gall disease in vineyards. Curr Top Microbiol Immunol. 2018.
2. Sawada H, Ieki H. Crown gall of kiwi caused by *Agrobacterium tumefaciens* in Japan. Plant Dis.
1992;76:212.
3. Young JM, Kerr A, Sawada H. Genus *Agrobacterium* Conn 1942, 359[AL]. In: Brenner DJ, Krieg NR,
Staley JT, Garrity GM, editors. Bergey's Manual of Systematics of Archaea and Bacteria. The
Proteobacteria. 2ed ed. New York: Springer-Verlag; 2005. p. 340-5.
4. Panagopoulos CG, Psallidas PG. Characteristics of Greek isolates of *Agrobacterium tumefaciens*
(E.F. Smith & Townsend) Conn. J Appl Bacteriol. 1973;36:233-40.
5. Kerr A, Panagopoulos CG. Biotypes of *Agrobacterium radiobacter* var. *tumefaciens* and their
biological control. J Phytopathol. 1977;90(2):172-9.
6. Panagopoulos CG, Psallidas PG, Alivizatos AS, editors. Studies on biotype 3 of *Agrobacterium*
radiobacter var. *tumefaciens*. 4th International conference on plant pathogenic bacteria; 1978;
Angers, France.
7. Süle S. Biotypes of *Agrobacterium tumefaciens* in Hungary. J Appl Bacteriol. 1978;44(2):207-13.
8. Bishop AL, Burr TJ, Mittak VL, Katz BH. A monoclonal antibody specific to *Agrobacterium*
tumefaciens biovar 3 and its utilization for indexing grapevine propagation material. Phytopathol.
1989;79:995-8.
9. Ophel K, Kerr A. *Agrobacterium vitis* sp. nov. for strains of *Agrobacterium* biovar 3 from
grapevines. Int J Syst Bacteriol. 1990;40(3):236-41.
10.Mousavi SA, Osterman J, Wahlberg N, Nesme X, Lavire C, Vial L, et al. Phylogeny of the *Rhizobium-*
Allorhizobium-Agrobacterium clade supports the delineation of *Neorhizobium* gen. nov. Syst Appl
Microbiol. 2014;37(3):208-15.
11.Mousavi SA, Willems A, Nesme X, de Lajudie P, Lindström K. Revised phylogeny of *Rhizobiaceae*:
Proposal of the delineation of *Pararhizobium* gen. nov., and 13 new species combinations. Syst Appl
Microbiol. 2015;38:84-90.
12.de Lajudie P, Laurent-Fulele E, Willems A, Torek U, Coopman R, Collins MD, et al. *Allorhizobium*
undicola gen. nov., sp. nov., nitrogen-fixing bacteria that efficiently nodulate *Neptunia natans* in
Senegal. Int J Syst Evol Microbiol. 1998;48(4):1277-90.
13.Young JM, Kuykendall LD, Martinez-Romero E, Kerr A, Sawada H. A revision of *Rhizobium* Frank
1889, with an emended description of the genus, and the inclusion of all species of *Agrobacterium*
Conn 1942 and *Allorhizobium undicola* de Lajudie et al. 1998 as new combinations: *Rhizobium*

radiobacter, R. rhizogenes, R. rubi, R. undicola and *R. vitis*. Int J Syst Evol Microbiol. 2001;51(Pt 1):89-
103.
14. Costechareyre D, Rhouma A, Lavire C, Portier P, Chapulliot D, Bertolla F, et al. Rapid and efficient
identification of *Agrobacterium* species by *recA* allele analysis: *Agrobacterium recA* diversity. Microb
Ecol. 2010;60(4):862-72.
15. Ormeño-Orrillo E, Servín-Garcidueñas LE, Rogel MA, González V, Peralta H, Mora J, et al.
Taxonomy of rhizobia and agrobacteria from the *Rhizobiaceae* family in light of genomics. Syst Appl
Microbiol. 2015;38(4):287-91.
16. Hördt A, López MG, Meier-Kolthoff JP, Schleuning M, Weinhold L-M, Tindall BJ, et al. Analysis of
1,000+ type-strain genomes substantially improves taxonomic classification of Alphaproteobacteria.
Front Microbiol. 2020;11(468).
17. Ramírez-Bahena MH, Vial L, Lassalle F, Diel B, Chapulliot D, Daubin V, et al. Single acquisition of
protelomerase gave rise to speciation of a large and diverse clade within the
Agrobacterium/Rhizobium supercluster characterized by the presence of a linear chromid. Mol
Phylogenet Evol. 2014;73:202-7.
18. Slater SC, Goldman BS, Goodner B, Setubal JC, Farrand SK, Nester EW, et al. Genome sequences of
three *Agrobacterium* biovars help elucidate the evolution of multichromosome genomes in bacteria.
J Bacteriol. 2009;191(8):2501-11.
19. Harrison PW, Lower RP, Kim NK, Young JP. Introducing the bacterial 'chromid': not a chromosome,
not a plasmid. Trends Microbiol. 2010;18(4):141-8.
20. Jumas-Bilak E, Michaux-Charachon S, Bourg G, Ramuz M, Allardet-Servent A. Unconventional
genomic organization in the alpha subgroup of the Proteobacteria. J Bacteriol. 1998;180(10):2749-55.
21. Tanaka K, Urbanczyk H, Matsui H, Sawada H, Suzuki K. Construction of physical map and mapping
of chromosomal virulence genes of the biovar 3 *Agrobacterium* (*Rhizobium vitis*) strain K-Ag-1. Genes
Genet Syst. 2006;81(6):373-80.
22. Konstantinidis KT, Tiedje JM. Towards a genome-based taxonomy for prokaryotes. J Bacteriol.
2005;187(18):6258-64.
23. Chun J, Rainey FA. Integrating genomics into the taxonomy and systematics of the *Bacteria* and
Archaea. Int J Syst Evol Microbiol. 2014;64(2):316-24.
24. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, et al. A standardized
bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nat Biotechnol.
2018;36(10):996-1004.
25. Wayne LG, Brenner DJ, Colwell RR, Grimont PaD, Kandler O, Krichevsky MI, et al. Report of the ad
hoc committee on reconciliation of approaches to bacterial systematics. Int J Syst Bacteriol. 1987;37.
26. Stackebrandt E, Goebel BM. Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA
sequence analysis in the present species definition in bacteriology. Int J Syst Bacteriol. 1994;44.
27. de Lajudie PM, Andrews M, Ardley J, Eardly B, Jumas-Bilak E, Kuzmanovic N, et al. Minimal
standards for the description of new genera and species of rhizobia and agrobacteria. Int J Syst Evol
Microbiol. 2019;69(7):1852-63.
28. Kuzmanović N, Biondi E, Bertaccini A, Obradović A. Genetic relatedness and recombination
analysis of *Allorhizobium vitis* strains associated with grapevine crown gall outbreaks in Europe. J
Appl Microbiol. 2015;119:786-96.
29. Richter M, Rossello-Mora R. Shifting the genomic gold standard for the prokaryotic species
definition. Proc Natl Acad Sci USA. 2009;106(45):19126-31.
30. Lassalle F, Planel R, Penel S, Chapulliot D, Barbe V, Dubost A, et al. Ancestral genome estimation
reveals the history of ecological diversification in *Agrobacterium*. Genome Biol Evol. 2017;9(12):3413-
31.
31. Lassalle F, Dastgheib SMM, Zhao F-J, Zhang J, Verbarg S, Frühling A, et al. Phylogenomics reveals
the basis of adaptation of *Pseudorhizobium* species to extreme environments and supports a
taxonomic revision of the genus. Syst Appl Microbiol. 2021;44(1):126-65.
32. Lassalle F, Veber P, Jauneikaite E, Didelot X. Automated reconstruction of all gene histories in
large bacterial pangenome datasets and search for co-evolved gene modules with Pantagruel.
bioRxiv. 2019:586495.

33. Contreras-Moreira B, Vinuesa P. GET_HOMOLOGUES, a versatile software package for scalable
and robust microbial pangenome analysis. Appl Environ Microbiol. 2013;79(24):7696-701.
34. McGuire RG, Rodriguez-Palenzuela P, Collmer A, Burr TJ. Polygalacturonase production by
Agrobacterium tumefaciens biovar 3. Appl Environ Microbiol. 1991;57(3):660-4.
35. Gamalero E, Glick BR. Bacterial modulation of plant ethylene levels. Plant Physiol. 2015;169(1):13-
22.
36. Bruto M, Prigent-Combaret C, Muller D, Moënne-Loccoz Y. Analysis of genes contributing to plant-
beneficial functions in plant growth-promoting rhizobacteria and related Proteobacteria. Scientific
Reports. 2014;4(1):6261.
37. Salomone JY, Szegedi E, Cobanov P, Otten L. Tartrate utilization genes promote growth of
Agrobacterium spp. on grapevine. Mol Plant-Microbe Interact. 1998;11(8):836-8.
38. Szegedi E. Host range and specific L(+)tartrate utilization of biotype 3 of *Agrobacterium*
tumefaciens. Acta Phytopathol Acad Sci Hung. 1985;20:17-22.
39. Salomone JY, Crouzet P, De Ruffray P, Otten L. Characterization and distribution of tartrate
utilization genes in the grapevine pathogen *Agrobacterium vitis*. Molecular plant-microbe
interactions : MPMI. 1996;9(5):401-8.
40. Szegedi E, Otten L, Czakó M. Diverse types of tartrate plasmids in *Agrobacterium tumefaciens*
biotype III strains. Mol Plant-Microbe Interact. 1992;5:435-8.
41. Crouzet P, Otten L. Sequence and mutational analysis of a tartrate utilization operon from
Agrobacterium vitis. J Bacteriol. 1995;177(22):6518-26.
42. Kuzmanović N, Ivanović M, Prokić A, Gašić K, Zlatković N, Obradović A. Characterization and
phylogenetic diversity of *Agrobacterium vitis* from Serbia based on sequence analysis of 16S-23S
rRNA internal transcribed spacer (ITS) region. Eur J Plant Pathol. 2014;140:757-68.
43. De Ley J. Phylogeny of procaryotes. Taxon. 1974;23:291–300.
44. De Ley J, Tijtgat R, De Smedt J, Michiels M. Thermal stability of DNA: DNA hybrids within the
genus *Agrobacterium*. J Gen Microbiol. 1973;78(2):241-52.
45. Popoff MY, Kersters K, Kiredjian M, Miras I, Coynault C. Taxonomic position of *Agrobacterium*
strains of hospital origin. Annales de microbiologie. 1984;135a(3):427-42.
46. Portier P, Fischer-Le Saux M, Mougel C, Lerondelle C, Chapulliot D, Thioulouse J, et al.
Identification of Genomic Species in *Agrobacterium* Biovar 1 by AFLP Genomic Markers. Appl Environ
Microbiol. 2006;72(11):7123-31.
47. Mougel C, Thioulouse J, Perriere G, Nesme X. A mathematical method for determining genome
divergence and species delineation using AFLP. Int J Syst Evol Microbiol. 2002;52(Pt 2):573-86.
48. Gan HM, Lee MVJ, Savka MA. High-quality draft genome sequence of the type strain of
Allorhizobium vitis, the primary causal agent of grapevine crown gall. Microbiol Resour Announc.
2018;7(9):e01045-18.
49. Rodriguez-Palenzuela P, Burr TJ, Collmer A. Polygalacturonase is a virulence factor in
Agrobacterium tumefaciens biovar 3. J Bacteriol. 1991;173(20):6547-52.
50. Brisset M-N, Rodriguez-Palenzuela P, Burr TJ, Collmer A. Attachment, chemotaxis, and
multiplication of *Agrobacterium tumefaciens* biovar 1 and biovar 3 on grapevine and pea. Appl
Environ Microbiol. 1991;57(11):3178-82.
51. Muñoz JA, Coronado C, Pérez-Hormaeche J, Kondorosi A, Ratet P, Palomares AJ. MsPG3, a
Medicago sativa polygalacturonase gene expressed during the alfalfa-*Rhizobium meliloti* interaction.
Proc Natl Acad Sci USA. 1998;95(16):9687-92.
52. Otten L, Crouzet P, Salomone JY, de Ruffray P, Szegedi E. *Agrobacterium vitis* strain AB3 harbors
two independent tartrate utilization systems, one of which is encoded by the Ti plasmid. Mol Plant-
Microbe Interact. 1995;8:138-46.
53. Ruffner HP. Metabolism of tartaric and malic acids in *Vitis* : A review - Part A. Vitis: Journal of
Grapevine Research. 2016;21:247-.
54. Pantelić MM, Zagorac DČD, Ćirić IŽ, Pergal MV, Relić DJ, Todić SR, et al. Phenolic profiles,
antioxidant activity and minerals in leaves of different grapevine varieties grown in Serbia. J Food
Compos Anal. 2017;62:76-83.

55. Bellés JM, Garro R, Fayos J, Navarro P, Primo J, Conejero V. Gentisic acid as a pathogen-inducible
signal, additional to salicylic acid for activation of plant defenses in tomato. Molecular Plant-Microbe
Interactions®. 1999;12(3):227-35.
56. Bellés JM, Garro R, Pallás V, Fayos J, Rodrigo I, Conejero V. Accumulation of gentisic acid as
associated with systemic infections but not with the hypersensitive response in plant-pathogen
interactions. Planta. 2006;223(3):500-11.
57. Vlot AC, Dempsey DA, Klessig DF. Salicylic acid, a multifaceted hormone to combat disease. Annu
Rev Phytopathol. 2009;47:177-206.
58. Rowe JL, Starnes GL, Chivers PT. Complex transcriptional control links NikABCDE-dependent nickel
transport with hydrogenase expression in *Escherichia coli*. J Bacteriol. 2005;187(18):6317.
59. Biggs J. Ecology and biological control of *Agrobacterium vitis*, the grapevine crown gall pathogen.
Adelaide: University of Adelaide; 1994.
60. Konstantinidis KT, Rossello-Mora R, Amann R. Uncultivated microbes in need of their own
taxonomy. The ISME journal. 2017;11(11):2399-406.
61. Konstantinidis KT, Tiedje JM. Prokaryotic taxonomy and phylogeny in the genomic era:
advancements and challenges ahead. Curr Opin Microbiol. 2007;10(5):504-9.
62. Qin Q-L, Xie B-B, Zhang X-Y, Chen X-L, Zhou B-C, Zhou J, et al. A proposed genus boundary for the
prokaryotes based on genomic insights. J Bacteriol. 2014;196(12):2210.
63. Barco RA, Garrity GM, Scott JJ, Amend JP, Nealson KH, Emerson D. A genus definition for *Bacteria*
and *Archaea* based on a standard genome relatedness index. mBio. 2020;11(1):e02475-19.
64. Farrand SK, van Berkum PB, Oger P. *Agrobacterium* is a definable genus of the family
Rhizobiaceae. Int J Syst Evol Microbiol. 2003;53(5):1681-7.
65. Lindström K, Young JPW. International Committee on Systematics of Prokaryotes Subcommittee
on the taxonomy of *Agrobacterium* and *Rhizobium*: Minutes of the meeting, 7 September 2010,
Geneva, Switzerland. Int J Syst Evol Microbiol. 2011;61(12):3089-93.
66. Oren A, Garrity GM. List of new names and new combinations previously effectively, but not
validly, published. Int J Syst Evol Microbiol. 2016;66(11):4299-305.
67. Oren A, Garrity G. Notification of changes in taxonomic opinion previously published outside the
IJSEM. Int J Syst Evol Microbiol. 2020;70(7):4061-90.
68. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20.
69. Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, et al. The Galaxy platform for
accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucl Acids Res.
2018;46(W1):W537-W44.
70. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A new genome
assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;19(5):455-77.
71. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30(14):2068-9.
72. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al. NCBI
prokaryotic genome annotation pipeline. Nucl Acids Res. 2016;44(14):6614-24.
73. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for functional
characterization of genome and metagenome sequences. J Mol Biol. 2016;428(4):726-31.
74. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, et al. The Pfam protein families
database in 2019. Nucl Acids Res. 2018;47(D1):D427-D32.
75. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference resource for
gene and protein annotation. Nucl Acids Res. 2016;44(Database issue):D457-D62.
76. Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA, et al. The MetaCyc database of
metabolic pathways and enzymes and the BioCyc collection of Pathway/Genome Databases. Nucl
Acids Res. 2013;42(D1):D459-D71.
77. Vinuesa P, Ochoa-Sánchez LE, Contreras-Moreira B. GET_PHYLOMARKERS, a software package to
select optimal orthologous clusters for phylogenomics and inferring pan-genome phylogenies, used
for a critical geno-taxonomic revision of the genus *Stenotrophomonas*. Front Microbiol. 2018;9(771).

78. Goris J, Konstantinidis K, Klappenbach J, Coenye T, Vandamme P, Tiedje J. DNA-DNA hybridization
values and their relationship to whole-genome sequence similarities. Int J Syst Evol Microbiol.
2007;57.
79. Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. Genome sequence-based species delimitation
with confidence intervals and improved distance functions. BMC Bioinformatics. 2013;14(1):1-14.
80. Palmer M, Steenkamp ET, Blom J, Hedlund BP, Venter SN. All ANIs are not created equal:
implications for prokaryotic species boundaries and integration of ANIs into polyphasic taxonomy. Int
J Syst Evol Microbiol. 2020;70(4):2937-48.
81. Pritchard L, Glover RH, Humphris S, Elphinstone JG, Toth IK. Genomics and taxonomy in
diagnostics for food security: soft-rotting enterobacterial plant pathogens. Analytical Methods.
2016;8(1):12-24.
82. Yoon S-H, Ha S-m, Lim J, Kwon S, Chun J. A large-scale evaluation of algorithms to calculate
average nucleotide identity. Antonie Van Leeuwenhoek. 2017;110(10):1281-6.
83. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of
90K prokaryotic genomes reveals clear species boundaries. Nat Commun. 2018;9(1):5114.
84. Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC, et al.
Microbial species delineation using whole genome sequences. Nucl Acids Res. 2015;43(14):6761-71.
85. Koonin EV, Wolf YI. Genomics of bacteria and archaea: the emerging dynamic view of the
prokaryotic world. Nucleic Acids Res. 2008;36(21):6688-719.
86. Tempé J, Petit A, Holsters M, Montagu M, Schell J. Thermosensitive step associated with transfer
of the Ti plasmid during conjugation: Possible relation to transformation in crown gall. Proceedings of
the National Academy of Sciences of the United States of America. 1977;74(7):2848-9.
87. Morton ER, Fuqua C. Laboratory maintenance of *Agrobacterium*. Current Protocols in
Microbiology. 2012;24(1):3D.1.-3D.1.6.
88. Schumann P, Maier T. Chapter 13 - MALDI-TOF mass spectrometry applied to classification and
identification of bacteria. In: Goodfellow M, Sutcliffe I, Chun J, editors. Methods Microbiol. 41:
Academic Press; 2014. p. 275-306.
89. Tóth EM, Schumann P, Borsodi AK, Kéki Z, Kovács AL, Márialigeti K. *Wohlfahrtiimonas*
chitiniclastica gen. nov., sp. nov., a new gammaproteobacterium isolated from *Wohlfahrtia magnifica*
(Diptera: Sarcophagidae). Int J Syst Evol Microbiol. 2008;58(Pt 4):976-81.
90. Bini F, Geider K, Bazzi C. Detection of *Agrobacterium vitis* by PCR using novel *virD2* gene-specific
primers that discriminate two subgroups. Eur J Plant Pathol. 2008;122(3):403-11.
91. Bini F, Kuczmog A, Putnoky P, Otten L, Bazzi C, Burr TJ, et al. Novel pathogen-specific primers for
the detection of *Agrobacterium vitis* and *Agrobacterium tumefaciens*. Vitis. 2008;47:181-9.
92. Fuller SL, Savory EA, Weisberg AJ, Buser JZ, Gordon MI, Putnam ML, et al. Isothermal amplification
and lateral-flow assay for detecting crown-gall-causing *Agrobacterium* spp. Phytopathol.
2017;107(9):1062-8.

Tables:

Table 1 Characteristics of 14 strains of the *All. vitis* species complex analyzed in this study and their genome sequence features

Strain	Species	Geographic origin	Year of isolation	Reference	Genome sequencing	Contigs (N)	N50 (Kb)	Size (Mb)	GC Content (%)	Gene[a]	CDS[a]	Accession number
K309[T]	*All. vitis sensu stricto*	Australia	1977	(9)	(48)	22	999	5.75	57.55	5188	5136	LMVL00000000.2
IPV-BO 6186	*All. vitis sensu stricto*	Italy	2006	(90)	This study	79	608	5.80	57.57	5250	5196	VOLK00000000.1
IPV-BO 7105	*All. vitis sensu stricto*	Italy	2007	(28)	This study	91	462	5.81	57.54	5214	5157	VOLJ00000000.1
KFB 239	*All. vitis sensu stricto*	Serbia	2010	(42)	This study	82	456	6.15	57.57	5545	5490	VOLI00000000.1
KFB 253	*All. vitis sensu stricto*	Serbia	2011	(42)	This study	70	401	5.81	57.56	5290	5235	VOLF00000000.1
S4[T]	*All. ampelinum*	Hungary	1981	(38)	(18)	CG[b]	CG[b]	6.32	57.47	5840	5770	CP000633.1-CP000639.1
IPV-BO 1861-5	*All. ampelinum*	Italy	1984	(91)	This study	110	971	5.67	57.63	5125	5072	VOLM00000000.1
IPV-BO 5159	*All. ampelinum*	Italy	2003	(91)	This study	160	269	6.47	57.56	5939	5883	VOLL00000000.1
KFB 243	*All. ampelinum*	Serbia	2011	(42)	This study	210	270	6.52	57.62	5963	5909	VOLH00000000.1
KFB 250	*All. ampelinum*	Serbia	2011	(42)	This study	135	356	6.47	57.60	5845	5790	VOLG00000000.1
KFB 254	*All. ampelinum*	Serbia	2011	(42)	This study	76	492	5.96	57.60	5433	5380	VOLE00000000.1

KFB 264	*All. ampelinum*	Serbia	2011	(42)	This study	75	714	5.92	57.52	5355	5299	VOLD00000000.1
V80/94	*All. ampelinum*	USA	Oregon	(92)	(92)	66	413	5.98	57.48	5467	5414	NBZE00000000.1
Av2	*Allorhizobium* sp.	Croatia	2006	(28)	This study	123	407	6.23	57.58	5713	5659	VOLN00000000.1

973   [a] Numbers based on Prokka annotations.

974   [b] CG, Complete genome.

975   **Table 2** Protologue for *Allorhizobium ampelinum* sp. nov.

Species name	*Allorhizobium ampelinum*
Genus name	*Allorhizobium*
Specific epithet	*ampelinum*
Species status	sp. nov.
Species etymology	am.pe.li'num. Gr. n. ampelos grapevine; Gr. adj. ampelinos and N.L. neut. adj. ampelinum of the vine
Designation of the type strain	S4
Strain collection numbers	DSM 112012^T, ATCC BAA-846^T
16S rRNA gene accession number	U28505.1
Genome accession number	GCF_000016285.1
Genome status	Complete
Genome size	6,320,946
GC mol %	57.47
Country of origin	Hungary
Region of origin	Orgovány, Bács-Kiskun county
Date of isolation	1981
Source of isolation	Aerial gall on 2-year-old woody grapevine (*Vitis vinifera* cv. 'Izsaki Sarfeher')
Sampling date	1981
Number of strains in study	8
Source of isolation of non-type strains	Grapevine
Growth medium, incubation conditions used for standard cultivation	Yeast mannitol agar (YMA) at 28 °C
Conditions of preservation	-80°C
Gram stain	Negative

Cell shape	Rod
Colony morphology	Colonies on YMA are white to cream coloured, circular, convex and glistening
Positive tests with BIOLOG	pH 6, D-Mannose, D-Galactose, 1% Sodium Lactate, Pectin, Rifamycin SV, Tetrazolium Blue, Potassium Tellurite
Negative tests with BIOLOG	pH5, N-Acetyl Neuraminic Acid, 4% NaCl, 8% NaCl, 3-Methyl Glucose, Inosine, Fusidic Acid, Troleandomycin, D-Serine, Minocycline, Guanidine HCl, Niaproof 4, p-Hydroxy-Phenylacetic Acid, Lithium Chloride, γ-Amino-Butryric Acid, α-Hydroxy-Butyric Acid, α-Keto-Butyric Acid, Propionic Acid, Sodium Butyrate
Positive tests with API	URE, ESC, PNG, GLU (assimilation), ARA, MNE, MAN, MLT, OX
Negative tests with API	NO3, TRP, GLU (fermentation), ADH, GEL, CAP, ADI, PAC
Variable tests with API	NAG, MAL, GNT, CIT
Commercial kits used	BIOLOG GEN3, API 20NE
Oxidase[a]	Positive
Positive tests[a]	Growth at 35°C, growth in nutrient broth supplemented with 2% NaCl, citrate utilization, production of acid from sucrose, production of alkali from tartrate
Negative tests[a]	Production of 3-ketolactose from lactose, acid-clearing on PDA with $CaCO_3$, production of reddish-brown pellicle at the surface of ferric ammonium citrate broth, motility at pH 7.0, acid from d-(+)-melezitose, acid production from 4-hydroxyphenylacetate
Known pathogenicity	Plant pathogenic

[a]These tests were performed for strains S4[T], KFB 243, KFB 250, KFB 254 and KFB 264 by (42), except for a test of acid

production in a medium containing 4-hydroxyphenylacetate conducted in this study. For strains that were not

included in our former study, test of production of alkali from tartrate was conducted in the present work.

Figure captions:

Fig. 1 Maximum-likelihood core-genome phylogeny of 69 strains belonging to the genus

Allorhizobium and other *Rhizobiaceae* members (part collapsed). The tree was estimated with IQ-

TREE from the concatenated alignment of 344 top-ranked genes selected using GET_PHYLOMARKERS

software. The numbers on the nodes indicate the approximate Bayesian posterior probabilities

support values (first value) and ultra-fast bootstrap values (second value), as implemented in IQ-

TREE. The tree was rooted using the *Mesorhizobium* spp. sequences as the outgroup. The scale bar

represents the number of expected substitutions per site under the best-fitting GTR+F+ASC+R6

model. "*Allorhizobium vitis*" clade is collapsed on the left tree and shown expanded on the right. The

matrix represents the distribution of ANIb values for the genomic sequences of the clade

corresponding to the *All. vitis* species complex, relative to the typical species delimitation threshold

of 95%. The same tree, but without collapsing clades, is presented in the Figure S1.

Fig. 2 Maximum-likelihood pan-genome phylogeny of 69 strains belonging to the genus *Allorhizobium*

and other *Rhizobiaceae* members (part collapsed). The tree was estimated with IQ-TREE from the

consensus (COGtriangles and OMCL clusters) gene presence/absence matrix containing 33,396

clusters obtained using GET_HOMOLOGUES software. The numbers on the nodes indicate the

approximate Bayesian posterior probabilities support values (first value) and ultra-fast bootstrap

values (second value), as implemented in IQ-TREE. The tree was rooted using the *Mesorhizobium*

spp. sequences as the outgroup. The scale bar represents the number of expected substitutions per

site under the best-fitting GTR2+FO+R5 model. The same tree, but without collapsing clades, is

presented in the Figure S3.

Fig. 3 Pan-genome analyses of 14 *All. vitis* species complex strains. **a)** Bar plot showing the

frequencies of orthologous clusters as predicted by the COGtriangles and OMCL algorithms. **b)** Pie

chart showing the relative sizes (cluster numbers) contained in the core, soft-core, shell, and cloud

genome compartments.

Supplementary Information

1005

Additional file 1: Fig. S1. Maximum-likelihood core-genome phylogeny of 69 strains belonging to the

genus *Allorhizobium* and other *Rhizobiaceae* members (uncollapsed). The tree was estimated with

IQ-TREE from the concatenated alignment of 344 top-ranked genes selected using

GET_PHYLOMARKERS software. The numbers on the nodes indicate the approximate Bayesian

posterior probabilities support values (first value) and ultra-fast bootstrap values (second value), as

implemented in IQ-TREE. The tree was rooted using the *Mesorhizobium* spp. sequences as the

outgroup. The scale bar represents the number of expected substitutions per site under the best-

fitting GTR+F+ASC+R6 model. The same tree, but with collapsed clades, is presented in Figure 1.

1014

Additional file 2: Fig. S2. Maximum-likelihood core-genome phylogeny of 103 strains belonging to

the genus *Allorhizobium* (including 34 additional strains of *All. vitis* species complex strains whose

sequences are available in GenBank but not associated to a published study) and other *Rhizobiaceae*

members. The tree was estimated with IQ-TREE from the concatenated alignment of 302 top-ranked

genes selected using GET_PHYLOMARKERS software. The numbers on the nodes indicate the

approximate Bayesian posterior probabilities support values (first value) and ultra-fast bootstrap

values (second value), as implemented in IQ-TREE. The tree was rooted using the *Mesorhizobium*

spp. sequences as the outgroup. The scale bar represents the number of expected substitutions per

site under the best-fitting GTR+F+ASC+R7 model. The matrix in the top-right corner represents the

distribution of ANIb values for genomic sequences of the clade corresponding to the *All. vitis* species

complex, relative to the typical species delimitation threshold of 95%.

1026

Additional file 3: Fig. S3. Maximum-likelihood pan-genome phylogeny of 69 strains belonging to the

genus *Allorhizobium* and other *Rhizobiaceae* members (uncollapsed). The tree was estimated with

IQ-TREE from the consensus (COGtriangles and OMCL clusters) pan-genome matrix containing 33,396

clusters obtained using GET_HOMOLOGUES software. The numbers on the nodes indicate the

approximate Bayesian posterior probabilities support values (first value) and ultra-fast bootstrap

values (second value), as implemented in IQ-TREE. The tree was rooted using the *Mesorhizobium*

spp. sequences as the outgroup. The scale bar represents the number of expected substitutions per

site under the best-fitting GTR2+FO+R5 model. The same tree, but with collapsed clades, is presented

in Figure 2.

1036

1037   **Additional file 4: Fig. S4.** Heatmap representation of the average nucleotide identity (ANIb) for TAR

1038   regions of *All. vitis* species complex strains. PyANI program Version 0.2.9

1039   (https://github.com/widdowquinn/pyani) was used to calculate ANIb values and generate the

1040   clustered heatmap.

1041

1042   **Additional file 5: Fig. S5.** Score-oriented dendrogram showing the similarity of the MALDI-TOF mass

1043   spectra of 14 *All. vitis* species complex strains studied. The dendrogram was created using the MALDI

1044   Biotyper Compass Explorer software (Bruker, Version 4.1.90).

1045

Additional file 6: Table S1. List of additional strains and GenBank/EMBL/DDBJ accession numbers for

their nucleotide sequences used in this study. **a)** List of 55 reference *Rhizobiaceae* strains and

GenBank/EMBL/DDBJ accession numbers for their nucleotide sequences used in this study; **b)** List of

additional 34 *All. vitis* species complex strains and GenBank/EMBL/DDBJ accession numbers for their

nucleotide sequences used in this study. Although available in the public nucleotide sequence

databases, these genome sequences have not yet been presented in peer-reviewed study by

sequence depositors.

1053

1054   **Additional file 7: Table S2.** Pairwise OGRI comparisons amongst 14 *Allorhizobium vitis* species

1055   complex strain genomes towards species delimitation. ANI values were computed with various

1056   implementation of the ANI metric: **a)** ANIb, **b)** ANIm, **c)** orthoANI and **d)** fastANI; **e)** dDDH values

1057   were computed with the GGDC.

1058

1059   **Additional file 8: Table S3.** Pairwise ANIb comparisons amongst extended set of 48 *Allorhizobium*

1060   *vitis* species complex strain genomes towards species delimitation.

1061

Additional file 9: Table S4. Clusters of contiguous clade-specific genes. Clusters were identified

amongst sets of genes deemed specific of the focal clade based on detection by either Pantagruel or

GET_HOMOLOGUES pipelines. **a)** Clusters of genes specific to *Allorhizobium vitis* species complex

(present in all *All. vitis sensu stricto*, *All. ampelinum* and *Allorhizobium* sp. Av2, and in no other

Allorhizobium spp.); **b)** Clusters of genes specific to *All. vitis sensu stricto* (present in all five tested

strains and in none of *All. ampelinum*); **c)** Clusters of genes specific to *All. ampelinum* (present in all

eight tested strains and in none of *All. vitis sensu stricto*).

1069

1070   **Additional file 10: Table S5.** Pairwise ANIb values between tartrate utilization (TAR) regions of

1071   *Allorhizobium vitis* species complex strains.

1072

1073   **Additional file 11: Table S6.** Tartrate utilization (TAR) region genotype and tartrate metabolism

1074   phenotype (production of alkali from L-tartaric acid) of *All. vitis* species complex strains.

1075

1076   **Additional file 12: Table S7.** Pairwise OGRI comparisons amongst 69 *Rhizobiaceae* strain genomes

1077   towards genus delimitation. **a)** Average amino acid identity (AAI); **b)** percentage of conserved

1078   proteins (POCP); **c)** genome-wide average nucleotide identity (gANI) and alignment fraction (AF), with

1079   AF values indicated in parentheses.

1080

1081   **Additional file 13: Supplementary Methods.** This file contains additional detailed methods

1082   description.