A *De Novo* regulation design shows an effectiveness in altering plant secondary

metabolism

Abstract

Introduction:

Transcription factors (TFs) and cis-regulatory elements (CREs) control gene

transcripts involved in various biological processes. We hypothesize that TFs and

CREs can be effective molecular tools for *De Novo* regulation designs to engineer

plants.

Objectives:

We selected two Arabidopsis TF types and two tobacco CRE types to design a *De*

Novo regulation and evaluated its effectiveness in plant engineering.

Methods:

G-box and MYB recognition elements (MREs) were identified in four *Nicotiana*

tabacum JAZs (*NtJAZs*) promoters. MRE-like and G-box like elements were

identified in one nicotine pathway gene promoter. TF screening led to select

Arabidopsis Production of Anthocyanin Pigment 1 (PAP1/MYB) and Transparent

Testa 8 (TT8/bHLH). Two *NtJAZ* and two nicotine pathway gene promoters were

cloned from commercial Narrow Leaf Madole (NL) and KY171 (KY) tobacco

cultivars. Electrophoretic mobility shift assay (EMSA), cross-linked chromatin

immunoprecipitation (ChIP), and dual luciferase assays were performed to test the

promoter binding and activation by PAP1 (P), TT8 (T), PAP1/TT8 together, and the

PAP1/TT8/Transparent Testa Glabra 1 (TTG1) complex. A DNA cassette was

designed and then synthesized for stacking and expressing PAP1 and TT8 together.

Three years of field trials were performed by following industrial and GMO

protocols. Gene expression and metabolic profiling were completed to characterize

plant secondary metabolism.

Results:

PAP1, TT8, PAP1/TT8, and the PAP1/TT8/TTG1 complex bound to and activated

NtJAZ promoters but did not bind to nicotine pathway gene promoters. The

engineered red P+T plants significantly upregulated four *NtJAZs* but downregulated

the tobacco alkaloid biosynthesis. Field trials showed significant reduction of five

tobacco alkaloids and four carcinogenic tobacco specific nitrosamines in most or all

cured leaves of engineered P+T and PAP1 genotypes.

Conclusion:

G-boxes, MREs, and two TF types are appropriate molecular tools for a *De Novo*

regulation design to create a novel distant-pathway cross regulation for altering plant

secondary metabolism.

Key words: Arabidopsis; alkaloid; anthocyanin; molecular tools; tobacco;

transcription factor

Introduction

Plant MYB and bHLH proteins are two large TF families with diverse regulations

involved in various biological processes [1-4]. To date, TFs that regulate the

anthocyanin biosynthesis have gained a comprehensive characterization in both model

and non-model plants. MYB and bHLH TFs together with a WD40 protein have been

demonstrated to form MBW complexes to regulate the biosynthesis of anthocyanins

[5-7]. In particular, the MBW complex formed by Arabidopsis Production of

Anthocyanin Pigment 1 (PAP1), Transparent Testa 8 (TT8), and Transparent Testa

Glabra 1 (TTG1) has been characterized to be a main regulatory complex. *PAP1*, *TT8*,

and *TTG1* encode a R2R3-MYB TF (namely MYB75) [8], bHLH TF (namely

bHLH42) [9], and a WD40 protein [10], respectively. Moreover, the functions of

these three genes in Arabidopsis and their homologs in other plants have been

intensively characterized in genetics and biochemistry [11]. PAP1 and TT8 bind to

MYB recognizing elements (MREs) and G-box elements in promoters of anthocyanin

pathway genes (such as *DFR*, *ANS* and *3-GT*, Fig. 1 A), respectively [12-14]. The

current dogma is that on the one hand, PAP1, TT8, and TTG1 form a master

regulatory MYB-bHLH-WD40 (MBW) complex to only activate the anthocyanin

biosynthesis in plants (Fig. 1 A) [6, 15-17]; on the other hand, Arabidopsis JAZ

repressors can interact with PAP1, TT8, and their MBW complex to negatively

regulate the anthocyanin biosynthesis in the absence of jasmonate (JA) [18, 19].

Moreover, *PAP1* has been used to create red crop varieties to engineer anthocyanins,

such as tobacco [20, 21], tomato [22, 23], hop [24], and canola [25]. All these studies

reported that PAP1 alone could activate the anthocyanin biosynthesis in different

tested plants. However, whether PAP1, its partner TT8, and their homologs can

regulate plant secondary metabolism remains open for studies.

Tobacco is an ideal model plant to study plant secondary metabolism, given that it

produces thousands of metabolites in the alkaloid, phenolic, and terpenoid families

[21, 26-28]. Nicotine is the main alkaloid of tobacco (*Nicotiana tabacum*) and

harmfully addictive [29-36]. Other harmful tobacco alkaloids include nornicotine,

anabasine, anatabine, and myosmine (Fig. 1 B). Tobacco specific nitrosamines

(TSNAs) are carcinogenic compounds in tobacco smoke. Five common TSNAs are

N-nitrosonornicotine (NNN), nicotine-derived nitrosamine ketone (NNK), N'-

nitrosoanatabine (NAT), N-nitrosoanabasine (NAB), and 4-(methylnitrosamino)-1-(3-

pyridyl)-1-butanol (NNAL) [37-39]. Given that NNN is a severe carcinogen, the Food

and Drug Administration has proposed a guideline that the average level of NNN

must be less than 1.0 ppm (dry weight) in all final smokeless tobacco products [40].

To reduce these harmful compounds, numerous achievements have been

accomplished in elucidating the biosynthesis of nicotine and other tobacco alkaloids

[41-47]. Nicotine is biosynthesized from two distinct pathways (Fig. 1 B), the steps of

which are catalyzed by enzymes encoded by 10 genes *AO*, *QS*, *QPT2*, *ODC*, *ADC*,

AIH, *PMT*, *MPO*, *A622* and *BBL*, (Fig. 1 B) [42, 48, 49]. In addition, *CYP82E4* and

its homologs have been characterized to involve the demethylation of nicotine to form

nornicotine [50-52]. The mechanisms of the anatabine and anabasine formation are

unknown [42]. One hypothesis is that these two alkaloids result from the BBL

catalysis of nicotinic acid. The other is that anabasine is derived from lysine, a

different pathway (Fig. 1 B) [42]. Four main TSNAs, NNK, NNN, NAT, and NAB

(Fig. 1 B), result from the nitrosation of nicotine, nornicotine, anatabine, and

anabasine during the leaf curing process, respectively [42]. Meanwhile, the regulation

of the nicotine pathway has gained an intensive elucidation. Myelocytomatosis

oncogene (MYC2) and APETALA 2/ethylene response factor (AP2/ERF) of tobacco

are two groups of transcription factors (TFs) that directly bind to promoters of

nicotine pathway genes [42, 48, 53-55]. Moreover, jasmonate (JA), a plant hormone,

essentially controls the nicotine biosynthesis (Fig. 1B) [56-59]. JA regulates the

nicotine biosynthesis via a co-receptor complex consisting of coronatine insensitive 1

(COI1), a Skp/Cullin/F-box (SCF^{COI1}) complex, and JA ZIM-domain (JAZ) repressor

[53-55, 60]. The tobacco genome has been reported to have 15 NtJAZs, in which

NtJAZ1, NtJAZ3, NtJAZ7, and NtJAZ10 are associated with the regulation of the

nicotine biosynthesis [54, 60, 61]. When tobacco roots lack isoleucine-JA (JA-Ile),

NtJAZ1 and NtJAZ3 bind to NtMYC2 and then turn off the nicotine biosynthesis;

when JA-Ile exists, it induces the interaction of SCF^{COI1} and JAZ to form a complex,

which leads to the ubiquitination of JAZs to release NtMYC2. The freed NtMYC2

then activates the biosynthetic pathway of nicotine (Fig. 1 B) [54, 55, 60]. Compared

with the appropriate characterization of NtJAZs' functions, how their transcription in

tobacco is regulated remains open for investigations. To date, based on those pathway

and regulatory genes, on the one hand, antisense and RNAi have been reported to

down-regulate gene expression (such as *BBL*, *ODC*, *QPT*, and nicotine demethylase

gene) to reduce nicotine, nornicotine, or TSNAs in leaves [49, 55, 62, 63]. On the

other hand, these multiple successes revealed a challenge that the reduction of

nicotine was traded with the increase of anatabine and other alkaloids [64, 65]. These

successful engineering data indicate that continuous efforts are necessary to diminish

TSNAs, nicotine, and other alkaloids simultaneously.

Since 2000 when more than 1500 TFs were annotated in the genome of *Arabidopsis*

thaliana in [66], 320,370 TFs had been identified from 156 sequenced plant genomes

and sequences of nine additional plants by 2017 [67]. Likewise, 21,997,501 TF bind

sites (motifs or regulatory elements) (TFBSs) were determined in 63 plant genomes

and nearly 2,493,577 TFBSs (cis-regulatory elements) were localized in the promoters

[68]. Given that the interaction of TFs and TFBSs regulates plant development,

growth, responses to stresses, and metabolisms, we hypothesize that the large

numbers of cis-regulatory elements and TFs are useful molecular tools for *de novo*

designs to create novel regulations for plant engineering. For proof of concept, herein,

we selected tobacco crop to test our hypothesis, given that as described above, the

success might enhance diminishing harmful nicotine, other tobacco alkaloids (OTAs),

and TSNAs for human health. As introduced, given that *NtJAZ1* and *NtJAZ3* encode

two main repressors of the nicotine biosynthesis, we hypothesized that their cis-

regulatory elements were appropriate molecular tools to screen other plant TFs for *de*

novo regulation designs and positive designs could be tested in the downregulation of

nicotine biosynthesis in tobacco plants. Then, we analyzed cis-regulatory elements of

all *NtJAZ* promoters and cloned MRE(s) and/or G-box(es) from four *NtJAZ* genes. In

addition, we analyzed and identified MRE-like and G-box like elements of one

nicotine pathway gene. Meanwhile, we analyzed Arabidopsis TFs that have a

regulatory function to bind to these types of elements. As a result, PAP1 and TT8, two

master regulators of Arabidopsis anthocyanin biosynthesis, were identified for this

design. Mechanistic, transgenic, transcriptional, and metabolic quantification, and

field farming practices showed that this design was successful to create a distant

pathway-cross regulation (DPCR) crossing from the anthocyanin to nicotine pathway.

The constructed DPCR showed an effectiveness in significantly reducing nicotine,

four OTAs, and four TSNAs in tobacco plants, simultaneously. Moreover, the

resulting data disclose new regulatory functions of both PAP1 and TT8 and new

activators of four *NtJAZs*.

Materials and methods

Tobacco varieties

Six genotypic tobacco (*Nicotiana tabacum*) cultivars were used in this study. Two

commercial dark tobacco varieties were Narrow Leaf Madole (NL) and KY171 for

smokeless products (Andersen et al., 1990; Pearce et al., 2015). Two red transgenic

genotypes were generated from NL and KY171 described below. Two others were *N.*

tabacum Xanthi (an oriental tobacco variety) and a red PAP1 tobacco variety that is a

novel homozygous red Xanthi variety (Fig. S22 B) created from the overexpression of

the *Arabidopsis PAP1* (MYB75) (Xie et al., 2006). *Nicotiana benthamiana* was

grown for dual-luciferase assay. Growth conditions in the greenhouse and phytotron

are listed in supporting materials.

Cloning of NtJAZ1, NTJAZ3, PMT2, and ODC2 promoters from Narrow Leaf Madole

and KY171 cultivars

The promoter sequences of tobacco *NtJAZ1, NtJAZ3, NtJAZ7, NtJAZ10, PMT2*, and

ODC2 were identified from the genomic sequences of the NT90 cultivar curated at

NCBI (https://www.ncbi.nlm.nih.gov/). The sequences identified were 1000 bp or

1500 bp. Based on these sequences, primers (Table S1) were designed for PCR to

amplify *NtJAZ1, NtJAZ3, PMT2*, and *ODC2* promoters from the genomic DNA of

wild type NL and KY171 tobacco plants, and then the amplified genomic DNA

fragments were cloned to pEASY-T1 for sequencing. Briefly, the DNA fragments

from PCR were purified using a gel-extraction Kit (Thermo Fisher, Waltham, MA) by

following the manufacturer's instruction and then ligated to the pEASY-T1 plasmid

using a T4-ligase system (10 µl reaction system: 1 µl T4 buffer, 1 µl T4-ligase, 1.0 µl

pEASY-T1 linear plasmid and 7.0 µl purified PCR product). The resulting

recombinant plasmid was transformed into the competent *E. coli* Top10 cells, which

were screened on agar-solidified LB medium supplemented with 50 mg/l ampicillin.

Positive colonies were selected for sequencing at Eton Bio (Durham, NC, USA). All

promoter sequences obtained were 1000 or 1500 bp (Figs. S1, S2, and S4). Analysis

of promoter sequences and identification of regulatory elements such as AC- and AC-

like elements (Figs. S1, S2, and S4) were completed with the PLACE

(http://www.dna.affrc.go.jp/PLACE/) and PlantPan (http://plantpan.itps.ncku.edu.tw/)

tools. The *NtJAZ1, NtJAZ3, PMT2*, and *ODC2* promoters were used for

electrophoretic mobility shift assay, dual-luciferase, and ChIP assays described below.

Electrophoretic mobility shift assay of PAP1 and TT8 binding to promoters

To perform electrophoretic mobility shift assay (EMSA), we designed gene-specific

primers (Supporting Table S1) for PCR to amplify the binding domains of both *PAP1*

and *TT8*. The amplified *PAP1* binding domain from the N-terminus was 534 bp,

which included the *R2R3 MYB* domain sequences for Gate-way cloning. The

amplified *TT8* binding domain franking the *bHLH* region was 252 bp, in which both

BamHI and SaclI restriction sites were added to two ends for further cloning. After

the confirmation of sequence accuracy, three expression vector systems, pRSF-Dute,

pet-Dute, and pDest-His-MBP were comparatively tested to induce soluble

recombinant proteins. We selected the pDest-His-MBP vector, a Gateway cloning

system, for the *R2R3* domain of *PAP1*. We found that the pRSF-Dute vector with a

His-tag in the N-terminus was appropriate for expressing the bHLH domain of TT8

after several vectors were tested. As reported previously[69], detailed steps are listed

in Method S1.

Dual luciferase assays

Four reporter and four effector vectors were developed to perform dual luciferase

assays that were completed to analyze the regulatory activity of PAP1 alone, TT8

alone, PAP1 and TT8 together, and PAP1/TT8/TTG1 (WD40) together in activating

NtJAZ1, *NtJAZ3*, *PMT2,* and *ODC2* promoters. First, the pGreenII-0800 vector was

used to develop four reporter vectors. Next, four effector vectors were developed from

the PK2GW7 vector [20], to which the ORFs of *GFP*, *PAP1*, *TT8*, and *TTG1* were

cloned. Each gene was driven by 35S promoter. Then, four reporter and four effector

vectors were introduced into *Agrobacterium tumefaciens* strain GV3101, respectively.

Last, the activated effector and reporter *Agrobacterium* GV3101 cells were used to

inoculate leaves of *N. benthamiana* for transient expression assay. Detailed steps are

described in Method S2.

Cross-linked chromatin immunoprecipitation (ChIP) assays

A gateway cloning was used to develop two plasmids, in which *GFP* was fused to the

3'-end of *PAP1* and *TT8* ORFs without their stop codon for genetic transformation. A

pair of primers (Supporting Table S1) was designed to amplify the *PAP1* ORF

without its stop codon. The resulting PCR product was purified as described above

and then cloned into the pDonr221 vector via a BP reaction. The *PAP1* fragment was

further cloned to the upstream of *GFP* in the destination vector pGWB5 via a LR

reaction. This cloning step generated a recombinant vector pGWB5-PAP1-GFP, in

which *PAP1-GFP* was driven by 35S promoter. The pGWB5-PAP1-GFP vector was

introduced into the competent *E. coli* Top10 cells. Positive colonies were selected on

LB medium containing 50 mg/l kanamycin. One colony was used for suspension

culture on the same shaker as described above. Cultured *E. coli* cells were harvested

to isolate the pGWB5-PAP1-GFP binary plasmid, which was further introduced into

competent *Agrobacterium* GV301 cells. To develop a binary vector expressing *TT8*, a

pair of primers (Supporting Table S1) was designed for PCR. The reverse primer was

designed to include a HA-tag sequence. The resulting PCR product was cloned to the

plasmid PK2GW7 as described above. Next, the resulting binary vector PK2GW7-

TT8-HA-tag was introduced into competent *Agrobacterium* GV301 cells.

Agrobacterium GV301 cells harboring *PAP-GFP* or *TT8-HA-tag* cDNA were

activated and then injected to leaves of 5-week old wild type NL and KY171

seedlings that were grown in the phytotron. After the injection, plants were allowed to

grow 72 hrs. The infected leaves were then harvested to liquid nitrogen to isolate

nuclei and ChIP experiments. Detailed steps are listed in Method S3.

Synthesis of T-DNA, development of binary constructs, and genetic transformation of

two dark tobacco varieties

PAP1 and *TT8* of *A. thaliana* encode a R2R3-MYB and bHLH TF, respectively. A

DNA cassette was designed and then synthesized to stack *PAP1* and *TT8* for their

coupled overexpression. From 5' to 3' end, the cassette was composed of *attL1*, the

PAP1 cDNA (NP_176057), *NOS* (terminator), *35S* promoter, the *TT8* cDNA

(CAC14865), and *attL2* (Fig. S7 A). The synthesized sequence was introduced to the

entry vector pUC57 and then cloned to the destination binary vector pK2GW7 by attL

× attR combination (LR) reaction with LR clonase. The resulting binary vector,

namely PAP1-TT8-pK2GW7 (Fig. S7 B), was further introduced into the competent

cells of *Agrobacterium tumefaciens* strain GV3101. A positive colony was selected

for genetic transformation of tobacco. In addition, the pK2GW7 vector was used for

genetic transformation as control. These constructs were transformed into two

globally commercial dark tobacco varieties, NL and KY171, respectively. Detailed

steps (Fig. S7 C) for transformation and selection of T0 transgenic plants followed our

protocols reported previously [20]. Transgenic plants from NL and KY171 were

labeled as P+T-NL and P+T-KY, respectively. More than 20 T0 transgenic plants

were obtained for each variety and grown in the greenhouse to select T0 seeds, 12 of

which were further selected on medium supplemented with antibiotics to screen

antibiotic resistant T1 progeny. A 3:1 ratio of seed germination to seed death

indicated one single copy of T-DNA that was in the genomic DNA of T0 plants. This

ratio was observed in the T1 progeny of T0 plants. In this study, we focused on lines

with one single copy of transgenes. A large number of seeds were obtained from the

T1 progeny of each line. Seeds from these types of T1 progeny were screened on an

agar-solidified MS medium supplemented with 50 mg/l kanamycin. If all seeds from a

T1 plant could germinate T2 seedlings, it indicated that this T1 plant was a

homozygous line. Accordingly, red homozygous T2 plants (Fig. S7 M) were obtained

and then used for growth in the greenhouse and field trial. In addition, PCR and qRT-

PCR were performed to demonstrate the presence of transgene in the genome of T0

and T2 transgenic plants and the expression of both *PAP1* and *TT8* in both P+T-NL

and P+T-KY plants.

DNA and RNA extractions

Genomic DNA was extracted using a DNeasy Plant Mini Kit (QIAGEN, Hilden,

Germany). RNA was extracted using Trizol extraction reagents. Fresh plant samples

were ground into fine powder in liquid nitrogen. Extraction steps were followed the

manufacturers' protocols. Detailed steps are listed in Method S4.

Polymerase chain reaction and quantitative reverse transcript-polymerase chain

reaction

All real-time quantitative RT-PCR (qRT-PCR) analyses were performed for wild-

type, vector control transgenic, and red transgenic plants. SYBR-Green PCR Master-

mix (Bio-Rad, Hercules, CA) was used for qRT-PCR experiments on a Step-One

Real-time PCR System (Thermo Fisher, Waltham, MA). Detailed steps are listed in

Method S5.

Assays of anthocyanin and proanthocyanidin in seedlings

Fresh leaves and roots (100 mg) of wild type, P+T-NL1, P+T-KY1, and vector control

seedlings grown hydroponically (Supporting Figure S9) were used to extract

anthocyanins and proanthocyanidins (PAs). Methods were followed our anthocyanin

[20] and proanthocyanidin assay protocols [70]. Detailed steps are listed in Method

S6.

Field trial of P+T-NL1 and P+T-KY1 T2 progeny and air curing

Homozygous T2 progeny of P+T-NL1 and P+T-KY1 were planted at the research

station in Oxford, North Carolina in 2015. In addition, NL and KY171 plants were

planted as controls. Prior to planting transgenic plants in the field, we applied for

GMO permits from USDA-APHIS for this field trial. North Carolina State University

was licensed to grow transgenic tobacco plants in the field. Every single step strictly

followed the protocol requested in the license. Field planting also followed the

protocol of commercial tobacco production in North Carolina. These steps included

seedling growth in a contained greenhouse, selection and design of field, planting,

phenotypic observation, field management, topping, harvest of leaves, air curing, and

cleanup of plant residues from the field (Table S2 and Supporting Figures S15-S19).

Detailed steps are listed in Method S7 (Table S2, Figs S15-S19).

Field trial of red PAP1 tobacco plants and flue-curing

PAP1 tobacco is the progeny of a homologous red PAP1 Xanthi plant (#292) that is

generated by the overexpression of *PAP1* [21]. The field trials for PAP1 tobacco were

conducted in 2011 and 2012. Field design, seed germination, plantation, field

management, and plant growth management were the same as described above for

P+T-NL1 and P+T-KY1 genotypes. The main differences were that leaves were

harvested at different dates and flue-cured in barns. Detailed steps are listed in

Method S8 (Fig. S22 A-E).

Determination of TSNAs in cured leaves of field-grown P+T-NL1 and P+T-KY1

tobacco plants with high performance liquid chromatography coupled with triple

quadrupole tandem mass spectrometer

Accurate quantification of tobacco specific nitrosamines (TSNA) was completed with

high performance liquid chromatography coupled with triple quadrupole tandem mass

spectrometer (HPLC-QQQ-MS/MS). This analysis was performed using industrial

protocols on an Agilent 1200 series HPLC System coupled with AB Sciex Triple

quadrupole mass spectrometer (API 5000 with Electrospray) at RJ Reynolds. Four

TSNAs, N-nitrosonornicotine (NNN), N-nitrosoanatabine (NAT), N-nitrosoanabasine

(NAB), and nicotine-derived nitrosamine ketone (4(methylnitrosamino)-1-(3-pyridyl)-

1-butanone, NNK) in air-cured leaves, were quantified. One hundred milligrams of

dry leaf powder were extracted in an aqueous ammonium acetate solution (100 mM

aqueous ammonium acetate containing four deuterium analogs for TSNA) and filtered

through disposable PVDF syringe filters into autosampler vials. The extract was

separated in a Phenomenex Gemini C18 column with 3.0 μm particle 2.0 x150 mm.

The injection volume was 2.0-5.0 μl. TSNAs were detected by multiple reaction

monitoring (MRM) of the precursor ion to a product ion specific for each compound.

Quantification was achieved using an internal standard calibration comprised of ten

points. A separate internal standard was used for each analyte by using a mixture of

four stable isotope-labeled analytes. Results are reported in units of ng/g (ppb). Data

were determined to be acceptable if the correlation coefficient for the calibration

curve was greater than 0.99, standards were within 85% to 115% of expected

concentration values, check solution values were within 75% to 125% of target, peak

shape and resolution were acceptable (based on historical data), and values for the QC

samples were within established control limits. The instrument parameters for MS,

HPLC gradient, the quadrupole mass spectrometer parameters, reagents and standards

are listed in in Supporting Tables S3-S6.

Determination of nicotine and other tobacco alkaloids in cured leaves of field grown

P+T-NL1 and P+T-KY1 tobacco plants by GC

Analysis of gas chromatograph (GC) equipped with a flame ionization detector (FID)

was performed using a standard industry method to quantify nicotine and other

tobacco alkaloids at RJ Reynolds. In this study, analysis of nicotine and other tobacco

alkaloids was performed on an Agilent 6890 gas chromatograph (GC) equipped with a

flame ionization detector (FID) and an Agilent 7683 automatic sampler. One hundred

milligrams of air-cured dry leaf powder were alkalinized in 2 mM sodium hydroxide

(NaOH). This solution was extracted with methyl-tertbutyl ether (MTBE) spiked with

an internal standard using a wrist-action shaker. The mixture was allowed to separate

to two layers. The resulting MTBE layer was transferred to an autosampler vial for

analysis of nicotine, anabasine, nornicotine, and myosmine with GC-FID.

Quantification was achieved using an internal standard calibration comprised of six

points. Data were determined to be acceptable if the correlation coefficient of the

calibration curve was greater than 0.998, the response factors (RF) were consistent,

the percent relative standard deviation (%RSD) for RF was equal or less than 5%, the

check solution values were within 10% of target, values for the quality control (QC)

samples were within established control limits, and chromatograms had appropriate

identification of peaks. The instrument parameters, the oven program, and reagents

and conditions used in alkaloid analysis are listed in Supporting Table S7-S9. The

results were reported as percentiles (%, g/g, dry weight).

Determination of TSNAs and tobacco alkaloids in flue-cured leaves of PAP1 plants

with HPLC-ESI-MS

HPLC-ESI-MS analysis was performed on a 2010EV LC/UV/ESI/MS instrument

(Shimadzu, Japan). One gram of powdered samples was extracted in 20 ml of

methanol in a flask placed on a shaker for one hour at room temperature. The samples

were centrifuged twice at 4000 rpm and aliquots of the supernatant were transferred to

a 2.0 ml glass vial. The samples were separated on an Eclipse XDB-C18 analytical

column (250 mm × 4.6 mm, 5 µm, Agilent, Santa Clara, CA, USA). The mobile phase

solvents were composed of 1% acetic acid in water (solvent A; HPLC-grade acetic

acid and LC–MS grade water) and 100% LC–MS grade acetonitrile (solvent B). To

separate metabolites, the following gradient solvent system was used with ratios of

solvent A to B: 90:10 (0–5 min), 90:10–88:12 (5–10 min), 88:12–80:20 (10–20 min),

80:20–75:25 (20–30 min), 75:25–65:35 (30–35 min), 65:35–60:40 (35–40 min),

60:40–50:50 (40–55 min), and 50:50–10:90 (55–60 min). The column was then

washed for 10 min with 10% solvent B. The flow rate and the injection volume were

0.4 ml/min and 20 µl, respectively. The total ion chromatograms of positive

electrospray ionization were recorded from 0 to 60 min by mass spectrum detector

and mass spectra were scanned and stored from m/z of 120–1,600 at a speed of 1,000

amu/s. Standards of NAB, NAT, NNAL, NNK, and NNN (Sigma, St-Louis, MO)

were used to establish standard curves with a coefficient value at least 98%. The peak

values were used to estimate the levels of nicotine, nornicotine, anabasine, and

anatabine.

Determination of nicotine and nornicotine in leaves and roots of seedlings with

HPLC-qTOF-MS

Seedlings of P+T-NL1, P+T-KY1, vector control, and wild type plants were

hydroponically grown in the phytotron. Leaves and roots of 45-day old seedlings

(Supporting Figure S9) were used to analyze nicotine and nornicotine in both leaves

and roots. Fresh tissues were ground into fine powder in liquid nitrogen and dried in a

45°C oven. One hundred mg of powdered samples was extracted with 2.0 ml

methanol. Other steps were as described above. The analysis of nicotine and

nornicotine was performed with HPLC-qTOF-MS on Agilent 6520 time-of-flight LC-

MS/MS instrument (Agilent Technologies, Santa Clara, CA, USA). Mobile solvents

and column used for separation of metabolites were solvent A and B described above.

The gradient solvent system was composed of ratios of solvent A to B: 95:5 (0–5

min), 95:5 to 90:10 (5–10 min), 90:10 to 85:15 (10–15 min), 85:15 to 45:55 (15–20

min), 45:55 to 25:75 (20–25 min), 25:75 to 95:5 (25–26 min). After the last gradient

step, the column was equilibrated and washed for 10 min with solvents A: B (95:5).

The flow rate was 0.4 ml/min. The injection volume of samples was 5.0 µl. The dry

gas flow and the nebulizer pressure were set at 12 l/min and at 50 psi, respectively.

Metabolites were ionized with the negative mode. The mass spectra were scanned

from 100 to 3000 m/z. The acquisition rate was three spectra per second. Other MS

conditions included fragmentor: 150 V, skimmer: 65 V, OCT 1 RF Vpp: 750 V, and

collision energy: 30 psi. Standard curves were developed for both nicotine and

nornicotine. The results were reported as mg/g, dry weight.

Statistical analysis

Metabolite values for each group of samples such as B1 were averaged from samples

of three plots. One-way ANOVA was used to evaluate the statistical significance of

the contents of nicotine, nornicotine, anabasine, NNN, NNK, NAT, and NAB

between red and wild type leaves. Standard deviation was used to show the variation

of metabolite contents. For gene expression, protein analysis, nicotine and nornicotine

contents in seedlings, all values are averaged from four or five biological replicates.

Student t-test was performed to evaluate statistical significance. P-values less than

0.05, 0.01, and 0.001 mean significant, very significant, and extremely significant.

Results

MRE and G-box elements identified in the promoters of NtJAZ1, 3, 7, and 10

NtJAZ1 and NtJAZ3 are two main repressors of nicotine biosynthesis [54, 55, 60]. In

recent, totally 15 *NtJAZs* were mined from the tobacco genomes, showing that

NtJAZ7, and *NtJAZ10* were also associated with the regulation of nicotine

biosynthesis [61]. We identified the promoter sequences of 13 *NtJAZs* from the

genome of NC90 tobacco curated at NCBI and named these as *NtJAZ1pro* through

NtJAZ13pro. One or 1.5 kb nucleotides in the proximal upstream from the start codon

(ATG) were obtained for sequence characterization. Sequence analysis revealed that

NtJAZ1pro, *NtJAZ3pro,* and *NtJAZ7pro* had both MYB recognizing element (MRE)

and G-box elements, while *NtJAZ10pro* only had G-box elements (Figs S1-S2). Next,

we cloned the promoter sequences of *NtJAZ1pro* and *NtJAZ3pro* from two

commercial smokeless dark tobacco cultivars, Narrow Leaf Madole (NL) and KY171

(KY). Their sequences were identical to those of *NtJAZ1pro* and *NtJAZ3pro* from

TN90 (Fig. S1), therefore, we used the same names in this report. Sequence analysis

further substantiated the MRE and G-box elements in *NtJAZ1pro* and *NtJAZ3pro*.

NtJAZ1pro has two MRE and two G-box elements (Fig. 2 A and Fig. S1). The two

MRE elements are MRE1: ACCCACC at position -180 and MRE2: ACCCCAC at

position -172. The two G-boxes are <u>CACGTG</u> at positions -232 and AACGTG at -

395. *NtJAZ3pro* has one MRE3 element: <u>AACTACC</u> at position -169 and one G-box:

<u>CACGTG</u> at position -960 (Fig. 2 A and Fig. S1).

EMSA shows the binding of PAP1 and TT8 to the NtJAZ1 and NtJAZ3 promoters in

vitro

Electrophoretic mobility shift assay (EMSA) was carried out to test whether PAP1

and TT8 could bind to the MRE and G-box elements of *NtJAZ1$_{pro}$ and NtJAZ3$_{pro}$ in*

vitro. The R2R3-MYB binding domain (amino acids 1-178) of Arabidopsis PAP1 was

cloned to the pDEST-HisMBP vector, in which it was fused with the His-MBP-Tag at

the N-terminus (Fig. S3 A). The bHLH binding domain (amino acid 359-443) of TT8

was cloned to the PRSF-Dute vector, in which it was fused with the His-Tag at the N-

terminus (Fig. S3 B). Two constructs were introduced to *E. coli* to induce

recombinant proteins, which were further purified with a His-tag purification system

(Fig. S3 B). Biotin probes were prepared for MRE1 (ACCCACC), MRE2

(ACCCCAC), MRE3 (AACTACC), and G-box (CACGTG). Both competitive and

non-competitive probes were used as controls for EMSA. The resulting data showed

that the recombinant R2R3 binding domain directly bound to three MRE element

probes and the recombinant bHLH binding domain directly bound to the G-box

elements (Fig. 2 B). The binding activity of both recombinant R2R3 and bHLH

binding domains was competed by competitive probes but not competed by non-

competitive probes (Fig. 2 B). These data demonstrate that both PAP1 and TT8 bind

to the *NtJAZ1$_{pro}$ and NtJAZ3$_{pro}$*.

ChIP-Q-PCR shows the binding of PAP1 and TT8 to the NtJAZ1 and NtJAZ3

promoters in vivo

ChIP-Q-PCR analysis was performed to examine whether PAP1 and TT8 could bind

to *NtJAZ1pro* and *NtJAZ3pro in vivo*. In ChIP experiments, *PAP1* was fused at the 5'-

end of *GFP* in the binary vector pGWB5 to obtain pGWB5-PAP1-GFP, which was

introduced into leaves of wild type NL and KY171 seedlings for transient expression

via injection of *Agrobacterium*. *TT8* was fused at the 5'-end of an *HA-tag* sequence in

the binary vector PK2GW7 to obtain PK2GW7-TT8-HA-tag, which was also

introduced into leaves of wild type NL and KY171 seedlings for transient expression

by injection of *Agrobacterium*. The IP experiments of PAP1 and TT8 were performed

by using GFP and HA monoclonal antibodies. The ChIP PCR results showed that the

amplified *NtJAZ1* and *NtJAZ3* promoter fragments were increased 5.0-40.4 times by

PAP1 and 6.1.8-37.5 times by TT8 (Fig. 2 C).

Dual-luciferase assay demonstrates the activation of NtJAZ1 and NtJAZ3 promoters

by PAP1, TT8, PAP1/TT8, and PAP1/TT8/TTG1

Dual-luciferase assay was completed to demonstrate whether PAP1 alone, TT8 alone,

PAP1 and TT8 together, and PAP1/TT8/TTG1 (WD40) together could activate the

NtJAZ1 and *NtJAZ3* promoters in tobacco. PAP1, TT8, and TTG1 were used as

effectors. The ORF sequences of *PAP1*, *TT8*, *TTG1*, and *GFP* were respectively

cloned into the PCB2004 plasmid to generate four constructs, in which each gene was

driven by a 35S-promoter (Fig. 2 D). The *NtJAZ1pro* and *NtJAZ3pro* were cloned into

the plasmid pGreenII-0800 to drive the luciferase gene to obtain *NtJAZ1pro*-luciferase

and *NtJAZ3pro*-luciferase reporters (Fig. 2 D). Effectors and reporters were incubated

for dual-luciferase assays. Promoters alone and a binary vector were used as controls.

The resulting data showed that compared with controls, PAP1 and TT8 alone

enhanced the activity of *NtJAZ1pro* 3.8 and 7.2 times, respectively, and increased the

activity of *NtJAZ3pro* 1.9 (slightly) and 2.1 (significantly) times, respectively (Fig. 2

E). PAP1 and TT8 together enhanced the activity of *NtJAZ1pro* and *NtJAZ3pro*

approximately 4.0 and 6.3 times (Fig. 2 E). Furthermore, PAP1, TT8, and TTG1

together increased 12.7 and 15.6 times of the activity of both *NtJAZ1pro* and

NtJAZ3pro, much higher than PAP1 alone, TT8 alone, and PAP1 and TT8 together

(Fig. 2 E). These results demonstrate that PAP1 and TT8 alone, PAP1 and TT8

together, PAP1-TT8-TTG1 complex transcriptionally activate both *NtJAZ1* and

NtJAZ3.

521
PAP1, TT8, PAP1/TT8, and PAP1/TT8/TTG1 do not activate promoters of ODC2 and

PMT of tobacco

NtODC2 and *NtPMT2* are two key upstream pathway genes of the nicotine

biosynthesis. We obtained 1.0 kb long sequences for these two genes' promoters from

the tobacco genome database curated at NCBI. The promoter sequence of *NtPMT2*

has an MRE-like element (AACAACC at position -530) and a G-box like (CACGTT

at position -186) element, while the sequences of the *NtODC2* promoter does not have

MRE and G-box elements (Fig. S4).

EMSA experiments were performed to test whether PAP1 and TT8 could bind to

MRE-like and G-box-like elements. The EMSA results indicated that the TT8 binding

domain could bind to the G-box-like (CACGTT) element. However, when a

competitive probe was added, the binding signal was lost (Fig. S5 A). Furthermore,

compared with the positive control G-box elements from *NtJAZ1* and *NtJAZ3*

promoters, the resulting data showed that the signal from G-box like element was

much weaker (Fig. S5 A). This result indicates that TT8 can weakly bind to the

promoter of *NtPMT2*. When the MRE-like element probe was prepared for EMSA,

the resulting data showed that PAP1 could not bind to the MRE-like probe

(AACAACC), while PAP1 strongly bound to the positive control MRE1 and MRE3

probes of *NtJAZ1* and *NtJAZ3* promoters (Fig. S5 B).

Dual-luciferase assay and ChIP-Q-PCR were performed to test whether PAP1 and

TT8 could activate the *ODC2* and *PMT2* promoters. The dual-luciferase assay data

did not show that PAP1 alone, TT8 alone, PAP1/TT8 together, and the

PAP1/TT8/TTG1 complex could activate the activity of the two promoters (Fig. S6

A). Meanwhile, the ChIP-Q-PCR data showed that PAP1 and TT8 could not enrich

binding elements of the two promoters *in vivo* (Fig. S6 B). These data indicate that

PAP1, TT8, PAP1/TT8, and PAP1/TT8/TTG1 cannot activate the promoters of these

two key pathway genes.

Engineering of red genotypic P+T-NL and P+T-KY tobacco plants

An expression cassette was synthesized to stack the ORF sequences of *PAP1* (P) and

TT8 (T) and then used to generate 20 T0 transgenic plants from each of Narrow Leaf

Madole (NL) and KY171 (KY), two major commercial varieties (Fig. S7 A, B, C, E,

and G-H). In addition, vector control transgenic plants were generated (Fig. S7 D and

F). The first 12 red T0 transgenic plants from the two cultivars were tagged as P+T-

NL1 to 12 and P+T-KY1 to 12, respectively. The first 12 T0 vector control transgenic

plants from the two cultivars were tagged as NL1 to 12 and KY1 to 12, respectively.

These T0 plants were selected to obtain seeds to screen T1 and homozygous T2

progeny (Fig. S7 I-M), during which the copy number of the transgenes was

determined. Five P+T-NL lines (1-5) and five P+T-KY lines (1-5) were confirmed to

contain one single copy of the stacked two transgenes. Quantitative RT-PCR showed

that homozygous T2 progeny highly expressed *PAP1* and *TT8* (Fig. S8). Furthermore,

qRT-PCR was performed to understand the expression of *CHS, CHI, F3H, F3'H,*

DFR, and *ANS* involved in the biosynthesis of both anthocyanin and proanthocyanidin

(PA) and the expression of *3-GT* involved in the glycosylation of anthocyanidins (Fig.

1 A) in T2 seedlings (Fig. S9). The resulting data showed that the expression levels of

these seven genes were up-regulated 2-1000 times (Fig. S10 AB) in pink roots and red

leaves of the P+T-NL and P+T-KY genotypes. Further UV spectrometry-based

measurements showed that the roots and leaves of analyzed P+TNL1 and P+T-KY1

genotypes produced anthocyanins, while those of control plants did not (Fig. S11). It

was noticeable that the levels of anthocyanins were significantly higher in leaves than

in roots of both P+T-NL1 and P+T-KY1 plants (Fig. S11). To understand this

differentiation, the expression of *ANR* specifically involved in the colorless PA

formation [71] was analyzed. The resulting data showed that the expression of *ANR*

was significantly activated in roots but not in leaves of the two red genotypes (Fig.

S10). To substantiate this, root extracts of both P+T-NL1 and P+T-KY1 seedlings

were treated with DMACA. A bluish color was immediately developed from the

reaction (Fig. S12 A-B), showing the presence of PAs. However, this bluish color was

not developed from the extracts of both P+T-NL1 and P+T-KY1 leaves and wild type

roots and leaves, indicating the lack of PAs. Further measurement of DMACA-treated

samples at 640 nm showed that the ABS values were 5-6 times higher in the root

extracts than in the leaf extracts of P+T-NL1 and P+T-KY1 and in the leaf and root

extracts of wild type (Fig. S12 C-D). These data showed that the less red pigmentation

in roots resulted from the conversion of red anthocyanidins to colorless PAs by ANR.

In addition, the less red pigmentation in roots might result from the lack of light in the

hydroponic condition and secretion of anthocyanins.

Upregulation of NtJAZ1, 3, 7, and 10 and downregulation of the nicotine biosynthesis

in red P+T-NL and P+T-KY plants

Both transcriptional and metabolic profiling were performed to characterize the

biosynthesis of nicotine in homozygous T2 seedlings (Fig. S9) of different genotypes

selected, including P+T-NL1, P+T-NL2, P+T-NL3, P+T-KY1, P+T- KY 2, and P+T-

KY3. Quantitative RT-PCR was performed to characterize the transcriptional profiles

of 16 genes (Fig. 1 B) involved in the nicotine biosynthesis. The expression levels of

NtJAZ1, NtJAZ3, *NtJAZ7,* and *NtJAZ10* were significantly increased in roots of both

P+T-NL and P+T-KY genotypes (Fig. 3 A-B and K-L, Fig. S13). The expression

levels of *PMT1*, *PMT2*, *MPO*, *QPT2*, *A622*, *BBL* and *ERF189* were significantly

reduced in roots of both P+T-NL (Fig. 3 D-J) and P+T-KY plants (Fig. 3 N-T). The

expression levels of *OCD2* was not altered in the P+T-NL1, 2, and 3 lines (Fig. 3 C)

but significantly decreased in the P+T-KY1, 2 and 3 lines (Fig. 3 M), indicating the

different responses of this gene expression in two commercial varieties to transgenes.

The expression levels of *MYC1a*, *MYC1b*, *MYC2a*, and *MYC2b* were not altered in

roots of both P+T-NL and P+T-KY plants (Fig. S14). HPLC-MS analysis showed that

the contents of nicotine, nornicotine, and the total amount of the two compounds in

roots and leaves of P+K-NL1, P+K-NL2, P+K-NL3, P+K-KY1, P+K-KY2, and P+K-

KY3 plants were significantly reduced 1.5-3.4, 1.3-2.5, and 1.9-3.3 fold, respectively

(Fig. 4).

Field trials and reduction of nicotine, nornicotine, anatabine, anabasine, myosmine,

and four TSNAs in red P+T-NL1 and P+T-KY1 tobacco plants

Field trials that follow the commercial production practices of tobacco are essential to

examine the effects of new varieties or genes on the contents of nicotine, other

tobacco alkaloids, and TSNAs. In addition, field farming practices were necessary to

test the agricultural significance of this *de novo* regulation design. The field practice

exactly followed the protocols of both industry tobacco and GMO tobacco growth.

We planted T2 homozygous P+T-NL1 and P+T-KY1 progeny and their

corresponding wild-type tobacco variety in the field at the research station in Oxford,

North Carolina (Fig. 5 A-C and Figs S15-S18). Tobacco leaves were air-cured in a

contained ventilation barn (Fig. 5 D, Figs S18-S19).

Quantification with GC-FID showed the significant reduction of nicotine, nornicotine,

anabasine, anatabine, and myosmine in most or all cured leaves of red P+T-NL1 and

P+T-KY1 tobacco plants compared to those of non-transgenic controls. The contents

of nicotine were significantly reduced by 45-51% in all cured leaf groups of P+T-NL1

(Fig. 6 A) and by 19-30% in upper three groups of cured leaves of P+T-KY1 (Fig. 6

B). The contents of nornicotine were significantly decreased by 39-44% and 30-40%

in all cured leaf groups of T-P-NL1 and P+T-KY1, respectively (Fig. 6 C and D). The

contents of anabasine, anatabine, and myosmine were significantly reduced by 27-

45%, 55-66%, and 19-25% in all cured leaf groups of P+T-NL1 (Fig. 6 E, G, and I),

respectively, and were significantly reduced by 27-45%, 3256%, and 29-35% in all

cured leaf groups of P+T-KY1, respectively (Fig. 6 F, H, and J).

Quantification with HPLC-QQQ-MS/MS showed that the contents of four

carcinogenic TSNAs were significantly diminished in most or all leaf groups of

P+TNL1 and P+T-KY1 plants. The content of NNN was significantly decreased by

63-74% to a level of 0.136-0.375 ppm in all four groups of cured P+T-NL1 leaves and

by 67-79% to a level of 0.208-0.552 ppm in three groups of cured P+T-KY1 leaves

(Fig. 7 A and B). The content of NNN in the B3 leaf group of P+T-KY1 was reduced

by 63% to 0.488 ppm with a slight significance. The content of NNK was reduced by

38-72% to a level of 0.1-0.173 ppm and by 30-74% to a level of 0.151-0.3 ppm in

four groups of cured P+T-NL1 and P+T-KY1 leaves, respectively, in which the

reduction in three leaf groups of each genotype was significant (Fig. 7 C and D). The

content of NAT was significantly decreased by 77-92% to a level of 0.093-0.159 ppm

and by 70-80% to a level of 0.159-0.285 ppm in all leaf groups of P+T-NL1 and P+T-

KY1, respectively (Fig. 7 E and F). The content of NAB was significantly decreased

by 72-80% to a level of 0.011-0.021 ppm and by 65-77% to a level of 0.015-0.031

ppm in all leaf groups of P+T-NL1 and P+T-KY1, respectively (Fig. 7 G and H).

Furthermore, the contents of NNN, NNK, NAT, and NAB were summed to obtain the

total TSNA contents in each leaf group. The total contents of four TSNAs in each

group of wild type NL and KY171 leaves were 1.6-2.2 ppm and 2.2-2.6 ppm, while

these values were significantly reduced to be 0.4-0.6 ppm and 0.6-1.0 ppm in each

group of P+T-NL1 and P+T-KY1 leaves (Fig. 7 I-J). The average contents of four

TSNAs in the entire plant leaves of wild type NL and KY171 were approximately

1.77 ppm and 2.25 ppm, while these values were significantly reduced to be 0.69 ppm

and 0.87 ppm in those of P+T-NL1 and P+T-KY1 (Fig. 7 K-L).

Furthermore, the total contents of tobacco alkaloids in each group of leaves were

summed from nicotine, four other alkaloids, and total TSNAs. The resulting data

showed that the contents of total alkaloids were significantly decreased by 44-51% in

four leaf groups of P+TNL1 and 22-35% in the upper three group leaves of P+T-KY1

(Fig. S20 A-B).

Field trials and reduction of nicotine, nornicotine, anatabine, anabasine, myosmine,

and four TSNAs in flue-cured leaves of the red PAP1 tobacco

Our red PAP1 tobacco genotype (Fig. S22 B) is a homozygous red variety generated

from Xanthi by the overexpression of *PAP1* [21]. Gene expression analysis showed

that the expression of *PAP1* upregulated *NtAn1a* and *NtAn1b* (Fig. S21), two *TT8*

homologs [72]. This result indicated that *PAP1* alone activated tobacco anthocyanin

biosynthesis via upregulating the endogenous *N. tabacum TT8* (*NtTT8*) homologs.

Therefore, to substantiate our *de novo* design, we used this genotype to test whether

PAP1 and activated NtTT8 homologs could reduce nicotine, other tobacco alkaloids,

and TSNAs.

Two years' field trials were completed. The field design (Fig. S22 A-B) and cropping

management were the same as described above. When leaves were harvested, they

were harvested into five groups (Fig. S22 C-D) and then flue-cured in barns (Fig. S22

E). Analysis with HPLC-ESI-MS obtained peak values that showed the reduction of

nicotine, nornicotine, anabasine, and anatabine. The levels of nicotine were

significantly or slightly reduced by 10-50% and 5-25% in each group of leaves in

2011 and in 2012, respectively (Fig. S23 A-B). The levels of nornicotine, anabasine,

and anatabine were significantly or slightly reduced in all or most group leaves by 25-

70% and 5-55%, 30-60% and 5-20%, and 5-60% and 10-40% in 2011 and 2012,

respectively (Fig. S23 A-B). In addition, the total cumulative peak level of each

alkaloid was summed from all leaf groups of the entire PAP1 tobacco plant. The

resulting data showed that the total cumulative levels of nicotine, nornicotine,

anabasine, and anatabine in the entire plant were significantly reduced by 30%, 45%,

50%, and 40% in 2011 and 10%, 20%, 10%, and 30% in 2012 respectively (Fig. S24

A-B).

The contents of NNN, NNK, NAB, and NAT were reduced in flue-cured leaf groups

(Fig. S25 A-B). The contents of NNN were reduced by 20-60% and 25-60% in all leaf

groups in two years. The average content of NNN was diminished to a level less than

0.5 ppm in most leaf groups (Fig. S25 A-B). The contents of NNK were reduced by 5-

50% in all leaf groups in 2011 and 5-60% in four of five leaf groups in 2012. The

contents of NAB were reduced by 20-70% and 15-50% in all leaf groups in two years.

The contents of NAT were reduced by 50-65-% and 35-70% in all leaf groups in two

years. The contents of each NNN, NNK, NAB, and NAT were averaged for all leaf

groups of each plant. The resulting data showed that the average contents of each

TSNA were significantly reduced by 25-60% (P-value < 0.05) in 2011 and 2012 (Fig.

S25 A-B). The contents of four TSNAs were summed for all leaf groups of each

plant. The resulting data showed that the total contents of four TSNAs were

significantly reduced by 57% and 43% (P-value < 0.05) in 2011 and 2012,

respectively (Fig. S26).

Discussion

Our findings indicate that cis-regulatory elements and TFs are useful molecular tools

for a *de novo* design to create a novel regulation of plant secondary metabolism.

Herein, based on the regulation mechanism elucidated (Fig. 2), we constructively term

the designed regulation to be a *Distant Pathway-Cross Regulation* (DPCR) of PAP1

and TT8. The rationale is that the pathway of nicotine does not exist in Arabidopsis

and two anthocyanin TFs are metabolically characterized to be distant from the

tobacco alkaloid pathway. The other reason is that our findings are distinct from the

general pleotropic effects that have been generally used to explain gene functions

without a mechanistic elucidation. This DPCR definition is characterized by the

mechanisms that PAP1, TT8, and their complex formed with WD40 negatively

regulate the tobacco alkaloid biosynthesis via activating *NtJAZ* repressor genes.

Accordingly, this definition distinguishes from the master regulatory function of

PAP1, TT8, and their MBW complex that activates the anthocyanin biosynthesis in

Arabidopsis [8, 14, 73, 74] or red tobacco shown here via directly activating most or

late pathway genes, such as *DFR*, *ANS*, and *3-GT* (Fig. S10). Based on these data, we

propose a model to characterize the DPCR created from the *de novo* design and the

regulation of tobacco alkaloid biosynthesis in red tobacco plants (Fig. 8). Given that

NtTTG1 is expressed in roots, leaves, and flowers, this model proposes that PAP1 and

TT8 likely anchor to the endogenous tobacco NtTTG1 (WD40) to form a

heterogeneous MBW complex, which has a dual regulatory function that regulates

tobacco metabolism. First, the MBW complex directly activates the anthocyanin

biosynthesis (function I), which is intensively elucidated. Second, the MBW complex

upregulates four *NtJAZ1*, *NtJAZ3*, *NtJAZ7*, and *NtJAZ10*, which increase NtJAZ

repressors (function II). Next, the increased NtJAZs bind to more MYC2 and block

the release of this TF. Then, the reduction of free MYC2 results in downregulating

EFR and six or seven nicotine pathway genes (Fig. 3). Finally, the biosynthesis of

nicotine, OTAs, and four TSNAs is significantly diminished in red tobacco plants

(Figs. 4, 6, and 7). In addition, based on the report that AtMYC2 solely regulates the

AtJAZ transcription via a feed-back regulation [75], this model proposes that

PAP1/TT8 and NtMYC2 co-activate the expression of NtJAZs in red tobacco plants.

When JA is absent, PAP1 and TT8 play the main regulatory role to activate the

expression of four *NtJAZs*. To date, in addition to tobacco, PAP1 and its homologs

have been introduced to a few crops to engineer anthocyanins for value-added traits,

such as tomato [22, 23], hop [24], canola [25], and rose [76]. Although whether and

how PAP1 and TT8 can regulate non-anthocyanin or non-flavonoid pathways in these

plants remains unknown, it was interesting that PAP1 transgenic rose plants were

reported to produce more than 6.5-time scent terpenoids [76]. Based on our findings,

it is interesting to investigate whether a similar DPCR occurs to increase scent

terpenoids in rose in the future. Moreover, different TF families are a large part of

plant genomes [77-80], our findings suggest that it is necessary to investigate DPCR

events of other TFs to fully understand their regulatory functions. In particular, as

shown in our data that carcinogenic TSNAs are significantly reduced in red tobacco

plants, investigations of potential DPCR of TFs likely create value-added traits for

crop improvement and human health.

This *de novo* regulation design with PAP1/TT8 TFs and *NtJAZs* shows a promising

application in enhancing efforts for the simultaneous reduction of most harmful

tobacco alkaloids. As introduced above, in the past decades, multiple previous studies

had been undertaken to develop technologies to reduce harmful tobacco alkaloids,

such as gene suppression or silencing technologies with anti-sense or RNAi of

pathway genes (Fig. 1 B), including *PMT* [65], *ODC* [64], *QPT* [63], *NDM* [81],

A622 [82], *BBL* [49], and others [42]. On the one hand, most studies successfully

reported the decrease of the contents of nicotine, OTAs, or TSNAs in tobacco plants

grown in the greenhouse, growth chambers, or field. On the other hand, these

previous successes revealed a challenge that all tobacco alkaloids could not be

diminished simultaneously. For example, the decrease of nicotine led to either the

increase of anatabine or non-reduction of other tobacco alkaloids [64, 65]. In addition,

the reduction of harmful compounds was relatively limited. For instance, an RNAi of

A622 could mainly reduce NNN and total TSNAs [81]. A gene silencing of *BBL* was

reported to mainly decrease the nicotine content [49]. Furthermore, all these successes

have showed that the reduction of nicotine and other alkaloids to a practical level

requires more studies, such as elucidation of metabolic regulation and unknown

genes. As intensively characterized, the regulation of the nicotine biosynthesis is via

three cohort layers, the presence/absence of an active JA form, the JA and JAZ

signaling pathway, and the release of MYC2 for activation of ERF189 and pathway

genes [42, 48, 55] (Fig. 1 B). To date, an important unanswered question is how

tobacco plants regulate *NtJAZs*. In Arabidopsis, the expression of *AtJAZs* is solely

activated by AtMYC2 via a feed-back regulation [75]. Based on this, although

NtMYC2 members have not been reported to regulate the transcription of *NtJAZs* in

tobacco plants, it can be postulated that NtMYC2 members may also perform a feed-

back regulation of *NtJAZs* (Fig. 8). Our *de novo* design with PAP1 and TT8 and MRE

and G-box elements was used to target four *NtJAZs*. As designed, PAP1, TT8, PAP1

and TT8 together, and their MBW complex bound and activated promoters of the four

NtJAZs (Fig. 2) and the stacked overexpression of *PAP1* and *TT8* led to the

upregulation of four *NtJAZs* (Fig. 3 A-B and K-L, Fig. S13). As the positive result,

the biosynthesis of nicotine was downregulated in red P+T genotypic plants (Figs. 3

and 4). The contents of nicotine, four OTAs, and four TSNAs were simultaneously

reduced in most or all cured leaf groups and entire plant leaves of P-T-NL, P-T-KY,

and PAP1 genotypes (Figs. 6 and 7, Figs. S20). The NNN contents were particularly

diminished to a level less than 0.55 ppm in most of red tobacco leaves (Fig. 7 A-B).

This low content is significant to human health, because NNN is a severe tobacco

carcinogen and its content in any smokeless tobacco products is limited to 1.00 pm

proposed by FDA. In addition, other TSNAs were reduced to a level lower than 0.5

ppm in most or all leaves of red P+T genotypic tobacco plants. To further substantiate

the function of this *de novo* regulation design, we tested our PAP1 red tobacco

variety, in which the expression of *PAP1* activated two *NtTT8* homologs (Fig. S21).

Although this variety was created from Xanthi, a low-nicotine and flue-cured cultivar,

the two years' field trials showed the reduction of nicotine, OTAs, and TSNAs in

PAP1 plants (Figs. S23-24 and S25-26). These data indicate that PAP1 can partner

with the endogenous tobacco NtTT8 homologs to form a DPCR to downregulate the

nicotine biosynthesis. Taken together, these findings provide a new platform to

eliminate tobacco carcinogens to a non-harmful level and to alter plant metabolism in

other crop plants for novel traits.

The plant kingdom has a large number of TFs that regulate plant development,

structure, responses to stresses, and plant metabolisms. In 2000, when the genome of

Arabidopsis thaliana was sequenced [83], gene annotation and sequence analysis

revealed more than 1500 TFs (accounting for more than 5% of annotated genes).

Since then, more TFs have been reported from numerous sequenced plant genomes,

such as rice [84, 85]and soybean[86], and multiple browser servers have been

developed to search TF families and their regulatory functions [87-93]. By 2017, 156

plant genomes were sequenced, including 16 from Chlorophytae, one from

Charophyta, one from Marchantiophyta, two from Byrophyta, one from

Lycopodiophyta, two from Coniferophyta, one from Basa Magnoliophyta, 38 from

monocots, and 95 Eudicots. [67]. These 156 genomes together with sequences of nine

additional plant species allowed the annotation of 320,370 plant TFs, which were grouped into 58 families [67]. Likewise, the plant genomes have a large number of binding sites (motifs) (BS), such as G-box and MRE in promoters, to which TFs bind. In recent, a public PlantRegMap server was developed from 63 plant species (out of 156 genome-sequenced plants) to search TFBSs and the interactions between TF and BS [68]. This PlantRegMap characterized a conserved regulatory landscape that included 21,997,501 TFBSs, 21,346 TFs, and more than two million interactions between them [68]. It can be speculated that as more plant species will be sequences, the number of plant TFs will be increased. To our knowledge, none of TFs and BSs has been used as molecular tools for a novel regulatory design to engineer better valuable traits of crops. Based on our first proof-of-concept study with PAP1 vs. MRE and TT8 vs. G-box that are effective molecular tools to design novel regulations to create a DPCR in tobacco plant, it can be speculated that the large number of plant TFs and BSs provide an extremely rich molecular tool source for novel regulatory designs, which will be fundamental for plant engineering.

Conclusion

The findings show that PAP1, TT8, MREs, and G-box elements are useful molecular tools to design a *De Novo* regulation, a distant pathway-cross regulation (DPCR) of plant secondary metabolism in plants. The findings unearth novel regulatory functions of PAP1 and TT8 that are two positive regulators of *NtJAZs* and negatively regulate tobacco alkaloid biosynthesis in red tobacco plants. This *De Novo* regulation design significantly reduces harmful nicotine, other tobacco alkaloids, and all TSNAs in all

or most leaves of tobacco plants. <mark>Our findings indicate that the great number of TFs</mark>

<mark>and BSs provide an extremely rich molecular tool source for novel regulatory designs,</mark>

<mark>which will be fundamental in plant engineering for value-added crops and products.</mark>

Conflict of Interest

Authors declare no conflict of interest.

Supporting information

Supporting materials and growth conditions, Method S1-S8, 26 supporting figures and

9 supporting tables.

26 Supporting Figure legends

Figure S1 Promoter sequences of *NtJAZ1* and *NtJAZ3* and identification of cis-

elements

Figure S2 Promoter sequences of *NtJAZ7* and *NtJAZ10* and identification of cis-

elements

Figure S3 Images showing expression and purification of the binding domains of

PAP1 and TT8 induced in *E. coli in vitro*

Figure S4 Promoter sequences of *NtPMT2* and *NtODC* and identification of cis-

elements.

Figure S5 Electrophoretic mobility shift assays (EMSAs) showing weak binding of

TT8 to G-box-like probe and no binding of PAP1 to MRE-like probe.

Figure S6 Dual-luciferase and Chip-qPCR experiments showing no activation of

NtPMT2 and *NtODC2* promoters bound by PAP1 and TT8 alone, PAP1 and TT8

together, and PAP1-TT8-WD complex.

Figure S7 Diagrams showing gene stacking design for DNA synthesis, binary

construct, and selection of T2 homozygous plants.

Figure S8 Comparison of *AtPAP1* and *AtTT8* transgene expressions in roots of wild-

type Narrow Leaf Madole (NL) and KY171, P+T transgenic NL and KY171, and

vector control transgenic plants.

Figure S9 Phenotypes of wild type, vector-control, and P+T-NL and P+T-KY tobacco

plants used for analysis of gene expression and metabolites.

Figure S10 Transcriptional upregulation of eight flavonoid pathway genes in red

transgenic P+T-NL and P+T-KY tobacco plants.

Figure S11 Anthocyanin levels in roots and leaves of wild type and red P+T

transgenic tobacco plants.

Figure S12 Proanthocyanidin formation in roots of red P+T transgenic tobacco plants

Figure S13 Comparison of *NtJAZ7* and *NtJAZ10* gene expression in roots of wild-type

Narrow Leaf Madole (NL) and KY171, P+T transgenic NL and KY171, and vector

control transgenic plants.

Figure S14 Comparison of *MYC1a*, *MYC1b, MYC2a* and *MYC2b* gene expression in

roots of wild-type (WT) Narrow Leaf Madole (NL) and KY171, P+T transgenic NL

and KY, and vector control transgenic plants.

Figure S15 Schematic diagram showing procedures for field farming practice and leaf

harvest from wild type and red transgenic tobacco plants.

Figure S16 Field design for farming practice of red P+T transgenic plants.

Figure S17 Planting, plant growth, and phenotypes of wild type and red P+T

transgenic tobacco plants in field from the first day to 75 days after planting.

Figure S18 Harvest of plants from field for air-curing and cleanup of field.

Figure S19 Comparison of leaf color after air-curing and grouping of leaves for

sampling.

Figure S20 Reduction of total tobacco alkaloids in each leaf group of red P+T-NL1

and P+T-KY1 plants compared to wild type ones.

Figure S21 Upregulation of *NtAn1a* and *NtAn1b* in roots of red PAP1 tobacco plants.

Figure S22 Field farming practice of PAP1 tobacco plants and leaf harvest.

Figure S23 Reduction of nicotine, nornicotine, anabasine, and anatabine in PAP1

tobacco leaves.

Figure S24 Reduction of total nicotine, nornicotine, anabasine, and anatabine in all

leaves of PAP1 tobacco plant.

Figure S25 Reduction of NNN, NNK, NAB, and NAT in flue-cured leaves of PAP1

tobacco compared to leaves of wild type Xanthi control.

Figure S26 Reduction of total TSNAs in flue-cured leaves of PAP1 tobacco.

Nine Supporting Table capitals

Table S1 Thermal cycle and sequences of primers designed for PCR, qRT-PCR,

QPCR, respectively.

Table S2 Float trays used for seedlings in the water nursery bed in the greenhouse.

The same trays were prepared for PAP1 tobacco plants.

Table S3 Instrument Parameters used in Tobacco Specific Nitrosamines analysis.

Table S4 HPLC Gradient used in Tobacco Specific Nitrosamines analysis.

Table S5 Typical Parameter Table Triple Quadrupole Mass Spectrometer used in

Tobacco Specific Nitrosamines analysis.

Table S6 Reagents and Standards in for Tobacco Specific Nitrosamines analysis.

Table S7 Instrument Parameters used in Tobacco Alkaloids analysis.

Table S8 Oven Program used for in Tobacco Alkaloids analysis.

Table S9 Reagents and Standards used for Tobacco Alkaloids analysis.

References

[1] H. Du, B.R. Feng, S.S. Yang, Y.B. Huang, Y.X. Tang, The R2R3-MYB Transcription Factor Gene Family in Maize, PLoS One 7(6) (2012) 12.

[2] C. Dubos, R. Stracke, E. Grotewold, B. Weisshaar, C. Martin, L. Lepiniec, MYB transcription factors in Arabidopsis, Trends Plant Sci 15(10) (2010) 573-581.

[3] A. Feller, K. Machemer, E.L. Braun, E. Grotewold, Evolutionary and comparative analysis of MYB and bHLH plant transcription factors, Plant J. 66(1) (2011) 94-116.

[4] K. Wei, H. Chen, Comparative functional genomics analysis of bHLH gene family in rice, maize and wheat, BMC Plant Biol. 18(1) (2018) 309.

[5] T. Saigo, T. Wang, M. Watanabe, T. Tohge, Diversity of anthocyanin and proanthocyanin biosynthesis in land plants, Curr. Opin. Plant Biol. 55 (2020) 93-99.

[6] M.-Z. Shi, D.-Y. Xie, Biosynthesis and metabolic engineering of anthocyanins in *Arabidopsis thaliana*, Recent Pat Biotechnol 8(1) (2014) 47-60.

[7] K. Springob, H. Nakajima, M. Yamazaki, K. Saito, Recent advances in the biosynthesis and accumulation of anthocyanins, Nat Prod Rep 20 (2003) 288-303.

[8] J.O. Borevitz, Y. Xia, J. Blount, R.A. Dixon, C. Lamb, Activation tagging identifies a conserved MYB regulator of phenylpropanoid biosynthesis, Plant Cell 12(12) (2000) 2383-2394.

[9] N. Nesi, I. Debeaujon, C. Jond, G. Pelletier, M. Caboche, L. Lepiniec, The *TT8* gene encodes a basic helix-loop-helix domain protein required for expression of *DFR* and *BAN* genes in *Arabidopsis* siliques, The Plant Cell 12 (2000) 1863-1878.

[10] A.R. Walker, P.A. Davison, A.C. Bolognesi-Winfield, C.M. James, N. Srinivasdan, T.L. Blundell, J.J. Esch, M.D. Marks, J.C. Gray, The *TRANSPARENT TESTA GLABRA 1* locus, which regulates trichome differentiation and anthocyanin biosynthesis in Arabidopsis, encodes a WD40 repeat protein, Plant Cell 11 (1999) 1377-1349.

[11] W. Xu, D. Grain, S. Bobet, J. Le Gourrierec, J. Thévenin, Z. Kelemen, L. Lepiniec, C. Dubos, Complexity and robustness of the flavonoid transcriptional regulatory network revealed by comprehensive analyses of MYB–bHLH–WDR complexes and their targets in Arabidopsis seed, New Phytol. 202(1) (2014) 132-144.

[12] J. Thévenin, C. Dubos, W. Xu, J. Le Gourrierec, Z. Kelemen, F. Charlot, F. Nogué, L. Lepiniec, B. Dubreucq, A new system for fast and quantitative analysis of heterologous gene expression in plants, New Phytol. 193(2) (2012) 504-512.

[13] W.J. Xu, D. Grain, J. Le Gourrierec, E. Harscoet, A. Berger, V. Jauvion, A. Scagnelli, N. Berger, P. Bidzinski, Z. Kelemen, F. Salsac, A. Baudry, J.M. Routaboul, L. Lepiniec, C. Dubos, Regulation of flavonoid biosynthesis involves an unexpected complex transcriptional regulation of TT8 expression, in Arabidopsis, New Phytol. 198(1) (2013) 59-70.

[14] Z.X. Zhu, H.L. Wang, Y.T. Wang, S. Guan, F. Wang, J.Y. Tang, R.J. Zhang, L.L. Xie, Y.Q. Lu, Characterization of the cis elements in the proximal promoter regions of the anthocyanin pathway genes reveals a common regulatory logic that governs pathway regulation, J. Exp. Bot. 66(13) (2015) 3775-3789.

[15] A. Baudry, M.A. Heim, B. Dubreucq, M. Caboche, B. Weisshaar, L. Lepiniec, TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and proanthocyanidin biosynthesis in *Arabidopsis thaliana*, Plant J 39(3) (2004) 366-380.

[16] N.A. Ramsay, B.J. Glover, MYB-bHLH-WD40 protein complex and the
evolution of cellular diversity, Trends Plant Sci 10(2) (2005) 63-70.
[17] M.Z. Shi, D.Y. Xie, Engineering of red cells of *Arabidopsis thaliana* and
comparative genome-wide gene expression analysis of red cells versus wild-type
cells, Planta 233(4) (2011) 787-805.
[18] T. Qi, S. Song, Q. Ren, D. Wu, H. Huang, Y. Chen, M. Fan, W. Peng, C. Ren, D.
Xie, The jasmonate-ZIM-domain proteins interact with the WD-Repeat/bHLH/MYB
complexes to regulate jasmonate-mediated anthocyanin accumulation and trichome
initiation in *Arabidopsis thaliana*, Plant Cell 23(5) (2011) 1795-1814.
[19] T.C. Qi, H. Huang, D.W. Wu, J.B. Yan, Y.J. Qi, S.S. Song, D.X. Xie,
Arabidopsis DELLA and JAZ proteins bind the WD-Repeat/bHLH/MYB complex to
modulate gibberellin and jasmonate signaling synergy, Plant Cell 26(3) (2014) 1118-
1133.
[20] X. He, Y. Li, D. Lawson, D.-Y. Xie, Metabolic engineering of anthocyanins in
dark tobacco varieties Physiol Plant 159 (2017) 2-12.
[21] D.-Y. Xie, S.B. Sharma, E. Wright, Z.-Y. Wang, R.A. Dixon, Metabolic
engineering of proanthocyanidins through co-expression of anthocyanidin reductase
and the PAP1 MYB transcription factor, Plant J 45(6) (2006) 895-907.
[22] E. Butelli, L. Titta, M. Giorgio, H.-P. Mock, A. Matros, S. Peterek, E.G.W.M.
Schijlen, R.D. Hall, A.G. Bovy, J. Luo, C. Martin, Enrichment of tomato fruit with
health-promoting anthocyanins by expression of select transcription factors, Nat
Biotech 26(11) (2008) 1301-1308.
[23] D.L. Zuluaga, S. Gonzali, E. Loreti, C. Pucciariello, E.D. Degl'Innocenti, L.
Guidi, A. Alpi, P. Perata, Arabidopsis thaliana MYB75/PAP1 transcription factor
induces anthocyanin production in transgenic tomato plants, Funct Plant Biol 35(7)
(2008) 606-618.
[24] A. Gatica-Arias, M.A. Farag, M. Stanke, J. Matousek, L. Wessjohann, G. Weber,
Flavonoid production in transgenic hop (*Humulus lupulus* L.) altered by
PAP1/MYB75 from *Arabidopsis thaliana* L, Plant Cell Rep 31(1) (2012) 111-119.
[25] X. Li, M.J. Gao, H.Y. Pan, D.J. Cui, M.Y. Gruber, Purple canola: Arabidopsis
PAP1 increases antioxidants and phenolics in *Brassica napus* leaves, J Agric Food
Chem 58(3) (2010) 1639-1645.
[26] D.M. Ma, S.V.S. Gandra, R. Manoharlal, C. La Hovary, D.Y. Xie, Untargeted
metabolomics of *Nicotiana tabacum* grown in United States and India characterizes
the association of plant metabolomes wiith natural climate and geography, Front.
Plant Sci. 10 (2019) 18.
[27] L.H. Nugroho, R. Verpoorte, Secondary metabolism in tobacco, Plant Cell,
Tissue and Organ Culture 68(2) (2002) 105-125.
[28] A. Rodgman, T.A. Perfetti, The Chemical Components of Tobacco and Tobacco
Smoke, CRC Press, Taylor & Francis Group, Boca Raton, London, New York, 2009.
[29] I. Arany, M. Taylor, T. Fulop, M. Dixit, Adverse effects of chronic nicotine
exposure on the kidney: Potential human health implications of experimental findings,
International Journal of Clinical Pharmacology and Therapeutics 56(11) (2018) 501-
506.
[30] S.A. Grando, Connections of nicotine to cancer, Nature Reviews Cancer 14(6)
(2014) 419-429.

[31] L. Greillier, A.B. Cortot, J. Viguier, L. Brignoli-Guibaudet, C. Lhomel, F.
Eisinger, J.F. Morere, S. Couraud, Perception of Lung Cancer Risk: Impact of
Smoking Status and Nicotine Dependence, Current Oncology Reports 20 (2018) 7.
[32] P.N. Lee, B.A. Forey, J.S. Hamling, A.J. Thornton, Environmental tobacco
smoke exposure and heart disease: A systematic review, World Journal of Meta-
Analysis 5(2) (2017) 14-40.
[33] T. Sanner, T.K. Grimsrud, Nicotine: carcinogenicity and effects on response to
cancer treatment - a review, Frontiers in Oncology 5 (2015) 10.
[34] A. Santoro, C. Tomino, G. Prinzi, P. Lamonaca, V. Cardaci, M. Fini, P. Russo,
Tobacco Smoking: Risk to Develop Addiction, Chronic Obstructive Pulmonary
Disease, and Lung Cancer, Recent Patents on Anti-Cancer Drug Discovery 14(1)
(2019) 39-52.
[35] A.N. Thorndike, N.A. Rigotti, A tragic triad: coronary artery disease, nicotine
addiction, and depression, Current Opinion in Cardiology 24(5) (2009) 447-453.
[36] J.W. Tidey, D.R. Davis, M.E. Miller, I. Pericot-Valverde, R.L. Denlinger-Apte,
D.E. Gaalema, Modeling nicotine regulation: A review of studies in smokers with
mental health conditions, Prev. Med. 117 (2018) 30-37.
[37] S.S. Hecht, Biochemistry, biology, and carcinogenicity of tobacco-specific N-
nitrosamines, Chemical Research in Toxicology 11(6) (1998) 559-603.
[38] S.S. Hecht, I. Stepanov, S.G. Carmella, Exposure and Metabolic Activation
Biomarkers of Carcinogenic Tobacco-Specific Nitrosamines, Accounts of Chemical
Research 49(1) (2016) 106-114.
[39] E. Konstantinou, F. Fotopoulou, A. Drosos, N. Dimakopoulou, Z. Zagoriti, A.
Niarchos, D. Makrynioti, D. Kouretas, K. Farsalinos, G. Lagoumintzis, K. Poulas,
Tobacco-specific nitrosamines: A literature review, Food Chem. Toxicol. 118 (2018)
198-203.
[40] FDA, Tobacco Product Standard for N-Nitrosonornicotine Level in Finished
Smokeless Tobacco Products, in: H.A.H. SERVICES (Ed.) Food and Drug
Administration, 2017.
[41] R.F. Dawson, An experimental analysis of alkaloid production in *Nicotiana* - The
origin of nornicotine, Am J Bot 32(7) (1945) 416-423.
[42] R.E. Dewey, J.H. Xie, Molecular genetics of alkaloid biosynthesis in Nicotiana
tabacum, Phytochemistry 94 (2013) 10-27.
[43] B. Ladesic, T.C. Tso, Biochemical studies on tobacco alkaloids. 6. Biosynthesis
of nicotine through nornicotine, Phytochemistry 3(4) (1964) 541-545.
[44] S. Mizusaki, Y. Tanabe, M. Noguchi, E. Tamaki, Phytochemical studies on
tobacco alkaloids. 16. Changes in activities of ornitheine decarboxylase, putrescine N-
methyltransferase and N-methyl-putrescine oxidase in tobacco roots in relation to
nicotine biosynthesis, Plant Cell Physiol 14(1) (1973) 103-110.
[45] W. Stepka, L.J. Dewey, Conversion of nicotine to nornicotine in harvested
tobacco - fate of methyl group, Plant Physiol 36(5) (1961) 592-&.
[46] T.C. Tso, R.N. Jeffrey, Studies on tobacco alkaloids. 1. Changes in nicotine and
nornicotine content in *Nicotiana*, Plant Physiol 31(6) (1956) 433-439.
[47] T.C. Tso, R.N. Jeffrey, Studies on tobacco alkaloids. 2. The formation of nicotine
and nornicotine in tobacco supplied with N15, Plant Physiol 32(2) (1957) 86-92.
[48] T. Shoji, T. Hashimoto, Smoking out the masters: transcriptional regulators for
nicotine biosynthesis in tobacco, Plant Biotechnol 30(3) (2013) 217-224.

[49] R.S. Lewis, H.O. Lopez, S.W. Bowen, K.R. Andres, W.T. Steede, R.E. Dewey,
Transgenic and mutation-based suppression of a berberine bridge enzyme-like (BBL)
gene family reduces alkaloid content in field-grown tobacco, PLoS One 10(2) (2015).
[50] M. Chakrabarti, S.W. Bowen, N.P. Coleman, K.M. Meekins, R.E. Dewey, B.
Siminszky, CYP82E4-mediated nicotine to nornicotine conversion in tobacco is
regulated by a senescence-specific signaling pathway, Plant Mol Biol 66(4) (2008)
415-427.
[51] L.B. Gavilano, B. Siminszky, Isolation and characterization of the cytochrome
P450 gene CYP82E5v2 that mediates nicotine to nornicotine conversion in the green
leaves of tobacco, Plant Cell Physiol 48(11) (2007) 1567-1574.
[52] R.S. Lewis, S.W. Bowen, M.R. Keogh, R.E. Dewey, Three nicotine demethylase
genes mediate nornicotine biosynthesis in *Nicotiana tabacum* L.: Functional
characterization of the CYP82E10 gene, Phytochemistry 71(17-18) (2010) 1988-
1998.
[53] K. De Boer, S. Tilleman, L. Pauwels, R. Vanden Bossche, V. De Sutter, R.
Vanderhaeghen, P. Hilson, J.D. Hamill, A. Goossens, APETALA2/ETHYLENE
RESPONSE FACTOR and basic helix-loop-helix tobacco transcription factors
cooperatively mediate jasmonate-elicited nicotine biosynthesis, Plant J 66(6) (2011)
1053-1065.
[54] T. Shoji, T. Hashimoto, Tobacco MYC2 regulates jasmonate-inducible nicotine
biosynthesis genes directly and by way of the NIC2-locus ERF genes, Plant Cell
Physiol 52(6) (2011) 1117-1130.
[55] H.B. Zhang, M.T. Bokowiec, P.J. Rushton, S.C. Han, M.P. Timko, Tobacco
transcription factors NtMYC2a and NtMYC2b form nuclear complexes with the
NtJAZ1 repressor and regulate multiple jasmonate-inducible steps in nicotine
biosynthesis, Mol Plant 5(1) (2012) 73-84.
[56] I.T. Baldwin, Z.P. Zhang, N. Diab, T.E. Ohnmeiss, E.S. McCloud, G.Y. Lynds,
E.A. Schmelz, Quantification, correlations and manipulations of wound-induced
changes in jasmonic acid and nicotine in *Nicotiana sylvestris*, Planta 201 (1997) 397-
404.
[57] S. Imanishi, K. Hashizume, M. Nakakita, H. Kojima, Y. Matsubayashi, T.
Hashimoto, Y. Sakagami, Y. Yamada, K. Nakamura, Differential induction by methyl
jasmonate of genes encoding ornithine decarboxylase and other enzymes involved in
nicotine biosynthesis in tobacco cell cultures, Plant Mol Biol 38(6) (1998) 1101-1111.
[58] T. Shoji, Y. Yamada, T. Hashimoto, Jasmonate induction of putrescine N-
methyltransferase genes in the root of *Nicotiana sylvestris*, Plant Cell Physiol 41(7)
(2000) 831-839.
[59] B.F. Xu, M.P. Timko, Methyl jasmonate induced expression of the tobacco
putrescine N-methyltransferase genes requires both G-box and GCC-motif elements,
Plant Mol Biol 55(5) (2004) 743-761.
[60] T. Shoji, T. Ogawa, T. Hashimoto, Jasmonate-induced nicotine formation in
tobacco is mediated by tobacco COI1 and JAZ genes, Plant Cell Physiol 49(7) (2008)
1003-1012.
[61] H.Y. Zhang, W.J. Li, D.X. Niu, Z.J. Wang, X.X. Yan, X.L. Yang, Y.F. Yang, H.
Cui, Tobacco transcription repressors NtJAZ: Potential involvement in abiotic stress
response and glandular trichome induction, Plant Physiol. Biochem. 141 (2019) 388-
397.

[62] K.D. DeBoer, H.L. Dalton, F.J. Edward, S.M. Ryan, J.D. Hamill, RNAi-
mediated down-regulation of ornithine decarboxylase (ODC) impedes wound-stress
stimulation of anabasine synthesis in *Nicotiana glauca*, Phytochemistry 86 (2013) 21-
28.
[63] J. Xie, S. W, W. Maksymowicz, J. W, C. K, C. Wx, C. C, K. J, C. Ma,
Biotechnology: A tool for reduced risk tobacco products - The nicotine experience
from test tube to cigarette pack, Recent Advances in Tobacco Science, 2004, pp. 17-
37.
[64] K.D. DeBoer, H.L. Dalton, F.J. Edward, J.D. Hamill, RNAi-mediated down-
regulation of ornithine decarboxylase (ODC) leads to reduced nicotine and increased
anatabine levels in transgenic *Nicotiana tabacum* L, Phytochemistry 72(4-5) (2011)
344-355.
[65] Y. Chintapakorn, J.D. Hamill, Antisense-mediated down-regulation of putrescine
N-methyltransferase activity in transgenic *Nicotiana tabacum* L.can lead to elevated
levels of anatabine at the expense of nicotine, Plant Mol.Biol. 53 (2003) 87-105.
[66] J.L. Riechmann, J. Heard, G. Martin, L. Reuber, C.Z. Jiang, J. Keddie, L. Adam,
O. Pineda, O.J. Ratcliffe, R.R. Samaha, R. Creelman, M. Pilgrim, P. Broun, J.Z.
Zhang, D. Ghandehari, B.K. Sherman, G.L. Yu, Arabidopsis transcription factors:
genome-wide comparative analysis among Eukaryotes, Science 290 (2000) 2105 -
2110.
[67] J. Jin, F. Tian, D.-C. Yang, Y.-Q. Meng, L. Kong, J. Luo, G. Gao, PlantTFDB
4.0: toward a central hub for transcription factors and regulatory interactions in plants,
Nucleic Acids Res 45(D1) (2016) D1040-D1045.
[68] F. Tian, D.-C. Yang, Y.-Q. Meng, J. Jin, G. Gao, PlantRegMap: charting
functional regulatory maps in plants, Nucleic Acids Res 48(D1) (2019) D1104-
D1113.
[69] M.Z. Li, Y.Z. Li, L.L. Guo, N.D. Gong, Y.Z. Pang, W.B. Jiang, Y.J. Liu, X.L.
Jiang, L. Zhao, Y.S. Wang, D.Y. Xie, L.P. Gao, T. Xia, Functional characterization of
tea (*Camellia sinensis*) MYB4a transcription factor using an integrative approach,
Front. Plant Sci. 8 (2017) 17.
[70] P. Wang, Y. Liu, L. Zhang, W. Wang, H. Hou, Y. Zhao, X. Jiang, J. Yu, H. Tan,
Y. Wang, D.-Y. Xie, L. Gao, T. Xia, Functional demonstration of plant flavonoid
carbocations proposed to be involved in the biosynthesis of proanthocyanidins, The
Plant Journal 101(1) (2020) 18-36.
[71] D.-Y. Xie, S.B. Sharma, N.L. Paiva, D. Ferreira, R.A. Dixon, Role of
anthocyanidin reductase, encoded by *BANYULS* in plant flavonoid biosynthesis,
Science 299(5605) (2003) 396-399.
[72] Y.H. Bai, S. Pattanaik, B. Patra, J.R. Werkman, C.H. Xie, L. Yuan, Flavonoid-
related basic helix-loop-helix regulators, NtAn1a and NtAn1b, of tobacco have
originated from two ancestors and are functionally active, Planta 234(2) (2011) 363-
375.
[73] A. Baudry, M. Caboche, L. Lepiniec, TT8 controls its own expression in a
feedback regulation involving TTG1 and homologous MYB and bHLH factors,
allowing a strong and cell-specific accumulation of flavonoids in *Arabidopsis*
thaliana, Plant J 46(5) (2006) 768-779.
[74] A. Gonzalez, M. Zhao, J.M. Leavitt, A.M. Lloyd, Regulation of the anthocyanin
biosynthetic pathway by the TTG1/bHLH/Myb transcriptional complex in
Arabidopsis seedlings, Plant J 53(5) (2008) 814-827.

[75] A. Chini, S. Fonseca, G. Fernandez, B. Adie, J.M. Chico, O. Lorenzo, G. Garcia-
Casado, I. Lopez-Vidriero, F.M. Lozano, M.R. Ponce, J.L. Micol, R. Solano, The JAZ
family of repressors is the missing link in jasmonate signalling, Nature 448(7154)
(2007) 666-+.
[76] M.M.B. Zvi, E. Shklarman, T. Masci, H. Kalev, T. Debener, S. Shafir, M.
Ovadis, A. Vainstein, PAP1 transcription factor enhances production of
phenylpropanoid and terpenoid scent compounds in rose flowers, New Phytol. (2012)
10.1111/j.1469-8137.2012.04161.x.
[77] K. Feng, X.L. Hou, G.M. Xing, J.X. Liu, A.Q. Duan, Z.S. Xu, M.Y. Li, J.
Zhuang, A.S. Xiong, Advances in AP2/ERF super-family transcription factors in
plant, Critical reviews in biotechnology 40(6) (2020) 750-776.
[78] P.S. Millard, B.B. Kragelund, M. Burow, R2R3 MYB transcription factors -
functions outside the DNA-binding domain, Trends Plant Sci 24(10) (2019) 934-946.
[79] M. Pireyre, M. Burow, Regulation of MYB and bHLH transcription factors: a
glance at the protein level, Mol Plant 8(3) (2015) 378-88.
[80] M.K. Skinner, A. Rawls, J. Wilson-Rawls, E.H. Roalson, Basic helix-loop-helix
transcription factor gene family phylogenetics and nomenclature, Differentiation;
research in biological diversity 80(1) (2010) 1-8.
[81] R.S. Lewis, A.M. Jack, J.W. Morris, V.J.M. Robert, L.B. Gavilano, B.
Siminszky, L.P. Bush, A.J. Hayes, R.E. Dewey, RNA interference (RNAi)-induced
suppression of nicotine demethylase activity reduces levels of a key carcinogen in
cured tobacco leaves, Plant Biotechnol J 6(4) (2008) 346-354.
[82] K.D. DeBoer, J.C. Lye, C.D. Aitken, A.K.K. Su, J.D. Hamill, The A622 gene in
Nicotiana glauca (tree tobacco): evidence for a functional role in pyridine alkaloid
synthesis, Plant Mol Biol 69(3) (2009) 299-312.
[83] T.A.G. Initiative, Analysis of the genome sequence of the flowering plant
Arabidopsis thaliana, Nature 408 (2000) 796-815.
[84] J. Yu, e. al., A draft sequence of the rice genome (*Oryza sativa* L. ssp *indica*),
Science 296 (2002) 79-92.
[85] S.A. Goff, e. al, A draft sequence of the rice genome (*Oryza sativa* L. ssp
japonica), Science 296 (2002) 92-100.
[86] M.Y. Kim, S. Lee, K. Van, T.H. Kim, S.C. Jeong, I.Y. Choi, D.S. Kim, Y.S. Lee,
D. Park, J. Ma, W.Y. Kim, B.C. Kim, S. Park, K.A. Lee, D.H. Kim, K.H. Kim, J.H.
Shin, Y.E. Jang, K. Do Kim, W.X. Liu, T. Chaisan, Y.J. Kang, Y.H. Lee, J.K. Moon,
J. Schmutz, S.A. Jackson, J. Bhak, S.H. Lee, Whole-genome sequencing and intensive
analysis of the undomesticated soybean (Glycine soja Sieb. and Zucc.) genome,
PNAS 107(51) (2010) 22032-22037.
[87] A.Y. Guo, X. Chen, G. Gao, H. Zhang, Q.H. Zhu, X.C. Liu, Y.F. Zhong, X.C.
Gu, K. He, J.C. Luo, PlantTFDB: a comprehensive plant transcription factor database,
Nucleic Acids Res 36 (2008) D966-D969.
[88] J. Jin, H. Zhang, L. Kong, G. Gao, J. Luo, PlantTFDB 3.0: a portal for the
functional and evolutionary study of plant transcription factors, Nucleic Acids Res
42(D1) (2013) D1182-D1187.
[89] A.Y. Guo, K. He, D. Liu, S.N. Bai, X.C. Gu, L.P. Wei, J.C. Luo, DATF: a
database of Arabidopsis transcription factors, Bioinformatics 21(10) (2005) 2568-
2569.

[90] K. Iida, M. Seki, T. Sakurai, M. Satou, K. Akiyama, T. Toyoda, A. Konagaya, K.
Shinozaki, RARTF: Database and tools for complete sets of Arabidopsis transcription
factors, DNA Res. 12(4) (2005) 247-256.
[91] G. Gao, Y.F. Zhong, A.Y. Guo, Q.H. Zhu, W. Tang, W.M. Zheng, X.C. Gu, L.P.
Wei, J.C. Luo, DRTF: a database of rice transcription factors, Bioinformatics 22(10)
(2006) 1286-1287.
[92] D.M. Riano-Pachon, S. Ruzicic, I. Dreyer, B. Mueller-Roeber, PlnTFDB: an
integrative plant transcription factor database, BMC Bioinformatics 8 (2007) 10.
[93] J. Fredslund, DATFAP: A database of primers and homology alignments for
transcription factors from 13 plant species, Bmc Genomics 9 (2008) 10.

1213

1214

1215 **Figure Legends**

1216

Figure 1 Biosynthetic pathways of nicotine, other tobacco alkaloids, and

anthocyanin and their regulations. A, the biosynthetic pathway of Arabidopsis

anthocyanin from the step catalyzed by CHS. *PAP1*, *TT8*, and *TTG1* encode MYB75,

bHLH, and WD40 to form a MYB-bHLH-WD40 complex that activates late pathway

genes. Gene abbreviations are *CHS* and *CHI*: chalcone synthase and isomerase, *F3H*:

flavonone-3-hydroxylase, *F3'H* and *F3'5'H*: flavonoid-3'-hydroxylase and

3'5'hydroxylase, *DFR*: dihydroflavonol reductase, *ANS*: anthocyanidin synthase, *3-*

GT: glycosyltransferase. *ANR* encodes anthocyanidin reductase, a key enzyme toward

proanthocyanidin biosynthesis. **B,** the biosynthetic pathway of nicotine, nornicotine,

and tobacco specific nitrosamines (TSNAs) and the regulation mechanism. MYC2

and EAR are two positive transcription factors that activate the expression of most

pathway genes. In the absence of isoleucine-jasmonate (Ile-JA), NtJAZ1 and NtJAZ3

repressors bind to MYC2 to inhibit the activation of the pathway genes (red lines). In

the presence of bioactive Ile-JA (green arrows), SCF^{COI1} perceives this plant hormone

to form a new JA-SCF^{COI1} complex to bind to NtJAZs and lead them to proteasomal

ubiquitination. The degradation of NtJAZs releases MYC2 to activate pathway gene

expression toward the biosynthesis of nicotine. Gene abbreviations are *AO*: aspartate

oxidase, *QS*: quinolinate synthase, *QPT2*: quinolinate phosphoribosyltransferase 2,

ADC: arginine decarboxylase, *AIH*: agmatine iminohydrolase, *ODC*: ornithine

decarboxylase, *MPT1*: N-methylputricine transferase 1, *MPO1*: methylputricine

oxidase 1, *A622*: an isoflavone reductase like enzyme, *BBL*: berberine bridge enzyme-

like protein, and *NND*: nicotine N-demethylase.

1239

Figure 2 Binding and activation of *NtJAZ1* and *NtJAZ3* promoters by PAP1 and

TT8 alone, PAP1 and TT8 together, and PAP1-TT8-WD40 complex. A, three

MYB response element (MRE) types and G-box were identified in the *NtJAZ1* and

NtJAZ3 promoters; **B,** electrophoretic mobility shift assay (EMSA) showed that PAP1

and TT8 bound to three MRE types and G-box of the NtJAZ1 and NtJAZ3 promoters,

respectively. Competitive and non-competitive probes were used for binding of both

TFs. 20× and 10×: concentrations of tested competitive and non-competitive probes

20 and 10 times as those MRE and G-box probes. **C,** fold change values from ChIP-

qPCR showed that both PAP1 and TT8 bound to the *NtJAZ1* and *NtJAZ3* promoters *in*

vivo. Approximately 150 bp *NtJAZ1* and *NtJAZ3* promoter fragments containing both

MRE and G-box were enriched with anti-HA antibodies in qRT-PCR analysis. The

regions of *NtJAZ1* and *NtJAZ3* promoters that do not contain a MRE and G-box were

used as negative controls. Green and red bars represent wild type and red tobacco

plants, respectively. **D,** schematic diagrams shows four effector constructs (PK2GW7-

GFP, PAP1, TT8 and TTG1) and two reporter constructs (pGreenII-0800-Jaz1pro and

Jaz3 pro) used for dual-luciferase assay. **E,** luciferase (LUC)/renilla (REN)

luminescence ratios from dual luciferase assays showed that PAP1 and TT8 alone

bound to and activated the promoters of both *NtJAZ1* and *NtJAZ3*, two TFs together

increased promoter activity, and two TFs and WD40 together increased the highest

activity. The promoters were fused to a firefly luciferase (reporter). The promoters of

NtJAZ1 and *NtJAZ3* genes were used in dual luciferase assay. Values represent the

mean ± S.D. (n=5). Asterisks on top of bars indicate that the values are significantly

higher than those of controls (*P<0.05, **P<0.01, ***P<0.001). Standard bars mean

standard errors.

1264

Figure 3 Upregulation of two *NtJAZs* and downregulation of nicotine pathway

genes and *ERF189* in roots of P+T transgenic seedlings

A-J, in roots of P+T-NL transgenic lines, the expression levels of *NtJAZ1* (**A**) and

NtJAZ3 (**B**) were significantly upregulated; the expression level of *ODC2* (**C**) was not

changed; while the expression levels of *PMT1* (**D**), *PMT2* (**E**) *MPO* (**F**), *QPT* (**G**),

A622 (**H**), *BBLs* (**I**) (primer pairs designed for all three BBLs), and *ERF189* (**J**) were

significantly downregulated. **K-T**, in roots of P+T-KY transgenic lines, the expression

levels of *NtJAZ1* (**K**) and *NtJAZ3* (**L**) were significantly upregulated, while, the

expression levels *ODC2* (**M**), *PMT1* (**N**), *PMT2* (**O**) *MPO* (**P**), *QPT* (**Q**), *A622* (**R**),

BBLs (**S**), and *ERF189* (**T**) were significantly downregulated. Data of three transgenic

lines are shown for both P+T-NL and P+T-KY genotypes. Wild type samples were

pooled from five plants. Data of three lines are shown for vector control transgenic

NL and KY171 plants. WT: wild type, P+T NL1, P+T NL2, P+T NL3: three lines of

stacked *PAP1* and *TT8* transgenic NL plants, VC-1, 2, and 3-NL: three vector control

transgenic NL lines, P+T KY1, P+T KY2, P+T KY3: three lines of stacked *PAP1* and

TT8 transgenic KY171 plants. VC-1, 2, and 3-KY: three vector control transgenic

KY171 lines.

Green and red bars represent wild type and vector control transgenic, and red P+T-NL

and P+T-KY plants, respectively. Values represent the mean ± S.D. (n=3). Asterisks

on top of the bars indicate that values are significantly lower or higher in red

transgenic lines than in wild type plants (*P<0.05, **P<0.01, ***P<0.001). Standard

bars mean standard errors.

1287

Figure 4 Significant reduction of nicotine and nornicotine contents in leaves and

roots of P+T-NL and P+T-KY seedlings. The contents of nicotine and nornicotine

were significantly reduced in roots and leaves of both transgenic P+T-NL and P+TKY

lines. **A-C:** reduction of nicotine (**A**), nornicotine (**B**), and total nicotine and

nornicotine (**C**) in leaves of three P+T-NL lines (1, 2, and 3); **D-F:** reduction of

nicotine (**D**), nornicotine (**E**), and total nicotine and nornicotine (**F**) in roots of three

P+T-NL lines (1, 2, and 3); **G-I:** reduction of nicotine (**G**), nornicotine (**H**), and total

nicotine and nornicotine (**I**) in leaves of three P+T-KY lines (1, 2 and 3); **J-L:**

reduction of nicotine (**J**), nornicotine (**K**), and total nicotine and nornicotine (**L**) in

roots of three P+T-NL lines (1, 2, and 3). Bars labeled with "a" and "b" means p-

value less than 0.05 and bars labelled with the same lowcase "a" or "b" means no

significant difference.

1300

Figure 5 Phenotypes of P+T-NL1 and P+T-KY1 plants versus their

corresponding wild-type tobacco plants in the field and of air-cured leaves. Field

farming practice of four genotypes was performed in Oxford, North Carolina. **A,**

phenotypes of 30-day and 60-day old WT-NL vs. red P+T-NL1 plants. **B**, phenotypes

of 30-day and 60-day old WT-KY171 vs. red P+T-KY1 plants. **C-D**, phenotypes of

topped plants (**C**) and leaves in air curing (**D**). Plant name abbreviations are WT-NL:

wild-type Narrow Leaf Madole (NL) variety, P+T-NL1: Stacked *PAP1* and *TT8*

transgenic NL line 1, WT-KY171: wild-type KY171 variety, and P+T-KY1: stacked

PAP1 and *TT8* transgenic KY line 1.

1310

Figure 6 Reduction of nicotine and four other alkaloids in cured leaves of red

P+T-NL1 and P+T-KY1 tobacco plants. A-J, reduction of nicotine (**A-B**),

nornicotine (**C-D**), anabasine (**E-F**), anatabine (**G-H**), and myosmine (**I-J**) in four leaf

groups of P+T-NL (**A, C, E, G, and I**) and P+T-KY plants (**B, D, F, H, and J**). **K-L**,

cartoons showing leaf groups harvested from both wild type (**K**) and transgenic red

plants (**L**).

Green and red bars represent wild type and red tobacco plants, respectively. B1, B2,

B3, and B4 labels represent the 1^{st}, 2^{nd}, 3^{rd}, and 4^{th} group of leaves from the base to

the top of tobacco plant, respectively. Each metabolite in each leaf group was

quantified from leaves harvested from 120 tobacco plants in three plots. One-way

ANOVA test included in the SAS software was used to evaluate statistical

significance. Bars labelled with lowcase "a" and "b" mean significant difference

between paired green and red bars (n=120, P-value less than 0.05). Percentages

labeled in the top of red bars mean reduction compared to green bars. The standard

bars mean standard deviation. Abbreviations, WT-NL: wild-type Narrow Leaf

Madole (NL) variety, P+T-NL1: Stacked *PAP1* and *TT8* transgenic NL line 1,

WTKY: wild-type KY171 variety, and P+T-KY1: stacked *PAP1* and *TT8* transgenic

KY line 1.

1329

1330

Figure 7 Reduction of four individual and total TSNAs in leaves of red P+T-NL1

and P+T-KY1 tobacco plants. A-H, reduction of NNN (**A-B**), NNK (**C-D**), NAT

(**E-F**), and NAB (**G-H**) in each leaf group of P+T-NL1 (**A, C, E,** and **G**) and

P+TKY1 (**B, D, F,** and **H**) plants; **I-J,** reduction of total TSNA contents summed from

four individual TSNAs of each leaf group of P+T-NL1 (**I**) and P+T-KY1 (**J**) plants;

K-L, reduction of total TSNA contents summed from four individual TSNAs of all

leaves of P+T-NL1 (**K**) and P+T-KY1 (**L**) plants.

Green and red bars represent wild type and red tobacco plants, respectively. B1, B2,

B3, and B4 labels represent the 1^{st}, 2^{nd}, 3^{rd}, and 4^{th} group of leaves from the base to

the top of plants, respectively. Each metabolite in each group of leaves was quantified

from 120 tobacco plants. One-way ANOVA test included in the SAS software was

used to evaluate statistical significance. Paired green and red bars labelled with

lowcase "a" and "b" mean significant difference (n=120, P-value less than 0.05).

Percentages labeled on the top of red bars indicate reduction levels compared to green

bars. Abbreviation, WT-NL: Narrow Leaf Madole, WT-KY171: wild type KY171,

P+T-NL: stacked PAP1 and TT8 transgenic NL line 1, and P+T-KY1: stacked PAP1

and TT8 transgenic KY171 line 1.

1348

Figure 8 A model showing the distant pathway-cross regulation (DPCR) created

for the downregulation of the biosynthesis of tobacco alkaloids in red tobacco

plants

Regulation pathways created in red tobacco plants are featured by red colored solid

arrows, "X" shape, and "T" blocks. The stacked *PAP1* and *TT8* as well as the

endogenous *NtTTG1* encode MYB, bHLH, and WD40 proteins, respectively, which

form a complex to regulate two biosynthetic pathways. The regulation pathway

(function I) is the activation of biosynthesis of anthocyanin. The regulation pathway

(function II) is the DPCR that the MYB-bHLH-WD40 complex upregulates the

endogenous *NtJAZs* leading to the reduction of MYC2 and the downregulation of

EAR and the nicotine biosynthesis. In addition, The JA-JAZ-MYC regulatory pathway

is indicated with green arrows starting with JA (greenish arrows). The feedback

regulation of NtMYC2 indicated by a brownish bending arrow is proposed to activate

the expression of NtJAZs. SCF: Skp/Cullin/F-box and COI1: coronatine insensitive 1.

1363

Figures

1366

Figure 1 Biosynthetic pathways of nicotine, other tobacco alkaloids, and

anthocyanin and their regulations. A, the biosynthetic pathway of Arabidopsis

anthocyanin from the step catalyzed by CHS. *PAP1*, *TT8*, and *TTG1* encode MYB75,

bHLH, and WD40 to form a MYB-bHLH-WD40 complex that activates late pathway

genes. Gene abbreviations are *CHS* and *CHI*: chalcone synthase and isomerase, *F3H*:

flavonone-3-hydroxylase, *F3'H* and *F3'5'H*: flavonoid-3'-hydroxylase and

3'5'hydroxylase, *DFR*: dihydroflavonol reductase, *ANS*: anthocyanidin synthase, *3-*

GT: glycosyltransferase. *ANR* encodes anthocyanidin reductase, a key enzyme toward

proanthocyanidin biosynthesis. Red lines and arrows highlight that the complex

upregulates those steps.

B, the biosynthetic pathway of nicotine, nornicotine, and tobacco specific

nitrosamines (TSNAs) and the regulation mechanism. MYC2 and EAR are two

positive transcription factors that activate the expression of most pathway genes. In

the absence of isoleucine-jasmonate (Ile-JA), NtJAZ1 and NtJAZ3 repressors bind to

MYC2 to inhibit the activation of the pathway genes (red lines). In the presence of

bioactive Ile-JA (green arrows), SCF^{COI1} perceives this plant hormone to form a new

JA-SCF^{COI1} complex to bind to NtJAZs and lead NtJAZs to proteasomal

ubiquitination. The degradation of NtJAZs releases MYC2 to activate pathway gene

expression toward the biosynthesis of nicotine. Gene abbreviations are *AO*: aspartate

oxidase, *QS*: quinolinate synthase, *QPT2*: quinolinate phosphoribosyltransferase 2,

ADC: arginine decarboxylase, *AIH*: agmatine iminohydrolase, *ODC*: ornithine

decarboxylase, *MPT1*: N-methylputricine transferase 1, *MPO1*: methylputricine

oxidase 1, *A622*: an isoflavone reductase like enzyme, *BBL*: berberine bridge enzyme-

like protein, and *NND*: nicotine N-demethylase.

Figure 2 Binding and activation of *NtJAZ1* and *NtJAZ3* promoters by PAP1 and TT8 alone, PAP1 and TT8 together, and PAP1-TT8-WD40 complex. A, three MYB recognizing element (MRE) types and G-box were identified in the *NtJAZ1* and *NtJAZ3* promoters; **B**, electrophoretic mobility shift assay (EMSA) showed that PAP1 and TT8 bound to three MRE types and G-box of the *NtJAZ1* and *NtJAZ3* promoters, respectively. Competitive and non-competitive probes were used for binding of both TFs. $20\times$ and $10\times$: concentrations of tested competitive and non-competitive probes 20 and 10 times as those MRE and G-box probes. **C**, fold change values from Chip-qPCR showed that both PAP1 and TT8 bound to the *NtJAZ1* and *NtJAZ3* promoters *in vivo*. Approximately 150 bp *NtJAZ1* and *NtJAZ3* promoter fragments containing both MRE and G-box were enriched with anti-HA antibodies in qRT-PCR analysis. The regions of *NtJAZ1* and *NtJAZ3* promoters that do not contain a MRE and G-box were used as negative controls. Green and red bars represent wild type and red tobacco plants, respectively. **D**, schematic diagrams shows four effector constructs (PK2GW7-GFP, PAP1, TT8 and TTG1) and two reporter constructs (pGreenII-0800-Jaz1pro and Jaz3pro) used for dual-luciferase assay. **E**, luciferase (LUC)/renilla (REN) luminescence ratios from dual luciferase assays showed that PAP1 and TT8 alone bound to and activated the promoters of both *NtJAZ1* and *NtJAZ3*, two TF together increased promoter activity, and two TFs and WD40 together increased the highest activity. The promoters were fused to a firefly luciferase (reporter). The promoters of *NtJAZ1* and *NtJAZ3* genes were used in dual luciferase assay. Values represent the mean $\pm$ S.D. (n=5). Asterisks on top of bars indicate that the values are significantly higher than those of controls (*$P<0.05$, **$P<0.01$, ***$P<0.001$). Standard bars mean standard errors.

Figure 3 Upregulation of two *NtJAZs* and downregulation of six or seven nicotine pathway genes and *ERF189* in roots of P+T transgenic seedlings shown by qRT-PCR

A-J, in roots of P+T-NL transgenic lines, the expression levels of *NtJAZ1* (**A**) and *NtJAZ3* (**B**) were significantly upregulated; the expression level of *ODC2* (**C**) was not changed; and

the expression levels of *PMT1* (**D**), *PMT2* (**E**) *MPO* (**F**), *QPT* (**G**), *A622* (**H**), *BBLs* (**I**) (primer pairs designed for all three BBLs), and *ERF189* (**J**) were significantly downregulated. **K-T**, in roots of P+T-KY transgenic lines, the expression levels of *NtJAZ1* (**K**) and *NtJAZ3* (l) were significantly upregulated, and the expression levels *ODC2* (**M**), *PMT1* (**N**), *PMT2* (**O**) *MPO* (**P**), *QPT* (**Q**), *A622* (**R**), *BBLs* (**S**), and *ERF189* (**T**) were significantly downregulated. Data of three transgenic lines are shown for both P+T-NL and P+T-KY genotypes. Wild type samples were pooled from five plants. Data of three lines are shown for vector control transgenic NL and KY171 plants. WT: wild type, P+T NL1, P+T NL2, P+T NL3: three lines of stacked *PAP1* and *TT8* transgenic NL plants, VC-1, 2, and 3-NL: three vector control transgenic NL lines, P+T KY1, P+T KY2, P+T KY3: three lines of stacked *PAP1* and *TT8* transgenic KY171 plants. VC-1, 2, and 3-KY: three vector control transgenic KY171 lines.

Green and red bars represent wild type and vector control transgenic, and red P+T-NL and P+T-KY plants, respectively. Values represent the mean ± S.D. (n=3). Asterisks on top of the bars indicate that values are significantly lower or higher in red transgenic lines than in wild type plants (*P<0.05, **P<0.01, ***P<0.001). Standard bars mean standard errors.

Figure 4 Significant reduction of nicotine and nornicotine contents in leaves and roots of P+T-NL and P+T-KY seedlings. The contents of nicotine and nornicotine were significantly reduced in roots and leaves of both transgenic P+T-NL and P+T-KY lines. **A-C:** reduction of nicotine (**A**), nornicotine (**B**), and total nicotine and nornicotine (**C**) in leaves of three P+T-NL lines (1, 2, and 3); **D-F:** reduction of nicotine (**D**), nornicotine (**E**), and total nicotine and nornicotine (**F**) in roots of three P+T-NL lines (1, 2, and 3); **G-I:** reduction of nicotine (**G**),

nornicotine (**H**), and total nicotine and nornicotine (**I**) in leaves of three P+T-KY lines (1, 2 and 3); **J-L:** reduction of nicotine (**J**), nornicotine (**K**), and total nicotine and nornicotine (**L**) in roots of three P+T-NL lines (1, 2, and 3). Bars labeled with "a" and "b" means p1314 value less than 0.05 and bars labelled with the same lowcase "a" or "b" means no 1315 significant difference.

1316

1316

1317

Figure 5 Phenotypes of P+T-NL1 and P+T-KY1 plants versus their

corresponding wild-type tobacco plants in the field and of air-cured leaves. Field

farming practice of four genotypes was performed in Oxford, North Carolina. **A**,

phenotypes of 30-day and 60-day old WT-NL vs. red P+T-NL1 plants. **B**, phenotypes

of 30-day and 60-day old WT-KY171 vs. red P+T-KY1 plants. **C-D**, phenotypes of

topped plants (**C**) and leaves in air curing (**D**). Plant name abbreviations are WT-NL:

wild-type Narrow Leaf Madole (NL) variety, P+T-NL1: Stacked *PAP1* and *TT8*

transgenic NL line 1, WT-KY171: wild-type KY171 variety, and P+T-KY1: stacked

PAP1 and *TT8* transgenic KY line 1.

1328

Figure 6 Reduction of nicotine and four other alkaloids in air-cured leaves of red

P+T-NL1 and P+T-KY1 tobacco plants. A-J, reduction of nicotine (**A-B**),

nornicotine (**C-D**), anabasine (**E-F**), anatabine (**G-H**), and myosmine (**I-J**) in four leaf

groups of P+T-NL (**A, C, E, G, and I**) and P+T-KY plants (**B, D, F, H, and J**). **K-L**,

cartoons showing leaf groups harvested from both wild type (**K**) and transgenic red

plants (**L**).

Green and red bars represent wild type and red tobacco plants, respectively. B1, B2,

B3, and B4 labels represent the 1^{st}, 2^{nd}, 3^{rd}, and 4^{th} group of leaves from the base to

the top of tobacco plant, respectively. Each metabolite in each leaf group was

quantified from leaves harvested from 120 tobacco plants in three plots. One-way

ANOVA test included in the SAS software was used to evaluate statistical

significance. Bars labelled with lowcase "a" and "b" mean significant difference

between paired green and red bars (n=120, P-value less than 0.05). Percentages

labeled in the top of red bars mean reduction compared to green bars. The standard

bars mean standard deviation. Abbreviations, WT-NL: wild-type Narrow Leaf

Madole (NL) variety, P+T-NL1: Stacked *PAP1* and *TT8* transgenic NL line 1,

WTKY: wild-type KY171 variety, and P+T-KY1: stacked *PAP1* and *TT8* transgenic

KY line 1.

1349

Figure 7 Reduction of four individual and total TSNAs in leaves of red P+T-NL1 and

P+T-KY1 tobacco plants. A-H, reduction of NNN (**A-B**), NNK (**C-D**), NAT (**E-F**), and

NAB (**G-H**) in each leaf group of P+T-NL1 (**A, C, E,** and **G**) and P+TKY1 (**B, D, F,** and **H**)

plants; **I-J,** reduction of total TSNA contents summed from four individual TSNAs of each

leaf group of P+T-NL1 (**I**) and P+T-KY1 (**J**) plants; **K-L,** reduction of average total TSNA

contents summed from four individual TSNAs of all leaves of P+T-NL1 (**K**) and P+T-KY1

(**L**) plants.

Green and red bars represent wild type and red tobacco plants, respectively. B1, B2, B3, and

B4 labels represent the 1^{st}, 2^{nd}, 3^{rd}, and 4^{th} group of leaves from the base to the top of plants,

respectively. Each metabolite in each group of leaves was quantified from 120 tobacco

plants. One-way ANOVA test included in the SAS software was used to evaluate statistical

significance. Paired green and red bars labelled with lowcase "a" and "b" mean significant

difference (n=120, P-value less than 0.05). Percentages labeled on the top of red bars indicate

reduction levels compared to green bars. Abbreviation, WT-NL: Narrow Leaf Madole, WT-

KY171: wild type KY171, P+T-NL: stacked PAP1 and TT8 transgenic NL line 1, and P+T-

KY1: stacked PAP1 and TT8 transgenic KY171 line 1.

1365

1368

Figure 8 A model showing the distant pathway-cross regulation (DPCR) created for the

downregulation of the biosynthesis of tobacco alkaloids in red tobacco plants

Regulation pathways created in red tobacco plants are featured by red colored solid arrows,

"X" shape, and "T" blocks. The stacked *PAP1* and *TT8* as well as the endogenous *NtTTG1*

encode MYB, bHLH, and WD40 proteins, respectively, which form a complex to regulate

two biosynthetic pathways. The regulation pathway (function I) is the activation of

biosynthesis of anthocyanin. The regulation pathway (function II) is the DPCR that the

MYB-bHLH-WD40 complex upregulates the endogenous *NtJAZs* leading to the reduction of

MYC2 and the downregulation of *EAR* and the nicotine biosynthesis. In addition, The JA-

JAZ-MYC regulatory pathway is indicated with green arrows starting with JA (greenish

arrows). The feedback regulation of NtMYC2 indicated by a brownish bending arrow is

proposed to activate the expression of NtJAZs. SCF: Skp/Cullin/F-box and COI1: coronatine

insensitive 1.

1382

1383