Unbiased population heterozygosity estimates from genome-wide sequence data

Thomas L Schmidt[1]*, Moshe Jasper[1], Andrew R Weeks[1,2], Ary A Hoffmann[1]

[1]*School of BioSciences, Bio21 Institute, University of Melbourne, Parkville, Victoria, Australia*
[2]*cesar Pty Ltd, Parkville, Victoria, Australia*
*Correspondence: Tom Schmidt, Bio21 Institute, University of Melbourne, Parkville, Victoria, Australia,
toms@unimelb.edu.au

Abstract

1. Heterozygosity is a metric of genetic variability frequently used to inform the management of threatened taxa. Estimating observed and expected heterozygosities from genome-wide sequence data has become increasingly common, and these estimates are often derived directly from genotypes at single nucleotide polymorphism (SNP) markers. While many SNP markers can provide precise estimates of genetic processes, the results of 'downstream' analysis with these markers may depend heavily on 'upstream' filtering decisions.

2. Here we explore the downstream consequences of sample size, rare allele filtering, missing data thresholds and known population structure on estimates of observed and expected heterozygosity using two reduced-representation sequencing datasets, one from the mosquito *Aedes aegypti* (ddRADseq) and the other from a threatened grasshopper, *Keyacris scurra* (DArTseq).

3. We show that estimates based on polymorphic markers only (i.e. SNP heterozygosity) are always biased by global sample size (N), with smaller N producing larger estimates. By contrast, results are unbiased by sample size when calculations consider monomorphic as well as polymorphic sequence information (i.e. genome-wide or autosomal heterozygosity). SNP heterozygosity is also biased when differentiated populations are analysed together and particularly when these have different sample sizes (n), while autosomal heterozygosity remains unbiased. We also show that when nucleotide sites with missing genotypes are included, observed and expected heterozygosity estimates diverge in proportion to the amount of missing data permitted at each site.

4. We make two recommendations for estimating genome-wide heterozygosity: (i) autosomal heterozygosity should be reported instead of (or in addition to) SNP heterozygosity; (ii) sites with any missing data should be omitted. This should facilitate comparisons within and across studies and between observed and expected measures of heterozygosity.

Key-words: Heterozygosity, Single Nucleotide Polymorphisms (SNPs), RADseq, DArTseq, Filtering, Conservation, Population Structure, Genetic Mixing

Introduction

The power provided by single nucleotide polymorphism (SNP) markers detected using genome-wide sequencing approaches is leading to their increased use in conservation genetic studies (Garner et al., 2016). SNPs are popular for investigating levels of genetic differentiation among remnant populations and for comparing levels and patterns of genetic variation within populations (e.g. Maroso, Franch, Dalla Rovere, Arculeo, & Bargelloni, 2016; Campbell et al., 2019) which provides information on the adaptive potential of populations (Ørsted, Hoffmann, Sverrisdóttir, Nielsen, & Kristensen, 2019) as well as patterns of inbreeding and relatedness (Mulvena et al., 2020). Results from SNP studies are being interpreted for use in management decisions that include genetic rescue, genetic mixing and founder selection in threatened species programs (e.g. Fitzpatrick et al., 2020). The relative ease of generating SNP genotypes is leading to their increased use by non-specialists, particularly through the availability of companies such as Diversity Arrays Technology (https://www.diversityarrays.com), which provide SNP genotypes through customised in-house processes (Gruber, Unmack, Berry, & Georges, 2017; Wright et al., 2019; Mulvena et al., 2020).

Considering the popularity of SNP markers, it is important to be aware of any biases inherent in their application to conservation genetics and elsewhere. While potential biases have been considered for the detection of structure between populations (Gruber et al. 2017; Linck & Battey, 2019; Wright et al., 2019), there has been less focus on the estimation of genetic variability within populations. These estimates are important because they link to the evolutionary potential of populations, which is typically higher in populations with greater genetic variability (Hoffmann, Sgrò, & Kristensen, 2017; Ørsted et al., 2019). Genetic variability of populations is therefore crucial when making genetic management decisions for threatened species (Weeks et al., 2011; Hoffmann et al., 2020).

Genetic variation in populations is measured in several ways, the most common of which are heterozygosity (observed and expected) and the proportion of nucleotide sites that are polymorphic. Heterozygosity is usually estimated from a substantial number of individuals sampled from each population, but with large quantities of sequence data fewer individuals may be needed (Nazareno, Bemmels, Dick, & Lohmann, 2017). Accurate heterozygosity estimates also require that the apparent diversity at a site is not related to errors introduced during sequencing or genotyping, the latter of which requires adequate coverage to ensure that both strands of a diploid individual are sequenced (Nielsen, Paul, Albrechtsen, & Song, 2011). While expected heterozygosity is estimated from allele frequencies, observed heterozygosity is estimated from individual genotypes directly and depends on both the amount of genetic variation in the population and the level of inbreeding, which increases homozygosity (Ritland, 1996). Inbreeding can thus be estimated by comparing observed heterozygosity to expected heterozygosity, with the latter expected to be relatively higher when there is inbreeding ($F_{IS} > 0$).

For heterozygosity, h_i, the observed heterozygosity for an individual at site i can be averaged across n sites as $\frac{\sum_{i=1}^{n} h_{i...n}}{n}$ and averaged across a sample of individuals to provide a population estimate. This can be calculated from variation at polymorphic sites only (i.e. SNP

heterozygosity) or at both polymorphic and monomorphic sites (i.e. genome-wide or autosomal heterozygosity). Both SNP heterozygosity and autosomal heterozygosity appear in the literature; most population-focussed studies tend to report SNP heterozygosity (e.g. Jones et al., 2012; Chen, Cosgrove, Bowman, Fitzpatrick, & Clark, 2016; Bock, Kantar, Caseys, Matthey-Doret, & Rieseberg, 2018; Mathur, Tomeček, Heniff, Luna, & DeWoody, 2019; Surbakti et al., 2020) although others use autosomal heterozygosity (e.g. Hohenlohe et al., 2010) which is the parameter reported in studies comparing individual genomes (e.g. Gopalakrishnan et al., 2017; Westbury, Petersen, Garde, Heide-Jørgensen, & Lorenzen, 2019). As SNP heterozygosities will be orders of magnitude larger than autosomal heterozygosities, the two parameters cannot be directly compared, though for studies of a single population autosomal heterozygosity can be converted to SNP heterozygosity by dividing the estimate by the proportion of polymorphic sites. Fig. S1 provides a visualisation of how observed heterozygosity is calculated using all sequence information (autosomal heterozygosity) and using polymorphic markers only (SNP heterozygosity).

This paper investigates how SNP heterozygosity and autosomal heterozygosity perform under variable conditions of sampling and filtering. These include local and global sample size, rare allele filtering, missing data thresholds and the analysis of multiple differentiated populations, all of which are common sources of variability within or between studies. We explore these questions with a pair of genome-wide datasets of the sort frequently used for assessing variation in wild populations. We focus initially on a ddRADseq dataset from one population of a common species and then consider a DArTseq dataset from a threatened species that covers multiple populations. We make some recommendations for assessing heterozygosity when study aims include comparisons of genetic variability across populations and with other studies.

Materials and Methods

Sequence data from the same population

We start by considering a single, well-mixed population. We use double digest restriction-site associated (ddRAD) sequence data obtained from 100 female *Aedes aegypti* mosquitoes sampled from a 0.125 km^2 area of Kuala Lumpur, Malaysia (Jasper, Schmidt, Ahmad, Sinkins, & Hoffmann, 2019). Note that as this ddRADseq dataset contains only females, and as *Ae. aegypti* mosquitoes do not have definable sex chromosomes but rather a small sex-determining region (Fontaine et al., 2017), we did not need to filter out genotypes at sex chromosomes.

We took subsamples from this population as follows.

Ten subsamples: We tested the effect of five population sample sizes (n = 10, 5, 4, 3, 2) on heterozygosity estimates by subsampling the 100 individuals, without replacement. We repeated the subsampling 10 times for each sample size n.

Nested subsamples: We tested the effect of six larger sample sizes (n = 50, 40, 30, 20, 10, 5) on heterozygosity estimates by subsampling the 100 individuals twice, without replacement. This can help indicate whether filtering choices produce similar patterns at large n as at small n. These subsamples were also used to test whether different filtering choices could produce variable heterozygosity estimates from the same sample of individuals. To reduce variation among subsamples of different size, we used a nested subsampling approach. The 100 individuals were randomly assigned to two groups, A and B, each of n = 50. Group A and Group B were then subsampled once at each n, but where each subsample could only include individuals that were included at the next highest n. For instance, the subsample at n = 30 could only contain individuals that were present in the n = 40 subsample to allow for a direct comparison between sample sizes.

Sequence data from multiple populations

We considered the issue of multiple populations being included in a comparison by reanalysing a set of four populations of *Keyacris scurra* (Key's Matchstick Grasshopper) taken from a larger set of sequencing data derived from a Diversity Arrays Technology (DArT) approach. *Keyacris scurra* has recently been listed as endangered and is currently restricted in range to refugia in south-eastern Australia. These four populations have experienced very low gene flow and are highly differentiated (pairwise F_{ST} = 0.14–0.28). The four populations were processed and sequenced together as part of the same project (Hoffmann et al., 2020). Note that no reference assembly is available for this species so the term "autosomal heterozygosity" here will also include sequence data from any differentiated sex chromosomes (or regions of chromosomes).

Sequence processing

For the ddRADseq dataset, aligned sequences were built into a Stacks.v2 (Catchen, Hohenlohe, Bassham, Amores, & Cresko, 2013) catalog with the program ref_map. For the DArTSeq dataset, sequence data were built into a *de novo* Stacks catalog using the program denovo_map, allowing for up to four mismatches within and between individuals. We analysed both datasets with the Stacks program "Populations", which was used to estimate observed and expected heterozygosity for a range of filtering settings described below.

Results

Estimates based on polymorphic sites (SNP heterozygosity)

Our first aim was to see how a dataset filtered with settings typically used for assessing genetic structure (i.e. variation between populations) might perform when used to estimate heterozygosity (i.e. variation within populations). Analysis of genetic structure will usually

consider only polymorphic sites (SNPs). When filtering SNPs, the normal approach is to combine the entire data set, remove sites not genotyped in a sufficient number of individuals (typically 70-95%), and then filter out sites with a minor allele frequency (MAF) or minor allele count (MAC) that is not met globally (e.g. Lemopoulos et al., 2019; Mathur et al., 2019; Mulvena et al., 2020). Simulations suggest that a MAC ≥ 3 is optimal for detecting population structure (Linck & Battey, 2019), to avoid bias from singletons and doubletons.

Population comparisons using filtered polymorphic sites: effects of sample size

We start with a simple comparison of how global (N) and local (n) sample size affects SNP heterozygosity estimates. For this we use the ten subsamples (n = 10, 5, 4, 3, 2), which we analyse first individually (i.e. with each subsample run in a separate Stacks run) and then together (i.e. where the ten subsamples are run in a single Stacks run). Filtering followed a standard approach, retaining a single SNP from each RAD locus (--write-single-snp) which had no more than 20% missing data and that had a MAC ≥ 3. Observed and expected heterozygosities were estimated from these filtered polymorphic sites.

SNP heterozygosity estimates are shown in Figs 1a-d and indicate how this type of filtering approach presents problems for comparing estimates across studies. In all cases, heterozygosity was larger when fewer samples were used for estimation. Observed and expected heterozygosity estimates are both strongly affected by sample size, and effects are stronger when subsamples are analysed individually.

The source of this issue is that heterozygosity is generally lower for SNPs with rare alleles (where most individuals are homozygous for the common allele) than for SNPs with common alleles. For instance, for the n = 3 subsamples analysed in individual runs (Fig. 1a, c) all SNPs have minor alleles at 0.5 frequency when MAC ≥ 3 is applied, leading to expected heterozygosity of 0.5. As additional samples are added, SNPs with rare alleles become more likely to be detected, leading to lower heterozygosity estimates (c.f. Fig. S1). As MAC filtering is applied globally, heterozygosity is lower when populations are analysed together (Fig. 1b, d) as global sample size is ten times larger in these runs. However, even in these runs there were clear differences between SNP heterozygosity estimates for n = 10 (N = 100) and n ≤ 5 (N ≤ 50). These problems should diminish if singletons and doubletons are included because rare alleles are then more likely to be detected in small samples, but they will nevertheless persist (c.f. Fig. S1).

To investigate these effects at n ≥ 5, we use the nested subsamples from Groups A and B, first analysing each subsample from Group A in individual runs, then analysing each pair of subsamples of equal n from A and B together (Fig. 2). Here we see that sample size bias is present throughout, and specifically shows a bias from global N (total sample size in the analysis) rather than local n (sample size of each population). Although these effects reduce as N increases, they are nevertheless evident even with n = 40 and N ≥ 80.

Fig. 1: Observed (blue) and expected (red) SNP heterozygosity estimated from ten subsamples of mosquitoes at each n, with MAC ≥ 3 filtering applied to the population subsamples in individual runs (a, c) or when analysed together (b, d). Note that for MAC ≥ 3, individual analysis of n = 2 subsamples is not possible.

Fig. 2: Observed (blue) and expected (red) SNP heterozygosity estimated from mosquito samples of different size (n) and filtered at MAC ≥ 3. We compared estimates for Group A only with those of Group A and Group B combined in one analysis. Shading reflects similarity among numbers.

Population comparisons using filtered polymorphic sites: missing data thresholds

We investigated effects of missing data thresholds on observed and expected SNP heterozygosities using the nested subsamples from Group A, filtered with thresholds of 0% (i.e. no missing data allowed), 10%, 20%, 30%, 40%, or 50%. Thus in each case variation in heterozygosity was assessed in a single population of size n (n = 50, 40, 30, 20, 10, 5).

We see a considerable effect of missing data thresholds on SNP heterozygosity (Fig. 3). Samples of larger n were more strongly affected by choice of missing data threshold, with stringent filtering tending to produce higher estimates. When 50 individuals were used, filtering with a 10% missing data threshold (a common parameter setting) produced an estimate for observed heterozygosity 1.22 times higher than filtering with a 30% threshold (also a common parameter setting). This effect was less pronounced for expected heterozygosity (1.09 times higher) at n = 50, and this trend began to reverse at lower n.

The higher observed heterozygosity estimates at 0% versus 50% thresholds might be expected if there is a correlation between errors and the presence of missing data at a site. In that case, errors at monomorphic sites with high missing data could be read as low frequency polymorphisms, pushing down heterozygosity estimates when N is large. We expect this problem would increase if singletons and doubletons are included, as more low-frequency errors would be included in the estimates (c.f. Fig. S4).

Finally, the large population sizes in *Ae. aegypti* indicate this Malaysian sample is unlikely to contain inbred individuals, and thus we do not expect observed and expected heterozygosities to differ substantially. We note that when estimates of observed and expected heterozygosity

are compared, these parameters are most similar when filtering with a 0% missing data threshold, and they start to diverge as this threshold is increased. Less stringent missing data thresholds may thus introduce artefacts of differential observed and expected heterozygosities, which may lead to incorrect inferences of local breeding patterns.

Fig. 3: Effect of missing data thresholds on observed (a) and expected (b) SNP heterozygosity estimates from the Group A mosquito subsamples filtered at MAC ≥ 3.

Population comparisons using all polymorphic sites

Considering these inconsistencies in SNP heterozygosity estimates when filtering datasets with 'typical' settings for genetic structure, we reran the above analyses with settings of MAC ≥ 1 and selecting all SNPs rather than one per RAD locus. These analyses thus considered variation at all polymorphic sites including those with singletons and doubletons. We used a maximum

missing data threshold of 0% (or as specified), to avoid potential artefacts caused by including sites with missing data (Fig. 3).

Including all polymorphic sites reduced the bias from sample size; however, similar patterns of strong bias were still observed (Figs S2-S3). Also, as predicted, missing data thresholds had a larger effect on estimates when singletons and doubletons were included (Fig. S4). This lends support to the expectation that correlations between errors and the degree of missing data at a site drive heterozygosity estimates down when missing data thresholds are less stringent.

In light of these inconsistencies SNP heterozygosity appears prone to bias, regardless of whether filtering follows a typical protocol used for genetic structure or when considering every polymorphic site. This bias is demonstrated in Fig. S1, which shows how, when calculating SNP heterozygosity, the numerator remains proportionate to the number of heterozygous sites regardless of sample size, but the denominator is consistently biased downwards. This downward bias should diminish for very large N but for rare populations or small budgets this cannot be solved by sequencing more individuals. This limits the potential for SNP heterozygosity estimates in one study to inform the results of other studies.

Estimates based on all polymorphic and monomorphic sites (autosomal heterozygosity)

Given the above challenges, we next explored how sample size and missing data affect autosomal heterozygosity, which considers both monomorphic and polymorphic nucleotides. We ran analyses on identical datasets to those used previously. For filtering, we used no MAC cut-off, and estimated heterozygosity across every site rather than every polymorphic site. In the output from the Stacks.v2 program "Populations", this corresponds to the entries in the "# All positions (variant and fixed)" subsection. We used a maximum missing data threshold of 0% (unless specified differently).

Population comparisons using all sites: effects of sample size

When considering variation at all nucleotide sites, observed and expected heterozygosity estimates are far less affected by N than SNP heterozygosity estimates (Figs 4, 5). Though there is some variability among subsamples when analysed individually, when analysed together estimates for n ≥ 10 for the multiple subsamples (Fig. 4) and nested subsamples (Fig. 5) indicated consistent observed heterozygosities of ~0.00039 and expected heterozygosities of ~0.00040. The similar estimates for these two parameters match expectations for this sample of Malaysian *Ae. aegypti*, where inbreeding is unlikely given the large size of mosquito populations and the spatial distribution of sampling.

Fig. 4: Observed (blue) and expected (red) autosomal heterozygosity estimated from ten subsamples of mosquitoes at each n, with the population subsamples analysed in individual runs (a, c) or analysed together (b, d).

These consistent estimates are expected when all sites are taken into account because for smaller samples the higher frequency of heterozygotes at polymorphic sites will be offset by the lower number of polymorphic sites overall. Heterozygosity estimates from a set of individuals thus correlate directly with population heterozygosity because sites are not first

filtered by polymorphism (c.f. Fig. S1). In this sense, autosomal heterozygosity is a parameter that is both more robust to variation in study design and also a more accurate measure of genetic variation which can be used in comparisons across studies and organisms (e.g. Westbury et al., 2019).

Fig. 5: Observed (blue) and expected (red) autosomal heterozygosity estimated from mosquito samples of different size (n). We compared estimates for Group A only with those of Group A and Group B combined in one analysis. Shading reflects similarity among numbers.

Population comparisons with all sites: effects of missing data thresholds

Missing data thresholds had a smaller effect on autosomal heterozygosity than on SNP heterozygosity (Fig. 6; c.f. Figs 3, S4). Nevertheless, a similar problematic pattern is clear in the divergence between observed and expected heterozygosities when sites with missing data are included. At n = 50, filtering with a 0% missing data threshold estimated expected heterozygosity as 1.08 times larger than observed heterozygosity, while for a 20% threshold it was 1.29 times larger.

Considering these results, we propose that heterozygosity estimation should exclude nucleotide sites that have any missing genotypes, as these may be more likely to contain errors or otherwise skew parameter estimates. While this filtering might at first appear overly stringent, autosomal heterozygosity is calculated using far more sites than SNP heterozygosity, and should normally be based on sufficient sites even after strict filtering. For example, when a 20% missing data threshold is used to estimate SNP heterozygosity (MAC ≥ 1), heterozygosity is estimated from 95,293 polymorphic sites. When a 0% missing data threshold is used to estimate autosomal heterozygosity (MAC ≥ 0), heterozygosity is estimated from 7,813,360 sites,

of which 17,968 are polymorphic. The specific number of nucleotides used in autosomal heterozygosity calculations may vary across studies, but these should accurately reflect the degree of variation across the genome.

Fig. 6: Effect of missing data thresholds on observed (a) and expected (b) autosomal heterozygosity estimates from the Group A mosquito samples.

Multiple population considerations

We first estimated heterozygosity for the four *K. scurra* populations using equal sample sizes of 10. Fig. S5 compares results of a 0% missing data threshold against a 20% threshold, showing that at 20%, the ratio of observed to expected heterozygosity is higher (x̄ = 1.156) than at 0% (x̄

= 1.091), supporting our previous findings that missing data may bias the ratio of heterozygosities. Accordingly, we used a 0% missing data threshold in the following analyses.

Population comparisons: local sample size variation

Although global sample size analysed above had little effect on autosomal heterozygosity when n ≥ 10, we have yet to consider differences in sample size among populations. We estimated heterozygosity for the four *K. scurra* populations with one population (Goulburn) set at either half (5,10,10,10) or double (10,5,5,5) the size of the other populations compared to an equal population size. We compared results for autosomal heterozygosity (Fig. 7a) and SNP heterozygosity (Fig. 7b).

Fig. 7: Estimates of observed (blue) and expected (red) autosomal heterozygosity (a, white text) and SNP heterozygosity (b, black text) in four *K. scurra* populations of equal or unequal sample size. Numbers in brackets indicate sample sizes for the four populations sequentially (Goulburn to Wallendbeen). Shading reflects similarity among numbers.

When 10 individuals are analysed from each population, the Goulburn, Hall and Wallendbeen populations all have similar autosomal heterozygosities, while Cooma is much lower (Fig. 7a). There was no noticeable effect from unequal sample size. However, unequal sample size had

clear effects on SNP heterozygosity (Fig. 7b). Our previous observation that variation in N had a global effect on SNP heterozygosity (Fig 1) was again evident, and here revealed that variation in n affected local estimates. This bias could lead to misinterpretation of relative SNP heterozygosities within studies when populations with different sample sizes are included in analyses.

Population comparisons: impact of population structure

Next, we investigate the effects of combining genetically differentiated populations in an analysis. In terms of mtDNA variation and DArT SNPs, Cooma was separate from the other populations and particularly Wallendbeen (Hoffmann et al., 2020). We estimated heterozygosity for each of the four *K. scurra* populations analysed individually (i.e. with each population in a separate Stacks run) and compared this to populations analysed together (i.e. with all populations in a single Stacks run). We used the same 0% missing data threshold as before, and compared results for autosomal heterozygosity (Fig. 8a) and SNP heterozygosity (Fig. 8b).

Autosomal heterozygosity is unaffected by whether differentiated populations are analysed individually or together (Fig. 8a). However, strong biases on SNP heterozygosity are evident (Fig. 8b). When the four populations are analysed individually, estimates are much higher than when analysed together. Additionally, the population at Cooma, which otherwise recorded the lowest heterozygosity of the four populations, has higher heterozygosity than all other populations when analysed by itself.

As allele frequencies vary among these populations, many variant sites will only be polymorphic in one or two populations and are monomorphic in the others. Thus when populations are analysed together, estimates for each population will include these variant sites that are locally monomorphic, leading to lower heterozygosity estimates than when analysed individually.

A similar explanation accounts for the sharp variation in estimates at Cooma. When analysed individually, Cooma recorded fewer polymorphic sites (3235) than the other populations (4693, 4734, 5182); this pattern was also observed when populations were analysed together. However, heterozygosity at these 3235 polymorphic sites was higher than at the other populations.

These findings show how SNP heterozygosity estimates represent different parameters when populations are analysed in individual runs compared to when they are analysed alongside other populations. When analysed individually, the SNP heterozygosity of a population is equal to autosomal heterozygosity multiplied by the proportion of sites that are polymorphic (c.f. Fig. S1). When analysed with other populations, the SNP heterozygosity of a population will be shaped by whichever other populations are included, as the structure between these populations will determine which locally monomorphic sites are called as SNPs. Accordingly, SNP heterozygosities, if they are to be reported, should probably be calculated from populations analysed individually, and the total number of variant sites should also be reported for each population to provide further context to the heterozygosity estimates.

Fig. 8: Estimates of observed (blue) and expected (red) autosomal heterozygosity (a, white text) and SNP heterozygosity (b, black text) in four *K. scurra* populations when estimated individually or when the four populations were combined. Shading reflects similarity among numbers.

Comparing heterozygosity estimates

A final consideration concerns how to interpret heterozygosity estimates across studies. We have proposed several guidelines for filtering data to allow cross-study comparisons. The most important of these is that heterozygosity estimates should be derived from variation at both monomorphic and polymorphic sites. Table 1 compares variation in SNP heterozygosity with variation in autosomal heterozygosity for the four *K. scurra* populations analysed individually. We did not compare populations when analysed together due to the confoundment of SNP heterozygosities in these analyses (Fig 8b).

For *K. scurra*, variation in autosomal heterozygosity is approximately twice as large as variation in SNP heterozygosity (Table 1). A large difference in SNP heterozygosity might not be detected even when comparing populations with very low and very high levels of genetic variation

because the exclusion of monomorphic sites in each population will reduce differences in genetic variability among the populations.

Table 1. Comparison of observed and expected SNP heterozygosity and autosomal heterozygosity in four *K. scurra* populations when estimated individually. Calculations are based on the results from Fig. 8. MAX/MIN is the ratio between the largest score and the smallest score.

	SNP heterozygosity		Autosomal heterozygosity	
	Observed	Expected	Observed	Expected
MAX/MIN	1.164	1.064	1.983	2.230
Coefficient of variance	0.063	0.024	0.236	0.267

Discussion

Comparisons of heterozygosity across populations and species are frequently used to inform management decisions in conservation programs. An example of a relevant management action involves selecting populations for genetic and evolutionary rescue, which aims to decrease levels of inbreeding and increase levels of genetic variation in target populations through targeted introductions of individuals from other populations (Whiteley, Fitzpatrick, Funk, & Tallmon, 2015; Hoffmann et al., 2020). The usefulness of source populations for conservation translocations is to a large extent determined by their genetic variability (Reid et al., 2016; Ørsted et al., 2019). Tracking changes in heterozygosity across time can also be a worthwhile means of tracking the genetic health of threatened populations (Mitrovski, Hoffmann, Heinze, & Weeks, 2008) and is particularly useful for determining outcomes of management interventions (Weeks et al., 2017). All of these objectives require that heterozygosity estimates are comparable across populations within a study and across different studies.

In this paper, we have shown that filtering genome-wide sequence data using optimal settings for detecting genetic structure will produce heterozygosity estimates that are poorly-suited to comparisons across studies. Specifically we show that heterozygosity estimates that consider only polymorphic sites (SNP heterozygosity) are always biased by sample size, with larger estimates resulting from smaller global sample size (N) and smaller local sample size (n) relative to other populations. Heterozygosity estimates that consider monomorphic and polymorphic sites (autosomal heterozygosity) do not suffer from these biases. We also found that when sites with missing data are included, observed and expected heterozygosity estimates diverge, with the divergence proportional to the amount of missing data permitted.

Following this, we propose two general guidelines that should help meaningful comparisons: (i) studies aiming to summarise population genetic variation should report autosomal heterozygosity, either by itself or alongside SNP heterozygosity; (ii) sites with missing data should be omitted from heterozygosity calculations. Although we have not explicitly investigated the importance of sequencing coverage, this is widely known to be critical for accurately identifying heterozygotes (Nielsen et al., 2011). This being the case, our findings that heterozygosity estimates can be consistent even at low n (Fig. 7a) point to the optimal design for heterozygosity being deep sequencing of a small number of individuals (perhaps 5-10) from each population, rather than shallower sequencing of many individuals. Previous work has found these numbers to be adequate (Nazareno, Bemmels, Dick, & Lohmann, 2017); although Nazareno et al (2017) pre-filtered their dataset globally to retain only polymorphic sites, they did not refilter for polymorphisms when subsampling this dataset and thus their results would not have been subject to sampling size biases within the study.

SNP heterozygosity is frequently the only measure of heterozygosity reported (e.g. Jones et al., 2012; Chen et al., 2016; Bock et al., 2018; Mathur et al., 2019; Surbakti et al., 2020). Although there may be cases where SNP heterozygosity is the appropriate parameter, it will be subject to biases from sample size that do not affect autosomal heterozygosity, making the latter a better 'default' choice for reporting variation in populations. As taxa of conservation interest are frequently rare or difficult to sample in large numbers, large sample sizes for all populations would be difficult to achieve. SNP heterozygosity also does not capture the actual level of overall variation in the genome. For instance, *K. scurra* from Cooma had a higher SNP heterozygosity estimate (when calculated at the population level independently) yet fewer polymorphic sites than the other populations; this low level of polymorphism was evident in the autosomal heterozygosity estimate. While we advocate autosomal heterozygosity as a default choice for reporting genome-wide averages, there may be circumstances where only one variable site or a specific set of variable sites are of interest. In this case, heterozygosity at this specific set of sites can also be estimated free of bias provided that these sites are not further filtered by polymorphism when analysed across populations. Sites should still be included in a heterozygosity estimate for a population even if the population is monomorphic for some of the sites.

Whole-genome sequencing studies frequently report autosomal rather than SNP heterozygosity (e.g. Gopalakrishnan et al., 2017; Westbury, Petersen, Garde, Heide-Jørgensen, & Lorenzen, 2019). However, this methodology has been less commonly applied in reduced-representation sequencing studies such as those using RADseq or DArTseq markers. One reason for this may be that whole-genome studies frequently analyse single individuals rather than a sample, and as SNP heterozygosity calculated from a single diploid individual will always be equal to 1 (c.f. Fig S1) this approach would not be considered. Another reason may be that reduced-representation protocols are commonly applied to wild populations of understudied organisms, where the first step is typically to analyse genetic structure among populations. Monomorphic sites are uninformative for most analyses of genetic structure and thus these will typically be removed during filtering, and this remaining set of SNPs will be used to estimate both genetic structure and heterozygosity. Here we propose that analysis of genetic structure (variation

between populations) and heterozygosity (variation within populations) be treated as distinct targets and filtered using different parameter settings.

When expected heterozygosity is higher than observed heterozygosity, it is often treated as evidence for local inbreeding (e.g. Hoffmann et al., 2020), an important parameter that can indicate a need for genetic intervention in threatened species (Ralls et al., 2018). However, Figs 3, 6, and S4 show that more permissive filtering thresholds for missing data may be skewing the ratio of heterozygosities, leading to $F_{IS} > 0$. Inferences of inbreeding in the presence of missing data may be confounded, as observed and expected heterozygosities quickly diverge when sites with missing data are included. These opposing effects of missing data filters on observed and expected heterozygosities are consistent with sequencing error rates being higher at nucleotide positions where there are some missing genotypes.

A possible explanation is that sequencing errors are more likely at monomorphic sites; these sites are more common, and more likely to be retained than errors at polymorphic sites which may introduce a third allele and be removed from the dataset (most filtering protocols retain only biallelic sites). Monomorphic sites with errors may therefore be coded as low-frequency SNPs. Including them could affect the observed and expected autosomal heterozygosity estimates differently depending on whether errors are mostly coded as homozygous or heterozygous. Sequencing errors affecting monomorphic sites could also account for observations on SNP heterozygosity (Figs 3, S4), where higher missing data thresholds result in a decrease in observed and expected heterozygosity (due to monomorphic loci being erroneous included as low-frequency SNPs).

Despite these issues, the *K. scurra* data also indicate clear instances where inbreeding levels differ across populations. We note that for *K. scurra* from Cooma, observed and expected heterozygosities were almost identical, while expected heterozygosities are 8-15% higher in the other populations, suggesting a situation where inbreeding in Cooma is low but genetic variability is also low. An accurate assessment of inbreeding in populations versus low genetic variation is important when making recommendations around genetic mixing of threatened populations, which can target both the masking of deleterious genes expressed as a consequence of inbreeding as well as problems from low levels of genetic variability (Weeks et al., 2011; Ralls et al., 2018; Hoffmann et al., 2020). It would be worth further evaluating optimal filtering strategies for assessing the ratio of observed and expected heterozygosities, using a set of populations where inbreeding is known to occur in some populations but not others.

This study has highlighted issues with estimating population heterozygosities from SNP data directly, and shown that autosomal heterozygosity estimates are more robust to the influence of sample size and are likely to be more comparable across studies. We provide general guidelines for estimating population heterozygosity from genome-wide sequence data that are usually different from guidelines for estimating other population genetic parameters such as gene flow, population structure, relatedness and effective population size. As in previous assessments (Linck & Battey, 2019), our results demonstrate that SNP datasets need to be carefully evaluated when they are used to obtain genetic parameters for populations that inform management decisions.

Acknowledgements

This research was supported by the Australian Research Council (Discovery Grant DP190100990), the Wellcome Trust (Grant no. 108508). and the National Health and Medical Research Council (Program Grant no. 1132412; Fellowship Grant no. 1118640). This research was facilitated by use of the Nectar Research Cloud.

Data Archiving

Aligned .bam files for 100 *Ae. aegypti* and 40 *K. scurra* will be made available through the NCBI SRA by time of publication.

Author Contributions

AAH and ARW conceived the study; TLS, MJ, AAH and ARW designed the methodology; TLS and MJ analysed the data; TLS, AAH and ARW led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

References

Bock, D. G., Kantar, M. B., Caseys, C., Matthey-Doret, R., & Rieseberg, L. H. (2018). Evolution of invasiveness by genetic accommodation. *Nature Ecology and Evolution*, *2*(6), 991–999. doi:10.1038/s41559-018-0553-z

Campbell, M. R., Vu, N. V., LaGrange, A. P., Hardy, R. S., Ross, T. J., & Narum, S. R. (2019). Development and Application of Single-Nucleotide Polymorphism (<scp>SNP</scp>) Genetic Markers for Conservation Monitoring of Burbot Populations. *Transactions of the American Fisheries Society*, *148*(3), 661–670. doi:10.1002/tafs.10157

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool set for population genomics. *Molecular Ecology*, *22*(11), 3124–40. doi:10.1111/mec.12354

Chen, N., Cosgrove, E. J., Bowman, R., Fitzpatrick, J. W., & Clark, A. G. (2016). Genomic Consequences of Population Decline in the Endangered Florida Scrub-Jay. *Current Biology*, *26*(21), 2974–2979. doi:10.1016/j.cub.2016.08.062

Fitzpatrick, S. W., Bradburd, G. S., Kremer, C. T., Salerno, P. E., Angeloni, L. M., & Funk, W. C. (2020). Genomic and Fitness Consequences of Genetic Rescue in Wild Populations. *Current Biology*, *30*(3), 517-522.e5. doi:10.1016/j.cub.2019.11.062

Fontaine, A., Filipović, I., Fansiri, T., Hoffmann, A. A., Cheng, C., Kirkpatrick, M., … Lambrechts, L. (2017). Extensive genetic differentiation between homomorphic sex chromosomes in the mosquito vector, *Aedes aegypti*. *Genome Biology and Evolution*, *9*(9), 2322–2335. doi:10.1093/gbe/evx171

Garner, B. A., Hand, B. K., Amish, S. J., Bernatchez, L., Foster, J. T., Miller, K. M., … Luikart, G. (2016,

February 1). Genomics in Conservation: Case Studies and Bridging the Gap between Data and Application. *Trends in Ecology and Evolution*. Elsevier Ltd. doi:10.1016/j.tree.2015.10.009

Gopalakrishnan, S., Samaniego Castruita, J. A., Sinding, M. H. S., Kuderna, L. F. K., Räikkönen, J., Petersen, B., … Gilbert, M. T. P. (2017). The wolf reference genome sequence (Canis lupus lupus) and its implications for Canis spp. population genomics. *BMC Genomics*, *18*(1), 495. doi:10.1186/s12864-017-3883-3

Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2017). DARTR: An R package to facilitate analysis of SNP data generated from reduced representation genome sequencing. *Molecular Ecology Resources*, *18*(3), 1–9. doi:10.1111/1755-0998.12745

Hoffmann, A. A., Sgrò, C. M., & Kristensen, T. N. (2017, July 1). Revisiting Adaptive Potential, Population Size, and Conservation. *Trends in Ecology and Evolution*. Elsevier Ltd. doi:10.1016/j.tree.2017.03.012

Hoffmann, A. A., White, V., Jasper, M., Yagui, H., Sinclair, S., & Kearney, M. (2020). An endangered flightless grasshopper with strong genetic structure maintains population genetic variation despite extensive habitat loss. *Authorea Preprints*. doi:10.22541/AU.160403053.38828134/V1

Hohenlohe, P. A., Bassham, S., Etter, P. D., Stiffler, N., Johnson, E. A., & Cresko, W. A. (2010). Population Genomics of Parallel Adaptation in Threespine Stickleback using Sequenced RAD Tags. *PLoS Genetics*, *6*(2), e1000862. doi:10.1371/journal.pgen.1000862

Jasper, M., Schmidt, T. L., Ahmad, N. W., Sinkins, S. P., & Hoffmann, A. A. (2019). A genomic approach to inferring kinship reveals limited intergenerational dispersal in the yellow fever mosquito. *Molecular Ecology Resources*, *19*(5), 1254–1264. doi:10.1111/1755-0998.13043

Jones, F. C., Chan, Y. F., Schmutz, J., Grimwood, J., Brady, S. D., Southwick, A. M., … Kingsley, D. M. (2012). A genome-wide SNP genotyping array reveals patterns of global and repeated species-pair divergence in sticklebacks. *Current Biology*, *22*(1), 83–90. doi:10.1016/j.cub.2011.11.045

Lemopoulos, A., Prokkola, J. M., Uusi-Heikkilä, S., Vasemägi, A., Huusko, A., Hyvärinen, P., … Vainikka, A. (2019). Comparing RADseq and microsatellites for estimating genetic diversity and relatedness — Implications for brown trout conservation. *Ecology and Evolution*, *9*(4), 2106–2120. doi:10.1002/ece3.4905

Linck, E., & Battey, C. J. (2019). Minor allele frequency thresholds strongly affect population structure inference with genomic data sets. *Molecular Ecology Resources*, *19*(3), 639–647. doi:10.1111/1755-0998.12995

Maroso, F., Franch, R., Dalla Rovere, G., Arculeo, M., & Bargelloni, L. (2016). RAD SNP markers as a tool for conservation of dolphinfish *Coryphaena hippurus* in the Mediterranean Sea: Identification of subtle genetic structure and assessment of populations sex-ratios. *Marine Genomics*, *28*, 57–62. doi:10.1016/j.margen.2016.07.003

Mathur, S., Tomeček, J. M., Heniff, A., Luna, R., & DeWoody, J. A. (2019). Evidence of genetic erosion in a peripheral population of a North American game bird: the Montezuma quail (*Cyrtonyx montezumae*). *Conservation Genetics*, *20*(6), 1369–1381. doi:10.1007/s10592-019-01218-9

Mitrovski, P., Hoffmann, A. A., Heinze, D. A., & Weeks, A. R. (2008). Rapid loss of genetic variation in an endangered possum. *Biology Letters*, *4*(1), 134–138. doi:10.1098/rsbl.2007.0454

Mulvena, S. R., Pierson, J. C., Farquharson, K. A., McLennan, E. A., Hogg, C. J., & Grueber, C. E. (2020). Investigating inbreeding in a free-ranging, captive population of an Australian marsupial. *Conservation Genetics*, *21*(4), 665–675. doi:10.1007/s10592-020-01278-2

Nazareno, A. G., Bemmels, J. B., Dick, C. W., & Lohmann, L. G. (2017). Minimum sample sizes for population genomics: an empirical study from an Amazonian plant species. *Molecular Ecology Resources*, *17*(6), 1136–1147. doi:10.1111/1755-0998.12654

Nielsen, R., Paul, J. S., Albrechtsen, A., & Song, Y. S. (2011). Genotype and SNP calling from next-generation sequencing data. *Nature Reviews Genetics*, *12*(6), 443–451. doi:10.1038/nrg2986

Ørsted, M., Hoffmann, A. A., Sverrisdóttir, E., Nielsen, K. L., & Kristensen, T. N. (2019). Genomic variation predicts adaptive evolutionary responses better than population bottleneck history. *PLoS Genetics*, *15*(6). doi:10.1371/journal.pgen.1008205

Ralls, K., Ballou, J. D., Dudash, M. R., Eldridge, M. D. B., Fenster, C. B., Lacy, R. C., … Frankham, R. (2018). Call for a Paradigm Shift in the Genetic Management of Fragmented Populations. *Conservation Letters*, *11*(2), e12412. doi:10.1111/conl.12412

Reid, N. M., Proestou, D. A., Clark, B. W., Warren, W. C., Colbourne, J. K., Shaw, J. R., … Whitehead, A. (2016). The genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild fish. *Science*, *354*(6317), 1305–1308. doi:10.1126/science.aah4993

Ritland, K. (1996). Estimators for pairwise relatedness and individual inbreeding coefficients. *Genetical Research*, *67*(2), 175–185. doi:10.1017/s0016672300033620

Surbakti, S., Parker, H. G., McIntyre, J. K., Maury, H. K., Cairns, K. M., Selvig, M., … Ostrander, E. A. (2020). New Guinea highland wild dogs are the original New Guinea singing dogs. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(39), 24369–24376. doi:10.1073/pnas.2007242117

Weeks, A. R., Heinze, D., Perrin, L., Stoklosa, J., Hoffmann, A. A., Van Rooyen, A., … Mansergh, I. (2017). Genetic rescue increases fitness and aids rapid recovery of an endangered marsupial population. *Nature Communications*, *8*(1). doi:10.1038/s41467-017-01182-3

Weeks, A. R., Sgro, C. M., Young, A. G., Frankham, R., Mitchell, N. J., Miller, K. A., … Hoffmann, A. A. (2011). Assessing the benefits and risks of translocations in changing environments: A genetic perspective. *Evolutionary Applications*, *4*(6), 709–725. doi:10.1111/j.1752-4571.2011.00192.x

Westbury, M. V., Petersen, B., Garde, E., Heide-Jørgensen, M. P., & Lorenzen, E. D. (2019). Narwhal Genome Reveals Long-Term Low Genetic Diversity despite Current Large Abundance Size. *IScience*, *15*, 592–599. doi:10.1016/j.isci.2019.03.023

Whiteley, A. R., Fitzpatrick, S. W., Funk, W. C., & Tallmon, D. A. (2015, January 1). Genetic rescue to the rescue. *Trends in Ecology and Evolution*. Elsevier Ltd. doi:10.1016/j.tree.2014.10.009

Wright, B. R., Grueber, C. E., Lott, M. J., Belov, K., Johnson, R. N., & Hogg, C. J. (2019). Impact of reduced-representation sequencing protocols on detecting population structure in a threatened marsupial. *Molecular Biology Reports*, *46*(5), 5575–5580. doi:10.1007/s11033-019-04966-6

Unbiased population heterozygosity estimates from genome-wide sequence data

Thomas L Schmidt, Moshe Jasper, Andrew R Weeks, Ary A Hoffmann

Supplementary Figures S1 – S5

a b c d e f g h i j k l m n o   Ho

a 1 $\frac{3}{15} = 0.20$

b $\frac{3}{3} = 1$

c 1 2 3 4 5 $\frac{13}{75} = 0.17$

d $\frac{13}{30} = 0.43$

e 1 2 3 4 5 6 7 8 9 10 $\frac{29}{150} = 0.19$

f $\frac{29}{90} = 0.32$

g 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 $\frac{61}{300} = 0.20$

h $\frac{61}{220} = 0.28$

Fig S1. Estimating heterozygosity in a hypothetical population of diploid individuals.

Analyses consider population samples of n = 1 (**a**,**b**), 5 (**c**,**d**), 10 (**e**,**f**) and 20 (**g**,**h**). Observed heterozygosity is calculated using all sites irrespective of polymorphism (autosomal heterozygosity: **a**,**c**,**e**,**g**) or using SNPs only (SNP heterozygosity: **b**,**d**,**f**,**h**). Individuals have been genotyped at 15 sites, and the circles indicate the unphased genotype at each site. When autosomal heterozygosity is calculated, the numerator and denominator remain proportionate to the number of sites evaluated (denominator) and the number of heterozygous sites (numerator) regardless of sample size. When SNP heterozygosity is calculated, the denominator is biased by whether SNPs with rare alleles are detected in the sample, which will remain an issue even at very large n. These biases will be present in both observed and expected heterozygosity. Note that in actual sequence data the proportion of monomorphic sites will be much higher than shown here, producing smaller autosomal heterozygosity estimates.

Fig S2. Observed (blue) and expected (red) SNP heterozygosity estimated from ten subsamples of mosquitoes at each n, with MAC ≥ 1 filtering applied to the population subsamples in individual runs (a, c) or when analysed together (b, d).

Observed				Expected		
n	**Group A only**	**Group A**	**Group B**	**Group A only**	**Group A**	**Group B**
50	0.16818	0.14005	0.14171	0.18127	0.14815	0.14880
40	0.17529	0.14171	0.14728	0.19102	0.14880	0.15780
30	0.18202	0.14728	0.15246	0.20110	0.15780	0.15813
20	0.19584	0.15246	0.15467	0.21869	0.15813	0.16737
10	0.22403	0.15467	0.16125	0.25537	0.16737	0.16777
5	0.27067	0.16125	0.17075	0.30818	0.16777	0.18551

Fig S3. Observed and expected SNP heterozygosity estimated from mosquito samples of different size taken from Group A and Group B and filtered at MAC ≥ 1. We compared estimates for Group A only with those of Group A and Group B combined in one analysis. Shading reflects the relative size of the heterozygosity estimates.

Fig S4. Effect of missing data thresholds on observed (a) and expected (b) SNP heterozygosity estimates from the Group A mosquito samples filtered at MAC ≥ 1.

Fig S5. Estimates of observed and expected autosomal heterozygosity in four *K. scurra* populations filtered at 0% missing vs 20% missing data. The mean heterozygosity ratio between observed and expected values at 0% is 1.092, and at 20% it is 1.156. Entries with similar shading are more similar to each other.